Search for sterile neutrinos at the DANSS experiment

I Alekseev^{a,b,c}, V Belov^d, V Brudanin^{b,d}, M Danilov^{e,*}, V Egorov^{d,f}, D Filosofov^d, M Fomina^d, Z Hons^{d,g}

S Kazartsev^{d,f}, A Kobyakin^{a,c}, A Kuznetsov^d, I Machikhiliyan^a, D Medvedev^d, V Nesterov^a, A Olshevsky^d,

N Pogorelov^a, D Ponomarev^d, I Rozova^d, N Rumyantseva^d, V Rusinov^a, E Samigulin^{a,b}, Ye Shevchik^d, M Shirchenko^d,

Yu Shitov^{d,h}, N Skrobova^{a,c,e}, A Starostin^a, D Svirida^a, E Tarkovsky^a, J Vlášek^{d,g}, I Zhitnikov^d, D Zinatulina^d

^aAlikhanov Institute for Theoretical and Experimental Physics NRC "Kurchatov Institute",

B. Cheremushkinskaya 25, Moscow, 117218, Russia

^bNational Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Kashirskoe highway 31, Moscow, 115409, Russia

^cMoscow Institute of Physics and Technology, 9 Institutskiy per., Dolgoprudny, Moscow Region, 141701, Russia

^d Joint Institute for Nuclear Research, Joliot-Curie 6, Dubna, Moscow region, 141980, Russia

^eLebedev Physical Institute of the Russian Academy of Sciences, 53 Leninskiy Prospekt, Moscow, 119991, Russia

^fDubna State University, Universitetskaya 19, Dubna, Moscow Region, 141982, Russia

^gCzech Technical University in Praque, Zikova 1903/4, 166 36 Praque 6, Czechia

¹Dubna State University, University and State 1, University 1, University

performed calibrations with radioactive sources and reported the ratio of numbers of observed to predicted events of 0.88 ± 0.05 [3]. This deficit is the so called "Gallium" anomaly" (GA). Mueller et al. [4] made new estimates of the reactor $\tilde{\nu}_e$ flux which is about 6% higher than experimental measurements at small distances. This deficit is the so called "Reactor antineutrino anomaly" (RAA). Both

neutrinos [9].

The survival probability of a reactor $\tilde{\nu}_e$ at short distances in the 4ν mixing scenario (3 active and 1 sterile neutrino) is described by a familiar expression

$$1 - \sin^2 2\theta_{14} \sin^2 \left(\frac{1.27 \Delta m_{14}^2 [\text{eV}^2] L[\text{m}]}{E_{\nu} [\text{MeV}]} \right).$$
(1)

The existence of sterile neutrinos would manifest itself in distortions of the $\tilde{\nu}_e$ energy spectrum at short distances. At longer distances these distortions are smeared out and

^{*}Corresponding author

Email address: danilov@lebedev.ru (M Danilov)

only the rate is reduced by a factor of $1 - \sin^2(2\theta_{14})/2$. Measurements at only one distance from a reactor core are not sufficient since the theoretical description of the $\tilde{\nu}_e$ energy distribution is considered not to be reliable enough. The most reliable way to observe such distortions is to measure the $\tilde{\nu}_e$ spectrum with the same detector at different distances. In this case, the shape and normalization of the $\tilde{\nu}_e$ spectrum as well as the detector efficiency are canceled out. Detector positions should be changed frequently enough in order to cancel out time variations of the detector and reactor parameters. The DANSS experiment uses this strategy and measures $\tilde{\nu}_e$ spectra at 3 distances from the reactor core centre: 10.7 m, 11.7 m, and 12.7 m to the detector centre. The detector positions are changed typically 3 times a week. Antineutrinos are detected by means of the Inverse Beta Decay (IBD) reaction

$$\tilde{\nu}_e + p \to e^+ + n \text{ with } E_{\tilde{\nu}} = E_{e^+} + 1.80 \text{ MeV.}$$
 (2)

2. The DANSS Detector

The DANSS detector is described elsewhere [10]. Here we mention only a few essential features. The DANSS spectrometer does not contain any flammable or other dangerous liquids and may therefore be located very close to the core of a $3.1 \, \mathrm{GW}_{\mathrm{th}}$ industrial power reactor at the Kalinin Nuclear Power Plant (KNPP) 350 km NW of Moscow. DANSS is installed under the reactor core. The reactor cauldron, cooling pond, concrete and other materials provide a good shielding equivalent to $\sim 50 \text{ m}$ of water, which removes the hadronic component of the cosmic background and reduces the cosmic muon flux by a factor of 6. The size of the reactor core is quite big (3.7 m in height and 3.2 m in diameter) which leads to the smearing of the oscillation pattern. This drawback is compensated by a high $\tilde{\nu}_e$ flux of $\sim 5 \times 10^{13} \tilde{\nu}_e/\text{cm}^2/\text{s}$ at a distance of 11 m.

DANSS is a highly segmented plastic scintillator detector with a total volume of 1 m^3 , surrounded with a composite shield of copper (Cu — 5 cm), borated polyethylene (CHB — 8 cm), lead (Pb — 5 cm) and one more layer of borated polyethylene (CHB — 8 cm) (see Fig. 1). It is surrounded on 5 sides (excluding bottom) by double layers of 3 cm thick scintillator plates to veto cosmic muons.

The basic element of DANSS is a polystyrene-based extruded scintillator strip $(1 \times 4 \times 100 \text{ cm}^3)$ with a thin $(\sim 0.2 \text{ mm})$ Gd-containing surface coating. The amount of Gd in the detector is $0.35 \%_{wt}$. The coating serves as a light reflector and a (n, γ) -converter simultaneously.

Light from the strip is collected with three wavelengthshifting (WLS) Kuraray fibers Y-11, \oslash 1.2 mm, glued into grooves along the strip. One (blind) end of each fiber is polished and covered with a mirror paint, which decreases the total longitudinal attenuation of a light signal to about 30%/m. This non-uniformity can be corrected using information from orthogonal strips or from the neutron capture position, which is typically ~ 10 cm away from the positron production point. The response non-uniformity across the strip is $\sigma = 7.8\%$ [11]. This non-uniformity of the response can't be corrected. It leads, together with the energy losses in the inactive strip reflective layers, to the deterioration of the energy resolution in comparison with naive estimates from the photoelectron statistics.

Each 50 parallel strips are combined into a module, so that the whole detector (2500 strips) is a structure of 50 intercrossing modules. Each module is viewed with a compact photomultiplier tube (PMT) (Hamamatsu R7600U-300) coupled to all 50 strips of the module via 100 WLS fibers, two per strip. PMTs are placed inside the shielding but outside the copper layer, which serves also as a module frame. In addition, to get a more precise energy and space pattern of an event, each strip is equipped with an individual Silicon PhotoMultiplier (SiPM) (MPPC S12825-050C(X) coupled to the strip via the third WLS fiber. The SiPM is fixed directly at the end of the strip using a plastic light connector. All signals are digitized with specially designed 12 bit, 125 MHz FADCs[12]. Only front-end electronics is placed inside the shielding but outside the Cu layer. All other electronics is placed outside of the detector shielding. One 6U VME board serves 64 channels. SiPMs (PMTs) register about 18 (20) photoelectrons (p.e.) per MeV. These numbers were obtained using measurements with cosmic muons and artificially driven LEDs. So the total number is 38 p.e./MeV. Parameterized strip response non-uniformities have been incorporated into the GEANT4 (Version 4.10.4) MC simulation of the detector. The MC simulation included also a spread in the light yields of different strips, dead channels, Poisson fluctuations in the number of p.e. at the first 2 PMT dinodes, the excess noise factor for SiPMs due to the optical cross-talk between pixels. The experimental energy resolution for cosmic muon signals in the scintillator strips is 15 % worse than that from the MC calculation. Therefore, the MC estimations are scaled up by the corresponding factor. Figure 2 shows the simulated DANSS response to a 4.125 MeV positron signal. The energy resolution is modest. This leads to additional smearing of the oscillation pattern, comparable with the smearing due to the large reactor core size.

Figure 3 shows the energy distribution of neutron capture signals from a 248 Cm source placed at the center of the detector. Two peaks correspond to the neutron capture by Gd and by protons. Fits of the high energy parts of the peaks give a resolution compatible with the MC simulations.

Figure 4 shows the energy distribution of signals from a 60 Co γ -source placed in the center of the detector. The observed energy resolution of $\frac{\sigma}{E} = 20.3\%$ is consistent with the MC expectations. The same is true in case of a 22 Na source. The visible energy in the detector from the radioactive sources is lower than the source energy due to losses in inactive strip reflective layers, dead channels and photons escaped from the detector.

SiPM gain calibration was performed using noise sig-

nals typically every 5 days. Calibration with cosmic muons of all strips in the whole detector was also performed once in \sim 5 days. A detailed description of the calibration procedure is presented elsewhere [13]. Figure 5 shows the number of p.e. detected by a SiPM in one strip versus the energy deposited by cosmic muons. The strip response is linear within 0.7% in the range (1.7—4.7) MeV. The energy measured by PMTs is proportional on average to the energy, measured by SiPMs. Therefore, the PMT energy response is also linear. Positrons with energies higher than 4.7 MeV typically deposit their energy in several strips. Therefore, the detector response should be linear for high energies as well.




Figure 1: Simplified cross section of a corner of the DANSS detector.




Figure 2: Reconstructed positron energy for 4.125 MeV positrons from MC simulations.

3. Data taking and analysis

The trigger of the experiment is produced when the digital sum of all PMT signals is above 0.7 MeV or the energy in the veto system is larger than 4 MeV. As a consequence, the IBD process appears in the data as two distinct events, prompt and delayed. For each trigger, waveforms for all SiPMs and PMTs are recorded in 512 ns windows.




Figure 3: Energy spectrum of the delayed signals measured with the $^{248}\mathrm{Cm}$ neutron source.




Figure 4: Detector response to the ⁶⁰Co source.

The data analysis is performed in several stages. The first one is the noise cleanup. At this step an average event time is calculated and all hits in SiPMs more than 15 ns away are rejected. Then single pixel SiPM hits which have no confirmation by the corresponding PMT are also rejected. After these two steps less than 2 noise SiPM hits per event survive out of about 40 in the original sample. At the next stage various characteristics of the event such as the total energy, the number of hits and so on are calculated. At this stage we also look for a continuous ionization cluster and calculate its visible energy. The visible energy of a positron cluster is converted using MC simulations into the deposited energy by taking into account average losses in the inactive reflective layers of the strips and dead channels. Sometimes photons from the positron annihilation produce signals in the strips attributed to the positron cluster. This leads to an increase of the visible energy. Such a shift is also corrected on average using MC simulations. A typical size of the total correction is $\sim 2\%$. The next step is a search for the time-correlated pairs of prompt-delayed events. We start with searching for an




Figure 5: Number of p.e. detected with SiPMs vs energy deposited by cosmic muons in a scintillator strip (top) and the deviation from a straight line fit (bottom).

event with more than 3.5 MeV energy deposit. This is a delayed event candidate unless it has the muon veto (more than 4 MeV energy in a veto counter or more than 1 hit in the veto counters with a low threshold of ~ 0.15 MeV or more than 20 MeV energy in the main detector). Then we look backward in time searching for a prompt event with more than 1 MeV in the positron cluster and no muon veto. An IBD candidate pair is considered found if the time difference between the prompt and delayed events is in the range $(2-50) \mu s$. For a valid pair we also require no event with the muon veto within 60 μ s before the prompt signal (within 200 μ s if E > 300 MeV is released in the main detector). No other event should occur within 45 μ s before and 80 μ s after the prompt event. The found pairs of prompt and delayed events form the experimental sample of IBD candidates. Similar to the experimental sample, the accidental coincidence sample is formed by looking for a prompt signal in 16 regions: 5, 10,..., 80 ms before the neutron candidate. This sample provides us with a modelindependent measure of the accidental background. Distributions for IBD candidates, the accidental background and their difference, which represents the IBD signal without accidental background are presented in figures 6 and 7.

Figure 6 shows the distance between positron and neutron candidates. The width of the signal distribution is determined by the neutron travel in the detector before the capture and the uncertainty of the neutron capture vertex reconstruction from the observed γ -flash. The distribution of the neutron capture time from the positron production moment is presented in Fig. 7. These two figures illustrate also the method used to get signal distributions: a parameter is plotted for the experimental data set of the IBD candidates and for the accidental background data set using the same selection criteria. A signal distribution is obtained as a difference of these two distributions. It is worth mentioning that the very procedure of the accidental background subtraction only negligibly increases the statistical errors in the signal distributions. This is because the accidental background is calculated using the 16 times larger sample of pairs. However, the final errors are determined by the initial counts in the IBD candidate sample, which include the accidental background. Therefore, several cuts are applied, in order to reduce the accidental background. These cuts are designed to be very soft with respect to the signal in order to avoid any distortions. All cuts were selected without looking at the final results. They have been fixed after collection of about 10% of the data. The cuts include the following requirements:

- The distance between positron and neutron candidates should be shorter than 45 cm (55 cm) for a positron candidate reconstructed in 2 (3) dimensions;
- The additional energy outside a positron cluster should be less than 1.8 MeV, and the most energetic hit should have the energy less than 0.8 MeV;
- The number of hits in the delayed event is higher than 3;
- The number of hits in the prompt event outside the positron cluster is less than 11;
- The most energetic hit of the positron cluster lies in a fiducial volume of the detector which does not include outer strips in the X and Y directions as well as 4 highest and 4 lowest strips in the vertical Z direction.

Positron energy spectra for 3 detector positions (top, middle, bottom) are shown in Fig. 8 with statistical errors only. The corresponding numbers of events are 338244, 247753 and 324938. The IBD counting rate is 4910 events per day in the top position. The positron energy does not include annihilation photons and hence is 1.02 MeV lower than the usually used prompt energy. The muon-induced background (discussed below) is subtracted. Three other reactors at the KNPP are 160 m, 334 m, and 478 m away from the DANSS detector. The IBD counting rate from these reactors is 0.6% of the IBD counting rate from the nearest reactor for the top detector position. This contribution is taken into account by the corresponding reduction of the normalization of the obtained spectra.

The energy spectrum of the background from neutrons produced by muons inside the veto system is obtained from events with the muon veto. The amount of this background is determined from a fit of the positron candidate energy spectrum during reactor off periods using the shape of the background determined from events with the muon veto. This procedure reduces uncertainties in the background shape to a negligible level. A possible small uncertainty in the background rate is taken into account during




Figure 6: Distance between positron and neutron reconstructed positions. Errors are smaller than sizes of points.




Figure 7: Time between prompt and delayed signals. Errors are smaller than sizes of points.

systematic error studies. This is the most important background. It constitutes 2.7 % of the IBD rate at the top detector position.

The energy spectrum of the background from fast neutrons produced outside the detector shielding is estimated by a linear extrapolation from a (10-16) MeV region to lower energies. This background calculation and subtraction is performed separately for the positron candidate energy spectra with and without muon veto. This background constitutes ~ 0.1% of the IBD signal in the (1-8) MeV region.

A background from ⁹Li and ⁸He produced by cosmic muons was estimated using the time distribution between cosmic events with the energy deposit in the detector bigger than 800 MeV and the IBD candidates. No sign of the exponential decay component with known ⁹Li decay time of 257.2 ms was found. The corresponding upper limit on this type of background is 5.4 events/day at the 90% confidence level.




Figure 8: Positron energy distributions measured at different detector positions. Statistical errors only.

The shape of the positron spectrum agrees roughly with the MC predictions based on the $\tilde{\nu}_e$ spectrum from [14, 4]. However, a quantitative comparison requires additional studies of calibration and systematic errors and improvements in the MC simulation of the detector. Since the results of the present analysis practically do not depend on the $\tilde{\nu}_e$ spectrum shape and normalization we postpone these studies till a forthcoming paper.

Figure 9 shows the ratio of positron energy spectra at the bottom and top detector positions.




Figure 9: Ratio of positron energy spectra measured at the bottom and top detector positions (statistical errors only). The dashed line is the prediction for 3ν case ($\chi^2 = 35.0$, 24 degrees of freedom). The solid curve is the prediction for the best fit in the 4ν mixing scenario ($\chi^2 = 21.9$, $\sin^2 2\theta_{14} = 0.05$, $\Delta m_{14}^2 = 1.4 \text{ eV}^2$). The dotted curve is the expectation for the optimum point from the RAA and GA fit [5] ($\chi^2 = 83$, $\sin^2 2\theta_{14} = 0.14$, $\Delta m_{14}^2 = 2.3 \text{ eV}^2$)

The exclusion area in the sterile neutrino parameter space was calculated using the Gaussian CL_s method [15] assuming only one type of sterile neutrinos. For a grid of points in the Δm_{14}^2 , $\sin^2 2\theta_{14}$ plane predictions for the ratio $R^{pre}(E)$ of positron spectra at the bottom and top detector positions were calculated. Calculations included the MC integration over the $\tilde{\nu}_e$ production point in the reactor core, $\tilde{\nu}_e$ detection point in the detector, and positron energy resolution. The $\tilde{\nu}_e$ production point distributions in the reactor core were provided by the KNPP for different time periods. These distributions were almost flat in the reactor core radius and height with a fast drop near the edges of the core. The distribution averaged over the campaign was used in the calculations. It was checked that this approximation practically did not influence the final results. The $\tilde{\nu}_e$ energy spectrum from [14, 4] averaged over the campaign was used for the calculations. However, the final result practically did not depend on the shape and normalization of the $\tilde{\nu}_e$ spectrum since only the ratio of measured positron spectra at the different positions was used in the analysis. This was checked by repeating the analysis using the spectrum from the beginning of the campaign.

The obtained theoretical prediction for a given point in the Δm_{14}^2 , $\sin^2 2\theta_{14}$ plane was compared with the prediction for the three neutrino case using the Gaussian CL_s method for the 90% confidence level (CL) exclusion area estimation. The difference in χ^2 for the two hypotheses $\Delta \chi^2 = \chi^2_{4\nu} - \chi^2_{3\nu}$ was used for the comparison. This difference in χ^2 has a Gaussian distribution with the mean value and width determined from the fit to the Asimov data set, an infinite statistics data sample with parameters fixed to the 4ν or 3ν hypotheses [15]. Confidence levels for the 4ν (CL_{4 ν}) and 3ν (CL_{3 ν}) hypotheses were calculated using the obtained parameters of the Gaussian distributions for these hypotheses and the observed value of $\Delta \chi^2$. These confidence levels provide a measure for the consistency with the 4ν and 3ν hypotheses correspondingly and $CL_s = CL_{4\nu}/CL_{3\nu}$. The 4ν hypothesis for a given point in the sterile neutrino parameter space is excluded at the 90% CL if $CL_s < 1-0.9$. The CL_s method is more conservative than the usual Confidence Interval method.

The χ^2 for each hypothesis was constructed using 24 data points R_i^{obs} in the (1—7) MeV positron energy range

$$\chi^{2} = \sum_{i=1}^{N} (R_{i}^{obs} - k \times R_{i}^{pre})^{2} / \sigma_{i}^{2}, \qquad (3)$$

where R_i^{obs} (R_i^{pre}) is the observed (predicted) ratio of $\tilde{\nu}_e$ counting rates at the two detector positions and σ_i is the statistical standard deviation of R_i^{obs} , and k is a normalization factor left free in the fit to compensate possible changes in the overall detector efficiency and in the $\tilde{\nu}_e$ production point distribution in the reactor core.

The oscillations due to the known neutrinos were neglected since at such short distances they do not change the $\tilde{\nu}_e$ spectrum in the studied energy range. The procedure was repeated for all points of the grid in order to get the whole exclusion area. Influence of systematic uncertainties in the parameters was estimated by repeating the analysis with different values of parameters. A point in the Δm_{14}^2 , sin² $2\theta_{14}$ plane was included into the final excluded area if it appeared in the excluded areas for all tested variations of the parameters. The following variations of the parameters were tested:

- The energy resolution ±10% in comparison with the MC predictions;
- A flat background which gives ±0.1% events at the top position of the detector which corresponds to 100 % variation of this background;
- A background with the energy distribution identical to the distribution of the background produced by cosmic muons inside the detector. The fraction of such background was $\pm 0.5\%$ of the IBD rate at the top position of the detector which corresponded to 15 % variation of this background;
- All possible combinations of changes listed above;
- The reduced range of the energies used in the fit to (1.5-6) MeV.

Figure 10 shows the obtained 90% CL excluded area in the Δm_{14}^2 , $\sin^2 2\theta_{14}$ plane. For some values of Δm_{14}^2 the obtained limits are more stringent than previous results[16, 17, 18]. It is important to stress that our results are based only on the comparison of the positron energy distributions at the two distances measured with the same detector. Therefore the results do not depend on the $\tilde{\nu}_e$ spectrum shape and normalization as well as on the detector efficiency. The excluded area covers a large fraction of regions indicated by the GA and RAA. In particular, the most preferred point $\Delta m_{14}^2 = 2.3 \text{ eV}^2$, $\sin^2 2\theta_{14} = 0.14[5]$ is excluded at more than 5 σ CL. In our analysis the point $\Delta m_{14}^2 = 1.4 \text{ eV}^2$, $\sin^2 2\theta_{14} = 0.05$ has the smallest $\chi^2 = 21.9$. The difference in χ^2 with the 3ν case is 13.1 which corresponds to ~ 3σ . The significance of this indication of the existence of the sterile neutrino will be studied taking into account systematic uncertainties after collection of more data this year.

Acknowledgments

The authors are grateful to the directorates of ITEP and JINR for constant support of this work. The authors appreciate the permanent assistance of the KNPP administration and Radiation and Nuclear Safety Departments. The detector construction was supported by the Russian State Corporation ROSATOM (state contracts H.4x.44.90.13.1119 and H.4x.44.9B.16.1006). The operation and data analysis became possible due to the valuable support from the Russian Science Foundation grant 17-12-01145.




Figure 10: 90% CL exclusion area in Δm_{14}^2 , $\sin^2 2\theta_{14}$ parameter space. The shaded area represents our analysis. Curves show allowed regions from neutrino disappearance experiments[5, 19], and the star is the best point from the RAA and GA fit[5].

References

References

- Fogli G. L. et al., Global analysis of neutrino masses, mixings, and phases: Entering the era of leptonic CP violation searches, Phys.Rev. D 86 (2012) 013012. doi:10.1103/PhysRevD.86. 013012.
- [2] Patrignani C. *et al.*, Review of particle physics, Chin. Phys. C 40 (2016) 100001.
- [3] Abdurashitov J. N. et al., Measurement of the response of a Ga solar neutrino experiment to neutrinos from a ³⁷Ar source, Phys.Rev. C 73 (2006) 045805. doi:10.1103/PhysRevC.73. 045805.
- [4] Mueller T. A. et al., Improved predictions of reactor antineutrino spectra, Phys.Rev. C 83 (2011) 054615. doi:10.1103/ PhysRevC.83.054615.
- [5] Mention G. *et al.*, The reactor antineutrino anomaly, Phys.Rev. D 83 (2011) 073006. doi:10.1103/PhysRevD.83.073006.
- [6] Athanassopoulos C. *et al.*, Evidence for $\tilde{\nu}_{\mu} \rightarrow \tilde{\nu}_{e}$ Oscillations from the LSND Experiment at the Los Alamos Meson Physics Facility, Phys.Rev.Lett. **77** (1996) 3082.
- [7] Aguilar-Arevalo A A *et al.*, Improved Search for $\tilde{\nu}_{\mu} \rightarrow \tilde{\nu}_{e}$ Oscillations in the MiniBooNE Experiment, Phys.Rev.Lett. **110** (2006) 161801.
- [8] Ade P. A. R. et al., Planck 2015 results XX. Constraints on inflation, Astron.Astrophys. 20 (2016) 594. doi:10.1051/ 0004-6361/201525898.
- [9] Abazajian K. N. *et al.*, Light sterile neutrinos: A white paper, FERMILAB-PUB-12-881-PPD.
- [10] Alekseev I. et al., DANSS: Detector of the reactor AntiNeutrino based on Solid Scintillator, JINST 11 (11) (2016) P11011. doi: 10.1088/1748-0221/11/11/P11011.
- [11] Alekseev I. et al., Nonuniformity of light collection from scintillator strips with wavelength sifting fibers of the DANSS detector, Instr.Exp.Tech. to be published.
- [12] Alekseev I. et al., Photodetectors signal digitization in the DANSS detector, Instr.Exp.Tech. to be published.
- [13] Alekseev I. *et al.*, The DANSS neutrino spectrometer: calibration of the detector, stability of the response and light yield studies, Lett.Elem.Part.Atom.Nucl. to be published.

- [14] Huber P., Determination of antineutrino spectra from nuclear reactors, Phys.Rev. C 84 (2011) 024617. doi:10.1103/ PhysRevC.84.024617.
- [15] Qian X. *et al.*, The Gaussian CL_s method for searches of new physics, Nucl.Inst.Meth. A **827** (2016) 63. doi:10.1016/j. nima.2016.04.089.
- [16] Achkar A. *et al.*, Search for neutrino oscillations at 15, 40 and 95 meters from a nuclear power reactor at Bugey, Nucl.Phys. B 434 (1995) 503. doi:10.1016/0550-3213(94)00513-E.
- [17] Adamson P. et al., Limits on Active to Sterile Neutrino Oscillations from Disappearance Searches in the MINOS, Daya Bay, and Bugey-3 Experiments, Phys.Rev.Lett. 117 (2016) 151801. doi:10.1103/PhysRevLett.117.151801.
- [18] Ko Y. J. *et al.*, Sterile neutrino search at NEOS Experiment, Phys.Rev.Lett. **118** (2017) 121802. doi:10.1103/PhysRevLett. 118.121802.
- [19] J. Kopp, P. A. N. Machado, M. Maltoni, and T. Schwetz, Sterile neutrino oscillations: the global picture, JHEP 1305 (2013) 050. doi:10.1007/JHEP05(2013)050.