

Beam-energy and centrality dependence of direct-photon emission from ultra-relativistic heavy-ion collisions

- A. Adare,¹⁴ S. Afanasiev,³⁴ C. Aidala,^{15, 44, 49, 50} N.N. Ajitanand,^{70, *} Y. Akiba,^{64, 65, †} R. Akimoto,¹³ H. Al-Bataineh,⁵⁸ J. Alexander,⁷⁰ M. Alfred,²⁷ A. Al-Jamel,⁵⁸ H. Al-Ta’ani,⁵⁸ A. Angerami,¹⁵ K. Aoki,^{37, 40, 64} N. Apadula,^{32, 71} L. Aphecetche,⁷² Y. Aramaki,^{13, 64} R. Armendariz,⁵⁸ S.H. Aronson,⁸ J. Asai,^{64, 65} H. Asano,^{40, 64} E.C. Aschenauer,⁸ E.T. Atomssa,^{41, 71} R. Averbeck,⁷¹ T.C. Awes,⁶⁰ B. Azmoun,⁸ V. Babintsev,²⁸ A. Bagoly,¹⁹ M. Bai,⁷ G. Baksay,²² L. Baksay,²² A. Baldissari,¹⁷ B. Bannier,⁷¹ K.N. Barish,⁹ P.D. Barnes,^{44, *} B. Bassalleck,⁵⁷ A.T. Basye,¹ S. Bathe,^{6, 9, 65} S. Batsouli,^{15, 60} V. Baublis,⁶³ F. Bauer,⁹ C. Baumann,⁵¹ S. Baumgart,⁶⁴ A. Bazilevsky,⁸ S. Belikov,^{8, 32, *} R. Belmont,^{14, 77} R. Bennett,⁷¹ A. Berdnikov,⁶⁷ Y. Berdnikov,⁶⁷ J.H. Bhom,⁸¹ A.A. Bickley,¹⁴ M.T. Bjorndal,¹⁵ D.S. Blau,^{39, 56} M. Boer,⁴⁴ J.G. Bosissevain,⁴⁴ J.S. Bok,^{57, 58, 81} H. Borel,¹⁷ K. Boyle,^{65, 71} M.L. Brooks,⁴⁴ D.S. Brown,⁵⁸ J. Bryslawskyj,^{6, 9} J. Bryslawskyj,⁹ D. Bucher,⁵¹ H. Buesching,⁸ V. Bumazhnov,²⁸ G. Bunce,^{8, 65} J.M. Burward-Hoy,⁴⁴ S. Butsyk,^{44, 57, 71} C.M. Camacho,⁴⁴ S. Campbell,^{15, 71} V. Canoa Roman,⁷¹ A. Caringi,⁵² P. Castera,⁷¹ J.-S. Chai,^{36, 73} B.S. Chang,⁸¹ W.C. Chang,² J.-L. Charvet,¹⁷ C.-H. Chen,^{65, 71} S. Chernichenko,²⁸ C.Y. Chi,¹⁵ J. Chiba,³⁷ M. Chiu,^{8, 15, 29} I.J. Choi,^{29, 81} J.B. Choi,^{11, *} S. Choi,⁶⁹ R.K. Choudhury,⁵ P. Christiansen,⁴⁶ T. Chujo,^{76, 77} P. Chung,⁷⁰ A. Churyn,²⁸ O. Chvala,⁹ V. Cianciolo,⁶⁰ Z. Citron,^{71, 79} C.R. Cleven,²⁴ Y. Cobigo,¹⁷ B.A. Cole,¹⁵ M.P. Comets,⁶¹ Z. Conesa del Valle,⁴¹ M. Connors,^{24, 65, 71} P. Constantin,^{32, 44} M. Csanad,¹⁹ T. Csorgo,^{20, 80} T. Dahms,⁷¹ S. Dairaku,^{40, 64} I. Danchev,⁷⁷ T.W. Danley,⁵⁹ K. Das,²³ A. Datta,⁴⁹ M.S. Daugherty,¹ G. David,^{8, 71} M.K. Dayananda,²⁴ M.B. Deaton,¹ K. Dehmelt,^{22, 71} H. Delagrange,^{72, *} A. Denisov,²⁸ D. d’Enterria,^{15, 41} A. Deshpande,^{65, 71} E.J. Desmond,⁸ K.V. Dharmawardane,⁵⁸ O. Dietzsch,⁶⁸ L. Ding,^{32, 71} J.H. Do,⁸¹ M. Donadelli,⁶⁸ L. D’Orazio,⁴⁸ J.L. Drachenberg,¹ O. Drapier,⁴¹ A. Drees,⁷¹ K.A. Drees,⁷ A.K. Dubey,⁷⁹ J.M. Durham,^{44, 71} A. Durum,²⁸ D. Dutta,⁵ V. Dzhordzhadze,^{9, 74} S. Edwards,^{7, 23} Y.V. Efremenko,⁶⁰ J. Egdemir,⁷¹ F. Ellinghaus,¹⁴ W.S. Emam,⁹ T. Engelmore,¹⁵ A. Enokizono,^{26, 43, 60, 64, 66} H. En’yo,^{64, 65} B. Espagnon,⁶¹ S. Esumi,⁷⁶ K.O. Eyser,^{8, 9} B. Fadem,⁵² W. Fan,⁷¹ N. Feege,⁷¹ D.E. Fields,^{57, 65} M. Finger,^{10, 34} M. Finger, Jr.,^{10, 34} F. Fleuret,⁴¹ S.L. Fokin,³⁹ B. Forestier,⁴⁵ Z. Fraenkel,^{79, *} J.E. Frantz,^{15, 59, 71} A. Franz,⁸ A.D. Frawley,²³ K. Fujiwara,⁶⁴ Y. Fukao,^{40, 64} S.-Y. Fung,⁹ T. Fusayasu,⁵⁴ S. Gadrat,⁴⁵ K. Gainey,¹ C. Gal,⁷¹ P. Gallus,¹⁶ P. Garg,^{4, 71} A. Garishvili,⁷⁴ I. Garishvili,^{43, 74} F. Gastineau,⁷² H. Ge,⁷¹ M. Germain,⁷² A. Glenn,^{14, 43, 74} H. Gong,⁷¹ X. Gong,⁷⁰ M. Gonin,⁴¹ J. Gosset,¹⁷ Y. Goto,^{64, 65} R. Granier de Cassagnac,⁴¹ N. Grau,^{3, 15, 32} S.V. Greene,⁷⁷ G. Grim,⁴⁴ M. Grosse Perdekamp,^{29, 65} T. Gunji,¹³ L. Guo,⁴⁴ H.-Å. Gustafsson,^{46, *} T. Hachiya,^{26, 64, 65} A. Hadj Henni,⁷² C. Haegemann,⁵⁷ J.S. Haggerty,⁸ M.N. Hagiwara,¹ K.I. Hahn,²¹ H. Hamagaki,¹³ J. Hamblen,⁷⁴ R. Han,⁶² J. Hanks,^{15, 71} H. Harada,²⁶ E.P. Hartouni,⁴³ K. Haruna,²⁶ M. Harvey,⁸ S. Hasegawa,³³ T.O.S. Haseler,²⁴ K. Hashimoto,^{64, 66} E. Haslum,⁴⁶ K. Hasuko,⁶⁴ R. Hayano,¹³ X. He,²⁴ M. Heffner,⁴³ T.K. Hemmick,⁷¹ T. Hester,⁹ J.M. Heuser,⁶⁴ H. Hiejima,²⁹ J.C. Hill,³² K. Hill,¹⁴ R. Hobbs,⁵⁷ A. Hodges,²⁴ M. Hohlmann,²² R.S. Hollis,⁹ M. Holmes,⁷⁷ W. Holzmann,^{15, 70} K. Homma,²⁶ B. Hong,³⁸ T. Horaguchi,^{13, 26, 64, 75, 76} Y. Hori,¹³ D. Hornback,⁷⁴ N. Hotvedt,³² J. Huang,⁸ S. Huang,⁷⁷ M.G. Hur,³⁶ T. Ichihara,^{64, 65} R. Ichimiya,⁶⁴ H. Iinuma,^{37, 40, 64} Y. Ikeda,^{64, 76} K. Imai,^{33, 40, 64} J. Imrek,¹⁸ M. Inaba,⁷⁶ Y. Inoue,^{64, 66} A. Iordanova,⁹ D. Isenhower,¹ L. Isenhower,¹ M. Ishihara,⁶⁴ T. Isobe,^{13, 64} M. Issah,^{70, 77} A. Isupov,³⁴ D. Ivanishchev,⁶³ Y. Iwanaga,²⁶ B.V. Jacak,⁷¹ M. Javani,²⁴ Z. Ji,⁷¹ J. Jia,^{8, 15, 70} X. Jiang,⁴⁴ J. Jin,¹⁵ O. Jinnouchi,⁶⁵ B.M. Johnson,^{8, 24} T. Jones,¹ K.S. Joo,⁵³ D. Jouan,⁶¹ D.S. Jumper,^{1, 29} F. Kajihara,^{13, 64} S. Kametani,^{13, 64, 78} N. Kamihara,^{64, 65, 75} J. Kamin,⁷¹ M. Kaneta,⁶⁵ S. Kaneti,⁷¹ B.H. Kang,²⁵ J.H. Kang,⁸¹ J.S. Kang,²⁵ H. Kanou,^{64, 75} J. Kapustinsky,⁴⁴ K. Karatsu,^{40, 64} M. Kasai,^{64, 66} T. Kawagishi,⁷⁶ D. Kawall,^{49, 65} M. Kawashima,^{64, 66} A.V. Kazantsev,³⁹ S. Kelly,¹⁴ T. Kempel,³² V. Khachatryan,⁷¹ A. Khanzadeev,⁶³ K.M. Kijima,²⁶ J. Kikuchi,⁷⁸ A. Kim,²¹ B.I. Kim,³⁸ C. Kim,³⁸ D.H. Kim,⁵³ D.J. Kim,^{35, 81} E. Kim,⁶⁹ E.-J. Kim,¹¹ H.J. Kim,⁸¹ K.-B. Kim,¹¹ M. Kim,⁶⁹ S.H. Kim,⁸¹ Y.-J. Kim,²⁹ Y.K. Kim,²⁵ Y.-S. Kim,³⁶ D. Kincses,¹⁹ E. Kinney,¹⁴ K. Kiriluk,¹⁴ Á. Kiss,¹⁹ E. Kistenev,⁸ A. Kiyomichi,⁶⁴ J. Klatsky,²³ J. Klay,⁴³ C. Klein-Boesing,⁵¹ D. Kleinjan,⁹ P. Kline,⁷¹ L. Kochenda,⁶³ V. Kochetkov,²⁸ Y. Komatsu,^{13, 37} B. Komkov,⁶³ M. Konno,⁷⁶ J. Koster,²⁹ D. Kotchetkov,^{9, 59} D. Kotov,^{63, 67} A. Kozlov,⁷⁹ A. Kral,¹⁶ A. Kravitz,¹⁵ F. Krizek,³⁵ P.J. Kroon,⁸ J. Kubart,^{10, 31} G.J. Kunde,⁴⁴ N. Kurihara,¹³ K. Kurita,^{64, 66} M. Kurosawa,^{64, 65} M.J. Kweon,³⁸ Y. Kwon,^{74, 81} G.S. Kyle,⁵⁸ R. Lacey,⁷⁰ Y.S. Lai,¹⁵ J.G. Lajoie,³² D. Layton,²⁹ A. Lebedev,³² Y. Le Bornec,⁶¹ S. Leckey,⁷¹ B. Lee,²⁵ D.M. Lee,⁴⁴ J. Lee,^{21, 73} K.B. Lee,³⁸ K.S. Lee,³⁸ M.K. Lee,⁸¹ S.H. Lee,^{32, 71} S.R. Lee,¹¹ T. Lee,⁶⁹ M.J. Leitch,⁴⁴ M.A.L. Leite,⁶⁸ M. Leitgab,²⁹ B. Lenzi,⁶⁸ Y.H. Leung,⁷¹ B. Lewis,⁷¹ N.A. Lewis,⁵⁰

X. Li,¹² X. Li,⁴⁴ X.H. Li,⁹ P. Lichtenwalner,⁵² P. Liebing,⁶⁵ H. Lim,⁶⁹ S.H. Lim,^{44,81} L.A. Linden Levy,^{14,29}
 A. Litvinenko,³⁴ H. Liu,^{44,58} M.X. Liu,⁴⁴ T. Liška,¹⁶ S. Lökö^s,^{19,20} B. Love,⁷⁷ D. Lynch,⁸ C.F. Maguire,⁷⁷
 T. Majoros,¹⁸ Y.I. Makdisi,^{7,8} M. Makek,^{79,82} A. Malakhov,³⁴ M.D. Malik,⁵⁷ A. Manion,⁷¹ V.I. Manko,³⁹
 E. Mannel,^{8,15} Y. Mao,^{62,64} H. Masui,⁷⁶ S. Masumoto,^{13,37} F. Matathias,^{15,71} L. Mašek,^{10,31} M.C. McCain,²⁹
 M. McCumber,^{14,44,71} P.L. McGaughey,⁴⁴ D. McGlinchey,^{14,23,44} C. McKinney,²⁹ N. Means,⁷¹ M. Mendoza,⁹
 B. Meredith,²⁹ Y. Miake,⁷⁶ T. Mibe,³⁷ A.C. Mignerey,⁴⁸ D.E. Mihalik,⁷¹ P. Mikeš,^{10,31} K. Miki,^{64,76} T.E. Miller,⁷⁷
 A. Milov,^{8,71,79} S. Mioduszewski,⁸ D.K. Mishra,⁵ G.C. Mishra,²⁴ M. Mishra,⁴ J.T. Mitchell,⁸ M. Mitrovski,⁷⁰
 G. Mitsuka,⁶⁵ Y. Miyachi,^{64,75} S. Miyasaka,^{64,75} A.K. Mohanty,⁵ S. Mohapatra,⁷⁰ H.J. Moon,⁵³ T. Moon,⁸¹
 Y. Morino,¹³ A. Morreale,⁹ D.P. Morrison,⁸ S.I. Morrow,⁷⁷ J.M. Moss,⁴⁴ S. Motschwiller,⁵² T.V. Moukhanova,³⁹
 D. Mukhopadhyay,⁷⁷ T. Murakami,^{40,64} J. Murata,^{64,66} A. Mwai,⁷⁰ T. Nagae,⁴⁰ S. Nagamiya,^{37,64} K. Nagashima,²⁶
 Y. Nagata,⁷⁶ J.L. Nagle,¹⁴ M. Naglis,⁷⁹ M.I. Nagy,^{19,80} I. Nakagawa,^{64,65} Y. Nakamiya,²⁶ K.R. Nakamura,^{40,64}
 T. Nakamura,^{26,64} K. Nakano,^{64,75} S. Nam,²¹ C. Nattrass,⁷⁴ A. Nederlof,⁵² J. Newby,⁴³ M. Nguyen,⁷¹
 M. Nihashi,^{26,64} T. Niida,⁷⁶ B.E. Norman,⁴⁴ R. Nouicer,^{8,65} T. Novák,²⁰ N. Novitzky,^{35,71} A.S. Nyanin,³⁹
 J. Nystrand,⁴⁶ C. Oakley,²⁴ E. O'Brien,⁸ S.X. Oda,¹³ C.A. Ogilvie,³² H. Ohnishi,⁶⁴ I.D. Ojha,⁷⁷ M. Oka,⁷⁶
 K. Okada,⁶⁵ O.O. Omiwade,¹ Y. Onuki,⁶⁴ J.D. Orjuela Koop,¹⁴ J.D. Osborn,⁵⁰ A. Oskarsson,⁴⁶ I. Otterlund,⁴⁶
 M. Ouchida,^{26,64} K. Ozawa,^{13,37,76} R. Pak,⁸ D. Pal,⁷⁷ A.P.T. Palounek,⁴⁴ V. Pantuev,^{30,71} V. Papavassiliou,⁵⁸
 B.H. Park,²⁵ I.H. Park,^{21,73} J. Park,⁶⁹ S. Park,^{64,69,71} S.K. Park,³⁸ W.J. Park,³⁸ S.F. Pate,⁵⁸ L. Patel,²⁴ M. Patel,³²
 H. Pei,³² J.-C. Peng,²⁹ W. Peng,⁷⁷ H. Pereira,¹⁷ D.V. Perepelitsa,^{14,15} V. Peresedov,³⁴ D.Yu. Peressounko,³⁹
 C.E. PerezLara,⁷¹ R. Petti,^{8,71} C. Pinkenburg,⁸ R.P. Pisani,⁸ M. Proissl,⁷¹ M.L. Purschke,⁸ A.K. Purwar,^{44,71}
 H. Qu,^{1,24} P.V. Radzevich,⁶⁷ J. Rak,^{32,35,57} A. Rakotozafindrabe,⁴¹ I. Ravinovich,⁷⁹ K.F. Read,^{60,74}
 S. Rembeczki,²² M. Reuter,⁷¹ K. Reygers,⁵¹ D. Reynolds,⁷⁰ V. Riabov,^{56,63} Y. Riabov,^{63,67} E. Richardson,⁴⁸
 D. Richford,⁶ T. Rinn,³² D. Roach,⁷⁷ G. Roche,^{45,*} S.D. Rolnick,⁹ A. Romana,^{41,*} M. Rosati,³² C.A. Rosen,¹⁴
 S.S.E. Rosendahl,⁴⁶ P. Rosnet,⁴⁵ Z. Rowan,⁶ P. Rukoyatkin,³⁴ J. Runchey,³² P. Ružička,³¹ V.L. Rykov,⁶⁴ S.S. Ryu,⁸¹
 B. Sahlueller,^{51,71} N. Saito,^{37,40,64,65} T. Sakaguchi,^{8,13,78} S. Sakai,⁷⁶ K. Sakashita,^{64,75} H. Sakata,²⁶ H. Sako,³³
 V. Samsonov,^{56,63} M. Sano,⁷⁶ S. Sano,^{13,78} M. Sarsour,²⁴ H.D. Sato,^{40,64} S. Sato,^{8,33,37,76} T. Sato,⁷⁶ S. Sawada,³⁷
 B.K. Schmoll,⁷⁴ K. Sedgwick,⁹ J. Seele,¹⁴ R. Seidl,^{29,64,65} A.Yu. Semenov,³² V. Semenov,^{28,30} A. Sen,^{24,32} R. Seto,⁹
 D. Sharma,^{71,79} T.K. Shea,⁸ I. Shein,²⁸ A. Shevel,^{63,70} T.-A. Shibata,^{64,75} K. Shigaki,²⁶ M. Shimomura,^{32,55,76}
 T. Shohjoh,⁷⁶ K. Shoji,^{40,64} P. Shukla,⁵ A. Sickles,^{8,29,71} C.L. Silva,^{32,44,68} D. Silvermyr,^{46,60} C. Silvestre,¹⁷
 K.S. Sim,³⁸ B.K. Singh,⁴ C.P. Singh,⁴ V. Singh,⁴ M.J. Skoby,⁵⁰ S. Skutnik,³² M. Slunečka,^{10,34} W.C. Smith,¹
 A. Soldatov,²⁸ R.A. Soltz,⁴³ W.E. Sondheim,⁴⁴ S.P. Sorensen,⁷⁴ I.V. Sourikova,⁸ F. Staley,¹⁷ P.W. Stankus,⁶⁰
 E. Stenlund,⁴⁶ M. Stepanov,^{49,58,*} A. Ster,⁸⁰ S.P. Stoll,⁸ T. Sugitate,²⁶ C. Suire,⁶¹ A. Sukhanov,⁸ J.P. Sullivan,⁴⁴
 J. Sun,⁷¹ Z Sun,¹⁸ Z. Sun,¹⁸ J. Sziklai,⁸⁰ T. Tabaru,⁶⁵ S. Takagi,⁷⁶ E.M. Takagui,⁶⁸ A. Takahara,¹³ A. Taketani,^{64,65}
 R. Tanabe,⁷⁶ K.H. Tanaka,³⁷ Y. Tanaka,⁵⁴ S. Taneja,⁷¹ K. Tanida,^{33,40,64,65,69} M.J. Tannenbaum,⁸ S. Tarafdar,^{4,77}
 A. Taranenko,^{56,70} P. Tarján,¹⁸ E. Tennant,⁵⁸ H. Themann,⁷¹ D. Thomas,¹ T.L. Thomas,⁵⁷ R. Tieulent,⁴⁷
 T. Todoroki,^{64,76} M. Togawa,^{40,64,65} A. Toia,⁷¹ J. Tojo,⁶⁴ L. Tomášek,³¹ M. Tomášek,^{16,31} Y. Tomita,⁷⁶
 H. Torii,^{26,64} R.S. Towell,¹ V-N. Tram,⁴¹ I. Tserruya,⁷⁹ Y. Tsuchimoto,^{13,26,64} T. Tsuji,¹³ S.K. Tuli,^{4,*}
 H. Tydesjö,⁴⁶ N. Tyurin,²⁸ B. Ujvari,¹⁸ C. Vale,^{8,32} H. Valle,⁷⁷ H.W. van Hecke,⁴⁴ M. Vargyas,^{19,80}
 E. Vazquez-Zambrano,¹⁵ A. Veicht,^{15,29} J. Velkovska,⁷⁷ R. Vértesi,^{18,80} A.A. Vinogradov,³⁹ M. Virius,¹⁶
 A. Vossen,²⁹ V. Vrba,^{16,31} E. Vznuzdaev,⁶³ M. Wagner,^{40,64} D. Walker,⁷¹ X.R. Wang,^{58,65} D. Watanabe,²⁶
 K. Watanabe,⁷⁶ Y. Watanabe,^{64,65} Y.S. Watanabe,¹³ F. Wei,^{32,58} R. Wei,⁷⁰ J. Wessels,⁵¹ S.N. White,⁸ N. Willis,⁶¹
 D. Winter,¹⁵ S. Wolin,²⁹ C.P. Wong,²⁴ C.L. Woody,⁸ R.M. Wright,¹ M. Wysocki,^{14,60} B. Xia,⁵⁹ W. Xie,^{9,65}
 C. Xu,⁵⁸ Q. Xu,⁷⁷ Y.L. Yamaguchi,^{13,64,65,71,78} K. Yamaura,²⁶ R. Yang,²⁹ A. Yanovich,²⁸ Z. Yasin,⁹ J. Ying,²⁴
 S. Yokkaichi,^{64,65} J.H. Yoo,³⁸ Z. You,^{44,62} G.R. Young,⁶⁰ I. Younus,^{42,57} H. Yu,⁵⁸ I.E. Yushmanov,³⁹ W.A. Zajc,¹⁵
 O. Zaudtke,⁵¹ A. Zelenski,⁷ C. Zhang,^{15,60} S. Zharko,⁶⁷ S. Zhou,¹² J. Zimamyi,^{80,*} L. Zolin,³⁴ and L. Zou⁹

(PHENIX Collaboration)

¹Abilene Christian University, Abilene, Texas 79699, USA

²Institute of Physics, Academia Sinica, Taipei 11529, Taiwan

³Department of Physics, Augustana University, Sioux Falls, South Dakota 57197, USA

⁴Department of Physics, Banaras Hindu University, Varanasi 221005, India

⁵Bhabha Atomic Research Centre, Bombay 400 085, India

⁶Baruch College, City University of New York, New York, New York, 10010 USA

⁷Collider-Accelerator Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

⁸Physics Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

- ⁹ University of California-Riverside, Riverside, California 92521, USA
¹⁰ Charles University, Ovocný trh 5, Praha 1, 116 36, Prague, Czech Republic
¹¹ Chonbuk National University, Jeonju, 561-756, Korea
¹² Science and Technology on Nuclear Data Laboratory, China Institute of Atomic Energy, Beijing 102413, People's Republic of China
¹³ Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan
¹⁴ University of Colorado, Boulder, Colorado 80309, USA
¹⁵ Columbia University, New York, New York 10027 and Nevis Laboratories, Irvington, New York 10533, USA
¹⁶ Czech Technical University, Zikova 4, 166 36 Prague 6, Czech Republic
¹⁷ Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France
¹⁸ Debrecen University, H-4010 Debrecen, Egyetem tér 1, Hungary
¹⁹ ELTE, Eötvös Loránd University, H-1117 Budapest, Pázmány P. s. 1/A, Hungary
²⁰ Eszterházy Károly University, Károly Róbert Campus, H-3200 Gyöngyös, Mátrai út 36, Hungary
²¹ Ewha Womans University, Seoul 120-750, Korea
²² Florida Institute of Technology, Melbourne, Florida 32901, USA
²³ Florida State University, Tallahassee, Florida 32306, USA
²⁴ Georgia State University, Atlanta, Georgia 30303, USA
²⁵ Hanyang University, Seoul 133-792, Korea
²⁶ Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan
²⁷ Department of Physics and Astronomy, Howard University, Washington, DC 20059, USA
²⁸ IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia
²⁹ University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA
³⁰ Institute for Nuclear Research of the Russian Academy of Sciences, prospekt 60-letiya Oktyabrya 7a, Moscow 117312, Russia
³¹ Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague 8, Czech Republic
³² Iowa State University, Ames, Iowa 50011, USA
³³ Advanced Science Research Center, Japan Atomic Energy Agency, 2-4 Shirakata Shirane, Tokai-mura, Naka-gun, Ibaraki-ken 319-1195, Japan
³⁴ Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia
³⁵ Helsinki Institute of Physics and University of Jyväskylä, P.O.Box 35, FI-40014 Jyväskylä, Finland
³⁶ KAERI, Cyclotron Application Laboratory, Seoul, Korea
³⁷ KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan
³⁸ Korea University, Seoul, 136-701, Korea
³⁹ National Research Center "Kurchatov Institute", Moscow, 123098 Russia
⁴⁰ Kyoto University, Kyoto 606-8502, Japan
⁴¹ Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France
⁴² Physics Department, Lahore University of Management Sciences, Lahore 54792, Pakistan
⁴³ Lawrence Livermore National Laboratory, Livermore, California 94550, USA
⁴⁴ Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA
⁴⁵ LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubière Cedex, France
⁴⁶ Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden
⁴⁷ IPNL, CNRS/IN2P3, Univ Lyon, Université Lyon 1, F-69622, Villeurbanne, France
⁴⁸ University of Maryland, College Park, Maryland 20742, USA
⁴⁹ Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003-9337, USA
⁵⁰ Department of Physics, University of Michigan, Ann Arbor, Michigan 48109-1040, USA
⁵¹ Institut für Kernphysik, University of Muenster, D-48149 Muenster, Germany
⁵² Muhlenberg College, Allentown, Pennsylvania 18104-5586, USA
⁵³ Myongji University, Yongin, Kyonggido 449-728, Korea
⁵⁴ Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan
⁵⁵ Nara Women's University, Kita-uoya Nishi-machi Nara 630-8506, Japan
⁵⁶ National Research Nuclear University, MEPhI, Moscow Engineering Physics Institute, Moscow, 115409, Russia
⁵⁷ University of New Mexico, Albuquerque, New Mexico 87131, USA
⁵⁸ New Mexico State University, Las Cruces, New Mexico 88003, USA
⁵⁹ Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701, USA
⁶⁰ Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
⁶¹ IPN-Orsay, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, BP1, F-91406, Orsay, France
⁶² Peking University, Beijing 100871, People's Republic of China
⁶³ PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia
⁶⁴ RIKEN Nishina Center for Accelerator-Based Science, Wako, Saitama 351-0198, Japan
⁶⁵ RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973-5000, USA
⁶⁶ Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan
⁶⁷ Saint Petersburg State Polytechnic University, St. Petersburg, 195251 Russia
⁶⁸ Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil
⁶⁹ Department of Physics and Astronomy, Seoul National University, Seoul 151-742, Korea
⁷⁰ Chemistry Department, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA

⁷¹Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York 11794-3800, USA

⁷²SUBATECH (Ecole des Mines de Nantes, CNRS-IN2P3, Université de Nantes) BP 20722-44307, Nantes, France

⁷³Sungkyunkwan University, Suwon, 440-746, Korea

⁷⁴University of Tennessee, Knoxville, Tennessee 37996, USA

⁷⁵Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan

⁷⁶Tomonaga Center for the History of the Universe, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

⁷⁷Vanderbilt University, Nashville, Tennessee 37235, USA

⁷⁸Waseda University, Advanced Research Institute for Science and Engineering, 17 Kikui-cho, Shinjuku-ku, Tokyo 162-0044, Japan

⁷⁹Weizmann Institute, Rehovot 76100, Israel

⁸⁰Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Hungarian Academy of Sciences (Wigner RCP, RMKI) H-1525 Budapest 114, POBox 49, Budapest, Hungary

⁸¹Yonsei University, IPAP, Seoul 120-749, Korea

⁸²Department of Physics, Faculty of Science, University of Zagreb, Bijenička c. 32 HR-10002 Zagreb, Croatia

(Dated: November 15, 2021)

The PHENIX collaboration presents first measurements of low-momentum ($0.4 < p_T < 3 \text{ GeV}/c$) direct-photon yields from Au+Au collisions at $\sqrt{s_{NN}}=39$ and 62.4 GeV. For both beam energies the direct-photon yields are substantially enhanced with respect to expectations from prompt processes, similar to the yields observed in Au+Au collisions at $\sqrt{s_{NN}}=200$. Analyzing the photon yield as a function of the experimental observable $dN_{\text{ch}}/d\eta$ reveals that the low-momentum ($>1 \text{ GeV}/c$) direct-photon yield $dN_{\gamma}^{\text{dir}}/d\eta$ is a smooth function of $dN_{\text{ch}}/d\eta$ and can be well described as proportional to $(dN_{\text{ch}}/d\eta)^{\alpha}$ with $\alpha \sim 1.25$. This new scaling behavior holds for a wide range of beam energies at the Relativistic Heavy Ion Collider and Large Hadron Collider, for centrality selected samples, as well as for different, $A+A$ collision systems. At a given beam energy the scaling also holds for high p_T ($> 5 \text{ GeV}/c$) but when results from different collision energies are compared, an additional $\sqrt{s_{NN}}$ -dependent multiplicative factor is needed to describe the integrated-direct-photon yield.

The measurement of direct photons is a unique tool to study the strongly coupled quark-gluon plasma (QGP) produced in heavy ion collisions and its “fireball” evolution to hadron resonance matter. Direct photons leave the interaction region unmodified, and provide information from different stages of the collision integrated over space and time [1–3]. Of particular interest are low momentum direct photons, which are thought to be radiated mostly by the QGP and the hot hadron gas.

In experiments direct photons are detected simultaneously with a large number of photons from final state hadron decays, mostly from π^0 and η mesons. The main experimental challenge is to subtract these decay contributions from the measurement to obtain the photons directly emitted from the collision. In addition to photons from the hot fireball, direct photons are also emitted from initial hard scattering processes, such as quark-gluon Compton scattering among the incoming partons. Disentangling this prompt component from the photons emitted from the fireball is the second experimental challenge.

Although a very promising observable, only recently PHENIX discovered first evidence of low p_T direct photons in central $A+A$ collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$ [4]. Since then PHENIX has published centrality dependent yields in Au+Au [5] as well as azimuthal asymmetries of direct photon emission [6, 7]. Results on yields are also available from STAR [8] for the same collision system and from ALICE [9, 10] for Pb+Pb collisions at $\sqrt{s_{NN}} = 2760 \text{ GeV}$.

A consistent picture emerges based on those data: a

large number of low p_T direct photons are radiated from the fireball, the yield has a strong centrality dependence and a significant azimuthal anisotropy expressed by the coefficients v_2 and v_3 of a Fourier expansion of the direct photon yield with respect to the reaction plane.

A large body of theoretical work on direct photon emission exists in the literature. Many model calculations are qualitatively consistent with the data, but a quantitative description remains difficult, primarily due to the simultaneous presence of large yields and azimuthal anisotropies [11–34].

In this publication we present low- p_T direct photon data from Au+Au collisions at $\sqrt{s_{NN}} = 39 \text{ GeV}$ and 62.4 GeV taken with the PHENIX experiment in 2010. We compare the spectra and the integrated yields to those from $p+p$ and Au+Au collisions at various centralities at $\sqrt{s_{NN}} = 200 \text{ GeV}$ [4, 5], and to those from Pb+Pb collisions at $\sqrt{s_{NN}} = 2760 \text{ GeV}$ [9]. This covers a factor of 70 in $\sqrt{s_{NN}}$ for colliding systems of comparable size.

The 39 and 62.4 GeV spectra are obtained from two data samples of minimum bias Au+Au collisions that have a total of 7.79×10^7 and 2.12×10^8 events, respectively. The minimum bias trigger and centrality selection is derived from data taken with the PHENIX beam-beam counters [35]. The data analysis uses the same techniques deployed for the analysis of the $\sqrt{s_{NN}} = 200 \text{ GeV}$ Au+Au data [5], which were taken in the same year under nearly identical conditions. Here we give a brief overview of the setup and data analysis, and refer to our previous publication for more details [5].

Photons are reconstructed through their conversion to




FIG. 1. Direct photon spectra normalized by $(dN_{ch}/d\eta)^{1.25}$ for Au+Au at 39 and 64.2 GeV (a) and (b) at 200 GeV [6]; panel (c) compares for different $A+A$ systems at different $\sqrt{s_{NN}}$ [9, 36]. Panels (a) and (b) also show $p+p$ data [5, 37–39]. All panels show pQCD calculations for the corresponding \sqrt{s} [16, 40]. All error bars are quadratic sum systematic and statistical uncertainties. Uncertainties on the $dN_{ch}/d\eta$ are not included.

e^+e^- pairs in the detector material, specifically the readout boards of the Hadron Blind Detector (HBD) [41] that are located at a radius of 60 cm from the collision point. The trajectories and momenta of the e^+ and e^- are determined by the central arm tracking detectors [42]. Each of the two central arms covers 90° in azimuth and a rapidity range of $|\eta| < 0.35$. A transverse momentum cut, $p_T > 200$ MeV/c, is applied to each trajectory. To identify trajectories as e^+ or e^- candidates, we require a minimum of three associated signals in the ring-imaging Čerenkov detector [43] and that the energy measured in the Electromagnetic Calorimeter (EMCal) [44] matches the measured momentum ($E/p > 0.5$).

All e^+ and e^- reconstructed in the same arm are matched to pairs. In the 2010 setup there is no tracking near the collision point, so the origin of the tracks is assumed. For each e^+e^- pair the mass is calculated twice: first assuming the pair originated at the event vertex (m_{vtx}), then assuming the e^+e^- is a conversion pair from the HBD readout boards (m_{HBD}). In the latter case, m_{HBD} will be consistent with zero, within a mass resolution of a few MeV/c², while m_{vtx} will be about 12 MeV/c². With a cut on both masses a sample of photon conversion is selected with a purity of about 99%. The combinatorial background is negligible, because the

conversion material, in radiation length $X/X_0 \sim 3\%$, is about 10 times thicker than materials closer to the vertex; and it is at a relatively large distance from the event vertex. The 1% contamination is mostly from π^0 Dalitz decays, $\pi^0 \rightarrow \gamma e^+e^-$, and from conversions in front of the HBD readout boards.

The direct photon content in the inclusive photon sample is determined by the ratio R_γ , which is the ratio of all emitted photons (γ^{incl}) to those from hadron decays (γ^{hadron}). R_γ is determined using a double ratio method:

$$R_\gamma = \frac{\gamma^{\text{incl}}}{\gamma^{\text{hadron}}} = \frac{\langle \varepsilon_\gamma f \rangle \left(N_\gamma^{\text{incl}} / N_\gamma^{\pi^0, \text{tag}} \right)_{\text{Data}}}{\left(\gamma^{\text{hadron}} / \gamma^{\pi^0} \right)_{\text{Sim}}}. \quad (1)$$

All quantities in this equation are functions of the conversion photon p_T^{ee} . The measured quantities are the number of detected conversion photons N_γ^{incl} and the subset of those that are tagged as π^0 decay photon $N_\gamma^{\pi^0, \text{tag}}$. The tagged photons $N_\gamma^{\pi^0, \text{tag}}$ are determined statistically in bins of the p_T^{ee} . Each conversion photon is paired with all showers with $E > 400$ MeV measured in the EMCal of the same arm. The invariant $e^+e^-\gamma$ mass is calculated and the counts above the combinatorial background in the π^0 mass peak give $N_\gamma^{\pi^0, \text{tag}}$. To convert

the ratio $N_\gamma^{incl}/N_\gamma^{\pi^0,tag}$ to $\gamma^{incl}/\gamma^{\pi^0}$ only $N_\gamma^{\pi^0,tag}$ needs to be corrected for the momentum averaged conditional acceptance-efficiency $\langle \varepsilon_\gamma f \rangle$ for the second decay photon to be reconstructed in the EMCAL. All other corrections to the numerator and denominator cancel [5]. Because rather loose cuts are applied to the EMCAL showers, $\langle \varepsilon_\gamma f \rangle$ is mostly determined by the π^0 decay kinematics and the detector geometry. Thus, $\langle \varepsilon_\gamma f \rangle$ can be calculated to a few percent accuracy using a Monte-Carlo simulation of π^0 decays in the acceptance. Photons from pions are determined from the measured π^0 spectra [45] and the decay kinematics. The spectrum of decay photons (γ^{hadron}) is derived from γ^{π^0} and the η/π^0 ratio [46], which is independent of collision system and energy, with additional contribution from heavier mesons of about 4%.

Once R_γ is established the direct photon spectrum can be calculated as:

$$\gamma^{\text{direct}} = (R_\gamma - 1) \gamma^{\text{hadron}}. \quad (2)$$

The uncertainty on γ^{hadron} , approximately 10% [5], cancels in R_γ (with that of γ^{π^0} in (1)) but has to be applied to γ^{direct} . The systematic uncertainties of the new data at 39 and 62.4 GeV are similar in magnitude to those for 200 GeV presented in [5]. For the integrated yield we treat every systematic uncertainty as p_T -correlated in the interest of consistency throughout the different data sets.

Figure 1 shows the invariant yield of direct photons normalized to $(dN_{ch}/d\eta)^{1.25}$, this normalization is discussed below. Panel (a) shows Au+Au minimum bias data at $\sqrt{s_{NN}} = 62.4$ and 39 GeV, panel (b) gives Au+Au data in three centrality classes at 200 GeV, and panel (c) compares data from different beam energies and systems. Below 3 GeV/c the 62.4 and 39 GeV data show a substantial direct photon yield, which are comparable in magnitude and spectral shape, albeit within large uncertainties. For 62.4 GeV we can also extract a direct photon signal for 0%-20% and 20%-40% centrality selection and find that the direct photon yield increases with centrality. All observations are similar to those already published for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [5].

For the 200 GeV data, the centrality dependence of direct photons was analyzed in terms of the number of nucleon participants N_{part} . It was found that the integrated photon yield increases as $(N_{\text{part}})^\alpha$. This approach is not useful for comparing results across $\sqrt{s_{NN}}$, because N_{part} saturates at the same value for similar size systems at different beam energies. Instead we choose $dN_{ch}/d\eta$, which is a measured observable, for the comparison.

Direct photon production at high p_T is dominated by hard scattering and scales with the number of binary collisions N_{coll} . To study the interplay of bulk and hard particle production, we plot N_{coll} versus $dN_{ch}/d\eta$ at midrapidity for several beam energies in Fig. 2. The PHENIX data are taken from [47] and the $\sqrt{s_{NN}} = 2760$ GeV data are from ALICE [48]. The data exhibit a remarkable scaling between N_{coll} and $dN_{ch}/d\eta$, which takes the form:

$$N_{\text{coll}} = \frac{1}{SY(\sqrt{s_{NN}})} \times \left(\frac{dN_{ch}}{d\eta} \right)^\alpha \quad (3)$$




FIG. 2. Number of binary collisions, N_{coll} , vs. $dN_{ch}/d\eta|_{\eta=0}$, for four beam energies. All data are simultaneously fitted by a power-law with $\alpha = 1.25$. The error bars show the uncertainty from of N_{coll} from the Glauber MC.

Here we introduce the specific yield SY , which increases logarithmically with $\sqrt{s_{NN}}$ as $SY(\sqrt{s_{NN}}) = 0.98 \log(\sqrt{s_{NN}}) - 1.83$. In contrast, the power α is independent of $\sqrt{s_{NN}}$. We fit all data simultaneously and find $\alpha = 1.25 \pm 0.02$. This finding combined with the results of [49] imply that $N_{\text{coll}} \sim (dN_{ch}/d\eta)^{1.25} \sim N_{qp}^{1.25}$, where N_{qp} is the number of quark participants.

Panel (b) of Fig. 1 shows that at high- p_T the direct photon yield normalized by $(dN_{ch}/d\eta)^{1.25}$ is the same for all Au+Au centrality selections and for $p+p$ at $\sqrt{s_{NN}} = 200$ GeV. This is expected, because at high p_T photons are predominantly produced in hard scattering and there is no nuclear modification to the spectra. Above 3–4 GeV the data agree well with a perturbative quantum chromodynamics (pQCD) calculations for a normalization and factorization scale of $\mu = 0.5p_T$ [40, 50], also shown in the figure. The pQCD calculation was normalized to the experimental $dN_{ch}/d\eta$ for $\sqrt{s} = 200$ GeV from [51]. The $p+p$ data were fitted with a modified power law function $a(1 + p_T^2/b)^c$ [36]. Below 2 GeV/c the Au+Au yield is significantly enhanced compared to that from $p+p$. Surprisingly, the yield from Au+Au continues to follow the N_{coll} scaling behavior independent of centrality. We find a universal momentum spectrum of direct photons after they are normalized to $N_{\text{coll}} = (dN_{ch}/d\eta)^{1.25}$.




FIG. 3. Integrated direct photon yield ($p_T > 1.0 \text{ GeV}/c$) vs. $dN_{\text{ch}}/d\eta$, for data sets shown in Fig. 1. The dashed line is a power law fit with a fixed $\alpha = 1.25$. Also shown is the integrated yield of the fit to $p+p$ data and from the pQCD calculations, both scaled by N_{coll} [16, 40].




FIG. 4. Integrated direct photon yield ($p_T > 5.0 \text{ GeV}/c$) vs. $dN_{\text{ch}}/d\eta$, for different data sets. The dashed lines show power law fits to the data with fixed slope of $\alpha = 1.25$. Integrated yields from pQCD calculations scaled by N_{coll} are also shown.

In panels (a) and (c) of Fig. 1 we see that for $p_T < 2 \text{ GeV}/c$ the same scaling occurs not only for different centrality selections, but also for different $\sqrt{s_{NN}}$ and collision systems. At higher p_T the expected difference with $\sqrt{s_{NN}}$ is observed. Like for $\sqrt{s_{NN}} = 200 \text{ GeV}$, at high p_T the 2760 GeV data are well described by the pQCD calculation, though only above 5–6 GeV rather than 3–4 GeV. Note that the extrapolated pQCD calculations for $p+p$ at different \sqrt{s} seem to converge to the same normalized yield at low p_T as well, but at a tenth of the $A+A$ yield.

We quantify direct photon emission by integrating the invariant yield above $p_T = 1.0 \text{ GeV}/c$ and $p_T = 5.0 \text{ GeV}/c$. The integrals with the lower threshold will be dominated by excess low p_T photons unique to $A+A$

collisions, while the higher threshold characterizes the production in the pQCD region. The results are shown in Figs. 3 and 4 as a function of $dN_{\text{ch}}/d\eta$. Also plotted are power-law functions $A(dN_{\text{ch}}/d\eta)^\alpha$ with fixed $\alpha = 1.25$ and normalization fitted to the data.

For the lower p_T threshold the integrated yields from $A+A$ (Fig. 3) show the scaling behavior with $(dN_{\text{ch}}/d\eta)^{1.25}$ observed in Fig. 1. The $A+A$ points are compared to the integrated yield for $\sqrt{s} = 200 \text{ GeV}$ $p+p$ obtained from the fit to the data, which is scale with N_{coll} to the corresponding $dN_{\text{ch}}/d\eta$ for each $\sqrt{s_{NN}} = 200 \text{ GeV}$ $A+A$ point. The width of the band is given by the combined uncertainties of the fit function and N_{coll} . It is parallel to the $A+A$ trend but lower by about an order of magnitude. Also shown are the scaled integrated yields from pQCD calculations for $\sqrt{s} = 62.4, 200$, and 2760 GeV . They are consistent with the shaded band independent of beam energy.

For the p_T threshold of $5 \text{ GeV}/c$ the integrated yields from $Au+Au$ and $p+p$ at 200 GeV follow the same trend (Fig. 4), and are consistent with the pQCD calculation. The 2760 GeV data follow the same trend, but the yields are significantly higher. The pQCD calculation is slightly below the data.

While convenient and related to N_{coll} , the functional form $A(dN_{\text{ch}}/d\eta)^\alpha$ does not lend itself to an obvious interpretation. It is also not unique; for instance the data can be equally well fitted by $A(dN_{\text{ch}}/d\eta) + B(dN_{\text{ch}}/d\eta)^{4/3}$. The important point is that $A+A$ data from different centralities and a wide range of collision energies can be empirically described in terms of $dN_{\text{ch}}/d\eta$ with just two parameters, suggesting some fundamental commonality in the underlying physics.

There are two main conclusions from the $A+A$ data and the analyses presented in this paper. (i) At a given beam energy the direct photon yield scales with $dN_{\text{ch}}/d\eta^{1.25}$ or N_{coll} for all observed p_T , independent of centrality. There seems to be no qualitative change in the photon sources and/or their relative contributions. (ii) From $\sqrt{s_{NN}} = 39$ to 2760 GeV the same scaling is observed for $p_T < 2 \text{ GeV}/c$. This suggests that the main sources contributing to this p_T range are very similar also across beam energies.

If thermal radiation is the source of low p_T direct photons, the similarity at the same $dN_{\text{ch}}/d\eta$ across beam energies and centralities for $p_T < 2 \text{ GeV}/c$, suggests that the bulk of the matter that emits the radiation is similar in terms of temperature and space time evolution. This would be natural, if most of the photons are emitted near the transition from QGP to hadronic gas.

While at high p_T the scaled yields in $p+p$ and $A+A$ are identical, they differ by a factor of 10 at low p_T . This implies that there must be a transition from the small $p+p$ yield to the enhanced $A+A$ -like low p_T yields in the $dN_{\text{ch}}/d\eta$ range of ≈ 2 to 20 , which will be accessible with the data taken by PHENIX with small systems $p+Au$,

d +Au, and ${}^3\text{He}$ +Au.

We thank the staff of the Collider-Accelerator and Physics Departments at Brookhaven National Laboratory and the staff of the other PHENIX participating institutions for their vital contributions. We acknowledge support from the Office of Nuclear Physics in the Office of Science of the Department of Energy, the National Science Foundation, Abilene Christian University Research Council, Research Foundation of SUNY, and Dean of the College of Arts and Sciences, Vanderbilt University (U.S.A), Ministry of Education, Culture, Sports, Science, and Technology and the Japan Society for the Promotion of Science (Japan), Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundação de Amparo à Pesquisa do Estado de São Paulo (Brazil), Natural Science Foundation of China (People's Republic of China), Croatian Science Foundation and Ministry of Science and Education (Croatia), Ministry of Education, Youth and Sports (Czech Republic), Centre National de la Recherche Scientifique, Commissariat à l'Énergie Atomique, and Institut National de Physique Nucléaire et de Physique des Particules (France), Bundesministerium für Bildung und Forschung, Deutscher Akademischer Austausch Dienst, and Alexander von Humboldt Stiftung (Germany), J. Bolyai Research Scholarship, EFOP, the New National Excellence Program (ÚNKP), NKFIH, and OTKA (Hungary), Department of Atomic Energy and Department of Science and Technology (India), Israel Science Foundation (Israel), Basic Science Research Program through NRF of the Ministry of Education (Korea), Physics Department, Lahore University of Management Sciences (Pakistan), Ministry of Education and Science, Russian Academy of Sciences, Federal Agency of Atomic Energy (Russia), VR and Wallenberg Foundation (Sweden), the U.S. Civilian Research and Development Foundation for the Independent States of the Former Soviet Union, the Hungarian American Enterprise Scholarship Fund, the US-Hungarian Fulbright Foundation, and the US-Israel Binational Science Foundation.

-
- * Deceased
 † PHENIX Spokesperson: akiba@rcf.rhic.bnl.gov
- [1] P. Stankus, “Direct photon production in relativistic heavy-ion collisions,” *Ann. Rev. Nucl. Part. Sci.* **55**, 517 (2005).
 - [2] G. David, R. Rapp, and Z. Xu, “Electromagnetic Probes at RHIC-II,” *Phys. Rept.* **462**, 176 (2008).
 - [3] O. Linnyk, E. L. Bratkovskaya, and W. Cassing, “Effective QCD and transport description of dilepton and photon production in heavy-ion collisions and elementary processes,” *Prog. Part. Nucl. Phys.* **87**, 50 (2016).
 - [4] A. Adare *et al.* (PHENIX Collaboration), “Enhanced production of direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and implications for the initial temperature,” *Phys. Rev. Lett.* **104**, 132301 (2010).
 - [5] A. Adare *et al.* (PHENIX Collaboration), “Centrality dependence of low-momentum direct-photon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **91**, 064904 (2015).
 - [6] A. Adare *et al.* (PHENIX Collaboration), “Observation of direct-photon collective flow in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions,” *Phys. Rev. Lett.* **109**, 122302 (2012).
 - [7] A. Adare *et al.* (PHENIX Collaboration), “Azimuthally anisotropic emission of low-momentum direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **94**, 064901 (2016).
 - [8] L. Adamczyk *et al.* (STAR Collaboration), “Direct virtual photon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Lett. B* **770**, 451 (2017).
 - [9] J. Adam *et al.* (ALICE Collaboration), “Direct photon production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2760$ GeV,” *Phys. Lett. B* **754**, 235 (2016).
 - [10] Daniel Lohner (ALICE Collaboration), “Measurement of Direct-Photon Elliptic Flow in Pb-Pb Collisions at $\sqrt{s_{NN}} = 2760$ GeV,” *Proceedings, Workshop for Young Scientists on the Physics of Ultrarelativistic Nucleus-Nucleus Collisions (Hot Quarks 2012): Copamarina, Puerto Rico, October 14-20, 2012*, *J. Phys. Conf. Ser.* **446**, 012028 (2013).
 - [11] H. van Hees, C. Gale, and R. Rapp, “Thermal Photons and Collective Flow at the Relativistic Heavy-Ion Collider,” *Phys. Rev. C* **84**, 054906 (2011).
 - [12] H. van Hees, Min He, and R. Rapp, “Pseudo-critical enhancement of thermal photons in relativistic heavy-ion collisions?” *Nucl. Phys. A* **933**, 256 (2015).
 - [13] M. Dion, J.-F. Paquet, B. Schenke, Clint Young, S. Jeon, and C. Gale, “Viscous photons in relativistic heavy ion collisions,” *Phys. Rev. C* **84**, 064901 (2011).
 - [14] C. Shen, U. W Heinz, J.-F. Paquet, and C. Gale, “Thermal photons as a quark-gluon plasma thermometer reexamined,” *Phys. Rev. C* **89**, 044910 (2014).
 - [15] C. Shen, J. F. Paquet, G. S. Denicol, S. Jeon, and C. Gale, “Thermal photon radiation in high multiplicity p +Pb collisions at the Large Hadron Collider,” *Phys. Rev. Lett.* **116**, 072301 (2016).
 - [16] J. F. Paquet, C. Shen, Gabriel S. Denicol, M. Luzum, B. Schenke, S. Jeon, and C. Gale, “Production of photons in relativistic heavy-ion collisions,” *Phys. Rev. C* **93**, 044906 (2016).
 - [17] E. L. Bratkovskaya, S. M. Kiselev, and G. B. Sharkov, “Direct photon production from hadronic sources in high-energy heavy-ion collisions,” *Phys. Rev. C* **78**, 034905 (2008).
 - [18] E. L. Bratkovskaya, “Phenomenology of photon and dilepton production in relativistic nuclear collisions,” *Proceedings, 24th International Conference on Ultra-Relativistic Nucleus-Nucleus Collisions (Quark Matter 2014): Darmstadt, Germany, May 19-24, 2014*, *Nucl. Phys. A* **931**, 194 (2014).
 - [19] O. Linnyk, W. Cassing, and E. L. Bratkovskaya, “Centrality dependence of the direct photon yield and elliptic flow in heavy-ion collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **89**, 034908 (2014).
 - [20] M. Chiu, T. K. Hemmick, V. Khachatryan, Andrey Leonidov, Jinfeng Liao, and Larry McLerran, “Production of Photons and Dileptons in the Glasma,” *Nucl. Phys. A* **900**, 16 (2013).
 - [21] L. McLerran and B. Schenke, “The Glasma, Photons and the Implications of Anisotropy,” *Nucl. Phys. A* **929**, 71

- (2014).
- [22] L. McLerran and B. Schenke, “A Tale of Tails: Photon Rates and Flow in Ultra-Relativistic Heavy Ion Collisions,” *Nucl. Phys. A* **946**, 158 (2016).
 - [23] J. Berges, K. Reygers, N. Tanji, and R. Venugopalan, “Parametric estimate of the relative photon yields from the plasma and the quark-gluon plasma in heavy-ion collisions,” *Phys. Rev. C* **95**, 054904 (2017).
 - [24] Akihiko Monnai, “Thermal photon v_2 with slow quark chemical equilibration,” *Phys. Rev. C* **90**, 021901 (2014).
 - [25] C.-H. Lee and I. Zahed, “Electromagnetic Radiation in Hot QCD Matter: Rates, Electric Conductivity, Flavor Susceptibility and Diffusion,” *Phys. Rev. C* **90**, 025204 (2014).
 - [26] Simon Turbide, R. Rapp, and C. Gale, “Hadronic production of thermal photons,” *Phys. Rev. C* **69**, 014903 (2004).
 - [27] K. Dusling and I. Zahed, “Thermal photons from heavy ion collisions: A spectral function approach,” *Phys. Rev. C* **82**, 054909 (2010).
 - [28] M. Heffernan, P. Hohler, and R. Rapp, “Universal Parametrization of Thermal Photon Rates in Hadronic Matter,” *Phys. Rev. C* **91**, 027902 (2015).
 - [29] O. Linnyk, V. Konchakovski, T. Steinert, W. Cassing, and E. L. Bratkovskaya, “Hadronic and partonic sources of direct photons in relativistic heavy-ion collisions,” *Phys. Rev. C* **92**, 054914 (2015).
 - [30] G. Basar, Dmitri Kharzeev, and Vladimir Skokov, “Conformal anomaly as a source of soft photons in heavy ion collisions,” *Phys. Rev. Lett.* **109**, 202303 (2012).
 - [31] G. Basar, Dmitri E. Kharzeev, and E. V. Shuryak, “Magneto-sanoluminescence and its signatures in photon and dilepton production in relativistic heavy ion collisions,” *Phys. Rev. C* **90**, 014905 (2014).
 - [32] B. Muller, Shang-Yu Wu, and Di-Lun Yang, “Elliptic flow from thermal photons with magnetic field in holography,” *Phys. Rev. D* **89**, 026013 (2014).
 - [33] A. Ayala, P. Mercado, and C. Villavicencio, “Magnetic catalysis of a finite size pion condensate,” *Phys. Rev. C* **95**, 014904 (2017).
 - [34] V. V. Goloviznin, A. V. Nikolskii, A. M. Snigirev, and G. M. Zinovjev, “Probing a confinement by direct photons and dileptons,” arXiv:1804.00559.
 - [35] M. Allen *et al.* (PHENIX Collaboration), “PHENIX inner detectors,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 549 (2003).
 - [36] A. Adare *et al.* (PHENIX Collaboration), “Low-momentum direct photon measurement in Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV,” arXiv:1805.xxxxx.
 - [37] A. L. S. Angelis *et al.* (CERN-Columbia-Oxford-Rockefeller, CCOR Collaboration), “Search for Direct Single Photon Production at Large p_T in Proton Proton Collisions at $\sqrt{s} = 62.4$ GeV,” *Phys. Lett. B* **94**, 106 (1980).
 - [38] A. L. S. Angelis *et al.* (CMOR Collaboration), “Direct Photon Production at the CERN ISR,” *Nucl. Phys. B* **327**, 541 (1989).
 - [39] T. Akesson *et al.* (Axial Field Spectrometer Collaboration), “High $p_T\gamma$ and π^0 Production, Inclusive and With a Recoil Hadronic Jet, in pp Collisions at $\sqrt{s} = 63$ GeV,” *Sov. J. Nucl. Phys.* **51**, 836 (1990), [Yad. Fiz. 51, 1314 (1990)].
 - [40] J. F. Paquet, (2017), private communication, uses nuclear PDF nCTEQ15-np and photon fragmentation function BFG-II.
 - [41] W. Anderson *et al.*, “Design, Construction, Operation and Performance of a Hadron Blind Detector for the PHENIX Experiment,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **646**, 35 (2011).
 - [42] K. Adcox *et al.* (PHENIX Collaboration), “PHENIX central arm tracking detectors,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 489 (2003).
 - [43] M. Aizawa *et al.* (PHENIX Collaboration), “PHENIX central arm particle ID detectors,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 508 (2003).
 - [44] L. Aphecetche *et al.* (PHENIX Collaboration), “PHENIX calorimeter,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 521 (2003).
 - [45] A. Adare *et al.* (PHENIX Collaboration), “Evolution of π^0 suppression in Au+Au collisions from $\sqrt{s_{NN}} = 39$ to 200 GeV,” *Phys. Rev. Lett.* **109**, 152301 (2012).
 - [46] A. Adare *et al.* (PHENIX Collaboration), “Neutral pion production with respect to centrality and reaction plane in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **87**, 034911 (2013).
 - [47] A. Adare *et al.* (PHENIX Collaboration), “Transverse energy production and charged-particle multiplicity at midrapidity in various systems from $\sqrt{s_{NN}} = 7.7$ to 200 GeV,” *Phys. Rev. C* **93**, 024901 (2016).
 - [48] Kenneth Aamodt *et al.* (ALICE Collaboration), “Centrality dependence of the charged-particle multiplicity density at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2760$ GeV,” *Phys. Rev. Lett.* **106**, 032301 (2011).
 - [49] S. S. Adler *et al.* (PHENIX Collaboration), “Transverse-energy distributions at midrapidity in p+p, d+Au, and Au+Au collisions at $\sqrt{s_{NN}} = 62.4 - 200$ GeV and implications for particle-production models,” *Phys. Rev. C* **89**, 044905 (2014).
 - [50] A. P., Michel Fontannaz, Jean-Philippe Guillet, Eric Pilon, and Monique Werlen, “A New critical study of photon production in hadronic collisions,” *Phys. Rev. D* **73**, 094007 (2006).
 - [51] C. Patrignani *et al.* (Particle Data Group Collaboration), “Review of Particle Physics,” *Chin. Phys. C* **40**, 100001 (2016).