

Bayesian inverse regression for supervised dimension reduction with small datasets

Xin Cai*, Guang Lin†, Jinglai Li‡

Abstract

We consider supervised dimension reduction problems, namely to identify a low dimensional projection of the predictors \mathbf{x} which can retain the statistical relationship between \mathbf{x} and the response variable y . We follow the idea of the sliced inverse regression (SIR) class of methods, which is to use the statistical information of the conditional distribution $\pi(\mathbf{x}|y)$ to identify the dimension reduction (DR) space and in particular we focus on the task of computing this conditional distribution. We propose a Bayesian framework to compute the conditional distribution where the likelihood function is obtained using the Gaussian process regression model. The conditional distribution $\pi(\mathbf{x}|y)$ can then be obtained directly by assigning weights to the original data points. We then can perform DR by considering certain moment functions (e.g. the first moment) of the samples of the posterior distribution. With numerical examples, we demonstrate that the proposed method is especially effective for small data problems.

1 Introduction

In many supervised learning problems, especially regression problems, one frequently has to deal with small data problems where the available data are insufficient to provide a robust regression. If doing regression directly in such problems, one often risks of certain degree of overfitting or being incorrectly regularized. In either case, the resulting regression model may lose its prediction accuracy. Extracting and selecting the important features or eliminating the redundant ones is a key step to avoid overfitting and improve the robustness of the regression task [9]. The feature extraction and selection thus constitutes of identifying a low dimensional subspace of the predictors \mathbf{x} which retains the statistical relationship between \mathbf{x} and the response y , i.e. a supervised dimension reduction problem. Mathematically such problems are often posed as to estimate the central dimension reduction (DR) subspace [5]. A very popular class of methods

*School of Mathematical Sciences, Shanghai Jiao Tong University, Shanghai, China.

†Department of Mathematics, Purdue University, West Lafayette, IN, USA.

‡Department of Mathematical Sciences, University of Liverpool, Liverpool L69 7ZL, UK.
Email: jinglai.li@liverpool.ac.uk.

estimate this central subspace by considering the statistics of the predictors \mathbf{x} conditional on the response y , and such methods include the sliced inverse regression (SIR) proposed in the seminal work [17], the sliced average variance estimation [6], and many of their variants, e.g. [5, 16, 18, 31, 3, 15, 20, 27, 19]. Some of the extensions and variants have been developed specifically for supervised machine learning problems, e.g., [10, 30, 12]. Asymptotic analysis of these methods has also been extensively studied [33, 11, 32]. The literature in this topic is vast and we refer to [21, 14] for a more comprehensive list of references. We note here that most of these methods adopt nonparametric formulation without assuming any specific relation between \mathbf{x} and y , and some exceptions do exist, such as the likelihood acquired direction (LAD) method [4, 3], which uses specified likelihoods function to compute the reduced dimensions.

In this work we shall focus on the SIR type of methods to identify the DR space. As we can see, many of this type of methods focus on the question: what statistical information of the conditional distribution $\pi(\mathbf{x}|y)$ should one use to obtain the DR subspace? In this work however we consider a different aspect of the problem: how to obtain the conditional distribution $\pi(\mathbf{x}|y)$ especially when the data set is small? In SIR and most its variants, the conditional moments are approximately estimated by slicing the data [17]. As will be demonstrated with numerical examples, the slicing method does not perform well if we have a very small data set. The main purpose of the work is to address the problem of computing the conditional distribution $\pi(\mathbf{x}|y)$. In particular we present a Bayesian approach which can provide not only the first or second moments, but the full conditional distribution $\pi(\mathbf{x}|y)$, and once the distribution is available one can use any desired method to estimate the DR subspace based on the conditional distribution. Just like [4, 3], our method also involves constructing the likelihood function $\pi(y|\mathbf{x})$, but a main difference here is that our method estimates the likelihood function with a Gaussian Process (GP) regression. Once the likelihood function is available, we can compute the posterior distribution $\pi(\mathbf{x}|y)$ from the likelihood function and a desired prior distribution. In this work we choose to use the first order moment of the conditional distribution (following SIR) to demonstrate the method, while noting that the method can be easily extended to second or higher conditional moments. It should also be noted that the works [28, 22, 26] also propose methods to compute the reduced dimensions in a Bayesian framework. These methods, however, are fundamentally different from the present one. First in all these existing methods, prior distributions need to be assumed on the DR subspace or the related parameters, while in the proposed method, we do not impose any prior assumptions on the DR subspace. Second in [22, 26], either the forward [26] or the inverse [22] conditional distribution is assumed to be in a mixture form, while our method uses a GP nonparametric framework to model the conditional distribution $\pi(y|\mathbf{x})$, which can provide more flexibility than a mixture model. Finally in all the model-based approaches, Bayesian or non-Bayesian, the DR directions are either sampled with Markov chain Monte Carlo (MCMC) or computed with Maximum likelihood estimation (MLE), and both approaches can be rather computationally intensive. The proposed method here does not need to perform either MCMC

or MLE, and its computational cost is about the same as the standard SIR.

To summarize, the main contribution of the work is to propose a Bayesian framework method that allows to directly generate samples from the conditional distribution $\pi(\mathbf{x}|y)$ for any value of y and by doing so it avoids slicing the samples as is done in SIR, which makes it particularly effective for problems with very small numbers of data. The method uses non-parametric GP model to construct the likelihood function, which can rather flexibly and efficiently characterize the relation between \mathbf{x} and y .

The rest of this paper is organized as follows. In section 2 we set up a formulation of dimension reduction and go through the basic idea of the classic dimension reduction approach SIR. The BIR is introduced explicitly in Section 3, including a Bayesian formulation and the tool GP we used, and an algorithm will be given. In section 4 we provide several numerical examples. Section 6 offers concluding remarks.

2 Dimension reduction and sliced inverse regression

2.1 Problem setup

We consider a generic supervised dimension reduction problem. Let \mathbf{x} be a p -dimensional random variable defined on R^p following a distribution $\pi_0(\mathbf{x})$, and suppose that we are interested in a scalar function of \mathbf{x} , which ideally can be written as,

$$y = f(\mathbf{b}_1^T \mathbf{x}, \mathbf{b}_2^T \mathbf{x}, \dots, \mathbf{b}_K^T \mathbf{x}, \epsilon), \quad (1)$$

where \mathbf{b}_k for $k = 1 \dots K$ are some p -dimensional vectors, and ϵ is small noise independent of \mathbf{x} . It should be clear that, when this model holds, the projection of the p -dimension variable \mathbf{x} onto the k dimensional subspace of R^p spanned by $\{\mathbf{b}_1, \dots, \mathbf{b}_K\}$, captures all the information of \mathbf{x} with respect to y , and if $K < p$, we can achieve the goal of data reduction by estimating the coefficients $\{\mathbf{b}_k\}_{k=1}^K$. In practice, both the explicit expression of f and the coefficients $\{\mathbf{b}_k\}_{k=1}^K$ are unknown, and instead we have a set of data pairs $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$ drawn from the joint distribution $\pi(\mathbf{x}, y)$ defined by π_0 and Eq. (1). Finding a set of $\{\mathbf{b}_k\}_{k=1}^K$ that satisfy the Eq. (1) from the given data set $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$ is the task of supervised dimension reduction. In what follows we shall refer to the coefficients $\{\mathbf{b}_k\}_{k=1}^K$ as dimension-reduction (DR) directions, and the linear space B spanned by the $\{\mathbf{b}_k\}_{k=1}^K$ as the DR subspace. For a more formal and generic description of the DR problem (in the Central DR Subspace and Sufficient Dimension Reduction framework) we refer to [5].

2.2 Sliced inverse regression

The SIR approach [17] estimates the DR directions based on the idea of inverse regression (IR). In contrast to the forward regression $E(y | \mathbf{x})$, IR regresses each coordinate of \mathbf{x} against y . Thus as y varies, $E(\mathbf{x} | y)$ draws a curve in R^p along

the y coordinate, whose center is located at $E(E(\mathbf{x}|y)) = E(\mathbf{x})$. For simplicity we shall assume that throughout this section \mathbf{x} is a standardized random variable: namely $E(\mathbf{x}) = 0$ and $\text{Cov}(\mathbf{x}) = I$. Under the following condition the IR curve $E(\mathbf{x}|y)$ is contained in the DR subspace B [17]:

Condition 2.1. For any $\beta \in R^p$, the conditional expectation $E(\beta^T \mathbf{x} | \mathbf{b}_1^T \mathbf{x}, \dots, \mathbf{b}_K^T \mathbf{x})$ is linear in $\mathbf{b}_1^T \mathbf{x}, \dots, \mathbf{b}_K^T \mathbf{x}$.

This condition is satisfied when the distribution of \mathbf{x} is elliptically symmetric [17]. An important implication of this property is that the covariance matrix $\text{Cov}[E(\mathbf{x}|y)]$ is degenerated in any direction orthogonal to the DR subspace B . We see, therefore, that the eigenvectors associated with the largest K eigenvalues of $\text{Cov}[E(\mathbf{x}|y)]$ are the DR directions. So the key of estimating the DR direction is to obtain the covariance of the conditional expectation of the data, $\text{Cov}[E(\mathbf{x}|y)]$.

One of the most popular approaches to estimate the covariance $\text{Cov}[E(\mathbf{x}|y_j)]$ is SIR. Simply put, SIR produces a crude estimate of $E(\mathbf{x}|y)$, by slicing the data $(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_n, y_n)$ into H partitions according to the value of y_j and then estimating $E(\mathbf{x}|y \in I_h), h = 1, \dots, H$ using the data inside the interval I_h for each $h = 1, \dots, H$. Finally one use the H samples to compute an estimate of the covariance matrix $\text{Cov}[E(\mathbf{x}|y)]$. A complete SIR scheme is described as follows:

1. Divide range of y into H slices, I_1, \dots, I_H . Let the proportion of the y_i that falls in slice I_h be \hat{p}_h , i.e.,

$$\hat{p}_h = \frac{1}{n} \sum_{j=1}^n \delta_h(y_j),$$

where $\delta_h(y_j)$ takes the values 0 or 1 depending on whether y_j falls into the h th slice I_h or not.

2. Within each slice, compute the sample mean of the \mathbf{x}_j 's, denoted by $\hat{\mathbf{x}}_h (h = 1, \dots, H)$:

$$\hat{\mathbf{x}}_h = \frac{1}{(n\hat{p}_h)} \sum_{y_j \in I_h} \mathbf{x}_j.$$

3. Compute the weighted covariance matrix

$$\hat{C} = \sum_{h=1}^H \hat{p}_h \hat{\mathbf{x}}_h \hat{\mathbf{x}}_h^T.$$

4. Perform eigenvalue decomposition of \hat{C} , and return the eigenvectors associated with the k largest eigenvalues as the estimated DR directions $\hat{\mathbf{b}}_1, \dots, \hat{\mathbf{b}}_K$.

As is mentioned in Section 1, the slicing treat is often not sufficiently accurate when the data set is small, and in what follows we shall provide an alternative to compute the covariance matrix.

3 Bayesian inverse regression

3.1 Bayesian formulation for $\pi(\mathbf{x}|y)$

First recall that in the SIR framework, a key step is to compute the covariance $\text{Cov}[E(\mathbf{x}|y)]$. A natural choice to estimate the covariance $\text{Cov}[E(\mathbf{x}|y)]$ is to use the sample covariance of the data points,

$$\hat{C} = \frac{1}{n-1} \sum_{j=1}^n (\hat{\mathbf{x}}_j - \bar{\mathbf{x}})(\hat{\mathbf{x}}_j - \bar{\mathbf{x}})^T, \quad \bar{\mathbf{x}} = \frac{1}{n} \sum_{j=1}^n \hat{\mathbf{x}}_j, \quad (2)$$

where $\hat{\mathbf{x}}_j$ is an estimate of $E(\mathbf{x}|y_j)$ for all $j = 1 \dots n$, and (y_1, \dots, y_n) are the data points. Next we need to compute $\hat{\mathbf{x}}_j$, the estimate of $E(\mathbf{x}|y_j)$, and we propose to do so in a Bayesian framework. Namely we formulate the problem as to compute the posterior distribution:

$$\pi(\mathbf{x}|y) \propto \pi(y|\mathbf{x})\pi(\mathbf{x}), \quad (3)$$

where $\pi(y|\mathbf{x})$ is the likelihood function and $\pi(\mathbf{x})$ is the prior of \mathbf{x} .

We consider the prior distribution $\pi(\mathbf{x})$ first. To start we note that the choice of prior does not affect the DR subspace as this subspace structure lies in the function $f(\mathbf{x}, \epsilon)$ in Eq. (1) rather than the distribution of \mathbf{x} . As such, in principle one may use any prior distribution. However, if the chosen $\pi(\mathbf{x})$ is too different from π_0 , the GP model constructed from the data (following $\pi_0(\mathbf{x})$) may not be accurate for the samples drawn according to $\pi(\mathbf{x})$, which in turn may hurt the accuracy of the posterior $\pi(\mathbf{x}|y)$. For this reason one should use a prior distribution that is close to π_0 , and a natural choice is to perform a density estimation for the data $\{\mathbf{x}_j\}_{j=1}^n$ and use the estimated density as the prior. For example, one may use Gaussian mixtures [24] to estimate the prior distribution from the data $\{\mathbf{x}_j\}_{j=1}^n$. On the other hand, when the dimension of \mathbf{x} is high, estimating the density of \mathbf{x} can be challenging. In this case, we can just use the original data points $\{\mathbf{x}_j\}_{j=1}^n$, where the prior is simply π_0 .

3.2 The GP regression

The next step is to construct the likelihood function $\pi(y|\mathbf{x})$ from data, which, as mentioned earlier, is done by using the GP regression model.

Simply speaking the GP regression performs a nonparametric regression in a Bayesian framework [29]. The main idea of the GP method is to assume that the function of interest $f(\mathbf{x}, \epsilon)$ is a realization from a Gaussian random field, whose mean is $\mu(\mathbf{x})$ and covariance is specified by a kernel function $k(\mathbf{x}, \mathbf{x}')$, namely,

$$\text{Cov}[f(\mathbf{x}), f(\mathbf{x}')] = k(\mathbf{x}, \mathbf{x}').$$

The kernel $k(\mathbf{x}, \mathbf{x}')$ is positive semidefinite and bounded.

Now given the data points $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$, we want to predict the value of y at a new point \mathbf{x} . Now we let $\mathbf{X} := [\mathbf{x}_1, \dots, \mathbf{x}_n]$, and $\mathbf{Y} = [y_1, \dots, y_n]$. Under the

GP assumption, it is easy to see that the joint distribution of (\mathbf{Y}, y) is Gaussian,

$$\begin{bmatrix} \mathbf{Y} \\ y \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu(\mathbf{X}) \\ \mu(\mathbf{x}) \end{bmatrix}, \begin{bmatrix} K(\mathbf{X}, \mathbf{X}) + \sigma_n^2 I & K(\mathbf{X}, \mathbf{x}) \\ K(\mathbf{x}, \mathbf{X}) & K(\mathbf{x}, \mathbf{x}) \end{bmatrix} \right), \quad (4)$$

where σ_n^2 is the variance of observation noise, I is an identity matrix, and the notation $K(\mathbf{A}, \mathbf{B})$ denotes the matrix of the covariance evaluated at all pairs of points in set \mathbf{A} and in set \mathbf{B} using the kernel function $k(\cdot, \cdot)$.

It follows immediately from Eq. (4) that the conditional distribution $\pi_{GP}(y|\mathbf{x}, \mathbf{X}, \mathbf{Y})$ is also Gaussian:

$$\pi_{GP}(y|\mathbf{x}, \mathbf{X}, \mathbf{Y}) = \mathcal{N}(\mu_{\text{pos}}, \sigma_{\text{pos}}^2), \quad (5a)$$

where the posterior mean and variance are,

$$\begin{aligned} \mu_{\text{pos}}(\mathbf{x}) &= \mu(\mathbf{x}) + k(\mathbf{x}, \mathbf{X})(k(\mathbf{X}, \mathbf{X}) + \sigma_n^2 I)^{-1}(\mathbf{Y} - \mu(\mathbf{X})), \\ \sigma_{\text{pos}}^2 &= k(\mathbf{x}, \mathbf{x}) - k(\mathbf{x}, \mathbf{X})(k(\mathbf{X}, \mathbf{X}) + \sigma_n^2 I)^{-1}k(\mathbf{X}, \mathbf{x}). \end{aligned}$$

There are also a number of technical issues in the GP model, such as choosing the kernel function and determining the hyperparameters. For detailed discussion of these matters, we refer the readers to [29]. In what follows we shall use the GP posterior as the likelihood function, i.e., letting $\pi(y|\mathbf{x}) = \pi_{GP}(y|\mathbf{x}, \mathbf{X}, \mathbf{Y})$.

3.3 Computing the posterior mean

Once we obtain the likelihood function and the prior, we can draw samples from the posterior distribution (3) with MCMC. An alternative strategy is to use the original data points in an importance sampling (IS) formulation. Namely if we choose $\pi = \pi_0$, the original data set $\{\mathbf{x}_j\}_{j=1}^n$ follow the prior distribution, and for each \mathbf{x}_j we can compute the weight $w_j = \pi(y|\mathbf{x}_j)$. Finally the weights w_1, \dots, w_n are normalized so that $\sum_{j=1}^n w_j = 1$. We thus obtain obtain a set of weighted samples $\{(\mathbf{x}_j, w_j)\}_{j=1}^n$ drawn from the posterior $\pi(\mathbf{x}|y)$. Now let $\{(\mathbf{x}_j, w_j)\}_{j=1}^n$ be a set samples draw from the posterior, and we can estimate $E(\mathbf{x}|y)$ as

$$\hat{\mathbf{x}} = \sum_{j=1}^n w_j \mathbf{x}_j. \quad (6)$$

We repeat this procedure for each y_j for $j = 1 \dots n$, and then use Eq. (2) to compute $\text{Cov}[E(\mathbf{x}|y)]$. Since we use a Bayesian method to estimate $E(\mathbf{x}|y)$, we refer to proposed method as Bayesian inverse regression (BIR). We reinstate here that BIR essentially provides a means to draw samples from the conditional distribution $\pi(\mathbf{x}|y)$, and its application is not limited to estimate $E(\mathbf{x}|y)$, and it is possible to make use of the conditional distribution in a different manner. We also note that, using the Bayesian framework, we avoid slicing the data. Another issue that should be mentioned here is how to select the number of reduced dimensions; since BIR is also a method based on the eigenvalue decomposition of $\text{Cov}[E(\mathbf{x}|y)]$, the methods used in [17] and related works, e.g., [8], can be used directly here. Finally we present the BIR algorithm with MCMC in Alg. 1 and that with the original data in Alg. 2.

Algorithm 1 The Bayesian inverse regression algorithm with MCMC

Require: $\{(\mathbf{x}_j, y_j)\}_{j=1}^n, n_{MC}$

Ensure: The estimated DR directions: $\hat{\mathbf{b}}_1, \dots, \hat{\mathbf{b}}_K$

- 1: Construct the GP model from data $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$: $\pi_{GP}(y|\mathbf{x}, \mathbf{X}, \mathbf{Y})$;
 - 2: **for** $j = 1$ to n **do**
 - 3: Draw n_{MC} samples from $\pi_{GP}(y_j|\mathbf{x}, \mathbf{X}, \mathbf{Y})\pi(\mathbf{x})$: $\{\mathbf{x}_i\}_{i=1}^{n_{MC}}$;
 - 4: Compute $\hat{\mathbf{x}}_j = \frac{1}{n_{MC}} \sum_{i=1}^{n_{MC}} \mathbf{x}_i$;
 - 5: Compute \hat{C} using Eq. (2) and $\{\hat{\mathbf{x}}_j\}_{j=1}^n$;
 - 6: Perform eigenvalue decomposition of \hat{C} ;
 - 7: Return the eigenvectors associated with the k largest eigenvalues as $\hat{\mathbf{b}}_1, \dots, \hat{\mathbf{b}}_K$.
-

Algorithm 2 The Bayesian inverse regression algorithm with the original data

Require: $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$

Ensure: The estimated DR directions: $\hat{\mathbf{b}}_1, \dots, \hat{\mathbf{b}}_K$

- 1: Construct the GP model from data $\{(\mathbf{x}_j, y_j)\}_{j=1}^n$: $\pi_{GP}(y|\mathbf{x}, \mathbf{X}, \mathbf{Y})$;
 - 2: **for** $j = 1$ to n **do**
 - 3: **for** $i = 1$ to n **do**
 - 4: Let $w_i = \pi_{GP}(y_j|\mathbf{x}_i, \mathbf{X}, \mathbf{Y})$;
 - 5: Renormalize $\{w_i\}_{i=1}^n$ so that $\sum_{i=1}^n w_i = 1$;
 - 6: Compute $\hat{\mathbf{x}}_j = \sum_{i=1}^n w_i \mathbf{x}_i$;
 - 7: Compute \hat{C} using Eq. (2) and $\{\hat{\mathbf{x}}_j\}_{j=1}^n$;
 - 8: Perform eigenvalue decomposition of \hat{C} ;
 - 9: Return the eigenvectors associated with the k largest eigenvalues as $\hat{\mathbf{b}}_1, \dots, \hat{\mathbf{b}}_K$.
-

4 Numerical examples

In this section we compare the performance of the proposed BIR method with three common methods, SIR, likelihood-based DR (LDR) [4], and the Localized SIR (LSIR), in two mathematical and two real-data examples. The first example uses data simulated from a mathematical function, with which we want to exam the scalability of the methods with respect to the dimensionality of the problem. The second one is also a mathematical example, and with this example we compare the performance of different methods affected by the non-ellipticity of the distribution of \mathbf{x} . Our last example is based on real data, in which we compare the performance of different methods in the small data situation. In the GP model used in all the examples, we set the prior mean $\mu(\mathbf{x}) = 0$, and choose the Automatic Relevance Determination (ARD) squared exponential kernel [29]:

$$k(\mathbf{x}, \mathbf{x}') = \sigma_0^2 \exp\left(-\frac{1}{2} \sum_{i=1}^P \frac{(x_i - x'_i)^2}{\lambda_i^2}\right), \quad (7)$$




Figure 1: The posterior mean of the TG prior.

where the hyperparameters $\sigma_0, \lambda_1, \dots, \lambda_d$, and the σ_n are determined by maximum likelihood estimation [29]. In the first two mathematical examples, we use BIR Algorithm 1 with 10000 MCMC samples, and in the two real data examples, we use BIR Algorithm 2.

4.1 Mathematical examples with increasing dimensions

First we consider a d -dimensional problem where \mathbf{x} follows a standard normal distribution. The data are simulated from the following functions:

$$f(\mathbf{x}, \epsilon) = x_1(x_2 + x_3) + 0.5\epsilon, \quad (8a)$$

$$f(\mathbf{x}, \epsilon) = \frac{x_1 + x_2 + x_3}{0.5 + (x_4 + x_5)^2} + 0.1\epsilon, \quad (8b)$$

where $\epsilon \sim N(0, 1)$. Both problems have two DR directions. In the regression content, a well known limitation of the GP method is that it can not handle high dimension, and so here we want to test the scalability of the BIR method with respect to dimensionality. To do so we perform experiments for various dimensions: $d = 10, 20, 30, 40, 50$, where we set the number of data points to be $n = 5d$, i.e., growing linear with respect to dimensionality. To evaluate the performance of the methods, we use the R^2 metric of accuracy used in [17] to measure the accuracy of the DR subspace and the DR directions.

We repeat all the tests for 100 times and report the average. Specifically, we show the R^2 -accuracy of the DR subspace B and the two DR directions in Fig. 1. We can see that the BIR method has the best performance in all the tests in the two examples, except one situation: $d = 10$ for function 8b. The R^2 accuracy for each dimensions provide more information on the results. Namely, for Function 8a, BIR performs better than all the other methods in




Figure 2: The data points of (x_1, x_2) for different values of b .

both of the directions. For function 8b, the accuracy of BIR is slightly lower than than SIR and LSIR for the first direction, but it achieves significantly higher accuracy on the 2nd dimension than all the other methods. Finally we want to note here that as the dimensionality increases, the performance of BIR does not decay evidently, suggesting that the method can handle rather high dimensional problems.

4.2 Mathematical examples with non-Gaussian distributions

In our second example, we want to test the performance of the methods when the distribution of \mathbf{x} is strongly non-Gaussian. We assume \mathbf{x} is a 10-dimensional variable and the data are generated as follows. First let $\mathbf{u} = (u_1, u_2)$ follow a two-dimensional standard normal distribution. We then perform the following transform:

$$x_1 = u_1, \quad x_2 = u_1 - bu_1^2, \quad (9)$$

where $b \geq 0$. Here as we can see by varying parameter b we can control how different the distribution from Gaussian. Data of y are generated from \mathbf{u} , and so the transformation used for generating \mathbf{x} does not affect the data of y . Here we consider two functions for generating y :

$$y = \frac{u_1}{0.5 + (u_2 + 1.5)^2} + 0.5\epsilon, \quad (10a)$$

$$y = \sin(5\pi u_1) + u_2^2 + 0.1\epsilon, \quad (10b)$$

where $\epsilon \sim N(0, 1)$. In this test, we choose five different values of b : $b = 0, 5, 10, 15, 20$ with sample size $n = 100$, and we show the scatter plots of the data points for all these cases in Fig. 2, where we can see that the resulting data points move apart from Gaussian as b increases. We plot the R^2 accuracy against the value of b in Fig. 3 for both functions. From the figure we can see that for function 10a, BIR clearly outperforms all the other methods for all the values of b , and for function 10b, the BIR also has the best performance in all the cases, with LDR being about the same at $b = 10$ and 20. The results demonstrate that the BIR method performs well for highly non-Gaussian distributions.




Figure 3: The R^2 -accuracy of the DR subspace plotted as a function of b , for function (10a) (left) and function (10b) (right) respectively.

4.3 Death rate prediction

Methods	$n = 15$	$n = 20$	$n = 25$	$n = 30$	$n = 35$	$n = 40$
w/o DR	.1832 (.2013)	.0855 (.0502)	.0551 (.0216)	.0460 (.0171)	.0425 (.0161)	.0380 (.0089)
LDR	- (-)	.0823 (.0518)	.0569 (.0207)	.0490 (.0184)	.0444 (.0173)	.0383 (.0089)
SIR	.4403 (1.2417)	.0982 (.0769)	.0653 (.0310)	.0548 (.0216)	.0525 (.0217)	.0430 (.0108)
LSIR	- (-)	.0876 (.0461)	.0648 (.0252)	.0557 (.0224)	.0485 (.0174)	.0429 (.0100)
BIR	.0484 (.0110)	.0451 (.0104)	.0465 (.0111)	.0481 (.0105)	.0466 (.0114)	.0468 (.0126)

Table 1: The mean and the standard deviation (in parenthesis) of MRRE for Example 3. The best results are marked in bold.

Our third example is to use pollution and related factors to predict the death rate [23, 1]. This is a regression problem with 15 predictors and 60 data points and we choose this example to test how the methods perform with very small number of data. We first apply the DR methods to select one feature (we have conducted tests with 2 and 3 features which does not improve the regression accuracy, and so we omit those results here) and then construct a standard linear regression model of the data in the reduced dimension. As a comparison, we also perform the regression directly without DR. To test the methods with different numbers of data, we perform the experiments with 15, 20, 25, 30, 35, 40 data points randomly selected from the data set and another randomly selected 20 data points used as the test set. In each experiment we can compute the mean relative regression error (MRRE) using the data in the test set. Specifically, suppose $\{(\mathbf{x}_i, y_i)\}_{i=1}^{n_t}$ is the training set and $f_r(\cdot)$ is the

Methods	min	max
w/o DR	.0289	.3374
LDR	.0281	.7913
SIR	.0362	.2534
LSIR	.0380	.2350
BIR	.0247	.1183

Table 2: The minimal and maximal relative regression error (RRE) in the 100 trials with 20 data points for the death rate example. The best results are marked in bold.

regression model, the MRRE is computed as,

$$\text{MRRE} = \frac{1}{n_t} \sum_{i=1}^{n_t} \frac{|y_i - f_r(\mathbf{x}_i)|}{y_i}.$$

We repeat all the experiments 100 times, and compute the mean and the standard deviation of the obtained MRRE, which is shown in Table 1. First we observe that for $n = 40$ all the methods can achieve rather good accuracy; as n decrease, the results of all the other methods become evidently worse, while that of BIR remains quite stable, suggesting that the BIR is especially effective in the small data case. It should be noted that for $n = 15$ LDR and LSIR fail to produce reasonable results due to numerical instability, and so we omit the results here. More importantly it can be seen from the table that starting from $n = 30$, the regression without DR actually has the best performance, suggesting that implementing DR is only necessary when the number of data points is below 30. In all the cases DR is genuinely needed, i.e., $n < 25$, the BIR method performs significantly better than all other methods. To further analyze the performance, we also compute the minimal and the maximal relative regression errors (RRE) for the 20 data-point case, and present the results in Table 2. Once again, we can see that the BIR method has the best results in both the minimal and the maximal cases.

4.4 Automobile data set

Our last example is the automobile data set in the UCI Machine Learning Repository [7]. The original data set contains 205 instances described by 26 attributes including 16 continuous and 10 categorical. We preprocess the data set in the following way: we neglect the 10 categorical attributes, and remove the instances with missing values, yielding a data set with 159 instances and 16 attributes. We select one of the 16 attributes as the response and the others as the predictors: specifically we want to predict the price of an automobile based the other 15 attributes of it. In this problem we first select one feature using the DR methods, and then perform a linear regression with the selected feature.

sample size	20	30	40	50	60	70	80	90	100
w/o DR	.377 (.153)	.254 (.072)	.210 (.035)	.203 (.0393)	.181 (.025)	.173 (.026)	.173 (.022)	.169 (.022)	.170 (.022)
LDR	.394 (.175)	.262 (.097)	.209 (.035)	.198 (.033)	.178 (.028)	.173 (.031)	.175 (.024)	.172 (.027)	.171 (.024)
SIR	.497 (.224)	.284 (.093)	.2358 (.049)	.217 (.044)	.199 (.035)	.192 (.034)	.193 (.032)	.189 (.030)	.187 (.029)
LSIR	.489 (.210)	.284 (.079)	.225 (.043)	.216 (.040)	.194 (.032)	.189 (.034)	.190 (.029)	.183 (.033)	.178 (.027)
BIR	.188 (.034)	.184 (.034)	.178 (.031)	.178 (.029)	.167 (.025)	.164 (.027)	.167 (.021)	.165 (.024)	.162 (.023)

Table 3: The mean and the standard deviation (in parenthesis) of MRRE for Example 4. The best results are marked in bold.

Methods	min	max
w/o DR	.1555	1.273
LDR	.170	1.134
SIR	.169	1.874
LSIR	.217	1.146
BIR	.113	0.287

Table 4: The minimal and maximal relative regression error (RRE) in the 100 trials with 20 data points for the automobile price example. The best results are marked in bold.

Just like the previous example, we want to examine the performance of the DR methods in the small-data setting, i.e., a setting where direct regression can not provide accurate results. To do so, we conduct the experiments with $n = 10, 20, \dots, 90, 100$ randomly selected samples and another 50 random samples used as the test set for all the cases. We repeat each experiment 100 times, and we compute the MRRE each time. The mean and the standard deviation of the MRRE results are reported in Table 3. From the data given in Table 3, we obtain rather similar conclusions as those of Example 3. Namely, the BIR method has the best MRRE of all the four methods used. In Table 4, we show the minimal and the maximal RRE for the 20 data-point case, and just like the results in Example 3, we find that the BIR method has the smallest RRE in both the minimal and the maximal cases.

5 Conclusions

We consider dimension reduction problems for regression and we propose a Bayesian approach for computing the conditional distribution $\pi(\mathbf{x}|y)$ and perform the dimension reduction. The method construct the likelihood function from the data with a GP regression model and MCMC to generate samples from the conditional distribution $\pi(\mathbf{x}|y)$. Numerical examples demonstrate that the proposed method is particularly effective for problems with very small data set. We restate here that, due to the use of GP model, BIR does not apply to problems with very high dimensions. Rather, we expect BIR can be useful for problems with moderately high dimensions (e.g., less than 100), and a very limited amount of data (e.g. hundreds or less). Finally, we also note that, in many practical problems gradient information is available, and DR methods which takes advantages of the gradient information have also been developed, e.g. [10, 2, 13]. In this case, we expect that the gradient information can also be used to enhance the performance of the BIR method, via, for example, Gradient-Enhanced Kriging [25], and we plan to investigate this problem in the future.

References

- [1] Samprit Chatterjee and Ali S Hadi. *Regression analysis by example*. John Wiley & Sons, 2015.
- [2] Paul G Constantine. *Active subspaces: Emerging ideas for dimension reduction in parameter studies*, volume 2. SIAM, 2015.
- [3] R Dennis Cook and Liliana Forzani. Likelihood-based sufficient dimension reduction. *Journal of the American Statistical Association*, 104(485):197–208, 2009.
- [4] R Dennis Cook, Liliana M Forzani, and Diego R Tomassi. Ldr: A package for likelihood-based sufficient dimension reduction. *Journal of Statistical Software*, 39(i03), 2011.

- [5] R Dennis Cook and Liqiang Ni. Sufficient dimension reduction via inverse regression: A minimum discrepancy approach. *Journal of the American Statistical Association*, 100(470):410–428, 2005.
- [6] R Dennis Cook and Sanford Weisberg. Sliced inverse regression for dimension reduction: Comment. *Journal of the American Statistical Association*, 86(414):328–332, 1991.
- [7] Dheeru Dua and Casey Graff. UCI machine learning repository, 2017.
- [8] Louis Ferré. Determining the dimension in sliced inverse regression and related methods. *Journal of the American Statistical Association*, 93(441):132–140, 1998.
- [9] Kenji Fukumizu, Francis R Bach, and Michael I Jordan. Dimensionality reduction for supervised learning with reproducing kernel hilbert spaces. *Journal of Machine Learning Research*, 5(Jan):73–99, 2004.
- [10] Kenji Fukumizu and Chenlei Leng. Gradient-based kernel method for feature extraction and variable selection. In *Advances in Neural Information Processing Systems*, pages 2114–2122, 2012.
- [11] Tailen Hsing and Raymond J Carroll. An asymptotic theory for sliced inverse regression. *The Annals of Statistics*, pages 1040–1061, 1992.
- [12] Minyoung Kim and Vladimir Pavlovic. Dimensionality reduction using covariance operator inverse regression. In *2008 IEEE Conference on Computer Vision and Pattern Recognition*, pages 1–8. IEEE, 2008.
- [13] Rémi Lam, Olivier Zahm, Youssef Marzouk, and Karen Willcox. Multifidelity dimension reduction via active subspaces. *arXiv preprint arXiv:1809.05567*, 2018.
- [14] Bing Li. *Sufficient dimension reduction: Methods and applications with R*. CRC Press, 2018.
- [15] Bing Li, Yuexiao Dong, et al. Dimension reduction for nonelliptically distributed predictors. *The Annals of Statistics*, 37(3):1272–1298, 2009.
- [16] Bing Li and Shaoli Wang. On directional regression for dimension reduction. *Journal of the American Statistical Association*, 102(479):997–1008, 2007.
- [17] Ker-Chau Li. Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association*, 86(414):316–327, 1991.
- [18] Lexin Li and Xiangrong Yin. Sliced inverse regression with regularizations. *Biometrics*, 64(1):124–131, 2008.
- [19] Qian Lin, Zhigen Zhao, and Jun S Liu. Sparse sliced inverse regression via lasso. *Journal of the American Statistical Association*, pages 1–33, 2019.

- [20] Yanyuan Ma and Liping Zhu. A semiparametric approach to dimension reduction. *Journal of the American Statistical Association*, 107(497):168–179, 2012.
- [21] Yanyuan Ma and Liping Zhu. A review on dimension reduction. *International Statistical Review*, 81(1):134–150, 2013.
- [22] Kai Mao, Feng Liang, and Sayan Mukherjee. Supervised dimension reduction using bayesian mixture modeling. In *Proceedings of the Thirteenth International Conference on Artificial Intelligence and Statistics*, pages 501–508, 2010.
- [23] Gary C McDonald and Richard C Schwing. Instabilities of regression estimates relating air pollution to mortality. *Technometrics*, 15(3):463–481, 1973.
- [24] Geoffrey McLachlan and David Peel. *Finite Mixture Models*. John Wiley & Sons, 2004.
- [25] Max D Morris, Toby J Mitchell, and Donald Ylvisaker. Bayesian design and analysis of computer experiments: use of derivatives in surface prediction. *Technometrics*, 35(3):243–255, 1993.
- [26] Brian J Reich, Howard D Bondell, and Lexin Li. Sufficient dimension reduction via bayesian mixture modeling. *Biometrics*, 67(3):886–895, 2011.
- [27] Kean Ming Tan, Zhaoran Wang, Tong Zhang, Han Liu, and R Dennis Cook. A convex formulation for high-dimensional sparse sliced inverse regression. *Biometrika*, 105(4):769–782, 2018.
- [28] Surya T Tokdar, Yu M Zhu, Jayanta K Ghosh, et al. Bayesian density regression with logistic gaussian process and subspace projection. *Bayesian analysis*, 5(2):319–344, 2010.
- [29] Christopher KI Williams and Carl Edward Rasmussen. *Gaussian processes for machine learning*. MIT Press, 2006.
- [30] Qiang Wu, Sayan Mukherjee, and Feng Liang. Localized sliced inverse regression. In *Advances in neural information processing systems*, pages 1785–1792, 2009.
- [31] Yingcun Xia, Howell Tong, Wai Keungxs Li, and Li-Xing Zhu. An adaptive estimation of dimension reduction space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64(3):363–410, 2002.
- [32] Li-Xing Zhu, Kai-Tai Fang, et al. Asymptotics for kernel estimate of sliced inverse regression. *The Annals of Statistics*, 24(3):1053–1068, 1996.
- [33] Li-Xing Zhu and Kai W Ng. Asymptotics of sliced inverse regression. *Statistica Sinica*, pages 727–736, 1995.