

On recursively defined combinatorial classes and labelled trees

Ali Chouria*, Vlad-Florin Drăgoi[†] and Jean-Gabriel Luque[‡]

April 14, 2020

Abstract

We define and prove isomorphisms between three combinatorial classes involving labeled trees. We also give an alternative proof by means of generating functions.

Keywords: Combinatorial classes, Catalan numbers, labeled trees, rooted trees, generating functions.

1 Introduction

Based on the theory of combinatorial species (see *e.g.* [1]), Flajolet and Sedgewick [4] wrote a reference book on combinatorial analysis. In particular in the first part, they provided a list of basic constructions for exponential generating functions. Mainly, complex combinatorial structures are obtained by combining the following three combinatorial classes: SET, SEQ, and CYC. For instance, they described *surjections* (SEQ(SET)), *set partitions* (SET(SET)), *alignments* (SEQ(CYC)) and *permutations* (SET(CYC)). In this article, we focus on the later one because it has the remarkable property of having the same generating function as the combinatorial class of sequences. More precisely, our starting point consists in giving an explicit bijection between the class of set-of-cycles and the class of sequences (see Section 3.4). Our goal is to study an example of a class defined inductively

*ali.chouria1@univ-rouen.fr. Laboratoire LITIS - EA 4108, Université de Rouen-Normandie, Avenue de l'Université - BP 8, 76801 Saint-Étienne-du-Rouvray, ISLAIB, University of Jendouba, Tunisia

[†]vlad-florin.dragoi@univ-rouen.fr. Department of Mathematics and Computer Sciences "Aurel Vlaicu" University of Arad, Romania

[‡]jean-gabriel.luque@univ-rouen.fr. Laboratoire LITIS - EA 4108, Université de Rouen-Normandie, Avenue de l'Université - BP 8, 76801 Saint-Étienne-du-Rouvray Cedex

by a combinatorial class equation. We chose the equation $\text{SET}(\text{CYC}(\mathcal{C})) = \mathcal{C}$ because the underlying combinatorics reveal a world rich in interpretation and provide fruitful perspectives. In particular, this equation reveals an isomorphism between sets of necklaces of planar labelled trees, forests of labelled trees, and rooted labelled trees (see Section 4). At the heart of the equation we study here are the Catalan numbers. They are involved in the enumeration of numerous classes of combinatorial objects of prime importance in computer science, *e.g.* Dyck paths, binary trees, non-crossing partitions *etc.* [15, 5, 4]. Notice that more than 60 possible enumerations are listed in [15] and the enumerated objects lead to several applications in computer science, such as sorting techniques based on binary trees [6]. In our case, Catalan numbers are representing ordered trees. The article ends with Section 4.4, where we propose other recursive tree-like combinatorial classes.

2 Background and notations

We recall here well known definitions and results concerning combinatorial classes and generating functions. The material contained in this section mainly refers to [4].

2.1 Combinatorial classes

In the most general context, a *combinatorial class* is a triplet $(\mathcal{O}, \mathcal{P}, \omega)$ where \mathcal{O} is the discrete set of the combinatorial objects we want to enumerate, \mathcal{P} is the discrete set of the properties in regard to which you want to enumerate our objects, and $\omega : \mathcal{O} \rightarrow \mathcal{P}$ is a map such that for every $p \in \mathcal{P}$ the preimage $\omega^{-1}(p)$ is finite, which is a minimal requirement in order to be able to enumerate the objects of \mathcal{O} with respect to the properties of \mathcal{P} .

We consider the restricted context where $\mathcal{P} = \mathbb{N}$ and the preimage of 0 contains only one element. More formally, a combinatorial class is a pair $\mathcal{C} = (\mathcal{O}, \omega)$ where $\omega : \mathcal{O} \rightarrow \mathbb{N}$ is such that $\text{card}(\omega^{-1}(n)) < \infty$ for any integer n . For the sake of simplicity, and when there is no ambiguity, we use the same name for a class and the set of its objects. Then, we denote by $|\mu|$ the *degree* (or *weight*) $\omega(\mu)$ of $\mu \in \mathcal{C}$ and we set $\mathcal{C}_n = \{\mu \in \mathcal{C} \mid \omega(\mu) = n\}$. If $\#(\omega^{-1}(0)) = 1$ then we denote by ϵ the unique element of \mathcal{C} of weight 0. We also set $\mathcal{C}^+ := \mathcal{C} \setminus \{\epsilon\}$.

2.2 Labelled combinatorial classes

Recall that the symmetric group \mathcal{S}_n is the group of bijections of $\{1, \dots, n\}$. It is a group of order $n!$ whose each element is denoted by the word of its images. For instance, the cycle sending 1 to 2, 2 to 3, and 3 to 1 is denoted by 231. Obviously, the set of permutations is closed by composition and all its elements are invertible.

Formally, a *labelled combinatorial class* is a combinatorial class endowed with a sequence $(\rho_n)_{n \in \mathbb{N}}$ of representations ρ_n of the symmetric group \mathcal{S}_n (*i.e.* an application associating a map $\rho_n(\sigma) : \mathcal{C}_n \rightarrow \mathcal{C}_n$ to each permutation $\sigma \in \mathcal{S}_n$ in such a way that $\rho_n(\sigma \circ \sigma') = \rho_n(\sigma) \circ \rho_n(\sigma')$). An equivalent way (see *e.g.* [4]) to define labelled combinatorial class

consists in considering that each element of \mathcal{C}_n is a graph whose vertices are labelled by numbers from 1 to n ; the image of a permutation by the underlying representation is just the permutation of the labels.

Let (\mathcal{C}, ω) and (\mathcal{C}', ω') be two labeled combinatorial class. If the sets \mathcal{C} and \mathcal{C}' are disjoint, then we define the class $\mathcal{C} \boxplus \mathcal{C}' = (\mathcal{C} \cup \mathcal{C}', \omega'')$ with $\omega''(e) = \omega(e)$ if $e \in \mathcal{C}$ and $\omega''(e) = \omega'(e)$ if $e \in \mathcal{C}'$. One extends to the case where $\mathcal{C} \cap \mathcal{C}' \neq \emptyset$ by replacing \mathcal{C}' by a copy which is disjoint of \mathcal{C} in the definition of $\mathcal{C} \boxplus \mathcal{C}'$.

We also define $\mathcal{C} \boxtimes \mathcal{C}'$, i.e., the combinatorial class such that the elements of $(\mathcal{C} \boxtimes \mathcal{C}')_n$ are the pairs (e, e') where e is obtained by relabeling an element of \mathcal{C}_i and e' is obtained by relabeling an element of \mathcal{C}'_j , with $i + j = n$ such that the set of the labels in (e, e') is $\{1, \dots, n\}$ and each relabeling preserves the initial order on the vertices. The degree of (e, e') in $\mathcal{C} \boxtimes \mathcal{C}'$ is the sum of the degree of the respective preimage of e and e' in \mathcal{C} and \mathcal{C}' . As a special case, for each labeled class \mathcal{C} , we denote $\mathcal{C}^\bullet = \bullet \boxtimes \mathcal{C}$, where \bullet is the class of the unique element of which, denoted also by \bullet , has degree 1.

2.3 The exponential generating function of a combinatorial class

The exponential generating function (EGF) of a combinatorial class \mathcal{C} is the exponential generating function of the numbers $C_n = \#(\mathcal{C}_n)$, in other words

$$S_{\mathcal{C}}(x) = \sum_{n \geq 0} C_n \frac{x^n}{n!} = \sum_{\mu \in \mathcal{C}} \frac{x^{|\mu|}}{|\mu|!}.$$

We say that two classes are *isomorphic* if their EGF are equal

$$\mathcal{C} \equiv \mathcal{C}' \Leftrightarrow S_{\mathcal{C}} = S_{\mathcal{C}'} \Leftrightarrow \forall n \in \mathbb{N}, C_n = C'_n.$$

Classically, we have [4]

$$S_{\mathcal{C} \boxplus \mathcal{C}'} = S_{\mathcal{C}} + S_{\mathcal{C}'} \text{ and } S_{\mathcal{C} \boxtimes \mathcal{C}'} = S_{\mathcal{C}} S_{\mathcal{C}'}.$$

2.4 Labelled sequences

If \mathcal{C}^+ is a labelled combinatorial class such that $\mathcal{C}_0^+ = \emptyset$ then we define, up to an equivalence, the class $\text{SEQ}(\mathcal{C}^+)$ of labelled sequences by the equation

$$\text{SEQ}(\mathcal{C}^+) \equiv [] \boxplus (\mathcal{C}^+ \boxtimes \text{SEQ}(\mathcal{C}^+)),$$

where $[]$ denotes the class having a single element ϵ which is degree 0.

It is easy to show that such a class exists and that its associated exponential generating function is

$$S_{\text{SEQ}(\mathcal{C}^+)}(x) = \frac{1}{1 - S_{\mathcal{C}^+}(x)}.$$

From a combinatorial point of view, the elements of $\text{SEQ}(\mathcal{C}^+)_n$ are k -tuple $[\mu_1, \dots, \mu_k]$ where each μ_i is obtained by an order preserving relabelling of an element of $\mathcal{C}_{j_i}^+$, in such

a way that the whole set of labels in $[\mu_1, \dots, \mu_k]$ is $\{1, \dots, n\}$ (as a consequence one has $\sum_{i=1}^k j_i = n$). So to any element $s = [\mu_1, \dots, \mu_k] \in \text{SEQ}(\mathcal{C}^+)_n$ we associate an ordered partition $\Pi = [\Pi_1, \dots, \Pi_k]$ of size n such that each Π_i is the set of the labels of μ_i .

Let us be a little more precise. A *labelled* list of elements of \mathcal{C}^+ is a list $L = [\mu_1, \mu_2, \dots, \mu_k] \in \text{SEQ}(\mathcal{C}^+)$ with each μ_i associated to a set Ω_i of non-negative integers with $\text{card}(\Omega_i) = |\mu_i|$ and $\Omega_i \cap \Omega_j = \emptyset$ for all i, j . A *standard* labelled list of elements of \mathcal{C}^+ is a labelled list L of elements of \mathcal{C}^+ such that the set of all the labels of L is $\{1, \dots, |L|\}$.

2.5 Labelled sets and labelled cycles

We consider the labelled combinatorial class $\text{SET}(\mathcal{C}^+)$ such that $\text{SET}(\mathcal{C}^+)_n$ is the quotient of the set $\text{SEQ}(\mathcal{C}^+)_n$ by the relation $[\mu_1, \dots, \mu_k] \equiv_S [\mu_{\sigma(1)}, \dots, \mu_{\sigma(k)}]$ for any permutation $\sigma \in \mathcal{S}_k$. Straightforwardly, each element of $\text{SET}(\mathcal{C}^+)$ can be represented by a set of (order preserved) relabelled elements of \mathcal{C}^+ . The exponential generating function

$$S_{\text{SET}(\mathcal{C}^+)}(x) = \exp\{S_{\mathcal{C}^+}(x)\}.$$

is easily deduced from the construction.

If one consider the equivalence relation generated by $[\mu_1, \dots, \mu_k] \equiv_C [\mu_2, \dots, \mu_k, \mu_1]$, the one obtains an other labelled combinatorial class $\text{CYC}(\mathcal{C}^+)$ whose elements can be represented by necklace of (order preserved) relabelled elements of \mathcal{C}^+ . We denote a necklace by $(\mu_1, \dots, \mu_k) = (\mu_2, \dots, \mu_k, \mu_1)$. Again, the generating series

$$S_{\text{CYC}(\mathcal{C}^+)}(x) = \log \left\{ \frac{1}{1 - S_{\mathcal{C}^+}(x)} \right\}.$$

is deduced from the construction.

3 Set partitions and related constructions

3.1 Three constructions based on set partitions

A *set partition* of size n is a set $\pi = \{\pi_1, \dots, \pi_k\}$ such that $\pi_1 \cup \dots \cup \pi_k = \{1, \dots, n\}$ and $\pi_i \cap \pi_j = \emptyset$ for any two indices $1 \leq i \neq j \leq k$. The set of set partitions \mathcal{S}_p endowed with the size is a combinatorial classes satisfying $\mathcal{S}_p \equiv \text{SET}(\text{SET}(\bullet)^+)$, and so, $S_{\mathcal{S}_p}(x) = \exp(\exp(x) - 1)$. The numbers $\text{Sp}_n = 1, 1, 2, 5, 15, 52, 203, 877, 4140 \dots$ are the well known Bell numbers (see sequence A000110 in [16]).

If \mathcal{C}^+ is a labeled combinatorial class such that $\mathcal{C}_0^+ = 0$ and $\mathcal{P} \equiv \text{SET}(\mathcal{C}^+)$ then the definitions above allows to associate to each element $p = \{p_1, \dots, p_k\} \in \mathcal{P}_n$ a set partitions $\pi(p) = \{\text{labels}(p_1), \dots, \text{labels}(p_k)\}$ of size n where for each $1 \leq i \leq k$, $\text{labels}(p_i)$ denotes the set of the labels of p_i .

An *ordered partition* of size n is a sequence $\Pi = [\Pi_1, \dots, \Pi_k]$ of non empty sets such that $\{\Pi_1, \dots, \Pi_k\}$ is a set partition of $\{1, \dots, n\}$. The set of ordered partitions \mathcal{O}_p endowed with the size is a combinatorial classes satisfying $\mathcal{O}_p \equiv \text{SEQ}(\text{SET}(\bullet)^+)$ and $S_{\mathcal{O}_p}(x) = \frac{1}{2 - \exp(x)}$.

The numbers $\mathcal{O}p_n = 1, 1, 3, 13, 75, 541, 4683, 47293, 545835, \dots$ are the Fubini numbers (see sequence A000670 in [16]). If $\mathcal{L} \equiv \text{SEQ}(\mathcal{C}^+)$ then the definitions above allows to associate to each element $\ell = [\ell_1, \dots, \ell_k] \in \mathcal{L}_n$ a set partition $\Pi(\ell) = [\text{labels}(\ell_1), \dots, \text{labels}(\ell_k)]$ of size n .

A *cyclic partition* of size n is a necklace $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_k)$ such that $\{\mathbf{p}_1, \dots, \mathbf{p}_k\}$ is a set partition of size n . The set of ordered partitions $\mathcal{C}p$ endowed with the size is a combinatorial classes satisfying $\mathcal{C}p \equiv \text{CYC}(\text{SET}(\bullet)^+)$ and $S_{\mathcal{C}p}(x) = \log\left(\frac{1}{2-\exp(x)}\right)$.

The numbers $\mathcal{C}p_n = 1, 1, 3, 13, 75, 541, 4683, 47293, 545835, \dots$ are listed in sequence A000670 [16]. If $\mathcal{N}e \equiv \text{CYC}(\mathcal{C}^+)$ then the above definitions allow us to associate to each element $c = (c_1, \dots, c_k) \in (\mathcal{N}e)_n$ a cyclic partition $\mathbf{p}(c) = (\text{labels}(c_1), \dots, \text{labels}(c_k))$ of size n .

3.2 An explicit isomorphism

From the generating series we have $\text{SET}(\mathcal{C}p) \equiv \mathcal{O}p$. Indeed, this equality translates in terms of generating function as $\exp\left(\log\left(\frac{1}{2-e^x}\right)\right) = \frac{1}{2-e^x}$. In order to understand a more general identity introduced later in the paper, we make explicit this bijection. Assume that $c = \{c^{(1)}, \dots, c^{(k)}\} \in \text{SET}(\mathcal{C}p)$. If $c^{(i)} = (c_1^{(i)}, \dots, c_{h_i}^{(i)})$ then we consider σ_i the only circular permutation on the indices $\{1, \dots, h_i\}$ such that $\min \bigcup_j \text{labels}(c_{(j)}^{(i)}) = \min \text{labels}(c_{\sigma_i^{-1}(1)}^{(i)})$. In other words, if $\ell_i = [\ell_1^{(i)}, \dots, \ell_{h_i}^{(i)}] = [c_{\sigma_i^{-1}(1)}^{(i)}, \dots, c_{\sigma_i^{-1}(h_i)}^{(i)}]$ then $\min \bigcup_j \text{labels}(\ell_{(j)}^{(i)}) = \min \text{labels}(\ell_1^{(i)})$. Now, consider the unique permutation $\rho \in \mathcal{S}_k$ such that

$$\min \text{labels}(c^{(\rho^{-1}(1))}) > \min \text{labels}(c^{(\rho^{-1}(2))}) > \dots > \min \text{labels}(c^{(\rho^{-1}(k))})$$

and set

$$\text{stol}(c) = [\ell_1^{(\rho(1))}, \dots, \ell_{h_{\rho(1)}}^{(\rho(1))}, \dots, \ell_k^{(\rho(k))}, \dots, \ell_{h_{\rho(k)}}^{(\rho(k))}] \in \mathcal{O}p. \quad (1)$$

For instance,

$$\begin{aligned} \text{stol}(\{(\{11\}, \{2, 5\}, \{10\}), (\{6\}, \{1, 3, 4\}, \{7, 9\}), (\{8, 12\})\}) = \\ \{(\{8, 12\}, \{2, 5\}, \{10\}, \{11\}, \{1, 3, 4\}, \{7, 9\}, \{6\})\}. \end{aligned}$$

Let $\ell = [\ell_1, \dots, \ell_k] \in \mathcal{O}p$ and $1 = i_0 \leq \dots \leq i_{h-1} < i_h = k + 1 \in \{1, \dots, k + 1\}$ be the set of indices satisfying

$$\min \bigcup_{i < i_{j+1}} \text{labels}(\ell_i) > \min \bigcup_{i \geq i_j} \text{labels}(\ell_i) \quad (2)$$

with h maximal.

For instance, the indices associated to $\{(\{8, 12\}, \{2, 5\}, \{10\}, \{11\}, \{1, 3, 4\}, \{7, 9\}, \{6\})\}$ are $1 \leq 2 \leq 5 < 8$. We define

$$\text{ltos}(\ell) = \{c_1, \dots, c_k\} \in \text{SET}(\mathcal{C}p), \quad (3)$$

where c_j denotes the necklace $(\ell_{i_{j-1}}, \dots, \ell_{i_j-1})$. For instance,

$$\begin{aligned} \mathbf{ltos}(\{\{8, 12\}, \{2, 5\}, \{10\}, \{11\}, \{1, 3, 4\}, \{7, 9\}, \{6\}\}) = \\ \{(\{1, 3, 4\}, \{7, 9\}, \{6\}), (\{2, 5\}, \{10\}, \{11\}), (\{8, 12\})\}. \end{aligned}$$

It is easy to check that $\mathbf{ltos}(\mathbf{stol}(c)) = c$ and $\mathbf{stol}(\mathbf{ltos}(\ell)) = \ell$. So we have

Proposition 1 *The map \mathbf{stol} is an isomorphism of combinatorial classes and \mathbf{ltos} is its reverse map.*

3.3 About Lyndon words

Recall that the free monoid (e.g. [9]) Σ^* on a set Σ is the monoid whose elements are all the finite sequences endowed with the catenation product \cdot that consists of pasting one sequence to the right of another. The empty sequence plays the role of the identity element. For instance, in the free monoid $\{a, b\}^*$ we have $[a, b, a, a, b] \cdot [b, a, a] = [a, b, a, a, b, b, a, a]$. In literature, brackets and commas are often omitted; the elements of a free monoid are then noted as juxtapositions of letters called words (the empty word, noted by ε , corresponds to the sequence $[\]$). The name of the free monoid comes from the fact that it fulfills the universal property, that is every monoid having a generating set in bijection with Σ is isomorphic to a quotient of Σ^* .

Any pair of sequences under the form $u \cdot v$ and $v \cdot u$ are said conjugate. In other words, the conjugates of a sequence are all its circular shift. This is obviously an equivalence relation that preserves the periods, i.e., the conjugate sequences of $u \cdot^k$ are exactly the sequences $v \cdot^k$ where v is conjugate to u . In terms of combinatorial class the free monoid is nothing but $\text{SEQ}(\Sigma)$ and its quotient by conjugation is $\text{CYC}(\Sigma)$.

Assume that the alphabet Σ is totally ordered by the order $<$. Then the free monoid is totally ordered with the lexicographic order \prec . The minimal element for the lexicographic order is the empty sequence $[\]$ and we have $[a] \cdot u \prec [b] \cdot v$ if $a < b$ or $a = b$ and $u \prec v$.

A *Lyndon words* a non periodic sequence which is minimal in its conjugacy class. Their name comes from the mathematician Roger Lyndon who studied them in 1954[10]. Nevertheless, it should be noted that they had been introduced a year earlier by Anatoly Shirshov [14]. Lyndon words play a very important role for understanding of free groups [2], free associative algebras, and free Lie algebras [13]. Readers may refer to [11] for a rather complete survey.

Among all the properties of Lyndon's words, one of the most interesting is that they play for the free monoid the same role as prime numbers play for integers. This property is that any sequence factorizes as a unique weakly decreasing catenation of Lyndon words [12]. In other words, the free monoid Σ^* is in bijection with the multisets of aperiodic sequences over σ . For instance, if we assume $a < b$ the sequence $u = [a, b, a, b, b, a, b, a, b, a, a, a, b, a, b, a]$ factorizes as $u = [a, b, a, b, b] \cdot [a, b] \cdot [a, b] \cdot [a, a, a, b, a, b] \cdot [a]$. This means that the sequence u is assimilated to the multiset $\{(a, a, a, b, a, b), (a, b, a, b, b), (a, b), (a, b), (a)\}$ (remark the multiplicity of (a, b)). It is interesting to note that this correspondence is precisely the one that is calculated when applying \mathbf{ltos} . Indeed, let $\ell = [\ell_1, \dots, \ell_k] \in \mathcal{Op}$, the alphabet $\Sigma = \{\ell_1, \dots, \ell_k\}$ is totally ordered by $\ell_i < \ell_j$ if and only if $\min \ell_i < \min \ell_j$. In fact, since each numbers of $\{1, \dots, n\}$ appears only

one time in the sequence, only the minimal elements the sets are relevant and all works as if our alphabet be $\{1, \dots, n\}$. For instance, $\{\{8, 12\}, \{2, 5\}, \{10\}, \{11\}, \{1, 3, 4\}, \{7, 9\}, \{6\}\}$ is assimilated to $[8, 2, 10, 11, 1, 7, 6]$. The indices of equation (2), except the larger which is not relevant, indicate where to catenate in order to apply the complete factorization. In our example, we found the indices $\{1, 2, 5, 8\}$ and, then, we have $[8, 2, 10, 11, 1, 7, 6] = [8] \cdot [2, 10, 11] \cdot [1, 7, 6]$. Notice that, since the components are two by two distinct, the bijection with multisets of cycles class sends the sequences we consider on set of necklaces. For instance, $[8] \cdot [2, 10, 11] \cdot [1, 7, 6] \sim \{(1, 7, 6), (2, 10, 11), (8)\}$. We recover the $\mathbf{ltos}(\ell)$ by replacing each integer by the set of which it is the minimum. In our example we have

$$\{(1, 7, 6), (2, 10, 11), (8)\} \rightarrow \{(\{1, 3, 4\}, \{7, 9\}, \{6\}), (\{2, 5\}, \{10\}, \{11\}), (\{8, 12\})\}.$$

Of course, this may seem like a very sophisticated way to revisit the bijection of the previous section. Nevertheless, this remark is valuable because it will allow us to link our constructions to notions of algebras (enveloping algebras, Hopf algebras, Lie algebras of primitive elements etc.) that we will explore in future works.

3.4 Set of cycles and sequences

Let \mathcal{C}^+ be a labeled combinatorial sequences such that $\mathcal{C}_0^+ = \emptyset$. We define $\mathcal{J} = \text{SET}(\text{CYC}(\mathcal{C}^+))$ and $\mathcal{S} = \text{SEQ}(\mathcal{C}^+)$.

Let us show that the map \mathbf{stol} allows us to compute an explicit isomorphism from \mathcal{J} to \mathcal{S} . We define $\mathbf{jtoset}_{\mathcal{C}^+} : \mathcal{J} \rightarrow \text{SET}(\mathcal{C}p)$ by

$$\mathbf{jtoset}_{\mathcal{C}^+}(\{(c_1^{(1)}, \dots, c_{h_1}^{(1)}), \dots, (c_1^{(k)}, \dots, c_{h_k}^{(k)})\}) = \{(\mathbf{labels}(c_1^{(1)}), \dots, \mathbf{labels}(c_{h_1}^{(1)})), \dots, (\mathbf{labels}(c_1^{(k)}), \dots, \mathbf{labels}(c_{h_k}^{(k)}))\}.$$

We define also $\mathbf{jtoseq}_{\mathcal{C}^+} : \mathcal{J} \rightarrow \mathcal{S}$ such that

$$\mathbf{jtoseq}_{\mathcal{C}^+}(\{(c_1^{(1)}, \dots, c_{h_1}^{(1)}), \dots, (c_1^{(k)}, \dots, c_{h_k}^{(k)})\}) = \ell$$

is the unique permutation of the vector $[c_1^{(1)}, \dots, c_{h_1}^{(1)}, \dots, c_1^{(k)}, \dots, c_{h_k}^{(k)}]$ such that $\Pi(\ell) = \mathbf{stol}(\mathbf{jtoset}_{\mathcal{C}^+}(\{(c_1^{(1)}, \dots, c_{h_1}^{(1)}), \dots, (c_1^{(k)}, \dots, c_{h_k}^{(k)})\}))$.

Let $c = \{(c_1^{(1)}, c_2^{(1)}), (c_1^{(2)}), (c_1^{(3)}, c_2^{(3)})\}$ be a set of labelled graphs as shown in Fig. 1
Then $\mathbf{jtoset}_{\mathcal{C}^+}(c)$ equals

$$\{(\{3, 7, 5, 20\}, \{9, 14, 17, 19, 11\}), (\{1, 18\}), (\{2, 8, 13\}, \{4, 16, 12, 15, 10, 6\})\}.$$

Furthermore, $\Pi(\ell) = \mathbf{stol}(\mathbf{jtoset}_{\mathcal{C}^+}(c)) =$

$$[\{3, 7, 5, 20\}, \{9, 14, 17, 19, 11\}, \{2, 8, 13\}, \{4, 16, 12, 15, 10, 6\}, \{1, 18\}].$$

Since \mathbf{stol} is one to one, the equality on generating functions allows us to deduce that $\mathbf{jtoseq}_{\mathcal{C}^+}$ is an isomorphism of combinatorial classes. The inverse bijection $\mathbf{seqtoj}_{\mathcal{C}^+} : \text{SET}(\mathcal{C}p) \rightarrow \mathcal{J}$ is defined by

$$\mathbf{seqtoj}_{\mathcal{C}^+}(\ell) = \{(c_1^{(1)}, \dots, c_{h_1}^{(1)}), \dots, (c_1^{(k)}, \dots, c_{h_k}^{(k)})\}$$




Figure 1: Five labelled graphs

where $[c_1^{(1)}, \dots, c_{h_1}^{(1)}, \dots, c_1^{(k)}, \dots, c_{h_k}^{(k)}]$ is the unique permutation of ℓ such that $\text{ltos}(\Pi(\ell)) = \text{jtoset}_{\mathcal{C}^+}(\{(c_1^{(1)}, \dots, c_{h_1}^{(1)}), \dots, (c_1^{(k)}, \dots, c_{h_k}^{(k)})\})$.

In the aforementioned example, we deduce the indices of the minimum elements in $\Pi(\ell)$ being $1 < 3 < 5$. Hence,

$$\begin{aligned} \text{ltos}(\Pi(\ell)) &= \{(\{1, 18\}), (\{2, 8, 13\}, \{4, 16, 12, 15, 10, 6\}), (\{3, 7, 5, 20\}, \{9, 14, 17, 19, 11\})\} \\ &= \text{jtoset}_{\mathcal{C}^+}(c). \end{aligned}$$

We summarize the results of this section in the following theorem

Theorem 1 *The maps which make commuting the following diagram are explicit isomorphisms of combinatorial classes*

$$\text{SET}(\text{CYC}(\mathcal{C}^+)) \begin{array}{c} \xrightarrow{\text{jtoset}_{\mathcal{C}^+}} \\ \xleftarrow{\text{seqto}_{\mathcal{C}^+}} \end{array} \text{SEQ}(\mathcal{C}^+). \quad (4)$$

4 Labelled and unlabelled trees

We illustrate the previous result by investigating the combinatorial classes \mathcal{R} satisfying

$$\text{SET}(\text{CYC}(\mathcal{R}^\bullet)) \equiv \text{SEQ}(\mathcal{R}^\bullet) \equiv \mathcal{R}. \quad (5)$$

This isomorphism can be translated into the following functional equation:

$$\exp \left\{ \log \left\{ \frac{1}{1 - xS_{\mathcal{R}}(x)} \right\} \right\} = \frac{1}{1 - xS_{\mathcal{R}}(x)} = S_{\mathcal{R}}(x). \quad (6)$$

This equation has a unique solution

$$S_{\mathcal{R}}(x) = \frac{1 - \sqrt{1 - 4x}}{2x} \quad (7)$$

which is also the ordinary generating function of the Catalan numbers $\mathbf{C}_n = \frac{1}{n+1} \binom{2n}{n}$ [15, 4]. Hence, \mathcal{R} is unique up to an isomorphism and

$$R_n = \frac{(2n)!}{(n+1)!}. \quad (8)$$

The sequence of R_n is

$$1, 1, 4, 30, 336, 5040, 95040, 2162160 \quad A001761[16].$$

4.1 Labelled trees from unlabelled trees

Definition 1 *A tree is a list of trees (possibly empty) connected to a node, called its root, by an edge (also called branch). Notice that this is a valid recursive definition which base case is a root together with an empty list. The degree $\omega(t)$ of a tree t is the number of its edges or, equivalently the number of its nodes which are not its root.*

Let \mathcal{D} be the set of trees. There are a finite number of trees having a given degree, so the pair (\mathcal{D}, ω) is a (unlabelled) combinatorial class. The number D_n is known to be the Catalan number C_n (see *e.g.* [15]). So the ordinary generating function of the class \mathcal{D} , *i.e.*,

$$S_{\mathcal{D}}^{ord}(x) = \sum_{n \geq 0} D_n x^n, \quad (9)$$

fulfills the same functional equation (6) as the exponential generating function of \mathcal{R} . So each \mathcal{R}_n is in one to one correspondence with $\mathcal{D}_n \times \mathcal{S}_n$. This suggests that one can exhibit an explicit realization of the class \mathcal{R} by labeling the nodes which are not the root of each tree $t \in \mathcal{D}$ by $\{1, \dots, \omega(t)\}$, without repetition and in any possible way.

4.2 Shifted structure

The class \mathcal{R}^\bullet is isomorphic to the class \mathcal{R}_r of trees with labeled root endowed with the weight ω_r counting the total number of nodes, including the root. More precisely, for a given n , a tree of $(\mathcal{R}_r)_{n+1}$ is obtained by labelling the root of a tree in \mathcal{R}_n with any of the possible value from the set $\{1, \dots, n+1\}$ and relabel, if necessary, the nodes with respect to the order induced by the initial permutation.

Example 1 *Let t be a rooted labelled tree in \mathcal{R}_5 , and its associated permutation is $\pi = (1, 2, 4, 5, 3)$. Then the set of all rooted labeled trees where the root is labeled obtained from t is given by the set of permutations $\{(6, 1, 2, 4, 5, 3), (5, 1, 2, 4, 6, 3), (4, 1, 2, 5, 6, 3), (3, 1, 2, 5, 6, 4), (2, 1, 3, 5, 6, 4), (1, 2, 3, 5, 6, 4)\}$.*

In terms of generating function, this operation leads to

$$S_{\mathcal{R}_r}(x) = S_{\mathcal{R}^\bullet}(x) = xS_{\mathcal{R}}(x) = \frac{1 - \sqrt{1 - 4x}}{2}$$

and so

$$(R_r)_n = n!C_{n-1} = \frac{(2n-2)!}{(n-1)!},$$

for any $n \geq 1$. The sequence of R_{r_n} is given by

$$1, 2, 12, 120, 1680, 30240 \quad A001813 [16].$$

We insists on the fact that at this point $(\mathcal{R}_r)_n$ counts trees having n nodes including the root.

4.3 Hanging trees in necklaces

A *labelled necklace of planar trees* is a necklaces on which trees are hung and all the nodes (comprising the roots) are labeled by $\{1, \dots, n\}$ where n is the total numbers of nodes (comprising roots). We denote by \mathcal{N} the set of such necklaces. The cyclic structure comes from the fact that a necklace is invariant by rotation. The weight $\omega_N(\mathbf{n})$ of a necklace \mathbf{n} is the total number of the nodes, comprising roots, of the trees it contains. The pair (\mathcal{N}, ω_N) is a labeled combinatorial class that satisfies

$$\mathcal{N} \equiv \text{CYC}(\mathcal{R}^\bullet). \quad (10)$$

We depict in Fig. 2 elements of \mathcal{N} that are equivalent under cyclic rotation.




Figure 2: Three rooted labeled trees equivalent under cyclic rotation

Notices that Labeled necklaces of rooted trees appear under the name of "planar labelled trees" in the work of Miloudi [8]. To be more precise, he studied combinatorial class which is straightforwardly isomorphic to \mathcal{N} and he proved $N_n = (2n - 3)!/(n - 1)!$ for any $n \geq 2$ and $N_1 = 1$. We recover this result from the interpretation of (10) in terms of generating function. Indeed,

$$S_{\text{CYC}(\mathcal{R}^\bullet)}(x) = \log \frac{1}{1 - \frac{1 - \sqrt{1 - 4x}}{2}} = \log \frac{2}{1 + \sqrt{1 - 4x}} = \log \frac{1 - \sqrt{1 - 4x}}{2x}.$$

Hence,

$$S_{\text{CYC}(\mathcal{R}^\bullet)}(x) = \log \mathcal{R}(x), \quad (11)$$

and the exact formula for N_n is obtained by expanding the function as a Taylor series. These numbers are also mentioned by Wolfdieter Lang in [16], see the sequence below

1, 1, 3, 20, 210, 3024, 55440, 1235520, 32432400 A006963.

We now have all the material to make explicit the isomorphisms suggested by (5). To this aim we consider jewellery boxes which are sets of necklaces and forests which

are sequences of trees. More formally, in terms of combinatorial classes we define $\mathcal{J} = \text{SET}(\mathcal{N})$ and $\mathcal{F} = \text{SEQ}(\mathcal{R}^\bullet)$. Let $\text{jtof} = \text{jtoseq}_{\mathcal{R}^\bullet} : \mathcal{J} \rightarrow \mathcal{F}$ and its inverse bijection $\text{ftoj} = \text{seqtoj}_{\mathcal{R}^\bullet} : \mathcal{F} \rightarrow \mathcal{J}$. An explicit isomorphism $\text{rtof} : \mathcal{R} \rightarrow \mathcal{F}$ is obtained by removing the root to any tree in \mathcal{R} . The reciprocal isomorphism $\text{ftor} : \mathcal{F} \rightarrow \mathcal{R}$ consists in connecting all the trees of a given sequence to an additional node called the root. All these constructions are summarized in the following result which is a corollary of Theorem 1.

Corollary 1 *The maps which make commuting the following diagram are explicit isomorphisms of combinatorial classes*

$$\mathcal{J} \begin{array}{c} \xrightarrow{\text{jtof}} \\ \xleftarrow{\text{ftoj}} \end{array} \mathcal{F} \begin{array}{c} \xrightarrow{\text{ftor}} \\ \xleftarrow{\text{rtof}} \end{array} \mathcal{R}. \quad (12)$$

In Fig. 4.3 we illustrate the bijections from (12) using two examples.




Figure 3: A set of three labelled necklaces of planar trees (leftmost box) in bijection with a forest of labelled trees (middle box) in bijection with a rooted labelled tree (rightmost box). In the upper part of the figure the cycles are $\{(\{1, 5, 2\}), (\{3, 4\}), (\{6\}, \{7\})\}$, whereas in the lower part the cycles are $\{(\{1, 5, 2\}), (\{3, 4\}), (\{6\}), (\{7\})\}$

4.4 Other recursive tree-like combinatorial classes

The tree-like structure constructed from sequences is the most rigid one. It involves rooted trees that are embedded in a plan in such a way that the branches are always pointing downwards and so, the order of the sequences of the subtrees is relevant. If we relax the constraint of the orientation of the branches then the trees become invariant by rotation.

In that context, a tree is a non-oriented graph without cycle with a privileged vertex called a root. Each root can be seen as a labeled necklace on which the subtrees are hanged, forming a sort of windmill (see Figure 4 for examples). In other words, a tree is either an isolated root or a root with a cycle of trees. The combinatorial class satisfies the isomorphism

$$\mathcal{W} \equiv \bullet \boxplus (\bullet \boxtimes \text{CYC}(\mathcal{W})), \quad (13)$$

and its generating function satisfies

$$S_{\mathcal{W}}(x) = x \left(\log \left(\frac{1}{1 - S_{\mathcal{W}}(x)} \right) + 1 \right). \quad (14)$$

Expanding both sides of the equation and identifying the coefficients, we get a system, the resolution of which allows us to obtain the first cases of the enumeration:

$$1, 2, 9, 68, 730, 10164, 173838, 3524688, 82627200, \dots \quad A000169 \quad [16]$$




Figure 4: The nine windmills of degree 3.

This is another example of a tree-like structure studied among others in [1]. Notice that no closed form for the generating function is known but there exists a formula for the coefficients as a combination of Stirling numbers of first kind,

$$W_n = \sum_{i=0}^n i! \binom{n}{i} s_{n-1,i}, \quad (15)$$

where $s_{n,i}$ denotes the (unsigned) Stirling number of first kind that counts the number of permutations of n objects with exactly i cycles. Indeed, from (14), $S_{\mathcal{W}}(x)$ is the inverse of $g(x) = \frac{x}{1 - \log(1-x)}$ for the composition. The Lagrange inversion theorem [7] is a classical combinatorial tools allowing us to compute the Taylor expansion of inverse function. In our case, the direct application of the Lagrange inversion Theorem implies that

$$W_n = \left(\frac{d}{dx} \right)^{n-1} \left(\frac{x}{g(x)} \right)^n \Big|_{x=0} = \left(\frac{d}{dx} \right)^{n-1} \left(1 + \log \left(\frac{1}{1-x} \right) \right)^n \Big|_{x=0} \quad (16)$$

is the coefficient of x^{n-1} in $\left(1 - \log \left(\frac{1}{1-x} \right) \right)^n$ multiplied by $(n-1)!$. Knowing that the exponential generating function of Stirling's numbers $s_{i,k}$ (k fixed) is $\frac{1}{k!} \log \left(\frac{1}{1-x} \right)^k =$

$\sum_i s_{i,k} \frac{x^i}{i!}$, an easy computation allows us to deduce (15) from (16).

The last example we consider is the one where no more order constraints are imposed on the sub-trees of the same node. In this context, a tree is a root with a (possibly empty) set of trees. The trees of this kind can be drawn as nesting of disjoint discs with numbered surfaces (see some examples in Figure 5). Notice that, in these examples nested disk configurations, of degree 3 are as numerous as the windmills of degree 3 (see Figure 4). Of course, this is not always the case and generally, there are fewer nested discs




Figure 5: The nine configurations of nested discs of degree 3.

configurations than windmills. For instance, there are two windmills of degree 4 the roots of which is labeled by 1 with three sub-windmills of degree 1 while there is only one nested discs configuration the root of which is labeled by 1 containing three discs (see Figure 6).




Figure 6: Two windmills and one nested discs configuration.

The combinatorial class satisfies the isomorphism

$$\mathcal{N}pt = \bullet \boxtimes \text{SET}(\mathcal{N}pt),$$

and its generating series satisfies the functional equation

$$S_{\mathcal{N}pt}(x) = xe^{S_{\mathcal{N}pt}(x)}.$$

Solving these equation, one finds

$$S_{\mathcal{N}pt}(x) = -W(-x),$$

where $W(x)$ denotes the Lambert W function that is the principal branch of the functional inverse of $x \rightarrow xe^x$ [3]. The Taylor expansion of $W(x)$ is obtained by applying Lagrange inversion Theorem and implies $Npt_n = n^{n-1}$. The sequence of the Npt_n 's can also be found in [16]:

$$1, 2, 9, 64, 625, 7776, 117649, 2097152, \dots \quad A000169.$$

References

- [1] F. Bergeron, G. Labelle, and P. Leroux, *Combinatorial Species and Tree-like Structures*, Encyclopedia of Mathematics and its Applications, M. Readdy, Trans. Cambridge: Cambridge University Press, 1997.
- [2] K. T. Chen, R. H. Fox, R. C. Lyndon, *Free differential calculus. IV. The quotient groups of the lower central series*, Ann. of Math. (2), 68 (1958), pp. 81-95.
- [3] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, *On the LambertW function*, Adv. Comput. Math. (5), 1 (1996), pp. 329–359.
- [4] P. Flajolet, and R. Sedgewick, *Analytic combinatorics*, Cambridge University Press, 2009.
- [5] F. Hivert, J. C. Novelli, and J. Y. Thibon, *The algebra of binary search trees*, Theoretical Computer Science, 339(1), 129-165, 2005.
- [6] D. E. Knuth, *The Art of Computer Programming, Volume 3: (2nd Ed.) Sorting and Searching*, Addison Wesley Longman Publishing Co., Inc., 1998.
- [7] J. L. Lagrange, *Nouvelle méthode pour résoudre les équations littérales par le moyen des séries*, Mémoires de l'Académie Royale des Sciences et Belles-Lettres de Berlin, (24), 251-326, 1770.
- [8] P. Leroux and B. Miloudi, *Généralisations de la Formule d'Otter*, Annales des sciences mathématiques du Québec, 16(1), 53-80, 1992.
- [9] M. Lothaire, *Combinatorics on words*, Cambridge Mathematical Library **17** Foreword by Roger Lyndon (2nd ed.), 1997.
- [10] R. C. Lyndon, *On Burnside's problem*, Transactions of the American Mathematical Society, 77 **2** (1954), 202-215.
- [11] C. Reutenauer, *Free lie algebras*, Handbook of Algebra, North-Holland **3** (2003), 887-903.
- [12] M.P. Schützenberger, *On a factorisation of free monoids*, Proc. Amer. Math. Soc., 16 (1965), pp. 21-24.
- [13] A. I. Shirshov, *On free Lie rings*, Mat. Sbornik N.S., 45 **87** (1958), 113-122.
- [14] A. I. Shirshov, *Subalgebras of free Lie algebras*, Mat. Sbornik N.S., 33 **75** (1953), 441-52.
- [15] R. P. Stanley, *Catalan Numbers*, Cambridge University Press, 2015.
- [16] OEIS Foundation Inc. (2020), The On-Line Encyclopedia of Integer Sequences, <http://oeis.org/>.