FINITELY BOUNDED HOMOGENEITY TURNED INSIDE-OUT

JAKUB RYDVAL

ABSTRACT. The classification problem for countable finitely bounded homogeneous structures is notoriously difficult, with only a handful of published partial classification results, e.g., for directed graphs. We introduce the Inside-Out correspondence, which links the classification problem, viewed as a computational decision problem, to the problem of testing the embeddability between reducts of countable finitely bounded homogeneous structures. On the one hand, the correspondence enables polynomial-time reductions from various decision problems that can be represented within the embeddability problem, e.g., the double-exponential square tiling problem. This leads to a new lower bound for the complexity of the classification problem: 2NEXPTIME-hardness. On the other hand, it also follows from the Inside-Out correspondence that the classification (decision) problem is effectively reducible to the (search) problem of finding a finitely bounded Ramsey expansion of a countable finitely bounded homogeneous structure. We subsequently prove that the closely related problem of homogenizability is already undecidable.

1. Introduction

A structure \mathfrak{B} over a finite relational signature is homogeneous if every isomorphism between two of its finite substructures extends to an automorphism of \mathfrak{B} . If B is additionally countable, then it is uniquely determined up to isomorphism by its age, i.e., the class of all finite structures embeddable into \mathfrak{B} [40]. It is well-known that there are uncountably many countable homogeneous structures already among directed graphs [?], and hence countable homogeneous structures are typically only studied under additional restrictions. A class K of finite structures over a finite relational signature τ is finitely bounded [48] if it consists of all finite τ -structures which do not embed any member of some finite set \mathcal{N} of finite τ -structures (bounds); we write $\mathcal{K} = \text{Forb}_e(\mathcal{N})$. A countable τ -structure is finitely bounded if its age has this property. By Fraïssé's theorem, a finitely bounded class K forms the age of an (up to isomorphism unique) countable finitely bounded homogeneous structure if and only if K has the Amalgamation Property (AP). The classification problem for countable finitely bounded homogeneous structures is notoriously difficult [3, 44, 33], with only a handful of published partial classification results, e.g., for directed graphs [31]. Viewed as a computational decision problem, it can be stated as follows:

INSTANCE: A finite set \mathcal{N} of finite structures over a finite relational signature τ . QUESTION: Does Forb_e(\mathcal{N}) have the AP?

In the present article, we study the computational complexity of the classification problem (for countable finitely bounded homogeneous structures). We first adopt a different but equivalent definition of finite boundedness, namely, in terms of

A conference version of this material appeared in the Proceedings of the 51st EATCS International Colloquium on Automata, Languages and Programming (ICALP), #150, 1-20, 2024 [52].

definability by a universal first-order sentence. The advantage is that now the inputs to the classification problem are more stable under basic modifications, e.g., their size does not increase exponentially if we add a fresh symbol to the signature. The downside is that the equivalence of the two definitions only holds up to a single-exponential blow-up in one direction. We elaborate on this below.

For a set of bounds \mathcal{N} , we define the size of \mathcal{N} as the sum of the sizes of all structures in \mathcal{N} , where the size of a structure is the sum of the cardinalities of the domain and all relations. For a universal first-order sentence Φ , we write $\operatorname{fm}(\Phi)$ for the class of its finite models. Given a finite set of bounds \mathcal{N} , we can obtain a universal first-order sentence Φ of size polynomial in the size of \mathcal{N} satisfying $\operatorname{fm}(\Phi) = \operatorname{Forb}_e(\mathcal{N})$ simply by describing each structure in \mathcal{N} up to isomorphism using a quantifier-free formula. However, given a universal sentence Φ , it can be the case that a smallest \mathcal{N} satisfying $\operatorname{fm}(\Phi) = \operatorname{Forb}_e(\mathcal{N})$ is of size single-exponential in the size of Φ . The reason is that obtaining \mathcal{N} from Φ is comparable to rewriting Φ in CNF. Nevertheless, we will see that the size difference between the two inputs does not have a significant impact on our results.

The classification problem can now be stated as follows:

INSTANCE: A universal first-order sentence Φ over a finite relational signature τ . QUESTION: Does fm(Φ) have the AP?

1.1. Lower bounds. We show that the classification problem admits an efficient reduction from the problem of testing the embeddability between reducts of countable finitely bounded homogeneous structures. Note that, using a standard compactness argument (e.g. König's tree lemma), the embeddability problem can be equivalently phrased as the containment problem for finitely bounded amalgamation classes up to taking reducts to a given subset of their signatures. This is the formulation that we will use for the remainder of the present article because it allows us to formulate our results in their full generality, see Theorem 1.1. The formulation of Theorem 1.1 below only covers the cases with the *Strong Amalgamation Property* (SAP) but we provide an additional auxiliary result, Theorem 1.2, showing that this restriction is without loss of generality. For a class \mathcal{K} of structures with a common signature σ and a subset $\tau \subseteq \sigma$, we write $\mathcal{K}|_{\tau}$ for the class of the τ -reducts of all structures in \mathcal{K} .

Theorem 1.1 (First Inside-Out lemma). There exists a polynomial-time computable function Ω mapping each pair Φ_1, Φ_2 of universal first-order sentences and each common subset τ of their signatures to a universal sentence $\Phi := \Omega(\Phi_1, \Phi_2, \tau)$ such that the following are equivalent:

- (1) $fm(\Phi)$ has the SAP;
- (2) $fm(\Phi)$ has the AP;
- (3) $\operatorname{fm}(\Phi_1)$ has the SAP and $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$.

Theorem 1.2. There exists a polynomial-time computable function Γ mapping each universal first-order sentence Φ to a universal first-order sentence $\Gamma(\Phi)$ over the signature of Φ expanded by a fresh binary symbol E such that:

- (1) $fm(\Phi)$ has the AP if and only if $fm(\Gamma(\Psi))$ has the SAP;
- (2) if τ is a common subset of the signatures of Φ_1 and Φ_2 , then

$$\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$$
 if and only if $\operatorname{fm}(\Gamma(\Phi_1))|^{\tau \cup \{E\}} \subseteq \operatorname{fm}(\Gamma(\Phi_2))|^{\tau \cup \{E\}}$.

The next corollary is a direct consequence of Theorem 1.1 and Theorem 1.2.

Corollary 1.3. The following four decision problems are polynomial-time equivalent:

- Given a universal first-order sentence Φ over a finite relational signature, decide whether fm(Φ) has the (S)AP;
- Given universal first-order sentences Φ_1 and Φ_2 over finite relational signatures σ_1 and σ_2 , respectively, and a subset $\tau \subseteq \sigma_1 \cap \sigma_2$, decide whether

$$\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$$
 and $\operatorname{fm}(\Phi_1)$ has the $(S)AP$;

The significance of our reduction lies in the fact that it enables further polynomial-time reductions from various decision problems that can be represented within such containment. As an example, we show how to efficiently reduce from the double-exponential square tiling problem to the containment problem for reducts of finitely bounded classes with the *Free Amalgamation Property* (FAP).

Theorem 1.4. The question whether $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$ holds for a given pair of universal first-order sentences Φ_1 and Φ_2 , such that $\operatorname{fm}(\Phi_1)$ and $\operatorname{fm}(\Phi_2)$ both have the FAP, and a given subset τ of their signatures is $\operatorname{2NEXPTIME}$ -hard.

Theorem 1.1 and Theorem 1.4 together yield the following lower bound for the complexity of the classification problem.

Corollary 1.5. The classification problem for countable finitely bounded homogeneous structures is 2NEXPTIME-hard.

Before we proceed further, we remark that Corollary 1.5 has a counterpart in the setting where inputs are specified by sets of bounds. Here the lower bound drops to NEXPTIME (Remark 5.2).

1.2. **Towards decidability.** The question of decidability of the classification problem has been considered many times in the context of the Lachlan-Cherlin programme for countable homogeneous structures [3, 44, 33], and is known to have a positive answer in the case of binary signatures [46, 17]. It also appears as an open problem in Bodirsky's book on infinite-domain constraint satisfaction [10].

In the light of the fact that the definition of finite boundedness using universal first-order sentences is equivalent to the standard one only up to a single-exponential blow-up in size of the input, it is important that we also provide evidence of provable relative increase in complexity compared to the case of binary signatures. Otherwise, it might as well be the case that our hardness results are only byproducts of the succinctness of the input specification.

An inspection of the decidability results mentioned above reveals that the classification problem for countable finitely bounded homogeneous structures over binary signatures is decidable in coNEXPTIME (Proposition 3.3). We do not expect this upper bound to be sharp, and the best lower bound for the binary case we currently have is Π_3^p -hardness (Proposition 3.4). We remark that the coNEXPTIME upper bound drops to Π_2^p if the inputs are specified by sets of bounds instead of universal first-order sentences (Theorem 15 in [6]).

We do not have any upper bound on the complexity of the classification problem for arbitrary finite signatures, but we have a result which *almost* shows that the classification problem admits an efficient reduction back to the containment problem for reducts of strong amalgamation classes (Theorem 1.6).

Theorem 1.6 (Second Inside-Out lemma). There exists a polynomial-time computable function Δ mapping each universal first-order sentence Φ over a finite

relational signature ρ to a pair of universal first-order sentences $\Phi_1 := \Delta(\Phi)_1$ and $\Phi_2 := \Delta(\Phi)_2$ over some finite relational signatures τ and σ , respectively, such that $\rho \subseteq \tau \subseteq \sigma$ and the following are equivalent:

- (1) $fm(\Phi)$ has the SAP;
- (2) $fm(\Phi_1)$ has the SAP;
- (3) $fm(\Phi_2)$ has the SAP;
- (4) $\operatorname{fm}(\Phi_1) \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$.

On an input Φ to the classification problem, we can compute $\Delta(\Phi)$ and forward it to an oracle for the containment problem for reducts of strong amalgamation classes. Note that the only issue with this reduction are false positives; an oracle for the containment problem might return yes on inputs where $fm(\Phi_1) \nsubseteq fm(\Phi_2)|^{\tau}$ but $fm(\Phi_1)$ or $fm(\Phi_2)$ does not have the SAP. We show that this issue can be eliminated under the assumption of a certain stronger version of a conjecture from structural Ramsey theory.

More specifically, we show that Theorem 1.6 combined with Theorem 1.2 can be used to effectively reduce from the classification decision problem to the search problem of finding a finitely bounded Ramsey expansion of a given finitely bounded strong amalgamation class. The question whether such an expansion always exists was left open in [23] in a considerably more general setting, see also Conjecture 1 in [56]. While the most general version of this question was answered negatively in [38], it is remains open for arbitrary amalgamation classes over finite relational signatures (see Question 7.1 in [38]). For finitely bounded amalgamation classes, the question was formulated as a conjecture in [10]. The next result implies the decidability of the classification problem under the assumption that every finitely bounded strong amalgamation class has a computable finitely bounded Ramsey expansion.

Theorem 1.7. There exists a non-deterministic exponential-time reduction from the classification (decision) problem for countable finitely bounded homogeneous structures to the (search) problem of finding a finitely bounded Ramsey expansion for a finitely bounded class with the SAP.

Note that it follows from Corollary 1.5 and Theorem 1.7 that there is no polynomial function bounding the sizes of finitely bounded Ramsey expansions of finitely bounded strong amalgamation classes. We attribute this to the fact that universal first-order sentences provide a more compact representation of finitely bounded classes than sets of bounds. In the latter regime, the lower bound in Corollary 1.5 drops to NEXPTIME (Remark 5.2) while the conditional upper bound stemming from Theorem 1.7 stays the same. On the other hand, if the classification problem turns out to be undecidable, then it follows from Theorem 1.7 that there is not even any computable function bounding the sizes of finitely bounded Ramsey expansions. However, purely on the basis of empirical evidence for the complexity of the containment problem [27, 18] and the usual sizes of finitely bounded Ramsey expansions [42], we would rather speculate that the classification problem is decidable. We leave the following question open.

Open question: Is the classification problem for countable finitely bounded homogeneous structures in 2NEXPTIME?

1.3. **Homogenizability.** If we relax the input to sentences of the logic *Strict NP* (SNP) [39] and the question to the existence of a structure with a finitely bounded

homogeneous expansion, then we obtain a different problem, a weaker form of homogenizability [2, 5, 36, 43].

INSTANCE: An SNP sentence Φ over a finite relational signature τ . QUESTION: Does there exist a universal first-order sentence Φ^+ such that

$$\operatorname{fm}(\Phi) = \operatorname{fm}(\Phi^+)|^{\tau}$$
 and $\operatorname{fm}(\Phi^+)$ has the AP?

A countable structure is (finitely-bounded) homogenizable if it is first-order interdefinable with (finitely bounded) homogeneous structure with a finite relational signature. Sufficient conditions for finitely-bounded homogenizability were provided by Hubička and Nešetřil [41], generalizing previous work of Cherlin, Shelah, and Shi [34]. We prove that the above relaxed version of the classification problem is already undecidable, even if the input SNP sentence comes from the Datalog fragment and uses at most binary relation symbols (Theorem 1.8). Our proof also covers the more specific question of homogenizability. As a byproduct of our proof, we also get the undecidability of some other properties for Datalog programs. This concerns, e.g., the questions whether a given Datalog program can be rewritten in the logic MMSNP, whether it solves some finite-domain CSP, or whether it defines a structure with a homogeneous Ramsey expansion in a finite relational signature.

To keep these results as general as possible, we formulate them for some reasonable promise relaxations of the original question, i.e., where a subclass and a superclass of the positive instances are being separated from each other with the promise that the input never belongs to the complement of the subclass within the superclass. Both items in Theorem 1.8 are formulated as a statement of the form "the question whether X or not even Y is hard." This is a compact way for writing that both X and Y (and every property in between) are hard, the formulation tacitly assumes that the inputs never satisfy "Y and not X."

Theorem 1.8. For a given a Datalog sentence Φ over a signature containing at most binary relation symbols, it is undecidable whether

- (1) Φ is logically equivalent to a monadic Datalog sentence, or Φ is not even logically equivalent to any GMSNP sentence;
- (2) Whether simultaneously $fm(\Phi)$ is:
 - the CSP of a finite structure,
 - the age of a finitely-bounded homogenizable structure,
 - the age of a reduct of a finitely bounded homogeneous Ramsey structure, or $fm(\Phi)$ is not even the CSP or the age of any ω -categorical structure.

We remark that we do not know how to prove the undecidability of these problems for any further restrictions to the input, e.g., for universal first-order sentences. Therefore, Theorem 1.8 merely shows that SNP sentences are an exceptionally bad choice of an input to the question (albeit one that is often used [5, 50]). To further emphasise this point, note that the definability of finite-domain CSPs in monadic Datalog is decidable if the input is specified by a structure parametrizing the CSP. This is a direct consequence of Theorem 7.4 in [7]. By Theorem 1.8, the same question is undecidable if the input is specified by an SNP sentence. For CSPs of reducts of finitely bounded homogeneous structures, there is currently no better specification for the input than by an SNP sentence.

The next corollary extracts the statement originally announced in the abstract.

Corollary 1.9. It is undecidable whether a given SNP sentence defines the age of homogenizable structure. The statement is true even if the SNP sentence comes from the Datalog fragment and uses at most binary relation symbols.

1.4. **Organisation of the article.** Section 2 provides some basic notions from logic necessary for the presentation of the proofs of our results, and Section 3 provides some basic information about the classification problem for countable finitely bounded homogeneous structures.

Section 4 contains the proofs of Theorem 1.2, Theorem 1.1, and Theorem 1.6. Our proofs of these results are elementary and require very little external knowledge.

Section 5 contains the proof of Theorem 1.4. The proof is by a polynomial-time reduction from the double-exponential square tiling problem. The idea of the proof is loosely inspired by the proof of 2NEXPTIME-hardness of the containment problem for the logic MMSNP from [27]. Despite the fact that every MMSNP-definable class is a reduct of a finitely bounded free amalgamation class [18], we cannot use the hardness proof from [27] directly. The reason is that the amount of symbols needed to obtain a finitely bounded expansion which is an amalgamation class might be exponential in the size of the input. This is essentially a consequence of the lower bounds on the arities of the relations of such an expansion obtained in [41].

Section 6 contains the proof of Theorem 1.7. The proof is by a straightforward application of a combinatorial tool known as the "canonisation lemma" [23, 21]. The idea of the proof is loosely inspired by the proof of decidability of the containment for classes of finite structures definable in the logic MMSNP from [18].

Section 7 contains the proof of Theorem 1.8. The proof is by a polynomial-time reduction from the problem of testing the regularity of context-free languages.

1.5. Connections to automata theory. As a part of the proof of Theorem 1.8, we encode context-free grammars and deterministic finite automata into finitely bounded classes. In the case of deterministic finite automata, and also in the case of regular grammars, this can be done so that the resulting class has the SAP (even FAP). We invite the reader to view at Figure 1, which contains a lattice of encodings between several categories which play a role in the proof of Theorem 1.8.

The encodings in Figure 1 are functorial w.r.t. containment and the labels on the edges describe their variance. The contravariance of some of the encodings explains why Theorem 1.1 does not immediately yield the undecidability of the classification problem for countable finitely bounded homogeneous structures even though the containment of regular languages in context-free languages is undecidable. Instead, we can only reduce from the containment of context-free languages in regular languages, which is not harder than the containment between regular languages [4]. Our results indicate that a more general version of this phenomenon might be happening on the level of reducts of finitely bounded classes with the SAP.

The general message we want to convey is that the relationship between finitely bounded classes with the SAP and arbitrary finitely bounded classes shares similarities with the relationship between regular grammars and context-free grammars. One key difference is that the regularity of context-free grammars can be tested in linear time, while testing the SAP for finitely bounded classes is 2NEXPTIME-hard.

1.6. Connections to constraint satisfaction. SNP is an expressive fragment of existential second-order logic and thus, by Fagin's Theorem, of the complexity class NP. Despite the name, SNP already has the full power of NP, in the sense




FIGURE 1. A lattice of functorial encodings between several categories used in the present article. The edge-labels provide additional information about the variance of the encodings w.r.t. containment.

that every problem in NP is equivalent to a problem in SNP under polynomial-time reductions [39]. In addition, this logic class has many connections to Constraint $Satisfaction\ Problems\ (CSPs)$. The constraint satisfaction problem of a relational structure \mathfrak{B} , denoted by $CSP(\mathfrak{B})$, is (the membership problem for) the class of all finite structures which homomorphically map to \mathfrak{B} . Every computational decision problem is polynomial-time equivalent to a CSP [14]. Many practically relevant problems like Boolean satisfiability or graph colorability can even be formulated as a finite-domain CSP, i.e., where the template \mathfrak{B} can be chosen finite. The basic link from SNP to CSP is that every sentence of the monotone fragment of this logic defines a finite disjoint union of CSPs of (possibly infinite) relational structures [10]. There are, however, some more nuanced connections, such as the one that led to the formulation of the Feder-Vardi conjecture, now known as the finite-domain CSP dichotomy theorem [59].

In their seminal work [39], Feder and Vardi showed that the *Monotone Monadic* fragment of SNP (MMSNP) exhibits a dichotomy between P and NP-completeness if and only if the seemingly less complicated class of all finite-domain CSPs exhibits such a dichotomy, they also conjectured the latter to be true. The logic class MMSNP contains all finite-domain CSPs, and many other interesting combinatorial problems, e.g., whether the vertices of a given graph can be 2-coloured without obtaining any monochromatic triangle [49]. The Feder-Vardi conjecture was confirmed in 2017 independently by Bulatov and Zhuk [29, 58].

There is a yet unconfirmed generalization of the Feder-Vardi conjecture, to the CSPs of reducts of countable finitely bounded homogeneous structures, formulated by Bodirsky and Pinsker in 2011 [22]. Here we refer to it as the *Bodirsky-Pinsker conjecture*. Roughly said, the condition imposed on the structures within the scope of the Bodirsky-Pinsker conjecture ensures that the CSP is in NP and that its template enjoys some of the universal-algebraic properties that have played an essential role in the proofs of the Feder-Vardi conjecture [8]. At the same time,

it covers CSP-reformulations of many natural problems in qualitative reasoning, as well as all problems definable in MMSNP. Every reduct of a countable finitely bounded homogeneous structure is uniquely described by an SNP sentence up to isomorphism, and its CSP is definable in monotone SNP.

Current approaches to the Bodirsky-Pinsker conjecture are problematic in the sense that they rely on the scarce pool of partial classification results for countable finitely bounded homogeneous structures, see, e.g., [16, 20, 45, 15]. Despite not having any direct consequences for the conjecture, our results provide evidence for the need for a fundamentally new language-independent approach to it. As an example, it is a folklore fact that from every finite structure $\mathfrak A$ over a finite relational signature one can construct in polynomial time a finite structure \mathfrak{B} over a finite binary relational signature such that $CSP(\mathfrak{A})$ and $CSP(\mathfrak{B})$ are polynomial-time equivalent [30, 39]. By Corollary 1.5 and Proposition 3.3, such a reduction is unlikely to exist for universal sentences representing finitely bounded homogeneous structures, unless it avoids the classification problem. Another piece of evidence is the fact that, for every Φ as in Theorem 1.1, the CSP of every structure whose age equals $fm(\Phi)$ is trivial, independently on whether or not $fm(\Phi)$ has the SAP (Remark 4.4). First steps towards a language-independent approach to the conjecture were taken in the recent works of Mottet and Pinsker [51] and Bodirsky and Bodor [11], but they do not fully address the issues stemming from the classification problem. We elaborate on this claim in the text below.

Besides CSPs, the classification problem is also relevant to other related areas of theoretical computer science such as verification of database-driven systems [26], sets with atoms [35], or description logics with concrete domains [47, 6].

1.7. Subtleties of the infinite-domain CSP. In 2016, Bodirsky and Mottet presented an elegant tool for lifting tractability from finite-domain constraint satisfaction to the infinite [19], hereby establishing the first general link between the Feder-Vardi and the Bodirsky-Pinsker conjecture. Since then, their method has been used numerous times to prove new or reprove old complexity classification results for infinite-domain CSPs. One prominent such example is the universal-algebraic proof of the complexity dichotomy for MMSNP [18]. Conveniently enough, every MMSNP sentence defines a finite union of CSPs of structures within the scope of the Bodirky-Pinsker conjecture [13], so the classification problem was not relevant in this context. There is a prospect that the methods from [19] will also prove useful in proving a complexity dichotomy for the even more general logic class Guarded Monotone SNP (GMSNP) introduced in [9]. Also GMSNP enjoys the above mentioned property of MMSNP [17], and hence avoids the the classification problem. However, outside of GMSNP there exists a regime where the methods from [19] definitely fall short, and where the classification problem becomes relevant.

Consider for instance the complexity dichotomy for temporal CSPs, i.e., for CSPs of structures with domain \mathbb{Q} and whose relations are definable by a Boolean combination of formulas of the form (x=y) or (x < y), obtained by Bodirsky and Kára in 2010 [16]. At the present time, these problems are already very well understood; tractable temporal CSPs can always be solved by an algorithm that repeatedly searches for a set of potential minimal elements among the input variables, where each instance of the search is performed using an oracle for a tractable finite-domain CSP. The latter is determined by the shape of the Boolean combinations.

E.g., in the case of

$$CSP(\mathbb{Q}; \{(x, y, z) \in \mathbb{Q}^3 \mid (x = y < z) \lor (y = z < x) \lor (z = x < y)\}),$$

solving the finite-domain CSP in question amounts to solving linear equations modulo 2 [16, 24]. It is known that the tractability results from [16] cannot be obtained using the reduction from [19].

In 2022, Mottet and Pinsker introduced the machinery of *smooth approximations* [51], vastly generalizing the methods in [19]. The last section of their paper is devoted to temporal CSPs, and the authors manage to reprove a significant part of the dichotomy on just a few pages. They achieve this by applying some of their general results to first-order expansions of $(\mathbb{Q}; <)$ and obtaining either NP-hardness for the CSP, or one of the two types of symmetry that played a fundamental role in the original proof from [16]. This symmetry can then be used to prove correctness of the reduction to a finite-domain CSP described above, but only under an explicit usage of the homogeneity of $(\mathbb{Q};<)$ (see Proposition 3.1 in [24] and the last section of [51]). In contrast to the methods in [19] which only use homogeneity as a blackbox, this approach can be described as language-dependent.

A similar situation occurs in the case of phylogeny CSPs [15], which capture decision problems concerning the existence of a binary tree satisfying certain constraints imposed on its leaves. Tractable phylogeny CSPs are strikingly similar to tractable temporal CSPs; they can always be solved by an algorithm that repeatedly searches for a subdivision of the input variables into two parts, representing the two different branches below the root of a binary tree, where each instance of this search is performed using an oracle for a tractable finite-domain CSP. However, for tractable phylogeny CSPs, already the homogeneity of the infinite-domain CSP template is both sufficient and necessary for proving the correctness of the reduction to the finite-domain CSP (Theorem 6.13 and Lemma 6.12 in [15]). We can therefore speak of a case of extreme language-dependency. Temporal and phylogeny CSPs are special cases of CSPs of structures obtainable from the universal homogeneous binary tree [12] by specifying relations using first-order formulas. Achieving a complexity dichotomy in this context will require a non-trivial combination of the methods from [16] and [15].

An optimal way of approaching the conjecture would be gaining a very good understanding of the class of reducts of finitely bounded homogeneous structures, e.g., through some sort of a classification. However, it is unclear how realistic this prospect is as model-theoretic properties often tend to be undecidable [32]. Our results point in the direction of decidability of the classification problem, but with a very impractical non-deterministic double-exponential lower bound. We remark that homogeneity is a vital part of the Bodirsky-Pinsker conjecture; this assumption can be weakened or strengthened but not dropped entirely, as otherwise we get a class that provably does not have a dichotomy [39, 10].

2. Preliminaries

Due to the technical nature of our results, we need to fix some notation. The set $\{1, \ldots, n\}$ is denoted by [n], and we use the bar notation for tuples. We extend the usual containment relation on sets to tuples by ignoring the ordering on the entries. E.g., we might write $x \in \bar{t}$ if x appears in an entry of a tuple \bar{t} .

A (relational) signature τ is a set of relation symbols, each $R \in \tau$ with an associated natural number called arity. We say that τ is binary if it consists of symbols of arity ≤ 2 . A (relational) τ -structure \mathfrak{A} consists of a set A (the domain) together with the relations $R^{\mathfrak{A}} \subseteq A^k$ for each $R \in \tau$ with arity k. An expansion of \mathfrak{A} is a σ -structure \mathfrak{B} with A=B such that $\tau\subseteq\sigma$, $R^{\mathfrak{B}}=R^{\mathfrak{A}}$ for each relation symbol $R \in \tau$. Conversely, we call \mathfrak{A} a reduct of \mathfrak{B} and denote it by $\mathfrak{B}|^{\tau}$. These two notions naturally extend to classes of structures over a common signature. The union of two τ -structures $\mathfrak A$ and $\mathfrak B$ is the τ -structure $\mathfrak A \cup \mathfrak B$ with domain $A \cup B$ and relations of the form $R^{\mathfrak{A} \cup \mathfrak{B}} := R^{\mathfrak{A}} \cup R^{\mathfrak{B}}$ for every $R \in \tau$. A disjoint union of \mathfrak{A} and $\mathfrak B$ is a union of two copies of $\mathfrak A$ and $\mathfrak B$ with disjoint domains. The *substructure* of a τ -structure $\mathfrak A$ on a subset $B\subseteq A$ is the τ -structure $\mathfrak B$ with domain B and relations $R^{\mathfrak{B}} = R^{\mathfrak{A}} \cap B^k$ for every $R \in \tau$ of arity k. The factor of a τ -structure \mathfrak{A} through an equivalence relation $E \subseteq A^2$ is the τ -structure \mathfrak{A}/E with domain A/E and relations $R^{\mathfrak{A}/E} = R^{\mathfrak{A}/E}$ for every $R \in \tau$ of arity k. We say that E is a relational congruence on \mathfrak{A} if the definitions of the relations of \mathfrak{A}/E do not depend on the choices of the representatives of the equivalence classes w.r.t. E.

A homomorphism $h\colon \mathfrak{A}\to \mathfrak{B}$ for τ -structures $\mathfrak{A},\mathfrak{B}$ is a mapping $h\colon A\to B$ that preserves each relation of \mathfrak{A} , i.e., if $\bar{t}\in R^{\mathfrak{A}}$ for some k-ary relation symbol $R\in \tau$, then $h(\bar{t})\in R^{\mathfrak{B}}$. We write $\mathfrak{A}\to \mathfrak{B}$ if \mathfrak{A} maps homomorphically to \mathfrak{B} . For a structure \mathfrak{A} over a finite relational signature τ , we define $\mathrm{CSP}(\mathfrak{A})$ as the class of all finite τ -structures which homomorphically map to \mathfrak{A} . An embedding is a homomorphism $h\colon \mathfrak{A}\to \mathfrak{B}$ that additionally satisfies the following condition: for every k-ary relation symbol $R\in \tau$ and $\bar{t}\in A^k$ we have $h(\bar{t})\in R^{\mathfrak{B}}$ only if $\bar{t}\in R^{\mathfrak{A}}$. We write $\mathfrak{A}\hookrightarrow \mathfrak{B}$ if \mathfrak{A} embeds to \mathfrak{B} . The age of \mathfrak{A} , denoted by $\mathrm{age}(\mathfrak{A})$, is the class of all finite structures which embed to \mathfrak{A} . An isomorphism is a surjective embedding. Two structures \mathfrak{A} and \mathfrak{B} are isomorphic if there exists an isomorphism from \mathfrak{A} to \mathfrak{B} . An automorphism is an isomorphism from \mathfrak{A} to \mathfrak{A} . The orbit of a tuple $\bar{t}\in A^k$ in \mathfrak{A} is the set $\{g(\bar{t})\mid g\text{ is an automorphism of }\mathfrak{A}\}$. A countable structure \mathfrak{A} is ω -categorical if, for every $k\geq 1$, there are finitely many orbits of k-tuples in \mathfrak{A} . Every homogeneous structure in a finite relational signature is ω -categorical, and so are the reducts of such structures.

For a first-order sentence Φ , we denote the class of all its finite models by $\operatorname{fm}(\Phi)$. We say that a first-order formula is k-ary if it has k free variables. For a first-order formula ϕ , we use the notation $\phi(\bar{x})$ to indicate that the free variables of ϕ are among \bar{x} . This does not mean that the truth value of ϕ depends on each entry in \bar{x} .

We assume that equality = as well as the nullary predicate symbol \bot for falsity are always available when building first-order formulas. Thus, $atomic\ \tau$ -formulas, or τ -atoms for short, over a relational signature τ are of the form \bot , (x=y), and $R(\bar{x})$ for some $R \in \tau$. If ϕ is a disjunction of possibly negated τ -atoms, then we define the $Gaifman\ graph$ of ϕ as the undirected graph whose vertex set consists of all free variables of ϕ and where two distinct variables x, y form an edge if and only if they appear jointly in a negative atom of ϕ .

A universal first-order τ -sentence is of the form $\forall \bar{x}. \phi(\bar{x})$ for a quantifier-free τ -formula ϕ . Let Φ be a universal τ -sentence whose quantifier-free part ϕ is in CNF. We call Φ Horn if every clause of ϕ is Horn, i.e., contains at most one positive disjunct. We call Φ complete (connected) if the Gaifman graph of each clause of ϕ is complete (connected). It is a folklore fact that, if Φ is complete (connected), then fm(Φ) is preserved by (disjoint) unions.

An SNP τ -sentence is a second-order sentence Φ of the form $\exists X_1,\ldots,X_n \forall \bar{x}. \phi$ where ϕ is a quantifier-free formula in CNF over $\tau \cup \{X_1,\ldots,X_n\}$. We call Φ monadic if X_i is unary for every $i \in [n]$; monotone if ϕ does not contain any positive τ -atoms (in particular no positive equality atoms); and guarded if, for every positive atom β there exists a negative atom α containing all variables of β . Note that all notions from the previous paragraph easily transfer to SNP sentences viewed as universal sentences in an extended signature. The monadic monotone and the guarded monotone fragments of SNP are denoted by MMSNP and GMSNP, respectively. The monotone Horn fragment of SNP is commonly known as the logic programming language Datalog. When we say that a Datalog program Φ solves the CSP of a structure \mathfrak{B} , we simply mean that $\operatorname{fm}(\Phi) = \operatorname{CSP}(\mathfrak{B})$.

3. The Classification Problem

We start by stating Fraïssé's theorem and fixing some additional terminology. Let K be a class of finite structures in a finite relational signature τ closed under isomorphisms and substructures. An amalgamation diagram for K is a pair of structures $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathcal{K}$ whose substructures on $B_1 \cap B_2$ are identical. An amalgamation diagram is one-point if $|B_1 \setminus B_2| = |B_2 \setminus B_1| = 1$. Per convention, in a one-point amalgamation diagram $\mathfrak{B}_1,\mathfrak{B}_2$, we denote by b_1 and b_2 the unique elements contained in $B_1 \setminus B_2$ and $B_2 \setminus B_1$, respectively. An amalgam for an amalgamation diagram $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathcal{K}$ is a structure $\mathfrak{C} \in \mathcal{K}$ for which there are embeddings $f_1: \mathfrak{B}_1 \hookrightarrow \mathfrak{C}$ and $f_2: \mathfrak{B}_2 \hookrightarrow \mathfrak{C}$ such that $f_1|_{B_1 \cap B_2} = f_2|_{B_1 \cap B_2}$. Then \mathcal{K} has the amalgamation property (AP) if every amalgamation diagram for K has an amalgam in \mathcal{K} . The strong version of the AP is when the amalgam can always be chosen so that $f_1(B_1) \cap f_2(B_2) = f_1(B_1 \cap B_2)$. Without loss of generality, we may always assume that $f_1(B_1) \cup f_1(B_2) \subseteq B_1 \cup B_2$ in the case of AP and $f_1(B_1) \cup f_1(B_2) = B_1 \cup B_2$ in the case of SAP. For a class K closed under isomorphisms and substructures, the AP is implied by the property of being closed under unions, also called free amalgams. In this case we say that K has the Free Amalgamation Property (FAP).

Theorem 3.1 (Fraïssé). For a class K of finite structures in a finite relational signature τ , the following are equivalent:

- K is the age of an up to isomorphism unique countable homogeneous τstructure;
- K is closed under isomorphisms, substructures, and has the AP.

Countable homogeneous structures arise as limit objects of well-behaved classes of finite structures in the sense of Fraïssé's theorem. For finitely bounded classes, the formulation of this theorem is particularly simple because the closure under isomorphisms and substructures is trivially true. Every countable homogeneous structure is uniquely described by its age up to isomorphism. Consequently, every countable finitely bounded homogeneous structure is uniquely described by finite set of bounds or, equivalently, by a universal first-order sentence.

It is known that the question whether a finitely bounded class has the AP can be tested algorithmically in the case where the signature is binary [46]. This decidability result is based on the following observation.

Proposition 3.2 ([46]). A class of finite relational τ -structures that is closed under isomorphisms and substructures has the AP if and only if it has the AP restricted to one-point amalgamation diagrams.

As a consequence of Proposition 3.2, if a finitely bounded class over a binary signature does not have the AP, then the size of a smallest counterexample to the AP is polynomial in the size of the set of bounds [17]. Such a counterexample can be non-deterministically guessed and verified using a coNP-oracle, which places the problem at the second level of the polynomial hierarchy (Theorem 15 in [6]). Note that this upper bound only applies to the case where the input is specified by a set of bounds. If the input is specified by a universal first-order sentence, then it might be the case that a smallest set of bounds witnessing finite boundedness is exponentially larger. Consequently, the algorithm from [17] only gives us a relatively weak upper bound for the case where the inputs are specified by universal sentences.

Proposition 3.3. Let Φ be a universal sentence over a finite binary relational signature τ . If $fm(\Phi)$ does not have the AP, then the size of a smallest counterexample to the AP is at most single-exponential in the size of Φ . Consequently, the question whether $fm(\Phi)$ has the AP is decidable in conexprise.

Proof. Suppose that $fm(\Phi)$ does not have the AP. By Proposition 3.2, we may assume that a counterexample to the AP for $fm(\Phi)$ is a one-point amalgamation diagram formed by two structures $\mathfrak{B}_1, \mathfrak{B}_2 \in \text{fm}(\Phi)$. Let k be the number of variables in Φ . We define \mathcal{N} as the set of all τ -structures \mathfrak{A} such that $\mathfrak{A} \not\models \Phi$, $|A| \leq k$, and A = [|A|]. Clearly, Forb_e(\mathcal{N}) = fm(Φ), and the size of \mathcal{N} is at most single-exponential in the size of Φ . By the proof of Theorem 4 in [17], we may assume that the size of \mathfrak{B}_1 and \mathfrak{B}_2 is polynomial in the size of \mathcal{N} . Hence, we may assume that it is at most single-exponential in the size of Φ . Now we must verify that $\mathfrak{B}_1, \mathfrak{B}_2 \in \text{fm}(\Phi)$. This can be done in time exponential in the size of Φ simply by evaluating the quantifier-free part of Φ on all possible inputs. Subsequently, we must verify that no amalgam $\mathfrak{C} \in \text{fm}(\Phi)$ of \mathfrak{B}_1 and \mathfrak{B}_2 can be obtained either by identifying b_1 and b_2 , or by adding (b_1, b_2) or (b_2, b_1) to some relations of $\mathfrak{B}_1 \cup \mathfrak{B}_2$. This can also be done in time exponential in the size of Φ because only single-exponentially many structures $\mathfrak C$ need to be checked. In sum, the existence of a counterexample can be tested in NEXPTIME, which is what we had to show. П

The upper bound provided by Proposition 3.3 is not entirely unreasonable since a smallest counterexample to the AP might be of size exponential in the size of the input sentence even if the signature is binary. Such situations arise, e.g., in the proof of the next proposition. We include it together with a full proof as a warm-up for the more technically involved arguments in the next sections.

Proposition 3.4. Given a universal sentence Φ over a finite binary relational signature, the question whether fm(Φ) has the AP is Π_3^p -hard.

Proof. We reduce from $\forall \exists \forall \text{-DNF}$, a basic complete problem for the complexity class Π_3^p [55]. Consider a general instance of $\forall \exists \forall \text{-DNF}$, which is of the form

$$(1) \qquad \forall x_1, \dots, x_k \exists x_{k+1}, \dots, x_\ell \forall x_{\ell+1}, \dots, x_m. \, \phi,$$

where ϕ is a disjunction of conjunctions of possibly negated propositional variables. We define the signature τ as follows. For every $i \in [m]$, τ contains the symbol X_i , which is binary for $i \in [\ell] \setminus [k]$ and unary otherwise. In addition, τ also contains the binary symbol P and the two unary symbols L, R. We define Φ as

$$\forall x, y_1, y_2 \big(L(y_1) \land R(y_2) \land P(x, y_1) \land P(x, y_2) \Rightarrow \phi' \big),$$

where ϕ' is the τ -formula obtained from ϕ by the following syntactical replacement of propositional variables by τ -atoms. For every $i \in [m]$ we replace each instance of the variable x_i by:

- $X_i(y_1)$ if $i \in [k]$,
- $X_i(y_1, y_2)$ if $i \in [\ell] \setminus [k]$, $X_i(x)$ if $i \in [m] \setminus [\ell]$.

" \Rightarrow " Suppose that (1) is satisfiable. Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an arbitrary one-point amalgamation diagram for $fm(\Phi)$. If $\mathfrak{B}_1 \cup \mathfrak{B}_2 \models \Phi$, then we are done because $\mathfrak{B}_1 \cup \mathfrak{B}_2$ is an amalgam for \mathfrak{B}_1 and \mathfrak{B}_2 . So suppose that instead $\mathfrak{B}_1 \cup \mathfrak{B}_2 \not\models \Phi$. Consider an evaluation of the quantifier-free part of Φ witnessing the fact that $\mathfrak{B}_1 \cup \mathfrak{B}_2 \not\models \Phi$. By definition, b_1 and b_2 do not appear together in any relation of $\mathfrak{B}_1 \cup \mathfrak{B}_2$ and $\mathfrak{B}_1, \mathfrak{B}_2 \models \Phi$. Thus, by the shape of Φ , it must be the case that x is assigned some element $b \in B_1 \cap B_2$, y_1 is assigned b_1 , and y_2 is assigned b_2 (or vice versa). For $i \in \{1, 2\}$, define $T_i := \{j \in [k] \mid b_i \in X_j^{\mathfrak{B}_i}\}$. Now, for a fixed $i \in \{1, 2\}$, consider the propositional assignment where, for every $j \in [k]$, the truth value of x_j is set to true if and only if $j \in T_i$. By the assumption that (1) is satisfiable, there exists $T_i \subseteq [\ell] \setminus [k]$ such that the following holds: if, for every $j \in [\ell] \setminus [k]$, we set x_j to true if and only if $j \in T'_i$, then every assignment of truth values to the remaining variables $x_{\ell+1}, \ldots, x_m$ satisfies the quantifier-free part of (1). We obtain an amalgam $\mathfrak{C} \in \text{fm}(\Phi)$ for \mathfrak{B}_1 and \mathfrak{B}_2 by adding, for every $i \in \{1,2\}$ and $j \in [\ell] \setminus [k]$, the pair (b_i, b_{3-i}) to $X_j^{\mathfrak{B}_1 \cup \mathfrak{B}_2}$ if and only if $j \in T_i'$.

"\(= "\) Suppose that (1) is not satisfiable. Then there exists $T \subseteq [k]$ together with an assignment of truth values to the variables x_1, \ldots, x_k witnessing the unsatisfiability of (1) such that precisely $\{x_i \mid i \in T\}$ are set to true in the assignment. We use T to construct a one-point amalgamation diagram $\mathfrak{B}_1, \mathfrak{B}_2 \in \text{fm}(\Phi)$ with no amalgam in fm(Φ). For $i \in \{1,2\}$, we set $B_i \coloneqq \{b_i\} \cup \{b_S \mid S \subseteq [m] \setminus [\ell]\}$. The relations are as follows. We have $L^{\mathfrak{B}_1} = \{b_1\}$, $R^{\mathfrak{B}_1} = \emptyset$, $L^{\mathfrak{B}_2} = \emptyset$, $R^{\mathfrak{B}_2} = \{b_2\}$. Moreover, for every $j \in T$, $S \subseteq [m] \setminus [\ell]$, and $i \in S$, we have $b_1 \in X_j^{\mathfrak{B}_1}$, $(b_S, b_1) \in P^{\mathfrak{B}_1}$, $b_S \in X_i^{\mathfrak{B}_1}$, $b_2 \in X_i^{\mathfrak{B}_2}$, $(b_S, b_2) \in P^{\mathfrak{B}_2}$, and $b_S \in X_i^{\mathfrak{B}_2}$. There are no other tuples in the relations of \mathfrak{B}_1 or \mathfrak{B}_2 . Note that $\mathfrak{B}_1, \mathfrak{B}_2 \in \text{fm}(\Phi)$ because $R^{\mathfrak{B}_1} = \emptyset$ and $L^{\mathfrak{B}_2} = \emptyset$. Clearly, no amalgam for \mathfrak{B}_1 and \mathfrak{B}_2 can be obtained by identifying b_1 and b_2 because $L^{\mathfrak{B}_1} = \{b_1\}$ and $R^{\mathfrak{B}_2} = \{b_2\}$. By the assumption that (1) is not satisfiable, there is also no strong amalgam for \mathfrak{B}_1 and \mathfrak{B}_2 satisfying Φ because the pair (b_1,b_2) cannot be present in any subset of the relations $X_{k+1}^{\mathfrak{B}_1\cup\mathfrak{B}_2},\ldots,X_{\ell}^{\mathfrak{B}_1\cup\mathfrak{B}_2}$.

Very little progress has been done on signatures containing symbols of arities larger than 2. In particular, it is not even known whether the AP is decidable for finitely bounded classes in general. The scenario where this is not the case is not entirely unrealistic since the closely related Joint Embedding Property (JEP) is undecidable already for finitely bounded classes of graphs [28].

4. Inside-Out correspondence

In this section, we prove three of the results announced in the first part of the introduction. We start with the proof of the following theorem, which was originally presented as Theorem 1.2

Theorem 4.1. There exists a polynomial-time computable function Γ mapping each universal first-order sentence Φ to a universal first-order sentence $\Gamma(\Phi)$ over the signature of Φ expanded by a fresh binary symbol E such that:

- (1) $\operatorname{fm}(\Phi)$ has the AP if and only if $\operatorname{fm}(\Gamma(\Psi))$ has the SAP;
- (2) if τ is a common subset of the signatures of Φ_1 and Φ_2 , then

$$\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$$
 if and only if $\operatorname{fm}(\Gamma(\Phi_1))|^{\tau \cup \{E\}} \subseteq \operatorname{fm}(\Gamma(\Phi_2))|^{\tau \cup \{E\}}$.

Proof. Let ϕ be the quantifier-free part of Φ , and let σ be its signature. For two tuples \bar{x}_1, \bar{x}_2 of the same arity k we use $E(\bar{x}_1, \bar{x}_2)$ as a shortcut for the formula stating that, for every $i \in [k]$, the i-the entry of \bar{x}_1 and the i-the entry of \bar{x}_2 are contained in an E-atom. We define $\Gamma(\Phi)$ as

$$\forall x, y, z \Big(E(x, x) \land \big(E(x, y) \Rightarrow E(y, x) \big) \land \big(E(x, y) \land E(y, z) \Rightarrow E(x, z) \big)$$

$$\land \bigwedge_{X \in \sigma} \forall \bar{x}_1, \bar{x}_2 \Big(E(\bar{x}_1, \bar{x}_2) \Rightarrow \big(X(\bar{x}_1) \Leftrightarrow X(\bar{x}_2) \big) \Big)$$

$$\land \forall \bar{x} \Big(\phi(\bar{x}) \lor \bigvee_{x_1, x_2 \in \bar{x}} \big(E(x_1, x_2) \land (x_1 \neq x_2) \big) \Big).$$

We start with (1). First, suppose that $\operatorname{fm}(\Gamma(\Phi))$ has the SAP. Let $\mathfrak{B}_1, \mathfrak{B}_2$ be a one-point amalgamation diagram for $\operatorname{fm}(\Phi)$. We convert it to a one-point amalgamation diagram $\mathfrak{B}'_1, \mathfrak{B}'_2$ for $\operatorname{fm}(\Gamma(\Phi))$ by letting E interpret as the diagonal relation. Since $\operatorname{fm}(\Phi)$ has the SAP, there exists a strong amalgam \mathfrak{C}' for $\mathfrak{B}'_1, \mathfrak{B}'_2$ in $\operatorname{fm}(\Gamma(\Phi))$. By the first two lines in the definition of $\Gamma(\Phi)$, we have that $E^{\mathfrak{C}'}$ is a relational congruence on \mathfrak{C}' . Hence, if $(b_1, b_2) \in E^{\mathfrak{C}'}$ holds for the two elements $b_1 \in B_1 \setminus B_2$ and $b_2 \in B_2 \setminus B_1$, then $\mathfrak{C}'/E^{\mathfrak{C}'}$ is isomorphic to both \mathfrak{B}'_1 and \mathfrak{B}'_2 . In that case an amalgam \mathfrak{C} for $\mathfrak{B}_1, \mathfrak{B}_2$ in $\operatorname{fm}(\Phi)$ can be obtained simply by gluing b_1 and b_2 . Otherwise a strong amalgam \mathfrak{C} for $\mathfrak{B}_1, \mathfrak{B}_2$ in $\operatorname{fm}(\Phi)$ can be obtained simply by taking the σ -reduct of \mathfrak{C}' .

Next, suppose that $\operatorname{fm}(\Phi)$ has the AP. Let $\mathfrak{B}_1, \mathfrak{B}_2$ be a one-point amalgamation diagram for $\operatorname{fm}(\Gamma(\Phi))$. Suppose that there exists $b \in B_1 \cap B_2$ such that $(b,b_1) \in E^{\mathfrak{B}_1}$ or $(b,b_2) \in E^{\mathfrak{B}_2}$. Then we obtain an amalgam \mathfrak{C} by adding tuples to the relations of $\mathfrak{B}_1 \cup \mathfrak{B}_2$ so that $E^{\mathfrak{C}}$ is a relational congruence on \mathfrak{C} . In this case, it is easy to see that $\mathfrak{C} \models \Gamma(\Phi)$. Otherwise, let $\mathfrak{B}'_1 \coloneqq \mathfrak{B}_1/E^{\mathfrak{B}_1}$ and $\mathfrak{B}'_2 \coloneqq \mathfrak{B}_2/E^{\mathfrak{B}_2}$. Since $E^{\mathfrak{B}_1}$ and $E^{\mathfrak{B}_2}$ are relational congruences on \mathfrak{B}_1 and \mathfrak{B}_2 , respectively, we have that $\mathfrak{B}'_1, \mathfrak{B}'_2$ is a one-point amalgamation diagram for $\operatorname{fm}(\Phi)$. Since $\operatorname{fm}(\Phi)$ has the AP, there exists an amalgam \mathfrak{C}' for $\mathfrak{B}'_1, \mathfrak{B}'_2$ in $\operatorname{fm}(\Phi)$. Without loss of generality, $C' \subseteq B'_1 \cup B'_2$. If $C' \neq B'_1 \cup B'_2$, then \mathfrak{C}' was obtained by gluing $[b_1]_{E^{\mathfrak{B}_1}}$ and $[b_2]_{E^{\mathfrak{B}_2}}$ in $\mathfrak{B}'_1 \cup \mathfrak{B}'_2$. Then we obtain \mathfrak{C} from $\mathfrak{B}_1 \cup \mathfrak{B}_2$ by adding (b_1,b_2) and (b_2,b_1) to the relation interpreting E. Since $(b,b_1) \notin E^{\mathfrak{B}_1}$ ans $(b,b_2) \notin E^{\mathfrak{B}_2}$ for every $b \in B_1 \cap B_2$, $E^{\mathfrak{C}}$ is a relational congruence on \mathfrak{C} . Now it is easy to see that \mathfrak{C} is a strong amalgam for $\mathfrak{B}_1 \cup \mathfrak{B}_2$ by adding tuples to relations so that, for every $X \in \sigma$, we have $\overline{t} \in X^{\mathfrak{C}}$ if and only if $[\overline{t}]_{E^{\mathfrak{B}_1 \cup \mathfrak{B}_2} \in X^{\mathfrak{C}'}$. Again it is easy to see that \mathfrak{C} is a strong amalgam for $\mathfrak{B}_1, \mathfrak{B}_2$ in $\operatorname{fm}(\Phi)$.

We continue with (2). Suppose that $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$. Let $\mathfrak A$ be an arbitrary structure from $\operatorname{fm}(\Gamma(\Phi_1))|^{\tau \cup \{E\}}$, and let $\mathfrak A_1$ be an arbitrary structure in $\operatorname{fm}(\Gamma(\Phi_1))$ whose $(\tau \cup \{E\})$ -reduct equals $\mathfrak A$. We set $\mathfrak A'_1 := \mathfrak A_1/E^{\mathfrak A_1}$. Clearly $\mathfrak A'_1 \models \Phi_1$. Let $\mathfrak A'$ be the τ -reduct of $\mathfrak A'_1$. Since $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$, there exists a σ -expansion $\mathfrak A'_2$ of $\mathfrak A'$ such that $\mathfrak A'_2 \models \Phi_2$. We define the $(\sigma \cup \{E\})$ -expansion $\mathfrak A_2$ of $\mathfrak A$ so that, for every $X \in \sigma$, we have $\overline{t} \in X^{\mathfrak A_2}$ if and only if $[\overline{t}]_{E^{\mathfrak A_1}} \in X^{\mathfrak A'_2}$. It is easy

to see that $\mathfrak{A}_2 \models \Gamma(\Phi_2)$, and hence $\mathfrak{A} \in \operatorname{fm}(\Gamma(\Phi_2))|^{\tau \cup \{E\}}$. Now suppose that $\operatorname{fm}(\Gamma(\Phi_1))|^{\tau \cup \{E\}} \subseteq \operatorname{fm}(\Gamma(\Phi_2))|^{\tau \cup \{E\}}$, and let $\mathfrak{A} \in \operatorname{fm}(\Phi_1)|^{\tau}$ be arbitrary. Let \mathfrak{A}_1 be an arbitrary structure in $\operatorname{fm}(\Phi_1)$ whose τ -reduct equals \mathfrak{A} . We define \mathfrak{A}_1' as the $(\sigma \cup \{E\})$ -expansion of \mathfrak{A}_1 where E interprets as the diagonal relation. Clearly $\mathfrak{A}_1' \models \Gamma(\Phi_1)$. Let \mathfrak{A}' be the $(\tau \cup \{E\})$ -reduct of \mathfrak{A}_1' . Since $\operatorname{fm}(\Gamma(\Phi_1))|^{\tau \cup \{E\}} \subseteq \operatorname{fm}(\Gamma(\Phi_2))|^{\tau \cup \{E\}}$, there exists a $(\sigma \cup \{E\})$ -expansion \mathfrak{A}_2' of \mathfrak{A}' such that $\mathfrak{A}_2' \models \Gamma(\Phi_2)$. We define \mathfrak{A}_2 as the σ -reduct of \mathfrak{A}_2' . It is easy to see that $\mathfrak{A}_2 \models \Phi_2$, and hence $\mathfrak{A} \in \operatorname{fm}(\Phi_2)|^{\tau}$.

Example 4.2. Let \mathcal{G} be the disjoint union of countably many copies of an undirected edge. Note that \mathfrak{G} is homogeneous, but its age does not have the SAP. It is easy to see that $age(\mathfrak{G}) = fm(\Phi)$ for a universal sentence Φ stating that each model is an undirected simple graph which does not embed any connected graph on three vertices. We give an explicit construction of strong amalgams for $fm(\Gamma(\Phi))$. Let $\mathfrak{B}_1, \mathfrak{B}_2$ be a one-point amalgamation diagram for $\operatorname{fm}(\Gamma(\Phi))$. There are the following two cases to consider. First, suppose that there exists $b \in B_1 \cap B_2$ such that $(b,b_1) \in E^{\mathfrak{B}_1}$ or $(b,b_2) \in E^{\mathfrak{B}_2}$. W.l.o.g., $(b,b_1) \in E^{\mathfrak{B}_1}$. If there exists $b' \in B_1 \cap B_2$ with an edge to b_1 , then it must also have an edge to b because $E^{\mathfrak{B}_1}$ is a relational congruence on \mathfrak{B}_1 . Note that, by the construction of $\Gamma(\Phi)$ and because $E^{\mathfrak{B}_2}$ is a relational congruence on \mathfrak{B}_2 , if there is an edge from b to b_2 , then $(b,b_2) \in E^{\mathfrak{B}_2}$. In this case we obtain \mathfrak{C} from $\mathfrak{B}_1 \cup \mathfrak{B}_2$ by adding (b_1, b_2) and (b_2, b_1) to $E^{\mathfrak{B}_1 \cup \mathfrak{B}_2}$. Otherwise, we set $\mathfrak{C} := \mathfrak{B}_1 \cup \mathfrak{B}_2$. Secondly, suppose that there is no $b \in B_1 \cap B_2$ such that $(b,b_1) \in E^{\mathfrak{B}_1}$ or $(b,b_2) \in E^{\mathfrak{B}_2}$. If there exists $b \in B_1 \cap B_2$ with an edge to both b_1 and b_2 , then we obtain \mathfrak{C} from $\mathfrak{B}_1 \cup \mathfrak{B}_2$ by adding (b_1, b_2) and (b_2, b_1) to $E^{\mathfrak{B}_1 \cup \mathfrak{B}_2}$. Otherwise, we set $\mathfrak{C} := \mathfrak{B}_1 \cup \mathfrak{B}_2$.

We continue with Theorem 4.3, which was originally presented as Theorem 1.1.

Theorem 4.3 (First Inside-Out lemma). There exists a polynomial-time computable function Ω mapping each pair Φ_1, Φ_2 of universal first-order sentences and each common subset τ of their signatures to a universal sentence $\Phi := \Omega(\Phi_1, \Phi_2, \tau)$ such that the following are equivalent:

- (1) $fm(\Phi)$ has the SAP;
- (2) $fm(\Phi)$ has the AP;
- (3) $\operatorname{fm}(\Phi_1)$ has the SAP and $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$.

Proof. Let σ_1 and σ_2 be the signatures gathering the remaining symbols in Φ_1 and Φ_2 . We denote the quantifier-free parts of Φ_1 and Φ_2 by $\phi_1(\bar{x}_1)$ and $\phi_2(\bar{x}_2)$, respectively. We first introduce two fresh unary symbols L, R and a fresh binary symbol P. Then, we introduce the signature σ'_2 containing, for each $R \in \sigma_2$ of arity k, a symbol R' of arity k + 2. We set

$$\begin{split} \Phi_1' &\coloneqq \forall \bar{x}_1 \Big(\phi_1(\bar{x}_1) \vee \bigvee_{x \in \bar{x}_1} L(x) \vee R(x) \Big), \\ \Phi_2' &\coloneqq \forall y_1, y_2, \bar{x}_2 \Big(L(y_1) \wedge R(y_2) \wedge \bigwedge_{x \in \bar{x}_2} P(y_1, x) \wedge P(y_2, x) \\ &\Rightarrow \bigvee_{(x_1, x_2) \in (\bar{x}_2, y_1, y_2), \{x_1, x_2\} \neq \{y_1, y_2\}} \Big(P(x_1, x_2) \wedge P(x_2, x_1) \Big) \\ &\vee \phi_2'(\bar{x}_2, y_1, y_2) \vee \bigvee_{x \in \bar{x}_2} L(x) \vee R(x) \Big), \end{split}$$

where ϕ_2' denotes the formula obtained from ϕ_2 by replacing each σ_2 -atom $X(\bar{t})$ by the σ_2' -atom $X'(\bar{t}, y_1, y_2)$. Then we set $\Phi := \Phi_1' \wedge \Phi_2'$ (and bring Φ into prenex-normal form).

" $(1) \Rightarrow (2)$ " This direction is trivial.

"(2) \Rightarrow (3)" For the first part, let $\mathfrak{B}_1, \mathfrak{B}_2$ be an arbitrary one-point amalgamation diagram for fm(Φ_1). Consider the $(\tau \cup \sigma_1 \cup \sigma_2' \cup \{L, R, P\})$ -expansions $\mathfrak{B}_1', \mathfrak{B}_2'$ of $\mathfrak{B}_1, \mathfrak{B}_2$ by empty relations, respectively, except that $(b_1, b_1) \in P^{\mathfrak{B}_1'}$ holds for the unique element $b_1 \in B_1 \setminus B_2$. Then $\mathfrak{B}_1', \mathfrak{B}_2' \models \Phi_1'$ is inherited directly from $\mathfrak{B}_1, \mathfrak{B}_2 \models \Phi_1$. Moreover, $\mathfrak{B}_1', \mathfrak{B}_2' \models \Phi_2'$ holds because L interprets as the empty relation in both \mathfrak{B}_1' and \mathfrak{B}_2' . Hence, $\mathfrak{B}_1', \mathfrak{B}_2'$ is a one-point amalgamation diagram for fm(Φ). Since fm(Φ) has the AP, there exists an amalgam \mathfrak{C}' for \mathfrak{B}_1' and \mathfrak{B}_2' satisfying Φ . We have that \mathfrak{C} is strong because $P^{\mathfrak{B}_1'} = \{(b_1, b_1)\}$ while $P^{\mathfrak{B}_2'} = \emptyset$. Without loss of generality, $C' = B_1 \cup B_2$. We define \mathfrak{C} as the $(\tau \cup \sigma_1)$ -reduct of \mathfrak{C}' . It is easy to see that $\mathfrak{C} \models \Phi_1$ because atoms with symbols from $\sigma_2' \cup \{P\}$ do not interact with the first clause in the quantifier-free part of Φ . We conclude that fm(Φ_1) has the SAP.

For the second part, let $\mathfrak{A} \in \operatorname{fm}(\Phi_1)|^{\tau}$ be arbitrary, and let \mathfrak{A}' be an arbitrary $(\tau \cup \sigma_1)$ -expansion of \mathfrak{A} satisfying Φ_1 . We define the one-point amalgamation diagram $\mathfrak{B}_1, \mathfrak{B}_2$ for $\operatorname{fm}(\Phi)$ as follows. The domains of \mathfrak{B}_1 and \mathfrak{B}_2 are $A \cup \{b_1\}$ and $A \cup \{b_2\}$, respectively, and the relations are inherited from \mathfrak{A}' , except that additionally $b_1 \in L^{\mathfrak{B}_1}$, $b_2 \in R^{\mathfrak{B}_2}$, and, for every $a \in A$, we have $(b_1, a) \in P^{\mathfrak{B}_1}$ and $(b_2, a) \in P^{\mathfrak{B}_2}$. Since \mathfrak{A}' satisfies Φ_1 , we get that $\mathfrak{B}_1 \models \Phi'_1$. Since $L^{\mathfrak{B}_1} = \{b_1\}$ and $R^{\mathfrak{B}_1} = \emptyset$, we get that $\mathfrak{B}_1 \models \Phi'_2$. In sum, $\mathfrak{B}_1 \models \operatorname{fm}(\Phi)$. Similarly we get that $\mathfrak{B}_2 \models \operatorname{fm}(\Phi)$. Hence, $\mathfrak{B}_1, \mathfrak{B}_2$ is a one-point amalgamation diagram for $\operatorname{fm}(\Phi)$. Since $\operatorname{fm}(\Phi)$ has the AP, there exists an amalgam \mathfrak{C} for \mathfrak{B}_1 and \mathfrak{B}_2 satisfying Φ . We have that \mathfrak{C} is strong because $L^{\mathfrak{B}_1} = \{b_1\}$ while $L^{\mathfrak{B}_2} = \emptyset$. Let \mathfrak{A}'' be the $(\tau \cup \sigma_2)$ -expansion of \mathfrak{A} such that, for every $X \in \sigma_2$, $\bar{t} \in X^{\mathfrak{A}''}$ if and only if $(\bar{t}, b_1, b_2) \in X'^{\mathfrak{C}}$. Since $\mathfrak{C} \models \operatorname{fm}(\Phi)$, we have that $\mathfrak{A}'' \models \Phi_2$, and hence $\mathfrak{A} \in \operatorname{fm}(\Phi_2)|^{\tau}$. We conclude that $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$.

"(3) \Rightarrow (1)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be a one-point amalgamation diagram for fm(Φ).

First, suppose that $b_1, b_2 \notin L^{\mathfrak{B}_1} \cup R^{\mathfrak{B}_1} \cup L^{\mathfrak{B}_2} \cup R^{\mathfrak{B}_2}$. We obtain \mathfrak{B}'_1 and \mathfrak{B}'_2 from \mathfrak{B}_1 and \mathfrak{B}_2 by taking their substructures on $B_1 \setminus (L^{\mathfrak{B}_1} \cup R^{\mathfrak{B}_1})$ and $B_2 \setminus (L^{\mathfrak{B}_2} \cup R^{\mathfrak{B}_2})$, respectively. Let \mathfrak{B}''_1 and \mathfrak{B}''_2 be the $(\tau \cup \sigma_1)$ -reducts of \mathfrak{B}'_1 and \mathfrak{B}'_2 , respectively. By the definition of Φ'_1 , we get that $\mathfrak{B}''_1, \mathfrak{B}''_2 \models \Phi_1$ because $\mathfrak{B}_1, \mathfrak{B}_2 \models \Phi'_1$. Since $fm(\Phi_1)$ has the SAP, there exists a strong amalgam \mathfrak{C}' for \mathfrak{B}''_1 and \mathfrak{B}''_2 satisfying Φ_1 . Without loss of generality, we have $C' = B'_1 \cup B'_2$. We obtain \mathfrak{C} by adding tuples to the relations of $\mathfrak{B}_1 \cup \mathfrak{B}_2$ so that $(b_1, b_2), (b_2, b_1) \in P^{\mathfrak{C}}$ and, for every $X \in \tau \cup \sigma_1$, we have $\overline{t} \in X^{\mathfrak{C}}$ whenever $\overline{t} \in X^{\mathfrak{C}'}$. Note that, by the second line of the definition of Φ'_2 , the quantifier-free part of Φ'_2 is trivially true on \mathfrak{C} for all assignments to the free variables whose range contains both b_1 and b_2 .

Second, suppose that $\{b_1, b_2\} \cap (L^{\mathfrak{B}_1} \cup R^{\mathfrak{B}_2} \cup L^{\mathfrak{B}_2} \cup R^{\mathfrak{B}_2}) \neq \emptyset$. We obtain \mathfrak{A} by first taking the substructure of \mathfrak{B}_1 (or equivalently \mathfrak{B}_2) on $(B_1 \cap B_2) \setminus (L^{\mathfrak{B}_1} \cup R^{\mathfrak{B}_1})$. Since $\mathfrak{B}_1 \models \Phi$, we have $\mathfrak{A} \in \text{fm}(\Phi_1)$. Let \mathfrak{A}' be the τ -reduct of \mathfrak{A} . Since $\text{fm}(\Phi_1)|^{\tau} \subseteq \text{fm}(\Phi_2)|^{\tau}$, there exists a $(\tau \cup \sigma_2)$ -expansion \mathfrak{A}'' of \mathfrak{A}' satisfying Φ_2 . We obtain \mathfrak{C} by adding tuples to the relations of $\mathfrak{B}_1 \cup \mathfrak{B}_2$ so that $(b_1, b_2), (b_2, b_1) \in P^{\mathfrak{C}}$ and, for every $X \in \tau \cup \sigma_2$, we have $(\bar{t}, b_1, b_2), (\bar{t}, b_2, b_1) \in X'^{\mathfrak{C}}$ whenever $\bar{t} \in X^{\mathfrak{A}'}$. By the definition of Φ_1' , the quantifier-free part of Φ_1' is trivially true on \mathfrak{C} for all assignments to the free variables whose range contains both b_1 and b_2 . Also, by the second line

of the definition of Φ'_2 , the quantifier-free part of Φ'_2 is trivially true on \mathfrak{C} for all assignments to the free variables whose range contains both b_1 and b_2 but it is not the case that y_1 is set equal b_1 and y_2 is set equal b_2 or vice versa.

It is now easy to check that $\mathfrak{C} \models \Phi$ and that the inclusion maps from \mathfrak{B}_1 and \mathfrak{B}_2 to \mathfrak{C} are embeddings. Hence $fm(\Phi)$ has the SAP.

Remark 4.4. If \mathfrak{B} is a structure whose age equals $fm(\Phi)$ for Φ as in Theorem 4.3, then $CSP(\mathfrak{B})$ is trivial. The reason is that $age(\mathfrak{B})$ contains the structure \mathfrak{C} whose domain has a single element and where each relation contains one tuple. Every structure over the signature of Φ has a homomorphism to \mathfrak{C} .

Finally, we prove Theorem 4.5, which was originally presented as Theorem 1.6.

Theorem 4.5 (Second Inside-Out lemma). There exists a polynomial-time computable function Δ mapping each universal first-order sentence Φ over a finite relational signature ρ to a pair of universal first-order sentences $\Phi_1 := \Delta(\Phi)_1$ and $\Phi_2 := \Delta(\Phi)_2$ over some finite relational signatures τ and σ , respectively, such that $\rho \subseteq \tau \subseteq \sigma$ and the following are equivalent:

- (1) $fm(\Phi)$ has the SAP;
- (2) $fm(\Phi_1)$ has the SAP;
- (3) $fm(\Phi_2)$ has the SAP;
- (4) $\operatorname{fm}(\Phi_1) \subseteq \operatorname{fm}(\Phi_2)|_{\tau}$.

Proof. Let $\phi(\bar{x})$ be the quantifier-free part of Φ . We first define the signatures τ and σ . The signature τ consists of all symbols from ρ together with two fresh unary symbols L and R. We write MIXED (\bar{x}) as a shortcut for the formula

$$\bigvee_{x_1, x_2 \in \bar{x}} \left(\neg L(x_1) \land \neg R(x_2) \right).$$

The signature σ contains all symbols from τ and additionally, for every k-ary symbol $X \in \rho$, a fresh k-ary symbol X'. We set

$$\begin{split} \Phi_1 &\coloneqq \forall \bar{x} \Big(\phi(\bar{x}) \vee \text{MIXED}(\bar{x}) \Big) \wedge \forall x \Big(L(x) \vee R(x) \Big), \\ \Phi_2 &\coloneqq \bigwedge_{X \in \rho} \forall \bar{y} \Big(\big(X(\bar{y}) \Leftrightarrow X'(\bar{y}) \big) \vee \big(\text{MIXED}(\bar{y}) \wedge \neg X(\bar{y}) \big) \\ &\wedge \forall \bar{x}. \, \phi'(\bar{x}) \wedge \forall x \Big(L(x) \vee R(x) \Big), \end{split}$$

where ϕ' is obtained from ϕ by replacing each atomic ρ -formula $X(\bar{x})$ with $X'(\bar{x})$.

- "(1) \Rightarrow (4)" Let $\mathfrak{A} \in \operatorname{fm}(\Phi_1)$ be arbitrary, and let \mathfrak{A}' be the ρ -reduct of \mathfrak{A} . We obtain the amalgamation diagram \mathfrak{B}_1 , \mathfrak{B}_2 for $\operatorname{fm}(\Phi)$ by defining \mathfrak{B}_1 and \mathfrak{B}_2 as the substructures of \mathfrak{A}' on $L^{\mathfrak{A}}$ and $R^{\mathfrak{A}}$, respectively. Since $\operatorname{fm}(\Phi)$ has the SAP, there exists a strong amalgam \mathfrak{C} for \mathfrak{B}_1 and \mathfrak{B}_2 satisfying Φ . We define the σ -expansion \mathfrak{A}'' of \mathfrak{A} such that, for every $X \in \rho$, we have $X^{\mathfrak{A}''} := X'^{\mathfrak{C}}$. Clearly $\mathfrak{A}'' \models \Phi_2$, and hence $\mathfrak{A} \in \operatorname{fm}(\Phi_2)|_{\tau}$.
- "(4) \Rightarrow (1)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an amalgamation diagram for $\operatorname{fm}(\Phi)$. We obtain the τ -structure \mathfrak{A} from $\mathfrak{B}_1 \cup \mathfrak{B}_2$ by setting $L^{\mathfrak{A}} := B_1$ and $R^{\mathfrak{A}} := B_2$. Since $\operatorname{fm}(\Phi_1) \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$, there exists a σ -expansion \mathfrak{A}' of \mathfrak{A} satisfying Φ_2 . We obtain a strong amalgam \mathfrak{C} for \mathfrak{B}_1 and \mathfrak{B}_2 in $\operatorname{fm}(\Phi)$ by taking the ρ -reduct of \mathfrak{A}' .
- "(2) \Rightarrow (1)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an amalgamation diagram for $\operatorname{fm}(\Phi)$. Consider the τ -expansions $\mathfrak{B}'_1, \mathfrak{B}'_2$ where L and R both interpret as the full unary relation. Clearly, $\mathfrak{B}'_1, \mathfrak{B}'_2$ is an amalgamation diagram for $\operatorname{fm}(\Phi_1)$. Since $\operatorname{fm}(\Phi_1)$ has the SAP, there

exists an amalgam \mathfrak{C}' for \mathfrak{B}'_1 and \mathfrak{B}'_2 satisfying Φ_1 . Then the ρ -reduct \mathfrak{C} of \mathfrak{C}' is a strong amalgam for \mathfrak{B}_1 and \mathfrak{B}_2 in $fm(\Phi)$.

- "(1) \Rightarrow (2)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an amalgamation diagram for $\operatorname{fm}(\Phi_1)$. We obtain two amalgamation diagrams $\mathfrak{B}_1', \mathfrak{B}_2'$ and $\mathfrak{B}_1'', \mathfrak{B}_2''$ for $\operatorname{fm}(\Phi)$ by taking the ρ -reducts of the substructures of $\mathfrak{B}_1, \mathfrak{B}_2$ on the subsets defined by L and R, respectively. Since $\operatorname{fm}(\Phi)$ has the SAP, there exist strong amalgams \mathfrak{C}' and \mathfrak{C}'' for the two amalgamation diagrams above. Without loss of generality, we have $C' = B_1' \cup B_2'$ and $C'' = B_1'' \cup B_2''$. We define \mathfrak{C} as the τ -expansion of $\mathfrak{C}' \cup \mathfrak{C}''$ where $L^{\mathfrak{C}} = B_1' \cup B_2'$ and $R^{\mathfrak{C}} = B_1'' \cup B_2''$. It is easy to check that \mathfrak{C} is a strong amalgam for $\mathfrak{B}_1, \mathfrak{B}_2$ in $\operatorname{fm}(\Phi_1)$.
- "(3) \Rightarrow (1)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an amalgamation diagram for $\operatorname{fm}(\Phi)$. Consider the σ -expansions $\mathfrak{B}'_1, \mathfrak{B}'_2$ where L and R both interpret as the full unary relation and the remaining symbols in σ interpret as their counterparts from ρ . Clearly, $\mathfrak{B}'_1, \mathfrak{B}'_2$ is an amalgamation diagram for $\operatorname{fm}(\Phi_2)$. Since $\operatorname{fm}(\Phi_2)$ has the SAP, there exists a strong amalgam \mathfrak{C}' for \mathfrak{B}'_1 and \mathfrak{B}'_2 satisfying Φ_2 . Then the ρ -reduct \mathfrak{C} of \mathfrak{C}' is a strong amalgam for \mathfrak{B}_1 and \mathfrak{B}_2 in $\operatorname{fm}(\Phi)$.
- "(1) \Rightarrow (3)" Let $\mathfrak{B}_1, \mathfrak{B}_2$ be an amalgamation diagram for $\operatorname{fm}(\Phi_2)$. Let $\mathfrak{B}'_1, \mathfrak{B}'_2$ be the ρ -structures with domains B'_1, B'_2 , respectively, and relations $X^{\mathfrak{B}'_i} := X'^{\mathfrak{B}'_i}$ ($i \in [2]$). By construction, $\mathfrak{B}''_1, \mathfrak{B}''_2$ is an amalgamation diagram for $\operatorname{fm}(\Phi)$. Since $\operatorname{fm}(\Phi)$ has the SAP, there exists a strong amalgam \mathfrak{C}' for \mathfrak{B}'_1 and \mathfrak{B}'_2 satisfying Φ . We obtain a strong amalgam \mathfrak{C} for \mathfrak{B}_1 and \mathfrak{B}_2 satisfying Φ by lifting the ρ -relations of \mathfrak{C}' to their σ -counterparts.

5. Hardness of Containment

The present section is fully devoted to the proof of the following theorem, originally presented in the introduction as Theorem 1.4.

Theorem 5.1. The question whether $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$ holds for a given pair of universal first-order sentences Φ_1, Φ_2 having τ as a common subset of their signatures and such that $\operatorname{fm}(\Phi_1)$ and $\operatorname{fm}(\Phi_2)$ both have the FAP is 2NEXPTIME-hard.

The proof of the theorem is by a polynomial-time reduction from the double-exponential square tiling problem. Consider the signature $\hat{\tau}$ consisting of the two binary symbols \hat{S}_1 and \hat{S}_2 . For every natural number $n \geq 1$, we define the the $\hat{\tau}$ -structure $\hat{\mathfrak{Q}}_n$ as follows. The domain is $[n]^2$, and the relations are

$$\begin{split} \hat{S}_{1}^{\hat{\mathfrak{Q}}_{n}} &:= \{ ((i,j), (i+1,j)) \mid i \in [n-1], j \in [n] \}, \\ \hat{S}_{2}^{\hat{\mathfrak{Q}}_{n}} &:= \{ ((i,j), (i,j+1)) \mid i \in [n], j \in [n-1] \}. \end{split}$$

The NP-complete square tiling problem [53] (see also [57]) can be stated as follows:

INSTANCE: A natural number n, a finite $\hat{\tau}$ -structure \mathfrak{T} , and $t \in T$.

QUESTION: Does there exist a homomorphism $h: \hat{\mathfrak{Q}}_n \to \mathfrak{T}$ with h(1,1) = t?

One can further increase the complexity by allowing a succinct encoding of the square. The input remains the same but now we ask for a square tiling with 2^{2^n} rows and columns, i.e., a homomorphism $h: \hat{\mathfrak{Q}}_{2^{2^n}} \to \mathfrak{T}$. Analogously to the natural complete problems based on Turing machines, this yields a decision problem which is complete for the complexity class 2NEXPTIME [53].

Proof of Theorem 5.1. Consider the signature τ consisting of the binary symbol O, the two 4-ary symbols S_1, S_2 , and the two (n+3)-ary symbols C_1, C_2 . The nodes of

the $2^{2^n} \times 2^{2^n}$ -grid will be encoded on pairs of first-order variables, but we remark in advance that the encoding is not coordinate-wise; the first (second) entry in a pair of first-order variables does not specify the first (second) coordinate of a node. The symbol O represents the initial pair (1,1), and the symbols S_1, S_2 represent vertical and horizontal successor relations, similarly as \hat{S}_1, \hat{S}_2 . The symbols C_1, C_2 will be used to encode functions $f: \{x_1, x_2\}^n \to \{x_1, x_2\}$ indexed by pairs (x_1, x_2) ; an atom $C_k(x_1, x_2, b_1, \ldots, b_n, b)$ with $b, b_1, \ldots, b_n \in \{x_1, x_n\}$ is to be interpreted as "the function value at (b_1, \ldots, b_n) is b." Note that there are 2^{2^n} -many such functions.

The universal first-order sentence Φ_1 verifies the 2^{2^n} -grid. In addition to all symbols from τ , its signature contains 2n-many symbols $\{B_k^i \mid i \in [n], k \in [2]\}$ of arity 6 and the 6 symbols $\{I_k, U_k, D_k \mid k \in [2]\}$ of arity 2(n+2). The purpose of these symbols is explained below. Before we proceed further, we introduce $SUCC(x_1, x_2, b_1, \ldots, b_n, c_1, \ldots, c_n)$ as a shortcut for the formula

$$\left(\bigvee_{j\in[n]}\bigwedge_{i\in[j-1]}\left(b_i=x_1\wedge c_i=x_1\vee b_i=x_2\wedge c_i=x_2\right)\right)$$
$$\wedge\left(b_j=x_1\wedge c_j=x_2\right)\wedge\left(\bigwedge_{i\in[n]\backslash[j]}\left(b_i=x_2\wedge c_i=x_1\right)\right).$$

The intuition is that, in terms of equalities with x_1 and x_2 , the tuple (c_1, \ldots, c_n) is a successor of (b_1, \ldots, b_n) when both tuples are read from left to right. For $k \in [2]$, we include the following (universally quantified) implications as conjuncts in Φ_1 .

We start with a conjunct ensuring that C_k specifies a function:

$$C_k(x_1, x_2, b_1, \dots, b_n, x_1) \wedge C_k(x_1, x_2, b_1, \dots, b_n, x_2)$$
$$\wedge \bigwedge_{i \in [n]} \left(b_i = x_1 \vee b_i = x_2 \right) \Rightarrow \bot.$$

Next, we include a conjunct ensuring that $O(x_1, x_2)$ encodes the initial pair (1,1):

$$O(x_1, x_2) \land \bigwedge_{i \in [n]} (b_i = x_1 \lor b_i = x_2) \Rightarrow C_k(x_1, x_2, b_1, \dots, b_n, x_1).$$

From the implication above, the encoding of the nodes of the grid becomes apparent; in each pair (x_1, x_2) , the first (second) entry serves as the bit zero (one), respectively, and the value in the k-th coordinate of the node is determined by the formula

$$\bigwedge_{b_1,\ldots,b_n\in\{x_1,x_2\}} C_k(b_1,\ldots,b_n,f(b_1,\ldots,b_n)),$$

for some function $f: \{x_1, x_2\}^n \to \{x_1, x_2\}.$

What is left is to ensure that every atom $S_k(x_1^1, x_2^1, x_1^2, x_2^2)$ represents a horizontal or vertical successor pair of grid nodes. This is where having existentially quantified second-order variables becomes essential. Recall that above we used SUCC as the successor predicate for numbers in 2^n encoded in binary as n-tuples over $\{x_1, x_2\}$. We proceed similarly for numbers in 2^{2^n} , but this time we encode them using 2^n -tuples of 2(n+1)-ary atomic formulas satisfied by the quadruple $(x_1^1, x_2^1, x_1^2, x_2^2)$.

We use the symbol I_k to mark the initial (synchronized) part of a pair of 2^n -tuples, i.e., where the function values in terms of C_k -atoms coincide. Furthermore, we use the symbol U_k for the upwards flipped bit, and the symbol D_k for the downwards flipped bit (carryover). First, we include two conjuncts enforcing the choice of the

upwards flipped bit $(k \in [2])$:

$$S_{k}(x_{1}^{1}, x_{2}^{1}, x_{1}^{2}, x_{2}^{2}) \Rightarrow I_{3-k}(x_{1}^{1}, x_{2}^{1}, x_{2}^{1}, \dots, x_{2}^{1}, x_{1}^{2}, x_{2}^{2}, x_{2}^{2}, \dots, x_{2}^{2})$$

$$\wedge \bigwedge_{i \in [n]} B_{k}^{i}(x_{1}^{1}, x_{2}^{1}, x_{1}^{1}, x_{1}^{2}, x_{2}^{2}, x_{1}^{2}) \vee B_{k}^{i}(x_{1}^{1}, x_{2}^{1}, x_{2}^{1}, x_{2}^{2}, x_{2}^{2}),$$

$$\bigwedge_{i \in [n]} \left(b_{i}^{1} = x_{1}^{1} \wedge b_{i}^{2} = x_{1}^{2} \vee b_{i}^{1} = x_{2}^{1} \wedge b_{i}^{2} = x_{2}^{2} \right) \wedge B_{k}^{i}(x_{1}^{1}, x_{2}^{1}, b_{i}^{1}, x_{1}^{2}, x_{2}^{2}, b_{i}^{2})$$

$$\Rightarrow U_{k}(x_{1}^{1}, x_{2}^{1}, b_{1}^{1}, \dots, b_{n}^{1}, x_{1}^{2}, x_{2}^{2}, b_{1}^{2}, \dots, b_{n}^{2}).$$

Then we include two conjuncts for the propagation of the synchronized part and the downwards flipped bit $(k \in [2])$:

$$\begin{split} \left(U_k(x_1^1, x_2^1, b_1^1, \dots, b_n^1, x_1^2, x_2^2, b_1^2, \dots, b_n^2) \vee I_k(x_1^1, x_2^1, b_1^1, \dots, b_n^1, x_1^2, x_2^2, b_1^2, \dots, b_n^2)\right) \\ &\wedge \operatorname{SUCC}(x_1^1, x_2^1, b_1^1, \dots, b_n^1, c_1^1, \dots, c_n^1) \wedge \operatorname{SUCC}(x_1^2, x_2^2, b_1^2, \dots, b_n^2, c_1^2, \dots, c_n^2) \\ &\Rightarrow I_k(x_1^1, x_2^1, c_1^1, \dots, c_n^1, x_1^2, x_2^2, c_1^2, \dots, c_n^2), \\ \left(U_k(x_1^1, x_2^1, b_1^1, \dots, b_n^1, x_1^2, x_2^2, b_1^2, \dots, b_n^2) \vee D_k(x_1^1, x_2^1, b_1^1, \dots, b_n^1, x_1^2, x_2^2, b_1^2, \dots, b_n^2)\right) \\ &\wedge \operatorname{SUCC}(x_1^1, x_2^1, c_1^1, \dots, c_n^1, b_1^1, \dots, b_n^1) \wedge \operatorname{SUCC}(x_1^2, x_2^2, c_1^2, \dots, c_n^2, b_1^2, \dots, b_n^2) \\ &\Rightarrow D_k(x_1^1, x_2^1, c_1^1, \dots, c_n^1, x_1^2, x_2^2, c_1^2, \dots, c_n^2). \end{split}$$

Finally, the following three conjuncts provide interpretation for $\{U_k, I_K, D_k\}$ -atoms in terms of the functions encoded by the C_k -atoms $(k \in [2])$:

$$\begin{split} U_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_1^2,x_2^2,b_1^2,\dots,b_n^2) &\Rightarrow \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_1^1)\right. \\ &\qquad \qquad \wedge C_k(x_1^2,x_2^2,b_1^2,\dots,b_n^2,x_2^2) \Big), \\ \\ I_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_1^2,x_2^2,b_1^2,\dots,b_n^2) &\Rightarrow \left(\left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_1^1)\right. \\ &\qquad \qquad \wedge C_k(x_1^2,x_2^2,b_1^2,\dots,b_n^2,x_1^2) \Big) \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1)\right. \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1)\right. \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1)\right. \\ \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1)\right) \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1)\right. \\ &\qquad \qquad \wedge \left(C_k(x_1^1,x_2^1,b_1^1,\dots,b_n^1,x_2^1,x_2^2,b_1^2,\dots,b_n^2)\right) \Big), \end{split}$$

The universal first-order sentence Φ_2 verifies the tile tiling. In addition to τ , its signature contains the binary symbol T_a for every $a \in T$. For $k \in [2]$, we include the following (universally quantified) implications as conjuncts in Φ_2 . First, we include the implication $O(x,y) \Rightarrow T_t(x,y)$, which ensures that the initial pair (1,1) is assigned the tile $t \in T$. Next, we include implications ensuring the horizontal and vertical consistency of the tiling $(k \in [2])$:

$$S_k(x_1^1, x_2^1, x_1^2, x_2^2) \Rightarrow \bigvee_{(a,b) \in \hat{S}_{\pi}^{r_{\pi}}} T_a(x_1^1, x_2^1) \wedge T_b(x_1^2, x_2^2),$$

and $T_a(x_1, x_2) \wedge T_b(x_1, x_2) \Rightarrow \bot$ for every pair $a, b \in T$ with $a \neq b$.

Note that every implication in the quantifier-free part of Φ_1 or Φ_2 is equivalent to a conjunction of (exponentially many) complete clauses. It follows that $\operatorname{fm}(\Phi_1)$ and $\operatorname{fm}(\Phi_2)$ both have the FAP.

" \Rightarrow " If there exists a homomorphism $h: \hat{\mathfrak{Q}}_{2^{2^n}} \to \mathfrak{T}$ satisfying h(1,1) = t, then $fm(\Phi_1) \subseteq fm(\Phi_2)$ is witnessed by adding (x,y) to T_a if and only if (x,y) encodes (i,j) in terms of C_k -atoms and h(i,j) = a.

" \Leftarrow " Let $\mathfrak{Q}_{2^{2^n}}$ be the τ -structure with domain $2^{2^n} \times 2^{2^n}$ such that the C_k -atoms satisfied by (i,j) correctly encode its value and the S_k -atoms correctly determine the horizontal and vertical successor relations. By construction, $\mathfrak{Q}_{2^{2^n}} \in \text{fm}(\Phi_1)$. Also, by construction, if $\mathfrak{Q}_{2^{2^n}} \in \text{fm}(\Phi_2)$, then there exists a homomorphism $h: \hat{\mathfrak{Q}}_{2^{2^n}} \to \mathfrak{T}$ satisfying h(1,1)=t.

Remark 5.2. With a slight modification, the proof of Theorem 5.1 can be used to prove NEXPTIME-hardness for the case where inputs are specified by sets of bounds instead of universal first-order sentences. The idea is as follows.

As stated in the proof of Theorem 5.1, each implication in Φ_1 or Φ_2 is equivalent to a conjunction of exponentially many complete clauses. By instantiating the equalities in the premises of the implications and imposing some additional equalities which were not relevant for the original construction, the number of free variables in the the said complete clauses can be bounded by four. Moreover, the tiles can be stored in binary, e.g., using a single symbol of arity $\lceil \log_2 |T| \rceil + 2$ instead of |T|-many symbols of arity 2. Now, on input (n, \mathfrak{T}, t) we compute Φ_1 and Φ_2 for the input $(\lceil \log_2 n \rceil, \mathfrak{T}, t)$ with the modification described above. We then define \mathcal{N}_1 and \mathcal{N}_2 as the sets of all σ_1 - and σ_2 -structures with domain [i] for $i \in [4]$ which do not satisfy Φ_1 and Φ_2 , respectively. By applying basic logarithm laws, we conclude that the sizes of \mathcal{N}_1 and \mathcal{N}_2 are polynomial in the size of the input to the tiling problem.

Due to the changes in the input, only the single-exponential square tiling problem can be represented in this way. The above idea also works in combination with Theorem 4.3, but the maximal size of a bound increases to 6.

6. Amalgamation through Canonisation

The present section is fully devoted to the proof of the following theorem, originally presented in the introduction as Theorem 1.7.

Theorem 6.1. There exists a non-deterministic exponential-time reduction from the classification (decision) problem for countable finitely bounded homogeneous structures to the (search) problem of finding a finitely bounded Ramsey expansion for a finitely bounded class with the SAP.

A homogeneous structure is called Ramsey if its age has the Ramsey property (RP). The precise definition of the Ramsey property is not essential to the present article and is therefore omitted. Let $\mathfrak A$ and $\mathfrak B$ be two structures. A function $f\colon A\to B$ is called canonical from $\mathfrak A$ to $\mathfrak B$ if, for every $m\in\mathbb N$, the componentwise action of f induces a well-defined function from the orbits of m-tuples in $\mathfrak A$ to the orbits of m-tuples in $\mathfrak B$. Below we state a simplified version of the canonisation lemma.

Theorem 6.2 ([23, 21]). Let \mathfrak{A} be an homogeneous Ramsey structure and let \mathfrak{B} be ω -categorical. If there exists an embedding from a reduct of \mathfrak{A} to a reduct of \mathfrak{B} , then there also exists such an embedding that is canonical from \mathfrak{A} to \mathfrak{B} .

Proof of Theorem 6.1. Recall the functions Γ and Δ from Theorem 4.1 and Theorem 4.5. For a given universal first-order sentence Φ and $i \in [2]$, we set $\Phi_i := \Delta_i(\Gamma(\Phi))$, and we choose τ as in Theorem 4.5. Suppose that there exists a function Θ mapping each universal first-order τ -sentence Φ to a universal first-order sentence

 $\Theta(\Phi)$ such that $\operatorname{fm}(\Phi) = \operatorname{fm}(\Theta(\Phi))|^{\tau}$ and $\operatorname{fm}(\Theta(\Phi))$ has the RP whenever $\operatorname{fm}(\Phi)$ has the SAP. For brevity, we write $\Phi_1^+ := \Theta(\Phi_1)$ and $\Phi_2^+ := \Theta(\Phi_2)$.

Let n be the largest arity of a symbol from the signature of Φ_1^+ or Φ_2^+ . For $i \in [2]$, we define $\operatorname{colors}(\Phi_i^+)$ as the set of all structures in $\operatorname{fm}(\Phi_i^+)$ with domain [n]. We call a mapping $\xi \colon \operatorname{colors}(\Phi_1^+) \to \operatorname{colors}(\Phi_2^+)$ a τ -recolouring from Φ_1^+ to Φ_2^+ if, for every $\mathfrak{C} \in \operatorname{colors}(\Phi_1^+)$, the τ -reducts of \mathfrak{C} and $\xi(\mathfrak{C})$ are identical, and the following extension ξ' is a well-defined mapping from $\operatorname{fm}(\Phi_1^+)$ to $\operatorname{fm}(\Phi_2^+)$: for every $\mathfrak{A} \in \operatorname{fm}(\Phi_1^+)$, the structure $\xi'(\mathfrak{A})$ on the same domain as \mathfrak{A} is obtained by replacing every isomorphic copy of $\mathfrak{C} \in \operatorname{colors}(\Phi_1^+)$ by an isomorphic copy of $\xi(\mathfrak{C}) \in \operatorname{colors}(\Phi_2^+)$. It is not hard to see that the existence of a τ -recolouring from Φ_1^+ to Φ_2^+ is decidable in non-deterministic exponential time in the combined size of Φ_1^+ and Φ_2^+ . We show that, under the assumption that $\operatorname{fm}(\Phi_1^+)$ and $\operatorname{fm}(\Phi_2^+)$ both have the RP, $\operatorname{fm}(\Phi_1)|^\tau \subseteq \operatorname{fm}(\Phi_2)|^\tau$ if and only if there is a τ -recolouring from Φ_1^+ to Φ_2^+ . This finishes the proof of the theorem.

Clearly, if there exists a τ -recolouring from Φ_1^+ to Φ_2^+ , then $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$. Indeed, for every $\mathfrak{A} \in \operatorname{fm}(\Phi_1)|^{\tau}$, we select an arbitrary expansion $\mathfrak{A}' \in \operatorname{fm}(\Phi_1^+)$, and then $\xi'(\mathfrak{A}') \in \operatorname{fm}(\Phi_2^+)$ witnesses $\mathfrak{A} \in \operatorname{fm}(\Phi_2)|_{\tau}$. The converse direction does not hold in general, but we will see that it does in our case. Suppose that $\operatorname{fm}(\Phi_1)|^{\tau} \subseteq \operatorname{fm}(\Phi_2)|^{\tau}$. By Theorem 3.1, for both $i \in [2]$, there exists a homogeneous structure \mathfrak{F}_i satisfying $age(\mathfrak{F}_i) = fm(\Phi_i^+)$. By compactness (e.g. König's tree lemma), there exists an embedding $e: \mathfrak{F}_1|^{\tau} \hookrightarrow \mathfrak{F}_2|^{\tau}$. By Theorem 6.2, we may assume that e is canonical from $\operatorname{Aut}(\mathfrak{F}_1)$ to $\operatorname{Aut}(\mathfrak{F}_2)$. It follows directly from the homogeneity of \mathfrak{F}_1 that e induces a well-defined mapping from $\operatorname{colors}(\Phi_1^+)$ to $\operatorname{colors}(\Phi_2^+)$ which is a τ -recolouring from Φ_1^+ to Φ_2^+ . More specifically, for every tuple \bar{t} in a relation of \mathfrak{F}_2 , the substructure of \mathfrak{F}_1 on $e^{-1}(\bar{t})$ is represented by some $\mathfrak{C} \in \operatorname{colors}(\Phi_1^+)$ up to isomorphism. Let \bar{t}' be any tuple over \mathfrak{F}_2 of the same arity as \bar{t} such that the function sending the *i*-th entry in $e^{-1}(\bar{t})$ to the *i*-th entry in $e^{-1}(\bar{t}')$ is an isomorphism between substructures of \mathfrak{F}_1 . By the homogeneity of \mathfrak{F}_1 , there exists an automorphism α of \mathfrak{F}_1 with $\alpha(e^{-1}(\bar{t})) = e^{-1}(\bar{t}')$. By the canonicity of e, there exists an automorphism β of \mathfrak{F}_2 such that $\bar{t}' = e(e^{-1}(\bar{t}')) = e(\alpha(e^{-1}(\bar{t}))) = \beta(e(e^{-1}(\bar{t}))) = \beta(\bar{t})$. This means that the assignment of a color $\xi(\mathfrak{C}) \in \operatorname{colors}(\Phi_2^+)$ based on e is indeed well-defined.

7. Undecidability of Homogenizability

The present section is fully devoted to the proof of the following theorem, originally presented in the introduction as Theorem 1.8.

Theorem 7.1. For a given a Datalog sentence Φ over a signature containing at most binary relation symbols, it is undecidable whether

- (1) Φ is logically equivalent to a monadic Datalog sentence, or Φ is not even logically equivalent to any GMSNP sentence;
- (2) Whether simultaneously $fm(\Phi)$ is:
 - the CSP of a finite structure,
 - the age of a finitely-bounded homogenizable structure,
 - the age of a reduct of a finitely bounded homogeneous Ramsey structure, or $fm(\Phi)$ is not even the CSP or the age of any ω -categorical structure.

As mentioned in Section 1.7, the logic GMSNP enjoys similar model-theoretic properties as MMSNP. We will use some of these properties (Theorem 7.2) in the proof of Theorem 7.1. As mentioned in the previous subsection, the definition of

the Ramsey property is not essential to the present article and is therefore omitted. The following theorem was proved in [17], except that the authors did not fully utilize the auxiliary result from [43] which guarantees the existence of a Ramsey expansion by a generic linear order.

Theorem 7.2 ([17]). Every GMSNP sentence is equivalent to a finite disjunction of connected GMSNP sentences. Every connected GMSNP sentence defines the age and the CSP of a reduct of a finitely bounded homogeneous Ramsey structure.

As usual, the *Kleene plus* and the *Kleene star* of a finite set of symbols Σ , denoted by Σ^+ and Σ^* , are the sets of all finite words over Σ of lengths ≥ 1 and ≥ 0 , respectively.

A context-free grammar (CFG) is a 4-tuple $\mathscr{G} = (N, \Sigma, P, S)$ where N is a finite set of non-terminal symbols, Σ is a finite set of terminal symbols, Σ is a finite set of production rules of the form $A \to w$ where $A \in N$ and $w \in (N \cup \Sigma)^+$, $S \in N$ is the start symbol. For $u, v \in (N \cup \Sigma)^+$ we write $u \to_G v$ if there are $x, y \in (N \cup \Sigma)^+$ and $(A \to w) \in P$ such that u = xAy and v = xwy. The language of \mathscr{G} is $L(\mathscr{G}) := \{w \in \Sigma^+ \mid S \to_{\mathscr{G}}^* w\}$, where $\to_{\mathscr{G}}^*$ denotes the transitive closure of $\to_{\mathscr{G}}$. Note that with this definition the empty word, i.e., the word ϵ of length 0, can never be an element of $L(\mathscr{G})$; some authors use a modified definition that also allows rules that derive ϵ , but for our purposes the difference is not essential. A context-free grammar is called (left-)regular if its production rules are always of the form $A \to a$ or $A \to Ba$ for non-terminal symbols A, B and a terminal symbol a. For a finite set Σ , we call a set $L \subseteq \Sigma^+$ regular if it is the language of a regular grammar with terminal symbols Σ .

Example 7.3. Consider the CFG $\mathcal G$ with a single terminal symbol a, non-terminal symbols S,A,B,C, and production rules $S\to a$, $S\to aa$, $S\to a$

The regularity problem for context-free languages can be stated as follows:

INSTANCE: A CFG \mathcal{G} . QUESTION: Is $L(\mathcal{G})$ regular?

It is well-known that this problem is undecidable, see, e.g., Theorem 6.6.6 in [54].

A deterministic finite automaton (DFA) is a 5-tuple $\mathcal{A} = (Q, \Sigma, \delta, q_0, F)$ where Q is a finite set of states, Σ is a finite set of input symbols, $\delta: Q \times \Sigma \to Q$ is a transition function, $q_0 \in Q$ is a distinguished starting state, and $F \subseteq Q$ is a distinguished set of final states. The language of \mathcal{A} is $L(\mathcal{A}) := \{a_1 \dots a_n \in \Sigma^+ \mid \delta(a_n, \dots \delta(a_2, \delta(a_1, q_0)) \dots) \in F\}$. Note that, as in the case of CFGs, the empty word ϵ can never be an element of $L(\mathcal{A})$ according to the present definition.

Let Σ be a finite set of symbols and L a subset of Σ^+ . The *Myhill-Nerode* equivalence relation on Σ^* , denoted by \sim_L , is defined by $w_1 \sim_L w_2$ if there is no $w \in \Sigma^*$ such that $|\{w_1w, w_2w\} \cap L| = 1$. The following correspondence is well-known.

Theorem 7.4 (Myhill-Nerode). For every finite set Σ and every $L \subseteq \Sigma^+$, the following are equivalent:

- L is regular;
- L is accepted by a DFA;
- \sim_L has finitely many classes.

Example 7.5. $\{a^mb^n \mid m, n \ge 1\}$ is regular while $\{a^nb^n \mid n \ge 1\}$ is not.

Let $\mathcal{G} = (N, \Sigma, P, S)$ be a CFG. The signature τ_{Σ} consists of the unary symbols I, T and the binary symbols R_a for every $a \in \Sigma$, and the signature τ_N consists of a binary symbol R_a for every element $a \in N$. For $a_1 \dots a_n \in \Sigma^+$, we set

$$\phi_{a_1...a_n}(x_1,...,x_{n+1}) := \bigwedge_{i \in [n]} R_{a_i}(x_i,x_{i+1}).$$

Let $\Phi_{\mathcal{E}}^+$ be the universal Horn sentence over the signature $\tau_{\Sigma} \cup \tau_N$ whose quantifier-free part contains, for every $(A, w) \in P$, the Horn clause

(2)
$$\phi_w(x_1, \dots, x_{|w|+1}) \Rightarrow R_A(x_1, x_{|w|+1}),$$

and additionally the Horn clause

(3)
$$I(x_1) \wedge R_S(x_1, x_2) \wedge T(x_2) \Rightarrow \bot.$$

Then $\Phi_{\mathcal{G}}$ is the connected Datalog sentence obtained from $\Phi_{\mathcal{G}}^+$ by existentially quantifying all symbols from τ_N upfront. This encoding of CFGs into Datalog programs is standard (Exercise 12.26 in [1]), and the correspondence provided by the next lemma can be shown via a straightforward induction. For a proof, we refer the reader to [25].

Lemma 7.6. For every τ_{Σ} -structure \mathfrak{A} , the following are equivalent:

- $\mathfrak{A} \models \Phi_{\mathscr{C}}$;
- for every $w \in L(\mathcal{G})$,

$$\mathfrak{A} \models \forall x_1, \dots, x_{|w|+1} \Big(I(x_1) \land \phi_w(x_1, \dots, x_{|w|+1}) \land T(x_{|w|+1}) \Rightarrow \bot \Big).$$

Next, let $\mathcal{A}=(Q,\Sigma,\delta,q_0,F)$ be a DFA. The signature τ_{Σ} is defined as before, and the signature τ_Q consists of the unary symbols R_q for every $q\in Q$ that is reachable from q_0 , i.e., there exists a word $a_1\ldots a_n\in \Sigma^+$ such that $q=\delta(a_n,\ldots\delta(a_2,\delta(a_1,q_0))\ldots)$. Note that q_0 is not necessarily reachable from itself. Let Φ_{sl}^+ be the universal Horn sentence over the signature $\tau_\Sigma\cup\tau_Q$ whose quantifier-free part contains: for every $a\in\Sigma$, the Horn clause

$$I(x_1) \wedge R_a(x_1, x_2) \Rightarrow R_{\delta(q_0, a)}(x_2),$$

for every $(q, a) \in Q \times \Sigma$ such that q is reachable from q_0 , the Horn clause

$$R_q(x_1) \wedge R_a(x_1, x_2) \Rightarrow R_{\delta(q,a)}(x_2),$$

and, for every $q \in F$ that is reachable from q_0 , the Horn clause

$$R_q(x_1) \wedge T(x_1) \Rightarrow \bot.$$

Then Φ_{sl} is the connected monadic Datalog sentence obtained from Φ_{sl}^+ by existentially quantifying all symbols from τ_Q upfront. The following lemma can be proved similarly as Lemma 7.6, and its proof is omitted.

Lemma 7.7. For every τ_{Σ} -structure \mathfrak{A} , the following are equivalent

- $\mathfrak{A} \models \Phi_{\mathscr{A}}$;
- for every $w \in L(\mathcal{A})$,

$$\mathfrak{A} \models \forall x_1, \dots, x_{|w|+1} \Big(I(x_1) \land \phi_w(x_1, \dots, x_{|w|+1}) \land T(x_{|w|+1}) \Rightarrow \bot \Big).$$

We are now ready to state the main lemma of this section, from which Theorem 7.1 follows immediately because the regularity problem for context-free languages is undecidable [54].

Lemma 7.8. Let \mathcal{G} be a context-free grammar. Then the following are equivalent:

- (1) $L(\mathcal{G})$ is regular.
- (2) $\Phi_{\mathscr{C}}$ is equivalent to connected monadic Datalog sentence.
- (3) $\Phi_{\mathscr{C}}$ is equivalent to a GMSNP sentence.
- (4) $fm(\Phi_{\mathscr{C}})$ is the CSP of a finite structure.
- (5) $\operatorname{fm}(\Phi_{\mathfrak{C}})$ is the CSP of an ω -categorical structure.
- (6) $fm(\Phi_{\mathcal{G}})$ is the age of a reduct of a finitely bounded homogeneous Ramsey structure.
- (7) $\operatorname{fm}(\Phi_{\mathscr{C}})$ is the age of an ω -categorical structure.
- (8) $fm(\Phi_{\mathfrak{B}})$ is the age of a finitely-bounded homogenizable structure.

Proof. " $(2) \Rightarrow (3)$ " This direction is trivial.

- "(4) \Rightarrow (5)" This direction is also trivial.
- "(6) \vee (8) \Rightarrow (7)" This direction is well-known and easy to see [40].
- "(3) \Rightarrow (6)" Since $\Phi_{\mathfrak{F}}$ is connected, $\operatorname{fm}(\Phi_{\mathfrak{F}})$ is preserved by disjoint unions. Since $\Phi_{\mathfrak{F}}$ is equivalent to a GMSNP sentence, by the first part of Theorem 7.2, $\Phi_{\mathfrak{F}}$ is equivalent to a disjunction $\Phi_1 \vee \cdots \vee \Phi_k$ of connected GMSNP sentences. We may assume that k is minimal with this property. If $k \geq 2$, then, by the minimality of k, there exist $\mathfrak{A}_1, \mathfrak{A}_2 \in \Phi_{\mathfrak{F}}$ such that $\mathfrak{A}_i \models \Phi_i \wedge \bigwedge_{j \in [k] \setminus \{i\}} \neg \Phi_j$ for $i \in \{1, 2\}$. But then the disjoint union of \mathfrak{A}_1 and \mathfrak{A}_2 does not satisfy $\Phi_{\mathfrak{F}}$, a contradiction. Hence, k = 1. Now the statement follows from the second part of Theorem 7.2.
- "(1) \Rightarrow (2) \land (4) \land (8)" By Theorem 7.4, there exists a DFA \mathscr{A} such that $L(\mathscr{G}) = L(\mathscr{A})$. By Lemma 7.6 and Lemma 7.7, we have $\operatorname{fm}(\Phi_{\mathscr{G}}) = \operatorname{fm}(\Phi_{\mathscr{A}})$. This implies item (2) because $\Phi_{\mathscr{A}}$ is in connected monadic Datalog. In fact, the sentence falls into an even stricter fragment which was called $\operatorname{caterpillar Datalog}$ in [37]. By Theorem 4.1 in [37], there exists a finite structure whose CSP equals $\operatorname{fm}(\Phi_{\mathscr{A}})$. This implies item (4). Finally, we show that $\operatorname{fm}(\Phi_{\mathscr{G}})$ is the age of a finitely-bounded homogenizable structure. Since the universal sentence $\Phi_{\mathscr{A}}^+$ is complete, the class $\operatorname{fm}(\Phi_{\mathscr{A}}^+)$ is closed under unions, and therefore has the AP. Let \mathfrak{H} be the homogeneous structure from Theorem 3.1 associated with $\operatorname{fm}(\Phi_{\mathscr{A}}^+)$. By the definition of $\operatorname{fm}(\Phi_{\mathscr{A}}^+)$, the signature τ_Q only contains unary symbols R_q for those $q \in Q$ which are reachable from q_0 . Let $Q_0 \subseteq Q$ be the set of states reachable from q_0 and, for every $q \in Q_0$, let $w_q \in \Sigma^+$ be an arbitrary word witnessing that q is reachable from q_0 . Note that, if $q_1, q_2 \in Q_0$ are distinct, then $w_{q_1} \neq w_{q_2}$. For every $w \in \Sigma^+$, consider the unary formula

$$\phi_w^I(x_1) := \exists x_2, \dots, x_{|w|+1} (I(x_{|w|+1}) \land \phi_w(x_{|w|+1}, \dots, x_1)).$$

We show that, for every $q \in Q_0$, the unary formulas $R_q(x)$ and $\phi_{w_q}^I(x)$ define the same relation in \mathfrak{H} . Then, for every $q \in Q_0$, the relation $R_q^{\mathfrak{H}}$ is first-order definable in the τ_{Σ} -reduct of \mathfrak{H} . This implies item (8). First, suppose that $\mathfrak{H} \models \phi_{w_q}^I(x)$ for some $x \in H$. Since w_q witnesses that q is reachable from q_0 and $\mathfrak{H} \models \Phi_{sl}^+$, it follows that $\mathfrak{H} \models R_q(x)$. Next, suppose that $\mathfrak{H} \models R_q(x)$ for some $x \in H$. Since \mathfrak{A} is deterministic, by the definition of Φ_{sl}^+ , there exists a structure \mathfrak{A} in the signature of \mathfrak{H} with domain $\{x_{|w_q|+1},\ldots,x_1\}$ that satisfies $\mathfrak{A} \models \Phi_{sl}^+ \wedge I(x_{|w_q|+1}) \wedge \phi_{w_q}(x_{|w_q|+1},\ldots,x_1)$ and whose substructure on $\{x_1\}$ is isomorphic to the substructure of \mathfrak{H} on $\{x_1\}$. Since $\mathfrak{A} \models \Phi_{sl}^+$,

there exists an embedding $e: \mathfrak{A} \hookrightarrow \mathfrak{H}$. Since \mathfrak{H} is homogeneous, there exists an automorphism f of \mathfrak{H} such that $f \circ e(x_1) = x$. Since automorphisms preserve first-order definable relations, it follows that $\mathfrak{H} \models \phi_{w_q}^I(x)$.

"(5) \vee (7) \Rightarrow (1)" Let \mathfrak{D} be an ω -categorical structure such that $\operatorname{fm}(\Phi_{\mathfrak{C}}) = \operatorname{age}(\mathfrak{D})$ or $\operatorname{fm}(\Phi_{\mathfrak{C}}) = \operatorname{CSP}(\mathfrak{D})$. Suppose, on the contrary, that $L(\mathfrak{C})$ is not regular. For every $w \in \Sigma^+$, consider the unary formula

$$\phi_w^T(x_1) := \exists x_2, \dots, x_{|w|+1} (\phi_w(x_1, \dots, x_{|w|+1}) \land T(x_{|w|+1})).$$

Since $L(\mathfrak{C})$ is not regular, by Theorem 7.4, $\sim_{L(\mathfrak{C})}$ has infinitely many classes. Let w_1, \ldots be the representatives of the classes. For every $i \geq 1$ let $R_i \subseteq D$ be the unary relation defined by $\phi^I_{w_i}(x_1)$ in \mathfrak{D} . For every $i \geq 1$, the formula $\phi^I_{w_i}(x_1)$ is satisfiable in \mathfrak{D} because it does not contain any T-atom. Since first-order formulas are preserved under automorphisms, for every $i \geq 1$, R_i is a non-empty union of unary orbits. By the definition of $\sim_{L(\mathfrak{C})}$, for every pair $i \neq j$, there exists $w \in \Sigma^*$ such that $|\{w_i w, w_j w\} \cap L(\mathfrak{C})| = 1$. By Lemma 7.6, exactly one of the relations R_i and R_j contains an element satisfying $\phi^T_w(x_1)$ in \mathfrak{D} . In other words, there exists an orbit that is contained in one of the two relations but not in the other. This can only be the case for all $i \neq j$ if there are infinitely many unary orbits, a contradiction to ω -categoricity. Therefore, $L(\mathfrak{C})$ is regular.

Acknowledgements

The author thanks Manuel Bodirsky, Simon Knäuer, Jakub Opršal, Paolo Marimon, and Michael Pinsker for inspiring discussions on the topic, and the anonymous reviewers of the conference version of the article for many helpful suggestions.

Funding

This research was funded in whole or in part by the Austrian Science Fund (FWF) [I 5948]. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

References

- Serge Abiteboul, Richard Hull, and Victor Vianu. Foundations of databases, volume 8. Addison-Wesley, 1995.
- [2] Ove Ahlman. Homogenizable structures and model completeness. Arch. Math. Log., 55(7-8):977-995, 2016.
- [3] Reza Akhtar and Alistair H. Lachlan. On countable homogeneous 3-hypergraphs. Arch. Math. Log., 34(5):331–344, 1995.
- [4] Peter RJ Asveld and Anton Nijholt. The inclusion problem for some subclasses of context-free languages. Theoretical computer science, 230(1-2):247-256, 2000.
- [5] Albert Atserias and Szymon Torunczyk. Non-homogenizable classes of finite structures. In Jean-Marc Talbot and Laurent Regnier, editors, 25th EACSL Annual Conference on Computer Science Logic, CSL 2016, August 29 - September 1, 2016, Marseille, France, volume 62 of LIPIcs, pages 16:1–16:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016.
- [6] Franz Baader and Jakub Rydval. Using model theory to find decidable and tractable description logics with concrete domains. J. Autom. Reason., 66(3):357–407, 2022.
- [7] Libor Barto, Jakub Bulín, Andrei Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. *Journal of the ACM (JACM)*, 68(4):1–66, 2021.
- [8] Libor Barto, Michael Kompatscher, Miroslav Olšák, Trung Van Pham, and Michael Pinsker. The equivalence of two dichotomy conjectures for infinite domain constraint satisfaction problems. In 32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS

- $2017,\ Reykjavik,\ Iceland,\ June\ 20-23,\ 2017,\ pages\ 1-12.$ IEEE Computer Society, 2017. available at arXiv:1612.07551.
- [9] Meghyn Bienvenu, Balder ten Cate, Carsten Lutz, and Frank Wolter. Ontology-based data access: A study through disjunctive datalog, csp, and MMSNP. ACM Trans. Database Syst., 39(4):33:1–33:44, 2014.
- [10] Manuel Bodirsky. Complexity of Infinite-Domain Constraint Satisfaction. Cambridge University Press, 2021.
- [11] Manuel Bodirsky and Bertalan Bodor. Canonical polymorphisms of ramsey structures and the unique interpolation property. In 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2021, Rome, Italy, June 29 July 2, 2021, pages 1–13. IEEE, IEEE, 2021.
- [12] Manuel Bodirsky, David Bradley-Williams, Michael Pinsker, and András Pongrácz. The universal homogeneous binary tree. J. Log. Comput., 28(1):133–163, 2018.
- [13] Manuel Bodirsky and Víctor Dalmau. Datalog and constraint satisfaction with infinite templates. J. Comput. Syst. Sci., 79(1):79–100, 2013. A preliminary version appeared in the proceedings of the Symposium on Theoretical Aspects of Computer Science (STACS'05).
- [14] Manuel Bodirsky and Martin Grohe. Non-dichotomies in constraint satisfaction complexity. In Luca Aceto, Ivan Damgård, Leslie Ann Goldberg, Magnús M. Halldórsson, Anna Ingólfsdóttir, and Igor Walukiewicz, editors, Automata, Languages and Programming, 35th International Colloquium, ICALP 2008, Reykjavik, Iceland, July 7-11, 2008, Proceedings, Part II - Track B: Logic, Semantics, and Theory of Programming & Track C: Security and Cryptography Foundations, volume 5126 of Lecture Notes in Computer Science, pages 184–196. Springer, July 2008.
- [15] Manuel Bodirsky, Peter Jonsson, and Van Trung Pham. The complexity of phylogeny constraint satisfaction problems. ACM Trans. Comput. Log., 18(3):23:1–23:42, 2017.
- [16] Manuel Bodirsky and Jan Kára. The complexity of temporal constraint satisfaction problems. J. ACM, 57(2):9:1–9:41, 2010.
- [17] Manuel Bodirsky, Simon Knäuer, and Florian Starke. ASNP: A tame fragment of existential second-order logic. In Marcella Anselmo, Gianluca Della Vedova, Florin Manea, and Arno Pauly, editors, Beyond the Horizon of Computability 16th Conference on Computability in Europe, CiE 2020, Fisciano, Italy, June 29 July 3, 2020, Proceedings, volume 12098 of Lecture Notes in Computer Science, pages 149–162. Springer, Springer, 2020.
- [18] Manuel Bodirsky, Florent R. Madelaine, and Antoine Mottet. A proof of the algebraic tractability conjecture for monotone monadic SNP. SIAM J. Comput., 50(4):1359–1409, 2021.
- [19] Manuel Bodirsky and Antoine Mottet. Reducts of finitely bounded homogeneous structures, and lifting tractability from finite-domain constraint satisfaction. In Martin Grohe, Eric Koskinen, and Natarajan Shankar, editors, Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '16, New York, NY, USA, July 5-8, 2016, pages 623–632. ACM, 2016. available at ArXiv:1601.04520.
- [20] Manuel Bodirsky and Michael Pinsker. Schaefer's theorem for graphs. Journal of the ACM (JACM), 62(3):1–52, 2015.
- [21] Manuel Bodirsky and Michael Pinsker. Canonical functions: a proof via topological dynamics. Contributions Discret. Math., 16(2):36–45, 2021.
- [22] Manuel Bodirsky, Michael Pinsker, and András Pongrácz. Projective clone homomorphisms. J. Symb. Log., 86(1):148–161, 2021.
- [23] Manuel Bodirsky, Michael Pinsker, and Todor Tsankov. Decidability of definability. J. Symb. Log., 78(4):1036–1054, 2013.
- [24] Manuel Bodirsky and Jakub Rydval. On the descriptive complexity of temporal constraint satisfaction problems. J. ACM, 70(1):2:1–2:58, 2023.
- [25] Manuel Bodirsky, Jakub Rydval, and André Schrottenloher. Universal horn sentences and the joint embedding property. Discret. Math. Theor. Comput. Sci., 23(2):1–15, 2021.
- [26] Mikolaj Bojanczyk, Luc Segoufin, and Szymon Torunczyk. Verification of database-driven systems via amalgamation. In Richard Hull and Wenfei Fan, editors, Proceedings of the 32nd ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems, PODS 2013, New York, NY, USA - June 22 - 27, 2013, pages 63-74. ACM, 2013.
- [27] Pierre Bourhis and Carsten Lutz. Containment in monadic disjunctive datalog, mmsnp, and expressive description logics. In Fifteenth International Conference on the Principles of Knowledge Representation and Reasoning, 2016.

- [28] Samuel Braunfeld. The undecidability of joint embedding and joint homomorphism for hereditary graph classes. *Discret. Math. Theor. Comput. Sci.*, 21(2):1–17, 2019. available at arXiv:1903.11932.
- [29] Andrei A. Bulatov. A dichotomy theorem for nonuniform csps. In Chris Umans, editor, 58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017, pages 319–330. IEEE, IEEE Computer Society, 2017.
- [30] Jakub Bulin, Dejan Delic, Marcel Jackson, and Todd Niven. A finer reduction of constraint problems to digraphs. *Log. Methods Comput. Sci.*, 11(4):1–33, 2015.
- [31] Gregory L. Cherlin. Homogeneous directed graphs. In Finite and Infinite Combinatorics in Sets and Logic, pages 81–95. Springer, 1993.
- [32] Gregory L. Cherlin. Forbidden substructures and combinatorial dichotomies: WQO and universality. Discret. Math., 311(15):1543-1584, 2011.
- [33] Gregory L. Cherlin. Homogeneous ordered graphs, metrically homogeneous graphs, and beyond, volume 1. Cambridge University Press, 2022.
- [34] Gregory L. Cherlin, Saharon Shelah, and Niandong Shi. Universal graphs with forbidden subgraphs and algebraic closure. Advances in Applied Mathematics, 22(4):454–491, 1999.
- [35] Lorenzo Clemente and Slawomir Lasota. Reachability analysis of first-order definable pushdown systems. In 24th EACSL Annual Conference on Computer Science Logic (CSL 2015). Schloss-Dagstuhl-Leibniz Zentrum für Informatik, 2015.
- [36] Jacinta Covington. Homogenizable relational structures. *Illinois Journal of Mathematics*, 34(4):731 743, 1990.
- [37] Péter L. Erdös, Claude Tardif, and Gábor Tardos. Caterpillar dualities and regular languages. SIAM J. Discret. Math., 27(3):1287–1294, 2013.
- [38] David M Evans, Jan Hubička, and Jaroslav Nešetřil. Automorphism groups and ramsey properties of sparse graphs. Proceedings of the London Mathematical Society, 119(2):515–546, 2019.
- [39] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: A study through datalog and group theory. SIAM J. Comput., 28(1):57–104, 1998.
- [40] Wilfrid Hodges. A shorter model theory. Cambridge University Press, Cambridge, 1997.
- [41] Jan Hubička and Jaroslav Nešetřil. Homomorphism and embedding universal structures for restricted classes. J. Multiple Valued Log. Soft Comput., 27(2-3):229–253, 2016. available at arXiv:0909.4939.
- [42] Jan Hubička and Jaroslav Nešetřil. All those ramsey classes (ramsey classes with closures and forbidden homomorphisms). Advances in Mathematics, 356:106791, 2019. available online: http://arxiv.org/abs/1606.07979.
- [43] Jan Hubička and Jaroslav Nešetřil. All those ramsey classes (ramsey classes with closures and forbidden homomorphisms). Advances in Mathematics, 356:1–89, 2019.
- [44] Julia F. Knight and Alistair H. Lachlan. Shrinking, stretching, and codes for homogeneous structures. In Classification Theory: Proceedings of the US-Israel Workshop on Model Theory in Mathematical Logic held in Chicago, Dec. 15–19, 1985, volume 1292, pages 192–229. Springer, Springer, Berlin, Heidelberg, 2006.
- [45] Michael Kompatscher and Trung Van Pham. A complexity dichotomy for poset constraint satisfaction. *Journal of Applied Logics*, 5(8):1663–1697, 2018.
- [46] Alistair H. Lachlan. Homogeneous structures. In Proceedings of the International Congress of Mathematicians, volume 1, pages 314–321, Berkeley, 1986. AMS.
- [47] Carsten Lutz and Maja Milicic. A tableau algorithm for description logics with concrete domains and general tboxes. J. Autom. Reason., 38(1-3):227-259, 2007.
- [48] Dugald Macpherson. A survey of homogeneous structures. Discret. Math., 311(15):1599–1634, 2011.
- [49] Florent R. Madelaine and Iain A. Stewart. Some problems not definable using structure homomorphisms. Ars Comb., 67:153–160, 2003.
- [50] Antoine Mottet. Promise and infinite-domain constraint satisfaction. In Aniello Murano and Alexandra Silva, editors, 32nd EACSL Annual Conference on Computer Science Logic, CSL 2024, February 19-23, 2024, Naples, Italy, volume 288 of LIPIcs, pages 41:1-41:19. Schloss-Dagstuhl-Leibniz Zentrum für Informatik, Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2024.

- [51] Antoine Mottet and Michael Pinsker. Smooth approximations and csps over finitely bounded homogeneous structures. In Christel Baier and Dana Fisman, editors, LICS '22: 37th Annual ACM/IEEE Symposium on Logic in Computer Science, Haifa, Israel, August 2 - 5, 2022, pages 36:1–36:13. ACM, 2022. available at arXiv:2011.03978.
- [52] Jakub Rydval. Homogeneity and Homogenizability: Hard Problems for the Logic SNP. In Karl Bringmann, Martin Grohe, Gabriele Puppis, and Ola Svensson, editors, 51st International Colloquium on Automata, Languages, and Programming (ICALP 2024), volume 297 of Leibniz International Proceedings in Informatics (LIPIcs), pages 150:1–150:20, Dagstuhl, Germany, 2024. Schloss Dagstuhl Leibniz-Zentrum für Informatik.
- [53] François Schwarzentruber. The complexity of tiling problems. CoRR, abs/1907.00102:1–12, 2019. arXiv:1907.00102.
- [54] Jeffrey O. Shallit. A second course in formal languages and automata theory. Cambridge University Press, 2008.
- [55] Larry J Stockmeyer. The polynomial-time hierarchy. Theoretical Computer Science, 3(1):1–22, 1976.
- [56] Van Thé and Lionel Nguyen. A survey on structural ramsey theory and topological dynamics with the kechris-pestov-todorcevic correspondence in mind. arXiv preprint arXiv:1412.3254, 2014.
- [57] Peter van Emde Boas. The convenience of tilings. In Complexity, Logic, and Recursion Theory, pages 331–363. CRC Press, 2019.
- [58] Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In 58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017, pages 331-342. IEEE Computer Society, 2017.
- [59] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. J. ACM, 67(5):30:1–30:78, 2020.

TU WIEN, VIENNA, AUSTRIA

 $Email\ address: {\tt jakub.rydval@tuwien.ac.at}$