

Investigation of proton structure function F_2^p at HERA in light of an analytical solution to Balitsky-Kovchegov equation

Ranjan Saikia ^{*1}, Pragyan Phukan ^{†2}, and Jayanta Kumar Sarma ^{‡1}

¹*HEP laboratory, Department of Physics, Tezpur University, Tezpur, Assam-784028, India*

²*Department of Physics, Moran College, Moranhat, Assam-785670, India*

Abstract

In this paper, the proton structure function $F_2^p(x, Q^2)$ at small- x is investigated using an analytical solution of the Balitsky-Kovchegov (BK) equation. In the context of the color dipole description of deep inelastic scattering (DIS), the structure function $F_2^p(x, Q^2)$ is computed by applying the analytical expression for the scattering amplitude $N(k, Y)$ derived from the BK solution. At transverse momentum k and total rapidity Y , the scattering amplitude $N(k, Y)$ represents the propagation of the quark-antiquark dipole in the color dipole description of DIS. Using the BK solution we extracted the integrated gluon density $xg(x, Q^2)$ and then compared our theoretical estimation with the LHAPDF global data fits NNPDF3.1sx and CT18. Finally, we investigated the behavior of $F_2^p(x, Q^2)$ in the kinematic region of $10^{-5} \leq x \leq 10^{-2}$ and $2.5 \text{ GeV}^2 \leq Q^2 \leq 60 \text{ GeV}^2$. Our predicted results for $F_2^p(x, Q^2)$ within the specified kinematic region are in good agreement with the recent high-precision data for $F_2^p(x, Q^2)$ from HERA (H1 Collaboration) and the LHAPDF global parametrization group NNPDF3.1sx.

Keywords: proton structure function, small- x , Balitsky-Kovchegov equation.

*Corresponding author: ranjans@tezu.ernet.in

†E-mail: phukan.pragyan@gmail.com

‡E-mail: jks@tezu.ernet.in

I Introduction

Understanding the substructure of the nucleon within matter is one of the fundamental research topics in high-energy particle physics. It is crucial to understand the structure of the nucleon in order to comprehend the fundamental structure of matter. With the introduction of high-energy accelerator facilities, we have been able to understand the substructure of the nucleon within the framework of quantum chromodynamics (QCD). Understanding the substructure of the nucleon has relied heavily on the structure functions of the nucleon. Deep inelastic scattering (DIS) experiments on hadrons by leptons have yielded significant data on the distribution of partons within hadrons in terms of the quark and gluon distributions. The DIS cross section is associated with the structure functions of the nucleon in relation to parton distributions. The measurements of the proton's structure functions $F_2^p(x, Q^2)$ and $F_L^p(x, Q^2)$ at HERA have begun a new era of parton density measurement within the nucleon [1–4]. These structure functions can be correlated with the momentum distributions of partons within the nucleon, and thus the parton distribution functions (PDFs). At high energies, or equivalently at small- x (Bjorken x) values, the gluon density dominates among partons, and hence the dominant contribution in $F_2^p(x, Q^2)$ and $F_L^p(x, Q^2)$ observations comes exclusively from gluons. Therefore, measurements of these structure functions at small- x are essential for calculating gluon distribution functions and visualising the overall hadronic wave function in high-density QCD.

The DGLAP (Dokshitzer–Gribov–Lipatov–Altarelli–Parisi) evolution equation [5–9], being the famous and well-established evolution equation, acts as a basic tool for the theoretical investigation of DIS structure functions. This equation has been successfully applied to address available HERA data in the moderate kinematical region of Bjorken's x ($x \geq 0.01$). At small- x , HERA data shows a steep rise in the behaviour of gluons in the region. This steep behaviour of gluons at small- x is well described by the famous BFKL (Balitsky-Fadin-Kuraev-Lipatov) evolution equation [10,11]. However, the rapid growth of gluons at small- x cannot continue indefinitely; otherwise, the physical cross-sections will violate unitarity and the Froissart-Martin bound [12]. As a result, both the DGLAP and BFKL evolution equations fail to account for implicit physics in high-density QCD. In order to maintain the unitarity of the theory, the infinite growth of gluons must be slowed down by certain processes. The phenomena of gluon recombination and saturation have provided the solution to the problems faced at small- x by the linear evolution equations [13–17]. The nonlinear phenomena of gluon recombination and saturation lead to nonlinear terms in the DGLAP and BFKL equations. The Balitsky-Kovchegov (BK) equation [18–21], nonlinearization of the BFKL equation, and mean-field approximation to the Jalilian-Marian-Iancu-McLerran-

Weigert-Leonidov-Kovner (JIMWLK) equation [22–25] describe the behaviour of gluon density in the small- x region. The BK equation is an integro-differential equation in coordinate space that can be transformed to momentum space to yield a partial differential equation useful for phenomenological studies at various high-energy accelerator facilities. This equation has been studied and solved numerically; however, due to its complexity, we have not seen an exact analytical solution to this equation to date. We recently proposed an approximate analytical solution to the BK equation that can be useful for phenomenology in the context of gluon saturation and hence small- x physics [26].

The collaborations at HERA contributed high-precision data for measuring the proton structure function F_2^p at different kinematical regions of x and Q^2 [27, 28]. Some recent works on the measurement of proton structure functions using various evolution equations and approaches are described in refs. [29–33], and the results correlate well with the experimental data. In ref. [34], for the first time, the reproduction of measurements of the DIS proton structure function at high energy from the color dipole description in momentum space was investigated. Using the knowledge of asymptotic solutions of the BK equation, the authors of ref. [34] measured the charm structure function F_2^c as a function of x for various values of Q^2 and compared their results favourably with those of the HERA experiment. At small- x , it has been shown at HERA that the contributions in PDFs comes exclusively from gluons. Thus, the determination of gluon density is essential in order to understand the overall hadronic wave function at small- x . We do not see any theoretical measurement of gluon density at small- x using BK evolution theory, which motivates the current work. We previously presented an analytical expression for the scattering amplitude $N(k, Y)$, which describes the propagation of the quark-antiquark dipole through the target color field in the color dipole model at various rapidities, Y [26]. In this work, we want to extract the gluon density $xg(x, Q^2)$ and analyze the proton structure function F_2^p at small- x by combining experimental data with QCD evolution theory using the analytical solution of the BK equation in momentum space. We compute the gluon density $xg(x, Q^2)$ and the proton structure function F_2^p at various kinematical regions of HERA and analyze their behavior at small- x .

The plan of the paper is as follows: In Section II, we relate the dipole-proton cross-section with forward scattering amplitude within the color dipole description of DIS in QCD. Then, using the analytical solution of the BK equation, we discuss how the structure function F_2^p of the proton can be obtained from the color dipole description. In Section III, we present the x evolution of the integrated gluon density $xg(x, Q^2)$ for two Q^2 values, viz. 35 GeV^2 and 100 GeV^2 using the BK evolution theory. Our results are compared with the LHAPDF global data fits NNPDF3.1sx [35] and CT18 [36]. Both have LHC and HERA data included in their analyses. Finally, the obtained numerical results on the proton structure function

F_2^p using the analytical solution of the BK equation and comparison with the data from the H1 Collaboration and LHAPDF global parameterization group NNPDF3.1sx [35] are shown. Section IV follows the discussion of the summary and conclusion of our work.

II Proton structure functions in the color dipole description

1 Color dipole description

In QCD, the color dipole description of DIS [37–39] has been a useful tool for various perturbative QCD (pQCD) calculations at small- x of high density QCD. It is especially convenient to work with the pQCD dipole description of DIS when it comes to the energy dependence of the scattering amplitude of any DIS event. The scattering process in the color dipole description of DIS can be factored into several steps where an incoming high-speed virtual photon (γ^*) after fluctuation from QCD vacuum changes to a quark-antiquark ($q\bar{q}$) dipole. The formed dipole then scatters off the target proton, eventually forming the final-state particles. In the color dipole description, following ref. [40], the total γ^*p cross section for the γ^*p scattering can be factored as

$$\sigma_{T,L}^{\gamma^*p} = \int d^2r \int_0^1 dz |\Psi_{T,L}(r, z, Q^2)|^2 \sigma_{q\bar{q}}(r, Y), \quad (1)$$

$\Psi_{T,L}$ represents the wave function for the virtual photon (γ^*) to fluctuate into the quark-antiquark ($q\bar{q}$) dipole with a transverse size of r . T, L represent the transverse and longitudinal polarisation states of the virtual photon, respectively, and z represents the longitudinal momentum fraction of the virtual photon carried by the quark (or antiquark). Q^2 represents the virtuality of the photon, while Y represents the total rapidity. The dipole-proton cross section $\sigma_{q\bar{q}}(r, Y)$, which is the imaginary component of the forward scattering amplitude of the dipole on the target proton, contains all the information on the hadronic interactions.

The dipole-proton cross section, $\sigma_{q\bar{q}}(r, Y)$, can be obtained from the elastic dipole-proton scattering amplitude $\mathcal{A}(r, Y)$ using the optical theorem as [41, 42]

$$\sigma_{q\bar{q}}(r, Y) = 2 \text{Im}\mathcal{A}(r, Y) = 2 \int d^2b N(r, Y, b) = \sigma_0 N(r, Y). \quad (2)$$

$N(r, Y, b)$ is the value for the imaginary part of the forward elastic dipole-proton scattering amplitude. Now, if one considers the target proton as a homogeneous disk of radius R_p , then the dipole-proton cross section can be related to the forward scattering amplitude $N(r, Y)$

by the following relation [34] :

$$\sigma_{q\bar{q}}(r, Y) = 2\pi R_p^2 N(r, Y). \quad (3)$$

The scattering amplitude $N(r, Y)$ will come from the solution of the BK equation.

After obtaining the γ^*p cross-section, one can obtain directly the proton structure function F_2^p from the γ^*p cross-section through the relation

$$F_2^p(x, Q^2) = \frac{Q^2}{4\pi^2\alpha_{em}} \left[\sigma_T^{\gamma^*p}(x, Q^2) + \sigma_L^{\gamma^*p}(x, Q^2) \right]. \quad (4)$$

2 F_2^p from the solution of the BK equation

Now, we see how the structure function F_2^p can be obtained using the scattering amplitude $N(r, Y)$ obtained from the analytical solution of the BK equation. The BK equation describes the high-energy evolution of the dipole-target proton scattering amplitude $N(r, Y)$ in the dipole description, which in turn describes the propagation of the quark-antiquark dipole through the target color field. We express the γ^*p cross-section in $N(k, Y)$ and hence the structure function of the proton in momentum space, as the asymptotic behaviour of the solution of the BK equation can be expressed naturally in momentum space. For that, let us transform $N(r, Y)$ to $N(k, Y)$ by the following simple Fourier transform

$$N(k, Y) = \frac{1}{2\pi} \int \frac{d^2r}{r^2} e^{ik \cdot r} N(r, Y) = \int_0^\infty \frac{dr}{r} J_0(kr) N(r, Y). \quad (5)$$

The proton structure function F_2^p in momentum space related to $N(k, Y)$ can be expressed as follows using the discussion from above and some algebraic calculations [34]:

$$F_2^p(x, Q^2) = \frac{Q^2 R_p^2 N_c}{4\pi^2} \int_0^\infty \frac{dk}{k} \int_0^1 dz |\tilde{\Psi}(k^2, z; Q^2)|^2 N(k, Y), \quad (6)$$

where the photon wave function $\tilde{\Psi}$ is now expressed in the momentum space and can be found in ref. [34] and N_c are the number of colors. The scattering amplitude $N(k, Y)$ comes from the solution of the BK equation. The scattering amplitude $N(k, Y)$ at total rapidity Y and transverse momentum k obeys the BK equation in momentum space as [20]

$$\partial_Y N = \bar{\alpha} \chi(-\partial_L) N - \bar{\alpha} N^2, \quad (7)$$

where $\bar{\alpha} = \frac{\alpha_s N_c}{\pi}$ and $\chi(\gamma) = 2\psi(1) - \psi(\gamma) - \psi(1 - \gamma)$ is the BFKL kernel, $\gamma = -\partial_L$, $L = \ln \frac{k^2}{k_0^2}$ with k_0 being some fixed low momentum scale. When expanding $\chi(\gamma)$ to second order, it

is seen [43–45] that with some variable transformation the BK equation can be rewritten as the FKPP (Fisher-Kolmogorov-Petrovsky-Piscounov) equation as [46, 47]

$$\partial_t u(t, x) = \partial_x^2 u(t, x) + u(t, x) - u^2(t, x). \quad (8)$$

Both the equations belong to the same universality class, and their solutions are of the same nature, i.e., travelling wave nature. We suggested an approximate analytical solution of the BK equation (7) in connection with the FKPP equation (8) using the homotopy perturbation method (HPM) in ref. [26]. Following ref. [26], the solution of the BK equation (7) in connection with the FKPP equation (8) is given by

$$N(k, Y) = \frac{Ne^Y}{1 - N + Ne^Y}. \quad (9)$$

N is the initial condition at $Y = 0$ i.e. $N(k, 0) = N$. This solution of the BK equation gives the scattering amplitude $N(k, Y)$ at any given rapidity $Y > 0$, once the initial condition N is known to us. For the initial condition N , we use the GBW (Golec-Biernat and Wusthoff) initial condition given by [48]

$$N(r, Y = 0) = 1 - \exp \left[- \left(\frac{r^2 Q_{s0}^2}{4} \right) \right]. \quad (10)$$

Q_{s0}^2 is the initial saturation momentum of gluons that can be fitted from the existing HERA data, and its value is 0.24 GeV^2 [49]. As we are dealing with the the BK equation in momentum space, the use of the GBW initial condition would be helpful as this can be transformed into momentum space simply results as

$$N(k, Y = 0) = \int \frac{d^2 r}{2\pi r^2} e^{ik \cdot r} N(r, Y = 0) = \frac{1}{2} \Gamma \left(0, \frac{k^2}{Q_{s0}^2} \right). \quad (11)$$

At large values of k^2/Q_{s0}^2 , the incomplete gamma function $\Gamma(0, k^2/Q_{s0}^2)$ behaves as

$$\Gamma \left(0, \frac{k^2}{Q_{s0}^2} \right) = \exp \left(- \frac{k^2}{Q_{s0}^2} \right). \quad (12)$$

Therefore, we can write the GBW initial condition (10) in momentum space as

$$N(k, Y = 0) = \frac{1}{2} \exp \left(- \frac{k^2}{Q_{s0}^2} \right). \quad (13)$$

Now, we use the above equation in equation (9) and replace the initial condition N with the

GBW initial condition in momentum space, which gives

$$N(k, Y) = \frac{e^{Y-k^2/Q_{s0}^2}}{1 - e^{-k^2/Q_{s0}^2} + e^{Y-k^2/Q_{s0}^2}}. \quad (14)$$

The scattering amplitude expression given above is an approximate analytical solution to the BK equation (7).

In light of the above discussions, we will now investigate the proton structure function F_2^p by combining experimental data from the HERA experiment with QCD evolution theory. To do this, we will use the analytical solution of the BK equation, which is expressed above.

III Results and discussions

To investigate the proton structure function F_2^p , we use the expression eqn.(6) with the analytical solution of the BK equation eqn.(14). We handle the expression eqn.(6) numerically for various inputs of x and Q^2 . In eqn.(6), $\tilde{\Psi}(k^2, z; Q^2)$ represents the probability of a virtual photon emitting a quark-antiquark pair with the momentum fraction z (quark) and $(1-z)$ (antiquark) of the virtual photon in the momentum space. Its expression is given by [34]

$$\begin{aligned} |\tilde{\Psi}(k^2, z; Q^2)|^2 = & \sum_q \left(\frac{4\bar{Q}_q^2}{k^2 + 4\bar{Q}_q^2} \right) e_q^2 \left\{ [z^2 + (1-z)^2] \right. \\ & \times \left[\frac{4(k^2 + \bar{Q}_q^2)}{k^2 \sqrt{k^2 + 4\bar{Q}_q^2}} \operatorname{arcsinh} \left(\frac{k}{2\bar{Q}_q} \right) + \frac{k^2 - 2\bar{Q}_q^2}{2\bar{Q}_q^2} \right] \\ & \left. + \frac{4Q^2 z^2 (1-z)^2 + m_q^2}{\bar{Q}_q^2} \times \left[\frac{k^2 + \bar{Q}_q^2}{\bar{Q}_q^2} - \frac{4\bar{Q}_q^4 + 2\bar{Q}_q^2 k^2 + k^4}{\bar{Q}_q^2 \sqrt{k^2(k^2 + 4\bar{Q}_q^2)}} \times \operatorname{arcsinh} \left(\frac{k}{2\bar{Q}_q} \right) \right] \right\}, \end{aligned} \quad (15)$$

where $\bar{Q}_q^2 = z(1-z)Q^2 + m_q^2$ and m_q is the mass of the quark of flavor q . Now, we see how the expression eqn.(15) will behave with transverse momentum k for different values of Q^2 with $m_q \rightarrow 0$ and $z = 1/2$ (as $0 < z < 1$). The probability distribution graph of the virtual photon to emit a quark-antiquark pair with $z = 1/2$ against transverse momentum k is depicted in Figure 1. The probability to emit a quark-antiquark pair increases as k increases until a certain point (peak point), after which it decreases as k increases. The peak point for every Q^2 value is the same, but the transverse momentum k corresponding to every peak is different for different Q^2 .

The solution (14) gives the propagation of the $q\bar{q}$ dipole in the color dipole description of QCD, which in turn gives the gluon content of the proton, i.e., the unintegrated gluon




Figure 1: The probability distribution of the virtual photon to emit a quark-antiquark pair as a function of k with $m_q \rightarrow 0$ and $z = \frac{1}{2}$ for various Q^2 .

density. To calculate conventional gluon density $xg(x, Q^2)$, we integrate the BK solution (14) over transverse momentum with the relation

$$xg(x, Q^2) = \int_0^{Q^2} \frac{dk}{k} N(k, Y). \quad (16)$$

The extracted results of gluon density from BK solution is compared with the global data fits NNPDF3.1sx and CT18. The results are shown in Figure 2.




Figure 2: x evolution of gluon density $xg(x, Q^2)$ extracted from the BK solution compared with the global data fits NNPDF3.1sx and CT18.

For the proton structure function F_2^p , we plug the expression given in eqn.(15) (with $m_q \rightarrow 0$ and $z = 1/2$) together with the BK solution given in eqn.(14) into eqn.(6). We set

$N_c = 3$ and $R_p \approx 0.831 \text{ fm} \approx 4.22 \text{ GeV}^{-1}$ in the expression eqn.(6) and solve the expression numerically for various Q^2 . The value of the R_p is taken from the recent study on the proton radius [50]. We analyse the expression eqn.(6) numerically and compare our results with the HERA measurements of the proton structure function from the H1 Collaboration [28] with constraints to the kinematic region: $10^{-5} \leq x \leq 10^2$ and $2.5 \leq Q^2 \leq 60 \text{ GeV}^2$. We also compare our results with the LHAPDF global parameterization group NNPDF3.1sx [35]. The reason for choosing the particular kinematic region $x \leq 10^{-2}$ is to describe the small- x behaviour of high-energy amplitude as the BK equation is only applicable at small- x . For the too high Q^2 range, we would need corrections from the DGLAP equation, which we cannot skip at too high Q^2 . The results are shown in Figure 3.




Figure 3: The results of the proton structure function F_2^p as a function of x at various Q^2 obtained in this work are compared with the data from the H1 Collaboration [28] and global data fit NNPDF3.1sx [35].

IV Summary and conclusion

In this work, we have investigated the proton structure function F_2^p in light of an analytical solution of the BK equation. Proton structure functions have been investigated by different collaborations at experimental facilities such as HERA and the LHC. On the phenomenological side, the proton structure functions have been studied by studying different QCD evolution equations and tested well at existing experimental facilities. To investigate the proton structure functions at small- x , the BK equation is most suitable for testing the experimental data on observables. In this work, we have stated our discussion on how we can obtain the proton structure function F_2^p from the color dipole description of the DIS in QCD. In the color dipole approach, the proton structure function F_2^p is obtained directly from the virtual photon-proton cross-section, which can be expressed in the dipole-proton scattering amplitude $N(k, Y)$. The dipole-proton scattering amplitude $N(k, Y)$ is obtained from the solution of the BK equation. Using the BK solution, we extracted the x evolution of the integrated gluon density $xg(x, Q^2)$ and compared our results with those of LHAPDF global data fits NNPDF3.1sx and CT18. Both NNPDF3.1sx and CT18 have HERA and LHC data included in their analyses. Finally, we computed the proton structure function F_2^p and compared it with the measurement of F_2^p at HERA from the H1 Collaboration and the LHAPDF global parameterization group NNPDF3.1sx. Our predicted results are in good agreement with the experimental results within the kinematic region we have constrained. Beyond that, we have to consider corrections to the DGLAP equation. We have also shown in this work how the virtual photon wave function Ψ would behave against transverse momentum k with $m_q \rightarrow 0$, $z = 1/2$ at various Q^2 . We found that the probability of the virtual photon emitting a quark-antiquark pair increases as k increases to a certain peak value, after which it starts to fall as k increases. We have seen that the maximum probability is the same for different Q^2 values, irrespective of the value of k at the maximum peak for different Q^2 values.

In this work, we have seen the ability of our BK solution to describe physics at small- x . We have successfully applied our BK solution to calculate the gluon density $xg(x, Q^2)$ and the proton's structure function F_2^p at small- x within the kinematic region we have constraint. Also, our results are testable at future experimental facilities such as the LHeC (Large Hadron electron Collider) [51, 52], EIC (Electron Ion Collider) [53], and the FCC-eh (Future Circular Collider electron-hadron) [54]. In these future experimental facilities, the measurement of the proton structure function will be performed at much lower values of x with increased precision. Nevertheless, we could investigate the proton structure function F_2^p using an analytical solution of the BK equation within the constrained region. We conclude that the analytical solution of the BK equation can serve as a convenient tool for

further studies at small- x and high-density QCD. We hope that the BK equation, together with future experimental facilities, will help us understand and explore phenomena inside hadrons at small- x in the near future.

References

- [1] M.Derrick et al., [ZEUS Collaboration], *Measurement of the proton structure function F_2 in ep scattering at HERA*, *Phys. Lett. B* **316** (1993) 412.
- [2] I. Abt et al., [H1 Collaboration], *Measurement of the proton structure function $F_2(x, Q^2)$ in the low x region at HERA*, *Nucl. Phys. B* **407** (1993) 515.
- [3] T. Ahmed et al., [H1 Collaboration], *A measurement of the proton structure function $F_2(x, Q^2)$* , *Nucl. Phys. B* **439** (1995) 471 [arXiv.hep-ex/9503001].
- [4] C. Adloff et al., [H1 Collaboration], *Determination of the longitudinal proton structure function $F_L(x, Q^2)$ at low x* , *Phys. Lett. B* **393** (1997) 452 [arXiv.hep-ex/9611017].
- [5] V. N. Gribov and L. N. Lipatov, *Deep inelastic electron scattering in perturbation theory*, *Phys. Lett. B* **37** (1971) 78.
- [6] V. N. Gribov and L. N. Lipatov, *e^+e^- pair annihilation and deep inelastic ep scattering in perturbation theory*, *Sov. J. Nucl. Phys.* **15** (1972) 75 [iNSPIRE-hep].
- [7] L. N. Lipatov, *The parton model and perturbation theory*, *Sov. J. Nucl. Phys.* **20** (1975) 94 [iNSPIRE-hep].
- [8] G. Altarelli and G. Parisi, *Asymptotic freedom in parton language*, *Nucl. Phys. B* **126** (1977) 298.
- [9] Y. L. Dokshitzer, *Calculation of the structure functions for deep inelastic scattering and e^+e^- annihilation by perturbation theory in quantum chromodynamics*, *Sov. Phys.-JETP* **46** (1997) 641 [iNSPIRE-hep].
- [10] E. A. Kuraev, L. N. Lipatov and Victor S. Fadin, *The pomeranchuk singularity in non-abelian gauge theories*, *Zh. Eksp. Teor. Fiz.* **72** (1977) 377 [iNSPIRE-hep].
- [11] I. I. Balitsky and L. N. Lipatov, *The pomeranchuk singularity in quantum chromodynamics*, *Sov. J. Nucl. Phys.* **28** (1978) 822 [iNSPIRE-hep].

- [12] M. Froissart, *Asymptotic behavior and subtractions in the mandelstam representation*, *Phys. Rev.* **123** (1961) 1053.
- [13] L. V. Gribov, E. M. Levin and M. G. Ryskin, *Singlet structure function at small x : Unitarization of gluon ladders*, *Nucl. Phys. B* **188** (1981) 555.
- [14] L. V. Gribov, E. M. Levin and M. G. Ryskin, *Semihard processes in QCD*, *Phys. Rep.* **100** (1983) 1.
- [15] A. H. Mueller and Jianwei Qiu, *Gluon recombination and shadowing at small values of x* , *Nucl. Phys. B* **268** (1986) 427.
- [16] A. H. Mueller, *Small- x behavior and parton saturation: A QCD model*, *Nucl. Phys. B* **335** (1990) 115.
- [17] W. Zhu and J. Ruan, *A new modified Altarelli-Parisi evolution equation with parton recombination in proton*, *Nucl. Phys. B* **559** (1999) 378 [[arXiv.hep-ph/9907330](#)].
- [18] I. Balitsky, *Operator expansion for diffractive high-energy scattering*, *AIP Conference Proceedings* **407** (1997) 953 [[arXiv.hep-ph/9706411](#)].
- [19] Y. V. Kovchegov, *Small- x F_2 structure function of a nucleus including multiple Pomeron exchanges*, *Phys. Rev. D* **60** (1999) 034008 [[arXiv.hep-ph/9901281](#)].
- [20] Y. V. Kovchegov, *Unitarization of the BFKL Pomeron on a nucleus*, *Phys. Rev. D* **61** (2000) 074018 [[arXiv.hep-ph/9905214](#)].
- [21] I. Balitsky, *Effective field theory for the small- x evolution*, *Phys. Lett. B* **518** (2001) 235 [[arXiv.hep-ph/0105334](#)].
- [22] I. Balitsky, *Operator expansion for high-energy scattering*, *Nucl. Phys. B* **463** (1996) 99 [[arXiv.hep-ph/9509348](#)].
- [23] Jalilian-Marian, A. Kovner, A. Leonidov, and H. Weigert, *The BFKL equation from the Wilson renormalization group*, *Nucl. Phys. B* **504** (1997) 415 [[arXiv.hep-ph/9701284](#)].
- [24] E. Iancu, A. Leonidov, and L. D. McLerran, *Nonlinear gluon evolution in the color glass condensate: I*, *Nucl. Phys. A* **692** (2001) 583 [[arXiv.hep-ph/0011241](#)].
- [25] H. Weigert, *Unitarity at small Bjorken x* , *Nucl. Phys. A* **703** (2002) 823 [[arXiv.hep-ph/0004044](#)].

- [26] R. Saikia, P. Phukan, and J. K. Sarma, *An analytical solution of Balitsky-Kovchegov equation using homotopy perturbation method*, *Int. J. Mod. Phys. A* **37** (2022) 2250190 [[arXiv.2204.10111](#)].
- [27] F. D. Aaron et. al., [H1 and ZEUS Collaborations], *Combined measurement and QCD analysis of the inclusive $e^\pm p$ scattering cross sections at HERA*, *JHEP* **01** (2010) 109 [[arXiv.0911.0884](#)].
- [28] V. Andreev et al., [H1 Collaboration], *Measurement of inclusive ep cross sections at high Q^2 at $\sqrt{s} = 225$ and 252 GeV and of the longitudinal proton structure function F_L at HERA*, *Eur. Phys. J. C* **74** (2014) 2814 [[arXiv.1312.4821](#)].
- [29] P. Phukan, M. Lalung, and J. K. Sarma, *Studies on gluon evolution and geometrical scaling in kinematic constrained unitarized BFKL equation: application to high-precision HERA DIS data*, *Eur. Phys. J. C* **79** (2019) 507 [[arXiv.1811.01648](#)].
- [30] P. Phukan, M. Lalung, and J. K. Sarma, *Small x phenomenology on gluon evolution through the BFKL equation in light of a constraint in multi-Regge kinematics*, *Communications in Theoretical Physics* **72**(2) (2020) 025201 [[arXiv.1906.06468](#)].
- [31] M. Lalung, P. Phukan and J. K. Sarma, *On analytical solutions of the DIS structure functions in the framework of GLR-MQ-ZRS equation*, *Results in Physics* **28** (2021) 104551 [[arXiv.1909.13591](#)].
- [32] G.R.Boroun and B.Rezaei, *An evaluation of the proton structure functions F_2 and F_L at small x* , *Phys. Lett. B* **816** (2021) 136274 [[arXiv.2010.15357](#)].
- [33] G.R.Boroun, *Non-linear corrections to the longitudinal structure function F_L from the parametrization of F_2 : Laplace transform approach*, *Eur. Phys. J. Plus* **137**, (2022) 371 [[arXiv.2109.08878](#)].
- [34] J. T. de Santana Amaral, M. A. Betemps, M. B. Gay Ducati, and G. Soyez, *γ^*p cross section from the dipole model in momentum space*, *Phys. Rev. D* **76** (2007) 094018 [[arXiv.hep-ph/0612091](#)].
- [35] R. D. Ball et al., [NNPDF Collaboration], *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[arXiv.1706.00428](#)].
- [36] T.-J. Hou et al., [CTEQ Collaboration], *New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC*, *Phys. Rev. D* **103** (2021) 014013 [[arXiv.1912.10053](#)].

- [37] N. N. Nikolaev and B. G. Zakharov, *Colour transparency and scaling properties of nuclear shadowing in deep inelastic scattering*, *Z. Phys. C* **49** (1991) 607.
- [38] A. H. Mueller, *Soft gluons in the infinite-momentum wave function and the BFKL pomeron*, *Nucl. Phys. B* **415** (1994) 373.
- [39] A. H. Mueller and B. Patel, *Single and double BFKL pomeron exchange and a dipole picture of high energy hard processes*, *Nucl. Phys. B* **425** (1994) 471 [[arXiv.hep-ph/9403256](#)].
- [40] A.M. Stasto, K. Golec-Biernat, and J. Kwiecinski, *Geometric scaling for the total γ^*p cross section in the low x region*, *Phys. Rev. Lett.* **86** (2001) 596 [[arXiv.hep-ph/0007192](#)].
- [41] V. Barone and E. Predazzi, *High-Energy Particle Diffraction*, Springer, 1 ed., (2002). ISBN 978-3540421078.
- [42] H. Kowalski, L. Motyka, and G. Watt, *Exclusive diffractive processes at HERA within the dipole picture*, *Phys. Rev. D* **74** (2006) 074016 [[arXiv.hep-ph/0606272](#)].
- [43] S. Munier and R. Peschanski, *Geometric Scaling as Traveling Waves*, *Phys. Rev. Lett.* **91** (2003) 232001 [[arXiv.hep-ph/0309177](#)].
- [44] S. Munier and R. Peschanski, *Traveling wave fronts and the transition to saturation*, *Phys. Rev. D* **69** (2004) 034008 [[arXiv.hep-ph/0310357](#)].
- [45] S. Munier and R. Peschanski, *Universality and tree structure of high-energy QCD* *Phys. Rev. D* **70** (2004) 077503 [[arXiv.hep-ph/0401215](#)].
- [46] R. A. Fisher, *The wave of advance of advantageous genes*, *Ann. Eugen.* **7** (1937) 355.
- [47] A. Kolmogorov, I. Petrovsky and N. Piscounov, *A study of the diffusion equation with increase in the amount of substance, and its application to a biological problem*, *Moscow Univ. Math. Bull.* **A1** (Engl. Transl) (1937) 1.
- [48] K. Golec-Biernat and M. Wüsthoff, *Saturation effects in deep inelastic scattering at low Q^2 and its implications on diffraction*, *Phys. Rev. D* **59** (1998) 014017 [[arXiv.hep-ph/9807513](#)].
- [49] J. L. Albacete, N. Armesto, J. G. Milhano, and C. A. Salgado, *Non-linear QCD meets data: A global analysis of lepton-proton scattering with running coupling BK evolution*, *Phys. Rev. D* **80** (2009) 034031 [[arXiv.0902.1112](#)].

- [50] W. Xiong et al., *A small proton charge radius from an electron–proton scattering experiment*, *Nature* **575** (2019) 147.
- [51] J L Abelleira Fernandez et al., [LHeC Collaboration], *A Large Hadron Electron Collider at CERN Report on the Physics and Design Concepts for Machine and Detector*, *J. Phys. G: Nucl. Part. Phys.* **39** (2012) 075001 [[arXiv.1206.2913](#)].
- [52] P. Agostini et al., *The Large Hadron–Electron Collider at the HL-LHC*, *J. Phys. G: Nucl. Part. Phys.* **48** (2021) 110501 [[arXiv.2007.14491](#)].
- [53] A. Accardi et al., *Electron-Ion Collider: The next QCD frontier*, *Eur. Phys. J. A* **52** (2016) 268.
- [54] A. Abada et al., [FCC Collaboration], *FCC Physics Opportunities*, *Eur. Phys. J. C* **79** (2019) 474.