Uniform error bound for PCA matrix denoising

Xin T. Tong, Wanjie Wang, and Yuguan Wang

August 29, 2024

Abstract

Principal component analysis (PCA) is a simple and popular tool for processing highdimensional data. We investigate its effectiveness for matrix denoising. We consider the clean data are generated from a low-dimensional subspace, but masked by independent highdimensional sub-Gaussian noises with standard deviation σ . Under the low-rank assumption on the clean data with a mild spectral gap assumption, we prove that the distance between each pair of PCA-denoised data point and the clean data point is uniformly bounded by $O(\sigma \log n)$. To illustrate the spectral gap assumption, we show it can be satisfied when the clean data are independently generated with a non-degenerate covariance matrix. We then provide a general lower bound for the error of the denoised data matrix, which indicates PCA denoising gives a uniform error bound that is rate-optimal. Furthermore, we examine how the error bound impacts downstream applications such as clustering and manifold learning. Numerical results validate our theoretical findings and reveal the importance of the uniform error.

1 Introduction

In the modern era, data is often referred to as "the new gold". Rich data with rapidly increasing statistical methods present us with powerful tools for extracting valuable information and explaining scientific problems. However, the process of collecting data inevitably introduces noise, which poses a significant challenge. While statistical methods typically exhibit stability in the presence of weak noise, they may struggle to perform well when the noise surpasses the signal present in clean data. This issue becomes particularly pronounced in the realm of high-dimensional data where each dimension of the data point is corrupted by noise. As the number of dimension grows, the overall noise also grows, which further exacerbates the curse of dimensionality [\[Donoho et al.,](#page-21-0) [2000\]](#page-21-0) when we try to analyze high-dimensional data given a small number of observations.

Consider *n* observed data points $Z_i = X_i + \xi_i \in \mathbb{R}^d$, where X_i is the underlying truth and ξ_i is an independent sub-Gaussian noise vector with covariance $\sigma^2 I$, $i \in [n]$. X_i , also called a *clean* data point, is unknown to us and algorithms are conducted on Z_i s. While the observed data Z_i usually fall into a high-dimensional space, we assume the intrinsic dimension of the clean data is low, i.e. it can be embedded into a low-dimensional subspace. It is common to assume that $||X_i||$'s are bounded by a constant; see [Von Luxburg et al. \[2008\]](#page-22-0). Without loss of generality, we assume $||X_i|| \leq 1$. In practice, this can be achieved by data normalization.

To evaluate the data quality of Z_i , we introduce the signal-to-noise ratio (SNR). SNR is a common metric that measures the relative strength of the signal when compared with the noise. In this simple setting, the SNR is given by:

$$
SNR = \max_{i \in [n]} \frac{\|X_i\|^2}{\mathbb{E}[\|\xi_i\|^2]} \le \frac{1}{\sigma^2 d}.
$$

When the dimension d increases, the SNR deteriorates and tends towards zero. With a low SNR, analyzing data directly based on Z_i will induce unsatisfactory results. Naturally, we seek to denoise Z_i first to improve the accuracy of data analysis. This procedure is known as matrix denoising in the literature (see for example [\[Donoho and Gavish, 2014\]](#page-20-0)), and we introduce it in the following section.

1.1 PCA for denoising

Suppose the clean data points X_i are distributed in a low-dimensional subspace with a dimension of r, where $r \ll d$. A direct idea to recover X_i is to use the singular value decomposition (SVD) of the noisy data matrix formed by Z_i .

We introduce the matrix form for the data points. Let $Z = [Z_1, \ldots, Z_n]^T$, $X = [X_1, \ldots, X_n]^T$ and $E = [\xi_1, \ldots, \xi_n]^T$ be the matrices formed by the noisy data points, clean data points, and pure noise, respectively. According to the assumption, $rank(X) = r$. We start from the following hypothetical denoising procedure and then propose the algorithm. Let the SVD of X be denoted as $X = U\Lambda V^T$, where $U \in \mathbb{R}^{n \times r}$, $\Lambda \in \mathbb{R}^{r \times r}$, and $V \in \mathbb{R}^{d \times r}$. The columns of V span the rdimensional subspace in which X lies, and VV^T is the projection operator onto this subspace. By applying this projection operator to Z_i , we obtain:

$$
VV^{T}Z_{i} = X_{i} + VV^{T}\xi_{i}, \text{ where } \mathbb{E}[\|VV^{T}\xi_{i}\|^{2}] = \sigma^{2} \text{tr}(VV^{T}) = r\sigma^{2}.
$$

Thus, when the underlying dimension $r = O(1)$ and $r \ll d$, the noise in the projection $VV^T Z_i$ is significantly weaker than the noise in Z_i . the associated SNR of the projected data $VV^T Z_i$ is of order $1/(\sigma^2 r)$, which can be sufficiently strong to yield accurate inference results. Therefore, by leveraging the SVD and performing the projection onto the low-dimensional subspace, we can effectively denoise the data and obtain accurate estimates.

In practice, there is no access to V . Therefore, we estimate it using the SVD of the noisy data matrix Z. The SVD of Z can be expressed as $Z = \hat{U} \hat{\Lambda} \hat{V}^T$, where typically Z has full rank due to the presence of noise. To focus on the most significant components of the data, we select only the first r columns of \hat{V} , denoted as \hat{V}_r , corresponding to the largest r singular values. By \hat{V}_r , we project the noisy data points Z_i onto the estimated subspace, resulting in the denoised estimates \hat{X}_i . The projection is given by:

$$
\hat{X}_i = \hat{V}_r \hat{V}_r^T Z_i, \quad i \in [n].
$$

The columns of \hat{V}_r can be interpreted as the r directions that capture the most variability of the data points Z_i . Therefore, they are often referred to as the principal directions, and the resulting \hat{X}_i are known as the principal components. This approach is commonly known as principal component analysis (PCA). We call it the PCA-denoising algorithm, presented in Table [1.](#page-1-0)

Utilizing PCA for noise reduction is not a new concept. It was first introduced in multivariate statistical analysis and then explored in various fields. For example, [Shepard \[1962\]](#page-22-1) introduced the use of PCA for multidimensional scaling and distance estimation. In the field of image processing, [Singh and Harrison \[1985\]](#page-22-2) applied PCA to denoise images. Discussions in Section [1.3](#page-2-0) provide more details of the related literature and results. We also refer interested readers to surveys and textbooks for more comprehensive lists [\[Jolliffe, 2005,](#page-21-1) [Abdi and Williams, 2010,](#page-20-1) [Chen et al., 2021\]](#page-20-2).

The denoised data X can be applied to various applications, such as empirical risk minimization, clustering, manifold learning and so on. The denoising step largely improves the performance of algorithms in these fields. More discussions can be found in our Section [4.](#page-14-0)

Algorithm 1 PCA-denoising

Input: Data $Z \in \mathbb{R}^{n \times d}$, dimension r. **Output:** Denoised data $\hat{X} \in \mathbb{R}^{n \times d}$. 1: Find the SVD of Z as $Z = \hat{U} \hat{\Lambda} \hat{V}^T$. 2: Denote the first r columns of \hat{V} as $\hat{V}_r \in \mathbb{R}^{d \times r}$. 3: Let $\hat{X} = Z \hat{V}_r \hat{V}_r^T$.

1.2 Our main interest and contribution

One crucial question for matrix denoising is assessing the accuracy of \hat{X}_i , i.e. the distance between the estimate \hat{X}_i and the clean data X_i for all $i \in [n]$. Most existing theoretical analysis of PCA focuses on the Frobenius distance between the two matrices \hat{X} and X, which is the squared root of the sum of squared ℓ_2 error among all data points. Our goal is to obtain a uniform error bound across all data points, which allows for individual statistical analysis on each sample. Specifically, we aim to establish the following $\ell_2 \to \ell_\infty$ or uniform error bound for the PCA-based denoising algorithm,

$$
\|\hat{X} - X\|_{2,\infty} := \max_{a \in \mathbb{R}^d, a \neq 0} \frac{\|(\hat{X} - X)a\|_{\infty}}{\|a\|_2} = \max_{i \in [n]} \|X_i - \hat{X}_i\| = O(\sigma \log n)
$$
(1)

with high probability, when $d = cn$ with c and σ being some absolute constants. Here the notation $O(\cdot)$ hides a factor that may depend on the low dimension r. The equivalence between $||\hat{X} - X||_{2,\infty}$ and $\max_{i \in [n]} ||X_i - \hat{X}_i||$ can be found in [Cape et al. \[2019a,](#page-20-3)[b\]](#page-20-4), which provides further theoretical insights into our results.

To understand our goal [\(1\)](#page-2-1), we consider a special low-dimensional case where the observed data Z_i also has a dimension of r, i.e., $d = r$. In this case, the noise E_i has a low dimension of r and $||E|| = O(\sigma^2 r)$ is small. Therefore, the dimension reduction is not necessary and we can directly use the observed data Z_i as an estimate of the clean data X_i . By a union bound, we can find the estimation error is $||Z - X||_{2,\infty} = ||E||_{2,\infty} = O(\sigma \log n)$, the same as [\(1\)](#page-2-1). Therefore, our goal [\(1\)](#page-2-1) implies that the PCA-denoising estimates \hat{X}_i achieve the same level of accuracy as in the low-dimensional case. In other words, the PCA-denoising step essentially removes the curse of dimensionality.

In this paper, we explore the estimate [\(1\)](#page-2-1) from several perspectives. Here is a summary of our main findings and the organisation of this paper:

- 1. In Section [2.1,](#page-5-0) Theorem [1](#page-5-1) establishes a general form of [\(1\)](#page-2-1) for arbitrary d and σ under the condition $||X_i|| \leq 1$. When assumptions $n \geq d$ and the r-th largest singular value of X that $\lambda_r(X) \geq c_X \sqrt{n}$ holds for an absolute constant c_X , we show that the estimate [\(1\)](#page-2-1) holds exactly. Our result does not impose any assumptions on the correlation structure of clean $data X_i$.
- 2. In Section [2.3,](#page-12-0) we investigate the sufficient conditions that the assumption $\lambda_r(X) \geq c_X \sqrt{n}$ holds. By the random matrix theory, we demonstrate that the covariance matrix of X with a non-zero r-th eigenvalue will suffice. We illustrate this assumption by a zigzag line example, which is motivated by temporal and spatial data sets.
- 3. Section [3](#page-14-1) presents a general lower bound on the signal-to-noise ratio and sample size n to ensure that the average error is no larger than any constant $\epsilon > 0$. The lower bound highlights that PCA-denoising has the rate-optimal signal-to-noise ratio and sample size requirement.
- 4. In Section [4,](#page-14-0) we demonstrate the practical implications of the uniform error bound $||\overline{X} X\|_{2,\infty}$ in various downstream applications. Assuming $\|X-X\|_{2,\infty} \leq \epsilon$, we provide performance guarantees for applications such as clustering and manifold learning.
- 5. Finally, in Section [5,](#page-17-0) we provide some numerical simulations to support our theoretical findings. We consider a clustering task on high-dimensional data sampled from two separated zigzag lines. For this data, PCA-denoising yields the uniform error bound [\(1\)](#page-2-1), and the denoised data enables efficient spectral clustering. We also show that data with a small "average error" is not sufficient to guarantee good clustering results for every sample in this task.

1.3 Related literature

Our findings reside at the intersection of PCA and matrix denoising, where plenty of related results exist in the literature. In this section, we will provide a brief overview of the relevant literature from the perspectives of PCA and matrix denoising. For ease of discussion, we would assume $n \geq d$ in this section, which is a common setting for many works in the discussion.

1.3.1 Comparing with PCA literature

Due to the wide range of applications for PCA, there is numerous of literature on its design and applications [\[Abdi and Williams, 2010\]](#page-20-1). The earliest works can be traced back to the 1960s [\[Rao,](#page-22-3) [1964,](#page-22-3) [Jolliffe, 1972\]](#page-21-2), where the discussions focus on multivariate statistical analysis. However, a rigorous understanding of PCA in high-dimensional settings emerged much later, mostly in the last 15 years. In the theoretical analysis of PCA, most studies have focused on the accuracy of subspace recovery, i.e. $\|\hat{V}_r\hat{V}_r^T - V V^T\|_*$. Here $\|\cdot\|_*$ is some operator norm, where ℓ_2 -operator norm is used in most classical settings. Denote $\mathcal{E}_V = V_r \hat{V}_r^T - V V^T$ for short. Using the eigenvector perturbation results like [Davis and Kahan \[1970\]](#page-20-5), [Wedin \[1972\]](#page-23-0), [Stewart and Sun \[1990\]](#page-22-4), [Bhatia](#page-20-6) [\[2013\]](#page-20-6), [Yu et al. \[2015\]](#page-23-1), and random matrix theory, an upper bound of the form $\|\mathcal{E}_V\|^2 \leq O(\sigma^2)$ can often be obtained, with additional conditions on X [\[Vu, 2011,](#page-23-2) [Chen et al., 2021\]](#page-20-2).

In particular, [Johnstone and Lu \[2009\]](#page-21-3) discussed the consistency of PCA recovery when the underlying dimension $r = 1$ and the single principal component is sparse. Assuming the existence of a spectral gap, [Cai and Zhang \[2018\]](#page-20-7) discussed the optimality of PCA recovery, in the context of the ℓ_2 operator norm $\|\mathcal{E}_V\|$ and Frobenius norm $\|\mathcal{E}_V\|_F$. It is worth mentioning that r can grow as $O(n)$ in this work. In [Zhang et al. \[2022\]](#page-23-3), the results were extended to the setting where the data distribution is heteroskedastic. Other than \mathcal{E}_V , V_r is also discussed in the literature. [Ding](#page-20-8) [\[2020\]](#page-20-8) studied the limiting distribution of \hat{V}_r . [Fan et al. \[2018\]](#page-21-4) and [Abbe et al. \[2020\]](#page-20-9) have shown the ℓ_{∞} error of estimated eigenvectors in \hat{V}_r is $O(1/n)$, assuming the true eigenvectors in V have ℓ_{∞} norm being $O(1/\sqrt{n})$. [Cape et al. \[2019b\]](#page-20-4) studied the distribution of $\hat{V}_r - VO$, where O is an orthogonal matrix, and establishes an error bound in $\|\cdot\|_{2,\infty}$ norm. In [Reiss and Wahl \[2020\]](#page-22-5), the error is studied when VV^T is replaced by a rank-r projection P that minimizes $\mathbb{E}||Z - PZ||^2$.

The model setting of this paper follows a similar line as these works. In particular, we assume the existence of a spectral gap of X , which enables the spectral perturbation analysis. Such a spectral gap is necessary in the existing works discussed above. One of the key steps in the proof, Proposition [1,](#page-10-0) can be interpreted as an application of the leave-one-out method discussed in [Chen](#page-20-2) [et al. \[2021\]](#page-20-2). The difference between this paper and these works mainly lies in the errors of interest, as explained below.

Using the upper bound on $\|\mathcal{E}_V\|$, we can straightforwardly obtain the denoising accuracy of PCA for a new data point $Z_{n+1} = x + \xi_{n+1}$, which is often referred as the test error in statistics [\[Hastie et al., 2009\]](#page-21-5). In particular, we can define and calculate

Test
$$
\hat{V}
$$
 := $\max_{x \in \text{span}(V), ||x|| = 1} \mathbb{E}_{\xi} [||\hat{V}_r \hat{V}_r^T (x + \xi_{n+1}) - x||^2]$
\n= $\max_{x \in \text{span}(V), ||x|| = 1} \mathbb{E}_{\xi} [||\mathcal{E}_V x||^2 + ||\hat{V}_r \hat{V}_r^T \xi_{n+1}||^2] = O(\sigma^2).$

This noise level is the same as our goal in (1) , if the $\log n$ factor is disregarded. However, in the proof, the independence between ξ_{n+1} and \hat{V}_r is required to obtain the bound on $\|\hat{V}_r\hat{V}_r^T\hat{\xi}_{n+1}\|^2$. Without this independence, such as in the calculation of $\|\hat{V}_r\hat{V}_r^T\hat{\zeta}_k\|^2$ for $k \in [n]$, this bound does not hold. In other words, bounding $\|\mathcal{E}_V\|$ does not directly lead to the desired bound [\(1\)](#page-2-1).

While Test_{\hat{V}} is already useful in many situations, it imposes limitations in practical applications. To ensure the independence condition between \hat{V}_r and the data points to be projected, one has to split the samples into two sets: one training data set to obtain the projection $V_r \hat{V}_r^T$, and another set where this projection is applied. Test_{\hat{V}} cannot be applied if the projection is on the training data set. This data-splitting approach leads to two problems: 1) It reduces the sample size, resulting in a loss of estimation accuracy, which is undesirable when the original dataset has a limited number of samples. 2) In many unsupervised learning tasks, statistical inference on the training data set itself is crucial. For example, we want to classify all data points in the clustering problem. Yet the clustering error on the training data cannot be evaluated using Test $_{\hat{V}}$.

In contrast, a uniform bound of the form [\(1\)](#page-2-1) guarantees the result without data splitting. We can obtain V_r from all samples and apply the projection to all samples, with a uniform error bound. It is more accurate with a larger sample size and allows us to carry out unsupervised operations on all the samples.

1.3.2 Comparing with matrix denoising literature

In the matrix denoising literature, the main interest is to find an estimate $\hat{X}(Z)$ so that the Frobenius norm $||\hat{X}(Z) - X||_F$ can be well bounded. Various approaches have been introduced to tackle this problem. [Donoho and Gavish \[2014\]](#page-20-0) considered using the minimizer of a regularized loss function $||Z - \hat{X}||_F^2 + \lambda ||\hat{X}||_*$, where $||\hat{X}||_*$ is the nuclear norm of \hat{X} . The approach has been proved to have a rate-optimal mean squared error in the Frobenius norm. The shrinkage method instead considers estimators of the form $\hat{X} = \hat{U} g(\hat{\Lambda}) \hat{V}^T$, where $g(\hat{\Lambda})$ is a diagonal matrix with diagonal entries being $g(\hat{\lambda}_i)$. The main interest is to find a good choice of g. Assuming the distribution of X is known, [Nadakuditi \[2014\]](#page-21-6) provided the optimal g for the Frobenius norm. When X is a spiked covariance matrix, most eigenvalues of the estimator \hat{X} are 1. Focusing on the spectrum of $\ddot{X} - X$, which has very few non-zeros, [Donoho et al. \[2018\]](#page-21-7) provided the optimal g under the ℓ_p norm of the spectrum for any general p. [Leeb \[2021\]](#page-21-8) extended this discussion to more applied scenarios, where some data can be missing or the singular vectors are sparse. Under the assumption that X is low-rank and rotation invariant (or the singular vectors are sparse), [Ding \[2020\]](#page-20-8) designed a stepwise SVD algorithm that could estimate the singular vectors by going through the data sequentially. [Montanari et al. \[2018\]](#page-21-9) discussed the scenario where the noise is from an unknown distribution and suggested a kernelized estimator to do shrinkage.

Compared with these methods, the PCA-denoising approach can be seen as a simple shrinkage method, with $g(\lambda_k) = 1_{k \leq r} \lambda_k$. It is much simpler and widely adopted in practice, with few requirements on the clean data X . Therefore, the theoretical analysis of it is worth special interest. For example, [Cai and Zhang \[2018\]](#page-20-7) has discussed the implication of PCA subspace recovery on matrix denoising, and we compared with it in Section [1.3.1.](#page-3-0)

All existing analyses of matrix denoising focus on bounding the error in the Frobenius norm (or ℓ_p generalization of it; see [Donoho et al. \[2018\]](#page-21-7)) which can also be seen as the average mean square error over all data points:

$$
\frac{1}{n}\sum_{i=1}^{n}||X_i - \hat{X}_i||^2 = \frac{1}{n}||X - \hat{X}||_F^2 = O(\sigma^2).
$$
\n(2)

This average error is at the same order as the uniform error bound in [\(1\)](#page-2-1) up to a logarithm term. But the error bound of form $||\hat{X} - X||_{2,\infty}$ provides a stronger mathematical guarantee as it captures the distribution of errors across all data points, while the error bound in Frobenius norm allows for some outliers with large individual errors. This distinction becomes crucial when analyzing the estimates on individual data points and nonlinear statistical models.

For instance, in the context of clustering, using the Frobenius error bound would only allow us to establish that the proportion of incorrectly classified data points tends to zero, i.e., the error rate goes to zero. However, it does not provide insights into the exact number of errors, which is often referred to as "strong consistency" in recent statistical literature [\[Abbe and Sandon, 2015,](#page-20-10) [Fan et al., 2018,](#page-21-4) [Hu and Wang, 2024\]](#page-21-10). On the other hand, a uniform error bound enables us to establish such strong consistency results, as demonstrated by Corollary [3](#page-15-0) in Section [4.1.](#page-14-2)

Such outliers with big errors may also cause challenges for the statistical models with nonlinear and local dependency on the data input, such as K-nearest-neighbor and neural networks. In Section [4.2,](#page-15-1) we demonstrate the error of manifold learning using our new uniform error bound, which extends the results in [Von Luxburg et al. \[2008\]](#page-22-0) to the settings where data is corrupted by high-dimensional noise. In Section [5,](#page-17-0) we provide numerical simulations on graphical Laplacian spectral clustering. There, we show that PCA-denoising leads to stable clustering results, while data with the same Frobenius error fails to guarantee satisfactory clustering outcomes.

The analysis technique of this paper differs from these works on matrix denoising, as it is closer to the techniques used in PCA analyses. This is reflected by the assumptions we made for X. In to the techniques used in FCA analyses. This is renected by the assumptions we made for λ . In particular, our result requires the r-th singular value of X to scale like \sqrt{n} . Such spectral gap requirement is not needed in [Donoho and Gavish \[2014\]](#page-20-0) and [Donoho et al. \[2018\]](#page-21-7). [Nadakuditi](#page-21-6) [\[2014\]](#page-21-6) and [Bao et al. \[2021\]](#page-20-11) assumed the existence of a spectral gap, but its characterization is sharper than the one used in this paper. On the other hand, such an assumption allows this paper

and Cai and Zhang $[2018]$ to discuss general cases where d and n diverge at different speeds, while most matrix denoising works focus on the scenario $n \times d$.

1.4 Notations

For any matrix A, we use A_i to denote the *i*-th row vector of A and $A_{i,j}$ to denote the (i, j) -th entry. Denote $\lambda_k(A)$ the k-th largest singular value of A. Denote I_k to be the $k \times k$ identity matrix. We denote the ℓ_2 -operator norm of A as $||A||$ and the Frobenius norm of A as $||A||_F$.

To generalize our discussion to other noise types, we introduce the notion of sub-Gaussian random variables. We say a random variable Z is sub-Gaussian $(0, \sigma^2, K_{\psi_2})$ if

$$
\mathbb{E}[Z] = 0, \quad \mathbb{E}[Z^2] = \sigma^2, \quad K_{\psi_2} \ge \sup_{p \ge 1} \mathbb{E}[|Z/\sigma|^p]^{1/p}.
$$

Given a constant C , we say C is absolute if it does not depend on any other constants. We write $C = C(x, y)$ if it only depends on constants x and y. In this paper, without further statements, we fix the sub-Gaussian parameter K_{ψ_2} and the rank r as absolute constants. For two series a_n and b_n , we say $a_n \lesssim b_n$ or $a_n = O(b_n)$ if there is a constant $C = C(r, K_{\psi_2})$ so that $\limsup_{n \to \infty} a_n/b_n \leq C$. Similarly, we have $a_n \geq b_n$. We say $a_n \geq b_n$ if there is a constant C, such that $a_n \leq Cb_n$ and $b_n \leq Ca_n$ when n is large enough. Finally, we use the notation $[N] := \{1, \ldots, N\}$ for any integer N .

2 Performance bounds for PCA denoising

2.1 Uniform bounds for PCA-denoising

In this section, we establish the upper bound for the uniform error $\|\hat{X} - X\|_{2,\infty}$, where \hat{X} is obtained from Algorithm [1.](#page-1-0) For notational simplicity, we write $\lambda_r = \lambda_r(X)$, which is the r-th largest singular value of X.

Theorem 1. Suppose $X \in \mathbb{R}^{n \times d}$ has rank r, where each row X_i^T is bounded by $||X_i|| \leq 1$. Suppose $E \in \mathbb{R}^{n \times d}$ has independent entries being sub-Gaussian $(0, \sigma^2, K_{\psi_2})$ distributed with $\sigma \leq 1$. Let \hat{X} be the denoised data matrix from Algorithm [1.](#page-1-0) Then there are constants $n_0, C_1 = C_1(K_{\psi_2})$ and $C_2 = C_2(K_{\psi_2}, r)$, so that if

$$
n > n_0, \quad \lambda_r > 1 + C_1 \sigma(\sqrt{n} + \sqrt{d}), \tag{3}
$$

the following holds with probability $1 - O(1/n)$,

$$
\begin{split} \|\hat{X} - X\|_{2,\infty} &\leq C_2 \min \left\{ \frac{\sigma(\sqrt{n} + \sqrt{d})}{\lambda_r} + \frac{n\sigma\sqrt{\log n}}{\lambda_r^2}, \frac{\sigma}{\lambda_r^2} (\sqrt{nd} + n\sqrt{\log n}) \right\} \\ &+ \frac{C_2 \sigma}{\lambda_r^4} \sqrt{\log n} (n + \sigma^2 n\sqrt{n} + \sigma^2 d\sqrt{n}) (1 + \sigma\sqrt{\log n}) (1 + \sigma(\sqrt{d} + \sqrt{\log n})). \end{split}
$$

Further, if $\lambda_r \geq c_X \sqrt{n}$ holds for an absolute constant $c_X > 0$, then the bound can be shortened as below for another $C_2 = C_2(K_{\psi_2}, r, c_X)$

$$
\|\hat{X} - X\|_{2,\infty} \le \begin{cases} C_2 \cdot \left(\sqrt{\frac{d}{n}} \sigma (1 + \frac{d}{n} \sigma^3 \sqrt{\log n} + \frac{d}{n} \sigma^4 \log n) + \sqrt{\log n} \sigma \right), & d \gtrsim n; \\ C_2 \sqrt{\log n} \sigma (1 + \sigma^4 \sqrt{\frac{d \log n}{n}}), & d \lesssim n. \end{cases}
$$

If we further assume $c_d = d/n$ and σ are absolute constants, then the bound can be shortened as below for another $C_2 = C_2(K_{\psi_2}, r, c_d, \sigma)$

$$
\|\hat{X} - X\|_{2,\infty} \le C_2 \sigma \log n.
$$

Theorem [1](#page-5-1) presents three estimation error bounds because of the trade-off between formula complexity and generality of settings, the first estimate being under the most general setting and the last estimate being under a strict but canonical setting. The first error bound allows the key constants n, σ , and d to be free from each other, which covers most settings. For example, it can cover the case $d \gg n$ in [Cai and Zhang \[2018\]](#page-20-7) and the case $d\sigma \approx 1$ in [Montanari et al. \[2018\]](#page-21-9). The error bound can be simplified under specific settings. The second error bound is a simplification when $\lambda_r \geq c_X \sqrt{n}$. Compared to the requirement $\lambda_r > 1 + C_1 \sigma(\sqrt{n} + \sqrt{d})$, this is a more strict condition since it avoids the case that σ decays when n increases. The last estimate achieves our goal in [\(1\)](#page-2-1). To achieve it, we assume $d \simeq n$ and $\sigma \simeq 1$, which is a canonical setting and can be found in for example [Donoho and Gavish \[2014\]](#page-20-0) and [Abbe et al. \[2020\]](#page-20-9).

Theorem [1](#page-5-1) has two assumptions to achieve the uniform bound on the error. The first assumption is the clean data X, that $\max_{i\in[n]} ||X_i|| \leq 1$. Such a condition is common in manifold analysis, such as Von Luxburg et al. $[2008]$. It can be achieved in any data set X by dividing each data point by a large constant $C \ge \max_{i \in [i]} ||X_i||$. This assumption also makes the discussed error the relative error. The second assumption is the spectral gap [\(3\)](#page-5-2). It is an essential requirement that the signal level in X should be no smaller than the noise level in E . Theorem [3](#page-14-3) below will show the necessity of such a requirement.

Before we provide the formal proof, it might be worth doing a naive one using $\|\hat{V}_r\hat{V}_r^T - VV^T\| =$ $O(\sigma)$ from the standard Davis–Kahan Theorem. One may attempt to approach the following bound (the authors tried this in the beginning)

$$
||X_i - \hat{X}_i|| = ||VV^T X_i - (\hat{V}_r \hat{V}_r^T)(X_i + \xi_i)||
$$

\n
$$
\leq ||(VV^T - \hat{V}_r \hat{V}_r^T)X_i|| + ||\hat{V}_r \hat{V}_r^T \xi_i|| = ||\hat{V}_r \hat{V}_r^T \xi_i|| + O(\sigma).
$$

The main issue comes from the term $\|\hat{V}_r \hat{V}_r^T \xi_i\|$. A simple bound that $\|\hat{V}_r \hat{V}_r^T \xi_i\| \leq \|\xi_i\| = O(\sigma)$ √ d) is too loose when $d \to \infty$. One may notice that the true projection VV^T causes $||VV^T \xi_i|| \approx$ $\sqrt{\text{tr}(V V^T)} = \sigma \sqrt{r}$ and expect a similar bound for $\|\hat{V}_r \hat{V}_r^T \xi_i\|$. However, obtaining such a bound is nontrivial because ξ_i is dependent on \hat{V}_r . This is exactly where the mathematical challenge lies. We introduce a proposition in Section [2.2](#page-10-1) to solve this problem by the leave-one-out trick.

Proof. Note that $U\Lambda O = XVO$ holds for any orthonormal matrix $O \in \mathbb{R}^{d \times d}$. Recall that $||X_i|| \le$ $1, \|\hat{V_r}\| \leq 1, \|V\| \leq 1.$ Therefore, we have

$$
||X_i - \hat{X}_i|| = ||VV^T X_i - (\hat{V}_r \hat{V}_r^T)(X_i + \xi_i)||
$$

\n
$$
\leq ||\hat{V}_r \hat{V}_r^T \xi_i|| + ||(VV^T - \hat{V}_r \hat{V}_r^T)X_i||
$$

\n
$$
\leq ||\hat{V}_r^T \xi_i|| + ||(VO - \hat{V}_r)\hat{V}_r^T X_i|| + ||VO(O^T V^T - \hat{V}_r^T)X_i||
$$

\n
$$
\leq \underbrace{||\hat{V}_r^T \xi_i||}_{(a)} + 2\underbrace{||VO - \hat{V}_r||}_{(b)}.
$$
\n(4)

We will discuss each part of the decomposition. We begin with part (b) by the Davis–Kahan Theorem, and then the much more challenging part (a).

Part (b). Consider the self-adjoint extension $\mathcal{E}(X) := \begin{bmatrix} 0, & X \\ x^T & 0 \end{bmatrix}$ X^T , 0 of X . First note that if $X = U\Lambda V^T$ is the SVD decomposition, $\mathcal{E}(X)$ has eigenvalue decomposition of form

$$
\mathcal{E}(X) = \begin{bmatrix} 0, & X \\ X^T, & 0 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} U, & -U \\ V, & V \end{bmatrix} \begin{bmatrix} \Lambda, & 0 \\ 0, & -\Lambda \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} U^T, & V^T \\ -U^T, & V^T \end{bmatrix}.
$$
 (5)

Since E has independent sub-Gaussian rows, Theorem 5.39 of [Vershynin \[2010\]](#page-22-6) indicates that there is a constant $C_1 = C_1(K_{\psi_2})$ so that with probability $1 - O(1/n^2)$,

$$
\|\mathcal{E}(X+E)-\mathcal{E}(X)\|=\|\mathcal{E}(E)\|\leq 2\|E\|\leq C_1\sigma(\sqrt{d}+\sqrt{n}).
$$

Therefore, by the Davis–Kahan Theorem [\[Davis and Kahan, 1970\]](#page-20-5), there exists an orthogonal matrix O and constant $C_2 = C_2(K_{\psi_2}, r)$, so that

$$
\|\hat{V}_r - VO\| \le \left\| \begin{bmatrix} \hat{U}_r \\ \hat{V}_r \end{bmatrix} - \begin{bmatrix} UO \\ VO \end{bmatrix} \right\| \le \frac{C_2 \sigma (\sqrt{d} + \sqrt{n})}{\lambda_r}.
$$
 (6)

An alternative way to bound part (b) is to note that V are the eigenvectors of $X^T X$ and \hat{V} are the eigenvectors of $Z^T Z = X^T X + \Delta$, where $\Delta = X^T E + E^T X + E^T E$. Then the difference

$$
\|\Delta\| \le \|X^T E + E^T X + E^T E\| \le 2\|X^T E\| + \|E^T E\|.
$$

We bound the right-hand side. Note that $X^T E = V \Lambda U^T E$, where each column of $U^T E / \sigma \in \mathbb{R}^{r \times d}$ is an independent, centered, and isotropic sub-Gaussian vector. By Theorem 5.39 of [Vershynin](#page-22-6) [\[2010\]](#page-22-6), with probability at least $1 - O(1/n^2)$, there are updated $C_1(K_{\psi_2})$ and $C_2(K_{\psi_2}, r)$, so that $||U^T E|| \le \sigma(\sqrt{d} + C_1\sqrt{r} + \sqrt{\log n}),$ and

$$
||XTE|| \le ||\Lambda|| ||UTE|| \le \frac{1}{3}\sigma\sqrt{n}C_2(\sqrt{d} + \sqrt{\log n}).
$$

Again, the same theorem can be applied to E, which shows that with probability at least $1 O(1/n^2),$ √ √

$$
||ETE|| \le C_1^2 \sigma^2 (\sqrt{n} + \sqrt{d} + \sqrt{2 \log n})^2 \le 4C_1^2 \sigma^2 (\sqrt{n} + \sqrt{d})^2.
$$

Combine them. It follows that we can update constant $C_2(K_{\psi_2}, r)$ so that

$$
\|\Delta\| \leq \frac{2}{3}C_2(\sigma\sqrt{nd} + \sigma\sqrt{n\log n}) + 4C_1^2\sigma^2(n + 2\sqrt{nd} + d)
$$

$$
\leq \frac{2}{3}C\sigma(\sqrt{nd} + \sqrt{n\log n} + \sigma n + \sigma\sqrt{nd} + \sigma d)
$$

$$
\leq C_2\sigma(\sqrt{nd} + \sqrt{n\log n} + \sigma n).
$$

The last step is achieved because we assume σ √ $\overline{d} \leq \lambda_r \leq ||X|| = \sqrt{n}$. Therefore, by the Davis-Kahan Theorem [\[Davis and Kahan, 1970\]](#page-20-5), there exists an orthogonal matrix O and updated $C_2(K_{\psi_2},r)$, so that

$$
\|\hat{V}_r - VO\| \le \frac{\|\Delta\|}{\lambda_r^2} \le \frac{C_2\sigma}{\lambda_r^2} (\sqrt{nd} + \sqrt{n\log n} + \sigma n). \tag{7}
$$

Since [\(6\)](#page-7-0) and [\(7\)](#page-7-1) both hold, it suffices to use the minimum of them as the upper bound of part (b).

Part (a). $\|\xi_i^T \hat{V}_r\|$ is the most challenging part because of the correlation between ξ_i and \hat{V}_r . If we ignore the correlation and bound it by $\|\xi_i\| \|\hat{V}_r - VO\|$, then the bound is of order $\sigma^2 \sqrt{d}$. For high-dimensional data, this d can be very large. Therefore, we need a delicate analysis of this term.

First, since the rows of E are independent sub-Gaussian distributed, by Theorem 5.39 in [Vershynin \[2010\]](#page-22-6), there is a constant $C_1 = C_1(K_{\psi_2})$, so that with probability at least $1 - O(1/n^3)$,

$$
|X_i^T \xi_i| \le \sigma (1 + \frac{1}{6}C_1(1 + \sqrt{\log n})) \le \frac{1}{3}C_1\sigma (1 + \sqrt{\log n}),
$$

$$
||\xi_i|| \le \sigma (\sqrt{d} + \frac{1}{6}C_1(1 + \sqrt{\log n})) \le \frac{1}{3}C_1\sigma (\sqrt{d} + \sqrt{\log n}),
$$

$$
||E|| \le \sigma (\sqrt{d} + \frac{1}{6}C_1(\sqrt{n} + \sqrt{\log n})) \le \frac{1}{3}C_1\sigma (\sqrt{d} + \sqrt{n}).
$$

We will define the event that all these hold as A . The following discussions are conducted when A happens. Let $\hat{\lambda}_k = \lambda_k(Z)$ be the k-th largest singular value of $Z = X + E$ and \hat{v}_k be the

corresponding right singular vector in \hat{V} . By Weyl's inequality, the singular values follow that, for $k \leq r$,

$$
\sqrt{n} + \frac{1}{3}C_1\sigma(\sqrt{n} + \sqrt{d}) \ge \hat{\lambda}_k \ge \lambda_r - \frac{1}{3}C_1\sigma(\sqrt{n} + \sqrt{d}), \quad \hat{\lambda}_{r+1} \le \frac{1}{3}C_1\sigma(\sqrt{n} + \sqrt{d}).
$$
 (8)

Consider the k-th entry of $\xi_i^T \hat{V}_r \in \mathbb{R}^r$. The k-th entry of $\xi_i^T \hat{V}_r$ is $\xi_i^T \hat{v}_k$. By definition of singular vectors, we have

$$
(X+E)^{T}(X+E)\hat{v}_{k} = \hat{\lambda}_{k}^{2}\hat{v}_{k}.
$$

Multiply ξ_i^T to both sides, and we have

$$
\hat{\lambda}_{k}^{2} \xi_{i}^{T} \hat{v}_{k} = \xi_{i}^{T} (X + E)^{T} (X + E) \hat{v}_{k}
$$
\n
$$
= \xi_{i}^{T} \left[(X_{i} + \xi_{i}) (X_{i} + \xi_{i})^{T} + \sum_{j \neq i} (X_{j} + \xi_{j}) (X_{j} + \xi_{j})^{T} \right] \hat{v}_{k}
$$
\n
$$
= \xi_{i}^{T} X_{i} X_{i}^{T} \hat{v}_{k} + \xi_{i}^{T} X_{i} \xi_{i}^{T} \hat{v}_{k} + \xi_{i}^{T} \xi_{i} X_{i}^{T} \hat{v}_{k} + ||\xi_{i}||^{2} \xi_{i}^{T} \hat{v}_{k} + \xi_{i}^{T} S_{i} \hat{v}_{k},
$$

where $S_i = \sum_{j \neq i} (X_j + \xi_j)(X_j + \xi_j)^T$. We rewrite this as

$$
(\hat{\lambda}_k^2 - \|\xi_i\|^2)\xi_i^T\hat{v}_k = \xi_i^T S_i\hat{v}_k + \xi_i^T X_i X_i^T \hat{v}_k + \xi_i^T X_i \xi_i^T \hat{v}_k + \xi_i^T \xi_i X_i^T \hat{v}_k. \tag{9}
$$

Now we analyze the terms on the right-hand side of [\(9\)](#page-8-0).

The first term $\xi_i^T S_i \hat{v}_k$ is the most challenging part since S_i is dependent on \hat{v}_k . Note that S_i is the version of $(X + E)^T (X + E)$ when sample *i* is deleted. Denote the eigenvalue decomposition of $S_i = \tilde{V} \tilde{\Lambda}^2 \tilde{V}^T$. Let \tilde{V}_r be the first r columns of \tilde{V} . Proposition [1](#page-10-0) in below states that if $y_k = \tilde{V}_r^T \hat{v}_k \in \mathbb{R}^r$, with probability $1 - O(1/n^2)$

$$
\|\hat{v}_k - \tilde{V}_r y_k\| = \|(I - \tilde{V}_r \tilde{V}_r^T)\hat{v}_k\| \le \frac{C_2}{\lambda_r^2} (1 + \sigma \sqrt{\log n}) (1 + \sigma(\sqrt{d} + \sqrt{\log n})).
$$
 (10)

We decompose \hat{v}_k into $\tilde{V}_r y_k$ and the remainder, and then it follows

$$
\xi_i^T S_i \hat{v}_k = \xi_i^T S_i \tilde{V}_r y_k + \xi_i^T S_i (\hat{v}_k - \tilde{V}_r y_k) = \xi_i^T \tilde{V}_r \tilde{\Lambda}_r^2 y_k + \xi_i^T S_i (\hat{v}_k - \tilde{V}_r y_k).
$$

Since \tilde{V} is based on S_i where sample i is deleted, so ξ_i is independent with \tilde{V}_r . Further, recall ξ_i is sub-Gaussian, using Hanson–Wright inequality (Theorem 1.1 in [Rudelson and Vershynin \[2013\]](#page-22-7)), there is an absolute constant $c > 0$ so that

$$
P(||\tilde{V}_r^T \xi_i||^2 - \sigma^2 r| > \sigma^2 t) \le 2 \exp\left(-c \min(\frac{t^2}{K_{\psi_2}^4 r}, \frac{t}{K_{\psi_2}^2})\right),\,
$$

We pick $t = \sqrt{r} K_{\psi_2}^2 \log n / \min\{c, \sqrt{c}\}\)$ then the probability will be less than $O(1/n^2)$ for sufficiently large *n*. In other words, we can update $C_2 = C_2(K_{\psi_2}, r)$

$$
\|\xi_i^T \tilde{V}_r\| \le \sigma C_2 (1 + \sqrt{\log n}),\tag{11}
$$

with $1 - O(1/n^2)$ probability. We assume [\(11\)](#page-8-1) takes place in the following discussion. Then consider $\tilde{\Lambda}_r^2$, which contains the eigenvalues of S_i . Because $S_i = (X + E)^T (I - e_i e_i^T)(X + E)$, its eigenvalues are dominated by the ones of $(X + E)^T (X + E)$. So with probability $1 - O(1/n^2)$, the eigenvalues of $\tilde{\Lambda}_r^2$ satisfy the following with some updated $C_1(K_{\psi_2})$ using [\(8\)](#page-8-2),

$$
\tilde{\lambda}_k \le \begin{cases}\n\hat{\lambda}_1^2 \le C_1 n & k \le r; \\
\hat{\lambda}_{r+1}^2 \le C_1 (\sqrt{n} + \sqrt{d})^2 \sigma^2, & k > r.\n\end{cases}
$$
\n(12)

Combine [\(11\)](#page-8-1), [\(12\)](#page-8-3) and $||y_k|| \leq 1$, we can update $C_1(K_{\psi_2})$ so that

$$
\|\xi_i^T \tilde{V}_r \tilde{\Lambda}_r^2 y_k\| \le \|\xi_i^T \tilde{V}_r\| \|S_i\| \le C_1 \sigma n \sqrt{\log n}.\tag{13}
$$

Now consider the term $\xi_i^T S_i(\hat{v}_k k - \tilde{V}_r y_k)$. Since S_i and ξ_i are independent, using Hanson–Wright inequality (Theorem 1.1 in [Rudelson and Vershynin \[2013\]](#page-22-7)), there is an updated $C_1(K_{\psi_2})$ so that with probability $1 - O(1/n^2)$,

$$
\begin{array}{rcl}\n\|\xi_i^T S_i\|^2 & \leq & \sigma^2 \text{tr}(S_i^2) + C_1 \sigma^2 (\|S_i^2\|_F + \|S_i^2\|) \log n \\
& \leq & C_1 \sigma^2 (\text{tr}(S_i^2) + \|S_i^2\|_F) \log n.\n\end{array}
$$

The term $||S_i^2|| = \lambda_1(S_i^2) \leq \text{tr}(S_i^2)$, so it is ignored. Now recall that $\text{tr}(S_i^2) = \sum_{i=1}^d \tilde{\lambda}_i^2$ and $||S_i^2||_F = (\text{tr}(S_i^4))^{1/2} = (\sum_{i=1}^d \tilde{\lambda}_i^4)^{1/2}$. Combining it with the results about $\tilde{\lambda}_i$ in [\(12\)](#page-8-3), we can update C_2 so that

$$
\begin{split} \|\xi_i^T S_i\| &\leq C_1 \sigma \sqrt{\sum_{k=1}^d \tilde{\lambda}_k^2 + \sqrt{\sum_{k=1}^d \tilde{\lambda}_k^4} \sqrt{\log n}} \\ &\leq 2C_1 \sigma \sqrt{\sum_{k=1}^d \tilde{\lambda}_k^2} \sqrt{\log n} \\ &\leq \frac{1}{2} C_2 \sigma \sqrt{n^2 + (\sqrt{n} + \sqrt{d})^4 \sigma^4 n} \sqrt{\log n} \\ &\leq C_2 \sigma \sqrt{\log n} (n + \sigma^2 n \sqrt{n} + \sigma^2 d \sqrt{n}). \end{split} \tag{14}
$$

Combining it with [\(10\)](#page-8-4) from Proposition [1,](#page-10-0) we have an updated $C_2(K_{\psi_2}, r)$, so that with probability $1 - O(1/n^2)$,

$$
\|\xi_i^T S_i(\hat{v}_k - \tilde{V}_r y_k)\| \le \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n})), \quad \Gamma := \frac{C\sigma}{\lambda_r^2} \sqrt{\log n} (n + \sigma^2 n \sqrt{n} + \sigma^2 d \sqrt{n}) (1 + \sigma \sqrt{\log n}).
$$

Consider the other three terms in (9) . Using the definition of event A , with probability $1 - O(1/n^2)$ and updated $C_1(K_{\psi_2})$, we have

$$
\begin{aligned} &\|\xi_i^T X_i X_i^T\| \le \|X_i^T \xi_i\| \le \sigma C_1 \sqrt{\log n}, \\ &\|\xi_i^T X_i \xi_i^T\| \le \xi_i^T \xi_i \|X_i\| \le \xi_i^T \xi_i \le C_1 \sigma^2 (\sqrt{d} + \sqrt{\log n})^2, \\ &\|\xi_i^T \xi_i X_i^T\| \le \xi_i^T \xi_i \|X_i\| \le \xi_i^T \xi_i \le C_1 \sigma^2 (\sqrt{d} + \sqrt{\log n})^2. \end{aligned}
$$

Combine all the four terms together in [\(9\)](#page-8-0), where $\xi_i^T S_i \hat{v}_k$ has been split into [\(13\)](#page-8-5) and $\Gamma(1+\sigma)$ √ Combine all the four terms together in (9), where $\xi_i^T S_i \hat{v}_k$ has been split into (13) and $\Gamma(1+\sigma(\sqrt{d}+\sqrt{d}))$ $\sqrt{\log n}$). Now we have

$$
(\hat{\lambda}_k^2 - ||\xi_i||^2)|\xi_i^T\hat{v}_k| \le \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n})) + C_1 d\sigma^2 + C_1 n\sigma\sqrt{\log n}.
$$

Also recall that when A takes place, $\hat{\lambda}_k^2 - ||\xi_i||^2 \geq \lambda_r^2/4$. Therefore, we can update $C_1(K_{\psi_2})$ so that √

$$
|\xi_i^T \hat{v}_k| \le \frac{4}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n})) + \frac{C_1 d\sigma^2 + nC_1 \sigma \sqrt{\log n}}{\lambda_r^2}.
$$
 (15)

Conclusion. The first error bound can be reached by removing the lower-order terms. Recall **Concrusion.** The first error bout
that $\sqrt{n} \ge \lambda_r \ge 1 + C_1 \sigma(\sqrt{n} + \sqrt{d}),$

$$
\frac{d\sigma^2}{\lambda_r^2} \le \frac{\sigma\sqrt{nd}}{C_1\lambda_r^2}, \quad \frac{d\sigma^2}{\lambda_r^2} \le \frac{\sigma\sqrt{d}}{C_1\lambda_r}, \quad \frac{\sigma}{\lambda_r^2}\sqrt{n\log n} \le \frac{\sigma}{\lambda_r^2}n\sqrt{\log n}, \quad \frac{\sigma}{\lambda_r^2}\sigma n \le \frac{\sigma}{\lambda_r^2}n\sqrt{\log n}.
$$

Next, if we fix $\lambda_r \geq c_X \sqrt{n}$ and $d \gtrsim n$, then our previous upper bound can be simplified because

$$
\frac{\sigma(\sqrt{n} + \sqrt{d})}{\lambda_r} + \frac{\sigma n \sqrt{\log n}}{\lambda_r^2} + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n}))
$$

$$
\lesssim \sqrt{\frac{d}{n}} \sigma(1 + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^4 d}{n} \log n) + \sqrt{\log n} \sigma.
$$

Similarly, if we fix $\lambda_r \geq c_X \sqrt{n}$ and $d \leq n$, then our previous upper bound can be simplified because

$$
\frac{\sigma}{\lambda_r^2}(\sqrt{nd} + n\sqrt{\log n}) + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n})) \lesssim \sqrt{\log n} \sigma (1 + \sigma^4 \sqrt{\frac{d \log n}{n}}).
$$

The detailed derivation of these simplifications can be found in Lemma 1 of [Tong et al. \[2024\]](#page-22-8). These conclude the second claim. For the third claim, simply note when $\sigma \leq 1$,

$$
\sqrt{\log n}\sigma (1 + \sigma^4 \sqrt{n^{-1}d\log n}) \le 2\log n\sigma.
$$

Finally, our claims work for every $i \in [n]$. Take the maximum over all $i \in [n]$, and with probability $1 - O(1/n)$, the bound holds. \Box

2.2 Leave-one-out eigenvector perturbation

In the proof of the main theorem, we use the leave-one-out method to deal with the most challenging term. This method considers leaving out one row or column of a random matrix and investigates how would the matrix change. It has been used to study the random matrix spectrum through Stieltjes transform [\[Tao, 2012\]](#page-22-9) and PCA eigenvector behavior [\[Chen et al., 2021,](#page-20-2) [O'Rourke et al., 2024\]](#page-22-10). Let \hat{v}_k be the k-th right singular vector of the data matrix without one specific sample and \tilde{V}_r be the right singular matrix of the original matrix. We show that \hat{v}_k and its projection in the column space of \tilde{V}_r differs by $O(\sigma\sqrt{d \log n}/\lambda_r^2)$, when [\(3\)](#page-5-2) holds. It delicately describes the contribution of one sample to the singular vectors. [Chen et al. \[2021\]](#page-20-2) studied the same problem and established a bound in (4.19) . The bound requires X and E to be symmetric, and $\|\hat{v}_k\|_{\infty} = O(1/\sqrt{d})$ to be effective. Our analysis does not rely on $\|\hat{v}_k\|_{\infty}$ or the symmetry, which is more general and allows arbitrary X . Such results can be useful in other applications. We rigorously present the result as the following proposition.

Proposition 1. Suppose $X \in \mathbb{R}^{n \times d}$ has rank r, where each row X_i^T is bounded by $||X_i|| \leq 1$. Suppose $E \in \mathbb{R}^{n \times d}$ has all entries being independent sub-Gaussian $(0, \sigma^2, K_{\psi_2})$ distributed. Let \tilde{Z} be the last n – 1 rows of $Z = X + E$. Denote the SVD of $X = U\Lambda V^T$, $Z = \hat{U}\hat{\Lambda}\hat{V}^T$ and $\tilde{Z} = \tilde{U}\tilde{\Lambda}\tilde{V}^T$. Let \hat{V}_r and \tilde{V}_r denote the first r columns of \hat{V} and \tilde{V} , respectively. Let λ_r be the smallest nonzero singular value of X. Then there are constants $n_0, C_1 = C_1(K_{\psi_2})$ and $C_2 = C_2(r, K_{\psi_2})$ so that if

$$
n > n_0, \quad \lambda_r > C_1 + C_1 \sigma(\sqrt{n} + \sqrt{d}),
$$

the following holds with probability at least $1 - O(1/n^2)$,

$$
\max_{k \le r} \|(I - \tilde{V}_r \tilde{V}_r^T)\hat{v}_k\| \le \begin{cases} C_1 \lambda_r^{-2}, & (\sqrt{d} + \sqrt{\log n})\sigma \le 2; \\ C_2 \lambda_r^{-2} \sigma (1 + \sigma \sqrt{\log n})(\sqrt{d} + \sqrt{\log n}), & (\sqrt{d} + \sqrt{\log n})\sigma > 2, \end{cases}
$$

and the following holds for both cases:

$$
\max_{k\leq r} \|(I-\tilde{V}_r\tilde{V}_r^T)\hat{v}_k\|\leq C_2\lambda_r^{-2}(1+\sigma\sqrt{\log n})(1+\sigma(\sqrt{d}+\sqrt{\log n})).
$$

To find out the difference between the original eigenvector and its projection in eigen-space formed by the leading r right singular vectors of the leave-one-out sample, we take advantage of the randomness of the perturbation. Roughly speaking, we project the eigenvector equation for \hat{v}_k to the eigen-space of interest and its orthogonal complement, where the latter is the difference of interest. By controlling the random effects of one sample on the eigen-space, we show that the difference has a small norm.

Proof. We discuss the problem for both the small noise case and the large noise case. We fix $k \leq r$ in the discussion. The claim for all k holds by a union bound.

Case 1: the small σ . Since the rows of E are independent sub-Gaussian distributed, by Theorem 5.39 in [Vershynin \[2010\]](#page-22-6), there is a constant $C_1 = C_1(K_{\psi_2})$, so that with probability at least $1 - O(1/n^3)$,

$$
\|\xi_1\| \le \sigma(\sqrt{d} + \frac{1}{6}C_1(1+\sqrt{\log n})) \le \frac{1}{3}C_1\sigma(\sqrt{d} + \sqrt{\log n}),
$$

$$
\|E\| \le \sigma(\sqrt{d} + \frac{1}{6}C_1(\sqrt{n} + \sqrt{\log n})) \le \frac{1}{3}C_1\sigma(\sqrt{d} + \sqrt{n}).
$$

We will define the event that both these hold as A . Our discussion assumes A holds. Denote

$$
\Delta = (X + E)^{T} (X + E) - (\tilde{X} + \tilde{E})^{T} (\tilde{X} + \tilde{E}) = (X_1 + \xi_1)(X_1 + \xi_1)^{T}.
$$

Since $||X_1|| \leq 1$,

$$
\|\Delta\|_F = \|X_1 + \xi_1\|^2 \le 2 + C_1^2 \sigma^2 (d + \log n).
$$

So by Weyl's inequality, the singular values follow that, for $k \leq r$,

$$
\hat{\lambda}_k \ge \lambda_r - \frac{1}{3}C_1\sigma(\sqrt{n} + \sqrt{d}), \quad \hat{\lambda}_{r+1} \le \frac{1}{3}C_1\sigma(\sqrt{n} + \sqrt{d}).\tag{16}
$$

Hence, by the Davis–Kahan Theorem [\[Davis and Kahan, 1970\]](#page-20-5), there is an orthogonal matrix O so that

$$
\|\hat{V}_r - \tilde{V}_r O\|_F \le \frac{4\|\Delta\|_F}{\hat{\lambda}_r^2 - \hat{\lambda}_{r+1}^2} \le \frac{8 + 4C_1^2 \sigma^2(d + \log n)}{\lambda_r(\lambda_r - \frac{2}{3}C_1 \sigma(\sqrt{n} + \sqrt{d}))} \le \frac{24(1 + C_1^2 \sigma^2(d + \log n))}{\lambda_r^2}.
$$

Let y_k be the k-th column of O, then $\hat{v}_k - \tilde{V}_r y_k$ is the k-th column of $\hat{V}_r - \tilde{V}_r O$. For any column, the ℓ_2 norm is bounded by the matrix Frobenius norm. Therefore, the result follows by

$$
\|(I-\tilde{V}_r\tilde{V}_r^T)\hat{v}_k\| = \min_{w_k \in \mathbb{R}^r} \|\hat{v}_k - \tilde{V}_rw_k\| \le \|\hat{v}_k - \tilde{V}_ry_k\| \le \|\hat{V}_r - \tilde{V}_rO\|_F, \quad \sigma(\sqrt{d} + \sqrt{\log n}) \le 2,
$$

and updating the value of C_1 .

Case 2: the large σ . This is a challenging case. We first introduce the notations. Recall the observed data is $Z = X + E$. We compare it with the case that sample 1 is missing, so we define $\tilde{Z} = \tilde{X} + \tilde{E}$ and the first sample is $Z_1 = X_1 + \xi_1$. The eigen-decomposition is $\tilde{Z}^T \tilde{Z} = \tilde{V} \tilde{\Lambda}^2 \tilde{V}^T$, where $\tilde{V} = [\tilde{V}_r, \tilde{V}_\perp].$

We define y_k and $y_{k,\perp}$ as follows:

$$
y_k = \tilde{V}_r^T \hat{v}_k, \quad y_{k,\perp} = \tilde{V}_\perp^T \hat{v}_k.
$$

Since $||(I - \tilde{V}_r \tilde{V}_r^T) \hat{v}_k|| = ||\tilde{V}_\perp \tilde{V}_\perp^T \hat{v}_k|| \le ||y_{k,\perp}||$, we are interested in bounding $||y_{k,\perp}||$. To do it, note that \hat{v}_k is the eigenvector of $Z^T Z$ corresponding to the eigenvalue $\hat{\lambda}_k$, so $Z^T Z \hat{v}_k = \hat{\lambda}_k^2 \hat{v}_k$. Decompose Z into \tilde{Z} and Z_1 , then it becomes

$$
(\tilde{Z}^T \tilde{Z} + Z_1 Z_1^T)(\tilde{V}_r y_k + \tilde{V}_\perp y_{k,\perp}) = \hat{\lambda}_k^2 (\tilde{V}_r y_k + \tilde{V}_\perp y_{k,\perp}).
$$

Multiplying by $\tilde{V}^T = [\tilde{V}_r, \tilde{V}_\perp]^T$ on both sides leads to:

$$
\begin{bmatrix}\n\tilde{\Lambda}_r^2 + \tilde{V}_r^T Z_1 Z_1^T \tilde{V}_r & \tilde{V}_r^T Z_1 Z_1^T \tilde{V}_\perp \\
\tilde{V}_\perp^T Z_1 Z_1^T \tilde{V}_r & \tilde{\Lambda}_\perp^2 + \tilde{V}_\perp^T Z_1 Z_1^T \tilde{V}_\perp\n\end{bmatrix}\n\begin{bmatrix}\ny_k \\
y_{k,\perp}\n\end{bmatrix} = \n\begin{bmatrix}\n\hat{\lambda}_k^2 y_k \\
\hat{\lambda}_k^2 y_{k,\perp}\n\end{bmatrix},
$$
\n(17)

where $\tilde{\Lambda}_r$ is the diagonal matrix consisting of the first r singular values of \tilde{Z} , and $\tilde{\Lambda}_\perp$ is the diagonal matrix consisting of the remaining singular values of Z .

Denote $a = \tilde{V}_r^T Z_1$ and $b = \tilde{V}_\perp^T Z_1$. From the second row of [\(17\)](#page-11-0), we have

$$
y_{k,\perp} = (\hat{\lambda}_k^2 I - (\tilde{\Lambda}_\perp^2 + bb^T))^{-1} ba^T y_k.
$$

So $||y_{k,\perp}||$ can be bounded by analyzing the terms $a, b, \hat{\lambda}_k$ and $\tilde{\Lambda}^2_{\perp}$.

Note that

$$
\mathbb{E}_{\xi_1}[\|\tilde{V}_r^T \xi_1\|^2] = \sigma^2 \text{tr}(\tilde{V}_r^T \tilde{V}_r) = \sigma^2 \|\tilde{V}_r\|_F^2 = \sigma^2 r, \quad \|\tilde{V}_r\| = 1.
$$

By the independence between \tilde{V} and $Z_1 = X_1 + \xi_1$, using Hanson–Wright inequality (Theorem 1.1 in [Rudelson and Vershynin \[2013\]](#page-22-7)), we find that

$$
P(||\tilde{V}_r^T \xi_1\|^2 - \sigma^2 r| > \sigma^2 t) \leq 2 \exp\left(-c \min(\frac{t^2}{K_{\psi_2}^4 r}, \frac{t}{K_{\psi_2}^2})\right).
$$

We pick $t = 3K_{\psi_2}^2 \log n/c$, then the probability is $O(1/n^3)$ for sufficiently large n. Likewise,

$$
P(||\tilde{V}_{\perp}^T \xi_1\|^2 - \sigma^2(d-r)| > \sigma^2 t) \leq 2 \exp\left(-c \min(\frac{t^2}{K_{\psi_2}^4(d-r)}, \frac{t}{K_{\psi_2}^2})\right).
$$

If we pick $t = 3K_{\psi_2}^2$ $\sqrt{d \log n}/\sqrt{c}$, the probability above will be $O(1/n)$. In conclusion, there is a constant $C_1 = C_1(K_{\psi_2}), C_2 = C_2(K_{\psi_2}, r)$, so that the following holds with probability at least $1 - O(1/n^3),$

$$
||a|| \le ||X_1|| + ||\tilde{V}_r^T \xi_1|| \le 1 + \frac{1}{2}C_1\sigma(\sqrt{r} + \sqrt{\log n}) \le C_2(1 + \sigma\sqrt{\log n}),
$$

$$
||b|| \le ||X_1|| + ||\tilde{V}_\perp^T \xi_1|| \le \frac{1}{6} + \frac{1}{6}C_1\sigma(\sqrt{d} + \sqrt{\log n}) \le \frac{1}{6}\lambda_r.
$$

Assuming A holds, we can combine these estimates with [\(16\)](#page-11-1) on $\hat{\lambda}_k$ and $\hat{\lambda}_{r+1}$, so that

$$
\|(\hat{\lambda}_k^2 I - \tilde{\Lambda}_\bot^2 + bb^T)^{-1}b\| \leq \frac{\|b\|}{\hat{\lambda}_r^2 - \hat{\lambda}_{r+1}^2 - \|b\|^2} \leq \frac{1 + C_1\sigma(\sqrt{d} + \sqrt{\log n})}{2\lambda_r^2 - \lambda_r^2}.
$$

Therefore, we have

$$
\|y_{k,\perp}\|\quad \leq \quad \|(\hat{\lambda}_k^2 I - \tilde{\Lambda}_\perp^2 + bb^T)^{-1}b\|\|a\|\|y_k\| \leq \frac{C_2}{\lambda_r^2}(1 + \sigma\sqrt{\log n})(1 + C_1\sigma(\sqrt{d} + \sqrt{\log n})).
$$

When σ √ $d \geq 2$, the latter factor can be further simplified. The final claim can be achieved with an updated value of C_2 .

 \Box

2.3 Condition on the minimum eigenvalue and zigzag lines

The main theorem in Section [2.1](#page-5-0) provides an upper bound on $||X - \hat{X}||_{2,\infty}$ when the smallest The main theorem in section 2.1 provides an upper bound on $\|\Lambda - \Lambda\|_{2,\infty}$ when the smallest nonzero singular value of the clean data matrix X is at the order of \sqrt{n} . Hence, it becomes pertinent to ask under what conditions this criterion can be satisfied.

We introduce a theorem below concerning low-rank random matrices. Let X be a random matrix with $cov(X_i) = \Sigma$, where Σ is a low-rank matrix. The smallest nonzero singular value of matrix with $cov(x_i) = \Sigma$, where Σ is a low-rank matrix. The smallest honzero singular value of X can be lower bounded by the smallest eigenvalue of Σ and \sqrt{n} . This result can be viewed as a straightforward extension of the standard random matrix theory as [\[Vershynin, 2010,](#page-22-6) Theorem 5.39].

Theorem 2. Suppose $X_i \in \mathbb{R}^d$ are i.i.d. samples from a distribution so that $||X_i|| \leq 1$ almost surely, with $\mathbb{E}[X_i] = 0$ and $cov(X_i) = \Sigma$ where $rank(\Sigma) = r$. Let $\lambda_i(\Sigma)$ be the *i*-th largest eigenvalue of Σ . If $n > r$, the following holds with probability at least $1 - e^{-ct^2}$,

$$
\sqrt{\lambda_r(\Sigma)}(\sqrt{n}-C\sqrt{r}-t) \leq \lambda_r(X) \leq \lambda_1(X) \leq \sqrt{n}.
$$

Here C and c are some constants that depend on $\lambda_r(\Sigma)$, but not on n and d.

Proof. Since $||X_i|| \leq 1$ almost surely, the Frobenius norm of X is bounded, and hence the spectral norm $||X|| = \lambda_1(X)$ can be bounded as follows:

$$
\lambda_1^2(X) \le ||X||_F^2 = \sum_{i=1}^n ||X_i||^2 \le n.
$$

Consider the lower bound of $\lambda_r(X)$. Denote the eigen-decomposition of Σ be $\Sigma = U_{\Sigma} \Lambda_{\Sigma} U_{\Sigma}^T$ where $U_{\Sigma} \in \mathbb{R}^{d \times r}$ and $\Lambda_{\Sigma} \in \mathbb{R}^{r \times r}$. We set $Y_i = \Lambda_{\Sigma}^{-1/2} U_{\Sigma}^T X_i \in \mathbb{R}^r$, then $||Y_i|| \leq [\lambda_r(\Sigma)]^{-1/2}$. Further, $\mathbb{E}[Y_i] = 0$, $cov(Y_i) = I_r$, and Y_i is sub-Gaussian distributed, where the constant K_{ψ_2} follows

$$
K_{\psi_2}(Y_i) = \sup_{\|y\|=1} \sup_{p\geq 1} \frac{\mathbb{E}[|\langle y, Y_i\rangle|^p]^{1/p}}{p^{1/2}} \leq \sup_{\|y\|=1} \sup_{p\geq 1} \frac{[\lambda_r(\Sigma)]^{-1/2}}{p^{1/2}} = [\lambda_r(\Sigma)]^{-1/2}.
$$

We can apply [\[Vershynin, 2010,](#page-22-6) Theorem 5.39] to Y and find constant C and c so that with probability $1 - 2 \exp(-ct^2)$, √ √

$$
\sqrt{n} - C\sqrt{r} - t \leq \lambda_r(Y).
$$

Finally, according to the definition of Y, the original matrix X of interest is $X = Y \Lambda_{\Sigma}^{1/2} U_{\Sigma}^{T}$. Therefore, the smallest singular value of X can be bounded by

$$
\lambda_r(X) \geq \lambda_r(Y) \sqrt{\lambda_r(\Sigma)} \geq \sqrt{\lambda_r(\Sigma)} (\sqrt{n} - C\sqrt{r} - t).
$$

The result is proved.

By Theorem [2,](#page-12-1) it suffices to compute the covariance of the clean data X_i and show its r-th eigenvalue is bounded from below. This can be done easily if X_i follows some known distributions such as Gaussian mixtures. Below, we discuss a scenario where the covariance computation is not so elementary.

For many temporal and spatial data sets [\[Cressie and Wikle, 2015,](#page-20-12) [Brunton and Kutz, 2022,](#page-20-13) [Khoo et al., 2024\]](#page-21-11), researchers expect the clean data to be generated from a one-dimensional curve, i.e. the function of time or location t. It is of interest to estimate the function x_t , where a popular approach is to use a piecewise linear approximation [\[Pavlidis and Horowitz, 1974,](#page-22-11) [Stone, 1961\]](#page-22-12). For the simplicity of exposition, we assume x_t is piecewise linear. In less mathematical terms, x_t is a zigzag line. Figure [1](#page-18-0) below shows two such objects in 3-dimensional space. For each zigzag line, there will be R time points $0 = t_1 < t_2 < t_3 < \ldots < t_R < 1 = t_{R+1}$, and

$$
x_s = x_{t_j} + (s - t_j)v_j, \quad s \in [t_j, t_{j+1}].
$$

Here $\{v_j, j = 1, \ldots, R\}$ is a group of d-dimensional vectors with unit-norm. We want to make a special note here that the number of segments R does not indicate the dimension of the sub-space dimension. For example, Figure [1](#page-18-0) shows two zigzag lines in a three-dimensional subspace $(r = 3)$ with any $R = 10$ segments.

Proposition 2. Suppose $X_i = x_{s_i}$ and $s_i \sim Unif[0,1]$ independently, $i \in [n]$. Suppose the linear subspace spanned by $\{v_j, j = 1, \ldots, R\}$ has dimension r and $\mathbb{E}[X_i] = 0$. If there is a constant $\rho > 0$ so that the segment endpoints $t_{i+1} - t_i \ge \rho$, then there exists a constant $c > 0$ independent of n and d, so that the following holds with probability $1 - O(1/n)$,

$$
c\sqrt{\rho^3/3 - \rho^4/4}\sqrt{n} \le \lambda_r(X) \le \lambda_1(X) \le \sqrt{n}.
$$

Proposition [2](#page-12-1) utilizes Theorem 2 to establish that the minimum singular value $\lambda_r(X) > c_X \sqrt{n}$ holds for the zigzag example, given that each segment has a minimum length $\rho > 0$ and no segment is degenerate.

In the case where there are degenerate segments, denoted by $\rho = o(1)$, we can still provide an upper bound on the error. Notably, Theorem [1](#page-5-1) demonstrates that the error of Algorithm [1](#page-1-0) can be controlled by $n\sigma/\lambda_r^2(X)$ for any arbitrary $\lambda_r(X)$. By incorporating Proposition [2](#page-13-0) into this result, we observe that the error tends to 0 as $\sigma \ll \rho^3$, even when $\rho = o(1)$.

 \Box

3 General lower bound for matrix denoising

In this section, we will show PCA-denoising is rate-optimal in terms of sample complexity and noise intensity. Theorem 3 of [Cai and Zhang \[2018\]](#page-20-7) has shown the optimality of PCA referring to the recovery of the low-dimensional subspace V_r . In its Remark 4, [Cai and Zhang \[2018\]](#page-20-7) claims that the result can be generalized to lower bounds for matrix denoising. Theorem 3 of [Montanari](#page-21-9) [et al. \[2018\]](#page-21-9) has studied the lower bound for matrix denoising under the assumption that $n \approx d$. Our result below is more general than these two works in some perspectives: we allow n and d to diverge at different speeds (in comparison with [Montanari et al. \[2018\]](#page-21-9); meanwhile their results don't involve hidden constants), and the estimator \hat{X} can be arbitrary [\(Cai and Zhang \[2018\]](#page-20-7) assumes \overline{X} is of rank at least r).

We consider a simplified zigzag model. Suppose we only have one segment v and the clean data is $X_i = (t_i + 1)v$ for $t_i \sim Unif(0, 1)$ independently. The observed data is $Z_i = X_i + \xi_i$ with noise ξ_i . A successful matrix denoising algorithm will generate $\hat{X} = (\hat{t} + 1)v^T$, which gives the direction v and the points of observation t_i , $i \in [n]$. We want to set up the general lower bound $||\overline{X} - X||_{2,\infty}$ for any estimator \overline{X} .

Theorem 3. Let $v \sim \mathcal{N}(0, d^{-1}I_d)$. Let $t_i \sim Unif(0, 1)$ independently, $i \in [n]$. Let $X_i = (t_i + 1)v$. Suppose $Z_i = X_i + \xi_i$ is observed, where $\xi_i \sim \mathcal{N}(0, \sigma^2 I_d)$, $i \in [n]$. Suppose for some $\epsilon \in (0, d/4)$, the noise level $\sigma > 4\epsilon/\sqrt{\Phi(-1)}$ or the sample size $n < d\sigma^2/(5\epsilon^2)$, then for any estimator \hat{X} , there is

$$
\mathbb{E}_v \mathbb{E}_{t_{[n]}, \xi_{[n]}} [\|\hat{X}_i - X_i\|_{2,\infty}^2] \ge \epsilon^2,
$$

where $t_{[n]} := (t_1, \ldots, t_n), \xi_{[n]} := (\xi_1, \ldots, \xi_n),$ and Φ is the cumulative distribution function of standard Gaussian.

Theorem [3](#page-14-3) establishes lower bounds for two crucial factors: the signal-to-noise ratio level and the sample size. To achieve a small error rate ϵ^2 , it is necessary to have both a low noise level $\sigma = O(\epsilon)$ and a large sample size $n = O(d\sigma^2/\epsilon^2)$. This explains why a spectral gap requirement in the form [\(3\)](#page-5-2) is necessary: Using Proposition [2,](#page-13-0) we have $\lambda_r(X) \ge c_X \sqrt{n}$ for some constant c_X , so the form [\(3\)](#page-5-2) is necessary: Using F10position 2, we have $\lambda_r(\Lambda) \leq c_X \sqrt{n}$ for some constant c_X , so (3) essentially requires $\sigma \sqrt{d} \lesssim \sqrt{n}$, while Theorem [3](#page-14-3) indicates the complementary case will lead to inaccurate denoising result. When $\sigma = C\epsilon$, the required sample size is $n \geq C d$, aligning with the assumption made in the last estimate in Theorem [1.](#page-5-1)

Proof of Theorem [3](#page-14-3) is presented in Section 3 of [Tong et al. \[2024\]](#page-22-8).

4 Applications of uniform denoising

In this section, we demonstrate the utility of having access to uniformly denoised data points in various downstream statistical learning applications. The main theme is to show that the learning results using uniformly denoised data are close to those achieved by clean data. In this section, we consider the clustering problem and manifold learning using the graphical Laplacian.

We also have results for empirical risk minimization (ERM) in the Supplementary Material[\[Tong et al., 2024\]](#page-22-8). ERM is a widely used approach for training statistical models [\[Hastie](#page-21-5) [et al., 2009,](#page-21-5) [Mei et al., 2018\]](#page-21-12). In ERM, the dataset consists of pairs (X_i, y_i) , where $X_i \in \mathbb{R}^d$ and y_i is typically a scalar. The prediction error is measured by a loss function $F(X_i, y_i, \theta)$, where the specific definition of F varies depending on the model. Under mild conditions, we show that the loss function based on uniformly denoised data \hat{X}_i is a close approximation of that based on the clean data X_i . Since the derivation is rather routine, we leave the details to the Section 4.1 of [Tong et al. \[2024\]](#page-22-8).

4.1 Clustering

Consider a clustering problem with clean data points $X_{[n]} \in \mathbb{R}^d$ in K clusters [\[Omran et al., 2007\]](#page-21-13). Denote \mathcal{C}_k as the index set of data points in the k-th cluster and the division is $\mathcal{C} = \{C_1, \dots, C_K\}$. K-means in [Hartigan and Wong \[1979\]](#page-21-14) is a very popular clustering algorithm. It aims to find the labels $\hat{\mathcal{C}}$ and centers $\hat{m} = \hat{m}_{[K]}$, so that the following is minimized

$$
f(\mathcal{C}, m) = \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} ||X_i - m_k||^2.
$$
 (18)

When the clean data X is available, k -means will achieve a good clustering result.

In practice, we only have access to noisy data $Z_i = X_i + \xi_i$. If we simply replace X_i in [\(18\)](#page-15-2) with Z_i , it is unlikely that we can get a good estimation. Using denoised data matrix \hat{X} , we would consider minimizing the loss function

$$
\hat{f}(\mathcal{C}, m) = \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} ||\hat{X}_i - m_k||^2,
$$
\n(19)

where \hat{X}_i comes from Algorithm [1.](#page-1-0) We expect the clustering result from \hat{f} to have a performance similar to that of using X directly. This is similar to existing consistency analysis of clustering in [Pollard \[1981\]](#page-22-13), [Rakhlin and Caponnetto \[2006\]](#page-22-14), and [Bubeck and Luxburg \[2009\]](#page-20-14), where the loss function convergence is of main interest.

To evaluate the clustering performance, the weak consistency focuses on the clustering error rate and the strong consistency focuses on the number of clustering errors. With the uniform error bound between \hat{X}_i and X_i , we can control the clustering error on each data point and conclude a strong consistency result in Proposition [3.](#page-15-0) Such results cannot be achieved by the error control on the Frobenius norm.

On the other hand, without further assumptions, it can be hard to show label consistency. This is mainly because k-means clustering result is non-unique and can have discontinuous dependence at data points near the boundary between two clusters. We remove such cases by further assuming there is a center $m_k \in \mathbb{R}^d$ for the k-th cluster, so that data points in \mathcal{C}_k are close to m_k . Such a requirement is standard in k-means related literature [\(Hartigan and Wong \[1979\]](#page-21-14), [Jin and Wang](#page-21-15) [\[2016\]](#page-21-15), [Jin et al. \[2017\]](#page-21-16)). A rigorous assumption is as follows.

Assumption 1. Suppose that under the division \mathcal{C} , the clusters satisfy the following with absolute constants c_0, δ, c_m :

- The cluster size does not degenerate: $|\mathcal{C}_k| \ge c_0 n$ for all $k \in [K]$.
- Data points in each cluster are close to a center: $||X_i m_k|| \leq \delta, \forall i \in \mathcal{C}_k$.
- The cluster centers are far apart: $||m_j m_k|| \geq c_m$, for all $1 \leq j \neq k \leq K$.

Proposition 3. Consider a clustering problem with noisy data points $Z_{[n]}$ and underlying truth C. Suppose $\|\hat{X} - X\|_{2,\infty} \leq \epsilon$. Let $\hat{\mathcal{C}}$ be the minimizer of the loss function $\hat{f}(\mathcal{C},m)$ defined in [\(19\)](#page-15-3). Then

• $\hat{\mathcal{C}}$ almost minimize the loss function with clean data $f(\mathcal{C}, m)$,

$$
f(\hat{\mathcal{C}}, \hat{m}) \le f(\mathcal{C}, m) + 4\epsilon.
$$

• Furthermore, suppose Assumption [1](#page-15-4) holds with $c_m > 2(\delta + \epsilon + 2\sqrt{2(\delta^2 + \epsilon^2)/c_0})$, then kmeans gives a perfect clustering result, i.e. the division $\hat{\mathcal{C}}$ is exactly the same with \mathcal{C} .

The proof can be found in the Supplementary Material[\[Tong et al., 2024\]](#page-22-8).

4.2 Manifold learning with graphical Laplacian

Graphical Laplacian is an important data processing tool in various applications. It is commonly used in spectral clustering, dimensionality reduction, and graph-based machine learning tasks that involve analyzing data on nonlinear manifolds. It enables users to capture the underlying graphical structures [\[Von Luxburg, 2007,](#page-22-15) [Singer, 2006,](#page-22-16) [Khoo et al., 2024\]](#page-21-11). To illustrate, consider a kernel function $k(\cdot, \cdot)$ as follows:

Assumption 2. The function $k(x, y)$ is symmetric and Lipschitz in both x and y. Furthermore, there is an absolute constant $K > 0$ so that $1/K \leq k(x, y) \leq K$ when $||x||, ||y|| \leq 1$.

A similar version of this assumption can be found in [Von Luxburg et al. \[2008\]](#page-22-0). A quick example that satisfies Assumption [2](#page-16-0) is the most commonly used Gaussian kernel, where $k(x, y)$ = $\exp(-\|x-y\|^2/(2b^2)).$

Suppose the clean data X_i are i.i.d. samples from a distribution p on the unit ball $\{x : ||x|| \leq 1\}$. The population normalized graphical Laplacian operator is defined as

$$
\mathcal{L}f(x) = \int (1 - \frac{k(x, y)}{\sqrt{d(x)d(y)}}) f(y)p(y)dy, \text{ where } d(x) = \int k(x, y)p(y)dy.
$$

 $\mathcal L$ can be used for different purposes. It can directly be used to optimize a certain utility function. Its eigenfunctions can be used to extract a low-dimensional representation of the manifold or to establish spectral clustering results.

In practice, we do not have access to $\mathcal L$ directly. We first build the empirical operator based on the clean data X_i . Define the kernel matrix A where $A_{i,j} = k(X_i, X_j)$, and the diagonal matrix D with $D_{i,i} = \sum_{j=1}^{n} A_{i,j}$. The normalized Laplacian is given by

$$
L = I - D^{-1/2} A D^{-1/2}.
$$
\n(20)

Under mild conditions, [Von Luxburg et al. \[2008\]](#page-22-0) has shown L converge to $\mathcal L$ as an operator, and so are its spectrum and eigenvalues. In particular, with k as a Gaussian kernel, if λ_0 is a simple eigenvalue of $\mathcal L$ with its eigen-function $||u(x)||_{\infty} = 1$, then Theorems 16 and 19 of [Von Luxburg](#page-22-0) [et al. \[2008\]](#page-22-0) indicate that L has an eigenvalue λ close to λ_0 , and the corresponding eigenvector v (of which the norm may not be 1) satisfies that

$$
\max_{i} |v_i - u(X_i)| = O(n^{-1/2}).
$$

Hence, v will carry similar information as u and can be used for further analysis such as clustering. These results provide a theoretical foundation for utilizing the eigenvectors of L in statistical inference.

In the context of this paper, it is natural to ask whether such a result can be extended to the case only $Z_i = X_i + E_i$ are observed. Intuitively, we will first apply Algorithm [1](#page-1-0) to obtain the denoised data \hat{X}_i , and achieve \hat{L} correspondingly. Define \hat{A} where $\hat{A}_{i,j} = k(\hat{X}_i, \hat{X}_j)$, \hat{D} where $\hat{D}_{i,i} = \sum_{j=1}^{n} \hat{A}_{i,j}$, and

$$
\hat{L} = (I - \hat{D}^{-1/2} \hat{A} \hat{D}^{-1/2}).
$$
\n(21)

We are interested in how different \hat{L} would be from L, in the sense of matrix ℓ_{∞} norm, the informative eigenvalues, and the corresponding eigenvectors. The following proposition addresses the result:

Proposition 4. Consider a kernel function that Assumption [2](#page-16-0) holds. Suppose the denoised matrix follows $||\hat{X} - X||_{2,\infty} \leq \epsilon$. Then there is a constant $C > 0$ independent of n and d so that in the ℓ_{∞} operator-norm

$$
\|\hat{L} - L\|_{\infty} \le C\epsilon.
$$

Furthermore, suppose L has a simple eigenvalue $\lambda_0 \neq 1$. Then there are N_0 , $\epsilon > 0$, and a constant $C > 0$ independent of n and d, so that when $n > N_0$, $\epsilon < \epsilon_0$, L, L has eigen-pair (v, λ) and (\hat{v}, λ) respectively, so that

$$
|\hat{\lambda} - \lambda| \le C\epsilon, \quad ||a\hat{v} - v||_{\infty} \le C\epsilon ||v||_{\infty},
$$

where $a = 1$ or -1 .

5 Numerical simulation

To support our theoretical results, we provide some numerical examples in this section. We consider a two-class clustering setting where the clean data are sampled from two separate lowdimensional zigzag lines embedded in a high-dimensional space. In detail, we first generate two zigzag lines, each consisting of 10 segments within a unit ball in \mathbb{R}^3 , as seen in Figure [1\(](#page-18-0)a). Then we embedded them into d-dimensional space by an orthonormal matrix $\Omega \in \mathbb{R}^{d \times 3}$ with d being some large number to be specified later. Denote the embedded zigzag lines as $x_0(t)$ and $x_1(t)$ for $0 \leq t \leq 1$. The clean data are generated as $X_i = \Omega x_{\ell(i)}(t_i)$, where $\ell(i) \sim Bernoulli(1/2)$ and $t_i \sim U[0,1], i \in [n]$. Here $\ell(i)$ can be regarded as the label of this sample and our goal is to recover it. Finally, we add the noise to the clean data, where $Z_i = X_i + \xi_i$, $\xi_i \sim \mathcal{N}(0, \sigma^2 I_d)$. The algorithm will be applied to the noisy data Z.

We first check the uniform error of PCA-denoised data by Algorithm [1.](#page-1-0) The error in Theorem [1](#page-5-1) requires the information of $\lambda_r(X)$, which is shown to be $O(\sqrt{n})$ in Proposition [2](#page-13-0) for the zigzag scenario. According to the generation procedure, the clean data X lies in the space with a dimension of $r = 3$, so we calculate $\lambda_3(X)$ for n ranging from 10^2 to 70^2 . The embedding Ω does not change the eigenvalue, and $\lambda_3(X) = \lambda_3(x)$ where $x = (x_{\ell(1)}(t_1), \dots, x_{\ell(n)}(t_n))$. We calculate it based on x so the dimension d does not affect the result. As seen in Figure [1\(](#page-18-0)b), $\lambda_3(X)$ scales It based on x so the dimension a does not affect the result. As seen in rigure $I(\nu)$, $\lambda_3(\Lambda)$ scales linearly with \sqrt{n} , which follows Proposition [2.](#page-13-0) Such a result indicates the simplified uniform error in Theorem [1](#page-5-1) holds, that $||X - \hat{X}||_{2,\infty} = O(\sigma + \sigma^4 \log n)$ when $d = c_d n$. We fix $n = 0.1d$ and let d range from 10^2 to 10^4 , where the added noise $\sigma \in \{0.0025, 0.05, 0.1\}$. Figure [1\(](#page-18-0)c) presents the uniform error of the noisy data Z and the denoised data \hat{X} by Algorithm [1.](#page-1-0) It can be found that uniform error of the noisy data χ and the denoised data X by Algorithm 1. It can be found that $\|\hat{X} - X\|_{2,\infty}/\sigma$ stays constant throughout the regime while $\|Z - X\|_{2,\infty}/\sigma$ grows as \sqrt{d} . Even in the high-dimensional case that $d = 10n$, the PCA-denoising algorithm can reduce the curse of dimension and reduce the uniform error to be $O(\sigma)$.

We then investigate the denoising effects on downstream statistical inference, particularly in the graphical Laplacian eigenvectors and spectral clustering. Let $L(B)$ denote the graphical Laplacian of any data matrix B . We find the eigenvector corresponding to the second smallest eigenvalue, which is called the Fiedler eigenvector. Denote it as $\eta_2(B)$. It is well known that the component signs of $\eta_2(B)$ can be used to infer cluster labels [\(Von Luxburg et al. \[2008\]](#page-22-0)). We are interested in the error of the Fiedler eigenvectors from different data sources and the associated clustering results.

To illustrate the importance of the uniform error, we consider three data matrices in this study: the clean data X, the PCA-denoised data \hat{X} , and a new data set \tilde{X} with errors not uniformly distributed. To make the comparison fair, the new matrix X has the same average error as X , i.e.

$$
\frac{1}{n}\sum_{i=1}^{n}||X_i - \hat{X}_i||^2 = \frac{1}{n}\sum_{i=1}^{n}||X_i - \tilde{X}_i||^2.
$$
 (22)

We design \tilde{X} in a way that some samples have a large error and other samples have no errors. In detail, we randomly select 10% of indices as \mathcal{I} , and define \tilde{X}_i as:

$$
\tilde{X}_i = \begin{cases} X_i, & \text{if } i \in \mathcal{I}^c; \\ \alpha X_i + (1 - \alpha)Z_i, & \text{if } i \in \mathcal{I}, \end{cases}
$$

where $\alpha \in (0,1)$ is a constant that ensures [\(22\)](#page-17-1). In other words, 90% of data in \tilde{X} are clean; the others have *i.i.d.* errors as $(1 - \alpha)\xi_i$, where α is chosen so that the average error is the same with \hat{X} . Comparisons between $\eta_2(\hat{X})$ and $\eta_2(\tilde{X})$ explain how uniform error is important in practice.

We fix $d = 5000$ and $n = 1000$. The bandwidth for graphical Laplacian is chosen as $b = \sqrt{0.005}$. We first compare the Fiedler eigenvectors from three data matrices X, \hat{X} , and \hat{X} at two noise levels $\sigma = 0.0075$ and $\sigma = 0.05$, shown in Figure [2\(](#page-19-0)a). We adjust the order of data samples so that samples from $x_0(t)$ come first and those from $x_1(t)$ are all at last. In the top panel of Figure [2\(](#page-19-0)a), we can see the sign $\eta_2(X)$ meets the label ℓ . For the two data with small noise $\sigma = 0.0075$, $\eta_2(\hat{X})$ and $\eta_2(\hat{X})$ are close to $\eta_2(X)$ at a large scale. The error of $\eta_2(\hat{X})$ is uniformly small but

Figure 1: Properties of data generated from two zig-zag lines. (a) The two zigzag lines where the clean data is generated from. They are separated by parallel hyperplanes with a distance of 0.2. (b) The third singular value of the clean data $X \in \mathbb{R}^{n \times 3}$, which scales linearly with the square root of the sample size n. (c) The normalized uniform errors of the noisy data Z , denoised data \hat{X} achieved at different dimension d. With a fixed standard deviation of the noise σ , the uniform errors obtained from the noisy data $Z \in \mathbb{R}^{n \times d}$ increase with the dimension d and sample size $n = 0.1d$. However, the uniform errors obtained from the denoised data \ddot{X} remain approximately constant. The results are consistent with different standard deviations, $\sigma = 0.025, 0.05,$ and 0.1.

 $\eta_2(X)$ has spiked errors. This is consistent with Proposition [4.](#page-16-1) When σ increases from 0.0075 to 0.05, the spiked error in $\|\tilde{X}_i - X_i\|$ is so large that the distance-based clustering algorithm fails. Meanwhile, the Fiedler eigenvector obtained by \hat{X} can still preserve the correct cluster labels. We then study the clustering results of the three data matrices for σ ranging from 10^{-3} to 1, shown in Figure [2\(](#page-19-0)b). We use the adjusted rand index (ARI) to measure the clustering accuracy, where $ARI = 1$ indicates a perfect clustering and $ARI = 0.5$ means a random guess. The performance of clustering using other data sources would of course depend on the strength of noise. Not surprisingly, because of the crash in estimating $\eta_2(X)$, using X fails when $\sigma \geq 0.05$ and using X gives a satisfactory clustering result till $\sigma \geq 0.1$.

Acknowledgments

The authors would like to thank Yuehaw Khoo for the throughout discussion and reading of this project. The authors also would like to thank the two anonymous referees, an Associate Editor and the Editor for their comments that improved the quality of this paper.

Figure 2: Spectral clustering with different data sources. (a) The Fiedler eigenvectors of the Laplacian formed by the clean data, the denoised data, and the "averaged denoised" data at two noise levels. The first row plots the second (Fiedler) eigenvector of the graphical Laplacian obtained by the clean data X, the denoised data \hat{X} , and averaged denoised data \hat{X} . They are denoted by $\eta_2(X), \eta_2(X)$, and $\eta_2(X)$, respectively. The second row plots the pointwise absolute difference between $(\eta_2(X), \eta_2(\hat{X}))$ and $(\eta_2(X), \eta_2(\tilde{X}))$. The standard deviation of the noise is chosen to be $\sigma = 0.0075$ and 0.05, corresponding to the two columns. (b) The adjusted rand indices (ARI) by spectral clustering using clean data X, denoised data \hat{X} , and the averaged denoised data \hat{X} when σ changes.

Funding

Xin T. Tong's research is supported by Singapore MOE grant Tier-1-A-8000459-00-00. Wanjie Wang's research is supported by Singapore MOE grant Tier-1-A-8001451-00-00. Yuguan Wang's research is supported by the Department of Statistics, University of Chicago.

References

- Emmanuel Abbe and Colin Sandon. Community detection in general stochastic block models: Fundamental limits and efficient algorithms for recovery. In the proc. of IEEE FOCS, pages 670–688, 2015. doi: https://doi.org/10.1109/FOCS.2015.47.
- Emmanuel Abbe, Jianqing Fan, Kaizheng Wang, and Yiqiao Zhong. Entrywise eigenvector analysis of random matrices with low expected rank. Ann. Stat., 48(3):1452–1474, 2020. doi: https: //doi.org/10.1214/19-AOS1854.
- Hervé Abdi and Lynne J Williams. Principal component analysis. Wiley Interdiscip. Rev. Comput. Stat., 2(4):433–459, 2010. doi: https://doi.org/10.1002/wics.101.
- Zhigang Bao, Xiucai Ding, and Ke Wang. Singular vector and singular subspace distribution for the matrix denoising model. Ann. Stat., $49(1):370-392$, 2021. doi: https://doi.org/10.1214/ 20-AOS1960.
- Rajendra Bhatia. Matrix Analysis, volume 169. Springer Science & Business Media, 2013. doi: http://dx.doi.org/10.1007/978-1-4612-0653-8.
- Steven L Brunton and J Nathan Kutz. Data-driven Science and Engineering: Machine Learning, Dynamical Systems, and Control. Cambridge University Press, 2022. doi: https://doi.org/10. 1017/9781108380690.
- Sébastien Bubeck and Ulrike von Luxburg. Nearest neighbor clustering: A baseline method for consistent clustering with arbitrary objective functions. J. Mach. Learn. Res., 10:657–698, 2009. doi: http://jmlr.org/papers/v10/bubeck09a.html.
- T Tony Cai and Anru Zhang. Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. Ann. Stat., $46(1):60-89$, 2018. doi: https://doi.org/ 10.1214/17-AOS1541.
- Joshua Cape, Minh Tang, and Carey E Priebe. The two-to-infinity norm and singular subspace geometry with applications to high dimensional statistics. Ann. Stat., 47(5):2405–2439, 2019a. doi: https://doi.org/10.1214/18-AOS1752.
- Joshua Cape, Minh Tang, and Carey E Priebe. Signal-plus-noise matrix models: eigenvector deviations and fluctuations. Biometrika, 106(1):243–250, 2019b. doi: https://doi.org/10.1093/ biomet/asy070.
- Yuxin Chen, Yuejie Chi, Jianqing Fan, Cong Ma, et al. Spectral methods for data science: A statistical perspective. Found. Trends Mach. Learn., 14(5):566–806, 2021. doi: http://dx.doi. org/10.1561/2200000079.
- Noel Cressie and Christopher K Wikle. *Statistics for Spatio-temporal Data*. John Wiley & Sons, 2015. ISBN ISBN: 978-0-471-69274-4.
- Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. SIAM J. Numer. Anal., 7(1):1–46, 1970. doi: https://doi.org/10.1137/0707001.
- Xiucai Ding. High dimensional deformed rectangular matrices with applications in matrix denoising. Bernoulli, 26(1):387–41, 2020. doi: https://doi.org/10.3150/19-BEJ1129.
- Jing Dong and Xin T Tong. Replica exchange for non-convex optimization. J. Mach. Learn. Res., 22(1):7826–7884, 2021. doi: https://www.jmlr.org/papers/v22/20-697.html.
- David Donoho and Matan Gavish. Minimax risk of matrix denoising by singular value thresholding. Ann. Stat., 42(6):2413–2440, 2014. doi: DOI:10.1214/14-AOS1257.
- David L Donoho, Matan Gavish, and Iain M Johnstone. Optimal shrinkage of eigenvalues in the spiked covariance model. Ann. Stat., $46(4)$:1742–1778, 2018. doi: https://doi.org/10.1214/ 17-AOS1601.
- David L Donoho et al. High-dimensional data analysis: The curses and blessings of dimensionality. AMS math challenges lecture, 1(2000):1–32, 2000. doi: https://api.semanticscholar.org/ CorpusID:5293263.
- Jianqing Fan, Weichen Wang, and Yiqiao Zhong. An l_{∞} eigenvector perturbation bound and its application to robust covariance estimation. J. Mach. Learn. Res., 18(207):1–42, 2018. doi: https://jmlr.org/papers/volume18/16-140/16-140.pdf.
- John A Hartigan and Manchek A Wong. Algorithm as 136: A k-means clustering algorithm. J. R. Stat. Soc. Ser. C Appl. Stat., 28(1):100–108, 1979. doi: http://dx.doi.org/10.2307/2346830.
- Trevor Hastie, Robert Tibshirani, Jerome H Friedman, and Jerome H Friedman. The Elements of Statistical Learning: Data Mining, Inference, and Prediction, volume 2. Springer, 2009.
- Y Hu and W Wang. Network-adjusted covariates for community detection. Biometrika, page asae011, 02 2024. doi: 10.1093/biomet/asae011.
- Jiashun Jin and Wanjie Wang. Influential features pca for high dimensional clustering. Ann. Stat., 44(6):2323–2359, 2016. doi: DOI:10.1214/15-AOS1423.
- Jiashun Jin, Zheng Tracy Ke, and Wanjie Wang. Phase transitions for high dimensional clustering and related problems. Ann. Stat., 45(5):2151–2189, 2017. doi: DOI:10.1214/16-AOS1522.
- Iain M Johnstone and Arthur Yu Lu. On consistency and sparsity for principal components analysis in high dimensions. JASA, 104(486):682–693, 2009. doi: https://doi.org/10.1198/jasa. 2009.0121.
- Ian Jolliffe. Principal component analysis. Encyclopedia of Statistics in Behavioral Science, 2005. doi: https://doi.org/10.1002/0470013192.bsa501.
- Ian T Jolliffe. Discarding variables in a principal component analysis. i: Artificial data. J. R. Stat. Soc. Ser. C Appl. Stat., 21(2):160–173, 1972. doi: https://doi.org/10.2307/2346488.
- Yuehaw Khoo, Xin T Tong, Wanjie Wang, and Yuguan Wang. Temporal label recovery from noisy dynamical data. $arXiv$ preprint $arXiv:2406.13635$, 2024. doi: https://doi.org/10.48550/arXiv. 2406.13635.
- William E Leeb. Matrix denoising for weighted loss functions and heterogeneous signals. SIMODS, 3(3):987–1012, 2021. doi: https://doi.org/10.1137/20M1319577.
- Song Mei, Yu Bai, and Andrea Montanari. The landscape of empirical risk for nonconvex losses. Ann. Stat., 46(6A):2747–2774, 2018. doi: https://www.jstor.org/stable/26542881.
- Scott Menard. Applied Logistic Regression Analysis. Number 106. Sage, 2002.
- Andrea Montanari, Feng Ruan, and Jun Yan. Adapting to unknown noise distribution in matrix denoising. arXiv preprint arXiv:1810.02954, 2018. doi: https://doi.org/10.48550/arXiv.1810. 02954.
- Raj Rao Nadakuditi. Optshrink: An algorithm for improved low-rank signal matrix denoising by optimal data-driven singular value shrinkage. IEEE Trans. Inf. Theory, 60(5):3002–3018, 2014. doi: 10.1109/TIT.2014.2311661.
- Mahamed GH Omran, Andries P Engelbrecht, and Ayed Salman. An overview of clustering methods. Intell. Data Anal., 11(6):583–605, 2007. doi: 10.3233/IDA-2007-11602.
- Sean O'Rourke, Van Vu, and Ke Wang. Matrices with gaussian noise: Optimal estimates for singular subspace perturbation. *IEEE Trans. Inf. Theory*, pages 1978–2002, 2024. doi: https: //doi.org/10.1109/TIT.2023.3331010.
- Theodosios Pavlidis and Steven L Horowitz. Segmentation of plane curves. IEEE Trans. Comput.,, 100(8):860–870, 1974. doi: 10.1109/T-C.1974.224041.
- David Pollard. Strong consistency of k-means clustering. Ann. Stat., 9(1):135–140, 1981. doi: https://doi.org/10.1214/aos/1176345339.
- Alexander Rakhlin and Andrea Caponnetto. Stability of k-means clustering. NeurIPS, 19, 2006. doi: https://proceedings.neurips.cc/paper\ files/paper/2006/file/ 58191d2a914c6dae66371c9dcdc91b41-Paper.pdf.
- C Radhakrishna Rao. The use and interpretation of principal component analysis in applied research. Sankhya A, 26(4):329–358, 1964. doi: https://www.jstor.org/stable/25049339.
- Markus Reiss and Martin Wahl. Nonasymptotic upper bounds for the reconstruction error of pca. Ann. Stat., 48(2):1098–1123, 2020. doi: https://doi.org/10.1214/19-AOS1839.
- Mark Rudelson and Roman Vershynin. Hanson-wright inequality and sub-gaussian concentration. Electron. Commun. Probab., 18(92):1–9, 2013. doi: https://doi.org/10.1214/ECP.v18-2865. URL <https://doi.org/10.1214/ECP.v18-2865>.
- George AF Seber and Alan J Lee. Linear Regression Analysis, volume 330. John Wiley & Sons, 2003.
- Shai Shalev-Shwartz and Shai Ben-David. Understanding Machine Learning: From Theory to Algorithms. Cambridge university press, 2014.
- Roger N Shepard. The analysis of proximities: multidimensional scaling with an unknown distance function. i. Psychometrika, 27(2):125–140, 1962. doi: https://doi.org/10.1007/BF02289630.
- Amit Singer. From graph to manifold laplacian: The convergence rate. Appl. Comput. Harmon. A., 21(1):128–134, 2006. doi: https://doi.org/10.1016/j.acha.2006.03.004.
- Ashbindu Singh and Andrew Harrison. Standardized principal components. Int. J. Remote Sens, 6(6):883–896, 1985. doi: https://doi.org/10.1080/01431168508948511.
- Gilbert W Stewart and Ji-guang Sun. *Matrix Perturbation Theory*. Academic Press, 1990.
- Henry Stone. Approximation of curves by line segments. Math. Comput., 15(73):40–47, 1961. doi: https://api.semanticscholar.org/CorpusID:121847482.
- Terence Tao. *Topics in Random Matrix Theory*, volume 132. American Mathematical Society, 2012.
- Xin T. Tong, Wanjie Wang, and Yuguan Wang. Supplementary material of "uniform error bound for pca matrix denoising". 2024.
- John W Tukey. A survey of sampling from contaminated distributions. contributions to probability and statistics. Essays in honor of Harold Hotelling, 1960.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. arXiv preprint arXiv:1011.3027, 2010. doi: https://doi.org/10.48550/arXiv.1011.3027.
- Ulrike Von Luxburg. A tutorial on spectral clustering. Stat. Comput., 17:395–416, 2007. doi: https://doi.org/10.1007/s11222-007-9033-z.
- Ulrike Von Luxburg, Mikhail Belkin, and Olivier Bousquet. Consistency of spectral clustering. Ann. Stat., 36(2):555–586, 2008. doi: 10.1214/009053607000000640.
- Van Vu. Singular vectors under random perturbation. Random Struct. Algor., 39(4):526–538, 2011. doi: https://doi.org/10.1002/rsa.20367.
- Grace Wahba et al. Support vector machines, reproducing kernel hilbert spaces and the randomized gacv. In Advances in Kernel Methods-Support Vector Learning, volume 6, pages 69–87. Citeseer, 1999. doi: 10.7551/mitpress/1130.003.0009.
- Per-Åke Wedin. Perturbation bounds in connection with singular value decomposition. BIT , 12: 99–111, 1972. doi: https://doi.org/10.1007/BF01932678.
- Yi Yu, Tengyao Wang, and Richard J Samworth. A useful variant of the davis–kahan theorem for statisticians. Biometrika, 102(2):315–323, 2015. doi: https://doi.org/10.1093/biomet/asv008.
- Anru R Zhang, T Tony Cai, and Yihong Wu. Heteroskedastic pca: Algorithm, optimality, and applications. Ann. Stat., 50(1):53–80, 2022. doi: https://doi.org/10.1214/21-AOS2074.

A Detailed verification in Theorem [1](#page-5-1)

Lemma [1](#page-5-1). Suppose the settings of Theorem 1 hold. If we fix $\lambda_r \ge c_X \sqrt{n}$ and $d \gtrsim n$,

$$
\frac{\sigma(\sqrt{n} + \sqrt{d})}{\lambda_r} + \frac{\sigma n \sqrt{\log n}}{\lambda_r^2} + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n}))
$$

$$
\lesssim \sqrt{\frac{d}{n}} \sigma(1 + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^4 d}{n} \log n) + \sqrt{\log n} \sigma.
$$

Otherwise, if we fix $\lambda_r \geq c_X \sqrt{n}$ and $d \lesssim n$,

$$
\frac{\sigma}{\lambda_r^2}(\sqrt{nd} + n\sqrt{\log n}) + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n})) \lesssim \sqrt{\log n} \sigma(1 + \sigma^4 \sqrt{\frac{d \log n}{n}}).
$$

Proof. Recall that $\sqrt{n} \geq \lambda_r \geq 1 + C_1 \sigma(\sqrt{n} +$ √ d),

$$
\frac{d\sigma^2}{\lambda_r^2} \le \frac{\sigma\sqrt{nd}}{C_1\lambda_r^2}, \quad \frac{d\sigma^2}{\lambda_r^2} \le \frac{\sigma\sqrt{d}}{C_1\lambda_r}, \quad \frac{\sigma}{\lambda_r^2}\sqrt{n\log n} \le \frac{\sigma}{\lambda_r^2}n\sqrt{\log n}, \quad \frac{\sigma}{\lambda_r^2}\sigma n \le \frac{\sigma}{\lambda_r^2}n\sqrt{\log n}.
$$

If we fix $\lambda_r \geq c_X \sqrt{n}$ and $d \gtrsim n$,

$$
\frac{\sigma(\sqrt{n} + \sqrt{d})}{\lambda_r} + \frac{\sigma n \sqrt{\log n}}{\lambda_r^2} + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n}))
$$
\n
$$
\leq \frac{\sqrt{d}\sigma}{\lambda_r} + \frac{\sigma n \sqrt{\log n}}{\lambda_r^2} + \frac{\sigma^2}{\lambda_r^4} \sqrt{\log n}(n\sqrt{d} + \sigma^2 n\sqrt{nd} + \sigma^2 d\sqrt{nd})(1 + \sigma \sqrt{\log n}) + \frac{\Gamma}{\lambda_r^2}
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma + \sqrt{\log n}\sigma + \frac{\sigma^2}{n^2} \sqrt{\log n}(n\sqrt{d} + \sigma^2 d\sqrt{nd})(1 + \sigma \sqrt{\log n}) + \frac{\Gamma}{n}
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma + \sqrt{\log n}\sigma + \frac{\sigma^2}{n} \sqrt{\log n} \sqrt{d} + \frac{\sigma^2}{n^2} \sqrt{n d \log n}\sigma^2 d + \frac{\sigma^3}{n} \log n \sqrt{d} + \frac{\sigma^3}{n^2} \log n\sigma^2 d\sqrt{nd} + \frac{\Gamma}{n}
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma(1 + \frac{\sigma}{\sqrt{n}}\sqrt{\log n} + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^2}{\sqrt{n}} \log n + \frac{\sigma^4 d}{n} \log n) + \sqrt{\log n}\sigma + \frac{\Gamma}{n}
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma(1 + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^4 d}{n} \log n) + \sqrt{\log n}\sigma + \frac{\sigma}{n^2} \sqrt{\log n}(n + \sigma^2 d\sqrt{n})(1 + \sigma \sqrt{\log n})
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma(1 + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^4 d}{n} \log n) + \sqrt{\log n}\sigma + \frac{\sigma}{n^2} \sqrt{\log n}(n + \sigma^2 d\sqrt{n} + \sigma \sqrt{\log n} + \sigma^3 d\sqrt{n \log n})
$$
\n
$$
\leq \sqrt{\frac{d}{n}}\sigma(1 + \frac{\sigma^3 d}{n} \sqrt{\log n} + \frac{\sigma^4 d}{n
$$

Similarly, if we fix $\lambda_r \geq c_X \sqrt{n}$ and $d \lesssim n$,

$$
\frac{\sigma}{\lambda_r^2}(\sqrt{nd} + n\sqrt{\log n}) + \frac{1}{\lambda_r^2} \Gamma(1 + \sigma(\sqrt{d} + \sqrt{\log n}))
$$
\n
$$
\lesssim \sqrt{\log n}\sigma + \frac{\sigma^2}{n^2}\sqrt{\log n}(n + \sigma^2 n\sqrt{n})(1 + \sigma\sqrt{\log n})(\sqrt{d} + \sqrt{\log n}) + \frac{\Gamma}{n}
$$
\n
$$
\lesssim \sqrt{\log n}\sigma(1 + \frac{\sigma}{n}(1 + \sigma^2\sqrt{n})(1 + \sigma\sqrt{\log n})(\sqrt{d} + \sqrt{\log n}))
$$
\n
$$
= \sqrt{\log n}\sigma\left(1 + \frac{\sigma\sqrt{d}}{n} + \frac{\sigma^3\sqrt{d}}{\sqrt{n}} + \frac{\sigma^2\sqrt{d\log n}}{n} + \frac{\sigma^4\sqrt{d\log n}}{\sqrt{n}} + \frac{\sigma\sqrt{\log n}}{n} + \frac{\sigma^3\sqrt{\log n}}{\sqrt{n}} + \frac{\sigma^2\log n}{n} + \frac{\sigma^4\log n}{\sqrt{n}}\right) + \frac{\Gamma}{n}
$$
\n
$$
\lesssim \sqrt{\log n}\sigma(1 + \sigma^4\sqrt{\frac{d\log n}{n}}) + \frac{\sigma}{n^2}\sqrt{\log n}(n + \sigma^2 n\sqrt{d} + \sigma\sqrt{\log n} + \sigma^3 n\sqrt{d\log n})
$$
\n
$$
\lesssim \sqrt{\log n}\sigma(1 + \sigma^4\sqrt{\frac{d\log n}{n}}).
$$

 $\hfill \square$

 \Box

B Proofs for zigzag line data objects

Lemma 2. Consider $A \in \mathbb{R}^{l \times m}$, $B \in \mathbb{R}^{m \times n}$, Suppose $l \ge \max\{m, n\}$ then

$$
\lambda_{\min\{m,n\}}(AB) \ge \lambda_m(A)\lambda_{\min\{m,n\}}(B).
$$

Proof. Denote the SVD of B as $B = U_B \Lambda_B V_B^T$. We discuss the case $m \ge n$ and $m \le n$ separately. Suppose $m \geq n$, then

$$
\lambda_n(AB) = \sqrt{\lambda_n(B^T A^T A B)} \ge \lambda_m(A) \sqrt{\lambda_n(B^T B)} = \lambda_m(A) \lambda_n(B).
$$

Suppose $m \leq n$, then let $U_B \in \mathbb{R}^{m \times m}$, $\Lambda_B \in \mathbb{R}^{m \times m}$, and $V_B \in \mathbb{R}^{n \times m}$. We see

$$
\lambda_m(AB) = \sqrt{\lambda_m(ABB^T A^T)} \ge \lambda_m(B)\sqrt{\lambda_m(AA^T)} = \lambda_m(A)\lambda_m(B).
$$

Proof of Proposition [2.](#page-13-0) According to the model, we rewrite $X_i^T = x_0^T + T_i^T W$, where $x_0 = \mathbb{E}[X_i]$ is a linear combination of $\{v_1, v_2, \cdots, v_R\}, T_i^T \in \mathbb{R}^{1 \times R}$, and $W \in \mathbb{R}^{R \times d}$ with the *i*-th row being v_i^T . The vector T_i is generated by $s_i \sim Unif(0,1)$, which can be further written as

$$
T_{i,j} = \begin{cases} t_{j+1} - t_j, & s_i \ge t_{j+1}; \\ s_i - t_j, & t_j < s_i \le t_{j+1}; \\ 0, & s_i < t_j. \end{cases}
$$

Therefore, $\sum_j T_{i,j} = s_i \leq 1$.

Consider the largest eigenvalue first. Since all v_i 's have unit norm and $s_i \sim Unif(0,1)$, so Consider the largest eigenvalue inst. Solventually $||X_i|| \leq 1$. Hence, $||X|| \leq \sqrt{n}$ by Theorem [2.](#page-12-1)

Now consider the smallest singular value. Again, by Theorem [2,](#page-12-1) we only need to show the r -th eigenvalue of $cov(X_i)$ is non-zero. Note that

$$
cov(X_i) = cov(W^T T_i) = W^T cov(T_i)W.
$$

Rewrite $W = SQ$, where $Q \in \mathbb{R}^{r \times d}$ contains the basis of the r-dimensional subspace spanned by $\{v_1, v_2, \dots, v_R\}$ and $S \in \mathbb{R}^{R \times r}$ contains the linear coefficients so that each row of SQ is v_i^T . Hence, $QQ^T = I_r$ and S has rank r. By Lemma [2](#page-25-0) we have

$$
\lambda_r(\text{cov}(X_i)) \ge \lambda_R(\text{cov}(T_i))\lambda_r^2(S)\lambda_r^2(Q) = \lambda_R(\text{cov}(T_i))\lambda_r^2(S).
$$

Since W has rank r, the r-th singular value of S is non-zero. Hence, it suffices to show $\lambda_R(\text{cov}(T_i))$ 0.

Now we investigate the R-th singular value of $cov(T_i)$, where

$$
\lambda_R(\text{cov}(T_i)) = \min_{a \in \mathbb{R}^R, ||a|| = 1} \text{var}[T_i^T a].
$$

Consider a vector a with $||a|| = 1$. Then $T_i^T a$ is

$$
T_i^T a = (s_i - t_j)a_j + \sum_{k < j} (t_{k+1} - t_k)a_k, \quad t_j < s_i \le t_{j+1}.
$$

Denote $m = \mathbb{E}[T_i^T a]$ and $b(j) = \sum_{k < j} (t_{k+1} - t_k) a_k - m$. Further define

$$
m_1(j) = \mathbb{E}[1_{s_i \in [t_j, t_{j+1}]}(s_i - t_j)] = (t_{j+1} - t_j)^2/2,
$$

$$
m_2(j) = \mathbb{E}[1_{s_i \in [t_j, t_{j+1}]}(s_i - t_j)^2] = (t_{j+1} - t_j)^3/3.
$$

Introduce these terms into $\text{var}(T_i^T a)$ and we have

$$
\begin{split} \text{var}[T_i'a] &= \sum_j \mathbb{E}[1_{s_i \in [t_j, t_{j+1}]}[(s_i - t_j)a_j + b(j)]^2] \\ &= \sum_j (b(j)^2 + 2b(j)m_1(j)a_j + m_2(j)a_j^2) \\ &\ge \sum_j (m_2(j) - (m_1(j))^2)a_j^2 \\ &\ge \min_{j \in [R]} [m_2(j) - (m_1(j))^2] \ge \frac{\rho^3}{3} - \frac{\rho^4}{4}. \end{split}
$$

The first inequality is obtained by optimizing over all possible $b(j)$. The second inequality is obtained since $\sum_j |a_j|^2 = 1$. The last inequality is obtained because $f(x) = x^3/3 - x^4/4$ is increasing on $x \in [0,1]$.

So, the result is proved by taking $c = \lambda_r(S)/2$.

$$
\Box
$$

C Proof for the statistical lower bound

Proof of Theorem [3.](#page-14-3) Consider any estimator \hat{X} based on $Z_{[n]}$ and denote it as $\hat{X}(Z_{[n]})$. We first set up the bound for the noise level σ and then discuss the sample size n.

Conditional on v, X_i and Z_i are independent of $Z_{[n]/i}$. So we consider a new problem where both $Z_{[n]}$ and v are known and the same estimator $\hat{X}_i(Z_{[n]})$. By the Blackwell thereom, the estimator can be improved by

$$
\tilde{X}_i(v, Z_i) = \mathbb{E}[\hat{X}_i(Z_{[n]})|v, Z_i]
$$

in the sense that

$$
\mathbb{E}[\|\tilde{X}_i(v, Z_i) - X_i\|^2] \le \mathbb{E}[\|\hat{X}_i(Z) - X_i\|^2] \le \epsilon^2.
$$
 (23)

Recall that the ℓ_2 error of $\tilde{X}_i(v, Z_i)$ is larger than the Bayes error. We consider the Bayes error. Since v is known, the estimation is $\hat{X}_i = (\hat{t}_i + 1)v$ and the uncertainty comes from \hat{t}_i only. Let P be the projection onto the direction of v and P_{\perp} the projection onto the complementary subspace. For notational simplicity, let $z_i = Z_i^T v / ||v||$ and $e_i = \xi_i^T v / ||v||$. The Bayes estimator of t_i is given by

$$
\hat{t}_{i} = \frac{\int_{0}^{1} t \exp(-\frac{1}{2\sigma^{2}}||Z_{i} - (t+1)v||^{2})dt}{\int_{0}^{1} \exp(-\frac{1}{2\sigma^{2}}||Z_{i} - (t+1)v||^{2})dt}
$$
\n
$$
= \frac{\int_{0}^{1} t \exp(-\frac{1}{2\sigma^{2}}||(z_{i}/||v||)v - (t+1)v||^{2}) \exp(-\frac{1}{2\sigma^{2}}||P_{\perp}Z_{i}||^{2})dt}{\int_{0}^{1} \exp(-\frac{1}{2\sigma^{2}}||(z_{i}/||v||)v - (t+1)v||^{2}) \exp(-\frac{1}{2\sigma^{2}}||P_{\perp}Z_{i}||^{2})dt}
$$
\n
$$
= \frac{\int_{0}^{1} t \exp(-\frac{||v||^{2}}{2\sigma^{2}}|(t+1) - z_{i}/||v|||^{2})dt}{\int_{0}^{1} \exp(-\frac{||v||^{2}}{2\sigma^{2}}|(t+1) - z_{i}/||v|||^{2})dt}
$$
\n
$$
= \frac{\int_{0}^{1} ((t+1) - z_{i}/||v||) \exp(-\frac{||v||^{2}}{2\sigma^{2}}|(t+1) - z_{i}/||v|||^{2})dt}{\int_{0}^{1} \exp(-\frac{||v||^{2}}{2\sigma^{2}}|(t+1) - z_{i}/||v|||^{2})dt} + \frac{z_{i}}{||v||} - 1
$$
\n
$$
= \frac{z_{i}}{||v||} - 1 + \frac{\sigma^{2}}{||v||^{2}} \frac{\phi(\sigma^{-1}(2||v|| - z_{i})) - \phi(\sigma^{-1}(||v|| - z_{i}))}{\phi(\sigma^{-1}(|||v|| - z_{i}))}
$$
\n
$$
= t_{i} + \frac{e_{i}}{||v||} + \frac{\sigma^{2}}{||v||^{2}} \frac{\phi(\sigma^{-1}(2||v|| - z_{i})) - \phi(\sigma^{-1}(||v|| - z_{i}))}{\phi(\sigma^{-1}(|||v|| - z_{i}))}
$$

The estimation error $|\hat{t}_i - t_i|$ consists of $e_i / ||v||$ and a complicated term. We now bound it. Consider a set $B = \{(t_i, e_i) : e_i(t_i - 0.5) > 0, |e_i| > \sigma\}$, then $P(B) = \Phi(-1)/2$, where Φ is the CDF of standard Gaussian. If $t_i > 0.5$, $e_i > 0$, then $z_i > 1.5||v||$ and so $\phi(\sigma^{-1}(2||v|| - z_i)) - \phi(\sigma^{-1}(||v|| - z_i))$ (z_i)) > 0. It suggests that

$$
|\hat{t}_i - t_i| \ge |e_i| / \|v\| > \sigma / \|v\|.
$$
 (24)

With a similar analysis, [\(24\)](#page-27-0) still holds when $t_i < 0.5$ and $\xi_i < 0$. As a conclusion, $|\hat{t}_i - t_i| \ge \sigma / ||v||$ at the occurrence of B.

Further if $||v|| \leq 2$, which happens with probability at least 3/4 by Markov inequality, then

$$
|\hat{t}_i - t_i| \ge \sigma/2,
$$

and the error follows

$$
\mathbb{E}[\|\hat{t}_i - t_i\|^2] \ge \mathbb{E}[\|\hat{t}_i - t_i\|^2 1_B] \ge \frac{1}{8}\sigma^2 P(B) = \frac{\Phi(-1)}{16}\sigma^2.
$$

Combine it with [\(23\)](#page-26-0), we need $\sigma \leq 4\epsilon/\sqrt{\Phi(-1)}$.

Next we consider the sample size n. Here we consider a new problem where both $t_{[n]}$ and $Z_{[n]}$ are observed but v is unknown. Then for any estimator $\hat{X}_i(Z_{[n]})$, we can design an estimator of v as

$$
\hat{v} = \frac{1}{n} \sum_{i=1}^{n} \frac{\hat{X}_i(Z_{[n]})}{t_i + 1}.
$$

Then the error is bounded by

$$
\mathbb{E} \|\hat{v} - v\|^2 = \mathbb{E} \left\| \frac{1}{n} \sum_{i=1}^n \frac{\hat{X}_i(Z_{[n]}) - X_i}{t_i + 1} \right\|^2 \le \mathbb{E}[\max_i \|X_i - \hat{X}_i(Z_{[n]})\|^2] \le \epsilon^2.
$$

Meanwhile, we know l^2 error of $\hat{v}_i(t_{[n]}, Z_{[n]})$ will be larger than the Bayes error of the Bayes estimator for v. The prior of v is $\mathcal{N}(0, \frac{1}{d}I_d)$ and the data $Z_{[n]}$ follows $Z_i \sim \mathcal{N}((t_i+1)v, \sigma^2 I_d)$ with given $t_{[n]}$. Therefore, the posterior distribution of v is also Gaussian, with mean and covariance

$$
\tilde{v} = \frac{\sum_{i=1}^{n} (t_i + 1)Z_i}{\sigma^2 d + \sum_{i=1}^{n} (t_i + 1)^2}, \quad \tilde{C} = \left(\frac{1}{d} + \frac{1}{\sigma^2} \sum_{i=1}^{n} (t_i + 1)^2\right)^{-1} I_d.
$$

The Bayes error is

$$
\mathbb{E}[\|\tilde{v} - v\|^2] = \mathbb{E}[\text{tr}(\tilde{C})] \ge \frac{d\sigma^2}{4n + \sigma^2/d}.
$$

Combine it with that $\mathbb{E}[\|\tilde{v}-v\|^2] \leq \epsilon^2$, we need $d\sigma^2/(4n+\sigma^2/d) \leq \epsilon^2$. Since we assume $\epsilon^2 \leq \frac{1}{16}d^2$, we have $\sigma^2/d \leq \frac{4}{15}n$, which further have $\mathbb{E}[[\|\tilde{v}-v\|^2]] \geq \frac{d\sigma^2}{5n^2}$. So $n < \frac{d\sigma^2}{5\epsilon^2}$ is impossible.

D Denoising applications

D.1 Empirical risk minimization for supervised learning

Empirical risk minimization (ERM) is a widely used approach for training statistical models [Hastie](#page-21-5) [et al. \[2009\]](#page-21-5), [Shalev-Shwartz and Ben-David \[2014\]](#page-22-17). In ERM, the dataset consists of pairs (X_i, y_i) , where $X_i \in \mathbb{R}^d$ and y_i is typically a scalar. The prediction error is measured by a loss function $F(X_i, y_i, \theta)$, where the specific definition of F varies depending on the model. The goal of ERM is to find the optimal model parameters θ^* that minimize the empirical prediction error as follows.

$$
\theta^* = \arg\min_{\theta} f(\theta), \quad f(\theta) := \frac{1}{n} \sum_{i=1}^n F(X_i, y_i, \theta). \tag{25}
$$

Some well-known examples are listed below:

• In standard linear regression [Seber and Lee \[2003\]](#page-22-18), $y_i \in \mathbb{R}$, $\theta \in \mathbb{R}^d$ and F is given by

$$
F(X_i, y_i, \theta) = \frac{1}{2} (\theta^T X_i - y_i)^2.
$$

• In linear regresion with Tuckey's biweight loss function [Tukey \[1960\]](#page-22-19),

$$
F(X_i, y_i, \theta) = \rho(\theta^T X_i - y_i), \quad \text{where } \rho(u) = \frac{c^2}{6} (1 - (1 - (u/c)^2)^3 1_{|u| < c}).
$$

Here c is some threshold constant. The introduction of the ρ function is to reduce the influence of possible outliers from data.

• In logistic regression [Menard \[2002\]](#page-21-17), $y_i \in \{+1, -1\}$ and the loss function F is given by

$$
F(X_i, y_i, \theta) = \log(1 + \exp(-y_i X_i^T \theta)).
$$

• In reproducing kernel Hilbert space regression [Wahba et al. \[1999\]](#page-23-4), we have a reproducing kernel $k(\cdot, \cdot)$ and its corresponding Hilbert space H. The goal is to find a function $f \in \mathcal{H}$ to minimize $L(f) = \frac{1}{n} \sum_{i=1}^{n} (f(x_i) - y_i)^2$. The kernel regression can also be viewed as a standard ERM problem. To do it, we parameterize $f_{\theta}(x) = \langle \theta, \varphi(x) \rangle$, where $\varphi(x)$ is the feature map and the kernel function $k(x, y) = \langle \varphi(x), \varphi(y) \rangle$. The loss function F is

$$
F(X_i, y_i, \theta) = \frac{1}{2} (\langle \theta, \varphi(X_i) \rangle - y_i)^2.
$$

• In an m-layered neural networks, we denote $\theta_i \in \mathbb{R}^d$ as the loading of the *i*-th layer. Hence, the parameters matrix is $\theta = [\theta_1, \dots, \theta_m] \in \mathbb{R}^{m \times d}$. The loss is given by

$$
F(X_i, y_i, \theta) = (y_i - \sigma(\theta_m^T \cdots \sigma(\theta_2^T \sigma(\theta_1^T X_i)) \cdots))^2,
$$

where the function $\sigma(\cdot)$ can be taken as various nonlinear functions, such as ReLu and sigmoid function.

In many applications, we do not have access to the clean data X_i , but only the noisy data $Z_i = X_i + \xi_i$. The high noise in Z_i will cause large error in the estimation of θ , if we use Z_i instead of X_i in F directly. It's natural to guess that the denoised data \hat{X}_i will lead to an estimator reasonably close to θ^* . Let $\hat{\theta}^*$ be the estimator, where

$$
\hat{\theta}^* = \arg\min \hat{f}(\theta), \quad \hat{f}(\theta) := \frac{1}{n} \sum_{i=1}^n F(\hat{X}_i, y_i, \theta). \tag{26}
$$

It is natural to ask how would $\hat{\theta}^*$ performs under the empirical risk function f in [\(25\)](#page-28-0), and how close would $\hat{\theta}^*$ be from θ^* . To answer these questions, we made the following assumptions.

Assumption 3. The loss function $F(x, y_i, \theta)$ is globally Lipschitz in x. That is, there are constants $L > 0$ and $\Delta > 0$, so that when $||z - x|| \leq \Delta$, there is

$$
|F(z, y_i, \theta) - F(x, y_i, \theta)| \le L ||z - x||.
$$

In general Assumption [3](#page-29-0) is easy to verify when the data are generated in bounded domains, even for complex nonlinear models such as neural networks. But it alone cannot guarantee θ^* is unique, so if we want to infer parameter error, we need some additional conditions.

Assumption 4. The empirical loss function f in [\(25\)](#page-28-0) has a unique minimizer θ^* and it is also a local minimum. That is, there are constants $\gamma, \delta, \lambda_f > 0$, so that $f(\theta) - f(\theta^*) \geq \gamma$ when $\|\theta - \theta^*\| \geq \delta$ and $\nabla^2 f(\theta) \succeq \lambda_f I$ when $\|\theta - \theta^*\| \leq \delta$.

Suppose f is strongly convex, Assumption [4](#page-29-1) holds immediately. This is also the most well understood ERM regime. Assumption [4](#page-29-1) also allows for general non-convex problems [Shalev-](#page-22-17)[Shwartz and Ben-David \[2014\]](#page-22-17). Similar version of it can be found in machine learning literature where finding θ^* is of interest [Dong and Tong \[2021\]](#page-20-15).

Proposition 5. Consider an ERM problem where Assumption [3](#page-29-0) holds. Suppose $\|\hat{X} - X\|_{2,\infty} \leq \epsilon$ where $L\epsilon < \Delta$. Let θ^* and $\hat{\theta}^*$ be the minimizers of f and \hat{f} , respectively. Then $f(\hat{\theta}^*)$ is close to the minimum $f(\theta^*)$, in the sense that

$$
|f(\hat{\theta}^*) - f(\theta^*)| \le 2L\epsilon.
$$

Suppose Assumption [4](#page-29-1) holds in addition with $L \in \gamma$, then the two minimizers are close,

$$
\|\hat{\theta}^* - \theta^*\|^2 \le 2L\epsilon/\lambda_f.
$$

Proposition [5](#page-29-2) indicates that the ERM training result using the PCA denoised data is as good the training result using clean data, assuming Assumption [3.](#page-29-0) If Assumption [4](#page-29-1) is also in place, then the learned parameters will also be close to each other. The proof can be found in the Supplementary Material.

Proof. For the first claim, simply note that,

$$
|\hat{f}(\theta) - f(\theta)| \leq \frac{1}{n} \sum_{i=1}^{n} |F(\hat{X}_i, y_i, \theta) - F(X_i, y_i, \theta)| \leq L\epsilon.
$$

Note that $f(\theta^*) \leq f(\hat{\theta}^*)$ and $\hat{f}(\hat{\theta}^*) \leq \hat{f}(\theta^*)$, and we have

$$
f(\hat{\theta}^*) \leq \hat{f}(\hat{\theta}^*) + |\hat{f}(\hat{\theta}^*) - f(\hat{\theta}^*)| \leq \hat{f}(\hat{\theta}^*) + L\epsilon \leq \hat{f}(\theta^*) + L\epsilon
$$

$$
\leq f(\theta^*) + |\hat{f}(\theta^*) - f(\theta^*)| + L\epsilon \leq f(\theta^*) + 2L\epsilon.
$$

This leads to our first claim.

For the second claim, note that by $|f(\hat{\theta}^*) - f(\theta^*)| \leq 2L\epsilon < \gamma$, we have $\|\theta^* - \theta\| \leq \delta$. Then by Taylor expansion, we have

$$
f(\hat{\theta}^*) = f(\theta^*) + \frac{1}{2}(\hat{\theta}^* - \theta^*)^T \nabla^2 f(\tilde{\theta})(\hat{\theta}^* - \theta^*) \ge f(\theta^*) + \frac{1}{2}\lambda_f \|\hat{\theta}^* - \theta^*\|^2.
$$

Combining it with $|f(\hat{\theta}^*) - f(\theta^*)| \leq 2L\epsilon$ again, it leads to

$$
\|\hat{\theta}^* - \theta^*\|^2 \le 4L\epsilon/\lambda_f.
$$

 \Box

D.2 Proofs for other applications

Proof of Proposition [3.](#page-15-0) For the first claim, simply note that,

$$
|\widehat{f}(\mathcal{C}, m) - f(\mathcal{C}, m)| \leq \frac{1}{n} \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} |||X_i - m_k||^2 - ||\widehat{X}_i - m_k||^2|
$$

$$
\leq \frac{1}{n} \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} (2\delta \epsilon + \epsilon^2) \leq 2\epsilon.
$$

The last inequality comes from the fact $\delta \leq 1$ because $||X_i|| \leq 1$ for all $i \in [n]$ and $\epsilon = o(1)$.

Recall that $||X_i|| \leq 1 + \epsilon$ and so likewise $|\hat{f}(\hat{\mathcal{C}}, \hat{m}) - f(\hat{\mathcal{C}}, \hat{m})| \leq 2\epsilon$. This further leads to

 $f(\hat{\mathcal{C}}, \hat{m}) \leq \hat{f}(\hat{\mathcal{C}}, \hat{m}) + 2\epsilon \leq \hat{f}(\mathcal{C}, m) + 2\epsilon \leq f(\mathcal{C}, m) + 4\epsilon.$

Consider the second claim. Note that given any division \tilde{C} , then the minimizer m of $f(\tilde{C}, m)$ and $\hat{f}(\tilde{\mathcal{C}}, m)$ can be found as

$$
m_k(\tilde{C}) = \frac{\sum_{i \in \tilde{C}_k} X_i}{|\tilde{C}_k|}, \quad \hat{m}_k(\tilde{C}) = \frac{\sum_{i \in \tilde{C}_k} \hat{X}_i}{|\tilde{C}_k|}.
$$

For the special case $\tilde{\mathcal{C}} = \hat{\mathcal{C}}$, we have $\hat{m}_k = \hat{m}_k(\hat{\mathcal{C}})$.

Now we consider $\hat{f}(\hat{\mathcal{C}}, \hat{m})$. According to the definition,

$$
\hat{f}(\hat{\mathcal{C}}, \hat{m}) \le \hat{f}(\mathcal{C}, \hat{m}(\mathcal{C})) \le \hat{f}(\mathcal{C}, m) = \frac{1}{n} \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} ||\hat{X}_i - m_k||^2
$$

$$
\le \frac{2}{n} \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} (||\hat{X}_i - X_i||^2 + ||X_i - m_k||^2)
$$

$$
\le 2\delta^2 + 2\epsilon^2.
$$

Now we consider the lower bound.

$$
\hat{f}(\hat{\mathcal{C}}, \hat{m}) = \frac{1}{n} \sum_{k \in [K]} \sum_{i \in \mathcal{C}_k} ||\hat{X}_i - \hat{m}_{j:i \in \hat{\mathcal{C}}_j}||^2
$$
\n
$$
\geq \frac{1}{n} \sum_{k} \sum_{i \in \mathcal{C}_k} (-||\hat{X}_i - m_k||^2 + \frac{1}{2}||m_k - \hat{m}_{j:i \in \hat{\mathcal{C}}_j}||^2)
$$
\n
$$
= -\hat{f}(\mathcal{C}, m) + \frac{1}{2n} \sum_{k} \sum_{i \in \mathcal{C}_k} ||m_k - \hat{m}_{j:i \in \hat{\mathcal{C}}_j}||^2.
$$
\n(27)

Combining the two bound above and we have

 $\overline{1}$

$$
\frac{1}{n}\sum_{k}\sum_{i\in\mathcal{C}_k}||m_k - \hat{m}_{j:i\in\hat{\mathcal{C}}_j}||^2 \le 2(\hat{f}(\mathcal{C},m) + \hat{f}(\hat{\mathcal{C}},\hat{m})) \le 8(\delta^2 + \epsilon^2).
$$

Now we compare $\hat{\mathcal{C}}$ and \mathcal{C} . To simplify the notations, we use $\ell(i)$ to denote the true label of node i so that $i \in \mathcal{C}_{\ell(i)}$ and use $\hat{\ell}(i)$ to denote the estimated label that $i \in \hat{\mathcal{C}}_{\hat{\ell}(i)}$. We need a projection $\pi : [K] \to [K]$ so that ℓ and $\pi(\ell)$ will match. The projection π is defined by matching the nearest centers:

$$
\pi(k) = \arg\min_{j} \|m_j - \hat{m}_k\|, \quad k \in [K].
$$
\n(28)

Then the overall distances between centers are

$$
c_0 \|m_k - \hat{m}_{\pi(k)}\|^2 \le \frac{1}{n} \sum_k \sum_{i \in C_k} \|m_k - \hat{m}_{\pi(k)}\|^2
$$

$$
\le \frac{1}{n} \sum_k \sum_{i \in C_k} \|m_k - \hat{m}_{\hat{\ell}(i)}\|^2 \le 8(\delta^2 + \epsilon^2).
$$
 (29)

Given $\pi(\hat{\ell})$, we want to show $\pi(\hat{\ell}) = \ell$. To prove it, we suppose there is a data point i where $\pi(\hat{\ell}(i)) \neq \ell(i) = k$. Define a new label vector ℓ' that differs from $\hat{\ell}$ on the data point i only, where $\ell'(i) = \pi^{-1}(k)$. Correspondingly we have C'. We want to show that C' will result in a smaller \hat{f} than $\hat{f}(\hat{\mathcal{C}}, \hat{m})$, and hence $\hat{\mathcal{C}}$ will not be a solution of k-means.

$$
\hat{f}(\mathcal{C}', \hat{m}(\mathcal{C}')) - \hat{f}(\hat{\mathcal{C}}, \hat{m}) \le \hat{f}(\mathcal{C}', \hat{m}) - \hat{f}(\hat{\mathcal{C}}, \hat{m})
$$
\n
$$
= \frac{1}{n} \sum_{j=1}^{n} ||\hat{X}_j - \hat{m}_{\ell'(j)}||^2 - \frac{1}{n} \sum_{j=1}^{n} ||\hat{X}_j - \hat{m}_{\hat{\ell}(j)}||^2
$$
\n
$$
= \frac{1}{n} ||\hat{X}_i - \hat{m}_{\ell'(i)}||^2 - \frac{1}{n} ||\hat{X}_i - \hat{m}_{\hat{\ell}(i)}||^2,
$$

where the last equality comes from that $\hat{\ell}(j) = \ell'(j)$ for all $j \neq i$. Further,

$$
\|\hat{X}_i - \hat{m}_{\ell'(i)}\| \le \|\hat{X}_i - m_{\ell(i)}\| + \|m_{\ell(i)} - \hat{m}_{\pi(\ell(i))}\|
$$

$$
\le \delta + \epsilon + 2\sqrt{2(\delta^2 + \epsilon^2)/c_0}.
$$

Recall that we assume $\pi(\hat{\ell}(i)) \neq \ell(i) = k$. Denote $j = \pi(\hat{\ell}(i))$, then

$$
\begin{aligned} \|\hat{X}_i - \hat{m}_{\hat{\ell}(i)}\| &\ge \|m_k - m_j\| - (\|\hat{X}_i - m_k\| + \|m_j - \hat{m}_{\hat{\ell}(i)}\|) \\ &\ge c_m - (\delta + \epsilon + 2\sqrt{2(\delta^2 + \epsilon^2)/c_0}) > \|\hat{X}_i - \hat{m}_{\ell'(i)}\|. \end{aligned}
$$

So $\hat{f}(\mathcal{C}',\hat{m}(\mathcal{C}')) < \hat{f}(\hat{\mathcal{C}},\hat{m})$ and ℓ' will be a strictly better solution to the k-means objective. This contradicts the definition of $\hat{\ell}$. So for all data points, $\pi(\hat{\ell}) = \ell$. \Box

Proof of Proposition [4.](#page-16-1) Consider $||\hat{L} - L||_{\infty}$ first. Since $||X - \hat{X}||_{2,\infty} \leq \epsilon$ and $k(x, y)$ is l-Lipschitz in x and y, so there is a constant $C_3 = C_3(l) > 0$ so that

$$
|(k(X_i, X_j) - k(\hat{X}_i, \hat{X}_j))|
$$

\n
$$
\leq |(k(X_i, X_j) - k(\hat{X}_i, X_j))| + |(k(\hat{X}_i, X_j) - k(\hat{X}_i, \hat{X}_j))| \leq C_3 \epsilon,
$$

and

$$
|D_{ii} - \hat{D}_{i,i}| = \left| \sum_{j \in [n]} (k(X_i, X_j) - k(\hat{X}_i, \hat{X}_j)) \right| \leq C_3 n\epsilon.
$$

Hence, for the normalized Laplacian, there is a $C_4 = C_4(K, l)$

$$
|\hat{L}_{i,j} - L_{i,j}| = |\frac{k(X_i, X_j)}{\sqrt{D_{i,i}D_{j,j}}} - \frac{k(\hat{X}_i, \hat{X}_j)}{\sqrt{\hat{D}_{i,i}\hat{D}_{j,j}}}| \le \frac{C_4\epsilon}{n}.
$$

Therefore, the matrix ℓ_{∞} and ℓ_1 norm follows that

$$
\|\hat{L} - L\|_1 = \|\hat{L} - L\|_{\infty} = \max_{i} \sum_{j} |\hat{L}_{i,j} - L_{i,j}| \le C_4 \epsilon.
$$

The first identity comes from the fact that L,\hat{L} are symmetric. We further present a result about $\|\hat{L}\|_{\infty}$ here. Since $\hat{D}_{i,i} \ge n/K$ and $\hat{A}_{i,j} \le K$,

$$
\|\hat{L}\|_{\infty} = \max_{i} \sum_{j} |\hat{L}_{i,j}| \le \max_{i} \{1 + \sum_{j} \frac{\hat{A}_{i,j}}{\sqrt{\hat{D}(i,i)\hat{D}(j,j)}}\} \le 1 + K^2.
$$
 (30)

Consider the eigenvalues and eigenvectors. Since λ_0 is simple, it has a neighborhood B of radius δ so that its intersection with the spectrum of $\mathcal L$ is $\{\lambda\}$. Theorem 15 of [Von Luxburg et al.](#page-22-0) [\[2008\]](#page-22-0) demonstrates that there is an N_0 , so that when $n > N_0$, the intersection of B with spectrum of L is $\{\lambda\}$. In other words, λ is a simple eigenvalue for L.

By Weyl's inequality, $|\lambda_i(\hat{L}) - \lambda_i(L)| \leq ||\hat{L} - L|| \leq C_4 \epsilon$, where the last inequality comes from $\|\hat{L}-L\| \leq \sqrt{\|\hat{L}-L\|_1 \|\hat{L}-L\|_{\infty}} \leq C_4 \epsilon$. Therefore, \hat{L} has an eigenvalue $\hat{\lambda}$ so that

$$
|\lambda - \hat{\lambda}| \le C_4 \epsilon.
$$

Let $\hat{P}v$ be the eigen-projection of v onto the one-dimensional space spanned by \hat{v} in l_{∞} norm, i.e.

$$
\hat{P}v = \hat{c}\hat{v}, \quad \text{where } \hat{c} = \arg\min_{c} \|v - c\hat{v}\|_{\infty}.
$$

Theorem 7 of [Von Luxburg et al. \[2008\]](#page-22-0) demonstrates that for sufficiently small ϵ , there is a constant C independent of d and n

$$
\|v-\hat{P}v\|_\infty\leq C(\|(L-\hat{L})v\|_\infty+\|v\|_\infty\|(L-\hat{L})\hat{L}\|_\infty)\leq C\epsilon\|v\|_\infty.
$$

Meanwhile, Proposition 18 of [Von Luxburg et al. \[2008\]](#page-22-0) shows that there is an $a \in \{\pm 1\}$ so that

$$
||v-a\hat v||_\infty\leq 2||v-\hat Pv||_\infty\leq 2C\epsilon ||v||_\infty.
$$

So the second claim is proved.

 \Box