arXiv:2312.11058v1 [math.AG] 18 Dec 2023 [arXiv:2312.11058v1 \[math.AG\] 18 Dec 2023](http://arxiv.org/abs/2312.11058v1)

HALF-INTEGRAL LEVELS

YIFEI ZHAO

To James Tao

Abstract. We construct equivalences among four notions associated to a reductive group scheme G: factorization super central extensions of the loop group of G by \mathbb{G}_m subject to a condition on the commutator, factorization super line bundles on the affine Grassmannian of G, rigidified sections of a quotient of 2-truncated K-theory over the Zariski classifying stack of G, and combinatorial data defined by Brylinski and Deligne in a conjectural extension of their classification theorem.

CONTENTS

Date: December 19, 2023.

The project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 427320536 – SFB 1442, as well as under Germany's Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics–Geometry–Structure.

 $2\,$ $\,$ YIFEI ZHAO $\,$

"Is this what it feels like to end?" "I do not know, for this is not our end."

—Kindred

INTRODUCTION

Let $k((t))$ denote the field of formal Laurent series with coefficients in a field k. Tate [\[Tat68\]](#page-48-0) discovered a remarkable central extension of its group of units $k((t))$ ^x by k^x :

$$
1 \to k^{\times} \to \mathcal{G}_{\text{Tate}} \to k((t))^{\times} \to 1. \tag{0.1}
$$

The preimage of $a \in k((t))^\times$ in $\mathscr{G}_{\text{Tate}}$ consists of nonzero elements of the relative determinant line det(ak[[t]] | k[[t]]), where $k[[t]] \subset k((t))$ is the lattice of formal Taylor series.

If we think of $k((t))^{\times}$ as the $k((t))$ -points of the algebraic group \mathbb{G}_m , then the following question arises: for a reductive group G, what are the "natural" central extensions of $G(k(\mathbf{t}))$ by k^* ?

Brylinski and Deligne [\[BD01\]](#page-46-1) parametrized a large class of central extensions of $G(k(\ell))$] by k^* using K-theory, as follows. Denote by \underline{K}_2 the Zariski sheafification of the second algebraic K-group. Starting with a central extension on the big Zariski site of $Spec(k)$:

$$
1 \to \underline{\mathrm{K}}_2 \to \mathrm{E} \to \mathrm{G} \to 1,\tag{0.2}
$$

evaluating at $Spec(k((t)))$ and pushing out along the tame symbol $\underline{K}_2(k((t))) \to k^*$, we find a central extension of $G(k(\ell))$ by k^{\times} . They went on to give a complete classification of central extensions [\(0.2\)](#page-1-1), valid over any regular base scheme of finite type over a field [\[BD01,](#page-46-1) Theorem 7.2. However, no central extension of \mathbb{G}_m by K_2 produces Tate's central extension [\(0.1\)](#page-1-2). This led Brylinski and Deligne to pose [\[BD01,](#page-46-1) Questions 12.13(iii)]:

"For $V = k[[t]]$, not all natural central extensions by k^{\times} are captured by 12.8. [...] We expect that 'natural' central extensions of $G(K)$ by k^{\times} are attached to data as follows: a Weyl group and Galois group invariant integer-valued symmetric bilinear form [...]"

Our first goal is to find an enlargement of the groupoid of central extensions of G by K_2 and prove that it meets the expectation of Brylinski and Deligne. To this end, we introduce a Zariski sheaf of connective spectra $\underline{\mathbf{K}}^{\text{super}}_{[1,2]}$ and establish the following result.

Theorem A (Theorem [2.2.3\)](#page-18-0). Let X be a regular scheme of finite type over a field and G be a reductive group X-scheme equipped with a maximal torus T. The following Picard groupoids are canonically equivalent:

- (1) rigidified sections of $\underline{K}_{[1,2]}^{\text{super}}$ $\sup_{[1,2]}^{\text{super}}$ over the Zariski classifying stack BG;
- (2) triples $(b, \widetilde{\Lambda}, \varphi)$ defined in [\[BD01,](#page-46-1) Questions 12.13(iii)].

Given a $k((t))$ -point of X, it is straightforward to produce from a rigidified section of $\underline{K}^{\text{super}}_{[1,2]}$ over BG a central extension of $G(k(\mathcal{t}))$ by k^* . In fact, the result will carry a canonical $\mathbb{Z}/2$ -grading, hence a "super central extension". This includes Tate's central extension in the special case $G = \mathbb{G}_m$.

The second goal of this article is to prove that this passage from K-theory to super central extensions of the loop group is reversible if one remembers an additional piece of structure called "factorization".

As observed by Beilinson and Drinfeld [\[BD04,](#page-47-0) [Dri06\]](#page-47-1), the construction of $\mathscr{G}_{\text{Tate}}$ globalizes, over any smooth curve X, to a *factorization super central extension* of the formal loop group $\mathscr{L}\mathbb{G}_m$. Intuitively speaking, this additional structure describes the behavior of $\mathscr{G}_{\text{Tate}}$ as one formal loop on X "factorizes" into two. The following result shows that factorization super central extensions of the loop group $\mathscr{L}G$, subject to a "tame commutator" condition which is automatic in characteristic zero, admit a parametrization parallel to Theorem [A.](#page-1-3)

Theorem B (Theorem [3.4.5\)](#page-31-0). Let X be a smooth curve over a field and G be a reductive group X-scheme. The following Picard groupoids are canonically equivalent:

- (1) factorization super central extensions of $\mathscr{L}G$ by \mathbb{G}_m with tame commutator;
- (1') factorization super line bundles over the affine Grassmannian Gr_G ;
- (2) triples $(b, \tilde{\Lambda}_+, \varphi)$ defined in [\[BD01,](#page-46-1) Questions 12.13(iii)] up to a "twist"—if G is equipped with a maximal torus.

Upon choosing a ϑ -characteristic, *i.e.* a square root $\omega^{1/2}$ of the canonical line bundle of the smooth curve X, the "twist" mentioned in (2) disappears. The Picard groupoids in Theorem [A](#page-1-3) then become canonically equivalent to those in Theorem [B,](#page-2-0) forming a commutative diagram:

$$
A(1) \xrightarrow{\cong} B(1)
$$
\n
$$
\downarrow^{\cong}
$$
\n
$$
B(1')
$$
\n
$$
\downarrow^{\cong}
$$
\n
$$
B(1')
$$
\n
$$
\downarrow^{\cong}
$$
\n
$$
A(2) \xrightarrow{\cong} B(2)
$$
\n(0.3)

In fact, the equivalences in [\(0.3\)](#page-2-1) are the "half-integral" generalizations of a family of equivalences which are valid without the choice of a ϑ -characteristic.

Corollary C (Corollary [3.4.7\)](#page-32-2). Let X be a smooth curve over a field and G be a reductive group X-scheme. The following Picard groupoids are canonically equivalent:

- (1) central extensions of G by \underline{K}_2 on the big Zariski site of X;
- (2) factorization central extensions of $\mathscr{L}G$ by \mathbb{G}_m with tame commutator;
- (3) factorization line bundles over Gr_G ;
- (4) triples (Q, Λ, φ) in [\[BD01,](#page-46-1) Theorem 7.2]—if G is equipped with a maximal torus.

This corollary already improves the current state of knowledge. Indeed, an equivalence between the Picard groupoids (1) and (3) was conjectured in Gaitsgory–Lysenko [\[GL18\]](#page-47-2) and established in [\[Gai20,](#page-47-3) [TZ21\]](#page-48-1) under the additional assumptions that G is split and a certain integer N_G is invertible in the ground field. The equivalence supplied by [C](#page-2-2)orollary C is valid for any reductive group X-scheme.

In the literature on covering groups in the equicharacteristic setting, the existence of factorization (super) line bundles over the affine Grassmannian Gr_{G} with favorable prop-erties is sometimes stated as an assumption, see e.g. [\[Lys16,](#page-48-2) [Lys17\]](#page-48-3) and [\[Laf18,](#page-48-4) §14]. The combination of Theorems [A](#page-1-3) and [B](#page-2-0) produces them unconditionally.

It is worth mentioning that Theorem [B](#page-2-0) is nontrivial already for $G = \mathbb{G}_m$. Indeed, fibers of the affine Grassmannian $\text{Gr}_{\mathbb{G}_m}$ over geometric points of X are highly nonreduced formal schemes. The groupoid of (super) line bundles over $\text{Gr}_{\mathbb{G}_m}$ does not appear to have a clean

description, but the equivalence $(1') \approx (2)$ of Theorem [B](#page-2-0) shows that the factorization ones do. Moreover, the equivalence $(1) \cong (1')$ shows that factorization (super) line bundles over Gr_G have canonical multiplicative structures over $\mathscr{L}G$. Unless G is simply connected, this assertion is not an obvious consequence of existing results.

From a differential geometric perspective, one could trace the conceptual origin of Corollary [C](#page-2-2) to works on Chern–Simons theory. Indeed, Dijkgraaf and Witten [\[DW90\]](#page-47-4) first recognized that the quantization parameter, or integral "level", of Chern–Simons theory for a compact Lie group G is best understood as an element of the reduced cohomology group $H_e^4(BG, \mathbb{Z})$. Suitably categorified, such an element transgresses to a central extension of the loop group of G by $U(1)$. A recent theorem of Waldorf [\[Wal17\]](#page-48-5) showed that this transgression procedure is reversible if one remembers the "fusion" structure of the target.

In the algebraic context, $H_e^4(BG, \mathbb{Z})$ should be replaced by the reduced weight-2 motivic cohomology group of BG, which classifies central extensions of G by K_2 via the isomorphism of [\[EKLV98\]](#page-47-5) (see also [\[Gai20,](#page-47-3) Theorem 6.3.5]):

$$
H_e^4(BG, \mathbb{Z}_{mot}(2)) \xrightarrow{\simeq} H_e^2(BG, \underline{K}_2). \tag{0.4}
$$

Hence, a central extension of G by \underline{K}_2 can be thought of as the algebraic notion of an *integral* level and [C](#page-2-2)orollary C provides four equivalent descriptions of it.^{[1](#page-3-1)} The equivalence (1) \leq (2) of Corollary [C](#page-2-2) is a direct analogue of Waldorf's theorem.

With this understanding, we propose to encode the algebraic notion of a *half-integral* level by the Picard groupoids in [\(0.3\)](#page-2-1). In fact, Dijkgraaf and Witten [\[DW90,](#page-47-4) §5] already observed that on *spin* manifolds, formally dividing a class in $H_e^4(BG, \mathbb{Z})$ by 2 sometimes leads to physically meaningful quantities. To interpret these "half-integral characteristic classes" as rigidified sections of $\underline{\mathbf{K}}_{[1,2]}^{\text{super}}$ over BG, we note that the natural inclusion of abelian groups below has a 2-torsion cokernel:

$$
H_e^2(BG, \underline{K}_2) \subset \pi_0 \Gamma_e(BG, \underline{K}_{[1,2]}^{\text{super}}). \tag{0.5}
$$

In the example of Tate's central extension, we have the equality $2 \cdot [\text{Tate}] = [c_1]^2$, where $[c_1]$ denotes the first Chern class of the universal line bundle over $\mathbf{B}\mathbb{G}_m$, so $[c_1]^2$ generates the abelian group $H_e^2(B\mathbb{G}_m, \underline{K}_2)$. Another example is the "critical level", *i.e.* Beilinson and Drinfeld's Pfaffian [\[BD91,](#page-46-2) §4], representing half of $[c_2]$ of the adjoint bundle over BG. It is half-integral precisely when the half sum of positive roots $\check{\rho}$ is not an integral weight. Halfintegral levels in our sense give rise to super conformal blocks on spin curves, as predicted by [\[DW90\]](#page-47-4), although we do not attempt to fully develop this notion here.

Let us now explain the structure of this article and comment on the proofs.

Structure of the article. This article is divided into two parts which can be read independently. The first part proves Theorem [A](#page-1-3) and the second part proves Theorem [B.](#page-2-0)

In §[1,](#page-4-2) we define $\underline{K}_{[1,2]}^{\text{super}}$ using a small but essential amount of homotopy theory. Namely, it is set to be the cofiber of a morphism of Zariski sheaves of connective spectra:

$$
\text{Sq}: \text{B\underline{K}}_1 \to \underline{\text{K}}_{[1,2]}.\tag{0.6}
$$

Here, \underline{K}_1 and $\underline{K}_{[1,2]}$ are the Zariski sheafified truncations of the K-theory spectrum. We also explain how to integrate sections of $\underline{K}_{[1,2]}^{\text{super}}$ over a global spin curve.

In §[2,](#page-15-0) we prove Theorem [A.](#page-1-3) The proof combines [\[BD01,](#page-46-1) Theorem 7.2] with our description of $\underline{K}_{[1,2]}$ obtained in §[1.](#page-4-2)

¹This algebraic notion is naturally associated to the chiral Wess–Zumino–Witten (WZW) model. We also mention that Henriques [\[Hen17\]](#page-47-6) proposed a definition of integral levels for the chiral WZW model via vertex algebras, while our notion is more directly related to chiral algebras, see [\[BD04,](#page-47-0) [Roz21\]](#page-48-6).

In §[3,](#page-23-1) we formulate Theorem [B.](#page-2-0) To define the notion of "tame commutator", we make essential use of the Contou-Carrère symbol over the Ran space, as constructed in Campbell– Hayash [\[CH21\]](#page-47-7). One of the phenomena we observe here is that the condition of having "tame commutator" is automatic in characteristic zero. This fact turns out to be equivalent to a new universality statement for the Contou-Carrère symbol.

Corollary D (Corollary [3.3.9\)](#page-30-1). Let X be a smooth curve over a field k with char(k) = 0. Then any pairing $\mathscr{L} \mathbb{G}_m \otimes \mathscr{L} \mathbb{G}_m \to \mathbb{G}_m$ compatible with factorization is an integral power of $the Contou-Carr\`ere symbol.$

We deduce this corollary from a surprising theorem of Tao [\[Tao21a\]](#page-48-7), which asserts that the presheaf $\text{Gr}_{\mathbb{G}_m}$ over the Ran space is *reduced* in a suitable sense, provided char(k) = 0. The assertion of Corollary [D](#page-4-3) is false if $char(k) > 0$. We do not use it in the proof of Theorem [B,](#page-2-0) which is valid in arbitrary characteristics.

In §[4,](#page-32-0) we prove Theorem [B.](#page-2-0) Our strategy is to first construct functors among the Picard groupoids in Theorem [B:](#page-2-0)

$$
(1) \to (1') \to (2). \tag{0.7}
$$

Our previous work [\[TZ21\]](#page-48-1) shows that the second functor is fully faithful. Here, we prove that the composition [\(0.7\)](#page-4-4) is an equivalence by exploiting the group structure inherent in (1). In our approach, each of the equivalences (1) \cong (2), (1') \cong (2) is established using special cases of the other in iteration, so we do not obtain one without the other.

Finally, we mention a shortcoming of this article: the top horizontal functor appearing in (0.3) is defined *ad hoc* as the composition of the other functors. It should have a conceptually transparent description as a "transgression" along the space of formal loops:

$$
\int_{(\mathring{D},\omega^{1/2})} : \Gamma_e(BG, \underline{K}_{[1,2]}^{\text{super}}) \to \text{Hom}(\mathscr{L}G, \text{Pic}^{\text{super}}), \tag{0.8}
$$

as in the differential geometric context, but we are unable to find such a description. One difficulty seems to be that we do not understand the behavior of Zariski-sheafified K-groups over singular spaces such as $\mathscr{L}G$. An attempt at defining [\(0.8\)](#page-4-5) as a "transgression" in the integral case, *i.e.* for sections of $K_2[2]$ over BG, was made in Kapranov–Vasserot [\[KV07\]](#page-48-8), but it relies on [\[KV07,](#page-48-8) Proposition 4.2.1] which is false as stated. A different strategy was carried out in Gaitsgory [\[Gai20\]](#page-47-3), but it requires the hypothesis that N_G be invertible in the ground field, which we wish to avoid.

Acknowledgements. James Tao has made the most important contibution to all problems considered in this article. This includes not only his published works on this topic [\[Tao21a,](#page-48-7) [Tao21b,](#page-48-9) [TZ21\]](#page-48-1), but also numerous ideas communicated to me during our collaboration. It is with profound gratitude and humility that I dedicate this article to him.

I thank Michael Finkelberg for fruitful e-mail exchanges and for his interest in Theorem [B](#page-2-0) in relation to [\[BDF](#page-47-8)⁺22]. I thank João Lourenço for the proof of Lemma [4.3.4.](#page-38-0)

Finally, I thank Dennis Gaitsgory for initiating me into factorization structures and for teaching me many things along the way.

Part 1. K-theory

1.
$$
\underline{\mathrm{K}}_{[1,2]}^{\mathrm{super}}
$$

The main goal of this section is to introduce the Zariski sheaf of connective spectra $\underline{K}^{\text{super}}_{[1,2]}$. The first section §[1.1](#page-5-0) reviews necessary notions concerning algebraic K-theory. In §[1.2,](#page-6-0) we give a "hands-on" description of the truncation $\underline{K}_{[1,2]}$. Using this description, we are able

to define $\underline{K}_{[1,2]}^{\text{super}}$ in §[1.3.](#page-9-0) The material of §[1.4](#page-11-0) is not needed in the sequal: its goal is to show that sections of $\underline{K}_{[1,2]}^{\text{super}}$ can be integrated over a global spin curve relative to a regular base scheme S to yield a super line bundle over S.

1.1. Connective K-theory.

1.1.1. Let Spc denote the ∞ -category of spaces. It is a symmetric monoidal ∞ -category under the Cartesian product.

Write $\text{Mon}_{\mathbb{E}_{\infty}}(\text{Spc})$ for the ∞ -category of \mathbb{E}_{∞} -monoids in Spc. It contains a full subcategory $\mathrm{Grp}_{{\mathbb E}_{\infty}}(\mathrm{Spc})$ consisting of grouplike ${\mathbb E}_{\infty}$ -monoids. The forgetful functor $\mathrm{Grp}_{{\mathbb E}_{\infty}}(\mathrm{Spc})\to$ $\text{Mon}_{\mathbb{E}_{\infty}}(\text{Spc})$ admits a left adjoint, called group completion:

$$
\Omega B: \mathrm{Mon}_{\mathbb{E}_{\infty}}(\mathrm{Spc}) \to \mathrm{Grp}_{\mathbb{E}_{\infty}}(\mathrm{Spc}).\tag{1.1}
$$

Let Sptr denote the ∞ -category of spectra. We use homotopical grading and denote by $\text{Sptr}_{\geq 0}$ the full subcategory of connective spectra.

There is a canonical equivalence of ∞ -categories [\[Lur17,](#page-48-10) Remark 5.2.6.26]:

$$
Grp_{\mathbb{E}_{\infty}}(Spc) \cong Sptr_{\geq 0}.
$$
 (1.2)

We shall also use without explicit mention the equivalence between Picard groupoids and 1-truncated connective spectra.

1.1.2. Let R be a commutative ring. Denote by $Vect(R)$ the category of finitely generated projective R-modules and $Vect(R)^\cong$ its maximal subgroupoid. The operation of direct sum equips $Vect(R)[°]$ with a symmetric monoidal structure. Its image under (1.1) is by definition the *connective* K-theory $K(R)$ of R.

We shall view K(R) either as a grouplike \mathbb{E}_{∞} -monoid or as a connective spectrum, using the canonical equivalence [\(1.2\)](#page-5-2).

Note that the unit of the adjunction between [\(1.1\)](#page-5-1) and the forgetful functor supplies a morphism of \mathbb{E}_{∞} -monoids:

$$
Vect(R)^{\cong} \to K(R), \quad \mathcal{E} \mapsto [\mathcal{E}]. \tag{1.3}
$$

1.1.3. We equip $\text{Mon}_{\mathbb{E}_{\infty}}(\text{Spc})$ and $\text{Gr}_{\mathbb{E}_{\infty}}(\text{Spc})$ with the canonical symmetric monoidal structure of [\[GGN15,](#page-47-9) Theorem 5.1]. With respect to these symmetric monoidal structures, [\(1.1\)](#page-5-1) is symmetric monoidal. Hence it lifts to a functor of \mathbb{E}_{∞} -monoids:

$$
\Omega B: \mathrm{Mon}_{\mathbb{E}_{\infty}}(\mathrm{Mon}_{\mathbb{E}_{\infty}}(\mathrm{Spc})) \to \mathrm{Mon}_{\mathbb{E}_{\infty}}(\mathrm{Grp}_{\mathbb{E}_{\infty}}(\mathrm{Spc})). \tag{1.4}
$$

The right adjoint of [\(1.1\)](#page-5-1), being lax symmetric monoidal, also lifts to a functor of \mathbb{E}_{∞} monoids and supplies the right adjoint of [\(1.4\)](#page-5-3), see [\[GGN15,](#page-47-9) Lemma 3.6].

The operation of tensor product upgrades $Vect(R)^{\cong}$ into an \mathbb{E}_{∞} -monoid in $Mon_{\mathbb{E}_{\infty}}(Spc)$. Thus K(R) acquires an \mathbb{E}_{∞} -monoid structure in $\text{Grp}_{\mathbb{E}_{\infty}}(\text{Spc})$ (*i.e.* K(R) is a connective \mathbb{E}_{∞} -spectrum) such that the unit [\(1.3\)](#page-5-4) is symmetric monoidal.

Remark 1.1.4. Informally, the symmetric monoidal structure on (1.3) says that for each pair of objects $\mathscr{E}_1, \mathscr{E}_2 \in \text{Vect}(\mathbb{R})^{\cong}$, $[\mathscr{E}_1 \otimes \mathscr{E}_2]$ is canonically equivalent to $[\mathscr{E}_1] \cdot [\mathscr{E}_2]$, together with the homotopy coherence data.

1.1.5. For any integer a, we write $K_{\geq a}(R)$ (resp. $K_{\leq a}(R)$) for the truncation $\tau_{\geq a}K(R)$ (resp. $\tau_{\leq a}K(R)$). For a pair of integers $a \leq b$, we write $K_{[a,b]} := \tau_{\geq a} \tau_{\leq b}K(R)$. We also use $K_a(R)$ to denote $\Omega^a K_{[a,a]}(R) \cong \pi_a K(R)$.

The association $S = Spec(R) \rightarrow K(R)$ defines a presheaf K of connective \mathbb{E}_{∞} -spectra on the category of affine schemes. Let \underline{K} denote its sheafification in the Zariski topology.

Zariski sheafification of the truncated presheaves above define $\underline{K}_{\geq a}$, $\underline{K}_{\leq a}$, $\underline{K}_{[a,b]}$, and \underline{K}_{a} . Since sheafification is t-exact, the forgetful functor from presheaves of spectra to sheaves of spectra is left t-exact. Hence $\underline{K}_{\leq a}$ is a-truncated as a presheaf of spectra, *i.e.* its value at any R has vanishing homotopy groups above degree a.

Remark 1.1.6. For example, the map sending $\mathscr{E} \in \text{Vect}(\mathbb{R})^{\cong}$ to its determinant line bundle $\det(\mathscr{E})$ induces an isomorphism of sheaves of Picard groupoids:

$$
\underline{K}_{[0,1]} \xrightarrow{\simeq} Pic^{\mathbb{Z}},
$$

where $Pic^{\mathbb{Z}}$ sends R to the Picard groupoid of \mathbb{Z} -graded line bundles on $Spec(R)$, see [\[BS17,](#page-47-10) Proposition 12.18].

1.2. The sheaf $K_{[1,2]}$.

1.2.1. The goal of this subsection is to give an explicit description of $\underline{K}_{[1,2]}$. More precisely, we consider the fiber sequence defined by truncation:

$$
B^2K_2(R) \to K_{[1,2]}(R) \to BK_1(R)
$$

for each ring R, which induces a fiber sequence of Zariski sheaves of connective spectra:

$$
B^2\underline{K}_2 \to \underline{K}_{[1,2]} \to B\underline{K}_1. \tag{1.5}
$$

Our description will be that of the fiber sequence [\(1.5\)](#page-6-1).

1.2.2. Denote by $Pic(R) \subset Vect(R)^{\cong}$ the full subcategory of line bundles over $Spec(R)$.

The map [\(1.3\)](#page-5-4) induces a morphism of the underlying *pointed spaces* Pic(R) $\rightarrow K(R)$ sending L to $[\mathscr{L}]$ – $[\mathscr{O}]$. Thus, we obtain a morphism of Zariski sheaves of pointed spaces, without changing the same notation:

$$
\text{Pic} \to \underline{\mathbf{K}}, \quad \mathcal{L} \to [\mathcal{L}] - [\mathcal{O}]. \tag{1.6}
$$

Since the class of $[\mathscr{L}] - [\mathscr{O}]$ in K₀(R) vanishes Zariski locally on Spec(R), the morphism [\(1.6\)](#page-6-2) factors through $\underline{K}_{\geq 1}$ and we may compose it with the truncation map $\underline{K}_{\geq 1} \to \underline{K}_{[1,2]}$ to obtain a map of Zariski sheaves of pointed spaces:

$$
s: \text{Pic} \to \underline{\mathbf{K}}_{[1,2]}.\tag{1.7}
$$

The description of $\underline{\mathbf{K}}_{[0,1]}$ via determinant (Remark [1.1.6\)](#page-6-3) shows that [\(1.7\)](#page-6-4) is a section of (1.5) on the underlying sheaves of pointed spaces, *i.e.* the composition of (1.7) with the truncation map $\underline{K}_{[1,2]} \rightarrow B \underline{K}_1$ is the canonical isomorphism Pic $\stackrel{\simeq}{\rightarrow} B \underline{K}_1$.

1.2.3. Let $\mathscr C$ be a site and A_1 , A_2 be sheaves of abelian groups over $\mathscr C$.

Consider the following two groupoids:

- (1) the groupoid of extensions A of $B(A_1)$ by $B^2(A_2)$ as sheaves of connective spectra, equipped with a section $s : B(A_1) \to A$ of the underlying sheaves of pointed spaces;
- (2) the (discrete) groupoid of anti-symmetric pairings $A_1 \otimes A_1 \rightarrow A_2$.

Let us construct a functor from (1) to (2) :

$$
\left\{\n\begin{array}{c}\n\text{B}(A_1) \\
\downarrow \text{ind} \\
B^2(A_2) \longrightarrow A \longrightarrow B(A_1)\n\end{array}\n\right\} \rightarrow\n\left\{\n\begin{array}{c}\n\text{anti-symmetric} \\
A_1 \otimes A_1 \rightarrow A_2\n\end{array}\n\right\}.
$$
\n(1.8)

Indeed, the section s defines a "cocycle" morphism of sheaves of spaces:

$$
B(A_1) \times B(A_1) \to B^2(A_2), \quad (x, y) \mapsto s(x + y) - s(x) - s(y). \tag{1.9}
$$

The morphism (1.9) is *bi-rigidified* in the following sense: it is equipped with trivializations along $B(A_1) \times e$ and $e \times B(A_1)$, which are isomorphic over $e \times e$.

To such a morphism, we may apply the loop space functor in the first, then the second factor, to obtain a map:

$$
A_1 \times A_1 \to A_2. \tag{1.10}
$$

Lemma 1.2.4. The map (1.10) is bilinear and anti-symmetric. The resulting functor (1.8) is an equivalence of groupoids.

1.2.5. The proof of Lemma [1.2.4](#page-7-1) proceeds by first giving an alternative definition of the functor [\(1.8\)](#page-6-6) which is evidently an equivalence, and then showing that it coincides with the construction above.

First, we observe that groupoid (1) is equivalent to the groupoid (1') of extensions A' of A_1 by $B(A_2)$ as sheaves of connective spectra, equipped with an E_1 -monoidal section $A_1 \rightarrow A'$. The equivalence is given by the functors Ω and B.

Put differently, the groupoid (1') consists of symmetric monoidal structures on the sheaf of associative monoids $B(A_2) \times A_1$ such that the inclusion and projection functors:

$$
B(A_2) \subset B(A_2) \times A_1 \to A_1
$$

are symmetric monoidal.

Such symmetric monoidal structures are in turn given by commutativity constraints on $B(A_2) \times A_1$ vanishing on $B(A_2)$, which are anti-symmetric bilinear pairings:

$$
A_1 \otimes A_1 \to A_2. \tag{1.11}
$$

Indeed, the commutativity constraint for two objects $x, y \in B(A_2) \times A_1$ is an isomorphism $x \otimes y \stackrel{\simeq}{\rightarrow} y \otimes x$ which depends only on the classes of x, y in A_1 . Such an isomorphism is the multiplication by a unique element of A_2 . The hexagon and inverse axioms translate to the bilinearity and anti-symmetry of the resulting map $A_1 \times A_1 \rightarrow A_2$.

The procedure above establishes an equivalence between the groupoid (1) and such pairings. Hence, it remains to prove that the pairing [\(1.11\)](#page-7-2) extracted from any object of the groupoid (1) concides with [\(1.10\)](#page-7-0). This follows from the observation in topology below.

1.2.6. Let M, N be \mathbb{E}_1 -monoids in Spc and let $s : M \to N$ be a morphism of pointed spaces. Then we may construct two maps of spaces:

$$
\Omega(M) \times \Omega(M) \xrightarrow[s_2]{s_1} \Omega^2(N). \tag{1.12}
$$

The map s_1 is given by applying Ω to the first, then the second factor, to the bi-rigidifed map of spaces:

$$
M \times M \to N, \quad (x, y) \mapsto s(x + y) - s(x) - s(y). \tag{1.13}
$$

The map s_2 uses the \mathbb{E}_1 -monoidal morphism $\Omega_s : \Omega(M) \to \Omega(N)$ induced from s and sends two loops $a, b \in \Omega(M)$ to the following loop in $\Omega(N)$:

$$
\begin{aligned} \mathbf{1} &\stackrel{\simeq}{\to} \Omega_s(a+b) - \left(\Omega_s(a) + \Omega_s(b)\right) \\ &\stackrel{\simeq}{\to} \Omega_s(b+a) - \left(\Omega_s(b) + \Omega_s(a)\right) \stackrel{\simeq}{\to} \mathbf{1}. \end{aligned} \tag{1.14}
$$

where the first and last isomorphisms are defined by the \mathbb{E}_1 -monoid structure on Ω_s and the middle one is defined by the braidings^{[2](#page-7-3)} on $\Omega(M)$ and $\Omega(N)$.

Claim: there is a homotopy equivalence $s_1 \stackrel{\simeq}{\rightarrow} s_2$.

²We use the term "braiding" to refer to the isomorphism $a \otimes b \stackrel{\simeq}{\to} b \otimes a$ in an E₂-monoid and the term "commutativity constraint" to refer to the same isomorphism in an \mathbb{E}_{∞} -monoid.

To prove the claim, we choose as our model for "spaces" topological spaces having the homotopy type of a CW complex, see [\[Lur18,](#page-48-11) 012Z] for its equivalence with the standard model using Kan complexes.

The monoidal operation on $\Omega(M)$ can be viewed as concatenation of loops and the braiding is given as follows. For two loops $a, b \in \Omega(M)$ we find a morphism:

$$
[0,1] \times [0,1] \to M, \quad (t_1, t_2) \mapsto a(t_1) + b(t_2), \tag{1.15}
$$

where the sum is the E_1 -monoid product on M. Then (1.15) can be viewed as a homotopy from its restriction to $([0,1] \times \{0\}) \cup (\{1\} \times [0,1])$ to its restriction to $(\{0\} \times [0,1]) \cup ([0,1] \times [0,1])$ {1}), exhibiting the braiding $a + b \stackrel{\simeq}{\rightarrow} b + a$ in $\Omega(M)$.

The same description holds for the braiding in $\Omega(N)$.

Now we come to the morphism s_1 . It carries a, b to the element of $\Omega^2(N)$ which is represented by a map $S^2 \rightarrow N$ fitting into the following diagram:

$$
[0,1] \times [0,1] \xrightarrow{(a,b)} M \times M
$$

$$
\downarrow \qquad \qquad \downarrow (1.13)
$$

$$
S^2 \longrightarrow N
$$
 (1.16)

where the left vertical map collapses the outer edges of the square.

Reading [\(1.16\)](#page-8-1) as a homotopy from its restriction to $([0,1] \times \{0\}) \cup (\{1\} \times [0,1])$ to its restriction to $({0} \times [0,1]) \cup ([0,1] \times {1})$ (both equivalent to the trivial loop in N), we see that it is precisely the loop [\(1.14\)](#page-7-5) defined by the braidings on $\Omega(M)$ and $\Omega(N)$.

The proof of the claim, and thus Lemma [1.2.4,](#page-7-1) is concluded.

1.2.7. We shall now use the equivalence of Lemma [1.2.4](#page-7-1) to classify the fiber sequence [\(1.5\)](#page-6-1) together with the distinguished section s [\(1.7\)](#page-6-4).

Namely, its image under the functor [\(1.8\)](#page-6-6) is an anti-symmetric form:

$$
\underline{\mathbf{K}}_1 \otimes \underline{\mathbf{K}}_1 \to \underline{\mathbf{K}}_2. \tag{1.17}
$$

Proposition 1.2.8. The map [\(1.17\)](#page-8-2) equals the product pairing $x, y \mapsto \{x, y\}$.

Proof. Let $\mathcal{L}_1, \mathcal{L}_2$ be sections of $B(K_1) \cong B\mathbb{G}_m$. The image of $(\mathcal{L}_1, \mathcal{L}_2)$ under the "cocycle" morphism, *i.e.* the special case of [\(1.9\)](#page-6-5) for $A_1 = \underline{K}_1$, $A_2 = \underline{K}_2$:

$$
B(\underline{K}_1) \times B(\underline{K}_1) \to B^2(\underline{K}_2)
$$
\n(1.18)

is the section in $B^2(\underline{K}_2)$ obtained from:

$$
([\mathcal{L}_1 \otimes \mathcal{L}_2] - [\mathcal{O}]) - ([\mathcal{L}_1] - [\mathcal{O}]) - ([\mathcal{L}_2] - [\mathcal{O}]) \tag{1.19}
$$

under the truncation map $\underline{K}_{\geq 2} \to B^2(\underline{K}_2)$.

Using the fact that [\(1.3\)](#page-5-4) is symmetric monoidal, the section [\(1.19\)](#page-8-3) is equivalent to ($[\mathscr{L}_1]$ − $[\mathscr{O}] \cdot ([\mathscr{L}_2] - [\mathscr{O}])$, where \cdot denotes multiplication on <u>K</u>.

The map Pic $\rightarrow \underline{K}_{\geq 1}$ induced from $\mathscr{L} \rightarrow [\mathscr{L}] - [\mathscr{O}]$ coincides with the map $B(\underline{K}_1) \rightarrow \underline{K}_{\geq 1}$ defined by truncation. Thus, (1.18) renders the following diagram commutative:

Here, the bottom horizontal arrow is the multiplicative structure on K and the vertical maps are defined by truncations. However, the only bi-rigidified morphism $B(\underline{K}_1) \times B(\underline{K}_1) \rightarrow B^2(K_2)$ rendering this diagram commutative is the product pairing $B^2(\underline{K}_2)$ rendering this diagram commutative is the product pairing.

Remark 1.2.9. Let us view the map defined by the product pairing $x, y \mapsto \{x, y\}$ as a morphism of Zariski sheaves of connective spectra:

$$
\operatorname{Pic} \otimes \operatorname{Pic} \to \operatorname{B}^2(\underline{\mathrm{K}}_2), \quad (\mathcal{L}_1, \mathcal{L}_2) \mapsto {\mathcal{L}_1, \mathcal{L}_2}.\tag{1.20}
$$

Then Proposition [1.2.8](#page-8-5) exihibits a canonical isomorphism of sections of $\underline{K}_{[1,2]}$:

$$
s(\mathcal{L}_1 \otimes \mathcal{L}_2) - s(\mathcal{L}_1) - s(\mathcal{L}_2) \cong \{\mathcal{L}_1, \mathcal{L}_2\},\tag{1.21}
$$

where $s: Pic \to \underline{K}_{[1,2]}$ is the morphism (1.7) of sheaves of pointed spaces. One may therefore view s as a "quadratic refinement" of the product pairing [\(1.20\)](#page-9-1).

1.2.10. Combining Proposition [1.2.8](#page-8-5) with the alternative construction of the pairing given in §[1.2.5,](#page-7-6) we obtain an explicit description of the loop space of (1.5) :

$$
B\underline{K}_2 \to \Omega(\underline{K}_{[1,2]}) \to \underline{K}_1. \tag{1.22}
$$

Namely, it splits as sheaves of \mathbb{E}_1 -monoids, given by [\(1.7\)](#page-6-4). The commutativity constraint on $\Omega(K_{[1,2]})$ is described by the anti-symmetric pairing:

$$
\underline{\mathbf{K}}_1 \otimes \underline{\mathbf{K}}_1 \to \underline{\mathbf{K}}_2, \quad x, y \mapsto \{x, y\}.
$$

1.2.11. Let R be a ring and \mathcal{M} be a line bundle over S := Spec(R). We shall temporarily work over the big Zariski site of Spec(R).

Multiplication by the object $\mathcal{M} \in K(R)$ induces a morphism of sheaves of connective spectra \cdot [\mathcal{M}] : <u>K</u> → <u>K</u>, hence a morphism:

$$
\cdot \left[\mathcal{M} \right] : \underline{\mathrm{K}}_{\left[1,2 \right]} \to \underline{\mathrm{K}}_{\left[1,2 \right]}.\tag{1.23}
$$

Since the image of $[\mathcal{M}]$ in $\Gamma(\text{Spec}(R),\underline{K}_0)$ is the multiplicative unit, multiplication by [M] induces the identity map on the homotopy sheaves \underline{K}_n for each $n \geq 0$. In particular, we obtain an automorphism of the triangles [\(1.5\)](#page-6-1):

B ²K² K[1,2] BK¹ B ²K² K[1,2] BK¹ id ⋅[M] id (1.24)

Claim: [\(1.23\)](#page-9-2) is the sum of the identity on $\underline{K}_{[1,2]}$ with the shearing map $\underline{BK}_1 \cong Pic \rightarrow$ $B^2\underline{K}_2$ defined by $\mathscr{L} \mapsto {\{\mathscr{L}, \mathscr{M}\}}$ (in the notation [\(1.20\)](#page-9-1)).

To see this, we may lift a section $\mathscr L$ of B \underline{K}_1 to $s(\mathscr L) = [\mathscr L] - [\mathscr O]$ of $\underline{K}_{[1,2]}$. The section $s(\mathcal{L})\cdot[\mathcal{M}]\cong[\mathcal{L}\otimes\mathcal{M}]-[\mathcal{M}]$ is then the sum of $s(\mathcal{L})$ with $\{\mathcal{L},\mathcal{M}\}$ by [\(1.21\)](#page-9-3).

1.3. The sheaf $K_{[1,2]}^{\text{super}}$ super
[1,2] **'**

1.3.1. Let Pic^{super} denote the Zariski sheaf of super *(i.e.* $\mathbb{Z}/2$ -graded) line bundles. As a sheaf of connective spectra, it coincides with the cofiber of the map:

$$
\underline{\mathbb{Z}} \to \text{Pic}^{\mathbb{Z}}, \quad n \mapsto (\mathcal{O}, 2n). \tag{1.25}
$$

We could suggestively denote Pic^{super} by $\underline{K}_{[0,1]}^{\text{super}}$, viewing [\(1.25\)](#page-9-4) as a morphism $\underline{K}_0 \to \underline{K}_{[0,1]}$ lifting the squaring map on K_0 .

The goal of this subsection is to introduce the Zariski sheaf $\underline{K}_{[1,2]}^{\text{super}}$ of connective spectra, defined as the cofiber of a map lifting the squaring map on $B\underline{K}_1$:

$$
\text{Sq}: \text{B\underline{K}}_1 \to \underline{\text{K}}_{[1,2]}.\tag{1.26}
$$

1.3.2. Let us first define [\(1.26\)](#page-10-0) as a morphism of sheaves of pointed spaces. To do so, we interpret $B\underline{K}_1$ as Pic and define [\(1.26\)](#page-10-0) by the formula:

$$
\text{Pic} \to \underline{\mathbf{K}}_{[1,2]}, \quad \mathcal{L} \mapsto [\mathcal{L}] - [\mathcal{L}^{-1}]. \tag{1.27}
$$

More precisely, the formula $\mathscr{L} \to [\mathscr{L}]-[\mathscr{L}^{-1}]$ defines a map $Pic(R) \to K(R)$, which induces [\(1.27\)](#page-10-1) upon Zariski sheafification and truncation as in §[1.2.2.](#page-6-7)

We argue that the structure of a morphism of connective spectra on (1.27) is unique, if it exists. Indeed, since [\(1.27\)](#page-10-1) has 1-connective source and target, it is equivalent to a morphism of sheaves of \mathbb{E}_1 -monoids:

$$
f: \mathbb{G}_m \to \Omega(\underline{\mathbf{K}}_{[1,2]}). \tag{1.28}
$$

Since $\Omega(\underline{K}_{[1,2]})$ is 1-truncated, an \mathbb{E}_{∞} -monoid structure on [\(1.28\)](#page-10-2) is equivalent to the condition that it preserves the commutativity constraint.

In particular, the following assertion involves no additional structure.

Proposition 1.3.3. The morphism of sheaves of pointed spaces [\(1.27\)](#page-10-1) lifts to a morphism of sheaves of connective spectra.

1.3.4. The proof of Proposition [1.3.3](#page-10-3) proceeds by explicitly identifying the morphism [\(1.28\)](#page-10-2) using the description of $\Omega(\underline{K}_{[1,2]})$ in §[1.2.10.](#page-9-5)

Namely, under the \mathbb{E}_1 -monoidal splitting $\Omega(\underline{K}_{1,2}) \cong B(\underline{K}_2) \times \underline{K}_1$, [\(1.28\)](#page-10-2) corresponds to two \mathbb{E}_1 -monoidal morphisms:

$$
f_1: \mathbb{G}_m \to \underline{\mathbf{K}}_1,
$$

$$
f_2: \mathbb{G}_m \to \mathbf{B}(\underline{\mathbf{K}}_2).
$$

Lemma 1.3.5. The following statements hold:

(1) f_1 is the squaring map $x \mapsto x^2$;

(2) f_2 is trivial as a morphism of sheaves of pointed spaces, and its \mathbb{E}_1 -monoid structure is defined by the automorphism of the trivial \underline{K}_2 -torsor:

$$
f_2(x) \otimes f_2(y) \stackrel{\simeq}{\to} f_2(xy), \quad 1 \mapsto 2 \cdot \{x, y\},\
$$

for each $x, y \in \mathbb{G}_m$.

Proof. As the section [\(1.7\)](#page-6-4) lifts the identity map on Pic $\cong B(\underline{K}_1)$, statement (1) follows from the isomorphism $\mathscr{L} \otimes (\mathscr{L}^{-1})^{-1} \cong \mathscr{L}^{\otimes 2}$.

For statement (2), we first apply the isomorphism [\(1.21\)](#page-9-3) to the pairs of sections $\mathscr{L}, \mathscr{L}^{-1} \in$ Pic and $\mathscr{L}, \mathscr{L} \in$ Pic to obtain isomorphisms in $\underline{K}_{[1,2]}$:

$$
s(\mathcal{L}) + s(\mathcal{L}^{-1}) \cong -\{\mathcal{L}, \mathcal{L}^{-1}\},
$$

$$
2 \cdot s(\mathcal{L}) \cong s(\mathcal{L}^{2}) - \{\mathcal{L}, \mathcal{L}\},
$$

where s denotes the seciton (1.7) .

Their difference yields an isomorphism in $\underline{K}_{[1,2]}$:

$$
[\mathcal{L}] - [\mathcal{L}^{-1}] \cong s(\mathcal{L}^2) - 2 \cdot {\mathcal{L}, \mathcal{L}}. \tag{1.29}
$$

In particular, this shows that f_2 is given by (−2) times the loop space of the self-pairing:

$$
\text{Pic} \to \text{B}^2(\underline{\text{K}}_2), \quad \mathcal{L} \to \{\mathcal{L}, \mathcal{L}\}.
$$
 (1.30)

According to [\[PR11,](#page-48-12) Theorem 2.5], the loop space of [\(1.30\)](#page-10-4) is the map $\mathbb{G}_m \to B(K_2)$ which is trivial as a morphism of sheaves of pointed spaces, with E_1 -monoid structure given, for any $x, y \in \mathbb{G}_m$, by the automorphism $1 \mapsto -\{x, y\}$ of the trivial $\underline{\mathrm{K}}_2$ -torsor. The desired conclusion follows.

Proof of Proposition [1.3.3.](#page-10-3) It suffices to prove that (1.28) preserves the commutativity constraint. In other words, given $x, y \in \mathbb{G}_m$, we must show that the following diagram of sections of $\Omega(\underline{K}_{[1,2]})$ commutes:

$$
f(x) \otimes f(y) \xrightarrow{\simeq} f(xy)
$$

\n
$$
\downarrow_{cf(x), f(y)} \qquad \qquad f(c_{x,y})
$$

\n
$$
f(y) \otimes f(x) \xrightarrow{\simeq} f(yx)
$$

\n(1.31)

Here, the horizontal morphisms are given by the \mathbb{E}_1 -monoid structure of f and the vertical morphisms are the commutativity constraints of $\Omega(K_{[1,2]})$, respectively \mathbb{G}_m (identity).

By Lemma [1.3.5,](#page-10-5) the commutativity of [\(1.31\)](#page-11-1) is equivalent to the following equality of sections of $\underline{\mathrm{K}}_2$:

$$
2 \cdot \{x, y\} = \{x^2, y^2\} + 2 \cdot \{y, x\}.
$$

This follows at once from the bilinearity and anti-symmetry of the pairing. \Box

Remark 1.3.6. From the proof of Proposition [1.3.3,](#page-10-3) we see that if we replace [\(1.27\)](#page-10-1) by the "obvious" lift of the squaring map $\mathscr{L} \mapsto 2 \cdot s(\mathscr{L})$, it would not define a morphism of sheaves of connective spectra.

1.3.7. Having constructed [\(1.27\)](#page-10-1), thus Sq [\(1.26\)](#page-10-0), as a morphism of sheaves of connective spectra, we define $\underline{K}_{[1,2]}^{\text{super}}$ to be the cofiber of Sq.

The following diagram summarizes four cofiber sequences of Zariski sheaves of connective spectra relevant for us:

$$
B(\underline{K}_1) \xrightarrow{\text{id}} B(\underline{K}_1)
$$
\n
$$
\downarrow s_q \qquad \downarrow 2
$$
\n
$$
B^2(\underline{K}_2) \longrightarrow \underline{K}_{[1,2]} \longrightarrow B(\underline{K}_1)
$$
\n
$$
\downarrow id \qquad \qquad \downarrow \qquad \qquad \downarrow
$$
\n
$$
B^2(\underline{K}_2) \longrightarrow \underline{K}_{[1,2]}^{\text{super}} \longrightarrow B(\underline{K}_1)/2
$$
\n
$$
(1.32)
$$

1.4. Integration on curves.

1.4.1. Given a quasi-compact and quasi-separated scheme S, we write $K(S)$ for the nonconnective K-theory spectrum of the stable ∞ -category Perf(S) [\[BGT13,](#page-47-11) §9]. The association $S \rightarrow K(S)$ is a Zariski sheaf of spectra [\[TT90,](#page-48-13) Theorem 8.1].

If S is regular, then the restriction of \bf{K} to the small Zariski site of S takes values in connective spectra, so $K(S)$ coincides with $\Gamma(S, K)$ [\[TT90,](#page-48-13) Proposition 6.8].

1.4.2. Let S be a regular affine scheme. Let $p : X_S \to S$ be a smooth, proper morphism of relative dimension 1 with connected geometric fibers.

The functor $\text{Perf}(X_S) \to \text{Perf}(S), \mathcal{E} \mapsto Rp_*\mathcal{E}$ induces a morphism of spectra $K(X_S) \to$ $K(S)$. By regularity, this amounts to a morphism of connective spectra:

$$
\Gamma(X_S, \underline{K}) \to \Gamma(S, \underline{K}).\tag{1.33}
$$

Lemma 1.4.3. For $n = 0, 1$, the morphism (1.33) fits into a commutative diagram:

$$
\Gamma(X_S, \underline{K}) \xrightarrow{(1.33)} \Gamma(S, \underline{K})
$$
\n
$$
\downarrow \qquad \qquad \downarrow \qquad (1.34)
$$
\n
$$
\Gamma(X_S, \underline{K}_{\leq n+1}) \to \Gamma(S, \underline{K}_{\leq n})
$$

where the vertical arrows are defined by truncation on \underline{K} .

Proof. We treat the case $n = 1$, as the case $n = 0$ is similar but simpler.

For $n = 1$, it suffices to trivialize the composition:

$$
\Gamma(X_S, \underline{K}_{\geq 3}) \longrightarrow \Gamma(X_S, \underline{K}) \stackrel{(1.33)}{\longrightarrow} \Gamma(S, \underline{K}) \longrightarrow \Gamma(S, \underline{K}_{\leq 1}),\tag{1.35}
$$

where the last arrow is defined by truncation on $\underline{\mathbf{K}}$.

Since $\underline{K}_{\leq 1} \cong Pic^{\mathbb{Z}}$ (Remark [1.1.6\)](#page-6-3) and S is regular, a section of $\Gamma(S, \underline{K}_{\leq 1})$ is trivialized once it is trivialized away from codimension ≥ 2 . Hence we may replace S by the spectrum of a discrete valuation ring R with field of fractions F.

In this case, X_S is Noetherian of Krull dimension 2, so for each $i \geq 0$, the complex $\Gamma(X_S, \underline{K}_i[i])$ is concentrated in cohomological degrees $\leq -i + 2$. Triviality of [\(1.35\)](#page-12-0) thus amounts to the *condition* that its induced map on H^{-1} below vanishes:

$$
H2(XS, K3) \rightarrow H0(S, K1) \cong R*.
$$
\n(1.36)

However, the formation of [\(1.35\)](#page-12-0) is of Zariski local nature on S, so [\(1.36\)](#page-12-1) fits into the commutative diagram:

$$
\begin{array}{ccc}H^2(X_S,\underline{K}_3)\,\rightarrow\,R^\times\\ \big\downarrow & & \big\downarrow\\ H^2(X_F,\underline{K}_3)\,\rightarrow\,F^\times\end{array}
$$

Here, $X_F \coloneqq X \times_S \text{Spec}(F)$ has Krull dimension 1, so $H^2(X_F, \underline{K}_3) = 0$ and (1.36) vanishes. \Box

1.4.4. We may now define a morphism:

$$
\int_{X_{\rm S}} : \Gamma(X_{\rm S}, \underline{\mathbf{K}}_{[1,2]}) \to \Gamma(\mathbf{S}, \mathrm{Pic}^{\mathbb{Z}}), \tag{1.37}
$$

to be the composition:

 $\Gamma(X_S, \underline{K}_{[1,2]}) \to \Gamma(X_S, \underline{K}_{\leq 2}) \to \Gamma(S, \underline{K}_{\leq 1}) \cong \Gamma(S, Pic^{\mathbb{Z}}).$

where the first morphism comes from the inclusion $\underline{K}_{[1,2]} \rightarrow \underline{K}_{\leq 2}$, and the second morphism is the bottom arrow of (1.34) for $n = 1$.

Comparing the cases $n = 1$ and $n = 0$ in [\(1.34\)](#page-12-2) shows that [\(1.37\)](#page-12-3) induces a morphism of fiber sequences:

$$
\Gamma(X_{\rm S}, B^2 \underline{K}_2) \rightarrow \Gamma(X_{\rm S}, \underline{K}_{[1,2]}) \rightarrow \Gamma(X_{\rm S}, B \underline{K}_1)
$$
\n
$$
\downarrow f_{X_{\rm S}} \qquad \qquad \downarrow f_{X_{\rm S}} \qquad \qquad \downarrow f_{X_{\rm S}} \qquad \qquad f_{X_{\rm S}} \qquad (1.38)
$$
\n
$$
\Gamma(S, \text{Pic}) \longrightarrow \Gamma(S, \text{Pic}^{\mathbb{Z}}) \longrightarrow \Gamma(S, \underline{\mathbb{Z}})
$$

Here, the rightmost vertical arrow has the following explict description: it associates to a line bundle over X_S its degree, viewed as a locally constant function over S.

Remark 1.4.5. If S is of finite type over a field, one may describe the first vertical functor in [\(1.38\)](#page-12-4) in terms of the Gersten resolution of \underline{K}_2 , see [\[Gai20,](#page-47-3) §2.4].

1.4.6. Let us now define the "super" variant of (1.37) , which requires a *spin structure* over X_S. From now on, we fix a square root $\omega^{1/2}$ of the relative canonical bundle $\omega_{X_S/S}$.

Define the morphism:

$$
\int_{(X_S,\omega^{1/2})} : \Gamma(X_S, \underline{K}_{[1,2]}) \to \Gamma(S, \text{Pic}^{\mathbb{Z}})
$$
\n(1.39)

to be the composition of [\(1.37\)](#page-12-3) with the multiplication $\cdot \left[\omega^{1/2}\right] : \underline{\mathrm{K}}_{[1,2]} \to \underline{\mathrm{K}}_{[1,2]}$ (*i.e.* the morphism [\(1.23\)](#page-9-2) for $\mathcal{M} \coloneqq \omega^{1/2}$).

The following observation shows that the $\omega^{1/2}$ -twisted integration morphism [\(1.39\)](#page-13-0) intertwines the squaring maps [\(1.25\)](#page-9-4) and [\(1.26\)](#page-10-0).

Lemma 1.4.7. The following diagram is canonically commutative:

$$
\Gamma(X_S, B\underline{K}_1) \xrightarrow{f_{X_S}} \Gamma(S, \underline{\mathbb{Z}}) \n\downarrow_{S_q} \qquad \qquad \downarrow_{n \to (\mathscr{O}, 2n)} \n\Gamma(X_S, \underline{K}_{[1,2]}) \xrightarrow{f_{(X_S, \omega^{1/2})}} \Gamma(S, \text{Pic}^{\mathbb{Z}})
$$
\n(1.40)

Proof. By construction, the lower circuit of [\(1.40\)](#page-13-1) sends a line bundle $\mathscr L$ over X_S to the Z-graded line bundle:

$$
\det(\mathrm{R}p_*(\mathscr{L} \otimes \omega^{1/2})) \otimes \det(\mathrm{R}p_*(\mathscr{L}^{-1} \otimes \omega^{1/2}))^{-1},
$$

This is the trivial line bundle by Grothendieck–Serre duality. It is placed in degree $2 \deg(\mathscr{L})$ by the Riemann–Roch formula.

1.4.8. Since X_S is regular, the Zariski cohomology group $H^2(X_S, \mathbb{G}_m)$ vanishes. Thus, taking cofibers of the vertical arrows in [\(1.40\)](#page-13-1) yields a morphism:

$$
\int_{(X_S,\omega^{1/2})} : \Gamma(X_S, \underline{K}_{[1,2]}^{\text{super}}) \to \Gamma(S, \text{Pic}^{\text{super}}). \tag{1.41}
$$

The morphisms of fiber sequences [\(1.24\)](#page-9-6) and [\(1.38\)](#page-12-4) induce a morphism of fiber sequences:

$$
\Gamma(X_S, B^2\underline{K}_2) \to \Gamma(X_S, \underline{K}_{[1,2]}^{super}) \to \Gamma(X_S, B(\underline{K}_1/2))
$$
\n
$$
\downarrow f_{X_S} \qquad \qquad \downarrow f_{(X_S, \omega^{1/2})} \qquad \qquad \downarrow f_{X_S}
$$
\n
$$
\Gamma(S, Pic) \longrightarrow \Gamma(S, Pic^{super}) \longrightarrow \Gamma(S, \underline{\mathbb{Z}}/2)
$$
\n
$$
(1.42)
$$

Here, the term $\Gamma(X_S, B(\underline{K}_1/2))$ is identified with the cofiber of the multiplication by 2 map on $\Gamma(X_S, B_{K_1})$, and the rightmost vertical arrow has the following description: it sends a line bundle over X_S to its degree mod 2.

1.4.9. We now explain how sections of $\underline{K}_{[1,2]}^{\text{super}}$ over the Zariski classifying stack of a split reductive group scheme define super conformal blocks, at least in the vacuum case.

Let $\mathcal{M}^{\text{spin}}$ denote the moduli stack of spin curves. Namely, an S-point of $\mathcal{M}^{\text{spin}}$ consists of a morphism $p: X_S \to S$ of smooth, proper morphism of relative dimension 1 with connected geometric fibers together with a square root $\omega^{1/2}$ of the relative canonical bundle.

Given an affine group scheme G, denote by BG the stack classifying Zariski locally trivial G-torsors.[3](#page-13-2)

³This is in accordance with the notation B used elsewhere in this section, but our BG is different from the usual stack classifying étale or fppf locally trivial G-torsors.

Denote by Bun_G the stack over $\mathscr{M}^{\text{spin}}$ whose S-points are triples $(X_S, \omega^{1/2}, P)$, where $(X_S, \omega^{1/2})$ is an S-point of $\mathcal{M}^{\text{spin}}$ and P is a G-bundle over X_S .

If G is split reductive, we shall define a functor:

$$
\Gamma(BG, \underline{K}_{[1,2]}^{\text{super}}) \to \Gamma(Bun_G, \text{Pic}^{\text{super}}). \tag{1.43}
$$

Given a section κ of $\underline{\mathbf{K}}_{[1,2]}^{\text{super}}$ over BG, we define the $\mathbb{Z}/2$ -graded quasi-coherent sheaf \mathbb{V}_{κ} over M^{spin} to be the pushforward along Bun_G $\rightarrow M^{\text{spin}}$ of the image of κ along [\(1.43\)](#page-14-0).

Construction of [\(1.43\)](#page-14-0). Consider an S-point $(X_S, \omega^{1/2}, P)$ of Bun_G where S is regular and affine. Etale locally over S , we may assume that P is Zariski locally trivial $[DSS5, Theorem$ 2], so it defines a morphism $P: X_S \rightarrow BG$.

Pulling back along P and applying [\(1.41\)](#page-13-3) yields a functor:

$$
\Gamma(\text{BG}, \underline{\mathbf{K}}^{\text{super}}_{[1,2]}) \to \Gamma(\text{S}, \text{Pic}^{\text{super}}). \tag{1.44}
$$

Since $Bun_G \to Spec(\mathbb{Z})$ is smooth, a super line bundle over Bun_G is equivalent to a compatible system of super line bundles over S, for all regular affine schemes S over BunG. Using functoriality of (1.44) in S, we obtain (1.43) .

Remark 1.4.10. Let us work over an algebraically closed field k.

If G is split simple and simply connected with a split maximal torus $T \subset G$, the Picard groupoid of sections of B²K₂ over BG rigidified along the base point $e : \text{Spec}(k) \to \text{BG}$ is discrete and isomorphic to the abelian group of Weyl-invariant quadratic forms on the cocharacter lattice Λ of T [\[BD01,](#page-46-1) Theorem 4.7].

This abelian group has a canonical generator: the Weyl-invariant quadratic form Q with $Q(\alpha) = 1$ at any short coroot α . To each integer κ , the quadratic form $\kappa \cdot Q$ thus defines a section of $B^2\underline{K}_2$ over BG.

The quasi-coherent sheaf \mathbb{V}_{κ} defined by this section, via the construction of §[1.4.9,](#page-13-4) is identified with the space of (vacuum) conformal blocks at level κ in the usual sense, see [\[BL94\]](#page-47-13). They are known to be finite locally free if $char(k) = 0$ [\[TUY89\]](#page-48-14).

We have not undertaken a serious investigation of \mathbb{V}_{κ} in the generality of §[1.4.9.](#page-13-4)

Remark 1.4.11. Let us note an analogue of [\(1.42\)](#page-13-5) for surfaces. Suppose that X is a proper smooth surface over an algebraically closed field k . Consider the composition:

$$
\int_{X} : \Gamma(X, B^{2}\underline{K}_{2}) \to H^{2}(X, \underline{K}_{2}) \cong CH^{2}(X) \xrightarrow{\deg} \mathbb{Z}.
$$
 (1.45)

Suppose that the dualizing sheaf $\omega_{X/k}$ admits a square root $\omega^{1/2}$. Claim: the morphism [\(1.45\)](#page-14-2) canonically extends to a morphism $\Gamma(X, \underline{K}_{[1,2]}^{\text{super}})$ $_{[1,2]}^{\text{super}}$ $\rightarrow \mathbb{Z}$.

This extension will be defined as an analogue of the morphism [\(1.41\)](#page-13-3) for surfaces. Namely, we note that [\(1.37\)](#page-12-3) has an analogue for surfaces: the map $\Gamma(X, \underline{K}_{[1,2]}) \to \mathbb{Z}$ induced from $\lbrack \mathcal{E} \rbrack \mapsto \chi(\mathrm{R}\Gamma(X,\mathcal{E})).$ To see that it factors through $\Gamma(X,\underline{K}_{[1,2]}^{\text{super}})$ $\sup_{[1,2]}^{\text{super}}$) when $\omega^{1/2}$ exists, we appeal to the equality:

$$
\chi(\mathrm{R}\Gamma(X,\mathscr{L}\otimes\omega^{1/2})) - \chi(\mathrm{R}\Gamma(X,\mathscr{L}^{-1}\otimes\omega^{1/2})) = 0,
$$
\n(1.46)

for every line bundle $\mathscr L$ over X, which follows from the Riemann–Roch formula:

$$
\chi(\mathrm{R}\Gamma(\mathrm{X},\mathscr{L}\otimes\omega^{1/2}))=\chi(\mathrm{R}\Gamma(\mathrm{X},\mathscr{O}))+\frac{1}{2}(\mathscr{L}\cdot\mathscr{L}-\omega^{1/2}\cdot\omega^{1/2}),
$$

where \cdot denotes the intersection pairing, *i.e.* the composition of (1.20) with (1.45) .

2. Brylinski–Deligne classification

In this section, we classify rigidified sections of $\underline{\mathbf{K}}_{[1,2]}^{\text{super}}$ over the Zariski classifying stack BG of a reductive group scheme G over a base scheme S, assumed regular and of finite type over a field. The main result is Theorem [2.2.3.](#page-18-0)

We begin in §[2.1](#page-15-1) with a classification of rigidified sections of $K_{[1,2]}$ over BG (Proposition [2.1.8\)](#page-16-0). Using tools developed in §[1.2,](#page-6-0) we reduce this result to the Brylinski–Deligne theorem [\[BD01,](#page-46-1) Theorem 7.2]. In §[2.2,](#page-17-0) we state the main result. The next subsection §[2.3](#page-19-0) is a technical interlude classifying central extensions of G by \mathbb{G}_m over an arbitrary base scheme. The results of §[2.1](#page-15-1) and §[2.3](#page-19-0) are combined in §[2.4](#page-21-0) to prove Theorem [2.2.3.](#page-18-0)

2.1. Classification: $\underline{\mathrm{K}}_{[1,2]}.$

2.1.1. Let S be a regular scheme of finite type over a field.

Let $G \rightarrow S$ be a reductive group scheme equipped with a maximal torus $T \subset G$. Cocharacters of T form an étale sheaf of abelian groups Λ over S.

Denote by $G_{\rm sc}$ the simply connected form of G. The preimage of T in $G_{\rm sc}$ is a maximal torus $T_{sc} \subset G_{sc}$, whose sheaf of cocharacters is denoted by Λ_{sc} . The algebraic fundamental group π_1 G may then be realized as $\Lambda/\Lambda_{\rm sc}$.

2.1.2. Denote by BG the stack of Zariski locally trivial G-torsor. Denote by $e : S \rightarrow BG$ the unit section. For a Zariski sheaf $\mathscr F$ of connective spectra, we write $\Gamma_e(BG,\mathscr F)$ for the fiber of the morphism:

$$
e^* : \Gamma(\text{BG}, \mathscr{F}) \to \Gamma(\text{S}, \mathscr{F}).
$$

We also denote by $\Gamma_e(BG,\mathscr{F})$ the presheaf over S whose section over an S₁-scheme is $\Gamma_e(\text{BG } \times_{\text{S}} \text{S}_1, \mathscr{F})$. It is a sheaf in the Zariski topology.

In this subsection, we describe $\underline{\Gamma}_e(BG, \underline{K}_{[1,2]})$ in terms of the combinatorics of G.

2.1.3. We first recall Brylinski and Deligne's description of $\underline{\Gamma}_{e}(\text{BG}, B^2 \underline{\text{K}}_2)$. Indeed, [\[BD01,](#page-46-1) Theorem 7.2] constructs a canonical equivalence of sheaves of Picard groupoids:

$$
\underline{\Gamma}_e(\text{BG}, \text{B}^2 \underline{\text{K}}_2) \xrightarrow{\simeq} \vartheta_\text{G}(\Lambda), \tag{2.1}
$$

where sections of $\vartheta_{\rm G}(\Lambda)$ are triples (Q, Λ, φ) defined below:

- (1) Q is a Weyl-invariant integral quadratic form on Λ ;
- (2) Λ is a central extension of Λ by \mathbb{G}_m , whose commutator pairing $\Lambda \otimes \Lambda \to \mathbb{G}_m$ equals $\lambda_1, \lambda_2 \mapsto (-1)^{b(\lambda_1, \lambda_2)}$ for $b(\lambda_1, \lambda_2) \coloneqq Q(\lambda_1 + \lambda_2) - Q(\lambda_1) - Q(\lambda_2);$
- (3) φ is an isomorphism between the restriction of $\tilde{\Lambda}$ to $\Lambda_{\rm sc}$ and the central extension induced from $Q_{\rm sc}$ (in the sense of Remark [2.1.4\)](#page-15-2), the restriction of Q to $\Lambda_{\rm sc}$.

The Picard groupoid structure on $\vartheta_G(\Lambda)$ is defined by sum in Q and Baer sum in Λ .

Remark 2.1.4. To be more explicit, [\[BD01,](#page-46-1) §3] first shows that $\underline{\Gamma}_e(\text{BT},\text{B}^2\underline{\text{K}}_2)$ is canonically equivalent to the sheaf of Picard groupoids $\vartheta(\Lambda)$ whose sections are pairs $(Q, \tilde{\Lambda})$, where Q is an integral quadratic form on Λ , and $\tilde{\Lambda}$ is as in (2).

Then [\[BD01,](#page-46-1) §4] shows that $\underline{\Gamma_e}(BG_{sc}, B^2\underline{K}_2)$ is the sheaf of discrete groupoids whose sections are Weyl-invariant integral quadratic forms on $\Lambda_{\rm sc}$. Restriction along T_{sc} ⊂ G_{sc} and applying the description of $\underline{\Gamma_e}(BT_{sc}, B^2 \underline{K}_2)$, we obtain a functor from Weyl-invariant quadratic forms on $\Lambda_{\rm sc}$ to $\vartheta(\Lambda_{\rm sc})$:

$$
Quad(\Lambda_{\rm sc})^{\rm W} \to \vartheta(\Lambda_{\rm sc}). \tag{2.2}
$$

In particular, any $Q_{sc} \in \text{Quad}(\Lambda_{sc})^W$ induces a central extension of Λ_{sc} by \mathbb{G}_m .

The functor from $\underline{\Gamma}_e(BG, B^2 \underline{K}_2)$ to $\vartheta_G(\Lambda)$ is given as follows. The pair $(Q, \tilde{\Lambda})$ is defined by its restriction along $T \subset G$, and the isomorphism φ comes from functoriality with respect to the commutative diagram:

$$
\begin{array}{ccc} T_{sc} & \subset & G_{sc} \\ \downarrow & & \downarrow \\ T & \subset & G \end{array}
$$

Remark 2.1.5. Both sides of [\(2.1\)](#page-15-3) are étale sheaves over S. Indeed, it is clear that $\vartheta_{\rm G}(\Lambda)$ satisfies étale descent. The étale descent of $\underline{\Gamma}_e(BG, B^2 \underline{K}_2)$ is established in [\[BD01,](#page-46-1) §2], logically prior to proving that [\(2.1\)](#page-15-3) is an equivalence.

2.1.6. Let us define an enlargement $\vartheta_{\rm G}^{\mathbb{Z}}(\Lambda)$ of $\vartheta_{\rm G}(\Lambda)$. Namely, a section of $\vartheta_{\rm G}^{\mathbb{Z}}(\Lambda)$ is a quadruple $(Q, \tilde{\Lambda}, \varphi, x)$ where $(Q, \tilde{\Lambda}, \varphi)$ is a section of $\vartheta_G(\Lambda)$ and:

(4)
$$
x : \Lambda \to \mathbb{Z}
$$
 is a character vanishing on Λ_{sc} (*i.e.* a character of $\pi_1 G$).

Therefore, as a sheaf of *pointed spaces*, $\vartheta_G^{\mathbb{Z}}(\Lambda)$ is the product $\vartheta_G(\Lambda) \times \underline{\text{Hom}}(\pi_1 \text{G}, \mathbb{Z})$. As a sheaf of Picard groupoids, we demand that it fits into a fiber sequence:

$$
\vartheta_{\mathcal{G}}(\Lambda) \to \vartheta_{\mathcal{G}}^{\mathbb{Z}}(\Lambda) \to \underline{\text{Hom}}(\pi_1 \mathcal{G}, \mathbb{Z}).
$$
\n(2.3)

Specifying the Picard groupoid structure on $\vartheta_{\rm G}^{\mathbb{Z}}(\Lambda)$ thus amounts to specifying a symmetric cocycle, *i.e.* a morphism of Picard groupoids:

$$
\underline{\text{Hom}}(\pi_1 G, \mathbb{Z}) \otimes \underline{\text{Hom}}(\pi_1 G, \mathbb{Z}) \to \vartheta_G(\Lambda), \tag{2.4}
$$

together with a null-homotopy of its precomposition with the anti-symmetrizer.

Construction of [\(2.4\)](#page-16-1). Given sections x_1, x_2 of $\underline{\text{Hom}}(\pi_1 \text{G}, \mathbb{Z})$, the morphism [\(2.4\)](#page-16-1) assigns to $x_1 \otimes x_2$ the triple $(Q, \tilde{\Lambda}, \varphi)$, where $Q(\lambda) = x_1(\lambda)x_2(\lambda)$, $\tilde{\Lambda}$ is the central extension defined by the cocycle $\lambda_1, \lambda_2 \mapsto (-1)^{x_1(\lambda_1)x_2(\lambda_2)}$, and φ is the identity automorphism of the trivial central extension of $\Lambda_{\rm sc}$ by \mathbb{G}_m .

In order to construct a null-homotopy of the image of $x_1 \otimes x_2 - x_2 \otimes x_1$, we need to trivialize the central extension of Λ by \mathbb{G}_m defined by the cocycle:

$$
\lambda_1, \lambda_2 \mapsto (-1)^{x_1(\lambda_1)x_2(\lambda_2) - x_2(\lambda_1)x_1(\lambda_2)}.
$$
\n
$$
(2.5)
$$

In other words, we need to find a map $q: \Lambda \to \mathbb{G}_m$ such that $q(\lambda_1 + \lambda_2)q(\lambda_1)^{-1}q(\lambda_2)^{-1}$ coincides with [\(2.5\)](#page-16-2). The desired map is set to be $q(\lambda) := (-1)^{x_1(\lambda)x_2(\lambda)}$.

Remark 2.1.7. By associating to each $\lambda \in \Lambda$ its fiber $\mathscr{L}^{\lambda} \subset \tilde{\Lambda}$ viewed as a \mathbb{G}_m -torsor, the central extension $\tilde{\Lambda}$ in a section $(Q, \tilde{\Lambda}, \varphi)$ of $\vartheta_G(\Lambda)$ can be viewed as a monoidal morphism $\Lambda \to \text{Pic}$ which preserves the commutativity constraint up to the factor $(-1)^b$.

Likewise, an object $(Q, \tilde{\Lambda}, \varphi, x)$ of $\vartheta_G^{\mathbb{Z}}(\Lambda)$ defines a monoidal morphism:

$$
\Lambda \to \mathrm{Pic}^{\mathbb{Z}}, \quad \lambda \mapsto (\mathscr{L}^{\lambda}, x(\lambda)),
$$

which preserves the commutativity constraint up to the factor $(-1)^{\tilde{b}}$ (viewed as a \mathbb{G}_m -valued bilinear form on Λ), where \tilde{b} is defined by:

$$
\lambda_1, \lambda_2 \mapsto b(\lambda_1, \lambda_2) + x(\lambda_1)x(\lambda_2). \tag{2.6}
$$

Proposition 2.1.8. There is a canonical equivalence of sheaves of Picard groupoids:

$$
\underline{\Gamma}_e(\text{BG}, \underline{\text{K}}_{[1,2]}) \xrightarrow{\simeq} \vartheta_G^{\mathbb{Z}}(\Lambda). \tag{2.7}
$$

It is related to the Brylinski–Deligne equivalence by a commutative diagram:

$$
\begin{array}{ccc}\n\underline{\Gamma}_e(BG, B^2 \underline{K}_2) & \subset & \underline{\Gamma}_e(BG, \underline{K}_{[1,2]}) \\
& \downarrow (2.1) & \downarrow (2.7) \\
\vartheta_G(\Lambda) & \subset & \vartheta_G^Z(\Lambda)\n\end{array}
$$

Proof. As a presheaf of pointed spaces, $\underline{\Gamma}_e(BG, \underline{K}_{[1,2]})$ is the direct product $\underline{\Gamma}_e(BG, B^2 \underline{K}_2) \times$ $\underline{\Gamma}_e(BG, B\underline{K}_1)$ thanks to the section [\(1.7\)](#page-6-4). In particular, $\underline{\Gamma}_e(BG, \underline{K}_{1,21})$ satisfies étale descent, see Remark [2.1.5.](#page-16-4)

The desired functor (2.7) is defined to be the Brylinski–Deligne equivalence (2.1) on the first factor and the canonical isomorphism:

$$
\underline{\Gamma}_e(BG, B\underline{K}_1) \xrightarrow{\simeq} \underline{\text{Hom}}(G, \underline{K}_1) \xrightarrow{\simeq} \underline{\text{Hom}}(\pi_1 G, \mathbb{Z})
$$
\n(2.8)

on the second factor.

It thus remains to lift this functor to one between Picard groupoids. We appeal to the description of $\underline{K}_{[1,2]}$ using the symmetric cocycle $B\underline{K}_1 \otimes B\underline{K}_1 \to B^2\underline{K}_2$ associated to the antisymmetric pairing $\{\cdot,\cdot\} : \underline{K}_1 \otimes \underline{K}_1 \to \underline{K}_2$ (Proposition [1.2.8\)](#page-8-5). Indeed, it suffices to construct an isomorphism between the $\underline{\Gamma_e}(\text{BG}, B^2\underline{K}_2)$ -valued pairing it induces on $\underline{\Gamma_e}(\text{BG}, B\underline{K}_1)$ and the pairing [\(2.4\)](#page-16-1):

$$
\underline{\Gamma}_e(BG, B\underline{K}_1) \otimes \underline{\Gamma}_e(BG, B\underline{K}_1) \xrightarrow{\{\cdot,\cdot\}} \underline{\Gamma}_e(BG, B^2\underline{K}_2)
$$
\n
$$
\downarrow (2.8) \qquad \qquad \downarrow (2.1)
$$
\n
$$
\underline{\text{Hom}}(\pi_1 G, \mathbb{Z}) \otimes \underline{\text{Hom}}(\pi_1 G, \mathbb{Z}) \xrightarrow{(2.4)} \vartheta_G(\Lambda)
$$
\n(2.9)

compatibly with null homotopies of their pre-composition with the anti-symmetrizer.

By definition of $\vartheta_G(\Lambda)$, it suffices to treat the case G = T as long as the isomorphism we construct is functorial in T.

In this case, any pair of characters x_1, x_2 of T defines under the top horizontal arrow of [\(2.9\)](#page-17-2) the central extension:

$$
1 \to \underline{K}_2 \to E \to T \to 1
$$

corresponding to the cocycle $T \otimes T \to \underline{K}_2$, $(t_1, t_2) \mapsto \{x_1(t_1), x_2(t_2)\}\.$ The null-homotopy the central extension defined by the cocycle $(t_1, t_2) \rightarrow \{x_1(t_1), x_2(t_2)\} - \{x_2(t_1), x_1(t_2)\}$ is exhibited by the map $T \to \underline{K}_2$, $t \mapsto \{x_1(t), x_2(t)\}$. These data correspond to the description of (2.4) (for G = T) under the equivalence of [\[BD01,](#page-46-1) Theorem 3.16].

2.2. Classification: $K_{[1,2]}^{\text{super}}$ super
[1,2] •

2.2.1. Let us define another enlargement $\vartheta_G^{\text{super}}(\Lambda)$ of $\vartheta_G(\Lambda)$ which fits into a fiber sequence of sheaves of Picard groupoids over S:

$$
\vartheta_{\mathrm{G}}(\Lambda) \to \vartheta_{\mathrm{G}}^{\mathrm{super}}(\Lambda) \to \underline{\mathrm{Hom}}(\pi_1 \mathrm{G}, \mathbb{Z}/2). \tag{2.10}
$$

Namely, an object of $\vartheta_G^{\text{super}}(\Lambda)$ is a triple $(b, \tilde{\Lambda}, \varphi)$ where:

- (1) b is a Weyl-invariant integral symmetric bilinear form on Λ , such that $b(\lambda, \lambda) \in 2\mathbb{Z}$ for any $\lambda \in \Lambda_{\rm sc}$;
- (2) $\tilde{\Lambda}$ is a central extension of Λ by \mathbb{G}_m , whose commutator pairing equals $\lambda_1, \lambda_2 \mapsto$ $(-1)^{b(\lambda_1,\lambda_2)+\epsilon(\lambda_1)\epsilon(\lambda_2)}$, where $\epsilon(\lambda) \coloneqq b(\lambda,\lambda) \mod 2$;
- (3) φ is an isomorphism between the restriction of $\tilde{\Lambda}$ to $\Lambda_{\rm sc}$ and the central extension induced by $Q_{\rm sc}$ as in §[2.1.3.](#page-15-4)

To define the Picard groupoid structure on $\vartheta_G^{\text{super}}(\Lambda)$, it is more natural to interpret $\tilde{\Lambda}$ as a monoidal morphism (see Remark [2.1.7\)](#page-16-5):

$$
\Lambda \to \text{Pic}^{\text{super}}, \quad \lambda \mapsto (\mathscr{L}^{\lambda}, \epsilon(\lambda)), \tag{2.11}
$$

which preserves the commutativity constraint up to the bilinear form $(-1)^b$. The Picard groupoid structure on $\vartheta_G^{\text{super}}(\Lambda)$ is induced from sum in b and the Picard groupoid structure of Pic super. In particular, it is *not* strictly commutative in general.

Let us construct the fiber sequence [\(2.10\)](#page-17-3). The inclusion of $\vartheta_G(\Lambda)$ in $\vartheta_G^{\text{super}}(\Lambda)$ sends (Q, $\tilde{\Lambda}, \varphi$) to the triple $(b, \tilde{\Lambda}, \varphi)$, where $b(\lambda_1, \lambda_2) := Q(\lambda_1 + \lambda_2) - Q(\lambda_1) - Q(\lambda_2)$. The second map $\vartheta_{\rm G}^{\rm super}(\Lambda) \to \text{Hom}(\pi_1\mathbf{G}, \mathbb{Z}/2)$ assigns to $(b,\tilde{\Lambda},\varphi)$ the homomorphism ϵ as in (2). Note that ϵ vanishes if and only if b comes from a quadratic form.

Remark 2.2.2. The sheaf of Picard groupoids $\vartheta_G^{\text{super}}(\Lambda)$ is introduced in [\[BD01,](#page-46-1) Questions 12.13(iii)]. For $G = T$ a torus, a variant of it has also appeared in [\[BD04,](#page-47-0) §3.10], where its sections are called ϑ -data.

Theorem 2.2.3. There is a canonical equivalence of sheaves of Picard groupoids:

$$
\underline{\Gamma}_e(\text{BG}, \underline{\mathbf{K}}_{[1,2]}^{\text{super}}) \xrightarrow{\simeq} \vartheta_G^{\text{super}}(\Lambda). \tag{2.12}
$$

It is related to the Brylinski–Deligne equivalence by a commutative diagram:

$$
\underline{\Gamma}_e(BG, B^2 \underline{K}_2) \subset \underline{\Gamma}_e(BG, \underline{K}_{[1,2]}^{\text{super}})
$$
\n
$$
\downarrow (2.1)
$$
\n
$$
\downarrow (2.12)
$$
\n
$$
\vartheta_G(\Lambda) \subset \vartheta_G^{\text{super}}(\Lambda)
$$
\n(2.13)

2.2.4. The proof of Theorem [2.2.3](#page-18-0) will occupy the remainder of this section. For now, we shall formulate a compatibility statement between the isomorphisms [\(2.7\)](#page-16-3) and [\(2.12\)](#page-18-1) (which will in fact be used to define [\(2.12\)](#page-18-1).)

To do so, we need to construct two morphisms of sheaves of Picard groupoids:

$$
Hom(\pi_1 G, \mathbb{Z}) \to \vartheta_G^{\mathbb{Z}}(\Lambda)
$$
\n(2.14)

$$
\vartheta_G^{\mathbb{Z}}(\Lambda) \to \vartheta_G^{\text{super}}(\Lambda). \tag{2.15}
$$

The morphism [\(2.14\)](#page-18-2) sends a character $x : \pi_1 \mathbb{G} \to \mathbb{Z}$ to the quadruple $(Q, \tilde{\Lambda}, \varphi, 2x)$ where $Q(\lambda) = -2x(\lambda)^2$, $\tilde{\Lambda}$ is the trivial central extension, and φ is the identity automorphism of the trivial central extension.

The morphism [\(2.15\)](#page-18-3) is the identity on the subgroupoid $\vartheta_G(\Lambda)$. To any character x in the additional factor Hom $(\pi_1 \mathbf{G}, \mathbb{Z})$, it assigns the triple $(b, \tilde{\Lambda}, \varphi)$ where $b(\lambda_1, \lambda_2) = x(\lambda_1)x(\lambda_2)$, Λ is the trivial central extension (*i.e.* the morphism [\(2.11\)](#page-18-4) is given by $\lambda \mapsto (\mathscr{O}, \epsilon(\lambda))$, and φ is the identity automorphism of the trivial central extension.

Lemma 2.2.5. The maps (2.14) , (2.15) thus defined are morphisms of Picard groupoids, and fit into a fiber sequence of such:

$$
Hom(\pi_1 G, \mathbb{Z}) \to \vartheta_G^{\mathbb{Z}}(\Lambda) \to \vartheta_G^{\text{super}}(\Lambda). \tag{2.16}
$$

Proof. We only verify that [\(2.16\)](#page-18-5) is indeed a fiber sequence. Let $(Q, \tilde{\Lambda}, \varphi, x)$ be an object in the fiber of [\(2.15\)](#page-18-3). Thus the induced symmetric form:

$$
\lambda_1, \lambda_2 \mapsto Q(\lambda_1 + \lambda_2) - Q(\lambda_1) - Q(\lambda_2) + x(\lambda_1)x(\lambda_2)
$$

must vanish. Setting $\lambda_1 = \lambda_2$, this implies that $x(\lambda) \in 2\mathbb{Z}$ for all $\lambda \in \Lambda$, so we may write $x = 2y$ for a character $y : \pi_1 \mathbf{G} \to \mathbb{Z}$ and there holds $\mathbf{Q}(\lambda) = -2y(\lambda)^2$. The fact that $(\mathbf{Q}, \tilde{\Lambda}, \varphi, x)$ lies

in the fiber also supplies us with a trivialization of $\tilde{\Lambda}$ compatible with φ . This yields an isomorphism between (Q, Λ, φ, x) and the image of y under [\(2.14\)](#page-18-2).

2.2.6. The compatibility statement asserts that [\(2.16\)](#page-18-5) coincides with the cofiber sequence defining $\underline{K}_{[1,2]}^{\text{super}}$ (see §[1.3.7\)](#page-11-3) evaluated at BG:

$$
\underline{\Gamma_e}(\text{BG}, \underline{\text{BK}}_1) \stackrel{\text{Sq}}{\to} \underline{\Gamma_e}(\text{BG}, \underline{\text{K}}_{[1,2]}) \to \underline{\Gamma_e}(\text{BG}, \underline{\text{K}}_{[1,2]}^{\text{super}})
$$
\n
$$
\downarrow \cong \qquad \qquad \downarrow (2.7) \qquad \qquad \downarrow (2.12)
$$
\n
$$
\text{Hom}(\pi_1 \text{G}, \mathbb{Z}) \xrightarrow{(2.14)} \vartheta_G^{\mathbb{Z}}(\Lambda) \xrightarrow{(2.15)} \vartheta_G^{\text{super}}(\Lambda)
$$
\n
$$
(2.17)
$$

The following statement can be verified without any knowledge of [\(2.12\)](#page-18-1).

Lemma 2.2.7. The left square in (2.17) is canonically commutative.

Proof. We use the expression (1.29) of the map Sq as the difference:

$$
B\underline{K}_1 \to \underline{K}_{[1,2]}, \quad \mathcal{L} \mapsto s(\mathcal{L}^2) - 2\{\mathcal{L}, \mathcal{L}\}.
$$
 (2.18)

Under the equivalence [\(2.7\)](#page-16-3), s corresponds to the natural inclusion of $Hom(\pi_1G, \mathbb{Z})$ in $\vartheta_{\rm G}^{\mathbb{Z}}(\Lambda)$, while the map $\mathscr{L} \mapsto {\{\mathscr{L},\mathscr{L}\}}$ corresponds to the restriction of (2.4) along the diagonal copy of $\text{Hom}(\pi_1\text{G},\mathbb{Z})$ (established in the proof of Proposition [2.1.8\)](#page-16-0). The map induced from [\(2.18\)](#page-19-2) upon taking $\underline{\Gamma}_e(BG, \cdot)$ is thus readily computed to be [\(2.14\)](#page-18-2).

2.3. Central extensions by K_1 .

2.3.1. In this subsection, we let S be an arbitrary base scheme and $G \rightarrow S$ be a reductive group scheme. Our goal is to classify central extensions of G by $K_1 \cong \mathbb{G}_m$.

When S is the spectrum of a field, this classification is obtained by Weissman [\[Wei11,](#page-48-15) Theorem 1.11]. We give a self-contained proof valid over any base scheme.

2.3.2. For a reductive group scheme $H \rightarrow S$, we write Rad(H) for the radical of H as defined in $[ABD⁺66, XXII, D^{\'et} function 4.3.6]$. Namely, it is the maximal torus of the center of H. The formation of Rad(H) is stable under base change and recovers the classical notion (maximal connected normal solvable subgroup) over a geometric point of S.

Given a central extension of a reductive group scheme H by \mathbb{G}_m (or any torus):

$$
1 \to \mathbb{G}_m \to \widetilde{H} \to H \to 1,\tag{2.19}
$$

we first observe that \widetilde{H} is representable by a reductive group scheme. Indeed, one checks directly that $\widetilde{H} \to S$ is smooth and its geometric fibers have vanishing unipotent radicals.

2.3.3. By functoriality of the algebraic fundamental group, we obtain a functor from the Picard groupoid of central extension of G by \mathbb{G}_m to that of extensions of π_1 G by Z as sheaves of abelian groups:

$$
\text{Hom}_{\mathbb{E}_1}(\mathcal{G}, \mathcal{B}\mathbb{G}_m) \to \text{Hom}_{\mathbb{Z}}(\pi_1 \mathcal{G}, \mathcal{B}\mathbb{Z}).\tag{2.20}
$$

Proposition 2.3.4. The functor (2.20) is an equivalence.

2.3.5. The Picard groupoids in (2.20) are of étale local nature on S, so we may assume the existence of a maximal torus $T \subset G$ in the proof of Proposition [2.3.4.](#page-19-4)

Since the G-conjugation extends along the map $G_{sc} \rightarrow G$, the quotient stack G/G_{sc} has a monoidal structure. As such, we have isomorphisms of monoidal stacks:

$$
G/G_{sc} \stackrel{\simeq}{\leftarrow} T/T_{sc} \stackrel{\simeq}{\rightarrow} \pi_1(G) \otimes \mathbb{G}_m. \tag{2.21}
$$

Here, the tensor product is understood in the derived sense and sheafified in the fppf, or equivalently the étale topology.

Proof of Proposition [2.3.4.](#page-19-4) In view of the isomorphisms (2.21) , it suffices to prove that the following two forgetful functors are equivalences:

$$
\text{Hom}_{\mathbb{Z}}(T/T_{\text{sc}}, B\mathbb{G}_m) \to \text{Hom}_{\mathbb{E}_1}(T/T_{\text{sc}}, B\mathbb{G}_m),\tag{2.22}
$$

$$
\mathrm{Hom}_{\mathbb{E}_1}(\mathrm{G}/\mathrm{G}_{\mathrm{sc}}, \mathrm{B}\mathbb{G}_m) \to \mathrm{Hom}_{\mathbb{E}_1}(\mathrm{G}, \mathrm{B}\mathbb{G}_m). \tag{2.23}
$$

Indeed, the left-hand-side of [\(2.22\)](#page-20-0) is identified with $\text{Hom}_{\mathbb{Z}}(\pi_1\text{G}, \text{BZ})$ by the vanishing of $\underline{\operatorname{Ext}}^1(-,\mathbb{G}_m)$ on the category of fppf sheaves of abelian groups.

Given a central extension of a reductive group scheme H by \mathbb{G}_m as in [\(2.19\)](#page-19-6), we have a short exact sequence of tori:

$$
1 \to \mathbb{G}_m \to \text{Rad}(\widetilde{H}) \to \text{Rad}(H) \to 1. \tag{2.24}
$$

Indeed, the fact that Rad(\overline{H}) \rightarrow Rad(H) is surjective can be checked on geometric fibers. Moreover, Rad (\widetilde{H}) contains \mathbb{G}_m since the latter is central, so the inclusion of the kernel $\text{Rad}(\widetilde{H}) \cap \mathbb{G}_m$ inside \mathbb{G}_m is an isomorphism.

We make two observations:

- (1) If H is a torus, then so is \widetilde{H} . This is because the map $Rad(\widetilde{H}) \rightarrow \widetilde{H}$ is an isomorphism by comparing (2.19) with (2.24) .
- (2) If H is semisimple, then we find an isomorphism $\mathbb{G}_m \stackrel{\simeq}{\to} \text{Rad}(\widetilde{H})$.

To prove that [\(2.22\)](#page-20-0) is an equivalence, it suffices to show that any central extension of a torus by \mathbb{G}_m is commutative. This follows from observation (1).

To prove that [\(2.23\)](#page-20-2) is an equivalence, we first write the left-hand-side as the groupoid of central extensions:

$$
1 \to \mathbb{G}_m \to \widetilde{G} \to G \to 1,\tag{2.25}
$$

equipped with a \widetilde{G} -equivariant splitting over $G_{\rm sc}$ for the adjoint action. Our task is to show that such a splitting exists uniquely.

To construct such a splitting, we may assume that G is simply connected in [\(2.25\)](#page-20-3). Let $\widetilde{G}_{der} \subset \widetilde{G}$ denote its derived subgroup. We claim that the composition:

$$
\widetilde{G}_{der} \subset \widetilde{G} \to G \tag{2.26}
$$

is an isomorphism.

It suffices to prove that (2.26) is a central isogeny, *i.e.* it is finite, flat, and surjective, with kernel contained in the center of \tilde{G}_{der} . The statement on the kernel is clear. The fact that [\(2.26\)](#page-20-4) is finite, finite, and surjective may be established smooth locally, so we base change along $\tilde{G} \rightarrow G$, where [\(2.26\)](#page-20-4) becomes the multiplication map:

$$
\widetilde{\mathbf{G}}_{\text{der}} \times \mathbb{G}_m \to \widetilde{\mathbf{G}}.
$$

However, by observation (2), this morphism is identified with the isogeny $\tilde{G}_{der} \times \text{Rad}(\tilde{G}) \rightarrow \tilde{G}$ of [\[ABD](#page-46-3)⁺66, XXII, 6.2.3].

The isomorphism (2.26) for G simply connected equips (2.25) with a section over $G_{\rm sc}$. It is unique since any two sections differ by a character $\mathrm{G}_\mathrm{sc} \to \mathbb{G}_m$ which is necessarily trivial. To see that this section is \tilde{G} -equivariant, it suffices to observe that the diagram:

is \widetilde{G} -equivariant, and any automorphism of $\widetilde{G} \times_G G_{\rm sc}$ preserves its derived subgroup. \Box

2.4. Proof of Theorem [2.2.3.](#page-18-0)

2.4.1. We return to the set-up of §[2.1.1.](#page-15-5) In particular, the base scheme S is assumed to be regular and of finite type over a field. In this subsection, we construct the equivalence [\(2.12\)](#page-18-1) and thereby prove Theorem [2.2.3.](#page-18-0)

We shall construct this equivalence in two stages: we first do it when π_1 G is torsionfree and satisfies a Galois cohomological condition. This step uses Proposition [2.1.8](#page-16-0) and Proposition [2.3.4.](#page-19-4) We then bootstrap the general case from this one, using the flasque resolution over general base due to González-Avilés [\[GA13\]](#page-47-14).

The fact that we have to play with Galois cohomology is because we do not know a priori that $\underline{\Gamma}_e(\text{BG}, \underline{\text{K}}_{[1,2]}^{\text{super}})$ $\binom{\text{super}}{[1,2]}$ satisfies étale descent.

2.4.2. Note that our hypothesis on S guarantees that every S-tori is isotrivial, *i.e.* split by a finite étale cover $[ABD^+66, X, Théorème 5.16]$. In particular, it makes sense for an S-torus to be quasi-trivial, see [\[CTS87,](#page-47-15) Definition 1.2].

Lemma 2.4.3. If π_1 G is the sheaf of cocharacters of a quasi-trivial torus, then both rows in [\(2.17\)](#page-19-1) are cofiber sequences of Zariski sheaves.

Proof. This assertion amounts to the Zariski local surjectivity of the two horizontal morphisms appearing in [\(2.17\)](#page-19-1):

$$
\begin{aligned} f_1&:\underline{\Gamma}_e\big(\mathrm{BG}, \underline{\mathrm{K}}_{[1,2]}\big) \to \underline{\Gamma}_e\big(\mathrm{BG}, \underline{\mathrm{K}}_{[1,2]}^{\mathrm{super}}\big),\\ f_2&:\vartheta^{\mathbb{Z}}_\mathrm{G}(\Lambda) \to \vartheta^{\mathrm{super}}_\mathrm{G}(\Lambda). \end{aligned}
$$

For f_2 , we note that comparing (2.3) with (2.10) leads to a Cartesian square:

$$
\vartheta_G^{\mathbb{Z}}(\Lambda) \longrightarrow \underline{\text{Hom}}(\pi_1 \mathbf{G}, \mathbb{Z})
$$
\n
$$
\downarrow f_2 \qquad \qquad \downarrow \text{mod } 2
$$
\n
$$
\vartheta_G^{\text{super}}(\Lambda) \rightarrow \underline{\text{Hom}}(\pi_1 \mathbf{G}, \mathbb{Z}/2)
$$
\n(2.27)

Claim: the "mod 2" morphism is surjective in the Zariski topology.

Indeed, Zariski locally on S, we may find a finite Galois cover $S_1 \rightarrow S$ which splits $\pi_1 G$. Denote by Γ the Galois group of S_1/S and M the Z-linear dual of the Γ-module associated to π_1 G at a geometric point of S. Then the problem amounts to the surjectivity of $M^{\Gamma} \rightarrow$ $(M/2)^{\Gamma}$, which follows from $H^{1}(\Gamma, M) = 0$ by quasi-triviality.

It follows that f_2 is also surjective in the Zariski topology.

For f_1 , the canonical maps in (1.32) induce a Cartesian square:

$$
\underline{\Gamma_e}(\text{BG}, \underline{\underline{K}}_{[1,2]}) \longrightarrow \underline{\Gamma_e}(\text{BG}, \underline{\underline{K}}_1)
$$
\n
$$
\downarrow f_1 \qquad \qquad \downarrow \text{mod } 2
$$
\n
$$
\underline{\Gamma_e}(\text{BG}, \underline{\underline{K}}_{[1,2]}^{\text{super}}) \longrightarrow \underline{\Gamma_e}(\text{BG}, \underline{\underline{K}}_1/2)
$$
\n(2.28)

Proposition [2.3.4](#page-19-4) implies that the "mod 2" morphism is identified with the one appearing in [\(2.27\)](#page-21-1). In particular, it is also surjective in the Zariski topology given the hypothesis on π_1 G. The same thus holds for f_1 .

2.4.4. Suppose that π_1 G is the sheaf of cocharacters of a quasi-trivial torus. By Lemma [2.2.7](#page-19-7) and Lemma [2.4.3,](#page-21-2) we may define a morphism fitting into (2.17) :

$$
\underline{\Gamma}_e(\text{BG}, \underline{\mathbf{K}}_{[1,2]}^{\text{super}}) \to \vartheta_G^{\text{super}}(\Lambda). \tag{2.29}
$$

It is an equivalence by Proposition [2.1.8.](#page-16-0)

2.4.5. For general G, we first introduce an auxiliary sheaf of Picard groupoids $\widetilde{\theta}_{G}^{super}(\Lambda)$, defined to be the fiber product:

$$
\begin{aligned}\n\widetilde{\vartheta}_{\mathcal{G}}^{\text{super}}(\Lambda) &\to \text{Quad}(\Lambda_{\text{sc}})^{\mathcal{W}} \\
&\downarrow \qquad \qquad \downarrow (2.2) \\
\vartheta^{\text{super}}(\Lambda) &\to \vartheta^{\text{super}}(\Lambda_{\text{sc}})\n\end{aligned} \tag{2.30}
$$

where the bottom horizontal map is defined by functoriality with respect to $\Lambda_{\rm sc} \to \Lambda$. Concretely, a section of $\widetilde{\theta}_{G}^{super}(\Lambda)$ is a triple $(b,\widetilde{\Lambda},\varphi)$ as in $\theta_{G}^{super}(\Lambda)$, but the Weyl-invariance on b is relaxed: it is only required to be Weyl-invariant over $\Lambda_{\rm sc}$.

Restrictions along T \subset G, G_{sc} \to G and applying the functor [\(2.29\)](#page-21-3) to T, G_{sc}, and T_{sc} produces a functor:

$$
\underline{\Gamma}_e(\text{BG}, \underline{\text{K}}_{[1,2]}^{\text{super}}) \to \widetilde{\vartheta}_{\text{G}}^{\text{super}}(\Lambda). \tag{2.31}
$$

2.4.6. Suppose that we have a central extension of reductive group S-schemes:

$$
1 \to T_1 \to \tilde{G} \to G \to 1,\tag{2.32}
$$

where T_1 is a torus with sheaf of cocharacters Λ_1 . Denote by \widetilde{T} the preimage of T in \widetilde{G} . It is a maximal torus with sheaf of cocharacters Λ .

By functoriality, we find two morphisms of presheaves of Picard groupoids:

$$
\underline{\Gamma}_e(BG, \underline{\mathbf{K}}^{\text{super}}_{[1,2]}) \to \lim_{[n]} \underline{\Gamma}_e(B(\widetilde{G} \times T_1^{\times n}), \underline{\mathbf{K}}^{\text{super}}_{[1,2]}),
$$
\n(2.33)

$$
\vartheta_{\mathcal{G}}^{\text{super}}(\Lambda) \to \lim_{[n]} \vartheta_{\widetilde{\mathcal{G}} \times T_{1}^{\times n}}^{\text{super}}(\widetilde{\Lambda} \oplus \Lambda_{1}^{\oplus n}), \tag{2.34}
$$

where the limits are taken over the simplicial category.

Lemma 2.4.7. The following statements hold:

- (1) the functor (2.33) is an equivalence;
- (2) the functor (2.34) is fully faithful.

Proof. Statement (1) follows from the fact that the induced map on Zariski classifying stacks $\widetilde{\mathrm{BG}} \to \mathrm{BG}$ is surjective in the Zariski topology.

To prove statement (2), we fit $\vartheta_G^{\text{super}}(\Lambda)$ into a fiber sequence of étale sheaves of Picard groupoids over S:

$$
\underline{\text{Hom}}_{\mathbb{Z}}(\Lambda, \text{B}\mathbb{G}_m) \to \vartheta_{\text{G}}^{\text{super}}(\Lambda) \to \Gamma^2(\check{\Lambda})^{\text{W}}.
$$
 (2.35)

Here, $\check{\Lambda}$ is the dual of $\Lambda,$ so $\Gamma^2(\check{\Lambda})^\mathrm{W}$ is the abelian group of Weyl-invariant symmetric bilinear forms on Λ . The second map in [\(2.35\)](#page-22-2) sends a triple $(b,\widetilde{\Lambda},\varphi)$ to b, so its fiber is precisely the Picard groupoid of symmetric monoidal morphisms $\Lambda \to B\mathbb{G}_m$.

The fully faithfulness will follow, if we know that the two outer terms in [\(2.35\)](#page-22-2) satisfy descent along $\Lambda \to \Lambda$. For $\underline{\text{Hom}}_{\mathbb{Z}}(\Lambda, \text{B}\mathbb{G}_m)$, this is because Λ is identified with colim $\text{Im}[\Lambda]$ $\Lambda_1^{\oplus n}$). For $\Gamma^2(\check{\Lambda})^W$, this is the elementary observation that a symmetric bilinear form on $\widetilde{\Lambda}$ descends to Λ if its restrictions to $\widetilde{\Lambda} \oplus \Lambda_1$ along the action and projection maps coincide. \Box

Remark 2.4.8. In fact, the functor [\(2.34\)](#page-22-1) is also an equivalence. This will be established in the course of the proof of Theorem [2.2.3](#page-18-0) below.

Proof of Theorem [2.2.3.](#page-18-0) The case where π_1 G is the sheaf of cocharacters of a quasi-trivial torus is already treated in §[2.4.4.](#page-21-4)

For general G, it remains to prove that the functor [\(2.31\)](#page-22-3) factors through an equivalence onto the full subgroupoid $\vartheta_{\mathcal{G}}^{\text{super}}(\Lambda)$.

To do so, we choose a central extension [\(2.32\)](#page-22-4) with the additional property that $\pi_1\tilde{G}$ is the sheaf of cocharacters of a quasi-trivial torus. Such central extensions exist, thanks to [\[GA13,](#page-47-14) Proposition 3.2].

Combining the equivalences for $\widetilde{G} \times T_1^{\times n}$ and Lemma [2.4.7,](#page-22-5) we obtain the following (solid) functors among Zariski sheaves of Picard groupoids:

$$
\underline{\Gamma_e}(\text{BG}, \underline{\mathbf{K}}^{\text{super}}_{[1,2]}) \xrightarrow{\simeq} \lim_{[n]} \underline{\Gamma_e}(\text{B}(\widetilde{G} \times \text{T}_1^{xn}), \underline{\mathbf{K}}^{\text{super}}_{[1,2]})
$$
\n
$$
\downarrow^{\simeq}
$$
\n
$$
\vartheta_G^{\text{super}}(\Lambda) \qquad \subset \qquad \lim_{[n]} \vartheta_{\widetilde{G} \times \text{T}_1^{xn}}^{\text{super}}(\widetilde{\Lambda} \oplus \Lambda_1^{\oplus n})
$$
\n(2.36)

Note that a symmetric bilinear form on Λ is Weyl-invariant if and only if its restriction to $\widetilde{\Lambda}$ is. Hence, the functor [\(2.31\)](#page-22-3) factors through the full subgroupoid $\vartheta_G^{\text{super}}(\Lambda)$, supplying the dashed arrow in [\(2.36\)](#page-23-2). It follows that all functors in (2.36) are equivalences. \Box

Corollary 2.4.9. Let G be a reductive group S-scheme. The Zariski sheaf of Picard groupoids $\underline{\Gamma}_e(\mathrm{BG},\underline{\mathrm{K}}^{\mathrm{super}}_{[1,2]}$ $\sum_{[1,2]}^{\text{super}}$) over S satisfies étale descent.

Proof. Working Zariski locally over S, we may assume that G admits a maximal torus T [\[ABD](#page-46-3)⁺66, XIV, Corollaire 3.20]. Let Λ denote its sheaf of cocharacters.

Theorem [2.2.3](#page-18-0) then implies that $\underline{\Gamma_e}(\text{BG}, \underline{K}_{[1,2]}^{\text{super}})$ $\binom{\text{super}}{[1,2]}$ is equivalent to $\vartheta_{\mathbf{G}}^{\text{super}}(\Lambda)$, which clearly satisfies étale descent. \square

2.4.10. We finish our study of $\underline{\Gamma}_e(BG, \underline{K}_{[1,2]}^{\text{super}})$ $\begin{bmatrix} \text{super} \\ \text{[1,2]} \end{bmatrix}$ by computing its homotopy sheaves. Let us assume that $T\subset G$ is a fixed maximal torus with sheaf of cocharacters $\Lambda.$

From the Cartesian square (2.30) , we obtain a long exact sequence of étale sheaves of abelian groups:

$$
\begin{aligned} 1\to \pi_1\widetilde{\vartheta}_{\mathrm{G}}^{\mathrm{super}}(\Lambda)\to \pi_1\vartheta^{\mathrm{super}}(\Lambda)\to \pi_1\vartheta^{\mathrm{super}}(\Lambda_{\mathrm{sc}})\\ \to \pi_0\widetilde{\vartheta}_{\mathrm{G}}^{\mathrm{super}}(\Lambda)\to \pi_0\vartheta^{\mathrm{super}}(\Lambda)\oplus\mathrm{Quad}(\Lambda_{\mathrm{sc}})^{\mathrm{W}}\to \pi_0\vartheta^{\mathrm{super}}(\Lambda_{\mathrm{sc}}). \end{aligned}
$$

The homotopy sheaves of $\vartheta^{\text{super}}(\Lambda)$ are easily computed: $\pi_1\vartheta^{\text{super}}(\Lambda)$ is isomorphic to $\underline{\text{Hom}}(\Lambda,\mathbb{G}_m)$, and $\pi_0\vartheta^{\text{super}}(\Lambda)$ is isomorphic to the sheaf of symmetric bilinear forms on Λ . Therefore, $\pi_1 \widetilde{\vartheta}_G^{\text{super}}(\Lambda)$ is isomorphic to $\underline{\text{Hom}}(\pi_1 G, \mathbb{G}_m)$, and $\pi_0 \widetilde{\vartheta}_G^{\text{super}}(\Lambda)$ is isomorphic to the sheaf of symmetric bilinear forms on Λ whose restriction to $\Lambda_{\rm sc}$ comes from a Weylinvariant quadratic form.

By definition of the full subgroupoid $\vartheta_G^{\text{super}}(\Lambda) \subset \widetilde{\vartheta}_G^{\text{super}}(\Lambda)$, we see that $b \in \pi_0 \widetilde{\vartheta}_G^{\text{super}}(\Lambda)$ belongs to $\pi_0 \vartheta_{\mathcal{G}}^{\text{super}}(\Lambda)$ if and only if it is Weyl-invariant.

Writing $\Gamma^2(\tilde{\Lambda})_{\rm sc}^W$ for the abelian sheaf of Weyl-invariant symmetric bilinear forms on Λ whose restriction to $\Lambda_{\rm sc}$ comes from a quadratic form, we obtain:

$$
\pi_i \underline{\Gamma}_e(BG, \underline{\mathbf{K}}^{\text{super}}_{[1,2]}) \cong \begin{cases} \Gamma^2(\check{\Lambda})_\text{sc}^\text{W} & i=0, \\ \underline{\text{Hom}}(\pi_1 G, \mathbb{G}_m) & i=1, \\ 0 & i \geq 2. \end{cases}
$$

Part 2. Loop groups

3. STATEMENTS

The goal of this section is to state the classification of factorization super central extensions of $\mathscr{L}G$: Theorem [3.4.5.](#page-31-0) The first two subsections §[3.1,](#page-24-0) §[3.2](#page-25-0) review the notions of factorization structure and loop groups. In $\S 3.3$, we use the Contou-Carrère symbol to define the notion of "tame commutator" and study its basic properties. In §[3.4,](#page-30-0) we state the classification theorem of factorization super central extensions of $\mathscr{L}G$ and briefly indicate the structure of its proof.

We work over a ground field k . Let X be a smooth curve over k .

3.1. Factorization.

3.1.1. Denote by Ran the presheaf whose S-points are nonempty finite subsets of $\text{Maps}(S, X)$. We shall write an S-point of Ran as $x^I = (x^i)_{i \in I}$, where I is a nonempty finite set.

Given an S-point x^I of Ran, we denote by Γ_{x^I} the sum of the graphs $\Gamma_{x^i} \subset S \times X$ over $i \in I$ as effective Cartier divisors. Let D_{x^I} be the completion of $S \times X$ along Γ_{x^I} and \tilde{D}_{x^I} be its open subscheme $D_{x^I} \backslash \Gamma_{x^I}$.

Two S-points x^I , x^J of Ran are called *disjoint* if $\Gamma_{x^I} \cap \Gamma_{x^J} = \emptyset$. Denote by $x^I \sqcup x^J$ the S-point of Ran given by their union.

Remark 3.1.2. For each nonempty finite set I, there is a tautological map $X^I \rightarrow \text{Ran}$, sending an S-point x^I of X^I to the associated finite subset of Maps(S, X). The presheaf Ran is identified with the colimit of presheaves:

$$
\underset{I}{\operatorname{colim}}(X^I)\xrightarrow{\simeq} \operatorname{Ran},
$$

indexed by the category of nonempty finite sets with surjections.

3.1.3. Let $\mathscr Y$ be a presheaf over Ran. Given an S-point x^I of Ran, we write $\mathscr Y_{x^I}$ for the base change of $\mathscr Y$ along x^{I} .

The presheaf $\mathscr Y$ is called *factorization* when we are supplied with a functorial system (in S) of isomorphisms for all disjoint pairs of S-points (x^I, x^J) of Ran:

$$
\varphi_{x^I, x^J}: \mathscr{Y}_{x^I \sqcup x^J} \xrightarrow{\simeq} \mathscr{Y}_{x^I} \times_S \mathscr{Y}_{x^J},\tag{3.1}
$$

satisfying the analogues of commutativity and associativity conditions. Namely, the following diagram commutes:

$$
\mathcal{Y}_{x^I \sqcup x^J} \xrightarrow{\varphi_{x^I, x^J}} \mathcal{Y}_{x^I} \times_S \mathcal{Y}_{x^J}
$$
\n
$$
\downarrow \cong \qquad \qquad \downarrow \cong
$$
\n
$$
\mathcal{Y}_{x^J \sqcup x^I} \xrightarrow{\varphi_{x^J, x^I}} \mathcal{Y}_{x^J} \times_S \mathcal{Y}_{x^I}
$$
\n
$$
(3.2)
$$

where the left vertical arrow comes from the equality $x^I \sqcup x^J = x^J \sqcup x^I$ as S-points of Ran and the right vertical arrow is the map swapping the two factors; the following diagram commutes for pairwise disjoint S-points $(x^{\text{I}}, x^{\text{J}}, x^{\text{K}})$ of Ran:

YxI⊔xJ⊔x^K Yx^I ×^S YxJ⊔x^K YxI⊔x^J ×^S Yx^K Yx^I ×^S Yx^J ×^S Yx^K ϕxI,xJ⊔x^K ϕxI⊔xJ,x^K id×ϕxJ,x^K ^ϕxI,x^J [×]id (3.3)

3.1.4. Let $\mathscr Y$ be a factorization presheaf such that $\mathscr Y_{x^I}$ satisfies fppf descent for each S-point x^{I} of Ran. (We do not impose fppf descent on $\mathscr Y$ because Ran itself does not satisfy étale descent, see [\[GL19,](#page-47-16) Warning 2.4.4].)

A factorization super line bundle over $\mathscr Y$ is a super line bundle $\mathscr L$ over $\mathscr Y$ equipped with functorial isomorphisms for all disjoint pairs of S-points (x^I, x^J) of Ran with respect to (3.1) :

$$
(\varphi_{x^I,x^J})^*(\mathcal{L}_{x^I} \boxtimes \mathcal{L}_{x^J}) \xrightarrow{\simeq} \mathcal{L}_{x^I \sqcup x^J},\tag{3.4}
$$

which are compatible with [\(3.2\)](#page-24-2) and [\(3.3\)](#page-24-3).

Let us spell out the compatibility with [\(3.2\)](#page-24-2). Denote by exch: $\mathscr{Y}_{x} \times_S \mathscr{Y}_{x} \to \mathscr{Y}_{x} \times_S \mathscr{Y}_{x}$ the map which exchanges the coordinates. The commutativity constraint of the Picard groupoid of super line bundles yields an isomorphism:

$$
\operatorname{exch}^*(\mathcal{L}_{x^J} \boxtimes \mathcal{L}_{x^I}) \xrightarrow{\simeq} \mathcal{L}_{x^I} \boxtimes \mathcal{L}_{x^J}.
$$
\n(3.5)

The compatbility states that the image of (3.5) under $(\varphi_{x^1,x^1})^*$, viewed as an isomorphism $(\varphi_{x^J},x^I)^*(\mathscr{L}_{x^J} \boxtimes \mathscr{L}_{x^I}) \stackrel{\simeq}{\rightarrow} (\varphi_{x^I},x^J)^*(\mathscr{L}_{x^I} \boxtimes \mathscr{L}_{x^J})$ by the commutativity of [\(3.2\)](#page-24-2), intertwines the isomorphisms [\(3.4\)](#page-25-2) attached to $(x^{\text{I}}, x^{\text{J}})$, respectively $(x^{\text{J}}, x^{\text{I}})$.

3.1.5. Let \mathcal{H} be a group factorization presheaf such that \mathcal{H}_{n} satisfies fppf descent for each S-point x^{I} of Ran.

A multiplicative super line bundle $\mathscr L$ over $\mathscr H$ is called *factorization* if it is equipped with a factorization structure which commutes with the multiplicative structure, *i.e.* (3.4) is an isomorphism of multiplicative line bundles over $\mathcal{H}_{x^I \sqcup x^J} \cong \mathcal{H}_{x^I} \times_S \mathcal{H}_{x^J}$.

Note that a multiplicative factorization super line bundle over $\mathscr H$ is equivalent to a super central extension of group presheaves over Ran:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \widetilde{\mathcal{H}} \to \mathcal{H} \to 1,\tag{3.6}
$$

equipped with a functorial homomorphism $\widetilde{\varphi}_{x^1,x^J}$ lifting the factorization isomorphism φ_{x^1,x^J} of $\mathscr H$ for each disjoint pair of S-points (x^I, x^J) of Ran:

$$
1 \to \mathbb{G}_{m,S} \times_S \mathbb{G}_{m,S} \to \widetilde{\mathcal{H}}_x \times_S \widetilde{\mathcal{H}}_x \to \mathcal{H}_x \times_S \mathcal{H}_x \to 1
$$

\n
$$
\downarrow (a,b) \to ab \qquad \qquad \downarrow \widetilde{\varphi}_{x^1,x^1} \qquad \qquad \downarrow \varphi_{x^1,x^1}
$$

\n
$$
1 \longrightarrow \mathbb{G}_{m,S} \longrightarrow \widetilde{\mathcal{H}}_{x^1 \sqcup x^1} \longrightarrow \mathcal{H}_{x^1 \sqcup x^1} \longrightarrow 1
$$

\n
$$
(3.7)
$$

which satisfies commutativity and associativity. The data (3.6) , (3.7) subject to these conditions are called a *factorization super central extension* of \mathscr{H} by $\mathbb{G}_{m, Ran}$. They form a Picard groupoid to be denoted by:

$$
\mathrm{Hom}_{\mathrm{fact}}(\mathscr{H},\mathrm{Pic}^{\mathrm{super}}).
$$

(We interpret them as homomorphisms $\mathscr{H} \to \text{Pic}^{\text{super}}$ compatible with factorization.)

Let us again be explicit about commutativity: (3.6) being a super central extension, each S-point (x^I, h^I) of $\widetilde{\mathscr{H}}$ carries a grading, viewed as a locally constant section of $\mathbb{Z}/2$ over S. Commutativity refers to the equality:

$$
\widetilde{\varphi}_{x^{\mathtt{I}},x^{\mathtt{J}}}(h^{\mathtt{I}},h^{\mathtt{J}})=(-1)^{\epsilon^{\mathtt{I}} \epsilon^{\mathtt{J}}}\widetilde{\varphi}_{x^{\mathtt{J}},x^{\mathtt{I}}}(h^{\mathtt{J}},h^{\mathtt{I}}),
$$

whenever h^{I} (resp. h^{J}) has grading ϵ^{I} (resp. ϵ^{J}).

3.2. Loop groups.

3.2.1. Let $Y \rightarrow X$ be an affine morphism of finite type.

Denote by $\mathscr{L}Y$ (resp. \mathscr{L}^+Y) the presheaf whose S-points are pairs (x^I, y^I) where x^I is an S-point of Ran and y^I is an X-morphism $\mathring{D}_{x^I} \to Y$ (resp. $D_{x^I} \to Y$). Note that \mathscr{L}^+Y is a closed subpresheaf of $\mathscr{L}Y$ and the structural morphism $\mathscr{L}Y \to \text{Ran}$ (resp. $\mathscr{L}^+Y \to \text{Ran}$) is indschematic (resp. schematic), see [\[KV04,](#page-48-16) 2.4-2.5].

Furthermore, $\mathscr{L}Y$ admits a canonical factorization structure. Indeed, for any disjoint pair of S-points $(x^{\mathrm{I}}, x^{\mathrm{J}})$ of Ran, there is a functorial isomorphism:

$$
\mathscr{L}_{x^I \sqcup x^J} Y \xrightarrow{\simeq} \mathscr{L}_{x^I} Y \times_S \mathscr{L}_{x^J} Y,
$$

induced from $\tilde{D}_{x^I\sqcup x^J} \cong \tilde{D}_{x^I}\sqcup \tilde{D}_{x^J}$, which is clearly commutative and associative. Analogously, \mathscr{L}^+ Y also admits a canonical factorization structure.

Since the association $Y \mapsto \mathscr{L}Y$ (resp. \mathscr{L}^+Y) preserves limits, it carries an affine group X-scheme G of finite type to a factorization group presheaf $\mathscr{L}G$ (resp. \mathscr{L}^+G) over Ran.

3.2.2. Let G be a smooth group X-scheme with connected geometric fibers. We also introduce the *affine Grassmannian* Gr_G as the presheaf whose S-points are triples (x^{I}, P, α) , where x^I is an S-point of X, P is a G-torsor over $S \times X$, and α is a trivialization of P over $S \times X\backslash \Gamma_{x}$. Then $Gr_G \to Ran$ is ind-schematic of ind-finite type; it is ind-proper when G is reductive [\[Zhu17,](#page-48-17) Theorem 3.1.3].

The factorization structure on Gr_G is defined by Beauville–Laszlo gluing [\[Zhu17,](#page-48-17) Theorem 3.2.1] and the canonical map $\mathscr{L}G \rightarrow Gr_G$ realizes the latter as the quotient $\mathscr{L}G/\mathscr{L}^+G$ in the étale topology [\[Zhu17,](#page-48-17) Proposition $3.1.9$].

3.2.3. For later purposes, we give a convenient description of $\mathscr{L}^+G \rightarrow \text{Ran}$ as an inverse limit of smooth affine group schemes relative to Ran.

Consider an S-point x^I of Ran. The morphism $\Gamma_{x^I} \to S$ is finite locally free. Denote by R_ΓG the Weil restriction along $\Gamma_{x^I} \to S$ of G (pulled back along $\Gamma_{x^I} \subset S \times X \twoheadrightarrow X$.) Then RΓG is representable by a smooth affine group S-scheme [\[BLR90,](#page-47-17) §7.6]. The evaluation map defines a short exact sequence:

$$
1 \to \mathcal{L}_{x}^{\geq 1} \mathbf{G} \to \mathcal{L}_{x}^{+} \mathbf{G} \to \mathbf{R}_{\Gamma} \mathbf{G} \to 1. \tag{3.8}
$$

More generally, we let $\Gamma_{x^I}^{(n)}$ (for $n \geq 0$) denote the *n*th order infinitesimal neighborhood of the closed immersion $\Gamma_{x^I} \subset S \times X$. Then $\Gamma_{x^I}^{(n)} \to S$ is finite locally free: writing $\mathscr I$ for the ideal sheaf defining Γ_{x^I} , we see that each $\mathscr{I}^n/\mathscr{I}^{n+1}$ is locally isomorphic to $\mathscr{O}_{S\times X}/\mathscr{I}$ as an \mathscr{O}_S -module. Let $R_{\Gamma^{(n)}} G$ be the Weil restrction along $\Gamma^{(n)}_{x^I} \to S$, which is again representable by a smooth affine group S-scheme. This gives us a limit presentation:

$$
\mathscr{L}_{x^I}^+G \xrightarrow{\simeq} \lim_n R_{\Gamma^{(n)}}G.
$$

Under the Tannakian formalism, the formula $\xi \mapsto 1 + \xi$ defines an isormorphism between the vector group S-scheme $\mathfrak{g} \otimes (\mathcal{I}^{n+1}/\mathcal{I}^{n+2})$ and the kernel of the evaluation map $R_{\Gamma^{(n+1)}} G \to$ $R_{\Gamma^{(n)}}G$. In particular, the group scheme $\mathscr{L}^{\geq 1}_{x^I}G$ in [\(3.8\)](#page-26-1) is an (infinite) iterated extension of vector group S-schemes.

3.3. Contou-Carrère.

3.3.1. For each integer $n \geq 1$, we shall define Tate central extension as a factorization super central extension:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \widetilde{\text{GL}}_n \to \mathscr{L}\text{GL}_n \to 1. \tag{3.9}
$$

Viewing $\widetilde{\mathrm{GL}}_n$ as a super line bundle over $\mathscr{L}\mathrm{GL}_n$, its fiber at an S-point $(x^{\mathrm{I}}, a^{\mathrm{I}})$ of $\mathscr{L}\mathrm{GL}_n$ is the super \mathscr{O}_S -module:

$$
\det (a^{\text{I}} \mathscr O^{\oplus n}_{{\rm D}_{{}_{\mathit{X}}}{\rm I}}\mid \mathscr O^{\oplus n}_{{\rm D}_{{}_{\mathit{X}}}{\rm I}}) \text{ with grading } \operatorname{rank} (a^{\text{I}} \mathscr O^{\oplus n}_{{\rm D}_{{}_{\mathit{X}}}{\rm I}}\mid \mathscr O^{\oplus n}_{{\rm D}_{{}_{\mathit{X}}}{\rm I}}) \text{ mod } 2,
$$

where det(L₁ | L₂) denotes the relative determinant of two lattices L₁, L₂ in the Tate \mathscr{O}_{S} module $\mathscr{O}_{\mathring{\mathcal{D}}}^{\hat{\oplus}n}$ $\mathbb{D}_{xI}^{\oplus n}$ and rank(L₁ | L₂) denotes their relative rank. (See [\[Dri06\]](#page-47-1) or [\[CH21,](#page-47-7) §3] for the definition of these notions).

The multiplicative structure of [\(3.9\)](#page-26-2) is defined by the canonical isomorphism:

$$
\det(a^Ib^I\mathscr O_{{\rm D}_{x^I}}^{\oplus n}\mid\mathscr O_{{\rm D}_{x^I}}^{\oplus n})\cong\det(a^I\mathscr O_{{\rm D}_{x^I}}^{\oplus n}\mid\mathscr O_{{\rm D}_{x^I}}^{\oplus n})\otimes\det(b^I\mathscr O_{{\rm D}_{x^I}}^{\oplus n}\mid\mathscr O_{{\rm D}_{x^I}}^{\oplus n}),
$$

for any S-points (x^I, a^I) and (x^I, b^I) of $\mathscr{L}GL_n$. The factorization isomorphism arises from the $\mathbb{Z}/2$ -graded multiplicativity of determinants with respect to direct sums:

$$
\det(a^I \mathcal{O}_{D_{x^I}}^{\oplus n} \oplus b^J \mathcal{O}_{D_{x^J}}^{\oplus n} | \mathcal{O}_{D_{x^I}}^{\oplus n} \oplus \mathcal{O}_{D_{x^J}}^{\oplus n}) \cong \det(a^I \mathcal{O}_{D_{x^I}}^{\oplus n} | \mathcal{O}_{D_{x^I}}^{\oplus n}) \otimes \det(b^J \mathcal{O}_{D_{x^J}}^{\oplus n} | \mathcal{O}_{D_{x^J}}^{\oplus n}),
$$

for S-points (x^I, a^I) , (x^J, b^J) of $\mathscr{L}GL_n$ with x^I, x^J disjoint.

3.3.2. Following [CH21, §4] , we define the *Contou-Carrère symbol* (or *tame symbol*) to be the commutator pairing of (3.9) for $n = 1$:

$$
\langle \cdot, \cdot \rangle : \mathscr{L} \mathbb{G}_m \otimes \mathscr{L} \mathbb{G}_m \to \mathbb{G}_{m, \text{Ran}}, \tag{3.10}
$$

Namely, $\langle \cdot, \cdot \rangle$ carries S-points $(x^{\text{I}}, a^{\text{I}})$, $(x^{\text{I}}, b^{\text{I}})$ of $\mathscr{L} \mathbb{G}_m$ to the element $(x^{\text{I}}, a^{\text{I}}\tilde{b}^{\text{I}}(\tilde{a}^{\text{I}})^{-1}(\tilde{b}^{\text{I}})^{-1})$ of $\mathbb{G}_{m,\text{Ran}}$, where \tilde{a}^{I} (resp. \tilde{b}^{I}) is a lift of a^{I} (resp. b^{I}) to $\widetilde{\mathbb{G}}_m$ which exists locally on S.

The pairing (3.10) is *factorization* in the following sense: given disjoint S-points x^{I} , x^{J} of Ran and lifts a^I , b^I (resp. a^J , b^J) of x^I (resp. x^J) to $\mathscr{L} \mathbb{G}_m$, there holds:

$$
\langle a^{\mathrm{I}}\sqcup a^{\mathrm{J}},b^{\mathrm{I}}\sqcup b^{\mathrm{J}}\rangle=\langle a^{\mathrm{I}},b^{\mathrm{I}}\rangle\langle a^{\mathrm{J}},b^{\mathrm{J}}\rangle.
$$

Furthermore, [\(3.10\)](#page-27-0) is perfect in the sense that its adjoint:

$$
\mathcal{L}\mathbb{G}_m \to \underline{\text{Hom}}(\mathcal{L}\mathbb{G}_m, \mathbb{G}_{m,\text{Ran}})
$$
\n(3.11)

is an isomorphism of factorization group presheaves [\[CH21,](#page-47-7) Corollary 5.4.1.1]. This pairing exhibits $\mathscr{L}^+\mathbb{G}_m$ as the Cartier dual of $\mathrm{Gr}_{\mathbb{G}_m}$ [\[CH21,](#page-47-7) Theorem 5.2.1].

3.3.3. More generally, let T be an X-torus with dual X-torus T, (3.11) induces an isomorphism between \mathscr{L} T and $\underline{\text{Hom}}(\mathscr{L} T, \mathbb{G}_{m,\text{Ran}})$.

In particular, for a pair of X-tori T_1 , T_2 with sheaves of cocharacters Λ_1 , Λ_2 , any bilinear form $b : \Lambda_1 \otimes \Lambda_2 \to \mathbb{Z}$ defines a factorization pairing:

$$
\langle \cdot, \cdot \rangle_b : \mathcal{L} \mathcal{T}_1 \otimes \mathcal{L} \mathcal{T}_2 \to \mathbb{G}_{m, \text{Ran}}, \tag{3.12}
$$

uniquely characterized by the property that its restriction along λ_1, λ_2 , viewed as homomorphisms from $\mathscr{L} \mathbb{G}_m$ to $\mathscr{L} \mathbb{T}_1$ (resp. $\mathscr{L} \mathbb{T}_2$), equals $b(\lambda_1, \lambda_2)(\cdot, \cdot)$.

Pairings $\mathscr{L} T_1 \otimes \mathscr{L} T_2 \to \mathbb{G}_{m, \text{Ran}}$ of the form $\langle \cdot, \cdot \rangle_b$ are called *tame*. Given morphisms $T'_1 \rightarrow T_1, T'_2 \rightarrow T_2$, a tame pairing $\mathscr{L} T_1 \otimes \mathscr{L} T_2 \rightarrow \mathbb{G}_{m, \text{Ran}}$ induces a tame pairing $\mathscr{L} T'_1 \otimes$ $\mathscr{L}T'_{2} \rightarrow \mathbb{G}_{m,\text{Ran}}$. The converse also holds for surjections of tori.

Lemma 3.3.4. Let $\langle \cdot, \cdot \rangle : \mathscr{L}T_1 \otimes \mathscr{L}T_2 \to \mathbb{G}_{m, \text{Ran}}$ be a factorization pairing. Given surjections of X-tori $T_1' \rightarrow T_1$, $T_2' \rightarrow T_2$, if the induced pairing $\mathscr{L} T_1' \otimes \mathscr{L} T_2' \rightarrow \mathbb{G}_{m, Ran}$ is tame, then so is $\langle \cdot, \cdot \rangle$.

3.3.5. Before proving Lemma [3.3.4,](#page-27-2) we shall make an observation.

Suppose that we are given X-tori T_1 and T_2 . *Claim:* all factorization morphisms of group presheaves over Ran below are trivial:

$$
\mathcal{L}^+T_1 \to \text{Gr}_{T_2};\tag{3.13}
$$

$$
Gr_{T_1} \to \mathcal{L}^+ T_2. \tag{3.14}
$$

For [\(3.13\)](#page-27-3), this is because $\mathscr{L}^+T_1 \to \text{Ran}$ is pro-smooth with connected geometric fibers, whereas $\text{Gr}_{T_2} \rightarrow \text{Ran}$ has formal geometric fibers. For [\(3.14\)](#page-27-4), this is because $\text{Gr}_{T_1} \rightarrow \text{Ran}$ is ind-proper, whereas $\mathscr{L}^+T_2 \to \text{Ran}$ is pro-affine. The combination of these two facts shows that any factorization bilinear pairing $\mathscr{L} T_1 \otimes \mathscr{L} T_2 \rightarrow \mathbb{G}_m$ induces, and is uniquely determined by a pairing $\mathscr{L}^+{\rm T}_1\otimes{\rm Gr}_{{\rm T}_2}\to\mathbb{G}_m.$

Proof of Lemma [3.3.4.](#page-27-2) Let \check{T}_2 (resp. \check{T}'_2) denote the X-torus dual to T_2 (resp. T'_2). By perfectness of the Contou-Carrère symbol, $\langle \cdot, \cdot \rangle$ is equivalent to a factorization morphism of group presheaves over Ran:

$$
\mathcal{L}T_1 \to \mathcal{L}\check{T}_2. \tag{3.15}
$$

We need to prove that for any pair of cocharacters λ_1 , λ_2 of T₁, T₂, the endomorphism φ of $\mathscr{L} \mathbb{G}_m$ defined by the composition:

$$
\mathscr{L}\mathbb{G}_m \stackrel{\lambda_1}{\to} \mathscr{L}\mathrm{T}_1 \stackrel{(3.15)}{\to} \mathscr{L}\mathrm{T}_2 \stackrel{\lambda_2}{\to} \mathscr{L}\mathbb{G}_m
$$

is given by $a \mapsto a^N$ for some integer N.

The hypothesis implies that this statement holds after composing with an endomorphism of $\mathscr{L} \mathbb{G}_m$ defined by *n*th power map $a \mapsto a^n$ for some integer $n \geq 1$.

By the observation of §[3.3.5,](#page-27-5) φ is uniquely determined by its restriction φ^+ to $\mathscr{L}^+\mathbb{G}_m$, whose image is also contained in $\mathscr{L}^+\mathbb{G}_m$. Note furthermore that for each S-point x^I of X^I , the restriction $\mathscr{L}_{x^I}^{\dagger} \mathbb{G}_m$ of $\mathscr{L}^{\dagger} \mathbb{G}_m$ is the extension of a group S-scheme of multiplicative type by an iterated extension of vector group S-schemes (§[3.2.3\)](#page-26-3).

In particular, φ^+ induces a homomorphism of short exact sequences:

$$
\begin{aligned} 0\to \mathscr{L}^{\geq 1}_{x^1}\mathbb{G}_m &\to \mathscr{L}^+_{x^1}\mathbb{G}_m \to \mathrm{R}_\Gamma\mathbb{G}_m \to 1\\ \downarrow_{\varphi_a^+} &\downarrow_{\varphi^+} &\downarrow_{\varphi_m^+}\\ 0\to \mathscr{L}^{\geq 1}_{x^1}\mathbb{G}_m &\to \mathscr{L}^+_{x^1}\mathbb{G}_m \to \mathrm{R}_\Gamma\mathbb{G}_m \to 1 \end{aligned}
$$

The fact that φ_m^+ is the N'th power map after composing with the *n*th power map shows that $n \mid N'$ and φ_m^+ is the Nth power map, for $N = N'/n$.

Since $\mathscr{L}_{x}^{\dagger} \mathbb{G}_m \to S$ is pro-smooth, it suffices to prove that φ^{\dagger} is the Nth power map on \bar{k} -points. In other words, given $f \in \bar{k}[[t]]^{\times}$ satisfying the equality:

$$
\varphi^+(f)^n = (f^N)^n \text{ in } \bar{k}[[t]]^{\times},
$$

we need to deduce the equality $\varphi^+(f) = f^N$. Setting $g = \varphi^+(f)/f^N$, we may write:

$$
g = 1 + \sum_{i \ge 1} a_i t^i \text{ in } \overline{k}[[t]]^{\times}.
$$

Claim: $g^n = 1$ implies $g = 1$. For char(\bar{k}) $\nmid n$, this holds because all *nth* roots of unity of $\bar{k}[[t]]$ are contained in \bar{k} . For char $(\bar{k}) | n$, this holds because the Frobenius is injective. \Box

Example 3.3.6. Suppose that k has characterstic $p > 0$. Let us define a factorization central extension whose commutator is not tame:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathscr{G} \to \mathscr{L}\mathbb{G}_m \to 1. \tag{3.16}
$$

Given a morphism Y \rightarrow S of k-presheaves, we write $\text{Fr}_{Y/S}: Y \rightarrow Y_{/S}^{(1)}$ for the pth power Frobenius of Y relative to S. Its formation is compatible with base change along S. Note that the presheaf $\mathscr{L} \mathbb{G}_m$ ⁽¹⁾ is canonically isomorphic to $\mathscr{L} \mathbb{G}_m$: an S-point of $\mathscr{L} \mathbb{G}_m$ ⁽¹⁾_{/R_i} /Ran is a pair (x^I, a) where x^I is an S-point of Ran and a is map $\mathring{D}_{\mathrm{Fr}^*_S(x^I)} \to \mathbb{G}_m$. However, $\mathring{D}_{Fr^*_{\mathcal{S}}(s^I)}$ is isomorphic to \mathring{D}_{x^I} since its formation depends only on the subset $|\Gamma_{x^I}|$ of $|S \times X|$. In particular, we may view $\text{Fr}_{\mathscr{L}\mathbb{G}_m/\text{Ran}}$ as an endomorphism of $\mathscr{L}\mathbb{G}_m$ over Ran.

The central extension [\(3.16\)](#page-28-1) is defined to be the presheaf of sets $\mathbb{G}_{m, Ran} \times_{Ran} \mathscr{L} \mathbb{G}_m$ whose group structure is defined by the cocycle:

$$
\mathscr{L}\mathbb{G}_m \otimes \mathscr{L}\mathbb{G}_m \to \mathbb{G}_{m,\text{Ran}}, \quad (a,b) \mapsto \langle \text{Fr}_{\mathscr{L}\mathbb{G}_m/\text{Ran}}(a),b \rangle,
$$

where $\langle \cdot, \cdot \rangle$ denotes the Contou-Carrère symbol. Since $\langle \cdot, \cdot \rangle$ is anti-symmetric, the commutator of [\(3.16\)](#page-28-1) is the pairing:

$$
\mathscr{L}\mathbb{G}_m \otimes \mathscr{L}\mathbb{G}_m \to \mathbb{G}_{m,\text{Ran}}, \quad (a,b) \mapsto \langle \text{Fr}_{\mathscr{L}\mathbb{G}_m/\text{Ran}}(a),b \rangle \langle a, \text{Fr}_{\mathscr{L}\mathbb{G}_m/\text{Ran}}(b) \rangle. \tag{3.17}
$$

Let us argue that this pairing is not tame over any geometric point $x : \text{Spec}(\bar{k}) \to X$. The choice of a uniformizer allows us to identify $\mathscr{L}_x \mathbb{G}_m$ with $\mathbb{G}_m((t))$. The morphism $\mathrm{Fr}_{\mathbb{G}_m((t))/\bar{k}}$ evaluates to the following map on R-points for any \bar{k} -algebra R:

$$
R((t))^{x} \to R((t))^{x}, \quad \sum_{n} a_{n} t^{n} \mapsto \sum_{n} (a_{n})^{p} t^{n}.
$$

The commutator [\(3.17\)](#page-29-0) is indeed $\langle \cdot, \cdot \rangle^{p+1}$ on \bar{k} -points. For a more general \bar{k} -algebra R, the Contou-Carrère pairing $\langle 1-a_1t, 1-b_{-1}t^{-1} \rangle$ equals $1-a_1b_{-1}$ for nilpotents $a_1, b_{-1} \in \mathbb{R}$ [\[APR04\]](#page-46-4). Taking R := $\bar{k}[\epsilon]/\epsilon^3$ with $a_1 := \epsilon x$, $b_{-1} := \epsilon y$ for $x, y \in \bar{k}^\times$ and equating the commutator $(1 - a_1^{\bar{p}}b_{-1})(1 - a_1b_{-1}^{\bar{p}})$ with $(1 - a_1b_{-1})^{p+1}$, we find $xy = 0$, which is impossible.

3.3.7. We now show that tameness is a positive characteristic phenomenon.

The assertion below relies on [\[Tao21a\]](#page-48-7) which uses the hypothesis char(k) = 0.

Proposition 3.3.8. Let T_1 , T_2 be a pair of X-tori. If char(k) = 0, then any factorization pairing $\mathscr{L} T_1 \otimes \mathscr{L} T_2 \rightarrow \mathbb{G}_{m, \text{Ran}}$ is tame.

Proof. Using the observations in §[3.3.5,](#page-27-5) it suffices to prove that any factorization pairing $\mathscr{L}^+\mathrm{T}_1\otimes\mathrm{Gr}_{\mathrm{T}_2}\to\mathbb{G}_{m,\mathrm{Ran}}$ is necessarily of the form $\langle\cdot,\cdot\rangle_b$ for some bilinear form $b:\Lambda_1\otimes\Lambda_2\to\mathbb{Z}$ (see §[3.3.3\)](#page-27-6).

Using the duality between Gr_{T_2} and $\mathscr{L}^+ \check{T}_2$ under the Contou-Carrère symbol, we reduce the statement to the special case $T_1 = T_2 = \mathbb{G}_m$.

For each I-tuple $\lambda^I = (\lambda^i)$ of integers, there is a closed immersion $\iota_{\lambda^I}: X^I \to \mathrm{Gr}_{\mathbb{G}_m, X^I}$ sending an S-point $x^I = (x^i)$ of X^I to the line bundle $\mathscr{O}(\sum_{i\in I}\Gamma_{x^i})$ over $S \times X$ equipped with its canonical trivialization of Γ_{x} . Consider the category of pairs (I, λ^I) where I is a nonempty finite set and λ^I is as above, where morphisms $(I, \lambda^I) \to (J, \lambda^J)$ are defined by surjections $\varphi: I \twoheadrightarrow J$ with $\lambda^j = \sum_{i \in \varphi^{-1}(j)} \lambda^i$ for each $j \in J$. The closed immersions ι_{λ^I} assemble into a morphism of presheaves over Ran:

$$
\mathrm{Gr}_{\mathbb{G}_m}^{\text{comb}} \coloneqq \operatornamewithlimits{colim}_{(\mathrm{I},\lambda^{\mathrm{I}})} \mathrm{X}^{\mathrm{I}} \to \mathrm{Gr}_{\mathbb{G}_m},
$$

which induces a bijection on field-valued points.

Claim: factorization pairings $\mathscr{L}^+\mathbb{G}_m \otimes \mathrm{Gr}_{\mathbb{G}_m}^{\text{comb}} \to \mathbb{G}_{m,\text{Ran}}$ are in bijection with sections of $\underline{\mathbb{Z}}$ over X. More precisely, locally on X, a generator is the colimit over (I, λ^I) of maps:

$$
f_{(\mathbf{I},\lambda^{\mathbf{I}})}:\mathscr{L}_{\mathbf{X}^{\mathbf{I}}}^{\mathbf{+}}\mathbb{G}_m \to \mathbb{G}_m, \quad (x^{\mathbf{I}},a) \mapsto \prod_{i\in\mathbf{I}}(a|_{\Gamma_{x^i}})^{\lambda^i},\tag{3.18}
$$

where $a|_{\Gamma_{x^i}}$ is the restriction of a to the closed subscheme $\Gamma_{x^i} \subset D_{x^I}$.

To prove the claim, we use the presentation [\(3.8\)](#page-26-1) of the group X^I -scheme $\mathscr{L}_{X^I}^* \mathbb{G}_m$ as an extension of $R_{\Gamma}\mathbb{G}_m$ by $\mathscr{L}^{\geq 1}_{X^1}\mathbb{G}_m$. Every character $\mathscr{L}^+_{X^1}\mathbb{G}_m \to \mathbb{G}_m$ must factor through $R_{\Gamma}\mathbb{G}_m$ and is uniquely determined by its restriction to the pairwise disjoint locus of X^I .

Given a factorization pairing $f': \mathcal{L}^+\mathbb{G}_m \otimes \mathrm{Gr}_{\mathbb{G}_m}^{\text{comb}} \to \mathbb{G}_{m,\text{Ran}}$, we obtain a system of maps indexed by (I, λ^I) :

$$
f'_{(I,\lambda^I)} : \mathscr{L}_{X^I}^+ \mathbb{G}_m \to \mathbb{G}_m. \tag{3.19}
$$

By the observation above, [\(3.19\)](#page-29-1) is uniquely determined by the case $I = \{1\}$. Moreover, f ′ $(\{1\},\lambda)$ is a character of $\mathrm{R}_{\Gamma}\mathbb{G}_m \cong \mathbb{G}_{m,\mathrm{X}}$, hence a section of $\underline{\mathbb{Z}}$ over X. Looking at I = $\{1,2\}$,

we see that the association $\lambda \mapsto f'_{\alpha}$ $\mathcal{L}_{(\{1\},\lambda)}'$ defines a group homomorphism $\underline{\mathbb{Z}} \to \underline{\mathbb{Z}}$. The group homomorphism corresponding to multiplication by n yields the nth power of (3.18) .

Under the hypothesis char(k) = 0, [\[Tao21a,](#page-48-7) Proposition 5.1.5] shows that any S-point of $Gr_{\mathbb{G}_m}$ admits a factorization $S \to S_0 \to Gr_{\mathbb{G}_m}$, where S_0 is reduced. Therefore, any pairing $\mathscr{L}^+\mathbb{G}_m \otimes \mathbb{G}_{\mathbb{G}_m} \to \mathbb{G}_{m,\text{Ran}}$ is uniquely determined by its values on reduced test schemes, hence on field-valued points. Consequently, any such pairing is uniquely determined by its restriction to $\mathscr{L}^+\mathbb{G}_m \otimes \mathrm{Gr}_{\mathbb{G}_m}^{\rm comb}$.

Corollary 3.3.9. If char(k) = 0, then any factorization pairing $\mathscr{L} \mathbb{G}_m \otimes \mathscr{L} \mathbb{G}_m \to \mathbb{G}_{m, \text{Ran}}$ is an integral power of the Contou-Carrère symbol.

Proof. This is a restatement of Proposition [3.3.8](#page-29-3) in the special case $T_1 = T_2 = \mathbb{G}_m$.

Remark 3.3.10. For any field k, we may view the proof of Proposition [3.3.8](#page-29-3) as establishing the implication $(1) \Rightarrow (2)$ between the following statements:

- (1) any S-point of $\text{Gr}_{\mathbb{G}_m}$ admits a factorization $S \to S_0 \to \text{Gr}_{\mathbb{G}_m}$ where S_0 is reduced;
- (2) any factorization pairing $\mathscr{L} \mathbb{G}_m \otimes \mathscr{L} \mathbb{G}_m \to \mathbb{G}_{m, \text{Ran}}$ is an integral power of the Contou-Carrère symbol.

Since (2) fails when char(k) > 0 (Example [3.3.6\)](#page-28-2), (1) must also fail when char(k) > 0. In other words, the hypothesis char(k) = 0 in [\[Tao21a,](#page-48-7) Theorem 1.2.1] is necessary.

Remark 3.3.11. It is known that the Contou-Carrère symbol is the universal Steinberg symbol over a point of X, cf. [\[GO15\]](#page-47-18). The universal property established in Corollary [3.3.9](#page-30-1) is of a different kind: instead of imposing the Steinberg relation, we impose compatibility with factorization.

3.4. Classification.

3.4.1. Let G be a reductive group X-scheme. Denote by $Rad(G)$ the radical of G, *i.e.* the maximal torus of the center \mathbb{Z}_{G} of G [\[ABD](#page-46-3)⁺66, XXII, Définition 4.3.6].

Our principal goal is to study factorization super central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m, Ran}$ subject to the followng property: the commutator of the induced factorization super central extension of $\mathscr{L} \text{Rad}(G)$ by $\mathbb{G}_{m,\text{Ran}}$ is tame in the sense of §[3.3.3.](#page-27-6) Such a factorization super central extension of $\mathscr{L}G$ by $\mathbb{G}_{m, \text{Ran}}$ is said to have *tame commutator*.

By Example [3.3.6](#page-28-2) and Proposition [3.3.8,](#page-29-3) the condition of having tame commutator is vacuous when $char(k) = 0$, but not so when $char(k) > 0$.

3.4.2. Suppose that G has a maximal torus T with sheaf of cocharacters Λ .

Write $G_{\rm sc}$ for the simply connected form of G with induced maximal torus $T_{\rm sc}$ and sheaf of cocharacters $\Lambda_{\rm sc}$.

Recall that any integral Weyl-invariant quadratic form $Q_{\rm sc}$ on $\Lambda_{\rm sc}$ defines a section of $\vartheta(\Lambda_{\rm sc})$ via [\(2.2\)](#page-15-6). The corresponding central extension $\Lambda_{\rm sc}$ of $\Lambda_{\rm sc}$ by \mathbb{G}_m can be viewed as a monoidal morphism:

$$
\nu_{Q_{sc}} : \Lambda_{sc} \to B\mathbb{G}_m \cong \text{Pic}.\tag{3.20}
$$

We define a morphism of pointed X-stacks by the formula:

$$
\nu_{Q_{sc},+}: \Lambda_{sc} \to \text{Pic}, \quad \lambda \mapsto \nu_{Q_{sc}}(\lambda) \otimes \omega_X^{Q_{sc}(\lambda)}.
$$
 (3.21)

3.4.3. To specify the additional structure of the map $\nu_{Q_{sc,+}}$ inherited from the monoidal structure of $\nu_{Q_{sc}}$, we introduce a piece of terminology.

Let Λ denote, temporarily, any étale sheaf of finite free Z-modules over X and $b : \Lambda \otimes \Lambda \to \mathbb{Z}$ be any symmetric bilinear form. A morphism of X-stacks $\nu_+ : \Lambda \to \text{Pic}^{\text{super}}$ is said to be ω -monoidal with respect to b if it is equipped with ismorphisms:

$$
\mathscr{O}_{X} \xrightarrow{\simeq} \nu_{+}(0);
$$

$$
\nu_{+}(\lambda_{1}) \otimes \nu_{+}(\lambda_{2}) \otimes \omega_{X}^{b(\lambda_{1},\lambda_{2})} \xrightarrow{\simeq} \nu_{+}(\lambda_{1} + \lambda_{2}),
$$

for each $\lambda_1, \lambda_2 \in \Lambda$, satisfying unitality and associativity. (This notion does not refer to the commutativity constraint of the Picard groupoid $\operatorname{Pic}^{\operatorname{super}}.)$

In this terminology, the map $\nu_{Q_{sc},+}$ is ω -monoidal with respect to the symmetric form b_{sc} associated to $Q_{\rm sc}$.

3.4.4. Denote by $\vartheta_{G,+}^{super}(\Lambda)$ the Picard groupoid of triples (b,ν_+,φ) , where:

- (1) b is a Weyl-invariant integral symmetric bilinear form on Λ , such that $b(\lambda, \lambda) \in 2\mathbb{Z}$ if $\lambda \in \Lambda_{\rm sc}$ —we write $Q_{\rm sc}$ for the corresponding quadratic form on $\Lambda_{\rm sc}$;
- (2) ν_{+} is a morphism $\Lambda \to \text{Pic}^{\text{super}}$ which is ω -monoidal with respect to b and commutes with the commutativity constraint up to the bilinear form $(-1)^b$;
- (3) φ is an isomorphism between the restriction of ν_{+} to $\Lambda_{\rm sc}$ and $\nu_{Q_{\rm sc},+}$ as ω -monoidal morphisms.

The relation between $\vartheta_{G,+}^{super}(\Lambda)$ and the Picard groupoid $\vartheta_{G}^{super}(\Lambda)$ defined in §[2.2.1](#page-17-4) is as follows. Given a ϑ -*characteristic* $\omega^{1/2}$, *i.e.* a line bundle over X equipped with an isomorphism $(\omega^{1/2})^{\otimes 2} \cong \omega_{X}$, we obtain an isomorphism (called the $\omega^{1/2}$ -shift):

$$
\vartheta_{\mathcal{G}}^{\text{super}}(\Lambda) \xrightarrow{\simeq} \vartheta_{\mathcal{G},+}^{\text{super}}(\Lambda), \quad (b,\nu,\varphi) \mapsto (b,\nu_{+},\varphi), \tag{3.22}
$$

where ν_+ is the ω -monoidal morphism $\lambda \mapsto \nu(\lambda) \otimes (\omega^{1/2})^{b(\lambda,\lambda)}$ and ν is the monoidal morphism $\Lambda \rightarrow \text{Pic}^{\text{super}}$ corresponding to $\widetilde{\Lambda}$, *i.e.* [\(2.11\)](#page-18-4).

Theorem 3.4.5. Let G be a reductive group X-scheme. The following Picard groupoids are canonically equivalent:

- (1) factorization super central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m, Ran}$ with tame commutator;
- (2) factorization super line bundles over Gr_G ;
- (3) $\vartheta_{G,+}^{super}(\Lambda) if G$ is equipped with a maximal torus T with sheaf of cocharacters Λ .
- (4) rigidified sections of K_{r_1, r_2}^{super} $\sum_{[1,2]}^{\text{super}}$ over the Zariski classifying stack of G -if X is equipped with a ϑ -characteristic.

3.4.6. The equivalences of Theorem [3.4.5](#page-31-0) are constructed in several stages. We briefly indicate the steps and explain where prior works are used.

First, we prove that factorization super central extensions of \mathscr{L}^+G by $\mathbb{G}_{m,Ran}$ are canon-ically trivial (Proposition [4.1.2\)](#page-32-3). By descent, we obtain the functor $(1) \rightarrow (2)$.

Next, the equivalence $(2) \approx (3)$ is essentially known when G is a torus or a semisimple, simply connected group scheme. The torus case is treated in [\[TZ21\]](#page-48-1) (using a substantial theorem of [\[Tao21b\]](#page-48-9)). The simply connected case reduces to Falting [\[Fal03\]](#page-47-19).

One may then define the functor $(2) \rightarrow (3)$ for any reductive group scheme G, by appealing to functoriality with respect to the commutative diagram:

We then proceed as follows. We first prove "by hand" that $(1) \rightarrow (2)$ is an equivalence for G a torus or a semisimple, simply connected group scheme. Then we prove that, for any reductive group scheme G, the composition $(1) \rightarrow (2) \rightarrow (3)$ is an equivalence, whereas the second functor is fully faithful. Here, we use an idea of Finkelberg–Lysenko [\[FL10\]](#page-47-20), an idea of Gaitsgory [\[Gai20\]](#page-47-3), and an argument from [\[TZ21\]](#page-48-1).

The equivalences $(1) \approx (2) \approx (3)$ are completed in Proposition [4.4.13.](#page-44-1)

Finally, the equivalence (3) \leq (4) is defined by combining Theorem [2.2.3](#page-18-0) with the $\omega^{1/2}$ -shift [\(3.22\)](#page-31-1). The composed functor $(1) \rightarrow (2) \rightarrow (3) \rightarrow (4)$ thus a priori depends on a maximal torus T, but it shall follow from the construction that this is not the case. By étale descent, we obtain the equivalence (1) \cong (4) without assuming the existence of a maximal torus.

Corollary 3.4.7. Let G be a reductive group X-scheme. The following Picard groupoids are canonically equivalent:

- (1) factorization central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m,\text{Ran}}$ with tame commutator;
- (2) factorization line bundles over Gr_G ;
- (3) $\vartheta_{\rm G, t}(\Lambda)$, if G is equipped with a maximal torus T with sheaf of cocharacters Λ ;
- (4) central extensions of G by K_2 on the big Zariski site of X.

Proof. Each Picard groupoid in Theorem [3.4.5](#page-31-0) admits a canonical functor to $\text{Hom}(\pi_1\text{G},\mathbb{Z}/2)$: for the Picard groupoids $(1), (2)$, these are the functors remembering the grading; for (3) , this is the second functor in (2.10) ; for (4) , this is the lower horizontal functor in (2.28) .

The equivalences of Theorem [3.4.5](#page-31-0) commute with these functors to $\text{Hom}(\pi_1\text{G},\mathbb{Z}/2)$. By restricting them to the fibers, we obtain the corollary. (Note that the restriction of [\(3.22\)](#page-31-1) is defined without the choice of $\omega^{1/2}$.)

4. Proofs

The goal of this section is to prove Theorem [3.4.5.](#page-31-0) We begin in §[4.1](#page-32-1) by constructing a functor from factorization super central extensions of $\mathscr{L}G$ to factorization super line bundles over Gr_G. The subsections §[4.2,](#page-34-0) §[4.3,](#page-37-0) §[4.4](#page-41-0) prove the equivalences (1) \approx (2) \approx (3) of Theorem [3.4.5](#page-31-0) for tori, simply connected groups, respectively all reductive groups. In §[4.5,](#page-44-0) we show that the equivalence (1) \leq (4) obtained by combining the equivalence (1) \leq (3) and Theorem [2.2.3](#page-18-0) is independent of the choice of a maximal torus.

We remain in the context of the previous section: we fix a ground field k and a smooth curve X.

4.1. Triviality over arc groups.

4.1.1. In this subsection, we let G denote a smooth affine group X-scheme with connected geometric fibers. Our goal is to prove the following result.

Proposition 4.1.2. Any factorization super central extension of \mathscr{L}^+G by $\mathbb{G}_{m, Ran}$ is canonically trivial.

4.1.3. Consider an S-point x^I of Ran. Recall that the restriction $\mathscr{L}_{x^I}^+$ G is an extension of R_{Γ} G by $\mathscr{L}_{x^I}^{\geq 1}$ G as a group S-scheme [\(3.8\)](#page-26-1).

We begin with a Lemma which allows us to reduce the problem to $R_\Gamma G$.

Lemma 4.1.4. If S is locally Noetherian and normal, then pullback along [\(3.8\)](#page-26-1) defines an equivalence between the groupoid of central extensions of $R_{\Gamma}G$ by $\mathbb{G}_{m,S}$ and that of $\mathscr{L}^+_{x^I}G$.

Proof. Given a central extension:

$$
1 \to \mathbb{G}_{m, \mathcal{S}} \to \mathscr{G} \to \mathscr{L}_{x^{\mathrm{I}}}^+ \mathcal{G} \to 1, \tag{4.1}
$$

we claim that [\(4.1\)](#page-32-4) admits a unique splitting over $\mathscr{L}^{\geq 1}_{x}G$, and its image is normal in \mathscr{G} . Then the association $\mathscr{G} \mapsto \mathscr{G}/\mathscr{L}^{\geq 1}_{x^I}G$ supplies the desired inverse functor.

Since $\mathscr{L}_{x^I}^{\geq 1}$ G is an iterated extension of vector group S-schemes, to show the existence and uniqueness of the splitting, it suffices to show that any central extension of $\mathbb{G}_{a,S}$ by $\mathbb{G}_{m,S}$ is canonically split.

This assertion follows from the observations below:

- (1) any S-morphism $\mathbb{G}_{a,S} \to \mathbb{G}_{m,S}$ is trivial—this uses the hypothesis that S is reduced;
- (2) the pullback $Pic(S) \rightarrow Pic(\mathbb{G}_{a,S})$ is an equivalence—this uses the hypothesis that S is locally Noetherian and normal [\[GD67,](#page-47-21) Corollaire 21.4.13, p.361].

To prove that the resulting splitting $\mathscr{L}^{\geq 1}_{x^I}G \to \mathscr{G}$ has normal image, it suffices to show that it commutes with the conjugation action of $\mathscr{L}_{x}^{\dagger}$ G. However, this follows from the uniqueness of the splitting. \Box

Proof of Proposition [4.1.2.](#page-32-3) For an integer $n \geq 1$, we denote by $\text{Ran}^{\leq n} \subset \text{Ran}$ the subfunctor whose S-points are finite subsets of Maps (S, X) of cardinality ≤ 2 .

Given a factorization super central extension:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathscr{G} \to \mathscr{L}^+G \to 1,\tag{4.2}
$$

we first observe that "super" is redundant since the group scheme \mathscr{L}^+_{x} ^IG \rightarrow S for any S-point x^I of Ran has connected geometric fibers. Next, we claim that a trivialization of (4.2) over $\text{Ran}^{\leq 2}$ compatible with the factorization morphism [\(3.7\)](#page-25-4) for $|I| = |J| = 1$ uniquely extends to a trivialization of [\(4.2\)](#page-33-0) over Ran.

For a nonempty finite set I, consider the tautological X^I -point x^I of Ran (Remark [3.1.2\)](#page-24-4). Denote by $U \subset X^I$ the open subset where $\Gamma_{x^{i_1}} \cap \Gamma_{x^{i_2}} \neq \emptyset$ for at most one pair of distinct elements $i_1, i_2 \in I$. Let $Z \subset X^I$ be its complement, a closed subset of codimension ≥ 2 . (It is empty if $|I| \le 2$.) The induced U-point $x^I|_U$ is a disjoint union of x^i (for $i \ne i_1, i_2$) with $x^{\{i_1, i_2\}}$. The factorization morphism for $\mathscr G$ and its trivialization over $\text{Ran}^{\leq 2}$ define a trivialization of $\mathscr{G}_{x^{\mathrm{I}}}$ over U ⊂ X^I.

Since X^I is smooth, \mathscr{G}_{x} ¹ is pulled back from a central extension of R_ΓG by $\mathbb{G}_{m,X}$ ¹ (Lemma [4.1.4\)](#page-32-5). Since R_ΓG is also smooth, the trivialization of \mathscr{G}_{x^I} extends uniquely along U $\subset X^I$ [\[Sta18,](#page-48-18) 031T]. Given a surjection of nonempty finite sets $I \rightarrow J$, we need to argue that the trivialization of \mathscr{G}_{x^I} restricts to the trivialization of \mathscr{G}_{x^J} . This statement reduces to the case $|I| = |J| + 1$, where the diagonal $X^J \subset X^I$ intersects nontrivially with U. Since the two trivializations agree over $X^J \cap U$, they must agree over X^J .

Finally, we construct the trivialization of (4.2) over $\text{Ran}^{\leq 2}$ compatible with factorization. Consider the tautological $X^{\{1,2\}}$ -point $x^{\{1,2\}}$ of Ran, with U ⊂ $X^{\{1,2\}}$ the complement of the diagonal. The closed immersions $\Gamma_{x_i} \subset \Gamma_{x_i}$ _{1,2} (for $i = 1, 2$) induce projection maps from R_ΓG to G. Moreoever, the base change of $R_\Gamma G$ along the diagonal of $X^{\{1,2\}}$ is an extension \widetilde{G} of G by a vector group scheme. These morphisms are summarized in the following diagram:

$$
\begin{array}{ccc}\widetilde{G} & \stackrel{\Delta}{\longrightarrow}R_{\Gamma}G \xrightarrow[\text{pr}_2]{\text{pr}_1} G\\ & & \downarrow & \\ X & \stackrel{\Delta}{\longrightarrow}X^{\{1,2\}}\xrightarrow[\text{pr}_2]{\text{pr}_1}X\end{array}
$$

where both compositions in the upper row are the canonical surjection $\tilde{G} \rightarrow G$.

The restriction of $\mathscr G$ along the tautological X-point of Ran is pulled back from a central extension of G by $\mathbb{G}_{m,X}$, to be denoted by \mathscr{G}_1 . Similarly, the restriction $\mathscr{G}_{x^{(1,2)}}$ is pulled back

from a central extension of R_ΓG, to be denoted by \mathscr{G}_2 . By factorization, the restriction of \mathscr{G}_2 to U ⊂ X^{1,2} is identified with pr[∗] $\mathscr{G}_1 \otimes pr_2^* \mathscr{G}_1$. This identification extends uniquely to an isomorphism between \mathscr{G}_2 with $\mathrm{pr}_1^*\mathscr{G}_1 \otimes \mathrm{pr}_2^*\mathscr{G}_1 \otimes \mathscr{O}(n\Delta)$ as line bundles over $R_\Gamma G$, for some $n \in \mathbb{Z}$. Restriction to the unit section $e: X^{\{1,2\}} \to R_{\Gamma}G$ tells us that $n = 0$, so we obtain an isomorphism of central extensions of R_ΓG by $\mathbb{G}_{m,X^{\{1,2\}}}$:

$$
\mathcal{G}_2 \xrightarrow{\simeq} \text{pr}_1^* \mathcal{G}_1 \otimes \text{pr}_2^* \mathcal{G}_1. \tag{4.3}
$$

Restriction of [\(4.3\)](#page-34-1) along the diagonal then yields an isomorphism $\mathscr{G}_1 \overset{\simeq}{\to} \mathscr{G}_1^{\otimes 2}$, *i.e.* a trivialization of \mathscr{G}_1 . The trivialization of \mathscr{G}_2 is deduced from [\(4.3\)](#page-34-1), so it is automatically compatible with factorization. \Box

4.1.5. Using Proposition [4.1.2,](#page-32-3) we construct a functor of Picard groupoids:

$$
\text{Hom}_{\text{fact}}(\mathcal{L}\mathbf{G}, \text{Pic}^{\text{super}}) \to \Gamma_{\text{fact}}(\text{Gr}_{\mathbf{G}}, \text{Pic}^{\text{super}}), \tag{4.4}
$$

where the target consists of factorization super line bundles over Gr_G .

Indeed, any factorization monoidal morphism $\mathscr{L}G \rightarrow Pic^{super}$ is trivial over \mathscr{L}^+G , so the monoidal structure yields its descent data to $\mathscr{L}G/\mathscr{L}^+G \cong Gr_G$.

One can make a stronger statement: this monoidal structure yields descent data to the local Hecke stack Hec_G ∶= $\mathscr{L}^+G\backslash\mathscr{L}G/\mathscr{L}^+G$, and the resulting factorization super line bundle over Hec_G is compatible with the "convolution structure" of Hec_G. To make this precise, we need the prestack:

$$
\operatorname{Hec}_{G}^{[2]} := \mathscr{L}^+G \backslash \mathscr{L}G \times^{\mathscr{L}^+G} \mathscr{L}G/\mathscr{L}^+G,
$$

equipped with three maps p_1 , m , p_2 to Hec_G: projection onto the first factor, multiplication, and projection onto the second factor. There is also a unit section $e : B(\mathscr{L}^+G) \to \text{Hec}_G$, where B stands for delooping relative to Ran. A factorization super line bundle $\mathscr L$ over Hec_G is *compatible with convolution* if there are additional isomorphisms:

$$
\mathscr{O}_{B(\mathscr{L}^+G)} \xrightarrow{\simeq} e^* \mathscr{L};\tag{4.5}
$$

$$
(p_1)^* \mathcal{L} \otimes (p_2)^* \mathcal{L} \xrightarrow{\simeq} m^* \mathcal{L}, \tag{4.6}
$$

which satisfy the conditions of an associative algebra and commute with factorization.

Let $Hom_{fact}(Hec_G, Pic^{super})$ denote the Picard groupoid of factorization super line bundles over Hec_G compatible with convolution. The descent procedure then yields an equivalence of Picard groupoids:

$$
\text{Hom}_{\text{fact}}(\mathscr{L}\mathbf{G}, \text{Pic}^{\text{super}}) \xrightarrow{\simeq} \text{Hom}_{\text{fact}}(\text{Hec}_{\mathbf{G}}, \text{Pic}^{\text{super}}). \tag{4.7}
$$

This assertion follows at once from two observations: the convolution structure on Hec_G is defined by the Čech nerve of $B(\mathcal{L}^+G) \to B(\mathcal{L}G)$ and any monoidal morphism $B(\mathcal{L}^+G) \to$ Pic^{super} compatible with factorization is canonically trivial (Proposition [4.1.2\)](#page-32-3).

4.2. Tori.

4.2.1. Let Λ be an étale sheaf of finite free Z-modules over X. Denote by $\vartheta_{+}^{super}(\Lambda)$ the Picard groupoid of pairs $(b, F₊)$, where:

- (1) b is an integral symmetric bilinear form on Λ ;
- (2) $F_+ : \Lambda \to \text{Pic}^{\text{super}}$ is an ω -monoidal morphism with respect to b and commutes with the commutativity constraint up to the bilinear form $(-1)^b$.

This is the special case of $\vartheta_{G,+}^{super}(\Lambda)$ defined in §[3.4.4,](#page-31-2) for G the X-torus $\Lambda \otimes \mathbb{G}_m$.

4.2.2. Let T be an X-torus with sheaf of cocharacters Λ. We shall define a functor from the Picard groupoid of factorization super line bundles over Gr_{T} to $\vartheta_{+}^{\text{super}}(\Lambda)$:

$$
\Gamma_{\text{fact}}(\text{Gr}_{T}, \text{Pic}^{\text{super}}) \to \vartheta_{+}^{\text{super}}(\Lambda). \tag{4.8}
$$

This is a variant of [\[BD04,](#page-47-0) §3.10.7], where objects of $\vartheta_{+}^{\text{super}}(\Lambda)$ are called ϑ -data.

For each I-tuple $\lambda^I = (\lambda_i)$ of elements of Λ , there is a closed immersion $\iota_{\lambda^I}: X^I \to \mathrm{Gr}_{T,X^I}$ sending an S-point $x^I = (x^i)$ of X^I to the T-torsor $\mathscr{O}(\lambda_i \Gamma_{x^i})$ equipped with the canonical trivialization off Γ_{x^I} .

To define [\(4.8\)](#page-35-0), we take a factorization super line bundle $\mathscr L$ over $\text{Gr}_{\mathcal T}$ and construct a pair (b, F_+) . Given $\lambda_1, \lambda_2 \in \Lambda$, the line bundle $(\iota_{\lambda_1, \lambda_2})^* \mathscr{L}$ is identified with $(\iota_{\lambda_1})^* \mathscr{L} \boxtimes (\iota_{\lambda_2})^* \mathscr{L}$ off the diagonal of X^2 by factorization—this identification extends to an isomorphism:

$$
(\iota_{\lambda_1,\lambda_2})^* \mathscr{L} \cong (\iota_{\lambda_1})^* \mathscr{L} \otimes (\iota_{\lambda_2})^* \mathscr{L} \otimes \mathscr{O}_{X^2}(b(\lambda_1,\lambda_2)\Delta), \tag{4.9}
$$

for a uniquely defined integer $b(\lambda_1, \lambda_2)$. The associativity and unitality of the factorization isomorphism implies that $\lambda_1, \lambda_2 \mapsto b(\lambda_1, \lambda_2)$ is a bilinear form. Commutativity of the factorization isomorphism implies that b is symmetric and that $b(\lambda, \lambda)$ mod 2 agrees with the grading on $(\iota_{\lambda})^* \mathscr{L}$. The definition of b is complete.

The second datum F_+ is set to be $F_+(\lambda) = (\iota_\lambda)^* \mathscr{L}$, with ω -monoidal structure given by restricting (4.9) along the diagonal. The fact that F_{+} commutes with the brading up to the factor $(-1)^{b(\lambda_1,\lambda_2)}$ follows from the fact that the isomorphism $\mathscr{O}_{X^2}(\Delta)|_{\Delta} \cong \omega_X$ is equivariant against the exchange map $X^2 \to X^2$, $(x^1, x^2) \mapsto (x^2, x^1)$ up to the factor (-1).

Proposition 4.2.3. The functor [\(4.8\)](#page-35-0) is an equivalence of Picard groupoids.

Proof. The proof is identical to that of [\[TZ21,](#page-48-1) Proposition 1.4].

4.2.4. Let T, T' be X-tori whose sheaves of cocharacters are
$$
\Lambda
$$
, respectively Λ' . Suppose that we are given a factorization bilinear pairing:

$$
\langle \cdot, \cdot \rangle : \mathcal{L}^+ \mathbf{T} \otimes \mathbf{G} \mathbf{r}_{\mathbf{T}'} \to \mathbb{G}_m. \tag{4.10}
$$

Delooping in the first variable and pulling back along the projection map $\text{Gr}_{T} \to B(\mathscr{L}^+\text{T})$, we obtain a morphism compatible with factorization:

$$
\langle \cdot, \cdot \rangle : \text{Gr}_{T} \times_{\text{Ran}} \text{Gr}_{T'} \to B\mathbb{G}_{m},\tag{4.11}
$$

or equivalently a factorization line bundle over $Gr_{T\times T'}$.

Consider the factorization pairing $\langle \cdot, \cdot \rangle_b$ defined by a bilinear form $b : \Lambda \otimes \Lambda' \to \mathbb{Z}$ as in §[3.3.3.](#page-27-6) The property of Contou-Carrère symbol shows that $\langle \cdot, \cdot \rangle_b$ induces a pairing \mathscr{L}^+ T ⊗ $\mathrm{Gr}_{T'} \to \mathbb{G}_m$, hence a factorization line bundle $\mathcal{O}(b)$ over $\mathrm{Gr}_{T \times T'}$. We may calculate its image under the equivalence of Proposition [4.2.3.](#page-35-2)

Lemma 4.2.5. The factorization line bundle $\mathcal{O}(b)$ corresponds under the equivalence of Proposition [4.2.3](#page-35-2) to the pair consisting of:

- (1) the quadratic form $\Lambda \oplus \Lambda' \to \mathbb{Z}$, $(\lambda, \lambda') \mapsto b(\lambda, \lambda')$;
- (2) the w-monoidal morphism $\Lambda \oplus \Lambda' \to \text{Pic}$ with $(\lambda, \lambda') \mapsto \omega_X^{b(\lambda, \lambda')}$ $\chi^{b(\lambda,\lambda)}$, whose ω -monoidal structure is the isomorphism:

$$
(-1)^{b(\lambda_2,\lambda'_1)}\cdot \mathrm{id}: \omega_{X}^{b(\lambda_1+\lambda_2,\lambda'_1+\lambda'_2)} \stackrel{\simeq}{\to} \omega_{X}^{b(\lambda_1,\lambda'_1)}\otimes \omega_{X}^{b(\lambda_2,\lambda'_2)}\otimes \omega_{X}^{b(\lambda_1,\lambda'_2)+b(\lambda_2,\lambda'_1)},
$$

for any pair of elements $(\lambda_1, \lambda'_1), (\lambda_2, \lambda'_2) \in \Lambda \oplus \Lambda'.$

Proof. Let $(\lambda')^I = (\lambda'_i)$ be an I-tuple of elements of Λ' . The restriction of $\mathcal{O}(b)$ along:

 $\left(\mathrm{id},\iota_{(\lambda')^{\mathrm{I}}}\right) : \mathrm{Gr}_{\mathrm{T},\mathrm{X}^{\mathrm{I}}} \to \mathrm{Gr}_{\mathrm{T},\mathrm{X}^{\mathrm{I}}} \times_{\mathrm{X}^{\mathrm{I}}} \mathrm{Gr}_{\mathrm{T}',\mathrm{X}^{\mathrm{I}}}$

is the line bundle over Gr_{T,X^I} whose fiber at an S-point (x^I, P_T, α) of Gr_{T,X^I} is given by $\otimes_{i\in I} (P_T|_{\Gamma_{x_i}})^{b(-,\lambda'_i)}$, where the superscript indicates inducing along the character $T \to \mathbb{G}_m$ defined by $b(-, \lambda'_i) : \Lambda \to \mathbb{Z}$.

In particular, for an I-tuple $\lambda^I = (\lambda_i)$ of elements of Λ , further restricting $(id, \iota_{(\lambda')^I})^* \mathcal{O}(b)$ along ι_{λ^I} yields the following line bundle over X^I :

$$
\bigotimes_{i \in I} (p_i^* \omega_X^{b(\lambda_i, \lambda'_i)} \otimes \bigotimes_{\substack{j \in I \\ j \neq i}} p_{ij}^* \mathscr{O}_{X^2}(b(\lambda_j, \lambda'_i) \Delta)), \tag{4.12}
$$

where $p_i: X^I \to X$ (resp. $p_{ij}: X^I \to X^2$) denotes the projection onto the factor labeled by i (resp. factors labeled by (i, j)).

Statement (1) follows by inspecting [\(4.12\)](#page-36-0) for $I = \{1, 2\}$, seeing that the quadratic form $(\lambda, \lambda') \mapsto b(\lambda, \lambda')$ has symmetric form $(\lambda_1, \lambda'_1), (\lambda_2, \lambda'_2) \mapsto b(\lambda_2, \lambda'_1) + b(\lambda_1, \lambda'_2)$. The first part of statement (2) follows by inspecting [\(4.12\)](#page-36-0) for $I = \{1\}$, the second part for $I = \{1, 2\}$, taking into account the fact that the isomorphism $\mathcal{O}_{X^2}(\Delta)|_{\Delta} \cong \omega_X$ is equivariant for the exchange map $X^2 \to X^2$, $(x^1, x^2) \mapsto (x^2, x^1)$ up to the factor (-1) . $\text{map } X^2 \to X^2, (x^1, x^2) \mapsto (x^2, x^1) \text{ up to the factor } (-1).$

4.2.6. Let T be an X-torus with sheaf of cocharacters Λ . We now complete the classification of factorization super central extensions of $\mathscr{L} T$ by $\mathbb{G}_{m,Ran}$ with tame commutator.

In light of the equivalence between factorization super line bundles over Gr_{T} and $\vartheta^{\mathrm{super}}_+(\Lambda)$ (Proposition [4.2.3\)](#page-35-2), it remains to prove the following assertion.

Proposition 4.2.7. The functor [\(4.4\)](#page-34-2) induces an equivalence between:

- (1) factorization super central extensions of \mathscr{L} T with tame commutator; and
- (2) factorization super line bundles over Gr_{T} .

Proof. Passing through the equivalence (4.7) , we replace (4.4) by the forgetful functor:

$$
Homfact(HecT, Picsuper) \to \Gammafact(GrT, Picsuper),
$$
\n(4.13)

defined via pullback along $\text{Gr}_{T} \rightarrow \text{Hec}_{T}$.

The desired equivalence amounts to showing that every factorization super line bundle $\mathscr L$ over $\text{Gr}_{\mathbf T}$ admits a unique collection of the following pieces of structure:

- (1) an \mathscr{L}^+ T-equivariance structure;
- (2) compatibility data with convolution, *i.e.* (4.5) and (4.6) , on the factorization super line bundle over Hec_{t} induced from structure (1),

subject to the tameness condition: the induced factorization super central extension of \mathscr{L} T has tame commutator.

Since the \mathscr{L}^+ T-action on Gr_T is trivial, we may view an \mathscr{L}^+ T-equivariance structure on $\mathscr L$ as a morphism:

$$
\mathcal{L}^+ T \times_{\text{Ran}} \text{Gr}_T \to \mathbb{G}_{m,\text{Ran}},\tag{4.14}
$$

linear in the first variable. On the other hand, given a factorization super central extension of LT with commutator $\langle \cdot, \cdot \rangle : \mathscr{L} \mathbb{T} \otimes \mathscr{L} \mathbb{T} \to \mathbb{G}_{m, \text{Ran}}$, the $\mathscr{L}^+ \mathbb{T}$ -equivariance structure on the induced factorization super line bundle over Gr_T is precisely the map $\mathscr{L}^+T \times Gr_T \to \mathbb{G}_{m, Ran}$ associated to $\langle \cdot, \cdot \rangle$ by restriction (*cf.* §[3.3.5\)](#page-27-5). By the tameness condition, the morphism [\(4.14\)](#page-36-1) must then be of the form $\langle \cdot, \cdot \rangle_{b_1}$ for some bilinear form:

$$
b_1: \Lambda \otimes \Lambda \to \mathbb{Z}.\tag{4.15}
$$

Let us now analyze the compatibility data with convolution. The existence and uniqueness of the unital structure [\(4.6\)](#page-34-5) follows from the canonical triviality of factorization super line bundles over Ran and the bilinearity of (4.15) . The multiplicative structure (4.5) amounts to an isomorphism of factorization super line bundles over $\mathscr{L} T \times^{\mathscr{L}^+ T} G r_T$ equivariant against the leftmost \mathscr{L}^+ T-action. Triviality of the \mathscr{L}^+ T-action on Gr_{T} yields an isomorphism:

$$
\mathscr{L} T \times^{\mathscr{L}^+ T} \mathbf{Gr}_T \cong \mathbf{Gr}_T \times_{\mathrm{Ran}} \mathbf{Gr}_T,
$$

Since the \mathscr{L}^+ T-equivariance is defined by b_1 , the multiplicative structure [\(4.5\)](#page-34-4) amounts to an isomorphism of factorization line bundles over $\text{Gr}_{T} \times \text{Gr}_{T}$:

$$
m^*(\mathcal{L}) \cong (\mathcal{L} \boxtimes \mathcal{L}) \otimes \mathcal{O}(b_1), \tag{4.16}
$$

where $\mathcal{O}(b_1)$ is the factorization line bundle associated to the form b_1 in §[4.2.4.](#page-35-3)

Under the equivalence of Proposition [4.2.3,](#page-35-2) $\mathscr L$ corresponds to a pair (b, F_+) . Applying Proposition [4.2.3](#page-35-2) to $T \times T$, we see that [\(4.16\)](#page-37-1) exists if and only if $b_1 = b$. Indeed, the quadratic form equates $Q(\lambda_1 + \lambda_2)$ with $Q(\lambda_1) + Q(\lambda_2) + b_1(\lambda_1, \lambda_2)$ for each $\lambda_1, \lambda_2 \in \Lambda$, using Lemma [4.2.5\(](#page-35-4)1). When $b_1 = b$, Lemma 4.2.5(2) yields a canonical isomorphism of ω -monoidal morphisms associated to the two sides of (4.16) . The isomorphism (4.16) thus defined is the unique one satisfying the cocycle condition. \Box

4.2.8. Consider any factorization super central extension of \mathscr{L} T with tame commutator:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathcal{F} \to \mathcal{L} \mathbf{T} \to 1. \tag{4.17}
$$

The equivalences of Proposition [4.2.3,](#page-35-2) Proposition [4.2.7](#page-36-3) show that [\(4.17\)](#page-37-2) is classified by a pair (b, F_+) in $\vartheta_{+}^{\text{super}}(\Lambda)$.

In the course of the proof of Proposition [4.2.7,](#page-36-3) we have also established the fact that the commutator pairing of (4.17) equals $\langle \cdot, \cdot \rangle_b$.

4.3. Simply connected groups.

4.3.1. Let G denote a semisimple and simply connected group X-scheme.

Since $Gr_G \to \text{Ran}$ (resp. $\mathscr{L}G \to \text{Ran}$) has connected geometric fibers, every factorization super line bundle over Gr_G (resp. factorization super central extension of $\mathscr{L}G$ by $\mathbb{G}_{m, \text{Ran}}$) is pure of even grading.

Proposition 4.3.2. If G is semisimple and simply connected, then the functor (4.4) is an equivalence between:

- (1) factorization central extensions of $\mathscr{L}G$; and
- (2) factorization line bundles over Gr_G .

4.3.3. Before proving Proposition [4.3.2,](#page-37-3) we define Schubert varieties in Gr_G as flat schematic morphisms to Ran. We give a detailed presentation because Lemma [4.3.4](#page-38-0) below was also used in the proof of [\[TZ21,](#page-48-1) Lemma 3.6] but the justification there is inadequate.

Let us assume that G contains a Borel subgroup and a maximal torus $T \subset B \subset G$. Denote by $\Lambda^+ \subset \Lambda$ the subsheaf of dominant cocharacters of T. For an I-tuple $\lambda^I = (\lambda^i)$ of elements of Λ^+ , we may view ι_{λ^I} of §[4.2.2](#page-35-5) as a closed immersion $X^I \to Gr_{G,X^I}$. Denote by $Gr_G^{\leq \lambda^I} \subset Gr_{G,X^I}$ the schematic image of the map $\mathscr{L}_{X^I}^*G \to Gr_{G,X^I}$ defined by acting on $\iota_{\lambda^I}(X^I)$.

Since G is semisimple and simply connected, $\text{Gr}_{G,X}$ is reduced and $\text{Gr}_{G,X}$ $\rightarrow X^I$ has connected geometric fibers. In particular, the above closed subschemes define an isomorphism of indschemes:

$$
\underset{\lambda^I}{\text{colim}} \operatorname{Gr}_{G}^{\leq \lambda^I} \xrightarrow{\simeq} \operatorname{Gr}_{G,X^I}.
$$
\n(4.18)

Lemma 4.3.4. For each I-tuple λ^I of elements of Λ^+ , the projection $p: Gr_G^{\leq \lambda^I} \to X^I$ is flat and the canonical map below is an isomorphism:

$$
\mathscr{O}_{X^I} \to Rp_* \mathscr{O}_{G_r \zeta^{\times I}}.\tag{4.19}
$$

Proof. For $|I| \leq 2$, flatness of p is established in [\[Zhu09,](#page-48-19) §1.2]. The argument below which applies to general I is explained to me by João Lourenço. We call a morphism $f: Y_1 \rightarrow Y_2$ of schemes *derived* $\mathcal{O}\text{-connected}$ if the induced map $\mathcal{O}_{Y_2} \to Rf_*\mathcal{O}_{Y_1}$ is an equivalence.

We begin by recalling some classical facts taking place over geometric points of X. Let x be a \bar{k} -point of X and $\text{Gr}_{G,x}$ the fiber of Gr_G . Let W denote the Weyl group of (G, T) and W_{aff} := $\Lambda \rtimes W$ its affinization. Write $I_x \subset \mathscr{L}_x^+$ for the Iwahori group scheme associated to the Borel B. The affine flag variety $\text{Fl}_{G,x} \coloneqq \mathscr{L}_x G/I_x$ has I_x -orbits parametrized by W_{aff} . For each $\lambda \in \Lambda^+$, the preimage of $\mathrm{Gr}_{G,x}^{\leq \lambda}$ along $\mathrm{Fl}_{G,x} \to \mathrm{Gr}_{G,x}$ concides with $\mathrm{Fl}_{G,x}^{\leq w(\lambda)}$, the closure of the I_x-orbit corresponding to the longest element $w(\lambda)$ in W λ W \subset W_{aff}.

The scheme $\text{Fl}_{G,x}^{\leq w(\lambda)}$ admits a Demazure resolution $D^{w(\lambda)}$ associated to any reduced expression of $w(\lambda)$:

$$
D^{w(\lambda)} \xrightarrow{\pi} \mathrm{Fl}_{G,x}^{\leq w(\lambda)} \to \mathrm{Gr}_{G,x}^{\leq \lambda}.
$$
\n(4.20)

By [\[Fal03,](#page-47-19) Theorem 8], the morphism π is derived $\mathcal{O}\text{-connected}$. Since $G/B \to Spec(\bar{k})$ is derived $\mathcal{O}\text{-connected}$ [\[Kem76\]](#page-48-20), the same holds for the composition [\(4.20\)](#page-38-1). We collect two consequences of this fact:

- (1) $\mathrm{Gr}_{G,x}^{\leq \lambda} \to \mathrm{Spec}(\bar{k})$ is derived $\mathcal{O}\text{-connected}$; indeed, this is because $D^{w(\lambda)} \to \mathrm{Spec}(\bar{k})$ is derived $\mathscr O$ -connected, being an iterated $\mathbb P^1$ -bundle.
- (2) the convolution map $\operatorname{Gr}_{G,x}^{\leq \lambda_1} \widetilde{\times} \operatorname{Gr}_{G,x}^{\leq \lambda_2} \to \operatorname{Gr}_{G,x}^{\leq \lambda_1 + \lambda_2}$ is derived $\mathscr O$ -connected; indeed, the source and target both admit rational resolutions in the sense of [\[Kov22,](#page-48-21) Definition 9.1], so this claim follows from [\[Kov22,](#page-48-21) Theorem 9.12(i)].

To prove that $p: Gr_G^{\leq \lambda^I} \to X^I$ is flat, we consider the global convolution map $m: \widetilde{Gr}_G^{\leq \lambda^I} \to$ $\text{Gr}_{G}^{\leq \lambda^{I}}$ over X^{I} , where the composition $\widetilde{\text{Gr}}_{G}^{\leq \lambda^{I}} \to X^{I}$ is evidently flat. Statement (2) implies that for each \bar{k} -point x^I of X^I , the base change m_{x^I} of m satisfies $R^i(m_{x^I})_*\mathscr{O} \cong 0$ for $i \ge 1$. By [Gör03, Proposition 3.13], $m_*\mathcal{O}$ is X^I-flat and its formation is compatible with base change along X^I. Since $\text{Gr}_{G}^{\leq \lambda^{I}}$ is reduced and m is surjective on \bar{k} -points, we see that $m_{*}\mathscr{O}$ coincides with the structure sheaf of $\text{Gr}_{G}^{\leq \lambda^{I}}$. This implies that p is flat and its geometric fibers are identified with the corresponding Schubert varieties. The derived $\mathscr O$ -connectedness [\(4.19\)](#page-38-2) then follows from its pointwise version, *i.e.* statement (1) above.

4.3.5. Lemma [4.3.4](#page-38-0) has the following consequence: given any S-point $x^I : S \to X^I$, a line bundle $\mathscr L$ over Gr_{G,r^I} descends to S if and only if it is trivial over all geometric fibers. Indeed, if $\mathscr L$ is trivial over all geometric fibers, then by Lemma [4.3.4](#page-38-0) and cohomology and base change, the derived pushforward of $\mathscr L$ to S yields the desired descent.

On the other hand, [\[Fal03,](#page-47-19) Theorem 7] proves that the Picard group of $\text{Gr}_{G,x}$, for any \bar{k} -point x of X, is isomorphic to \mathbb{Z}^r , where r denotes the number of simple factors of G, with $(1,\cdots,1)\in\mathbb{Z}^r$ corresponding to an ample line bundle over $\mathrm{Gr}_{\mathbf{G},x}.$

Proof of Proposition [4.3.2.](#page-37-3) The problem is of étale locally nature on X, so we may assume that G contains a Borel subgroup and a maximal torus $T \subset B \subset G$.

Let $\mathscr L$ be a factorization line bundle over Gr_G . According to §[4.1.5,](#page-34-6) it suffices to prove that $\mathscr L$ admits a unique $\mathscr L^+G$ -equivariance structure and the induced factorization line bundle over Hec_{G} admits unique compatibility data with respect to convolution.

Consider an S-point (x^I, g) of $\mathscr{L}_{X^I}^*$ G. The action by g defines an automorphism act_g of Gr_{G,x^I} . There is a unique isomorphism:

$$
(\text{act}_g)^* \mathcal{L} \xrightarrow{\simeq} \mathcal{L} \tag{4.21}
$$

extending the identity over the unit section $e : S \to \text{Gr}_{G,x^I}$. Indeed, this is because the difference $(\text{act}_g)^* \mathscr{L} \otimes \mathscr{L}^{-1}$ is trivial along geometric fibers, so we may apply the observation of $\S4.3.5$. The uniqueness of (4.21) implies that it satisfies the cocycle condition.

The compatibility data with respect to convolution consist of isomorphisms [\(4.5\)](#page-34-4) and [\(4.6\)](#page-34-5). The second isomorphism is clear. The first one amounts to an isomorphism of line bundles over:

$$
\operatorname{Gr}_G \widetilde{\times} \operatorname{Gr}_G \coloneqq \mathscr{L} G \times^{\mathscr{L}^+G} \operatorname{Gr}_G,
$$

compatible with the left \mathscr{L}^+ G-equivariance. This isomorphism is constructed in the same way as (4.21) , by reducing to geometric fibers over Ran.

4.3.6. Suppose that G contains a maximal torus T with sheaf of cocharacters Λ . The Weyl group W acts naturally on Λ . Restricting along T \subset G and applying Proposition [4.2.3,](#page-35-2) each factorization line bundle over Gr_G defines a pair (b, F_+) with $b(\lambda, \lambda) \in 2\mathbb{Z}$, hence quadratic form $Q : \lambda \mapsto b(\lambda, \lambda)/2$ on Λ .

By [\[TZ21,](#page-48-1) Proposition 2.5], the quadratic form Q is Weyl-invariant and this procedure defines an equivalence of Picard groupoids between factorization line bundles over Gr_G and Weyl-invariant quadratic forms $\text{Quad}(\Lambda, \mathbb{Z})^W$ on Λ . (Evaluation on short coroots belonging to each simple factor of G defines an isomorphism $\mathrm{Quad}(\Lambda, \mathbb{Z})^{\mathrm{W}} \cong \mathbb{Z}^{\oplus r}.$

In summary, all of the Picard groupoids below are canonically equivalent when G is semisimple and simply connected:

$$
\text{Hom}_{\text{fact}}(\mathscr{L}G, \text{Pic}) \stackrel{\simeq}{\to} \text{Hom}_{\text{fact}}(\mathscr{L}G, \text{Pic}^{\text{super}})
$$
\n
$$
\downarrow^{\cong} \qquad \qquad \downarrow^{\cong}
$$
\n
$$
\Gamma_{\text{fact}}(\text{Gr}_G, \text{Pic}) \stackrel{\simeq}{\longrightarrow} \Gamma_{\text{fact}}(\text{Gr}_G, \text{Pic}^{\text{super}})
$$
\n
$$
\downarrow^{\cong}
$$
\n
$$
\text{Quad}(\Lambda, \mathbb{Z})^W
$$
\n(4.22)

In particular, composing these equivalences with the restriction along $T \subset G$ and the functor [\(4.8\)](#page-35-0), we obtain a functor:

$$
Quad(\Lambda, \mathbb{Z})^W \to \vartheta_+^{super}(\Lambda). \tag{4.23}
$$

In [\[TZ21,](#page-48-1) §2.4.7], we have verified that the image of Q is the pair (b, F_+) , where F_+ is the ω -twist of the monoidal morphism F_Q defined in §[3.4.2.](#page-30-2)

Remark 4.3.7. The equivalences in (4.22) remain valid when factorization central extensions of $\mathscr{L}G$ (resp. factorization line bundles over Gr_G) are replaced by central extensions of \mathscr{L}_x G (resp. line bundles over $\text{Gr}_{G,x}$) for any geometric point x of X.

4.3.8. Let us relax the hypothesis and let G be any reductive group X-scheme. We shall now use our knowledge about the simply connected case to perform a commutator calculation.

Denote by G_{sc} the simply connected form of G and G_{ad} the adjoint form of G. The G_{ad} -action on G by conjugation extends to a G_{ad} -action on G_{sc} , which we still refer to as the conjugation action.

Consider any factorization central extension:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathcal{G}_{\text{sc}} \to \mathcal{L}G_{\text{sc}} \to 1. \tag{4.24}
$$

Claim: the conjugation $\mathscr{L}G_{ad}$ -action on $\mathscr{L}G_{sc}$ extends uniquely to \mathscr{G}_{sc} .

Indeed, let (x^I, g) be an S-point of $\mathscr{L}G_{ad}$. Action by g defines an automorphism act_g of \mathscr{L}_{x} G_{sc}. Viewing \mathscr{G}_{sc} as a multiplicative line bundle over $\mathscr{L}G_{sc}$, we shall argue that there is a unique isomorphism:

$$
(\text{act}_g)^* \mathscr{G}_{\text{sc},x^{\text{I}}} \xrightarrow{\simeq} \mathscr{G}_{\text{sc},x^{\text{I}}},\tag{4.25}
$$

compatible with the multiplicative structure of $\mathscr{G}_{sc,x}$. According to §[4.3.5,](#page-38-3) it suffices to show that the two sides of [\(4.25\)](#page-40-0) are isomorphic on geometric fibers over S. This statement holds by Remark [4.3.7,](#page-39-2) seeing that $(\text{act}_g)^*$ induces the identity map on $\text{Quad}(\Lambda, \mathbb{Z})^W$.

4.3.9. Suppose that G contains a maximal torus T with sheaf of cocharacters Λ . Write T_{sc}, $T_{\rm ad}$ for the induced maximal tori in $G_{\rm sc}$, $G_{\rm ad}$, with sheaves of cocharacters $\Lambda_{\rm sc}$, $\Lambda_{\rm ad}$.

The $\mathscr{L}G_{\text{ad}}$ -action on \mathscr{G}_{sc} constructed in §[4.3.8](#page-39-3) restricts to an $\mathscr{L}T_{\text{ad}}$ -action, and we obtain a factorization pairing:

$$
\mathscr{L}T_{\text{ad}} \otimes \mathscr{L}T_{\text{sc}} \to \mathbb{G}_{m,\text{Ran}}, \quad (t_{\text{ad}}, t_{\text{sc}}) \mapsto (t_{\text{ad}}\tilde{t}_{\text{sc}}t_{\text{ad}}^{-1}) \cdot \tilde{t}_{\text{sc}}^{-1}, \tag{4.26}
$$

where $t_{\rm sc}$ is an arbitrary lift of $t_{\rm sc}$ to $\mathscr{G}_{\rm sc}$, which exists locally.

To compute this pairing, we recall that \mathscr{G}_{sc} is classified by a Weyl-invariant quadratic form $Q_{\rm sc}$ on $\Lambda_{\rm sc}$. Since $\Lambda_{\rm ad}$ is canonically dual to the root lattice, the formula:

$$
(\lambda, \alpha) \mapsto Q_{\rm sc}(\alpha) \langle \lambda, \check{\alpha} \rangle
$$

for each $\lambda \in \Lambda_{\text{ad}}$ and coroot $\alpha \in \Lambda_{\text{sc}}$ yields a pairing $b_1 : \Lambda_{\text{ad}} \otimes \Lambda_{\text{sc}} \to \mathbb{Z}$.

Lemma 4.3.10. The factorization pairing [\(4.26\)](#page-40-1) equals $\langle \cdot, \cdot \rangle_{b_1}$ (in the notation of §[3.3.3\)](#page-27-6).

Proof. The problem is of étale local nature on X, so we may assume that G is split and T is a split maximal torus. Each coroot α induces a morphism $f_{\alpha} : SL_2 \to G_{sc}$, sending the upper-triangular unipotent matrices to the root subgroup $U_{\alpha} \subset G_{\rm sc}$ and restricts to α on the diagonal torus $\mathbb{G}_m \subset SL_2$.

The central extension [\(4.26\)](#page-40-1) restricts along f_{α} to the $Q_{sc}(\alpha)$ -multiple of the factorization central extension:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \widetilde{\text{SL}}_2 \to \mathcal{L}\text{SL}_2 \to 1,\tag{4.27}
$$

defined by restricting the Tate central extension \widetilde{GL}_2 along $\mathscr{L}SL_2 \subset \mathscr{L}GL_2$. Indeed, factorization central extensions of $\mathscr{L}SL_2$ are uniquely determined by their quadratic forms (§[4.3.6\)](#page-39-4), hence by the commutator of their restrictions to $\mathscr{L}\mathbb{G}_m$, so we conclude by §[4.2.8](#page-37-4) and our definition of the Contou-Carrère symbol.

By functoriality of the construction, it suffices to show that [\(4.26\)](#page-40-1) equals the Contou-Carrère pairing for G = SL_2 and $\mathscr{G}_{\rm sc}$ being the central extension [\(4.27\)](#page-40-2). Note that the T_{ad} $(=(\mathbb{G}_m)$ -action on G $(=\mathrm{SL}_2)$ extends to the inner action of \mathbb{G}_m on GL_2 as the subgroup:

$$
\mathbb{G}_m \subset \text{GL}_2, \quad a \mapsto \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}.
$$
 (4.28)

Using the group structure on \widetilde{GL}_2 , we extend the induced inner $\mathscr{L} \mathbb{G}_m$ -action on $\mathscr{L} GL_2$ to an action on $\widetilde{\text{GL}}_2$. This action must restrict to the T_{ad}-action on the subgroup $\widetilde{\text{SL}}_2 \subset \widetilde{\text{GL}}_2$, by the uniqueness of the latter (§[4.3.8\)](#page-39-3). Therefore, it remains to prove that the commutator pairing in $\widetilde{\text{GL}}_2$ between the subtorus [\(4.28\)](#page-40-3) and the subtorus:

$$
\mathbb{G}_m \subset \text{GL}_2, \quad a \mapsto \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}
$$

is the Contou-Carrère symbol. This assertion holds because \widetilde{GL}_2 restricts to the Tate central extension $\tilde{\mathbb{G}}_m$ of $\mathscr{L}\mathbb{G}_m$ along [\(4.28\)](#page-40-3).

4.4. Descent.

4.4.1. Let $\mathscr X$ and $\mathscr H$ be factorization group presheaves whose base changes along any S-point of Ran are fppf sheaves.

An action of $\mathscr H$ on $\mathscr K$ as group presheaves is *compatible with factorization* if for any disjoint S-points x^I, x^J of Ran, the $\mathscr{H}_{x^I \sqcup x^J}$ -action on $\mathscr{K}_{x^I \sqcup x^J}$ coincides with the $\mathscr{H}_{x^I} \times \mathscr{H}_{x^J}$ action on $\mathcal{K}_x \times \mathcal{K}_x$ under the factorization isomorphisms of \mathcal{H} and \mathcal{K} . When this happens, the group presheaf $\mathscr K \rtimes \mathscr H$ over Ran inherits a factorization structure.

Consider a factorization super central extension:

$$
1 \to \mathbb{G}_m \to \widetilde{\mathcal{K}} \to \mathcal{K} \to 1.
$$

Suppose that $\widetilde{\mathscr{K}}$ is equipped with an \mathscr{H} -action which is trivial on A. Then we say that the \mathscr{H} -action on $\widetilde{\mathscr{K}}$ is *compatible with factorization* if for disjoint S-points x^I, x^J of Ran, the $\mathscr{H}_{x^I\sqcup x^J}$ -action on $\widetilde{\mathscr{K}}_{x^I\sqcup x^J}$ coincides with the induced $\mathscr{H}_{x^I}\times \mathscr{H}_{x^J}$ -action on the quotient:

$$
\widetilde{\mathscr{K}_{x^{\mathrm{I}}}} \times \widetilde{\mathscr{K}_{x^{\mathrm{J}}}} \twoheadrightarrow \widetilde{\mathscr{K}_{x^{\mathrm{I}} \sqcup x^{\mathrm{J}}}}.
$$

The following lemma is a variant of [\[BD01,](#page-46-1) Construction 1.7].

Lemma 4.4.2. Let \mathcal{K} , \mathcal{H} be as in §[4.4.1](#page-41-1) with \mathcal{H} acting on \mathcal{K} compatibly with factorization. The following categories are equivalent:

- (1) factorization super central extensions of $\mathscr{K} \rtimes \mathscr{H}$ by $\mathbb{G}_{m, Ran}$;
- (2) triples $(\widetilde{\mathcal{K}}, \widetilde{\mathcal{H}}, \alpha)$, where $\widetilde{\mathcal{K}}$ (resp. $\widetilde{\mathcal{H}}$) is a factorization super central extension of $\mathscr K$ (resp. H) by $\mathbb{G}_{m, \text{Ran}}$, and α is an $\mathscr H$ -action on $\widetilde{\mathscr K}$ which is trivial on $\mathbb{G}_{m, \text{Ran}}$, compatible with factorization, and induces the given $\mathscr H$ -action on $\mathscr K$.

Proof. The functor (1) \Rightarrow (2) is given by restricting a factorization central extension of $\mathscr{K} \rtimes \mathscr{H}$ along the group sub-presheaves $\mathscr{K} \subset \mathscr{K} \rtimes \mathscr{H}$, $\mathscr{H} \subset \mathscr{K} \rtimes \mathscr{H}$ to obtain $\widetilde{\mathscr{K}}$, $\widetilde{\mathscr{H}}$, and observing that the $\widetilde{\mathscr{H}}$ -action on $\widetilde{\mathscr{K}}$ factors through \mathscr{H} .

The functor $(2) \Rightarrow (1)$ is given by forming the central extension $\widetilde{\mathscr{K}} \rtimes \widetilde{\mathscr{H}}$ of $\mathscr{K} \rtimes \mathscr{H}$ by \mathbb{G}_m , Ran $\times \mathbb{G}_m$, Ran using the action α , and pushing out along the product map on \mathbb{G}_m , Ran. \square

Proposition 4.4.3. Let G be a reductive group X-scheme with a maximal torus T. Denote by $G_{\rm sc}$ the simply connected form of G, equipped with the conjugation T-action. The following categories are canonically equivalent:

- (1) factorization super central extensions of $\mathscr{L}G_{\mathrm{sc}} \rtimes \mathscr{L}T$ by $\mathbb{G}_{m,\mathrm{Ran}}$;
- (2) pairs $(\mathscr{G}_{\text{sc}},\mathscr{T})$, where \mathscr{G}_{sc} (resp. $\mathscr{T})$ is a factorization super central extension of $\mathscr{L}G_{\text{sc}}$ (resp. $\mathscr{L}T$) by $\mathbb{G}_{m, \text{Ran}}$.

Proof. We appeal to the equivalence of Lemma [4.4.2.](#page-41-2) It suffices to prove that given any pair $(\mathscr{G}_{sc}, \mathscr{T})$ in (2), there is a unique \mathscr{L} T-action on \mathscr{G}_{sc} which is trivial on $\mathbb{G}_{m, Ran}$, compatible with factorization, and induces the conjugation action on $\mathscr{L}G_{sc}$. This is established in §[4.3.8](#page-39-3) when T is replaced by G_{ad} , but the argument for uniqueness carries over.

4.4.4. Under the equivalence of Proposition [4.4.3,](#page-41-3) we shall write the factorization super central extension of $\mathscr{L}G_{\mathrm{sc}} \rtimes \mathscr{L}T$ induced from $(\mathscr{G}_{\mathrm{sc}}, \mathscr{T})$ as follows:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathscr{G}_{\text{sc}} \widetilde{\rtimes} \mathcal{F} \to \mathscr{L}G_{\text{sc}} \rtimes \mathscr{L}T \to 1.
$$

It is by construction the pushout of $\mathscr{G}_{\text{sc}} \rtimes \mathscr{T}$ along the product map on $\mathbb{G}_{m,\text{Ran}}$.

Lemma 4.4.5. Let $\widetilde{G} \rightarrow G$ be a surjection of reductive group X-schemes whose kernel is a torus. The induced morphism $\mathscr{L}\widetilde{G} \to \mathscr{L}G$ is surjective in the topology generated by fpqc and proper covers.

Proof. We shall deduce this from two statements:

- (1) $\mathscr{L}^+ \widetilde{G} \rightarrow \mathscr{L}^+ G$ is surjective in the fpqc topology;
- (2) $Gr_{\tilde{C}} \rightarrow Gr_G$ is surjective in the topology generated by proper covers.

Let us prove the lemma assuming both statements. Indeed, the morphism $\mathscr{L} \widetilde{G} \to \mathscr{L} G$ factors as:

$$
\mathscr{L} \widetilde{G} \to \mathscr{L} G \times_{\mathrm{Gr}_G} \mathrm{Gr}_{\widetilde{G}} \to \mathscr{L} G.
$$

By statement (2), the second morphism is surjective in the proper topology. We claim that the first morphism is surjective in the fpqc topology. Indeed, consider an S-point (g, x) of $\mathscr{L}G \times_{\text{Gr}_G} \text{Gr}_{\widetilde{G}}$. Since $\mathscr{L}G \to \text{Gr}_{\widetilde{G}}$ is surjective in the étale topology, we may lift x to an S₁-point \tilde{g} of $\mathscr{L}\tilde{G}$ over some étale cover S₁ \rightarrow S. The image of \tilde{g} in $\mathscr{L}G$ differs from g by an S_1 -point h of \mathscr{L}^+G . Statement (1) allows us to lift h to an S_2 -point \tilde{h} in $\mathscr{L}^+\tilde{G}$ over some fpqc cover $S_2 \rightarrow S_1$, which we may then use to modify \tilde{g} to obtain a lift of (g, x) .

To prove statement (1), we consider an S-point x^I of X^I with graph $\Gamma_{x^I} \subset S \times X$. The base change $\mathscr{L}_{x^I}^+ \widetilde{G} \to \mathscr{L}_{x^I}^+ G$ is the inverse limit of morphisms:

$$
R_{\Gamma^{(n)}}\widetilde{G} \to R_{\Gamma^{(n)}}G,\tag{4.29}
$$

which are the Weil restrictions of $\tilde{G} \to G$ along the finite locally free morphisms $\Gamma_{x}^{(n)} \to S$, see §[3.2.3.](#page-26-3) Since $\widetilde{G} \rightarrow G$ is affine, smooth, and surjective, the same holds for [\(4.29\)](#page-42-0). Hence $\mathscr{L}_{x}^{\dagger} \widetilde{G} \to \mathscr{L}_{x}^{\dagger} G$ is affine, *flat*, and surjective.

We now turn to statement (2). Since the formation of the affine Grassmannian is compatible with étale base change, we may assume that the kernel of $\widetilde{G} \rightarrow G$ is a *split* torus T. This implies that $\mathscr{L}\widetilde{G} \to \mathscr{L}G$ is surjective on field-valued points. Indeed, given any k-field F, the graph of an F-point of X^I is the disjoint union of schemes isomorphic to $Spec(F((t))),$ but the map $\widetilde{G}(F((t))) \to G(F((t)))$ is surjective because $H^1(F((t)), T) = 0$. In particular, any F-point of Gr_G lifts to Gr_G^a after a finite extension $F \subset F_1$.

On the other hand, the morphism $Gr_{\tilde{G}} \rightarrow Gr_G$ is ind-proper because \tilde{G} is reductive. Since Gr_G is of finite presentation, taking schematic points of Gr_G puts us in the following situation: an affine k-scheme S of finite type, an ind-proper S-indscheme Y, such that $Y \rightarrow S$ is surjective on field-valued points up to finite extension. We claim that some closed subscheme $Y_i \subset Y$ surjects onto S. Then $Y_i \to S$ is a proper cover which lifts to Y.

Let us prove the claim. Since S is of finite type over k , we reduce to the case where S is irreducible. Then its generic point lifts to some Y_i after a finite extension. Since $Y_i \rightarrow S$ is proper, its image contains the closure of the generic point which is all of S.

4.4.6. Let G denote a reductive group X-scheme with a maximal torus T. Denote by G_{sc} the simply connected form of G and $T_{sc} \subset G_{sc}$ the preimage of T. The T-action on G by conjugation extends to G_{sc} . There is a short exact sequence:

$$
1 \to T_{\rm sc} \to G_{\rm sc} \rtimes T \to G \to 1,\tag{4.30}
$$

where the first map is the anti-diagonal embedding $t \mapsto (t, t^{-1})$. Furthermore, its image is central in $G_{sc} \rtimes T$.

The exact sequence [\(4.30\)](#page-42-1) induces an exact sequence of factorization group presheaves:

$$
1 \to \mathcal{L}T_{\rm sc} \to \mathcal{L}G_{\rm sc} \rtimes \mathcal{L}T \to \mathcal{L}G, \tag{4.31}
$$

where the last map is surjective in the topology generated by fpqc and proper covers. Since perfect complexes satisfy derived proper descent [\[Cho22,](#page-47-23) Theorem 1.8] and loop groups are classical [\[GR14,](#page-47-24) Theorem 9.3.5], central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m,Ran}$ are equivalent to those of $\mathscr{L}G_{\mathrm{sc}} \times \mathscr{L}T$ by $\mathbb{G}_{m,\text{Ran}}$ equipped with a splitting over $\mathscr{L}T_{\mathrm{sc}}$ whose image is normal. (This idea is due to Gaitsgory, cf. [\[Gai20,](#page-47-3) Corollary 5.2.7].)

4.4.7. Let (G, T) be as above. Appealing to Proposition [4.4.3,](#page-41-3) we obtain an equivalence of Picard groupoids between:

- (1) factorization super central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m,\text{Ran}}$; and
- (2) triples $(\mathcal{T}, \mathscr{G}_{sc}, \varphi)$, where \mathcal{T} and \mathscr{G}_{sc} are factorization super central extensions:

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathcal{F} \to \mathcal{L} \mathbf{T} \to 1,\tag{4.32}
$$

$$
1 \to \mathbb{G}_{m,\text{Ran}} \to \mathscr{G}_{\text{sc}} \to \mathscr{L} \mathbf{G}_{\text{sc}} \to 1,\tag{4.33}
$$

and φ is an isomorphism of their pullbacks to $\mathscr{L} T_{\text{sc}}$, subject to the normality con $dition$ that the section $\mathscr{L} T_{sc}\to \mathscr{G}_{sc} \widetilde{\rtimes} \mathscr{T}$ induced from φ has normal image.

Lemma 4.4.8. A factorization super central extension of $\mathscr{L}G$ by $\mathbb{G}_{m, Ran}$ has tame commutator if and only if its restriction to \mathscr{L} does.

Proof. Note that any factorization (super) central extension of $\mathscr{L}G_{sc}$ by $\mathbb{G}_{m, Ran}$ has tame commutator (Lemma [4.3.10\)](#page-40-4). The claim now follows from Lemma [3.3.5,](#page-27-5) seeing that Rad(G) \times
T_{sc} \rightarrow T is an isogeny of X-tori. $T_{\rm sc} \rightarrow T$ is an isogeny of X-tori.

4.4.9. Consider a triple $(\mathcal{T}, \mathcal{G}_{\text{sc}}, \varphi)$ as in §[4.4.7](#page-43-0) and assume that \mathcal{T} has tame commutator. In particular, this implies that its commutator is $\langle \cdot, \cdot \rangle_b$ where b is the symmetric bilinear

form appearing in the classifying data of \mathcal{T} , see §[4.2.8.](#page-37-4) Under this assumption, we shall make the normality condition of $\S 4.4.7(2)$ $\S 4.4.7(2)$ $\S 4.4.7(2)$ explicit.

Lemma 4.4.10. If $\mathscr T$ has commutator $\langle \cdot, \cdot \rangle_b$, then the normality condition holds if and only if b is Weyl-invariant.

Proof. Let Λ (resp. $\Lambda_{\rm ad}$, $\Lambda_{\rm sc}$) denote the sheaf of cocharacters of T (resp. T_{ad}, T_{sc}). Note that \mathscr{T} has commutator $\langle \cdot, \cdot \rangle_b$ while $\mathscr{G}_{\rm sc}$ defines the pairing $\langle \cdot, \cdot \rangle_b$ via its $\mathscr{L}T_{\rm ad}$ -action (see Lemma [4.3.10\)](#page-40-4). The existence of φ implies that b and b₁ agree on $\Lambda_{\rm sc} \otimes \Lambda_{\rm sc}$. In particular, the restriction of b to $\Lambda_{\rm sc}$ comes from a Weyl-invariant quadratic form Q.

We shall prove that the normality condition holds if and only if b and b_1 coincide over $\Lambda \otimes \Lambda_{\rm sc}$. This latter condition means that for each $\lambda \in \Lambda$ and root $\alpha \in \Lambda_{\rm sc}$, there holds:

$$
b(\lambda, \alpha) = Q(\alpha)\langle \check{\alpha}, \lambda \rangle.
$$
 (4.34)

The equality (4.34) is equivalent to the Weyl-invariance of b.

Let us now analyze the normality condition. The section induced from φ has the following description on S-points:

$$
\mathscr{L}T_{sc} \to \mathscr{G}_{sc} \tilde{\rtimes} \mathscr{T}, \quad t_{sc} \mapsto (\tilde{t}_{sc}, \varphi(\tilde{t}_{sc})^{-1}), \tag{4.35}
$$

where $\tilde{t}_{\rm sc}$ is any lift of $t_{\rm sc}$ to $\mathscr{G}_{\rm sc}$ and $\varphi(\tilde{t}_{\rm sc})$ is its image in \mathscr{T} under φ . Since $\mathscr{L}T_{\rm sc}$ is a central subgroup of $\mathscr{L}G_{\text{sc}} \rtimes \mathscr{L}T$, the image of [\(4.35\)](#page-43-2) is normal if and only if it is central. This condition translates to the following equality in $\mathscr{G}_{\mathrm{sc}}\tilde{\mathfrak{A}}\mathscr{T}$:

$$
(g_{\rm sc}, t) \cdot (\tilde{t}_{\rm sc}, \varphi(\tilde{t}_{\rm sc})^{-1}) \cdot (g_{\rm sc}, t)^{-1} = (\tilde{t}_{\rm sc}, \varphi(\tilde{t}_{\rm sc})^{-1}), \tag{4.36}
$$

for all S-points $(g_{\rm sc}, t)$ of $\mathscr{L}G_{\rm sc} \rtimes \mathscr{L}T$ and $t_{\rm sc}$ of $\mathscr{L}T_{\rm sc}$.

The left-hand-side of [\(4.36\)](#page-43-3) computes to $(\langle (t, \tilde{t}_{\rm sc})_{b_1} \tilde{t}_{\rm sc}, \langle t, \varphi(\tilde{t}_{\rm sc})^{-1} \rangle_b \varphi(\tilde{t}_{\rm sc})^{-1})$. Its equality with the right-hand-side amounts to the equality:

$$
\langle t,\tilde{t}_{\rm sc}\rangle_{b_1}=\langle t,\varphi(\tilde{t}_{\rm sc})\rangle_b,
$$

for all S-points t of \mathscr{L} T and t_{sc} of \mathscr{L} T_{sc}, *i.e.* the agreement of b and b_1 over $\Lambda \otimes \Lambda_{\text{sc}}$.

4.4.11. Define the Picard groupoid $\widetilde{\theta}_{G,+}^{super}(\Lambda)$ by the Cartesian diagram:

$$
\begin{aligned}\n\widetilde{\vartheta}_{G,+}^{\text{super}}(\Lambda) &\rightarrow \text{Quad}(\Lambda_{\text{sc}}, \mathbb{Z})^{\text{W}} \\
&\downarrow \qquad \qquad \downarrow (4.23) \\
\theta_{+}^{\text{super}}(\Lambda) &\rightarrow \theta_{+}^{\text{super}}(\Lambda_{\text{sc}})\n\end{aligned}
$$

Then $\vartheta_{G,+}^{super}(\Lambda)$ can be viewed as the full subgroupoid of $\tilde{\vartheta}_{G,+}^{super}(\Lambda)$ consisting of objects whose images in $\theta_{+}^{super}(\Lambda)$ have a *Weyl-invariant* form *b*.

Pulling back along $T \subset G$, $G_{\rm sc} \subset G$, and using the compatibility over $T_{\rm sc}$, we obtain a functor:

$$
\Gamma_{\text{fact}}(\text{Gr}_G, \text{Pic}^{\text{super}}) \to \tilde{\vartheta}_{G,+}^{\text{super}}(\Lambda). \tag{4.37}
$$

4.4.12. We are now ready to establish the equivalence $(1) \approx (2) \approx (3)$ in Theorem [3.4.5.](#page-31-0) We shall do so using the equivalence of §[4.4.7,](#page-43-0) together with an argument from [\[TZ21\]](#page-48-1) showing that factorization super line bundles over Gr_G embed fully faithfully into $\widetilde{\vartheta}_{G,+}^{\text{super}}(\Lambda)$.

Denote by:

$$
\mathrm{Hom}^{\mathrm{tame}}_{\mathrm{fact}}(\mathscr{L} G, \mathrm{Pic}^{\mathrm{super}}) \subset \mathrm{Hom}_{\mathrm{fact}}(\mathscr{L} G, \mathrm{Pic}^{\mathrm{super}})
$$

the full subgroupoid of factorization super central extensions of $\mathscr{L}G$ by $\mathbb{G}_{m, Ran}$, characterized by the property of having tame commutator.

Proposition 4.4.13. Let G be a reductive group X-scheme equipped with a maximal torus T with sheaf of cocharacters Λ . The functors [\(4.4\)](#page-34-2) and [\(4.37\)](#page-44-2) induce equivalences among the following Picard groupoids:

$$
\text{Hom}^{\text{tame}}_{\text{fact}}(\mathscr{L}\mathbf{G}, \text{Pic}^{\text{super}}) \xrightarrow{\simeq} \Gamma_{\text{fact}}(\text{Gr}_{\mathbf{G}}, \text{Pic}^{\text{super}}) \xrightarrow{\simeq} \vartheta_{\mathbf{G},+}^{\text{super}}(\Lambda). \tag{4.38}
$$

Proof. The functors [\(4.4\)](#page-34-2) and [\(4.37\)](#page-44-2) a priori define:

$$
\mathrm{Hom}^{\mathrm{tame}}_{\mathrm{fact}}(\mathscr{L} G, \mathrm{Pic}^{\mathrm{super}}) \to \Gamma_{\mathrm{fact}}(\mathrm{Gr}_G, \mathrm{Pic}^{\mathrm{super}}) \to \widetilde{\vartheta}^{\mathrm{super}}_{G,+}(\Lambda),
$$

The composition is an equivalence onto $\vartheta_{G,+}^{super}(\Lambda)$ —this is the equivalence of §[4.4.7](#page-43-0) restricted to the subgroupoid characterized by the tameness condition, as we see from Lemma [4.4.8](#page-43-4) and Lemma [4.4.10.](#page-43-5) Hence it suffices to prove that [\(4.37\)](#page-44-2) is fully faithful and its essential image is contained in $\vartheta_{G,+}^{super}(\Lambda)$.

The fully faithfulness is the content of [\[TZ21,](#page-48-1) §3.2]. The assertion that its image lies in $\vartheta_{G,+}^{super}(\Lambda)$ amounts to establishing the Weyl-invariance of the bilinear form b associated to any factorization super line bundle over Gr_G . Since $Rad(G) \times T_{sc} \rightarrow T$ is a Weyl-equivariant isogeny of X-tori, the statement can be proved when G is replaced by $Rad(G) \times G_{sc}$. In this case, we claim that the external product:

$$
\boxtimes: \Gamma_{\mathrm{fact}}(\mathrm{Gr}_{\mathrm{Rad}(G)}, \mathrm{Pic}^{\mathrm{super}}) \times \Gamma_{\mathrm{fact}}(\mathrm{Gr}_{G_{\mathrm{sc}}}, \mathrm{Pic}) \rightarrow \Gamma_{\mathrm{fact}}(\mathrm{Gr}_{\mathrm{Rad}(G) \times G_{\mathrm{sc}}}, \mathrm{Pic}^{\mathrm{super}})
$$

is an equivalence of Picard groupoids. Indeed, this follows from the fiberwise characterization of line bundles over $Gr_{Rad(G)\times G_{sc}}$ which descend to $Gr_{Rad(G)}$ (§[4.3.5\)](#page-38-3).

4.5. Poor man's transgression.

4.5.1. Let G be a reductive group X-scheme. Denote by BG the X-stack classifying Zariski locally trivial G-torsors.

The following result completes the equivalence (1) \cong (4) in Theorem [3.4.5.](#page-31-0)

Proposition 4.5.2. Fix a ϑ -characteristic $\omega^{1/2}$ over X. There is a canonical equivalence of Picard groupoids:

$$
\int_{(\mathring{D},\omega^{1/2})} : \Gamma_e(BG, \underline{K}_{[1,2]}^{\text{super}}) \xrightarrow{\simeq} \text{Hom}^{\text{tame}}_{\text{fact}}(\mathscr{L}G, \text{Pic}^{\text{super}}). \tag{4.39}
$$

Furthermore, if G is equipped with a maximal torus T with sheaf of cocharacters Λ . Then the following diagram is canonically commutative:

$$
\Gamma_e(\text{BG}, \underline{\underline{K}}^{\text{super}}_{[1,2]}) \xrightarrow{\int (\underline{\underline{b}}, \underline{\underline{u}}^{1/2})} \text{Hom}^{\text{tame}}_{\text{fact}}(\mathscr{L}\text{G}, \text{Pic}^{\text{super}}) \xrightarrow{\downarrow} (4.38)
$$
\n
$$
\varphi_G^{\text{super}}(\Lambda) \xrightarrow{\qquad (3.22)} \vartheta_{\text{G},+}^{\text{super}}(\Lambda) \tag{4.40}
$$

4.5.3. In an ideal world, we would define [\(4.39\)](#page-45-0) by a "transgression" on K-theory and verify the commutativity of [\(4.40\)](#page-45-1), but we do not know how to do so.

In what follows, we offer a poor man's substitute: we first fix a maximal torus contained in Borel subgroup $T \subset B \subset G$ (which exists étale locally on X) and *define* [\(4.39\)](#page-45-0) as the com-position of the three equivalences in [\(4.40\)](#page-45-1). We denote this functor by Φ _(T,B) to emphasize its *a priori* dependence on (T, B) .

Then we prove that $\Phi_{(T,B)}$ is canonically independent of (T,B) , *i.e.* given two pairs $T_i \in B_i \subset G$ (for $i = 1, 2$) of maximal tori contained in Borel subgroups, there is a canonical isomorphism of functors:

$$
\alpha_{(T_1,B_1),(T_2,B_2)} : \Phi_{(T_1,B_1)} \xrightarrow{\simeq} \Phi_{(T_2,B_2)}.
$$
\n(4.41)

satisfiying the cocycle condition for three such pairs (T_i, B_i) $(i = 1, 2, 3)$.

These canonical isomorphisms allow us to glue the functors $\Phi_{(\text{T},B)}$ over an étale cover of X, which yields the equivalence [\(4.39\)](#page-45-0).

Remark 4.5.4. The definition of $\Phi_{(T,B)}$ uses only T. However, the isomorphism [\(4.41\)](#page-45-2) depends on B_1 and B_2 , so we prefer keep the Borel subgroups in the notation.

Proof of Proposition [4.5.2.](#page-45-3) It suffices to construct [\(4.41\)](#page-45-2) satisfying the cocycle condition.

Suppose that t is an X-point of T. The inner automorphism int_t : G \rightarrow G, $g \mapsto tgt^{-1}$ preserves $T \subset G$ and induces a commutative diagram:

$$
\Gamma_e(\text{BG}, \underline{\underline{K}}^{\text{super}}_{[1,2]}) \stackrel{(2.12)}{\longrightarrow} \vartheta^{\text{super}}_G(\Lambda)
$$
\n
$$
\downarrow \text{int}^*_{t} \qquad \qquad \downarrow \text{int}^*_{t}
$$
\n
$$
\Gamma_e(\text{BG}, \underline{\underline{K}}^{\text{super}}_{[1,2]}) \stackrel{(2.12)}{\longrightarrow} \vartheta^{\text{super}}_G(\Lambda)
$$
\n(4.42)

Rigidified sections of $\underline{K}^{\text{super}}_{[1,2]}$ over BG are equivalent to monoidal functors $G \to \Omega(\underline{K}^{\text{super}}_{[1,2]}$ $_{\left[1,2\right]}^{\text{super}}$). Since the target has a *symmetric* monoidal structure, int_t^* acts as the identity on the groupoid of such monoidal functors.

On the other hand, the right vertical functor int_t^* in [\(4.42\)](#page-45-4) carries a triple $(b, \tilde{\Lambda}, \varphi)$ to the triple $(b, \tilde{\Lambda}, \varphi \cdot t^b)$, where $t^b : \Lambda \to \mathbb{G}_m$ denotes the character sending $\lambda \in \Lambda$ to the character $b(\lambda,-): \mathcal{T} \to \mathbb{G}_m$ evaluated at t, and $\varphi \cdot t^b$ is the sum of φ with the restriction of t^b to $\Lambda_{\rm sc}$. The 2-isomorphism rendering [\(4.42\)](#page-45-4) commutative evalutes to the isomorphism:

$$
(b,\widetilde{\Lambda},\varphi)\xrightarrow{\simeq}\text{int}^*_t(b,\widetilde{\Lambda},\varphi)\xrightarrow{\simeq}(b,\widetilde{\Lambda},\varphi\cdot t^b),
$$

induced by the automorphism of the central extension $\widetilde{\Lambda}$ defined by t^b . These calculations are performed using the description of the commutator in [\[BD01,](#page-46-1) Proposition 3.13].

The situation is parallel for the functor (4.38) : an X-point t of T induces a commutative diagram:

$$
\text{Hom}^{\text{tame}}_{\text{fact}}(\mathscr{L}\mathbf{G}, \text{Pic}^{\text{super}}) \stackrel{(4.38)}{\longrightarrow} \vartheta^{\text{super}}_{\mathbf{G},+}(\Lambda)
$$
\n
$$
\downarrow^{\text{int}^*}_{\text{int}^*_{\mathbf{G}}} \downarrow^{\text{int}^*}_{\text{int}^*_{\mathbf{G}}}(\Lambda)
$$
\n
$$
\text{Hom}^{\text{tame}}_{\text{fact}}(\mathscr{L}\mathbf{G}, \text{Pic}^{\text{super}}) \stackrel{(4.38)}{\longrightarrow} \vartheta^{\text{super}}_{\mathbf{G},+}(\Lambda)
$$
\n
$$
(4.43)
$$

where the left vertical functor is isomorphic to the identity, the right vertical functor sends $(b,\widetilde{\Lambda}_{+},\varphi)$ to $(b,\widetilde{\Lambda}_{+},\varphi \cdot t^{b})$, and the 2-isomorphism rendering [\(4.43\)](#page-46-5) commutative is given by the automorphism t^b of $\widetilde{\Lambda}_+$. These calculations are performed using the description of the commutator in §[4.2.8.](#page-37-4)

Suppose now that $T_i \subset B_i \subset G$ (for $i = 1, 2$) are a pair of maximal tori contained in Borel subgroups and g is an X-point of G with $\text{int}_q(T_2) = T_1$, $\text{int}_q(B_2) = B_1$. Denote by Λ_i the sheaf of cocharacters of T_i .

The inner automorphism \int_{q} gives rise to a commutative diagram:

$$
\Gamma_e(BG, \underline{K}_{[1,2]}^{\text{super}}) \stackrel{(2.12)}{\longrightarrow} \vartheta_G^{\text{super}}(\Lambda_1) \simeq \vartheta_{G,+}^{\text{super}}(\Lambda_1) \stackrel{(4.38)}{\longleftarrow} \text{Hom}_{\text{fact}}^{\text{tame}}(\mathscr{L}G, \text{Pic}^{\text{super}})
$$
\n
$$
\downarrow \text{int}_{g}^* \qquad \qquad \downarrow \text{int}_{g}^* \qquad \qquad \downarrow \text{int}_{g}^* \qquad \qquad \downarrow \text{int}_{g}^* \qquad (4.44)
$$
\n
$$
\Gamma_e(BG, \underline{K}_{[1,2]}^{\text{super}}) \stackrel{(2.12)}{\longrightarrow} \vartheta_G^{\text{super}}(\Lambda_2) \simeq \vartheta_{G,+}^{\text{super}}(\Lambda_1) \stackrel{(4.38)}{\longleftarrow} \text{Hom}_{\text{fact}}^{\text{tame}}(\mathscr{L}G, \text{Pic}^{\text{super}})
$$

where the middle equivalences are defined by $\omega^{1/2}$ -shift [\(3.22\)](#page-31-1). Since the outer vertical functors are equivalent to the identity, the 2-isomorphism rendering [\(4.44\)](#page-46-6) commutative defines an isomorphism of functors:

$$
\alpha_g: \Phi_{(\mathrm{T}_1, \mathrm{B}_1)} \xrightarrow{\simeq} \Phi_{(\mathrm{T}_2, \mathrm{B}_2)}.\tag{4.45}
$$

Claim: α_g depends only on the pairs (T_1, B_1) and (T_2, B_2) (as opposed to g). Indeed, any other choice of an X-point of G conjugating (T_2, B_2) into (T_1, B_1) differs from g by an X-point of T_1 , so the claim is equivalent to the following assertion: for (T_1, B_1) = $(T_2, B_2) = (T, B)$, the isomorphism α_t is the identity on $\Phi_{(T,B)}$. However, this follows from the description of the 2-isomorphisms in [\(4.42\)](#page-45-4) and [\(4.43\)](#page-46-5).

Finally, we set [\(4.41\)](#page-45-2) to be the isomorphism [\(4.45\)](#page-46-7) for any X-point g conjugating (T_2, B_2) into (T_1, B_1) , which exists étale locally over X—these choices glue thanks to the independence of α_g on g. The cocycle condition follows from the equality $\alpha_{g_1g_2} = \alpha_{g_1} \cdot \alpha_{g_2}$ \Box

REFERENCES

- [ABD+66] Michael Artin, Jean-Etienne Bertin, Michel Demazure, Alexander Grothendieck, Pierre Gabriel, Michel Raynaud, and Jean-Pierre Serre, Schémas en groupes. Séminaire de géométrie algébrique du Bois Marie 1962–64 (SGA 3), Séminaire de Géométrie Algébrique de l'Institut des Hautes $\text{Études Scientificues}, \text{Institut des Hautes Études Scientificues}, \text{Paris}, 1963/1966.}$
- [APR04] Greg W. Anderson and Fernando Pablos Romo, Simple proofs of classical explicit reciprocity laws on curves using determinant groupoids over an Artinian local ring, Comm. Algebra 32 (2004), no. 1, 79–102. MR 2036223
- [BD91] Alexander Beilinson and Vladimir Drinfeld, Quantization of Hitchin's integrable system and Hecke eigensheaves, <https://math.uchicago.edu/~drinfeld/langlands/QuantizationHitchin.pdf>, 1991.
- [BD01] Jean-Luc Brylinski and Pierre Deligne, Central extensions of reductive groups by \mathbf{K}_2 , Publ. Math. Inst. Hautes Etudes Sci. (2001), no. 94, 5–85. MR 1896177 ´

- [BD04] Alexander Beilinson and Vladimir Drinfeld, Chiral algebras, American Mathematical Society Colloquium Publications, vol. 51, American Mathematical Society, Providence, RI, 2004. MR 2058353
- [BDF+22] Alexander Braverman, Gurbir Dhillon, Michael Finkelberg, Sam Raskin, and Roman Travkin, Coulomb branches of noncotangent type (with appendices by Gurbir Dhillon and Theo Johnson-Freyd), <https://arxiv.org/abs/2201.09475>, 2022.
- [BGT13] Andrew J. Blumberg, David Gepner, and Gonçalo Tabuada, A universal characterization of higher algebraic K-theory, Geom. Topol. 17 (2013), no. 2, 733–838. MR 3070515
- [BL94] Arnaud Beauville and Yves Laszlo, Conformal blocks and generalized theta functions, Comm. Math. Phys. 164 (1994), no. 2, 385–419. MR 1289330
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, Néron models, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990. MR 1045822
- [BS17] Bhargav Bhatt and Peter Scholze, Projectivity of the Witt vector affine Grassmannian, Invent. Math. 209 (2017), no. 2, 329–423. MR 3674218
- [CH21] Justin Campbell and Andreas Hayash, Geometric class field theory and cartier duality, <https://arxiv.org/abs/1710.02892>, 2021.
- [Cho22] Chang-Yeon Chough, Spectral excision and descent for perfect complexes, <https://arxiv.org/abs/2210.13679>, 2022.
- [CTS87] Jean-Louis Colliot-Thélène and Jean-Jacques Sansuc, Principal homogeneous spaces under flasque tori: applications, J. Algebra 106 (1987), no. 1, 148–205. MR 878473
- [Dri06] Vladimir Drinfeld, Infinite-dimensional vector bundles in algebraic geometry: an introduction, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 263– 304. MR 2181808
- [DS95] Vladimir Drinfeld and Carlos Simpson, B-structures on G-bundles and local triviality, Math. Res. Lett. 2 (1995), no. 6, 823–829. MR 1362973
- [DW90] Robbert Dijkgraaf and Edward Witten, Topological gauge theories and group cohomology, Comm. Math. Phys. 129 (1990), no. 2, 393–429. MR 1048699
- [EKLV98] Hélène Esnault, Bruno Kahn, Marc Levine, and Eckart Viehweg, The Arason invariant and mod 2 algebraic cycles, J. Amer. Math. Soc. 11 (1998), no. 1, 73–118. MR 1460391
- [Fal03] Gerd Faltings, Algebraic loop groups and moduli spaces of bundles, J. Eur. Math. Soc. (JEMS) 5 (2003), no. 1, 41–68. MR 1961134
- [FL10] Michael Finkelberg and Sergey Lysenko, Twisted geometric Satake equivalence, J. Inst. Math. Jussieu 9 (2010), no. 4, 719–739. MR 2684259
- [GA13] Cristian D. González-Avilés, Flasque resolutions of reductive group schemes, Cent. Eur. J. Math. 11 (2013), no. 7, 1159–1176. MR 3085137
- [Gai20] Dennis Gaitsgory, Parameterization of factorizable line bundles by K-theory and motivic cohomology, Selecta Math. (N.S.) 26 (2020), no. 3, Paper No. 44, 50. MR 4117995
- [GD67] A. Grothendieck and J. A. Dieudonné, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas, Quatrième partie, Inst. Hautes Études Sci. Publ. Math. (1967), no. 32, 361. MR 238860
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus, Universality of multiplicative infinite loop space machines, Algebr. Geom. Topol. 15 (2015), no. 6, 3107–3153. MR 3450758
- [GL18] Dennis Gaitsgory and Sergey Lysenko, Parameters and duality for the metaplectic geometric Langlands theory, Selecta Math. (N.S.) 24 (2018), no. 1, 227–301, References are to the corrected version, available at: <https://lysenko.perso.math.cnrs.fr>. MR 3769731
- [GL19] Dennis Gaitsgory and Jacob Lurie, Weil's conjecture for function fields. Vol. 1, Annals of Mathematics Studies, vol. 199, Princeton University Press, Princeton, NJ, 2019. MR 3887650
- [GO15] S. O. Gorchinskiĭ and D. V. Osipov, A higher-dimensional Contou-Carrère symbol: local theory, Mat. Sb. 206 (2015), no. 9, 21–98. MR 3438594
- [Gör03] Ulrich Görtz, On the flatness of local models for the symplectic group, Adv. Math. 176 (2003), no. 1, 89–115. MR 1978342
- [GR14] Dennis Gaitsgory and Nick Rozenblyum, DG indschemes, Perspectives in representation theory, Contemp. Math., vol. 610, Amer. Math. Soc., Providence, RI, 2014, pp. 139–251. MR 3220628
- [Hen17] André Henriques, The classification of chiral WZW models by $H^4_+(\text{BG}, \mathbb{Z})$, Lie algebras, vertex operator algebras, and related topics, Contemp. Math., vol. 695, Amer. Math. Soc., Providence, RI, 2017, pp. 99–121. MR 3709708
- [Kem76] George R. Kempf, Linear systems on homogeneous spaces, Ann. of Math. (2) 103 (1976), no. 3, 557–591. MR 409474
- [Kov22] Sándor J Kovács, Rational singularities, <https://arxiv.org/abs/1703.02269>, 2022.
- [KV04] Mikhail Kapranov and Eric Vasserot, Vertex algebras and the formal loop space, Publ. Math. Inst. Hautes Etudes Sci. (2004), no. 100, 209–269. MR 2102701 ´
- [KV07] Mikhail Kapranov and Éric Vasserot, Formal loops. II. A local Riemann-Roch theorem for determinantal gerbes, Ann. Sci. École Norm. Sup. (4) 40 (2007) , no. 1, 113–133. MR 2332353
- [Laf18] Vincent Lafforgue, Chtoucas pour les groupes réductifs et paramétrisation de Langlands globale, J. Amer. Math. Soc. 31 (2018), no. 3, 719–891. MR 3787407
- [Lur17] Jacob Lurie, Higher algebra, <https://people.math.harvard.edu/~lurie/papers/HA.pdf>, 2017, preprint.
- [Lur18] Jacob Lurie, Kerodon, <https://kerodon.net>, 2018.
- [Lys16] Sergey Lysenko, Twisted geometric Satake equivalence: reductive case, <https://arxiv.org/abs/1411.6782>, 2016.
- [Lys17] Sergey Lysenko, Twisted Whittaker models for metaplectic groups, Geom. Funct. Anal. 27 (2017), no. 2, 289–372. MR 3626614
- [PR11] Bjorn Poonen and Eric Rains, Self cup products and the theta characteristic torsor, Math. Res. Lett. 18 (2011), no. 6, 1305–1318. MR 2915483
- [Roz21] Nick Rozenblyum, Connections on moduli spaces and infinitesimal Hecke modifications, <https://arxiv.org/abs/2108.07745>, 2021.
- [Sta18] The Stacks Project Authors, Stacks Project, <https://stacks.math.columbia.edu>, 2018.
- [Tao21a] James Tao, $\text{Gr}_{G,\text{Ran}(X)}$ is reduced, <https://arxiv.org/abs/2011.01553>, 2021.
- [Tao21b] James Tao, n-excisive functors, canonical connections, and line bundles on the Ran space, Selecta Math. (N.S.) 27 (2021), no. 1, Paper No. 2, 75. MR 4198527
- [Tat68] John Tate, Residues of differentials on curves, Ann. Sci. École Norm. Sup. (4) 1 (1968), 149–159. MR 227171
- [TT90] R. W. Thomason and Thomas Trobaugh, Higher algebraic K-theory of schemes and of derived categories, The Grothendieck Festschrift, Vol. III, Progr. Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, pp. 247–435. MR 1106918
- [TUY89] Akihiro Tsuchiya, Kenji Ueno, and Yasuhiko Yamada, Conformal field theory on universal family of stable curves with gauge symmetries, Integrable systems in quantum field theory and statistical mechanics, Adv. Stud. Pure Math., vol. 19, Academic Press, Boston, MA, 1989, pp. 459–566. MR 1048605
- [TZ21] James Tao and Yifei Zhao, Central extensions by \mathbf{K}_2 and factorization line bundles, Math. Ann. 381 (2021), no. 1-2, 769–805. MR 4322626
- [Wal17] Konrad Waldorf, Transgressive loop group extensions, Math. Z. 286 (2017), no. 1-2, 325–360. MR 3648501
- [Wei11] Martin H. Weissman, Managing metaplectiphobia: covering p-adic groups, Harmonic analysis on reductive, p-adic groups, Contemp. Math., vol. 543, Amer. Math. Soc., Providence, RI, 2011, pp. 237–277. MR 2798431
- [Zhu09] Xinwen Zhu, Affine Demazure modules and T-fixed point subschemes in the affine Grassmannian, Adv. Math. 221 (2009), no. 2, 570–600. MR 2508931
- [Zhu17] , An introduction to affine Grassmannians and the geometric Satake equivalence, Geometry of moduli spaces and representation theory, IAS/Park City Math. Ser., vol. 24, Amer. Math. Soc., Providence, RI, 2017, pp. 59–154. MR 3752460