Thermodynamic bounds on generalized transport: From single-molecule to bulk observables

Cai Dieball¹ and Aljaž Godec^{1, *}

¹Mathematical bioPhysics Group, Max Planck Institute for Multidisciplinary Sciences, 37077 Göttingen, Germany

We prove that the transport of any differentiable scalar observable in *d*-dimensional nonequilibrium systems is bounded from above by the total entropy production scaled by the amount the observation "stretches" microscopic coordinates. The result—a time-integrated generalized speed limit—reflects the thermodynamic cost of transport of observables, and places underdamped and overdamped stochastic dynamics on equal footing with deterministic motion. Our work allows for stochastic thermodynamics to make contact with bulk experiments, and fills an important gap in thermodynamic inference, since microscopic dynamics is, at least for short times, underdamped. Requiring only averages but not sample-to-sample fluctuations, the proven transport bound is practical and applicable not only to single-molecule but also bulk experiments where only averages are observed, which we demonstrate by examples. Our results may facilitate thermodynamic inference on molecular machines without an obvious directionality from bulk observations of transients probed, e.g. in time-resolved X-ray scattering.

A complete thermodynamic characterization and understanding of systems driven far from equilibrium remains elusive. Central to non-equilibrium thermodynamics is the total entropy production ΔS_{tot} , which reflects the displacement from equilibrium [1]. ΔS_{tot} embodies the entropy change in both, the system and the environment coupled to it, and is a measure of the violation of time-reversal symmetry [1, 2]. Most importantly, the thermodynamic cost of non-equilibrium processes (stationary, transient, or even explicitly time-dependent) at temperature T is in fact $T\Delta S_{tot}$, which can thus be seen as the "counterpart" of free energy in equilibrium. The total entropy production allows, for example, to quantify the efficiency of molecular motors [3-5] and gain insight into the energetic budget of human red blood cells [6]. While most works focus on non-equilibrium steady-state (NESS) dynamics, transient processes that approach equilibrium states such as, e.g., protein folding [7-15] or thermal relaxation [16], are also characterized by $T\Delta S_{\rm tot}$ which then corresponds to an excess free energy [16–19]. Even more involved are transients towards NESS, relevant in e.g., the packaging of viral DNA [20], red blood cell flickering [21], enzymatically facilitated topological relaxation of DNA [22], or nanoparticle model systems for interrogating the fundamental laws of stochastic thermodynamics [23].

Despite its importance, the inference of ΔS_{tot} from experimental observations is far from simple, as it requires access to *all* dissipative degrees of freedom is the system, which is typically precluded by the fact that one only has access to some observable. Notably, neither the microscopic dynamics nor the projection underlying the observable are typically known, and often one can only observe transients.

To overcome these intrinsic limitations of experiments, diverse bounds (i.e., inequalities) on the entropy production have been derived, in particular thermodynamic uncertainty relations (TURs) [24–37] and speed limits [38– 48]. Such bounds provide conceptual insight about manifestations of irreversible behavior, and from a practical perspective they allow to infer a bound on ΔS_{tot} from measured trajectories, more precisely from the sampleto-sample fluctuations or speed of observables.

These results remain incomplete from several perspectives. First, their validity typically hinges on the assumption that the microscopic dynamics is overdamped or even a Markov-jump process. This is unsatisfactory because microscopic dynamics is, at least on short time scales, underdamped and the TUR does not hold for underdamped dynamics [49] (see, however, progress in [50– 53] and recently [54]). Similarly, thermodynamic speed limits have so far seemingly not been derived for underdamped dynamics. While inertial effects may not be important in colloidal systems [55], they are indeed relevant for, e.g., protein dynamics [56–59], and are known to invalidate overdamped theories even on long time scales [49]. Even the basic formulation of path-wise thermodynamics of inertial systems is fundamentally different, and in the analysis of densities or integrals over stochastic trajectories one cannot eliminate short-time inertial effects. Second, dissipative processes are often, especially in molecular machines without an obvious directionality (e.g. molecular chaperones [60]), mediated by intricate collective (and often fast) open-close motions visible in transients that are difficult to resolve even with advanced single-molecule techniques [61]. Experiments providing more detailed structural information, such as time-resolved X-ray scattering techniques [7–10, 12–15], are available, but probe bulk behavior for which the existing bounds do not apply. There is thus a pressing need to close the gaps and to cover underdamped dynamics and tap into bulk observations.

Here, we present thermodynamic bounds on the generalized transport of observables in systems evolving according to (generally time-inhomogeneous) over-

^{*} agodec@mpinat.mpg.de

damped, underdamped, or even deterministic dynamics, all treated on an equal footing, generalizing existing results on transport [4, 46]. Technically, the results may be classified as a time-integrated version of generalized speed limits and bring several conceptual and practical advantages. As a demonstration, we use the bounds for thermodynamic inference based on both, single-molecule and bulk (e.g. scattering) observables.

Rationale.—Consider the simplest case of a Newtonian particle with position x_{τ} and velocity v_{τ} at time τ dragged through a viscous medium against the (Stokes) friction force $F_{\gamma} = -\gamma v_{\tau}$ with friction constant γ causing a transfer of energy into the medium. The dissipated heat between times 0 and t is $\Delta Q_{\gamma} = \gamma \int_0^t v_{\tau} dx_{\tau} = \gamma \int_0^t v_{\tau}^2 d\tau$ and gives rise to entropy production in the medium [1]. Since deterministic dynamics does not produce entropy otherwise, we have $T\Delta S_{\text{tot}} = \Delta Q_{\gamma}$ in $\tau \in [0, t]$. This imposes a thermodynamic bound on transport via the Cauchy-Schwarz inequality $(x_t - x_0)^2 = (\int_0^t v_{\tau} 1 d\tau)^2 \leq \int_0^t v_{\tau}^2 d\tau \int_0^t 1^2 d\tau'$, yielding $T\Delta S_{\text{tot}} \geq \gamma (x_t - x_0)^2 / t$ with equality for constant velocity. Therefore, for given t and γ a minimum energy input ΔQ_{γ} is required to achieve a displacement $x_t - x_0$. The intuition that transport requires dissipation extends to general dynamics and scalar observables as follows.

Main result.—The transport of any differentiable scalar observable $z_{\tau} \equiv z(\mathbf{x}_{\tau}, \tau)$ (see [62] for $z_{\tau} \equiv z(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau)$) on a time interval [0, t] in *d*-dimensional generally underdamped and time-inhomogeneous dynamics $(\mathbf{x}_{\tau}, \mathbf{v}_{\tau})$ is bounded from above by $T\Delta S_{\text{tot}}$ as

$$T\Delta S_{\text{tot}} \geq \frac{\left\langle z_t - z_0 - \int_0^t \partial_\tau z_\tau d\tau \right\rangle^2}{t\mathcal{D}^z(t)}$$
$$\mathcal{D}^z(t) \equiv \frac{1}{t} \int_0^t \langle [\nabla_{\mathbf{x}} z_\tau]^T \boldsymbol{\gamma}^{-1}(\tau) \nabla_{\mathbf{x}} z_\tau \rangle d\tau, \qquad (1)$$

where γ is a positive definite, possibly time-dependent, symmetric friction matrix, $\langle \cdot \rangle$ denotes an ensemble average over non-stationary trajectories, and $\mathcal{D}^{z}(t)$ is a fluctuation-scale function of the observable that determines how much the observation z stretches microscopic coordinates \mathbf{x} . While ΔS_{tot} for stochastic dynamics differs from $\Delta Q_{\gamma}/T$, the bound (1) remains valid in the whole spectrum from Newtonian to underdamped and overdamped stochastic dynamics. The inequality saturates for $\nabla_{\mathbf{x}} z_{\tau} = c \gamma \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau)$ for any constant c and $\boldsymbol{\nu}$ defined in Eq. (4). Note that the application of Eq. (1) does not require knowledge of the parameters of the underlying motion. One only needs to infer the average in the numerator and \mathcal{D}^{z} . To infer the latter, different strategies are presented towards the end of the Letter.

Setting $z_{\tau} = x_{\tau}$ for d = 1, Eq. (1) includes the above deterministic case. In the overdamped limit, Eq. (1) complements the Benamou-Brenier formula that bounds transport in terms of a Wasserstein distance [48, 63], and special cases of the bound correspond to existing overdamped speed limits [4, 5, 46] (in particular, Ref. [46] already contains an overdamped version of \mathcal{D}^z , see [62] for a detailed connection to the existing literature).

The bound (1) characterizes the thermodynamic cost of transport and may be employed in thermodynamic inference. By only requiring the mean but not sampleto-sample fluctuations, the bound (1) is simpler than the TUR and allows to infer ΔS_{tot} from transients of bulk observables, probed, e.g., in time-resolved scattering experiments [7–10, 12–15, 64]. A disadvantage of this simplicity is that it is not useful for stationary states, unless they are translation invariant or periodic. The observable z_{τ} can represent a measured projection, whose functional form may be known (e.g., in X-ray scattering) or unknown (e.g., a reaction coordinate of a complex process). For optimization of thermodynamic inference z_{τ} may be chosen a posteriori and τ -dependent.

Outline.—First we describe the setup and discuss different notions of ΔS_{tot} from deterministic via underdamped to overdamped dynamics. Next we present examples in the context of single-molecule versus bulk Xray scattering experiments, as well as higher-order transport in stochastic heat engines. We then explain how to interpret and infer the fluctuation-scale function $\mathcal{D}^{z}(t)$. We conclude with a perspective.

Setup.—Let γ , **m** be $d \times d$ positive definite, symmetric friction and mass matrices with square root $\sqrt{\gamma}\sqrt{\gamma}^T = \gamma$. The full dynamics $\mathbf{x}_{\tau}, \mathbf{v}_{\tau} \in \mathbb{R}^d$ evolve according to [65]

$$d\mathbf{x}_{\tau} = \mathbf{v}_{\tau} d\tau$$
(2)
$$d\mathbf{v}_{\tau} = \mathbf{m}^{-1} \left[\mathbf{F}(\mathbf{x}_{\tau}, \tau) d\tau - \gamma \mathbf{v}_{\tau} d\tau + \sqrt{2k_{\rm B}T\gamma} d\mathbf{W}_{\tau} \right],$$

which in the overdamped limit reduce to

$$d\mathbf{x}_{\tau}^{\mathrm{od}} = \boldsymbol{\gamma}^{-1} \mathbf{F}(\mathbf{x}_{\tau}^{\mathrm{od}}, \tau) d\tau + \sqrt{2k_{\mathrm{B}}T\boldsymbol{\gamma}^{-1}} d\mathbf{W}_{\tau}, \quad (3)$$

where $\mathbf{F}(\mathbf{x}, \tau)$ is a force field and \mathbf{W}_{τ} the *d*-dimensional Wiener process. We allow $\boldsymbol{\gamma}(\tau)$ (and later also $T(\tau)$) to depend on time but suppress this dependence to simplify notation. We define the *local mean velocity* $\boldsymbol{\nu}$ of the probability density ρ in phase space as $\boldsymbol{\nu}^{\text{Newton}} = \mathbf{v}$ and

$$\boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \equiv \mathbf{v} + \mathbf{m}^{-1} k_{\mathrm{B}} T \frac{\nabla_{\mathbf{v}} \rho(\mathbf{x}, \mathbf{v}, \tau)}{\rho(\mathbf{x}, \mathbf{v}, \tau)}$$
$$\boldsymbol{\nu}^{\mathrm{od}}(\mathbf{x}, \tau) \equiv \boldsymbol{\gamma}^{-1} \left[\mathbf{F}(\mathbf{x}, \tau) - k_{\mathrm{B}} T \frac{\nabla_{\mathbf{x}} \rho^{\mathrm{od}}(\mathbf{x}, \tau)}{\rho^{\mathrm{od}}(\mathbf{x}, \tau)} \right].$$
(4)

The definition for overdamped dynamics is standard [1], whereas in the underdamped setting $\boldsymbol{\nu}$ is only the "irreversible" part of the probability current divided by density [51]. Note that $\mathbf{v}_{\tau} = d\mathbf{x}_{\tau}/d\tau$ does not exist for overdamped dynamics as $\langle |d\mathbf{x}_{\tau}/d\tau| \rangle \rightarrow \infty$ for $d\tau \rightarrow 0$ but $\langle d\mathbf{x}_{\tau} \rangle/d\tau = \boldsymbol{\gamma}^{-1}\mathbf{F}(\mathbf{x},\tau)$ is well-behaved. The overdamped limit is loosely speaking obtained for $\boldsymbol{\gamma}^{-1}\mathbf{m} \rightarrow \mathbf{0}$, whereby details of this limit depend on \mathbf{F} [66].

There are two differences between Newtonian and stochastic dynamics: (i) the energy exchange between system and bath counteracts friction, and (ii) changes in ρ give rise to a change in Gibbs entropy of a stochastic




FIG. 1. Particle x_t in a harmonic trap displaced from x = 1 to x = 0 at time 0. (a) The particle's mean position $\langle x_0 \rangle = 1$ moves towards the new center of the trap, whereby oscillations occur for small damping. The probability density around $\langle x_\tau \rangle$ in this example is a Gaussian of constant width (see [62] for details and parameters). (b) Quality factor Q of the transport bound for the simplest observable $z_\tau = x_\tau$, and quality factor of the TUR for the current $J_t = x_t - x_0$ in the overdamped case. Full saturation Q = 1 at all times can be achieved for overdamped and underdamped dynamics as described below.

system, thus contributing to the total entropy production as $\Delta S_{\text{tot}} = \Delta S_{\text{sys}} + \Delta S_{\text{bath}}$. In all cases considered we can write the total entropy production as [1, 2, 51]

$$T\Delta S_{\text{tot}} = \int_0^t d\tau \langle \boldsymbol{\nu}^T(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \boldsymbol{\gamma} \boldsymbol{\nu}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \rangle \ge 0. \quad (5)$$

Within this setup, an *educated guess* and stochastic calculus alongside the Cauchy-Schwarz inequality delivers the announced bound (1) (see proof in [62]).

Example 1: Colloid in displaced trap.— Consider a bead trapped in a harmonic potential displaced from position 1 to 0 at time $\tau = 0$. Knowing γ and observing only the mean particle transport $\langle x_t - x_0 \rangle$ we infer the entropy production from Eq. (1) to be $T\Delta S_{\rm tot} \geq$ $T\Delta S_{\text{tot}}^{\text{bound}} \equiv \gamma \langle x_t - x_0 \rangle^2 / t$. We inspect the quality of the bound, $Q \equiv \Delta S_{\text{tot}}^{\text{bound}} / \Delta S_{\text{tot}} \in [0, 1]$, as a function of time (see Fig. 1). For underdamped dynamics Q tends to 3/4 at short times due to inertia (see [62] for derivation). Using $z_{\tau} = x\nu(\tau)$ (for this example, ν turns out to be independent of x, v) we achieve saturation for all times, which is easily understood from our proof (see [62]). Saturation for this example was also achieved via the transient correlation TUR [37]. However, the present approach is expected to be numerically more stable and requires less statistics (see also Fig. S1 in [62]), since a determination of fluctuations and derivatives of observables is not required. Moreover, the simplest version $z_{\tau} = x_{\tau}$ outperforms the transient TUR for the simplest current $J_t = x_t - x_0$ (dashed line in Fig. 1b). This may be interpreted as the magnitude of $x_t - x_0$ entering (1) being more relevant than its precision entering the TUR for this example. In contrast to the TUR [49], we also have the advantage that Eq. (1) holds for underdamped dynamics.

A disadvantage of Eq. (1) is that it is not useful for steady-state dynamics, since there $\langle z_t - z_0 \rangle = 0$. An exception are spatially periodic systems treated as NESS [3, 4]. A particular example are overdamped Brownian clocks [67] where for given ΔS_{tot} the TUR limits *precision*

Example 2: Scattering experiments.—Since Eq. (1) only requires averages, it is applicable beyond singlemolecule probes to bulk experiments, i.e., experiments on samples of many molecules probing mean properties, e.g., scattering techniques. The recent surge in the development of time-resolved X-ray scattering on proteins [7–15] renders our bound particularly useful. Here, transients may be excited by a pressure [68, 69] or temperature [70] quench, or one directly monitors slow kinetics [71]. One typically observes the structure factor $S(\mathbf{q}) \equiv \frac{1}{N} \sum_{j,k=1}^{N} \langle e^{-i\mathbf{q} \cdot (\mathbf{r}_{t}^{j} - \mathbf{r}_{t}^{k})} \rangle$ [55, 72], where the sum runs over all scatterers (atoms, particles, etc.). This also applies to interacting colloid suspensions, where $S(\mathbf{q})$ is the Fourier transform of the pair correlation function [55]. An even simpler observable is the radius of gyration, $R_g^2 \equiv \frac{1}{N} \sum_{j=1}^N \langle (\mathbf{r}_j - \bar{\mathbf{r}})^2 \rangle$ [11, 55, 73], where $\bar{\mathbf{r}} \equiv \frac{1}{N} \sum_{j=1}^N \mathbf{r}_j$ is the center of mass. R_g^2 reflects the (statistical) size of molecules and is easily inferred from small q via Guinier's law, $S(|\mathbf{q}|) \stackrel{|\mathbf{q}|\to 0}{=} S(0) e^{-|\mathbf{q}|^2 R_g^2/3}$ [55, 73].

We consider the structure factor averaged over spatial dimensions S(q) (see [62] for the vector version). For simplicity assume that γ is a known scalar. We observe how S(q) changes over time. From Eq. (1) we can derive the bounds (see [62])

$$T\Delta S_{\text{tot}} \ge \frac{3\gamma [S_t(q) - S_0(q)]^2}{q^2 \int_0^t [N - S_\tau(2q)] d\tau} \ge \frac{3\gamma [S_t(q) - S_0(q)]^2}{q^2 t \max_\tau [N - S_\tau(2q)]}$$
$$T\Delta S_{\text{tot}} \ge \frac{N [R_g^2(t) - R_g^2(0)]^2}{4 \int_0^t d\tau R_g^2(\tau)/\gamma} \ge \frac{N [R_g^2(t) - R_g^2(0)]^2}{4t \max_\tau [R_g^2(\tau)]/\gamma}.$$
 (6)

The second bound in each line is a simplification when the maximum over time is known, and it is not necessary to measure for all τ . The quality of these bounds for entropy production of internal degrees of freedom during relaxation is shown in Fig. 2 for a solution of (b,c) Rouse polymers upon a temperature quench (see Fig. 2b,c; probing, e.g., the thermal relaxation asymmetry [16, 19, 74]) and confined nano-crystals upon a structural transformation (see Fig. 2d,e), respectively. Dissipative processes occur on distinct length scales, leading to changes of $S_t(q)$ at distinct q (Fig. 2b,d). In both cases, the sharpness of the bounds (6) depends on q, giving insight into the participation of distinct modes in dissipation. The q-dependence can in turn be used for optimization of inference. For small q modes we recover the bound in terms of $R_q^2(\tau)$. We recover ~ 30 % of $\Delta S_{\rm tot}$ in the quenched polymer solution and ~ 15 % for the transforming nano-crystal. This is in fact a lot considering that we only measure a 1d projection of 3(N-1)internal degrees of freedom. Note that here the large contribution from low q-modes is intuitive, since the temperature change mostly affects the overall statistical size of the polymer.




FIG. 2. (a) Sketch of a scattering setup with (b,c) Rouse polymers with N = 10 beads subject to a temperature quench from $T_0 = 2T$ to T at t = 0, and (d,e) harmonically confined "nano-crystal" with N = 16 with Hookean neighborinteractions subject to a quench in rest positions (see [62] for model details and parameters). (b,d) Structure factors and (c,e) corresponding quality factors [see Eq. (6); "simpler" bound contains max_{τ} instead of integral].




FIG. 3. Schematic of (a) a Stirling heat engine as realized in Ref. [75] and (b) a Carnot heat engine with adiabatic heating and cooling [78] as realized in Ref. [76]. Note that the stiffness is constant ("isochoric") during heating and cooling in (a) but *not* in (b). Red denotes hot and blue cold temperatures $T(\tau)$ of the environment (not to be confused with $T_{\rm p}$).

The bound (1) will be useful for many bulk experiments beyond scattering. Consider, for example, a bulk measurement of the mean FRET efficiency for a pair of donor and acceptor chromophores with Förster distance R_0 attached to some macromolecule, $E_t = \langle (1 + [(\mathbf{x}_t^{\mathrm{don}} - \mathbf{x}_t^{\mathrm{acc}})/R_0]^6)^{-1} \rangle$, where the simplest bound reads $T\Delta S_{\mathrm{tot}} \geq R_0^2 \gamma (E_t - E_0)^2/8t$ (for details see [62]).

Example 3: Higher-order transport. — Consider now a centered transient process x_{τ} , i.e., with constant mean $\langle x_t - x_0 \rangle = 0$. There is no mean transport. However, there is higher order transport, in the simplest case $\langle x_t^2 - x_0^2 \rangle \neq 0$. Setting $z_{\tau} = x_{\tau}^2$ we then have $T\Delta S_{\text{tot}} \geq \gamma \langle x_t^2 - x_0^2 \rangle^2 / 4 \int_0^t d\tau \langle x_{\tau}^2 \rangle$. A concrete example are Brownian heat engines [75–77] in Fig. 3, where an "effective temperature" in a parabolic trap with stiffness $\kappa(\tau)$ is defined as $T_{\rm p}(\tau) \equiv \kappa(\tau) \langle x_{\tau}^2 \rangle / k_{\rm B}$ [77]. In this scenario, where the medium temperature varies in time, the left-hand side of the transport bound (1) becomes

$$T\Delta S_{\text{tot}} \to \Delta(TS_{\text{tot}}) \equiv \int_0^{\tau} d\tau T(\tau) S_{\text{tot}}(\tau), \text{ yielding [79]}$$
$$\Delta(TS_{\text{tot}}) \ge \frac{\left[T_{\text{p}}(t) - T_{\text{p}}(0) - \int_0^t T_{\text{p}}(\tau) \frac{\partial_{\tau}\kappa(\tau)}{\kappa(\tau)} d\tau\right]^2}{4\int_0^t T_{\text{p}}(\tau)\kappa(\tau) d\tau/k_{\text{B}}\gamma}.$$
 (7)

During an isochoric heating step ($\kappa = \text{constant}$ such that $\partial_{\tau}\kappa(\tau) = 0$) highlighted in Fig. 3a, no work is performed [75], and the dissipation $\Delta(TS_{\text{tot}})$ of an efficient engine should be minimal. Thus, to achieve a given $T_{\text{p}}(t) - T_{\text{p}}(0)$ [80] for minimal $\Delta(TS_{\text{tot}})$, assuming that we can saturate the bound (as we can in the overdamped case), we either need long t or must maximize $\int_0^t d\tau T_{\text{p}}(\tau)$, implying substantial heating at the beginning of the time interval [0, t]. For time-dependent $\kappa(\tau)$ (see Fig. 3b or the Carnot engine [76]), the bound in Eq. (7) is more complicated since all terms contribute when $\partial_{\tau}\kappa(\tau) \neq 0$. However, the bound still serves as a fundamental limit that can be evaluated for any given protocol. Note that the results equally apply to underdamped heat engines (as theoretically considered in, e.g., [81]).

Of mathematical interest for higher-order transport is the moment-generating function $\phi_{\tau}(\mathbf{q}) \equiv \langle e^{-\mathbf{q}\cdot\mathbf{x}_{\tau}} \rangle$, giving

$$T\Delta S_{\text{tot}} \ge \frac{[\phi_t(\mathbf{q}) - \phi_0(\mathbf{q})]^2}{\int_0^t d\tau \phi_\tau(2\mathbf{q}) \mathbf{q}^T \boldsymbol{\gamma}^{-1}(\tau) \mathbf{q}},$$
(8)

which quantifies how changes of the q-th mode of the probability density contribute to ΔS_{tot} and will be useful for proving future bounds.

Interpretation and handling of $\mathcal{D}^{z}(t)$.—The interpretation of $\mathcal{D}^{z}(t)$ is intuitive: for a process \mathbf{x}_{τ} with given cost ΔS_{tot} , the transport $\langle z_{t} - z_{0} \rangle$ will be larger for a function z that stretches \mathbf{x} more. This stretching is rescaled by $\nabla_{\mathbf{x}} z$ in the quadratic form $\mathcal{D}^{z}(t)$ since ΔS_{tot} does not depend on z.

For simple marginal observations, $z(\mathbf{x}) = x_i$, we have that $\mathcal{D}^z = \gamma^{-1}{}_{ii}$. Moreover, if we only observe z_τ but not \mathbf{x}_{τ} , we can often bound $\mathcal{D}^{z}(t)$ in terms of $\langle z_{\tau} \rangle$ or in terms of constants, such as in the case of R_q^2 and $S_t(q)$ before (e.g., it suffices to know that z_{τ} has bounded derivatives). In the challenging case where we only observe z_{τ} but do not know the function $z(\mathbf{x})$ (i.e., it is some unknown projection or reaction coordinate), we can, given sufficient time resolution, determine or estimate \mathcal{D}^z as follows: For overdamped dynamics we have $t\mathcal{D}^z(t) = \int_0^t d\tau \frac{\operatorname{var}[dz_\tau]}{2k_{\rm B}Td\tau}$ (for steady-state dynamics see [82], where \mathcal{D}^z recently appeared in a correlation inequality). For underdamped dynamics, scalar m, γ , and $z_{\tau} = z(\mathbf{x}_{\tau})$ (no explicit time dependence) we can obtain $\mathcal{D}^{z}(t)$ if we know the momentum relaxation time m/γ via $\frac{\operatorname{var}\left(d\left[\frac{d}{d\tau}z(\mathbf{x}_{\tau})\right]\right)}{2k_{\mathrm{B}}Tdt}$ = $\frac{\gamma^2}{m^2} \langle \gamma^{-1} [\nabla_{\mathbf{x}} z(\mathbf{x}_{\tau})]^2 \rangle$. If the system relaxes to equilibrium we can obtain γ/m from observations of z_{τ} via equilibrium measurements of $\operatorname{var}(d[\frac{d}{d\tau} z(\mathbf{x}_{\tau})])$ and $\operatorname{var}(dz(\mathbf{x}_{\tau}))$ (see [62]). If the system relaxes to a NESS, we can upper bound \mathcal{D}^z by using that $m/\gamma \leq t_{\rm rel}$ with the relaxation time of the system determined, e.g. from correlation func-tions $t_{\rm rel}^{-1} = -\lim_{t\to\infty} t^{-1} \ln[\langle z_t z_0 \rangle - \langle z_t \rangle \langle z_0 \rangle]$ (see [62]).

Conclusion.—We proved an inequality upperbounding transport of any differentiable scalar observable in a general *d*-dimensional non-equilibrium system in terms of the total entropy production and fluctuation-scale function that "corrects" for the amount the observation stretches microscopic coordinates. We explained how to saturate the bound. The result. classifiable as a time-integrated generalized speed limit, may be understood as a thermodynamic cost of transport of observables and allows for inferring a lower bound on dissipation, thus complementing the TUR and existing speed limits. The bound places underdamped and overdamped stochastic as well as deterministic systems on equal footing. This fills an important gap, because microscopic dynamics is—at least on short time scales—underdamped, and the TUR does not hold for underdamped dynamics. In particular short-time TURs for overdamped dynamics [29, 30, 83] are expected to fail.

- U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, Rep. Prog. Phys. 75, 126001 (2012).
- [2] U. Seifert, Entropy production along a stochastic trajectory and an integral fluctuation theorem, Phys. Rev. Lett. 95, 040602 (2005).
- [3] U. Seifert, Stochastic thermodynamics: From principles to the cost of precision, Physica (Amsterdam) 504A, 176 (2018).
- [4] M. P. Leighton and D. A. Sivak, Dynamic and thermodynamic bounds for collective motor-driven transport, Phys. Rev. Lett. 129, 118102 (2022).
- [5] M. P. Leighton and D. A. Sivak, Inferring subsystem efficiencies in bipartite molecular machines, Phys. Rev. Lett. 130, 178401 (2023).
- [6] I. Di Terlizzi, M. Gironella, D. Herraez-Aguilar, T. Betz, F. Monroy, M. Baiesi, and F. Ritort, Variance sum rule for entropy production, Science 383, 971 (2024).
- [7] D. Sato, T. Hikima, and M. Ikeguchi, Time-resolved small-angle X-ray scattering of protein cage assembly, in *Protein Cages* (Springer US, 2023) p. 211.
- [8] M. Roessle, E. Manakova, I. Lauer, T. Nawroth, J. Holzinger, T. Narayanan, S. Bernstorff, H. Amenitsch, and H. Heumann, Time-resolved small angle scattering: kinetic and structural data from proteins in solution, J. Appl. Cryst. 33, 548 (2000).
- [9] S. Akiyama, S. Takahashi, T. Kimura, K. Ishimori, I. Morishima, Y. Nishikawa, and T. Fujisawa, Conformational landscape of cytochrome c folding studied by microsecond-resolved small-angle x-ray scattering, Proc. Natl. Acad. Sci. U.S.A. 99, 1329 (2002).
- [10] J. Stamatoff, An approach for time-resolved x-ray scattering, Biophys. J. 26, 325 (1979).
- [11] Y. Yamada, T. Matsuo, H. Iwamoto, and N. Yagi, A compact intermediate state of calmodulin in the process of target binding, Biochem. 51, 3963 (2012).
- [12] H. S. Cho, F. Schotte, V. Stadnytskyi, and P. Anfinrud, Time-resolved X-ray scattering studies of proteins, Curr. Opin. Struct. Biol. **70**, 99 (2021).

By only requiring averages, the transport bound is statistically less demanding and applicable to both, singlemolecule as well as bulk experiments. This is attractive in the context of time-resolved X-ray scattering on biomolecules, as it will allow thermodynamic inference from bulk observations of controlled transients [68–71]. This may facilitate thermodynamic inference on molecular machines without an obvious directionality such molecular chaperones [60], which remains challenging even with most advanced single-molecule techniques [61]. The bound allows for versatile applications and generalizations to vectorial observables \mathbf{z} and adaptations for Markov-jump dynamics.

Acknowledgments.—Financial support from the European Research Council (ERC) under the European Union's Horizon Europe research and innovation program (grant agreement No 101086182 to AG) is gratefully acknowledged.

- [13] D. Arnlund, L. C. Johansson, C. Wickstrand, A. Barty, G. J. Williams, E. Malmerberg, J. Davidsson, D. Milathianaki, D. P. DePonte, R. L. Shoeman, D. Wang, D. James, G. Katona, S. Westenhoff, T. A. White, A. Aquila, S. Bari, P. Berntsen, M. Bogan, T. B. van Driel, R. B. Doak, K. S. Kjær, M. Frank, R. Fromme, I. Grotjohann, R. Henning, M. S. Hunter, R. A. Kirian, I. Kosheleva, C. Kupitz, M. Liang, A. V. Martin, M. M. Nielsen, M. Messerschmidt, M. M. Seibert, J. Sjöhamn, F. Stellato, U. Weierstall, N. A. Zatsepin, J. C. H. Spence, P. Fromme, I. Schlichting, S. Boutet, G. Groenhof, H. N. Chapman, and R. Neutze, Visualizing a protein quake with time-resolved X-ray scattering at a freeelectron laser, Nat. Methods 11, 923 (2014).
- [14] H. S. Cho, N. Dashdorj, F. Schotte, T. Graber, R. Henning, and P. Anfinrud, Protein structural dynamics in solution unveiled via 100-ps time-resolved x-ray scattering, Proc. Natl. Acad. Sci. U.S.A. 107, 7281 (2010).
- [15] M. Cammarata, M. Levantino, F. Schotte, P. A. Anfinrud, F. Ewald, J. Choi, A. Cupane, M. Wulff, and H. Ihee, Tracking the structural dynamics of proteins in solution using time-resolved wide-angle X-ray scattering, Nat. Methods 5, 881 (2008).
- [16] M. Ibáñez, C. Dieball, A. Lasanta, A. Godec, and R. A. Rica, Heating and cooling are fundamentally asymmetric and evolve along distinct pathways, Nat. Phys. 20, 135 (2024).
- [17] J. L. Lebowitz and P. G. Bergmann, Irreversible Gibbsian ensembles, Ann. Phys. 1, 1 (1957).
- [18] S. Vaikuntanathan and C. Jarzynski, Dissipation and lag in irreversible processes, Europhys. Lett. 87, 60005 (2009).
- [19] A. Lapolla and A. Godec, Faster uphill relaxation in thermodynamically equidistant temperature quenches, Phys. Rev. Lett. **125**, 110602 (2020).
- [20] Z. T. Berndsen, N. Keller, S. Grimes, P. J. Jardine, and D. E. Smith, Nonequilibrium dynamics and ultraslow relaxation of confined DNA during viral packaging, Proc. Natl. Acad. Sci. U.S.A. 111, 8345 (2014).

- [21] H. Turlier, D. A. Fedosov, B. Audoly, T. Auth, N. S. Gov, C. Sykes, J.-F. Joanny, G. Gompper, and T. Betz, Equilibrium physics breakdown reveals the active nature of red blood cell flickering, Nat. Phys. 12, 513 (2016).
- [22] T. Stuchinskaya, L. A. Mitchenall, A. J. Schoeffler, K. D. Corbett, J. M. Berger, A. D. Bates, and A. Maxwell, How do type II topoisomerases use ATP hydrolysis to simplify DNA topology beyond equilibrium? Investigating the relaxation reaction of nonsupercoiling type II topoisomerases, J. Mol. Biol. **385**, 1397 (2009).
- [23] J. Gieseler and J. Millen, Levitated nanoparticles for microscopic thermodynamics-a review, Entropy 20, 326 (2018).
- [24] A. C. Barato and U. Seifert, Thermodynamic uncertainty relation for biomolecular processes, Phys. Rev. Lett. 114, 158101 (2015).
- [25] T. R. Gingrich, J. M. Horowitz, N. Perunov, and J. L. England, Dissipation bounds all steady-state current fluctuations, Phys. Rev. Lett. 116, 120601 (2016).
- [26] T. R. Gingrich, G. M. Rotskoff, and J. M. Horowitz, Inferring dissipation from current fluctuations, J. Phys. A: Math. Theor. 50, 184004 (2017).
- [27] T. Van Vu and Y. Hasegawa, Uncertainty relations for underdamped Langevin dynamics, Phys. Rev. E 100, 032130 (2019).
- [28] J. M. Horowitz and T. R. Gingrich, Thermodynamic uncertainty relations constrain non-equilibrium fluctuations, Nat. Phys. 16, 15 (2019).
- [29] S. K. Manikandan, D. Gupta, and S. Krishnamurthy, Inferring entropy production from short experiments, Phys. Rev. Lett. **124**, 120603 (2020).
- [30] S. Otsubo, S. Ito, A. Dechant, and T. Sagawa, Estimating entropy production by machine learning of short-time fluctuating currents, Phys. Rev. E 101, 062106 (2020).
- [31] K. Liu, Z. Gong, and M. Ueda, Thermodynamic uncertainty relation for arbitrary initial states, Phys. Rev. Lett. **125**, 140602 (2020).
- [32] T. Koyuk and U. Seifert, Thermodynamic uncertainty relation for time-dependent driving, Phys. Rev. Lett. 125, 260604 (2020).
- [33] C. Dieball and A. Godec, Mathematical, thermodynamical, and experimental necessity for coarse graining empirical densities and currents in continuous space, Phys. Rev. Lett. **129**, 140601 (2022).
- [34] C. Dieball and A. Godec, Coarse graining empirical densities and currents in continuous-space steady states, Phys. Rev. Res. 4, 033243 (2022).
- [35] A. Dechant and S.-i. Sasa, Continuous time reversal and equality in the thermodynamic uncertainty relation, Phys. Rev. Res. 3, L042012 (2021).
- [36] A. Dechant and S.-i. Sasa, Improving thermodynamic bounds using correlations, Phys. Rev. X 11, 041061 (2021).
- [37] C. Dieball and A. Godec, Direct route to thermodynamic uncertainty relations and their saturation, Phys. Rev. Lett. 130, 087101 (2023).
- [38] K. Brandner, K. Saito, and U. Seifert, Strong bounds on onsager coefficients and efficiency for three-terminal thermoelectric transport in a magnetic field, Phys. Rev. Lett. **110**, 070603 (2013).
- [39] N. Shiraishi, K. Funo, and K. Saito, Speed limit for classical stochastic processes, Phys. Rev. Lett. **121**, 070601 (2018).
- [40] N. Shiraishi and K. Saito, Information-theoretical bound

of the irreversibility in thermal relaxation processes, Phys. Rev. Lett. **123**, 110603 (2019).

- [41] G. Falasco and M. Esposito, Dissipation-time uncertainty relation, Phys. Rev. Lett. 125, 120604 (2020).
- [42] K. Yoshimura and S. Ito, Thermodynamic uncertainty relation and thermodynamic speed limit in deterministic chemical reaction networks, Phys. Rev. Lett. **127**, 160601 (2021).
- [43] E. Aurell, C. Mejía-Monasterio, and P. Muratore-Ginanneschi, Optimal protocols and optimal transport in stochastic thermodynamics, Phys. Rev. Lett. 106, 250601 (2011).
- [44] E. Aurell, K. Gawedzki, C. Mejía-Monasterio, R. Mohayaee, and P. Muratore-Ginanneschi, Refined second law of thermodynamics for fast random processes, J. Stat. Phys. 147, 487–505 (2012).
- [45] E. Potanina, C. Flindt, M. Moskalets, and K. Brandner, Thermodynamic bounds on coherent transport in periodically driven conductors, Phys. Rev. X 11, 021013 (2021).
- [46] A. Dechant and S.-i. Sasa, Entropic bounds on currents in Langevin systems, Phys. Rev. E 97, 062101 (2018).
- [47] S. Ito and A. Dechant, Stochastic time evolution, information geometry, and the Cramér-Rao bound, Phys. Rev. X 10, 021056 (2020).
- [48] T. Van Vu and K. Saito, Thermodynamic unification of optimal transport: Thermodynamic uncertainty relation, minimum dissipation, and thermodynamic speed limits, Phys. Rev. X 13, 011013 (2023).
- [49] P. Pietzonka, Classical pendulum clocks break the thermodynamic uncertainty relation, Phys. Rev. Lett. 128, 130606 (2022).
- [50] L. P. Fischer, H.-M. Chun, and U. Seifert, Free diffusion bounds the precision of currents in underdamped dynamics, Phys. Rev. E **102**, 012120 (2020).
- [51] C. Kwon and H. K. Lee, Thermodynamic uncertainty relation for underdamped dynamics driven by timedependent protocols, New J. Phys. 24, 013029 (2022).
- [52] R.-S. Fu and T. R. Gingrich, Thermodynamic uncertainty relation for Langevin dynamics by scaling time, Phys. Rev. E 106, 024128 (2022).
- [53] S. Lee, D.-K. Kim, J.-M. Park, W. K. Kim, H. Park, and J. S. Lee, Multidimensional entropic bound: Estimator of entropy production for Langevin dynamics with an arbitrary time-dependent protocol, Phys. Rev. Res. 5, 013194 (2023).
- [54] J. Lyu, K. J. Ray, and J. P. Crutchfield, Entropy production by underdamped Langevin dynamics (2024).
- [55] J. K. G. Dhont, An introduction to dynamics of colloids, Studies in Interface Science (Elsevier, 1996).
- [56] J. A. McCammon, Protein dynamics, Rep. Prog. Phys. 47, 1 (1984).
- [57] J. C. Smith, Protein dynamics: comparison of simulations with inelastic neutron scattering experiments, Q. Rev. Biophys. 24, 227 (1991).
- [58] G. R. Kneller, Inelastic neutron scattering from damped collective vibrations of macromolecules, Chem. Phys. 261, 1 (2000).
- [59] F. N. Brünig, J. O. Daldrop, and R. R. Netz, Pairreaction dynamics in water: Competition of memory, potential shape, and inertial effects, J. Phys. Chem. B 126, 10295–10304 (2022).
- [60] A. Mashaghi, G. Kramer, D. C. Lamb, M. P. Mayer, and S. J. Tans, Chaperone action at the single-molecule level, Chem. Rev. 114, 660–676 (2013).

- [61] C. Ratzke, B. Hellenkamp, and T. Hugel, Four-colour fret reveals directionality in the hsp90 multicomponent machinery, Nat. Commun. 5, 4192 (2014).
- [62] See Supplemental Material at [...] including references [????].
- [63] J.-D. Benamou and Y. Brenier, A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem, Numer. Math. 84, 375 (2000).
- [64] I. Josts, Y. Gao, D. C. Monteiro, S. Niebling, J. Nitsche, K. Veith, T. W. Gräwert, C. E. Blanchet, M. A. Schroer, N. Huse, A. R. Pearson, D. I. Svergun, and H. Tidow, Structural kinetics of msba investigated by stopped-flow time-resolved small-angle x-ray scattering, Structure 28, 348 (2020).
- [65] H. Risken, *The Fokker-Planck Equation* (Springer Berlin Heidelberg, 1989).
- [66] G. Wilemski, On the derivation of Smoluchowski equations with corrections in the classical theory of Brownian motion, J. Stat. Phys. 14, 153 (1976).
- [67] A. C. Barato and U. Seifert, Cost and precision of Brownian clocks, Phys. Rev. X 6, 041053 (2016).
- [68] K. Dave and M. Gruebele, Fast-folding proteins under stress, Cell. Mol. Life. Sci. 72, 4273 (2015).
- [69] J. Woenckhaus, R. Köhling, P. Thiyagarajan, K. C. Littrell, S. Seifert, C. A. Royer, and R. Winter, Pressurejump small-angle x-ray scattering detected kinetics of staphylococcal nuclease folding, Biophysical Journal 80, 1518 (2001).
- [70] J. Kubelka, Time-resolved methods in biophysics. 9. laser temperature-jump methods for investigating biomolecular dynamics, Photochem. Photobiol. Sci. 8, 499 (2009).
- [71] B. Vestergaard, M. Groenning, M. Roessle, J. S. Kastrup, M. v. de Weert, J. M. Flink, S. Frokjaer, M. Gajhede, and D. I. Svergun, A helical structural nucleus is the primary elongating unit of insulin amyloid fibrils, PLoS Biol. 5, e134 (2007).
- [72] C. Dieball, D. Krapf, M. Weiss, and A. Godec, Scattering fingerprints of two-state dynamics, New J. Phys. 24, 023004 (2022).
- [73] D. I. Svergun and M. H. J. Koch, Small-angle scattering studies of biological macromolecules in solution, Rep. Prog. Phys. 66, 1735 (2003).
- [74] C. Dieball, G. Wellecke, and A. Godec, Asymmetric thermal relaxation in driven systems: Rotations go opposite ways, Phys. Rev. Res. 5, L042030 (2023).
- [75] V. Blickle and C. Bechinger, Realization of a micrometresized stochastic heat engine, Nat. Phys. 8, 143 (2011).
- [76] I. A. Martínez, É. Roldán, L. Dinis, D. Petrov, J. M. R. Parrondo, and R. A. Rica, Brownian Carnot engine, Nat. Phys. **12**, 67 (2015).
- [77] I. A. Martínez, E. Roldán, L. Dinis, and R. A. Rica, Colloidal heat engines: a review, Soft Matter 13, 22 (2017).
- [78] I. A. Martínez, E. Roldán, L. Dinis, D. Petrov, and R. A. Rica, Adiabatic processes realized with a trapped brownian particle, Phys. Rev. Lett. **114**, 120601 (2015).
- [79] Alternatively we can also derive a bound for ΔS_{tot} but we opt for $\Delta(TS_{\text{tot}})$ for a "direct" energetic interpretation.
- [80] More precisely, a given $\gamma [T_{\rm p}(t) T_{\rm p}(0)]^2 / \kappa$.
- [81] A. Dechant, N. Kiesel, and E. Lutz, Underdamped stochastic heat engine at maximum efficiency, EPL (Europhysics Letters) 119, 50003 (2017).
- [82] A. Dechant, J. Garnier-Brun, and S.-i. Sasa, Thermodynamic bounds on correlation times, Phys. Rev. Lett. 131,

[83] S. Otsubo, S. K. Manikandan, T. Sagawa, and S. Krishnamurthy, Estimating time-dependent entropy production from non-equilibrium trajectories, Commun. Phys. 5, 11 (2022).

Supplemental Material for: Thermodynamic bounds on generalized transport: From single-molecule to bulk observables

Cai Dieball and Aljaž Godec

Mathematical bioPhysics Group, Max Planck Institute for Multidisciplinary Sciences, 37077 Göttingen, Germany

In this Supplemental Material we provide derivations and additional details for results and examples shown in the Letter. We provide a detailed account and comparison to existing results, the detailed derivation of the main result, and details and parameters complementing the examples in Figs. 1 and 2. Moreover, we explain in detail how to obtain \mathcal{D}^z from short-time fluctuations.

CONTENTS

I.	. Comparison to existing literature	2
	A. General discussion	$2 \\ 2$
	B. Spatially periodic systems	Z
II.	. Proof of the main result	4
	A. Sketch of proof	4
	B. Full proof	4
	C. Overdamped limit	5
	D. Generalization to velocity dependent variables	6
III.	. Details on the example in Fig. 1 in the Letter	6
	A. Saturation	7
	B. Short-time limit	8
	C. TUR applied to underdamped dynamics	8
IV.	. Derivation of Eq. (6) in the Letter	8
	A. Radius of gyration	9
	B. Structure factor	10
V.	. Details on the example in Fig. 2 in the Letter	11
	A. Computing the structure factor	11
	B. Computation of entropy production for the temperature quench	12
	C. Computation of entropy production for the structure quench	13
VI.	. Details on the handling of D^z	13
	A. Overdamped dynamics	13
	B. Underdamped dynamics	14
VII.	FRET bound	14
	References	14

I. COMPARISON TO EXISTING LITERATURE

A. General discussion

The transport bound presented in Eq. (1) in the Letter reads

$$T\Delta S_{\text{tot}} \geq \frac{\left\langle z_t - z_0 - \int_0^t \partial_\tau z_\tau d\tau \right\rangle^2}{t\mathcal{D}^z(t)}$$
$$\mathcal{D}^z(t) \equiv \frac{1}{t} \int_0^t \langle [\nabla_{\mathbf{x}} z_\tau]^T \boldsymbol{\gamma}^{-1}(\tau) \nabla_{\mathbf{x}} z_\tau \rangle d\tau, \tag{S1}$$

where the observable $z_{\tau} \equiv z(\mathbf{x}_{\tau}, \tau)$ is a scalar function of \mathbf{x}_{τ} and τ . This equation is proven below for underdamped stochastic dynamics, which includes overdamped dynamics as a special case.

Taking the limit $t \to 0$ without explicit time-dependence, i.e., $z_{\tau} \equiv z(\mathbf{x}_{\tau})$, Eq. (S1) simplified to the speed limit

$$\left(\frac{\mathrm{d}}{\mathrm{d}\tau}\langle z_{\tau}\rangle\right)^{2} \leq T\dot{S}_{\mathrm{tot}}(\tau)\langle [\nabla_{\mathbf{x}}z_{\tau}]^{T}\boldsymbol{\gamma}^{-1}(\tau)\nabla_{\mathbf{x}}z_{\tau}\rangle, \qquad (S2)$$

where $\dot{S}_{tot}(\tau) \equiv \frac{d}{d\tau} \Delta S_{tot}[0,\tau]$ is the entropy production rate. This result was derived for overdamped dynamics with diffusion matrix $\mathbf{D} = k_{\rm B}T\gamma^{-1}$ in Ref. [1]. However, it was neither extended to underdamped dynamics, nor identified to apply to bulk experiments. The importance of this extension to underdamped dynamics becomes particularly clear in this form, since speed limits inherently address the short-time behavior $\frac{d}{d\tau}\langle z_{\tau}\rangle$, which (even despite the averaging) differs substantially between for overdamped and underdamped dynamics (see $t \to 0$ in Fig. 1b in the Letter and divergence in Fig. S2). Also note that in the underdamped setting (colored lines in Fig. 1b in the Letter), the maximal quality of the bound is not at $t \to 0$, which (besides being more directly applicable) is a strong argument for preferring the time-integrated version over the speed limit (the latter corresponds to $t \to 0$).

Further restricting to one-dimensional space and $z_{\tau} = x_{\tau}$, i.e., to the bound

$$\left(\frac{\mathrm{d}}{\mathrm{d}\tau}\langle x_{\tau}\rangle\right)^{2} \leq D\dot{S}_{\mathrm{tot}}(\tau)/k_{\mathrm{B}}.$$
(S3)

This transport bound corresponds in periodic systems (see also following section) to speed limits from Refs. [2, 3] which were applied to gain important theoretical insights into the efficiency and energetics of molecular motors.

The bound (S1) complements the Benamou-Brenier formula [4] that applies to overdamped Markov observables [5, 6] by allowing for the more realistic underdamped dynamics, general projected (non-Markovian) observables z_{τ} , and time-dependent diffusivity. While the Benamou-Brenier formula gives a lower bound on the entropy production from the difference in initial and final probability densities, we directly target more easily measurable observables.

B. Spatially periodic systems

Systems that are periodic in space (such as, e.g., Brownian motion on a ring or an interval with periodic boundary conditions) are paradigmatic examples for non-equilibrium steady states [7], in particular to model molecular motors [2, 8]. Such systems can either be viewed as transient (e.g., with angle on the ring in $(-\infty, \infty)$]) or as steady states (e.g., angle in $[0, 2\pi)$). This complementary picture allows to apply both, the transport bound and the thermodynamic uncertainty relation (TUR) [9], to such systems.

In Fig. S1 we evaluate for simulated overdamped Langevin motion

$$dx_{\tau} = [D\partial_x \ln p_s(x_{\tau}) + \nu_s(x_{\tau})]d\tau + \sqrt{2D}dW_{\tau}, \qquad (S4)$$

with $D = k_{\rm B}T/\gamma$ and periodic functions $p_{\rm s}$ and $\nu_{\rm s}$ different versions of the TUR and transport bound, given by the equations (see Refs. [10, 11] for correlation-TUR (CTUR))

$$J_t^x \equiv \int_{\tau=0}^{\tau=t} 1 \circ \mathrm{d}x_\tau = x_t - x_0$$
$$J_t^\nu \equiv \int_{\tau=0}^{\tau=t} \nu(x_\tau) \circ \mathrm{d}x_\tau$$




FIG. S1. Evaluation of the bounds in Eq. (S5) for 800 trajectories (big black crosses) for the Langevin motion (S4) with $k_{\rm B}T = 1, t = 1800, D = 0.01, p_{\rm s}(x) = f(x) / \int_0^L f(x') dx'$ with $f(x) = 1.7 + \cos(2\pi x/L) + 0.3\sin(4\pi x/L) + 0.2\cos(8\pi x/L)$ and periodicity $L = 5, \nu_{\rm s}(x) = 0.01/p_{\rm s}(x)$ simulated with Euler time step dt = 0.04. In practical applications, most parameters are not known. Therefore, for the evaluation of Eq. (S5), $\nu_s(x)$ is inferred from the trajectories by inferring p_s from a histogram projected on [0, L] and subsequently fitting the product $\nu_{s}(x)p_{s}(x)$ which must be constant in one-dimensional steady-state dynamics [12]. D is obtained from the (overdamped) short time fluctuations $D = \operatorname{var}(dx_{\tau})/2d\tau$ averaged over all times $\tau \in [0, t]$. To address the fluctuations of the result, a bootstrapping procedure has been employed where 25 times the quality factor is derived from a random subset of 25 % of the trajectories (colored crosses).

$$\begin{aligned}
\rho_t^{\nu} &\equiv \int_0^t \nu^2(x_{\tau}) \mathrm{d}\tau \\
z_{\tau}^{\nu} &\equiv \int_0^{x_{\tau}} \nu(x) \mathrm{d}x \\
Q_{\mathrm{TUR}\,x/\nu} &\equiv \frac{2\langle J_t^{x/\nu} \rangle^2}{\mathrm{var}(J_t^{x/\nu})} \frac{k_{\mathrm{B}}}{\Delta S_{\mathrm{tot}}} \\
Q_{\mathrm{CTUR}\,\nu} &\equiv \frac{2\langle J_t^{\nu} \rangle^2}{\mathrm{var}(J_t^{\nu}) \left[1 - \frac{\mathrm{cov}(J_t^{\nu}, \rho_t^{\nu})^2}{\mathrm{var}(J_t^{\nu}) \mathrm{var}(\rho_t^{\nu})}\right]} \frac{k_{\mathrm{B}}}{\Delta S_{\mathrm{tot}}} \\
Q_{\mathrm{transport}\,x} &\equiv \frac{\langle x_t - x_0 \rangle^2}{Dt} \frac{k_B}{\Delta S_{\mathrm{tot}}} \\
Q_{\mathrm{transport}\,\nu} &\equiv \frac{\langle z_t^{\nu} - z_0^{\nu} \rangle^2}{D\langle (\partial_x z_\tau^{\nu})^2 \rangle} \frac{k_B}{\Delta S_{\mathrm{tot}}}.
\end{aligned}$$
(S5)

For this example, the transport bound $Q_{\text{transport }x}$ performs better than the TUR. However, there are also overdamped periodic examples where the TUR performs performs better [2]. As expected, the absence of variances in the transport bound leads to a statistically more robust inference as can be seen from the spread of the colored crosses in Fig. S1 (i.e. the statistical noise due to undersampling is substantially smaller when using the transport bound). $Q_{\text{TUR}\nu}$ was in included in the comparison since it is known to approach 1 close to equilibrium [13] and for short times [14]. However, it does not exceed $Q_{\text{TUR }x}$ in this example.

l

The analytical value for the CTUR with the optimal choice of J_t^{ν} , ρ_t^{ν} is $Q_{\text{CTUR}\nu} = 1$ [10, 11]. This is roughly achieved for 800 long trajectories. However, with "only" 200 trajectories the result still fluctuates a lot. The saturated transport bound with the choice of z_{τ}^{ν} is also very close to the analytical value of 1 and fluctuates much less, i.e., this bound is accessible from less trajectories as compared to the TUR.

Compared to the TUR, the transport bound has the additional advantage of being valid for transients and underdamped motion, i.e., the inference will not fail in case these underlying assumptions are not justified (one only has to be careful what assumptions one wants to employ for evaluating \mathcal{D}^z). For an illustration of the failure of the overdamped TUR see Fig. S2. in Sec. III C (transient example; see Ref. [15] for a steady-state example; see Supplemental Material of Ref. [11] for an example where the overdamped steady-state TUR fails for overdamped transients).

II. PROOF OF THE MAIN RESULT

Here, we prove our main result, i.e., the transport bound in Eq. (1) in the Letter. We first give a short sketch (which only serves to give an optional overview and may be skipped) and afterwards write the full proof. Finally, we will also show how the proof dramatically simplifies in the special case of overdamped dynamics.

A. Sketch of proof

To prove Eq. (1) in the Letter for an observable z_{τ} , we use the Cauchy-Schwarz inequality for stochastic integrals similarly as in [11] but generalized to underdamped dynamics. Recalling γ and $d\mathbf{W}_{\tau}$ and the $\boldsymbol{\nu}$ from Eqs. (2) and (4) in the Letter, respectively, we define the stochastic integrals $A_t \equiv \int_{\tau=0}^{\tau=t} \boldsymbol{\nu}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \cdot \sqrt{\gamma} d\mathbf{W}_{\tau}$ and $B_t \equiv \int_{\tau=0}^{\tau=t} \{\nabla_{\mathbf{x}} z_{\tau}\} \cdot \sqrt{\gamma}^{-1^T} d\mathbf{W}_{\tau}$ for which we can show that $\langle A_t^2 \rangle = T\Delta S_{\text{tot}}$ and $\langle B_t^2 \rangle = t\mathcal{D}^z(t)$, where ΔS_{tot} and $D^z(t)$ are defined in Eqs. (5) and (1) in the Letter, respectively. The Cauchy-Schwarz inequality thus implies $T\Delta S_{\text{tot}} \geq \langle A_t B_t \rangle^2 / t\mathcal{D}^z(t)$. To complete the proof, we compute $\langle A_t B_t \rangle = \langle \int_0^t \boldsymbol{\nu}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \cdot \{\nabla_{\mathbf{x}} z_{\tau}\} d\tau \rangle$ via rewriting $\{\nabla_{\mathbf{x}} z_{\tau}\} = \gamma \mathbf{m}^{-1} \nabla_{\mathbf{v}} \{\nabla_{\mathbf{v}}^T \mathbf{m} \gamma^{-1} \nabla_{\mathbf{x}} z_{\tau}\}$ [with \mathbf{m} from Eq. (2) in the Letter], integrating by parts in $\nabla_{\mathbf{v}}$, and substituting the Klein-Kramers equation (see Eq. (S6) below) in the form $\nabla_{\mathbf{v}} \cdot \mathbf{m}^{-1} \gamma \boldsymbol{\nu} \rho = [\nabla_{\mathbf{x}} \cdot \mathbf{v} + \nabla_{\mathbf{v}} \cdot \mathbf{m}^{-1} \mathbf{F}(\mathbf{x}, \tau) + \partial_{\tau}]\rho(\mathbf{x}, \mathbf{v}, \tau)$, which upon further integrations by parts and simplifications yields $\langle A_t B_t \rangle = \int_0^t d\tau \langle \mathbf{v}_{\tau} \cdot \nabla_{\mathbf{x}} z_{\tau} \rangle = \langle \int_0^t d\tau (\frac{d}{d\tau} - \partial_{\tau}) z_{\tau} \rangle = \langle z_t - z_0 - \int_0^t \partial_{\tau} z_{\tau} d\tau \rangle$, completing the proof of Eq. (1) in the Letter.

 $\langle z_t - z_0 - \int_0^t \partial_\tau z_\tau d\tau \rangle$, completing the proof of Eq. (1) in the Letter. From the proof we immediately know how to achieve saturation by saturating the Cauchy-Schwarz inequality, allowing optimal inference. Saturation occurs for $\nabla_{\mathbf{x}} z_\tau = c \gamma \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau)$ for any constant *c*. While one may not always be able to choose *z* this way, one should aim to approach this for saturation. If d = 1 (with natural boundaries) and $\boldsymbol{\nu}$ does not depend on \mathbf{v} , saturation is always feasible.

B. Full proof

Consider underdamped stochastic dynamics as in Eq. (2) in the Letter. Consider positive definite symmetric matrices **m** and $\gamma(\tau)$ (for simplicity denote the latter by γ). Using the "local mean velocity" ν as in Eq. (4) in the Letter we can write the underdamped Klein-Kramers dynamics as [16]

$$\boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \equiv \mathbf{v}_{\tau} + k_{\rm B} T \mathbf{m}^{-1} \frac{\nabla_{\mathbf{v}} \rho(\mathbf{x}, \mathbf{v}, t)}{\rho(\mathbf{x}, \mathbf{v}, t)}$$
$$\partial_{\tau} \rho(\mathbf{x}, \mathbf{v}, t) = \left(-\nabla_{\mathbf{x}} \cdot \mathbf{v} + \mathbf{m}^{-1} \nabla_{\mathbf{v}} \cdot \left[-\mathbf{F}(\mathbf{x}, \tau) + \boldsymbol{\gamma}(\tau) \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \right] \right) \rho(\mathbf{x}, \mathbf{v}, t) \,. \tag{S6}$$

Recalling the setup of the proof as presented above, and using the white noise property/Wiener correlation $\langle dW^i_{\tau} dW^j_{\tau'} \rangle = \delta(\tau - \tau') d\tau d\tau'$ we have

$$A_{t} \equiv \int_{\tau=0}^{\tau=t} \boldsymbol{\nu}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \cdot \sqrt{\gamma} d\mathbf{W}_{\tau}$$

$$B_{t} \equiv \int_{\tau=0}^{\tau=t} \{\nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)\} \cdot \sqrt{\gamma}^{-1^{T}} d\mathbf{W}_{\tau}$$

$$\langle A_{t}^{2} \rangle = \int_{0}^{t} d\tau \langle \boldsymbol{\nu}^{T}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \gamma \boldsymbol{\nu}(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau) \rangle = T \Delta S_{\text{tot}}([0, t])$$

$$\langle B_{t}^{2} \rangle = \int_{0}^{t} d\tau \left\langle \{\nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)\}^{T} \gamma^{-1} \{\nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)\} \right\rangle \equiv t \mathcal{D}^{z}(t) .$$
(S7)

Using the Einstein summation convention, we can check the matrix differential operator identity for a constant (i.e. **x**-independent) invertible $d \times d$ matrix **G**

$$(\mathbf{G}\nabla_{\mathbf{v}})[\mathbf{v}\cdot\mathbf{G}^{-1}\nabla_{\mathbf{x}}z(\mathbf{x},\tau)] = \mathbf{e}_i G_{ij}\partial_{v_j}v_n(\mathbf{G}^{-1})_{nm}\partial_{x_m}z(\mathbf{x},\tau) = \mathbf{e}_i G_{in}(\mathbf{G}^{-1})_{nm}\partial_{x_m}z(\mathbf{x},\tau) = \nabla_{\mathbf{x}}z(\mathbf{x},\tau).$$
(S8)

Using this identity for the choice $\mathbf{G}^{-1} = \mathbf{m}\gamma^{-1}$, integrating by parts, and substituting Eq. (S6) for $\nabla_{\mathbf{v}} \cdot \mathbf{m}^{-1}\gamma\boldsymbol{\nu}$, we obtain

$$\begin{aligned} -\langle A_t B_t \rangle &= -\int_0^t \mathrm{d}\tau \int \mathrm{d}\mathbf{x} \int \mathrm{d}\mathbf{v} \left\{ \mathbf{G} \nabla_{\mathbf{v}} [\mathbf{v} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau)] \right\} \cdot \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \rho(\mathbf{x}, \mathbf{v}, \tau) \\ &= \int_0^t \mathrm{d}(\mathbf{x}, \mathbf{v}, \tau) \{ \mathbf{v} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau) \} \nabla_{\mathbf{v}} \cdot \mathbf{G}^T \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \rho(\mathbf{x}, \mathbf{v}, \tau) \\ &= \int_0^t \mathrm{d}(\mathbf{x}, \mathbf{v}, \tau) \{ \mathbf{v} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau) \} \mathbf{m}^{-1} \nabla_{\mathbf{v}} \cdot \boldsymbol{\gamma} \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \rho(\mathbf{x}, \mathbf{v}, \tau) \\ &= \int_0^t \mathrm{d}\tau \int \mathrm{d}\mathbf{x} \int \mathrm{d}\mathbf{v} \{ \mathbf{v} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau) \} \left[\nabla_{\mathbf{x}} \cdot \mathbf{v} + \mathbf{m}^{-1} \nabla_{\mathbf{v}} \cdot \mathbf{F}(\mathbf{x}, \tau) + \partial_{\tau} \right] \rho(\mathbf{x}, \mathbf{v}, \tau) \\ &= \langle \mathbf{v}_t \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_t, t) - \mathbf{v}_0 \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_0, 0) \rangle \\ &- \mathbf{m} \int_0^t \mathrm{d}\tau \left\langle (\mathbf{v}_\tau \cdot \nabla_{\mathbf{x}}) [\mathbf{v}_\tau \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau)] + \mathbf{m}^{-1} \mathbf{F}(\mathbf{x}, \tau) \rangle \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \tau) + \mathbf{v}_\tau \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} \partial_{\tau} z(\mathbf{x}, \tau) \right\rangle. \end{aligned}$$

Note that

$$\left\langle \mathbf{v}_t \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_t, t) - \mathbf{v}_0 \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_0, 0) \right\rangle = \left\langle \int_0^t \mathrm{d}\tau \frac{\mathrm{d}}{\mathrm{d}\tau} [\mathbf{v}_\tau \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_\tau, \tau)] \right\rangle \,, \tag{S10}$$

and use the equation of motion in Eq. (S6) to see that

$$\left\langle \frac{\mathrm{d}}{\mathrm{d}\tau} [\mathbf{v}_{\tau} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)] \right\rangle = \left\langle \mathbf{m}^{-1} \left(\mathbf{F}(\mathbf{x}_{\tau}, \tau) - \boldsymbol{\gamma} \mathbf{v}_{\tau} \right) \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau) + \mathbf{v}_{\tau} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} [\mathbf{v}_{\tau} \cdot \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau) + \partial_{\tau} z(\mathbf{x}_{\tau}, \tau)] \right\rangle.$$
(S11)

Since ${\bf G}$ is independent of ${\bf x}$ we have

$$(\mathbf{v}_{\tau} \cdot \nabla_{\mathbf{x}})[\mathbf{v}_{\tau} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)] = (\mathbf{v}_{\tau} \cdot \mathbf{G}^{-1} \nabla_{\mathbf{x}})[\mathbf{v}_{\tau} \cdot \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}, \tau)], \qquad (S12)$$

which allows to conclude that

$$-\langle A_t B_t \rangle = -\int_0^t \mathrm{d}\tau \, \langle \mathbf{v}_\tau \cdot \nabla_{\mathbf{x}} z(\mathbf{x}_\tau, \tau) \rangle \;. \tag{S13}$$

Moreover, using

$$\frac{\mathrm{d}}{\mathrm{d}\tau} z(\mathbf{x}_{\tau},\tau) = \mathbf{v}_{\tau} \cdot \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau},\tau) + \partial_{\tau} z(\mathbf{x}_{\tau},\tau) , \qquad (S14)$$

we obtain

$$\langle A_t B_t \rangle = \left\langle z(\mathbf{x}_t, t) - z(\mathbf{x}_0, 0) - \int_0^t \mathrm{d}\tau \partial_\tau z(\mathbf{x}_\tau, \tau) \right\rangle \,. \tag{S15}$$

With Eq. (S7) this proves via Cauchy-Schwarz $\langle A_t B_t \rangle^2 \leq \langle A_t^2 \rangle \langle B_t^2 \rangle$ the transport bound

$$\frac{\left\langle z(\mathbf{x}_t, t) - z(\mathbf{x}_0, 0) - \int_0^t \mathrm{d}\tau \partial_\tau z(\mathbf{x}_\tau, \tau) \right\rangle^2}{t \mathcal{D}^z(t)/k_\mathrm{B}T} \le T\Delta S_{\mathrm{tot}} \,. \tag{S16}$$

C. Overdamped limit

Although the underdamped results already contain the overdamped and Newtonian cases as limiting results, it is interesting to see that the derivation in the overdamped case greatly simplifies. Here, we have that $\nabla_{\mathbf{x}} \cdot \boldsymbol{\nu}(\mathbf{x}, \tau) \rho(\mathbf{x}, \tau) = -\partial_{\tau} \rho(\mathbf{x}, \tau)$ [16] such that Eq. (S16) directly follows from

$$\langle A_t B_t \rangle = \int_0^t d\tau \int d\mathbf{x} \left\{ \nabla_{\mathbf{x}} z(\mathbf{x}, \tau) \right\} \cdot \boldsymbol{\nu}(\mathbf{x}, \tau) \rho(\mathbf{x}, \tau) = \int_0^t d\tau \int d\mathbf{x} z(\mathbf{x}, \tau) \partial_\tau \rho(\mathbf{x}, \tau) = \left\langle z(\mathbf{x}_t, t) - z(\mathbf{x}_0, 0) - \int_0^t d\tau \partial_\tau z(\mathbf{x}_\tau, \tau) \right\rangle.$$
 (S17)

D. Generalization to velocity dependent variables

For optimal inference in the case of complete observation of $(\mathbf{x}_{\tau}, \mathbf{v}_{\tau})$ it is also useful to generalize Eq. (S1) to velocity-dependent observables $z(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau)$. We now show how one can generalize the proof starting with $-\langle A_t B_t \rangle = -\int_0^t d\tau \int d\mathbf{x} \int d\mathbf{v} \{ \nabla_{\mathbf{x}} z(\mathbf{x}, \mathbf{v}, \tau) \} \cdot \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \rho(\mathbf{x}, \mathbf{v}, \tau)$ as in Eq. (S9) where $\boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau) \equiv \mathbf{v}_{\tau} + k_{\rm B} T \mathbf{m}^{-1} \frac{\nabla_{\mathbf{v}} \rho(\mathbf{x}, \mathbf{v}, t)}{\rho(\mathbf{x}, \mathbf{v}, \tau)}$. While the first term $(\mathbf{v}_{\tau} \text{ in } \boldsymbol{\nu}(\mathbf{x}, \mathbf{v}, \tau))$ will similarly as in Eqs. (S9)-(S15) yield $\langle z(\mathbf{x}_t, \mathbf{v}_t, t) - z(\mathbf{x}_0, \mathbf{v}_0, 0) - \int_0^t d\tau [\partial_{\tau} z(\mathbf{x}_{\tau}, \mathbf{v}_{\tau} + \dot{\mathbf{v}} \partial_{\mathbf{v}} z(\mathbf{x}_{\tau}, \mathbf{v}_{\tau}, \tau)] \rangle^2$. While $\dot{\mathbf{v}}$ contains the white noise part that takes very high values, its occurrence in the average value should not be problematic for sufficient statistics since the high values average out for any finite time resolution. Therefore this expression will remain operationally accessible if enough trajectories are measured and one freely chooses z_{τ} to optimize inference.

The second term of $\boldsymbol{\nu}$ contributes in the presence of **v**-dependence in z_{τ} as

$$k_{\rm B}T \int_0^t \mathrm{d}\tau \int \mathrm{d}\mathbf{x} \int \mathrm{d}\mathbf{v} \left\{ \nabla_{\mathbf{x}} z(\mathbf{x}, \mathbf{v}, \tau) \right\} \cdot \mathbf{m}^{-1} \nabla_{\mathbf{v}} \rho(\mathbf{x}, \mathbf{v}, t)$$
$$= -k_{\rm B}T \int_0^t \mathrm{d}\tau \int \mathrm{d}\mathbf{x} \int \mathrm{d}\mathbf{v} \rho(\mathbf{x}, \mathbf{v}, t) \nabla_{\mathbf{v}} \cdot \mathbf{m}^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}, \mathbf{v}, \tau) \,. \tag{S18}$$

This term did not contribute before since we had $\nabla_{\mathbf{v}} z_{\tau} = 0$. However, this term can also be obtained from trajectories by computing the operationally accessible integral $\int d(\nabla_{\mathbf{x}} z) \cdot d\mathbf{v}$ where only the noise part contributes giving rise to the term in Eq. (S18) (see Ref. [17]).

Following these lines, one can generalize and apply the transport bound even to velocity dependent observables. Note however, that the result becomes slightly more complicated, and that much of the intuition on spatial transport under an external friction is lost in the case of explicit velocity-dependence.

III. DETAILS ON THE EXAMPLE IN FIG. 1 IN THE LETTER

Here, we give details on the example of the displaced trap considered in Fig. 1 in the Letter. Note that the overdamped case of this example including application of the TUR can be found in [11]. In Figure 1 we consider the parameters a = 5, m = 1 and y = 1. In the overdamped case, γ cancels out in the quality factor.

The underdamped equations of motion for this example for a linear force originating from a harmonic potential around 0 with a stiffness a read

$$d\begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} = -\underbrace{\begin{bmatrix} 0 & -1 \\ a\frac{\gamma}{m} & \frac{\gamma}{m} \end{bmatrix}}_{\equiv \mathbf{A}} \begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} d\tau + \underbrace{\frac{\sqrt{2k_{\mathrm{B}}T\gamma}}{m} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}}_{\equiv \boldsymbol{\sigma}} \mathbf{W}_{\tau}$$
$$\mathbf{D} = \frac{\boldsymbol{\sigma}\boldsymbol{\sigma}^{T}}{2} = \frac{k_{\mathrm{B}}T\gamma}{m^{2}} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$
(S19)

The steady-state covariance matrix Σ_s and time-dependent covariance $\Sigma(t)$ obey the Lyapunov equations (see, e.g., Supplemental Material of Ref. [18] for a short derivation)

$$\mathbf{A}\boldsymbol{\Sigma}_{s} + \boldsymbol{\Sigma}_{s}\mathbf{A}^{T} = 2\mathbf{D} = \boldsymbol{\sigma}\boldsymbol{\sigma}^{T}$$
$$\boldsymbol{\Sigma}(t) = \boldsymbol{\Sigma}_{s} + e^{-\mathbf{A}t}[\boldsymbol{\Sigma}(0) - \boldsymbol{\Sigma}_{s}]e^{-\mathbf{A}^{T}t}.$$
(S20)

We assume the particle to be equilibrated in the trap around $\langle x_0 \rangle = y$ before displacing the trap at time t = 0 to position x = 0. Therefore we have

$$\boldsymbol{\Sigma}(0) = \boldsymbol{\Sigma}_{\mathrm{s}} = k_{\mathrm{B}} T \begin{bmatrix} \frac{1}{a\gamma} & 0\\ 0 & \frac{1}{m} \end{bmatrix}, \qquad (S21)$$

which can be also obtained from equipartition.

We see from Eq. (S20) that $\Sigma(\tau) = \Sigma_s$ for all $\tau \ge 0$, i.e., the covariance remains unchanged, and in particular there will be no coupling of positions and velocities, and the density reads

$$\rho(x,v,\tau) = \frac{\sqrt{a\gamma m}}{2\pi} \exp\left[-\frac{a\gamma(x-\langle x_{\tau}\rangle)^2 + m(v-\langle v_{\tau}\rangle)^2}{2k_{\rm B}T}\right].$$
(S22)

The mean starting from $\langle x_0 \rangle = y$ and $\langle v_0 \rangle = 0$ is governed by

$$\begin{bmatrix} \langle x_\tau \rangle \\ \langle v_\tau \rangle \end{bmatrix} = \exp(-\mathbf{A}\tau) \begin{bmatrix} y \\ 0 \end{bmatrix} .$$
 (S23)

The decisive term for the entropy production is [see Eqs. (4) and (5) in the Letter]

$$T\Delta S_{\text{tot}} = \gamma \int_{0}^{t} d\tau \left\langle \left[v_{\tau} + \frac{k_{\text{B}}T}{m} \frac{\partial_{v}\rho(x_{\tau}, v_{\tau}, \tau)}{\rho(x_{\tau}, v_{\tau}, \tau)} \right]^{2} \right\rangle$$
$$= \gamma \int_{0}^{t} d\tau \left\langle \left[v_{\tau} - \frac{k_{\text{B}}T}{m} \frac{2m(v_{\tau} - \langle v_{\tau} \rangle)}{2k_{\text{B}}T} \right]^{2} \right\rangle$$
$$= \gamma \int_{0}^{t} d\tau \left\langle v_{\tau} \right\rangle^{2} . \tag{S24}$$

Note that in this simple example, the local mean velocity is not local, i.e., it does not depend on x_{τ} . In the overdamped case we have $\nu(\tau) = -aze^{-a\tau}$ (see Supplemental Material of Ref. [11]), but in the underdamped case the eigenvalues of **A** can become complex (signaling oscillations)

$$\mathbf{A} = \begin{bmatrix} 0 & -1\\ a\frac{\gamma}{m} & \frac{\gamma}{m} \end{bmatrix}$$
$$0 = \lambda^2 - \frac{\gamma}{m}\lambda + a\frac{\gamma}{m}$$
$$\lambda_{1,2} = \frac{\gamma}{2m} \left[1 \pm \sqrt{1 - \frac{4am}{\gamma}} \right].$$
(S25)

We obtain oscillations for $4am > \gamma$, i.e., for weak damping.

A. Saturation

We know from the Cauchy-Schwarz proof of Eq. (S16) that we obtain saturation for $\gamma \nu(x, v, \tau) = c \partial_x z(x, \tau)$, i.e., in this case the optimal observable z_{opt} is given by

$$\gamma\nu(x,v,\tau) = \gamma\nu(\tau) \stackrel{\text{as in Eq. (S24)}}{=} \gamma\langle v_{\tau} \rangle \equiv \gamma f(\tau)$$

$$z_{\text{opt}}(x,\tau) = \gamma x \langle v_{\tau} \rangle = \gamma x \underbrace{\left(\begin{bmatrix} 0\\1 \end{bmatrix} \cdot \exp(-\mathbf{A}\tau) \begin{bmatrix} y\\0 \end{bmatrix} \right)}_{=f(\tau)}.$$
(S26)

Since the Cauchy-Schwarz inequality becomes saturated for this choice, we know that the transport bound Eq. (S16) is saturated, allowing optimal inference of ΔS_{tot} .

This predicted saturation can be checked for this case by computing

$$T\Delta S_{\text{tot}} = \gamma \int_{0}^{t} d\tau f^{2}(\tau)$$
$$t\mathcal{D}^{z_{\text{opt}}}(t) = \gamma \int_{0}^{t} d\tau f^{2}(\tau) = T\Delta S_{\text{tot}}$$
$$\left\langle z_{\text{opt}}(\mathbf{x}_{t}, t) - z_{\text{opt}}(\mathbf{x}_{0}, 0) - \int_{0}^{t} d\tau \partial_{\tau} z_{\text{opt}}(\mathbf{x}_{\tau}, \tau) \right\rangle = \gamma \langle x_{t} \rangle f(t) - \gamma \langle x_{0} \rangle \underbrace{f(0)}_{=0} - \gamma \int_{0}^{t} d\tau \langle x_{\tau} \rangle \partial_{\tau} f(\tau) =$$
$$= \gamma \int_{0}^{t} d\tau f(\tau) \partial_{\tau} \langle x_{\tau} \rangle = \gamma \int_{0}^{t} d\tau f^{2}(\tau) = T\Delta S_{\text{tot}} .$$
(S27)

Hence we indeed have saturation since the transport bound (S16) here reads $\Delta S_{\text{tot}}^2 / \Delta S_{\text{tot}} \stackrel{\text{here}}{\leq} \Delta S_{\text{tot}}$.

B. Short-time limit

In Fig. 1b we see that the short-time limit of the quality factor is 3/4. This can be confirmed analytically as follows,

$$\begin{bmatrix} \langle x_{\tau} \rangle \\ \langle v_{\tau} \rangle \end{bmatrix} = \exp(-\mathbf{A}\tau) \begin{bmatrix} y \\ 0 \end{bmatrix} = \begin{bmatrix} y \\ 0 \end{bmatrix} - \tau \begin{bmatrix} 0 & -1 \\ a\frac{\gamma}{m} & \frac{\gamma}{m} \end{bmatrix} \begin{bmatrix} y \\ 0 \end{bmatrix} + \mathcal{O}(\tau^2) = \begin{bmatrix} y \\ -\frac{\tau a z \gamma}{m} \end{bmatrix} + \frac{\tau^2}{2} \begin{bmatrix} -\frac{\gamma a y}{m} \\ \frac{\gamma^2 a y}{m^2} \end{bmatrix} + \mathcal{O}(\tau^3) \,. \tag{S28}$$

Thus, the short-time entropy production reads

$$T\Delta S_{\rm tot} = \frac{\gamma^3 t^3 a^2 y^2}{3m^2} + \mathcal{O}(t^4) \,. \tag{S29}$$

The short-time transport reads

bound
$$= \frac{\gamma}{t} (\langle x_t \rangle - y)^2 = \frac{\gamma}{t} \frac{t^4}{4} \frac{\gamma^2 a^2 y^2}{m^2} + \mathcal{O}(t^4) = \frac{3}{4} T \Delta S_{\text{tot}} + \mathcal{O}(t^4),$$
 (S30)

such that the short-time quality factor is indeed 3/4.

C. TUR applied to underdamped dynamics

The TUR [9], and in particular the transient version [19], only applies to overdamped systems. Though there are versions for underdamped dynamics these are not practical to use and in particular not operationally accessible, since, e.g., the bound does not directly follow from measured trajectories but instead requires variations of many different parameters in the system [20, 21]. However, the overdamped transient TUR with corresponding quality factor (for simplicity we take the special case $J_t = x_t - x_0$)

$$Q_{\text{TUR transient}} = \frac{2(t\partial_t \langle x_t - x_0 \rangle)^2}{\operatorname{var}(x_t - x_0)} \frac{k_{\text{B}}}{\Delta S_{\text{tot}}}, \qquad (S31)$$

fails for underdamped dynamics.

To illustrate that, we again consider the example of the displaced trap and compute $\operatorname{var}(x_t - x_0) = \operatorname{var}(x_t^2) + \operatorname{var}(x_0^2) + 2\operatorname{cov}(x_t, x_0) = 2\left[\frac{k_{\mathrm{B}}T}{a\gamma} - \operatorname{cov}(x_t, x_0)\right]$, which similarly to the above follows from

$$d\left\langle \begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} \begin{bmatrix} x_0 & v_0 \end{bmatrix} \right\rangle = -\mathbf{A} \left\langle \begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} \begin{bmatrix} x_0 & v_0 \end{bmatrix} \right\rangle d\tau$$
$$\left\langle \begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} \begin{bmatrix} x_0 & v_0 \end{bmatrix} \right\rangle - \left\langle \begin{bmatrix} x_{\tau} \\ v_{\tau} \end{bmatrix} \right\rangle \left\langle \begin{bmatrix} x_0 & v_0 \end{bmatrix} \right\rangle = e^{-\mathbf{A}\tau} \mathbf{\Sigma}_{s}$$
$$\operatorname{cov}(x_{\tau}, x_0) = \left(e^{-\mathbf{A}\tau} \mathbf{\Sigma}_{s} \right)_{xx}$$
$$\operatorname{var}(x_t - x_0) = 2 \left[\left(\mathbf{1} - e^{-\mathbf{A}t} \right) \mathbf{\Sigma}_{s} \right]_{xx} .$$
(S32)

For Eq. (S31), we also require the expression

$$\partial_t \langle x_t - x_0 \rangle = \partial_t \langle x_t \rangle = \partial_t \left(\begin{bmatrix} 1\\0 \end{bmatrix} \cdot \exp(-\mathbf{A}t) \begin{bmatrix} y\\0 \end{bmatrix} \right) = -\left(\begin{bmatrix} 1\\0 \end{bmatrix} \cdot \exp(-\mathbf{A}t)\mathbf{A} \begin{bmatrix} y\\0 \end{bmatrix} \right) \,. \tag{S33}$$

Using these results, we plot the quality factor (S31) for the underdamped dynamics displaced trap in Fig. S2a. We see that the bound $Q \leq 1$ heavily violated, i.e., that the overdamped transient TUR (unlike the transport bound) is not valid for underdamped dynamics. This is partially explained due to the different time-evolution of the total entropy production, see Fig. S2b. However, for small friction the bound $Q \leq 1$ for the transient TUR is even violated for longer times where ΔS_{tot} almost agrees which is due to the different character of the dynamics.

IV. DERIVATION OF EQ. (6) IN THE LETTER

Here, we apply and rearrange the transport bound for the radius of gyration and structure factor, reproducing the bounds presented in Eq. (6) in the Letter.




FIG. S2. (a) Quality factor in Eq. (S31) evaluated for underdamped dynamics with different friction (same parameters as in Fig. 1 in the Letter). Values above 1 (dashed gray line) show that the overdamped TUR bound is violated for this underdamped example. (b) Time-evolution of the entropy production up to time t. As expected, underdamped and overdamped dynamics give rise to very different short-time behavior.)

A. Radius of gyration

First, to approach the radius of gyration consider the following observable z, which yields $NR_g^2 = \langle z_\tau \rangle$,

$$\vec{c} \equiv \frac{1}{N} \sum_{k=1}^{N} \vec{x}_{k}$$
$$z(\mathbf{x}) \equiv \sum_{j=1}^{N} (\vec{x}_{j} - \vec{c})^{2}$$
$$NR_{g}^{2}(t) = \langle z(\mathbf{x}_{t}) \rangle, \qquad (S34)$$

where the \vec{x}_j are the 3d position vectors of the individual beads. Here, we denote 3d vectors by arrows and boldface notation is only used for the 3N dimensional vectors that contain all bead positions.

To compute \mathcal{D}^z we need to compute derivatives, i.e., compute

$$\vec{\nabla}_l z(\mathbf{x}) = 2(\vec{x}_l - \vec{c}) - \frac{2}{N} \sum_{j=1}^N (\vec{x}_j - \vec{c}) \,. \tag{S35}$$

For the full 3N-dimensional gradient we obtain

$$\frac{[\nabla z(\mathbf{x})]^2}{4} = z(\mathbf{x}) - \frac{1}{N} \sum_{j,l=1}^N (\vec{x}_j - \vec{c}) \cdot (\vec{x}_l - \vec{c})$$
$$= z(\mathbf{x}) - \frac{1}{N} \left[\sum_{j=1}^N (\vec{x}_j - \vec{c}) \right]^2$$
$$= z(\mathbf{x}).$$
(S36)

This yields the transport bound for R_g^2 in Eq. (6) in the Letter. To further generalize, note that for time-dependent $\gamma(\tau)$, the factor $1/\gamma$ has to be kept inside the time integral. Moreover, note that if the beads have different γ_j we may formulate the bound in terms of the minimum of these γ_j .

B. Structure factor

Consider the following z_{τ} and its derivative to approach $S_t(\vec{q})$ and the corresponding $\mathcal{D}^z(t)$,

$$z(\mathbf{x}) \equiv \frac{1}{N} \sum_{j,k=1}^{N} \cos[\vec{q} \cdot (\vec{x}_j - \vec{x}_k)]$$

$$S_t(\vec{q}) = \langle z(\mathbf{x}_t) \rangle$$

$$\vec{\nabla}_j z(\mathbf{x}) = -\frac{2\vec{q}}{N} \sum_{k=1}^{N} \sin[\vec{q} \cdot (\vec{x}_j - \vec{x}_k)]$$

$$A_N \equiv \frac{N^2}{4\vec{q}^2} [\nabla z(\mathbf{x})]^2 = \sum_{j,k,l=1}^{N} \sin[\vec{q} \cdot (\vec{x}_j - \vec{x}_l)] \sin[\vec{q} \cdot (\vec{x}_j - \vec{x}_l)] .$$
(S37)

In order to express the derivative in observable terms, consider the expression

$$B_N \equiv \frac{N^2 - N z_{2\vec{q}}(\mathbf{x})}{4} = \frac{1}{4} \sum_{k,l=1}^N \left(1 - \cos[2\vec{q} \cdot (\vec{x}_k - \vec{x}_l)]\right) = \frac{1}{2} \sum_{j,k=1}^N \sin^2[\vec{q} \cdot (\vec{x}_k - \vec{x}_l)] \,. \tag{S38}$$

Introduce the notation $s_{jk} = -s_{kj} = \sin[\vec{q} \cdot (\vec{x}_j - \vec{x}_k)]$ and note that from relabeling indices we get

$$6(NB_N - A_N) = 3N \sum_{k,l=1}^{N} s_{kl}^2 - 6 \sum_{j,k,l=1}^{N} s_{jl} s_{jk}$$

$$= \sum_{j,k,l=1}^{N} \left(s_{jl}^2 + s_{lk}^2 + s_{kj}^2 + 2s_{jl} s_{lk} + 2s_{lk} s_{kj} + 2s_{kj} s_{jl} \right)$$

$$= \sum_{j,k,l=1}^{N} \left(s_{jl} + s_{lk} + s_{kj} \right)^2$$

$$\ge 0, \qquad (S39)$$

with equality for $q \to 0$. This implies an upper bound of $t\mathcal{D}^z = 4\vec{q}^2 \int_0^t d\tau \langle A_N(\tau) \rangle / N^2 \leq 4\vec{q}^2 \int_0^t d\tau \langle B_N(\tau) \rangle / N$ that implies the vector-version of the bound

$$T\Delta S_{\text{tot}} \ge \frac{\gamma [S_t(\vec{q}) - S_0(\vec{q})]^2}{\vec{q}^2 \int_0^t [N - S_t(2\vec{q})] d\tau}$$
(S40)

Note that this bound can be improved by using that we consider 3d space without assuming isotropy. Namely, by considering the above derivation for the observable

$$z_{q}(\mathbf{x}) \equiv \frac{1}{3} \left[z_{\vec{q}_{1}}(\mathbf{x}) + z_{\vec{q}_{2}}(\mathbf{x}) + z_{\vec{q}_{3}}(\mathbf{x}) \right]$$

$$S_{t}(q) \equiv \frac{1}{3} \left[S_{t}(\vec{q}_{1}) + S_{t}(\vec{q}_{2}) + S_{t}(\vec{q}_{3}) \right], \qquad (S41)$$

where we choose $\vec{q_i} = q\vec{e_i}$ where the $\vec{e_i}$ span an orthonormal basis [e.g., the vectors $(1, 0, 0)^T$, $(0, 1, 0)^T$, $(0, 0, 1)^T$], i.e., we average the structure factor over the three spatial dimensions. This allows to derive the extra factor of 3 compared to Eq. (S40) to obtain the bound in Eq. (6) in the Letter,

$$T\Delta S_{\text{tot}} \ge \frac{\gamma [S_t(\vec{q}) - S_0(\vec{q})]^2}{\vec{q}^2 \int_0^t [N - S_t(2\vec{q})] \mathrm{d}\tau},$$
(S42)

since the sum of the $\vec{q_i}$ values in $(\nabla z)^2$ gives rise to an extra factor of 3 originating from $[(\vec{q_1} + \vec{q_2} + \vec{q_3})/3]^2 = (1,1,1) \cdot (1,1,1)/9 = 1/3$.

As for R_g^2 , to further generalize, note that for time-dependent $\gamma(\tau)$, the factor $1/\gamma$ has to be kept inside the time integral. Moreover note that if the beads have different γ_j we may formulate the bound in terms of the minimum of these γ_j .

Note that we can obtain the R_g -bound in Eq. (6) in the Letter from Eq. (S42) using Guinier's law for $q \to 0$, which means that we can obtain the value of the R_g -bound from the $S_t(q)$ -bound (see $q \to 0$ in Fig. 2c,e in the Letter).

V. DETAILS ON THE EXAMPLE IN FIG. 2 IN THE LETTER

A. Computing the structure factor

The time-resolved structure factor $S_t(q)$ and the corresponding quality factor of the bounds for the examples considered in Fig. 2 in the Letter are computed using normal mode analysis (the time integral in the denominator of the bound and integrals that averaged over rotated lattices are computed numerically).

The overdamped Rouse model considered in Figs. 2b,c in the Letter can be written as a 3N dimensional Langevin equation for the vector $\mathbf{r} = (\vec{r}^1, \dots, \vec{r}^N)$ that reads (see [18, 22] for similar calculations)

$$\mathrm{d}\mathbf{r}_t = -\kappa \mathbf{k} \mathbf{r}_t \mathrm{d}t + \sqrt{2D} \mathrm{d}\mathbf{W}_t,\tag{S43}$$

with $D = k_{\rm B}T/\gamma$, trap stiffness κ , and the connectivity matrix **k** is a $3N \times 3N$ matrix that reads ($\mathbb{1}_3$ is the 3d unit matrix and all terms not shown are 0)

To model the nano-crystal, we consider the same harmonic model (i.e., linear drift) but for the deviations $\delta \vec{r}^{j}$ from the rest positions, adapt the connectivity as shown in the schematic in Fig. 2a in the Letter (see also [22] on how to adapt connectivity), and also add a confinement by adding $\kappa_{\text{confine}} \mathbb{1}_{3N}$ to $\kappa \mathbf{k}$ in the drift term that keeps the beads closer to the prescribed mean positions, and avoids center-of-mass diffusion of $\delta \mathbf{R}$.

The models are solved via normal mode analysis, i.e., by working in the basis that diagonalizes $\mathbf{A} = \kappa \mathbf{k}$ (or for the nano-crystal $\mathbf{A} = \kappa \mathbf{k} + \kappa_{\text{confine}} \mathbb{1}_{3N}$).

The parameters chosen for Fig. 2 in the Letter in units of $k_{\rm B} = 1$ are in (b,c) stiffness $\kappa = 1$, D = 1, quench from $T_0 = 2$ to T = 1, and in (d,e) stiffness $\kappa = 10$, confinement $\kappa_{\rm confine} = 10$, rest positions spaced by 1 on cubic lattice and quenched to rectangular lattice with spacings 1, 1.2, 0.7 in x, y, z-direction.

Now change the basis to work in normal modes, i.e., define \mathbf{Q} and $\mathbf{X} = \mathbf{Q}^T \mathbf{R}$ that fulfill $\mathbf{Q}^T \mathbf{A} \mathbf{Q} = \text{diag}(a_l)$, such that we have $d\vec{X}_t^l = -a_l \vec{X}_t^l dt + \sqrt{2D} d\vec{W}_t^l$ where the $W_t^{l,i}$ are mutually independent Wiener processes (see also Appendix F of [22]). This orthogonal change of basis exists since \mathbf{A} is symmetric.

For Gaussian displacements we have (see also [22])

$$S_{t}(\vec{q}) \equiv \frac{1}{N} \sum_{j,k=1}^{N} \left\langle e^{-i\vec{q} \cdot (\vec{r}_{t}^{j} - \vec{r}_{t}^{k})} \right\rangle = \frac{1}{N} \sum_{j,k=1}^{N} \exp\left[-i\vec{q} \cdot \left\langle \vec{r}_{t}^{j} - \vec{r}_{t}^{k} \right\rangle - \frac{\vec{q}^{2}}{6} \operatorname{var}\left(\vec{r}_{t}^{j} - \vec{r}_{t}^{k} \right) \right].$$
(S45)

Consider positions of beads $\vec{r}^{j} = \vec{r}^{j} + \delta \vec{r}^{j}$ with a fixed (deterministic) rest position \vec{r}^{j} and random fluctuations $\delta \vec{r}^{j}$. This includes the Rouse model since if we set $\vec{r}^{j} = 0$ and $\kappa_{\text{confine}} = 0$ (then we will have center-of-mass diffusion but this can be neglected in the comoving frame). Using properties of Langevin dynamics with linear drift in the **Q**-rotated basis (see Appendix F of [22]), we obtain for the Rouse chain where $\langle \vec{r}_{t}^{j} - \vec{r}_{t}^{k} \rangle = 0$ and we quench the temperature (entering the variance) that

$$\vec{r}_t^{\,j} = \sum_{\alpha=1}^N Q_{j\alpha} \vec{X}_t^{\alpha}$$
$$d\vec{X}_t^{\alpha} = -a_\alpha \vec{X}_t^{\alpha} dt + \sqrt{2D} d\vec{W}_t^{\alpha}$$
$$V_\alpha(t) \equiv \langle (\vec{X}_t^{\alpha})^2 \rangle = \frac{3D}{a_\alpha} \left[1 + \left(\frac{T_0}{T} - 1 \right) e^{-2a_\alpha t} \right] \quad \text{for } \alpha \neq 1$$
$$\langle \vec{X}_t^{\alpha} \cdot \vec{X}_t^{\varphi} \rangle = 0 \quad \text{for } \alpha \neq \varphi$$
$$\left\langle (\vec{r}_t^{\,j})^2 \right\rangle = \sum_{\alpha=1}^N V_\alpha(t) Q_{j\alpha}^2$$

$$\left\langle (\vec{r}_t^{\,j} - \vec{r}_t^{\,k})^2 \right\rangle = \sum_{\alpha=2}^N V_\alpha(t) \left[Q_{j\alpha}^2 + Q_{k\alpha}^2 - 2Q_{j\alpha}Q_{k\alpha} \right]$$
$$S_t(\vec{q}) \equiv \frac{1}{N} \sum_{j,k=1}^N \left\langle e^{-i\vec{q}\cdot(\vec{r}_t^{\,j} - \vec{r}_t^{\,k})} \right\rangle = \frac{1}{N} \sum_{j,k=1}^N e^{-\vec{q}^{\,2} \left\langle (\vec{r}_t^{\,j} - \vec{r}_t^{\,k})^2 \right\rangle/6} \,. \tag{S46}$$

On the other hand, for the nano-crystal where we do not quench temperature (and thus also not the variances), but instead the rest positions $\{\overline{\vec{r^j}}\}_{j\in[1,N]}$ (which we write as a vector in 3N-dimensional space $\overline{\mathbf{R}}$), we have

$$\begin{split} \delta \vec{r}_t^{j} &= \sum_{\alpha=1}^N Q_{j\alpha} \vec{X}_t^{\alpha} \\ \vec{\mu}_{\alpha}(t) &\equiv \langle \vec{X}_t^{\alpha} \rangle = e^{-a_\alpha t} \vec{\mu}_{\alpha}(0) \quad \text{for all } \alpha \\ \left\langle \vec{r}_t^{j} - \vec{r}_t^{k} \right\rangle &= \vec{r}^{j} - \vec{r}^{k} + \left\langle \delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right\rangle \\ - \mathrm{i} \vec{q} \cdot \left\langle \delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right\rangle &= \sum_{\alpha=1}^N (Q_{j\alpha} - Q_{k\alpha}) e^{-a_\alpha t} \vec{\mu}_{\alpha}(0) \\ \boldsymbol{\mu}(0) &= Q^T (\mathbf{\bar{R}}_{\mathrm{before quench}} - \mathbf{\bar{R}}) \\ \mathrm{var} \left(\delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right) &= \mathrm{var} \left(\sum_{\alpha=1}^N (Q_{j\alpha} - Q_{k\alpha}) \vec{X}_t^{\alpha} \right) \\ &= \sum_{\alpha, \varphi=1}^N (Q_{j\alpha} - Q_{k\alpha}) (Q_{j\varphi} - Q_{k\varphi}) \mathrm{cov}_{\mathrm{scalar}} \left(\vec{X}^{\alpha}, \vec{X}^{\varphi} \right) \\ V_{\alpha}(t) &\equiv \mathrm{var} \left(\vec{X}_t^{\alpha} \right) &= \frac{3D}{a_\alpha} \left[1 + \left(\frac{V_{\alpha}(0)}{3D/a_\alpha} - 1 \right) e^{-2a_\alpha t} \right] \quad \text{for all } \alpha \\ V_{\alpha}(0) &= 3D/a_\alpha \text{ for structure quench, such that here } V_{\alpha}(t) = V_{\alpha}(0) \\ \mathrm{cov}_{\mathrm{scalar}} \left(X_t^{\alpha}, \vec{X}_t^{\varphi} \right) &= \delta_{\alpha\varphi} V_{\alpha}(t) \\ \Rightarrow \quad \mathrm{var} \left(\delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right) &= \sum_{\alpha=1}^N V_{\alpha}(t) \left[Q_{j\alpha}^2 + Q_{k\alpha}^2 - 2Q_{j\alpha}Q_{k\alpha} \right] \\ S_t(\vec{q}) &= \frac{1}{N} \sum_{j,k=1}^N \exp \left[-\mathrm{i} \vec{q} \cdot \left(\vec{r}^{j} - \vec{r}^{k} \right) \right] \exp \left[-\mathrm{i} \vec{q} \cdot \left\langle \delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right\rangle - \frac{\vec{q}^2}{6} \mathrm{var} \left(\delta \vec{r}_t^{j} - \delta \vec{r}_t^{k} \right) \right] . \quad (847) \end{aligned}$$

This is numerically implemented for the described quench in $\overline{\vec{r}^{j}}$, and afterwards, averaged over all directions since the crystals are randomly oriented in the solution (which is equivalent to an average over all directions of \vec{q})

$$S_t(q) = \frac{1}{4\pi} \int_{|\vec{q}|=q} S_t(\vec{q}) \,. \tag{S48}$$

From these results for $S_t(q)$ we compute the bounds in Eq. (6) in the Letter. We could directly compute R_g^2 from $\langle (\vec{r}_t^j - \vec{r}_t^k)^2 \rangle$ from the normal mode analysis, but instead we simply determine the value of the R_g^2 bound form the $q \to 0$ limit of the $S_t(q)$ bound.

B. Computation of entropy production for the temperature quench

To evaluate the quality factor Q, we need to divide by the bound by the actual entropy production. As a shortcut to computing the entropy production we may use that the transient entropy production of systems approaching an

13

equilibrium state is given by the difference in excess free energy (since in this case, the total entropy production equals the non-adiabatic entropy production, see [18]), which in turn is computed from a Kullback-Leibler divergence [23].

Note that we do not consider the entropy production of the center-of-mass diffusion ($\alpha = 1$ mode) since it does not enter $S_t(\vec{q})$, and since we may assume that this diffusion is already equilibrated in the sample (in which case it does not produce any entropy). Thus, we have for the excess free energy [23, Eq. (6)]

$$EFE(t) = \frac{3k_{\rm B}}{2} \sum_{\alpha=2}^{N} \left\{ \left(\frac{T_0}{T} - 1\right) e^{-2a_{\alpha}t} - \ln\left[1 + \left(\frac{T_0}{T} - 1\right) e^{-2a_{\alpha}t}\right] \right\}$$
$$= \frac{3k_{\rm B}}{2} \sum_{\alpha=2}^{N} \left\{ \left(\frac{T_0}{T} - 1\right) e^{-2a_{\alpha}t} - T\ln\left[1 + \left(\frac{T_0}{T} - 1\right) e^{-2a_{\alpha}t}\right] \right\}$$
$$\Delta S_{\rm tot}[0, t] = \Delta S_{\rm tot}[0, \infty] - \Delta S_{\rm tot}[t, \infty] = EFE(0) - EFE(t) .$$
(S49)

C. Computation of entropy production for the structure quench

Again, the entropy production is computed from the Kullback-Leibler divergences, but this time for Gaussians with the same variance but different mean values, where we have

$$2D_{KL}[P(t)||P(\infty)] = (\boldsymbol{\mu}_t - \boldsymbol{\mu}_\infty)^T \boldsymbol{\Sigma}_\infty^{-1} (\boldsymbol{\mu}_t - \boldsymbol{\mu}_\infty).$$
(S50)

For the considered quench we have, see Eq. (S47), (note that we need to include $\alpha = 1$ here since $\kappa_{\text{confine}} \neq 0$ implies that there is no center-of-mass diffusion)

$$\mu_{\infty} = \mathbf{0}$$

$$\mu_{0} = Q^{T}(\bar{\mathbf{r}}_{before quench} - \bar{\mathbf{r}})$$

$$\mu_{t} = \exp(-\operatorname{diag}(\mathbf{a})t)\mu_{0}$$

$$\boldsymbol{\Sigma}_{\infty}^{-1} = \operatorname{diag}(\mathbf{a})/D$$

$$\mathrm{EFE}(t)/T = D_{KL}[P(t)||P(\infty)] = \frac{1}{2D}\sum_{\alpha=1}^{N} a_{\alpha} \exp(-2a_{\alpha}t)\vec{\mu}_{\alpha}(0)^{2}$$

$$\Delta S_{tot} = \mathrm{EFE}(0)/T - \mathrm{EFE}(t)/T = \frac{1}{2D}\sum_{\alpha=1}^{N} a_{\alpha}[1 - \exp(-2a_{\alpha}t)]\vec{\mu}_{\alpha}(0)^{2}.$$
(S51)

VI. DETAILS ON THE HANDLING OF D^{z}

If we measure the full dynamics x_{τ} and know, or even choose, z_{τ} , then we can directly evaluate \mathcal{D}^z . Moreover, if we know z_{τ} but do *not* measure x_{τ} , we can often express \mathcal{D}^z in terms of $\langle z_{\tau} \rangle$, or bound it by this and constants (as for R_q^2 and $S_t(q)$ above).

However, in the challenging case that we only measure z_{τ} but do *not* know what function of the underlying dynamics x_{τ} it is (this would, e.g., be the case if we measure the dynamics of a macromolecule along some reaction coordinate), then we can often obtain \mathcal{D}^z from the short-time fluctuations of z_{τ} .

A. Overdamped dynamics

For overdamped dynamics, we obtain $\mathcal{D}^{z}(t)$ directly from the from short-time fluctuations of z_{τ} (see also Ref. [24]),

$$dz_{\tau} = \nabla_{\mathbf{x}} z_{\tau} \cdot d\mathbf{x}_{\tau} + \dot{z}_{\tau} dt = k_{\rm B} T \nabla_{\mathbf{x}} z_{\tau} \cdot \gamma^{-1} d\mathbf{W}_{\tau} + O(d\tau)$$

$$k_{\rm B} T \left\langle \nabla_{\mathbf{x}} z_{\tau} \cdot \gamma^{-1} \nabla_{\mathbf{x}} z_{\tau} \right\rangle = \frac{\operatorname{var}[dz_{\tau}]}{2d\tau}$$

$$k_{\rm B} T t \mathcal{D}^{z}(t) = \int_{0}^{t} d\tau \frac{\operatorname{var}[dz_{\tau}]}{2d\tau} .$$
(S52)

B. Underdamped dynamics

For simplicity, assume γ and m to be scalars and z_{τ} to not explicitly depend on time. Then, we can write

$$\frac{\mathrm{d}}{\mathrm{d}\tau} z(\mathbf{x}_{\tau}) = \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \mathbf{v}_{\tau}$$
$$\mathrm{d} \left[\frac{\mathrm{d}}{\mathrm{d}\tau} z(\mathbf{x}_{\tau}) \right] = \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \mathrm{d} \mathbf{v}_{\tau} + \mathrm{d} [\nabla_{\mathbf{x}} z(\mathbf{x}_{\tau})] \cdot \mathbf{v}_{\tau}$$
$$= \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \frac{\sqrt{2k_{\mathrm{B}}T\gamma}}{m} \mathrm{d} \mathbf{W}_{\tau} + O(\mathrm{d}t)$$
$$\frac{\mathrm{var} \left(\mathrm{d} \left[\frac{\mathrm{d}}{\mathrm{d}\tau} z(\mathbf{x}_{\tau}) \right] \right)}{2k_{\mathrm{B}}T \mathrm{d}t} = \left(\frac{\gamma}{m} \right)^{2} \left\langle \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \gamma^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \right\rangle . \tag{S53}$$

Hence, if we know γ/m , or can determine it from another experiment, we can again obtain $t\mathcal{D}^z(t)$ from observations of z_{τ} .

If we can observe z_{τ} while the system settles into a steady state, we can even infer γ/m from z_{τ} itself. We here distinguish two cases. First, if the dynamics approach equilibrium, then we can obtain γ/m from comparing the equilibrium fluctuations $\operatorname{var}_{eq}(dz)/dt^2 \sim \langle \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \gamma^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \rangle \gamma/m$ (see Supplemental Material of Ref. [24]) to the previously mentioned equilibrium result for var $\left(d \left[\frac{d}{d\tau} z(\mathbf{x}_{\tau}) \right] \right) \sim \langle \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \gamma^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \rangle (\gamma/m)^2$.

the previously mentioned equilibrium result for var $\left(d \left[\frac{d}{d\tau} z(\mathbf{x}_{\tau}) \right] \right) \sim \left\langle \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \cdot \gamma^{-1} \nabla_{\mathbf{x}} z(\mathbf{x}_{\tau}) \right\rangle (\gamma/m)^2$. Second, if approach a non-equilibrium steady state, we can still bound the thermalization time $\tau_{\text{thermalization}} = m/\gamma$ since it cannot be faster than the relaxation time scale $\tau_{\text{relax}}^{-1} = -\lim_{t \to \infty} t^{-1} \ln C_{zz}(t) \geq \tau_{\text{thermalization}}$, where $C_{zz}(t) = \langle z_t z_0 \rangle - \langle z_t \rangle \langle z_0 \rangle$

VII. FRET BOUND

For the FRET efficiency E_t measured between donor and acceptor chromophores at positions \vec{x}_1 and \vec{x}_2 with Förster radius R_0 we can compute

$$E_{t} = \left\langle \left[1 + \left(\frac{\vec{x}_{1} - \vec{x}_{2}}{R_{0}} \right)^{6} \right]^{-1} \right\rangle = \left\langle z(\mathbf{x}) \right\rangle$$

$$z(\mathbf{x}) = \left[1 + \left(\frac{\vec{x}_{1} - \vec{x}_{2}}{R_{0}} \right)^{6} \right]^{-1} \leq 1$$

$$\vec{\nabla}_{1}z(\mathbf{x}) = -z^{2}(\mathbf{x})6 \left(\frac{\vec{x}_{1} - \vec{x}_{2}}{R_{0}} \right)^{5} \frac{\vec{\nabla}_{1} \cdot \vec{x}_{1}}{R_{0}} = -\frac{2z^{2}(\mathbf{x})}{R_{0}} \left(\frac{\vec{x}_{1} - \vec{x}_{2}}{R_{0}} \right)^{5}$$

$$[\nabla z(\mathbf{x})]^{2} = [\vec{\nabla}_{1}z(\mathbf{x})]^{2} + [\vec{\nabla}_{2}z(\mathbf{x})]^{2} = \frac{8z^{4}(\mathbf{x})}{R_{0}^{2}} \left(\frac{\vec{x}_{1} - \vec{x}_{2}}{R_{0}} \right)^{10} = \frac{8z^{4}(\mathbf{x})}{R_{0}^{2}} \left(\frac{1}{z(\mathbf{x})} - 1 \right)^{5/3} = \frac{8z^{7/3}(\mathbf{x})}{R_{0}^{2}} [1 - z(\mathbf{x})]^{5/3} \leq \frac{8}{R_{0}^{2}}.$$
(S54)

Using this simple approximation, we obtain from the transport bound (S16)

$$T\Delta S_{\text{tot}} \ge R_0^2 \gamma [E_t - E_0]^2 / 8t$$
. (S55)

This bound can be easily improved, e.g., by bounding z and 1 - z above by the maximally measured values instead of by 1.

^[1] A. Dechant and S.-i. Sasa, Entropic bounds on currents in Langevin systems, Phys. Rev. E 97, 062101 (2018).

^[2] M. P. Leighton and D. A. Sivak, Dynamic and thermodynamic bounds for collective motor-driven transport, Phys. Rev. Lett. 129, 118102 (2022).

^[3] M. P. Leighton and D. A. Sivak, Inferring subsystem efficiencies in bipartite molecular machines, Phys. Rev. Lett. 130, 178401 (2023).

 ^[4] J.-D. Benamou and Y. Brenier, A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem, Numer. Math. 84, 375 (2000).

- [5] T. Van Vu and K. Saito, Thermodynamic unification of optimal transport: Thermodynamic uncertainty relation, minimum dissipation, and thermodynamic speed limits, Phys. Rev. X 13, 011013 (2023).
- [6] A. Dechant, Minimum entropy production, detailed balance and Wasserstein distance for continuous-time Markov processes, J. Phys. A: Math. Theor. 55, 094001 (2022).
- [7] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, Rep. Prog. Phys. 75, 126001 (2012).
- [8] U. Seifert, Stochastic thermodynamics: From principles to the cost of precision, Physica (Amsterdam) 504A, 176 (2018).
 [9] A. C. Barato and U. Seifert, Thermodynamic uncertainty relation for biomolecular processes, Phys. Rev. Lett. 114, 158101
- (2015).
- [10] A. Dechant and S.-i. Sasa, Improving thermodynamic bounds using correlations, Phys. Rev. X 11, 041061 (2021).
- [11] C. Dieball and A. Godec, Direct route to thermodynamic uncertainty relations and their saturation, Phys. Rev. Lett. 130, 087101 (2023).
- [12] C. W. Gardiner, Handbook of Stochastic Methods for Physics, Chemistry, and the Natural Sciences, 2nd ed. (Springer-Verlag, Berlin New York, 1985).
- [13] S. Pigolotti, I. Neri, É. Roldán, and F. Jülicher, Generic properties of stochastic entropy production, Phys. Rev. Lett. 119, 140604 (2017).
- [14] S. K. Manikandan, D. Gupta, and S. Krishnamurthy, Inferring entropy production from short experiments, Phys. Rev. Lett. 124, 120603 (2020).
- [15] P. Pietzonka, Classical pendulum clocks break the thermodynamic uncertainty relation, Phys. Rev. Lett. 128, 130606 (2022).
- [16] H. Risken, *The Fokker-Planck Equation* (Springer Berlin Heidelberg, 1989).
- [17] J. Lyu, K. J. Ray, and J. P. Crutchfield, Entropy production by underdamped Langevin dynamics (2024).
- [18] C. Dieball, G. Wellecke, and A. Godec, Asymmetric thermal relaxation in driven systems: Rotations go opposite ways, Phys. Rev. Res. 5, L042030 (2023).
- [19] A. Dechant and S.-i. Sasa, Current fluctuations and transport efficiency for general Langevin systems, J. Stat. Mech., 063209 (2018).
- [20] C. Kwon and H. K. Lee, Thermodynamic uncertainty relation for underdamped dynamics driven by time-dependent protocols, New J. Phys. 24, 013029 (2022).
- [21] S. Lee, D.-K. Kim, J.-M. Park, W. K. Kim, H. Park, and J. S. Lee, Multidimensional entropic bound: Estimator of entropy production for Langevin dynamics with an arbitrary time-dependent protocol, Phys. Rev. Res. 5, 013194 (2023).
- [22] C. Dieball, D. Krapf, M. Weiss, and A. Godec, Scattering fingerprints of two-state dynamics, New J. Phys. 24, 023004 (2022).
- [23] A. Lapolla and A. Godec, Faster uphill relaxation in thermodynamically equidistant temperature quenches, Phys. Rev. Lett. 125, 110602 (2020).
- [24] A. Dechant, J. Garnier-Brun, and S.-i. Sasa, Thermodynamic bounds on correlation times, Phys. Rev. Lett. 131, 167101 (2023).