New method for the solution of the two-body Dirac equation for the positronium bound states

E.M. Tursunov[∗](#page-0-0)

Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan and National University of Uzbekistan, 100174 Tashkent, Uzbekistan

Sh.G. Norbutaev[†](#page-0-1)

Institute of Nuclear Physics, Academy of Sciences, 100214, Ulugbek, Tashkent, Uzbekistan

B.A. Fayzullaev[‡](#page-0-2)

National University of Uzbekistan, 100174 Tashkent, Uzbekistan

Abstract

A new theoretical method is developed for the solution of the two-body bound-state Dirac equation for positronium. Only Coulomb potential was included in the Dirac Hamiltonian. It is shown that the two-body Dirac Hamiltonian can be written in the Hermitian matrix form of the 4×4 size and diagonalized in the momentum-state representation. Numerical results for the energy spectrum of the para- and ortho-positronium ground states performed within the variational method using the harmonic oscillator basis functions are in good agreement with a high-precision finite-element method of T.C. Scott et al. After the Fourier transformation into the coordinatestate representation the bound state wave functions of the para-Ps and ortho-Ps do not contain any singularity at the origin in contrast to above mentioned method. The weights of the large-small and small-large components of the ground state wave functions are estimated to be of order 10^{-6} , while the weight of the small-small component is of order 10^{-12} .

[∗] tursune@inp.uz

[†] norbutaev@inp.uz

[‡] bfayzullaev@nuu.uz

I. INTRODUCTION

A new generation of the high sensitivity and multi-photon total-body positron emission tomography systems opens perspectives for clinical applications of positronium (Ps) in medicine [\[1\]](#page-11-0). On the other hand, Ps is the most important object for the precision tests of the violation of the discrete symmetries such as C (charge conjugation), P (spatial parity), T (time reversal) and their combinations $[2-8]$. The spin-singlet bound states of the electron e^- and its antiparticle, the positron e^+ present para-positronium (p-Ps), at the same time the spin-triplet bound states present the ortho-positronium $(o-Ps)$.

The most important spectroscopic properties of the positronium, such as energy-level intervals are reproduced with a high precision at order α^6 within the non-relativistic QED $[9-13]$ $[9-13]$ based on the solutions of the Schrödinger bound-state equation. The explicit relativistic invariance built into QED is essential for particle physics but leads to unnecessary complications when QED is applied to Coulombic bound states, which deals with the energy scales small compared to m_ec^2 [\[2](#page-11-1)].

The two-body Dirac equation (TBDE) for positronium was studied in Ref. [\[14](#page-11-5)] using only the Coulomb interaction potential. In Ref. [\[15](#page-11-6)] the sixteen-component, no-pair Dirac-Coulomb-Breit equation, derived from the Bethe-Salpeter equation, is solved in a variational procedure using Gaussian-type basis functions for positronium, muonium, hydrogen atom, and muonic hydrogen. Another type of relativistic Schrödinger equation was obtained and solved based on the Pauli reduction of the TBDE [\[16,](#page-11-7) [17](#page-11-8)].

Recent studies of properties of anomalous bound states within the TBDE formalism [\[18\]](#page-11-9) have demonstrated that there are still many open questions for the structure of positronium. It was shown that the radial components of the Positronium bound-state wave-function [\[14\]](#page-11-5) contain a singularity near the origin and differ from their Pauli approximations. This difference is due to mixing of the ground-state wave function with the anomalous boundstate wave functions near the origin. The solutions of the Bethe-Salpeter equation do not show any mixing with anomalous states and do not have a singularity near the origin [\[19\]](#page-11-10).

The question is whether a bound state solution of the two-body Dirac equation with a pure Coulomb interaction potential can be free of mixing with an anomalous state. The aim of present work is to study this question within an alternative method. We solve the TBDE for the positronium bound states within a new theoretical method. The method is based

on the Hermitian Dirac Hamiltonian in the form of 4×4 matrix, each element of which represents a four-component operator. The solution of the TBDE for the positronium state with total momentum J and its projection M with the spin-orbital couplings $\vec{J} = \vec{I} + \vec{S}$ and $\vec{S} = \vec{S}_1 + \vec{S}_2$, $S_1 = S_2 = 1/2$, can be expressed in the form of a column of four components. Each of the large-large (LL), large-small (LS), small-large (SL) and smallsmall (SS) components of the eigen function represents a four-component spinor. The radial parts of these components are expressed as linear combination of the basic functions. This form of the Hermitian Dirac Hamiltonian contains a term $\vec{\sigma}_{\vec{p}},$ responsible for the coupling of the main large-large component with the large-small and small-large components. Numerical calculations have been performed within the variational method on the harmonic oscillator basic functions. The matrix elements of the most important operators including $\vec{\sigma}_{\vec{p}}$, are calculated in momentum state representation and are all real numbers in contrast to the coordinate space representation. As a results we come to the system of algebraic equations which is solved numerically [\[20](#page-11-11)[–23](#page-12-0)].

In Section II we give the main formalism of the method. Section III deals with numerical results and the conclusions are presented in the last Section.

II. SOLUTION OF THE TWO-BODY BOUND STATE DIRAC EQUATION

A. Two-body Dirac equation for positronium bound states

The Dirac Hamiltonian for the two-body bound system interacting via Coulomb potential $(\hbar = c = 1)$ is written in the form [\[14\]](#page-11-5)

$$
\hat{\mathbf{H}} = \vec{\alpha}_1 \times \mathbf{I}_4^{(2)} \cdot \vec{\mathbf{p}}_1 + \mathbf{I}_4^{(1)} \times \vec{\alpha}_2 \cdot \vec{\mathbf{p}}_2 + \beta_1 \times \mathbf{I}_4^{(2)} m_1 + \mathbf{I}_4^{(1)} \times \beta_2 m_2 + \mathbf{I}_{16} V(r_{12}), \tag{1}
$$

where the Dirac matrices

$$
\vec{\alpha} = \beta \vec{\gamma} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}, \qquad \beta = \gamma^0 = \begin{pmatrix} \mathbf{I}_2 & 0 \\ 0 & -\mathbf{I}_2 \end{pmatrix}, \tag{2}
$$

and \mathbf{I}_n is the unit $n \times n$ matrix operator, $V(r_{12})$ is the Coulomb potential. Index 1 (2) of the operator means that this operator acts on the wave function of the first (second) particle. In order to write the Hamiltonian in the relative coordinates we introduce the radius-vectors of the center of mass \vec{R} and the relative-motion \vec{r} of the positron-electron two-body system:

$$
\vec{\mathbf{R}} = \frac{1}{2}(\vec{\mathbf{r}}_1 + \vec{\mathbf{r}}_2), \quad \vec{\mathbf{r}} = \vec{\mathbf{r}}_1 - \vec{\mathbf{r}}_2.
$$

With the assumption $\vec{P}_R = 0$, one can write the total Dirac Hamiltonian of the relative motion in terms of the relative momentum \vec{p} and coordinate \vec{r} :

$$
\hat{\mathbf{H}} = \left\{ \vec{\boldsymbol{\alpha}}_1 \times \mathbf{I}_4^{(2)} - \mathbf{I}_4^{(1)} \times \vec{\boldsymbol{\alpha}}_2 \right\} \cdot \vec{\mathbf{p}} + \beta_1 \times \mathbf{I}_4^{(2)} m_1 + \mathbf{I}_4^{(1)} \times \beta_2 m_2 + \mathbf{I}_{16} V(r). \tag{3}
$$

.

Using the standard representation of the Dirac gamma matrices, the kinetic part of the Hamiltonian can be written in the form of symmetric matrix:

$$
\left\{\vec{\alpha}_1 \times \mathbf{I}_4^{(2)} - \mathbf{I}_4^{(1)} \times \vec{\alpha}_2\right\} \cdot \vec{\mathbf{p}} = \begin{pmatrix}\n0 & -\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \vec{\mathbf{p}} & \vec{\sigma}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & 0 \\
-\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \vec{\mathbf{p}} & 0 & 0 & \vec{\sigma}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} \\
\vec{\sigma}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & 0 & 0 & -\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \vec{\mathbf{p}} \\
0 & \vec{\sigma}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & -\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \vec{\mathbf{p}} & 0\n\end{pmatrix}
$$
\n(4)

Then the total Hermitian Dirac Hamiltonian can be written in the form of 4×4 matrix, each element of which represents the 4×4 matrix operator acting on the four-component spinor:

$$
\hat{\mathbf{H}} = \begin{pmatrix}\n\boldsymbol{\delta}_1 + \mathbf{I}_4 V(r_{12}) & -\mathbf{I}_2^{(1)} \times \vec{\boldsymbol{\sigma}}_2 \cdot \vec{\mathbf{p}} & \vec{\boldsymbol{\sigma}}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & 0 \\
-\mathbf{I}_2^{(1)} \times \vec{\boldsymbol{\sigma}}_2 \cdot \vec{\mathbf{p}} & \boldsymbol{\delta}_2 + \mathbf{I}_4 V(r_{12}) & 0 & \vec{\boldsymbol{\sigma}}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} \\
\vdots & \vdots & \ddots & \vdots \\
\boldsymbol{\sigma}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & 0 & -\boldsymbol{\delta}_2 + \mathbf{I}_4 V(r_{12}) & -\mathbf{I}_2^{(1)} \times \vec{\boldsymbol{\sigma}}_2 \cdot \vec{\mathbf{p}} \\
0 & \vec{\boldsymbol{\sigma}}_1 \times \mathbf{I}_2^{(2)} \cdot \vec{\mathbf{p}} & -\mathbf{I}_2^{(1)} \times \vec{\boldsymbol{\sigma}}_2 \cdot \vec{\mathbf{p}} & -\boldsymbol{\delta}_1 + \mathbf{I}_4 V(r_{12})\n\end{pmatrix},
$$
\n(5)

where the diagonal elements read $\boldsymbol{\delta}_1 = \mathbf{I}_2^{(1)} \times \mathbf{I}_2^{(2)}$ $\delta_2^{(2)} (m_1 + m_2), \quad \boldsymbol{\delta}_2 = \textbf{I}_2^{(1)} \times \textbf{I}_2^{(2)}$ $2^{(2)}(m_1 - m_2),$ $V(r_{12}) = \alpha$ r . The identity operators $I_2^{(1)}$ $\mathbf{I}_{2}^{(1)},\;\mathbf{I}_{2}^{(2)}$ $_2^{(2)}$ of the 2×2 size act only on the spin function of the first and second particles, respectively. The above form of the Hermitian Dirac Hamiltonian matrix differs from the non-Hermitian matrix Hamiltonian of Ref. [\[14\]](#page-11-5). However, they coincide with each other in the 16×16 matrix form.

It can be proven that the operators $-\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \hat{\vec{p}}$ and $\vec{\sigma}_1 \times \mathbf{I}_2^{(2)}$ $_{2}^{(2)}\cdot\hat{\vec{p}}$ of the Dirac Hamiltonian Eq.[\(5\)](#page-3-0), where $\hat{\vec{p}} = \vec{p}/p$ is the unit momentum vector, while acting on the spherical tensor $\mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1,\vec{\xi}_2;\hat{\vec{p}})$, can change the spin S and the orbital momentum ℓ of the system by one:

$$
(-\mathbf{I}_{2}^{(1)} \times \vec{\sigma}_{2} \cdot \hat{\vec{p}}) \mathcal{Y}_{(S_{1},S_{2})S\ell}^{JM}(\vec{\xi}_{1},\vec{\xi}_{2};\hat{\vec{p}}) = (-1)^{\ell+\ell^{+}} \sum_{h,S'}[h] ([S][S'])^{\frac{1}{2}} \left\{ \begin{array}{ll} S_{1} & S_{2} & S \\ \ell & J & h \end{array} \right\}
$$

\n
$$
\cdot \left\{ \begin{array}{ll} S_{1} & S_{2} & S' \\ \ell^{+} & J & h \end{array} \right\} \mathcal{Y}_{(S_{1},S_{2})S'\ell^{+}}^{JM}(\vec{\xi}_{1},\vec{\xi}_{2};\hat{\vec{p}}),
$$

\n
$$
(\vec{\sigma}_{1} \times \mathbf{I}_{2}^{(2)} \cdot \hat{\vec{p}}) \mathcal{Y}_{(S_{1},S_{2})S\ell}^{JM}(\vec{\xi}_{1},\vec{\xi}_{2};\hat{\vec{p}}) = (-1)^{\ell+\ell^{+}+1} \sum_{h,S'} (-1)^{S+S'}[h] ([S][S'])^{\frac{1}{2}} \left\{ \begin{array}{ll} S_{1} & S_{2} & S \\ \ell & J & h \end{array} \right\}
$$

\n
$$
\cdot \left\{ \begin{array}{ll} S_{1} & S_{2} & S' \\ \ell^{+} & J & h \end{array} \right\} \mathcal{Y}_{(S_{1},S_{2})S'\ell^{+}}^{JM}(\vec{\xi}_{1},\vec{\xi}_{2};\hat{\vec{p}}),
$$

\n
$$
(7)
$$

where $\ell^{\pm} = \ell \pm 1$, $S' = 0$ and/or 1, and $\ell^{\pm} + \vec{S}' = \vec{J}$. The last equations Eq.[\(6\)](#page-4-0) and Eq.[\(7\)](#page-4-1) contain the most important result for the structure of positronium within the two-body Dirac equation formalism.

B. Variational method on a harmonic-oscillator basis

The solution of the two-body Dirac equation for the positronium state with total momentum J and its projection M, where $\vec{\mathbf{J}} = \vec{\boldsymbol{\ell}} + \vec{\mathbf{S}}$ and $\vec{\mathbf{S}} = \vec{\mathbf{S}}_1 + \vec{\mathbf{S}}_2$,

$$
\left(\hat{\mathbf{H}} - E\right) \Psi^{JM}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{r}}) = 0. \tag{8}
$$

can be expressed as a column of four functions

$$
\Psi^{JM}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) = \begin{pmatrix} \Psi_{LL}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) \\ \Psi_{LS}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) \\ \Psi_{SL}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) \\ \Psi_{SS}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) \end{pmatrix},
$$
\n(9)

where each of the large-large (LL), large-small (LS), small-large (SL) and small-small (SS) components are four-component spinors and expressed as linear combination of the harmonic-oscillator basic functions with corresponding spin-angular parts. For the fixed total momentum J, possible spin-orbital couplings include terms with $\ell = J - 1$, $\ell = J$, and $\ell = J + 1$, and spins $S = 0$ and $S = 1$. From Eq.[\(6\)](#page-4-0) and the form of the Hamiltonian of Eq.[\(5\)](#page-3-0) one can find that the main LL- and the last SS- components of the eigen-state contain contributions from the same spin-orbital (ℓ, S) channels and differ from the LS- and SL- components containing contributions from channels (ℓ^{\pm}, S') , whose orbital components differ by one. In particular, in the ground state of the para-Ps with $J^{\pi} = 0^{+}$ the LL- and SS- components consist of only $(\ell, S) = (0, 0)$ spin-orbital channel, while the LS- and SLcomponents contain contributions only from the $(\ell^{\pm}, S') = (1, 1)$ channel. In the case of the ortho-Ps with $J^{\pi} = 1^{+}$, the main LL- and smallest SS-components contain contributions from the $(\ell, S) = (0, 1)$ and $(2, 1)$ spin-orbital channels, while the LS- and SL-components contain contributions from the $(\ell^{\pm}, S') = (1, 0)$ and $(1, 1)$ channels.

For the solution of the Dirac equation Eq.[\(8\)](#page-4-2) a column of probe wave functions of Eq.[\(9\)](#page-4-3) is expanded in a complete set of orthonormal harmonic oscillator states with which we calculate the matrix elements of the Hamiltonian of Eq.[\(5\)](#page-3-0).

The harmonic oscillator basis functions $\Psi_{n\ell S}^{JM}(\vec{\xi}_1,\vec{\xi}_2;\vec{r})$ with the variational oscillator parameter a_0 :

$$
\Psi_{n\ell S}^{JM}(\vec{\xi_1}, \vec{\xi_2}; \vec{\mathbf{r}}) = R_{n\ell}(a_0; r) \cdot \mathcal{Y}_{\ell S}^{JM}(\vec{\xi_1}, \vec{\xi_2}; \hat{\vec{\mathbf{r}}}), \tag{10}
$$

where $\mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1,\vec{\xi}_2;\hat{\vec{r}})$ is the spherical tensor $\mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1,\vec{\xi}_2;\hat{\vec{r}}) = \left\{ \left\{ \chi^{S_1}(\vec{\xi}_1) \times \chi^{S_2}(\vec{\xi}_2) \right\} \right\}$ $\left. \begin{array}{c} S \end{array} \right\} \times \mathbf{Y}_\ell(\hat{\vec{\mathbf{r}}}) \bigg\} \bigg\}^J_{\Lambda}$ \overline{M} , with $S_1 = S_2 = \frac{1}{2}$ $\frac{1}{2}$. The radial part of the basis function is presented as

$$
R_{nl}(a_0; r) = \left[\frac{2(n!)}{a_0^3 \Gamma(n+\ell+\frac{3}{2})}\right]^{\frac{1}{2}} \left(\frac{r}{a_0}\right)^{\ell} \exp\left(-\frac{r^2}{2a_0^2}\right) L_n^{\ell+\frac{1}{2}} \left(r^2/a_0^2\right).
$$
 (11)

where $L_n^{\ell+\frac{1}{2}}(r^2)$ are the Laguerre polynomials:

$$
L_n^{\ell+\frac{1}{2}}(r^2) = \sum_{k=0}^n \frac{(-1)^k}{k!} \frac{\Gamma(n+\ell+3/2)}{(n-k)!\Gamma(k+\ell+3/2)} \cdot r^{2k}.
$$

Accordingly, the harmonic oscillator basis functions $\Psi_{n\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \vec{r})$ are normalized to unity. The Fourier transformation of the coordinate-space basis functions is the same state multiplied by a phase [\[20](#page-11-11), [21](#page-11-12)]:

$$
\Psi_{n\ell S}^{JM}(\vec{\mathbf{p}}) = (2\pi)^{-3/2} \int d^3 r \Psi_{n\ell S}^{JM}(\vec{\mathbf{r}}) \exp(-i\vec{\mathbf{p}}\vec{\mathbf{r}}) = (-i)^{2n+\ell} R_{n\ell}(a_0; p) \cdot \mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \hat{\vec{\mathbf{p}}}), \quad (12)
$$

where the harmonic oscillator wave functions in momentum representation is defined by the equation

$$
R_{nl}(a_0; p) = \left[\frac{2(n!)a_0^3}{\Gamma(n+\ell+\frac{3}{2})}\right]^{\frac{1}{2}}(pa_0)^{\ell} \exp\left(-\frac{p^2a_0^2}{2}\right)L_n^{\ell+\frac{1}{2}}(p^2a_0^2).
$$
 (13)

The matrix elements of the Hamiltonian Eq.[\(5\)](#page-3-0) are calculated in the momentum state representation [\[20,](#page-11-11) [21\]](#page-11-12), because in this case the overlaps are all real, in contrast to the coordinate space representation, where the contribution of the operators $-\mathbf{I}_2^{(1)} \times \vec{\sigma}_2 \cdot \hat{\vec{p}}$ and $\vec{\boldsymbol{\sigma}}_1\!\times\!\textbf{I}^{(2)}_2$ $^{(2)}_{2} \cdot \hat{\vec{p}}$ are purely imaginary. Therefore, in momentum space, the four-component spinors Ψ_{LL} , Ψ_{LS} , Ψ_{SL} , Ψ_{SS} of Eq.[\(9\)](#page-4-3) are expanded in a complete set of harmonic oscillators with corresponding coefficients according to

$$
\Psi_{LL}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{p}}) = \sum_{n\ell S} a_{n\ell S} |n\ell SJM\rangle_{\mathbf{p}-space}
$$

$$
\Psi_{LS}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{p}}) = \sum_{n\ell S} b_{n\ell S} |n\ell SJM\rangle_{\mathbf{p}-space}
$$

$$
\Psi_{SL}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{p}}) = \sum_{n\ell S} c_{n\ell S} |n\ell SJM\rangle_{\mathbf{p}-space}
$$

$$
\Psi_{SS}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{p}}) = \sum_{n\ell S} d_{n\ell S} |n\ell SJM\rangle_{\mathbf{p}-space}
$$
with
$$
|n\ell SJM\rangle_{\mathbf{p}-space} = R_{n\ell}(a_0; p) \left\{ \left\{ \chi^{S_1}(\vec{\xi}_1) \times \chi^{S_2}(\vec{\xi}_2) \right\}_S \times \mathbf{Y}_{\ell}(\hat{\vec{\mathbf{p}}}) \right\}_{JM}.
$$

A minimization of the energy is performed with the oscillator parameter a_0 . After the inverse Fourier transformation, we again come to the solution of the Dirac equation in the coordinate space:

$$
\Psi_{LL}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{r}}) = \sum_{n\ell S} i^{2n+\ell} a_{n\ell S} R_{n\ell}(a_0; r) \mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \hat{\vec{\mathbf{r}}})
$$

$$
\Psi_{LS}(\vec{\xi}_1, \vec{\xi}_2; \vec{\mathbf{r}}) = \sum_{n\ell S} i^{2n+\ell} b_{n\ell S} R_{n\ell}(a_0; r) \mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \hat{\vec{\mathbf{r}}})
$$

$$
\Psi_{SL}(\vec{\xi}_1, \vec{\xi}_2; \vec{r}) = \sum_{n\ell S} i^{2n+\ell} c_{n\ell S} R_{n\ell}(a_0; r) \mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \hat{\vec{r}})
$$

$$
\Psi_{SS}(\vec{\xi}_1, \vec{\xi}_2; \vec{r}) = \sum_{n\ell S} i^{2n+\ell} d_{n\ell S} R_{n\ell}(a_0; r) \mathcal{Y}_{\ell S}^{JM}(\vec{\xi}_1, \vec{\xi}_2; \hat{\vec{r}})
$$

III. NUMERICAL RESULTS

Numerical calculations have been performed with the value of the fine-structure constant $\alpha = 1/137$ for comparison with the high-precision results of the finite-element method [\[14](#page-11-5)]. From the form of the Dirac Hamiltonian in Eq.[\(5\)](#page-3-0) one can find that the LS- and SLcomponents of the solution of the two-body Dirac equation are identical both for the para-Ps and ortho-Ps. For the ground and excited states of the para-positronium with $J = 0$ the main large-large (LL) and small-small (SS) components consist of only $(\ell, S) = (0, 0)$ spin-orbital channel, while the large-small (LS) and small-large (SS) components consist of only $(\ell, S) = (1, 1)$ channel. In the case of the ortho-Ps, the large-large (LL) and smallsmall (SS) components contain contributions from the $(\ell, S) = (0, 1)$ and $(2,1)$ spin-orbital channels, while the large-small (LS) and small-large (SL) components contain contributions from $(\ell, S) = (1, 0)$ and $(1,1)$ channels.

In Fig[.1](#page-8-0) and Fig[.2](#page-9-0) the radial wave-functions of the large-large (LL), large-small (LS) and small-small (SS) components of the para-Ps and ortho-Ps ground states are presented, respectively. In the upper index of the radial function notations the corresponding spinorbital channels are indicated as $\Psi_{LL}^{(\ell,S)}$, $\Psi_{LS}^{(\ell,S)}$, and $\Psi_{SS}^{(\ell,S)}$. As can be noted, the behaviors of the radial wave functions of all the components of the para-Ps are close to the main parts of the corresponding components of the ortho-Ps. In particular, the corresponding LL-components are almost identical. The values of the SS-component of the para-Ps are very close to the absolute values of the main SS-component $\Psi_{SS}^{(0,1)}(r)$ of the ortho-Ps. The LS-components of the ortho-Ps corresponding to different spin-orbital channels $\Psi_{LS}^{(1,0)}(r)$ and $\Psi_{LS}^{(1,1)}(r)$ are of the same form, but have opposite signs and differ by about 40% in the absolute values.

Another important result is that all the large-large, large-small , small-large and smallsmall components of the ground state wave functions for the para-Ps and ortho-Ps do not have any singularity around zero. This results is in contrast with the results of the method, employed in Refs.[\[14](#page-11-5)], where the ground state solution of the two-body Dirac equation

FIG. 1. Radial wave-functions of the large-large (LL), large-small (LS) and small-small (SS) components of the para-Ps ground state.

contains a singularity at the origin due to the mixing with the anomalous bound state. This problem was analyzed recently in details in Refs. [\[18,](#page-11-9) [19](#page-11-10)].

In Table [I](#page-9-1) a convergence of the numerical results of calculations for the energy $E/2m_e$, binding energy E_b , the sum of the weights of the LS- and SL- components $W = ||\Psi_{(L,S)}||^2 +$ $||\Psi_{(S,L)}||^2$ in the para- and ortho-positronium ground-states are presented in respect to the maximal oscillator quantum number of basis states n_{max} . The optimal value of the oscillator length $a_0 = 3.3 \times 10^5 \text{ GeV}^{-1}$. As can be seen from the table, all the parameters are well converged with increasing n_{max} . As noted above, the further increase of n_{max} results in the collapse of the variational basis. From the table one can see that the final result for the weights of the sum of the large-small and small-large components are identical for the para-Ps and ortho-Ps and estimated to be 6.656E-6. The weight of the small-small component is of order 10[−]¹² in the both para-Ps and ortho-Ps ground states.

In Table [II](#page-10-0) the energies of the lowest states of the para- and ortho-Ps are compared with the results of the finite-element method of high accuracy [\[14\]](#page-11-5). Of course, our variational method on the harmonic-oscillator basis is not very accurate as the finite-element method

FIG. 2. Radial wave-functions of the large-large (LL), large-small (LS) and small-small (SS) components of the ortho-Ps ground state.

TABLE I. Convergence of the numerical results for $E/2m_e$, E_b , the sum of weights of the LS- and SL- components in the para- and ortho-positronium ground-states in respect to n_{max} .

State	n_{max}	$E/2m_e$	E_b (eV)	W	
1S_0	10	0.9999935056580695	6.637203529 7.467E-6		
$(p-Ps)$	20	0.9999933597022965	6.786370016 6.760E-6		
	30	0.9999933453955095 6.800991522 6.667E-6			
	35	0.9999933438924752 6.802527620 6.659E-6			
	37	0.9999933433568433 6.803075034 6.656E-6			
3S_1	10	0.9999935056580694	6.63720352917.467E-6		
$(o-Ps)$	20	0.9999933597022961	6.786370017 6.756E-6		
	30	0.9999933453955091	6.800991522 6.666E-6		
	35	0.9999933438924745	6.802527620 6.660E-6		
	37	0.9999933433568923	6.803074984 6.656E-6		

TABLE II. Energies of the lowest states of the para- and ortho-Ps in comparison with the results of the finite-element method [\[14\]](#page-11-5).

(ℓ, S, J)	$E/2m_e$					$E/2m_e$ [14]					
$(0,0,0)$ 0.999 993 343 356 843 0.999 993 340 148 538 880											
$(0,1,1)$ 0.999 993 343 356 892 0.999 993 340 148 552 498											
$(1,1,0)$ 0.999 998 342 584 549 0.999 998 335 009 885 854											
(1,0,1)						$[0.999998342513543]0.999998335017278391$					

due-to the collapse of the variational basis at some value of n_{max} . The binding energies of the p-Ps and o-Ps ground states in our calculations are estimated to be 6.803075 eV, which should be compared with the values of 6.806403 eV for the para-Ps and 6.806354 eV for the ortho-Ps of Ref.[\[14\]](#page-11-5). This means that the precision of the energy calculations is of order 0.003 eV, which is good enough for the application of corresponding wave functions to the study of different processes with para- and ortho-positronium.

IV. SUMMARY

A new theoretical method was developed to solve the two-body Dirac equation for paraand ortho-positronium bound states. Only Coulomb potential was included into the Dirac Hamiltonian. It is shown that the two-body Dirac Hamiltonian can be written in the Hermitian form and solved in the momentum state representation. Numerical results for the energy spectrum of the para- and ortho-positronium lowest bound states performed within the variational method using the harmonic oscillator basis functions are in good agreement with a high-precision finite-element method. It was shown that the para- and ortho-Ps ground state wave functions after the Fourier transformation to the coordinate space representation do not contain any singularity at the origin in contrast to the traditional bound-state solution in the literature. The weights of the sum of the large-small and small-large components were estimated to be of order 10[−]⁶ , while the weight of the small-small components are of order 10^{-12} for the both para- and ortho-positronium ground states.

ACKNOWLEDGEMENTS

The authors thank D. Baye for valuable discussions and comparison with the results of the Lagrange-mesh method, E. Czerwinski, T.C. Scott and A.M. Rakhimov for useful discussions of the presented results.

- [1] S.D. Bass, S. Mariazzi, P. Moskal, E. Stepien, Rev. Mod. Phys. 95, 021002 (2023).
- [2] G.S. Adkins, D.B. Cassidy, J. Perez-Rios, Phys. Rep. 975, 1-61 (2022).
- [3] P.A. Vetter, S. J. Freedman, Phys. Rev. A 66, 052505 (2002).
- [4] A. Rubbia, Int. J. Mod. Phys. 19, 3961 (2004)
- [5] T. Yamazaki, T. Namba, S. Asai, T. Kobayashi, Phys. Rev. Lett. 104, 083401 (2010).
- [6] C Bartram and R Henning, J. Phys.: Conf. Ser. **1342**, 012106 (2020).
- [7] T.-E. Haugen, E. A. George, O. Naviliat-Cuncic, and P. A. Voytas, EPJ Web of Conferences 282, 01003(2023).
- [8] P. Moskal, E. Czerwinski, J. Raj et al. Nat. Commun. 15, 78 (2024).
- [9] W.E. Caswell, G.P. Lepage, Phys. Lett. B 167, 437 (1986).
- [10] A. Czarnecki, K. Melnikov, A. Yelkhovsky, Phys. Rev. Lett. 82, 311 (1999).
- [11] K. Pachucki, S.G. Karshenboim, Phys. Rev. Lett. 80, 2101 (1998).
- [12] A. Czarnecki, K. Melnikov, A. Yelkhovsky, Phys. Rev. A 59, 4316 (1999).
- [13] P. Labelle, Phys. Rev. D **58**, 093013 (1998).
- [14] T.C. Scott, J. Shertzer, R. A. Moore, Phys. Rev. A 45, 4393 (1992)
- [15] D. Ferenc and E. Matyus, Phys. Rev. A bf 107, 052803 (2023).
- [16] H.W. Crater, C.Y. Wong, P. Van Alstine, Phys. Rev. D 74, 054028 (2006).
- [17] H.W. Crater, Jin-Hee Yoon, Cheuk-Yin Wong, Phys. Rev. D 79, 034011 (2009).
- [18] C.W. Patterson, Phys. Rev. A **100**, 062128 (2019).
- [19] C.W. Patterson, Phys. Rev. A **107**, 042816 (2023).
- [20] Th. Gutsche and D. Robson, Phys. Lett. B 229, 333 (1989).
- [21] Th. Gutsche. A chiral potential model for the light-quark baryons. Ph.D thesis, Florida State University, 249 p. (1987).
- [22] E.M. Tursunov, J. Phys. G 36, 095006 (2009).

[23] E.M. Tursunov, and S. Krewald, Phys. Rev. D 90, 074015 (2014).