

Multilinear Fractional Integral Operators with Generalized Kernels

Yan Lin, Yuhang Zhao and Shuhui Yang*

Abstract In this article, we introduce a class of multilinear fractional integral operators with generalized kernels that are weaker than the Dini kernel condition. We establish the boundedness of multilinear fractional integral operators with generalized kernels on weighted Lebesgue spaces and variable exponent Lebesgue spaces, as well as the boundedness of multilinear commutators generated by multilinear fractional integral operators with generalized kernels and BMO functions. Even when the generalized kernels condition goes back to the Dini kernel condition, the conclusions on the commutators remain new.

1 Introduction

Serving as a generalization of the Hilbert transform to higher-dimensional Euclidean spaces, the Riesz transform, as a classical singular integral operator, is defined by setting, for any $f \in \mathcal{S}(\mathbb{R}^n)$ (the space of all Schwartz functions on \mathbb{R}^n) and $x \in \mathbb{R}^n$,

$$R_j(f)(x) := c_n \text{ p.v. } \int_{\mathbb{R}^n} \frac{x_j - y_j}{|x - y|^{n+1}} f(y) dy, \quad 1 \leq j \leq n,$$

where $c_n := \Gamma(\frac{n+1}{2})/\pi^{\frac{n+1}{2}}$. The Riesz transforms have wide applications in fields such as image processing, signal processing, and partial differential equations, and are commonly used for tasks such as edge detection and texture analysis [16, 24, 41, 42].

The Riesz potential, which plays a crucial role in harmonic analysis, is also known as the fractional integral operator. Let $0 < \alpha < n$ and the Riesz potential is defined by setting, for any $f \in \mathcal{S}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$I_\alpha(f)(x) := \frac{1}{\gamma(\alpha)} \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-\alpha}} dy,$$

where $\gamma(\alpha) := \pi^{\frac{n}{2}} 2^\alpha \Gamma(\frac{\alpha}{2})/\Gamma(\frac{n-\alpha}{2})$. Riesz potentials have important applications in fields such as signal processing, partial differential equations, and probability theory, and have recently been used in image encryption [17].

Due to the different definitions of these two types of operators, they play very different and important roles in image processing. By comparing the above two operators, apart from the coefficients, we can easily observe two differences.

2020 Mathematics Subject Classification. 42B25, 26A33, 42B35

Key words and phrases. multilinear fractional integral operators with generalized kernel, multilinear commutators, variable exponent Lebesgue spaces, weighted Lebesgue spaces

This work was partially supported by the National Natural Science Foundation of China No. 12071052 and Basic Research Foundation of China University of Mining and Technology (Beijing) – Cultivation of Top Innovative Talents for Doctoral No. BBJ2023058.

*Corresponding author, E-mail: yangshuhui@student.cumtb.edu.cn

- (i) The Riesz transform is a principal value integral, while the Riesz potential is not.
- (ii) The Riesz potential involves an additional parameter α compared to the Riesz transform.

Yabuta [40] introduced non-convolution ω -type Calderón-Zygmund operators which are more general than the Riesz transform. In 2014, Lu and Zhang [29] introduced multilinear Calderón-Zygmund operators with Dini's type kernels and established the boundedness of these operators and multilinear commutators on weighted Lebesgue spaces. In 2016, Zhang and Sun [43] established the boundedness of multilinear iterated commutators generated by multilinear Calderón-Zygmund operators with Dini's type kernels and BMO functions on weighted Lebesgue spaces.

Recently, Wu and Zhang [37] proposed multilinear fractional integral operators with Dini's type kernels. They established the boundedness of these operators on weighted Lebesgue spaces and variable exponent Lebesgue spaces, whose weight functions belong to $A_{\vec{P},q}$. Suppose that $\omega(t) : [0, \infty) \rightarrow [0, \infty)$ and $\omega(t)$ is a non-negative, non-decreasing function with $0 < \omega(1) < \infty$. For any $a > 0$,

- (i) ω is said to satisfy the Dini(a) condition, denoted as $\omega \in \text{Dini}(a)$, if

$$|\omega|_{\text{Dini}(a)} := \int_0^1 \omega^a(t) \frac{dt}{t} < \infty.$$

- (ii) ω is said to satisfy the \log^m -Dini(a) condition if the following inequality holds

$$\int_0^1 \omega^a(t) (1 + \log t^{-1})^m \frac{dt}{t} < \infty,$$

where $m \in \mathbb{Z}^+ := \{1, 2, 3, 4, \dots\}$.

It is straightforward to verify that the \log^m -Dini(a) condition is more stronger than the Dini(a) condition, and if $0 < a_1 < a_2 < \infty$, then $\text{Dini}(a_1) \subset \text{Dini}(a_2)$. In particular, if $\omega \in \text{Dini}(1)$, then

$$\sum_{j=0}^{\infty} \omega(2^{-j}) \sim \int_0^1 \omega(t) \frac{dt}{t} < \infty.$$

And if $\omega \in \log\text{-Dini}(1)$, meaning

$$\int_0^1 \omega(t) (1 + \log t^{-1}) \frac{dt}{t} < \infty,$$

then $\omega \in \text{Dini}(1)$ and

$$\sum_{k=1}^{\infty} k\omega(2^{-k}) \sim \int_0^1 \omega(t) (1 + \log t^{-1}) \frac{dt}{t} < \infty.$$

Definition 1.1. (see [37]) Let $0 < \alpha < mn$ and $K_\alpha(x, y_1, \dots, y_m)$ be a locally integrable function defined away from the diagonal $x = y_1 = \dots = y_m$ in $(\mathbb{R}^n)^{m+1}$, which is called the m -linear fractional integral kernel of type $\omega(t)$, if for some $A > 0$, it satisfies the following size condition

$$|K_\alpha(x, \vec{y})| \leq \frac{A}{\left(\sum_{j=1}^m |x - y_j|\right)^{mn-\alpha}}, \quad (1.1)$$

and the smoothness condition

$$|K_\alpha(x, \vec{y}) - K_\alpha(x', \vec{y})| \leq \frac{A}{\left(\sum_{j=1}^m |x - y_j|\right)^{mn-\alpha}} \omega\left(\frac{|x - x'|}{\sum_{j=1}^m |x - y_j|}\right), \quad (1.2)$$

where $|x - x'| \leq \frac{1}{2} \max_{1 \leq j \leq m} |x - y_j|$, and

$$\begin{aligned} & |K_\alpha(x, y_1, \dots, y_j, \dots, y_m) - K_\alpha(x, y_1, \dots, y'_j, \dots, y_m)| \\ & \leq \frac{A}{\left(\sum_{j=1}^m |x - y_j|\right)^{mn-\alpha}} \omega\left(\frac{|y_j - y'_j|}{\sum_{j=1}^m |x - y_j|}\right), \end{aligned} \quad (1.3)$$

where $|y_j - y'_j| \leq \frac{1}{2} \max_{1 \leq j \leq m} |x - y_j|$.

We say $T_\alpha : \mathcal{S}(\mathbb{R}^n) \times \dots \times \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ an m -linear fractional integral operator with kernel of type $\omega(t)$, $K_\alpha(x, \vec{y})$, if

$$T_\alpha(\vec{f})(x) := \int_{(\mathbb{R}^n)^m} K_\alpha(x, \vec{y}) \prod_{j=1}^m f_j(y_j) d\vec{y}, \quad (1.4)$$

where $x \notin \cap_{j=1}^m \text{supp} f_j$ and each $f_j \in \mathcal{S}(\mathbb{R}^n)$, $j = 1, \dots, m$.

Lin and Xiao [27] introduced a kind of multilinear singular integral operators with generalized kernels and they established the weighted norm inequalities on the product of weighted Lebesgue spaces. Inspired by [27] and [37], we focus on a class of generalized fractional integral kernels by attenuating the condition (1.2). For $k \in \mathbb{Z}^+$,

$$\begin{aligned} & \left(\int_{Q(x, 2^{k+2}\sqrt{mn}|x-x'|)^m \setminus Q(x, 2^{k+1}\sqrt{mn}|x-x'|)^m} |K_\alpha(x, \vec{y}) - K_\alpha(x', \vec{y})|^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\ & \leq C C_k 2^{k(\alpha - \frac{mn}{p_0})} |x - x'|^{\alpha - \frac{mn}{p_0}}, \end{aligned} \quad (1.5)$$

where $Q(x, 2^{k+2}\sqrt{mn}|x - x'|)$ denotes the cube centered at x with the sidelength $2^{k+2}\sqrt{mn}|x - x'|$, (p_0, p'_0) is a pair of positive numbers satisfying $\frac{1}{p_0} + \frac{1}{p'_0} = 1$, $1 < p_0 < \infty$ and C_k is a positive constant depending on k .

Definition 1.2. Let $0 < \alpha p'_0 < mn$ and T_α be an m -linear fractional integral operator defined by (1.4). Then T_α is called an m -linear fractional integral operator with generalized kernel if the following conditions are satisfied.

- (1) The kernel function satisfies the size condition (1.1) and the smoothness condition (1.5).
- (2) For fixed $1 \leq s_1, \dots, s_m \leq p'_0$ with $\frac{1}{s} = \frac{1}{s_1} + \dots + \frac{1}{s_m}$, T_α maps $L^{s_1} \times \dots \times L^{s_m} \rightarrow L^{\frac{sn}{n-s\alpha}, \infty}$.

Remark 1.1. As a matter of fact, when $C_k = \omega(2^{-k})$, it is straightforward to deduce that the condition (1.2) implies the condition (1.5) for any $1 < p_0 < \infty$. Denote $\Omega_k := Q(x, 2^k\sqrt{mn}|x - x'|)$.

$$\begin{aligned} & \left(\int_{Q(x, 2^{k+2}\sqrt{mn}|x-x'|)^m \setminus Q(x, 2^{k+1}\sqrt{mn}|x-x'|)^m} |K_\alpha(x, \vec{y}) - K_\alpha(x', \vec{y})|^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\ & \leq \left(\int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} \left[\frac{A}{\left(\sum_{j=1}^\infty |x - y_j|\right)^{mn-\alpha}} \omega\left(\frac{|x - x'|}{\sum_{j=1}^\infty |x - y_j|}\right) \right]^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\ & \leq \left(\int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} \left[\frac{A}{\left(\max_{1 \leq j \leq m} |x - y_j|\right)^{mn-\alpha}} \omega\left(\frac{|x - x'|}{\max_{1 \leq j \leq m} |x - y_j|}\right) \right]^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \end{aligned}$$

$$\begin{aligned}
&\leq \left(\int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} \left[\frac{A}{(2^k |x - x'|)^{mn-\alpha}} \omega(2^{-k}) \right]^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\
&\leq C \omega(2^{-k}) 2^{-k(mn-\alpha)} |x - x'|^{-(mn-\alpha)} \left(\int_{(\Omega_{k+2})^m} d\vec{y} \right)^{\frac{1}{p_0}} \\
&= C \omega(2^{-k}) 2^{-k(mn-\alpha)} |x - x'|^{-(mn-\alpha)} (2^{k+2} \sqrt{mn} |x - x'|)^{\frac{mn}{p_0}} \\
&\leq C \omega(2^{-k}) 2^{k(\alpha - \frac{mn}{p_0})} |x - x'|^{\alpha - \frac{mn}{p_0}} \\
&\leq C C_k 2^{k(\alpha - \frac{mn}{p_0})} |x - x'|^{\alpha - \frac{mn}{p_0}}.
\end{aligned}$$

Remark 1.2. when $\alpha = 0$, the m -linear fractional integral operator with generalized kernel corresponds precisely to the multilinear Calderón-Zygmund operator with generalized kernel studied by Yang, et al [39], as well as Gao, et al [18].

The article is structured as follows. In Section 2, we establish the boundedness of multilinear fractional integral operators with generalized kernels and multilinear commutators on weighted Lebesgue spaces. Section 3 focuses on demonstrating the boundedness of multilinear fractional integral operators and multilinear commutators on variable exponent Lebesgue spaces.

Throughout this article, the letter C always denotes a constant independent of the main parameters involved, whose value may vary from line to line. A cube $Q \subset \mathbb{R}^n$ always refers to a cube with sides parallel to the coordinate axes, and we denote its side length by $l(Q)$. For $t > 0$, the notation tQ represents the cube with the same center as Q and side length $l(tQ) = tl(Q)$. Denote by $|S|$ the Lebesgue measure and χ_S represents the characteristic function for a measurable set $S \subset \mathbb{R}^n$. $B(x, r)$ denotes the ball centered at x with radius r . The notation $X \sim Y$ indicates that there exists a constant $C > 0$ such that $C^{-1}Y \leq X \leq CY$. For any index $1 < q(x) < \infty$, we denote its conjugate index by $q'(x) := \frac{q(x)}{q(x)-1}$. Occasionally, we use the notation $\vec{f} = (f_1, \dots, f_m)$, $T(\vec{f}) := T(f_1, \dots, f_m)$, $d\vec{y} := dy_1 \cdots dy_m$ and $(x, \vec{y}) := (x, y_1, \dots, y_m)$ for convenience. For a set E and a positive integer m , we use the notation $(E)^m := \underbrace{E \times \cdots \times E}_m$ sometimes.

2 The Boundedness on Weighted Lebesgue Spaces

The arrangement of this Section is as follows. In Subsection 2.1, we present certain definitions and symbols that will be utilized later on. Subsection 2.2 is dedicated to establishing the pointwise estimate for the sharp maximal function. Subsection 2.3 addresses the boundedness of multilinear fractional integral operators with generalized kernels and multilinear commutators on weighted Lebesgue spaces.

2.1 Definitions and Lemmas

Definition 2.1. (see [31]) Let $f \in L^1_{loc}(\mathbb{R}^n)$ and denote by M the usual Hardy-Littlewood maximal operator. For a cube $Q \subset \mathbb{R}^n$ and $\delta > 0$, the maximal function M_δ is defined by

$$M_\delta(f)(x) := [M(|f|^\delta)(x)]^{\frac{1}{\delta}} = \left(\sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)|^\delta dy \right)^{\frac{1}{\delta}}.$$

Let $M^\#$ be the standard sharp maximal function, that is

$$M^\# f(x) := \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y) - f_Q| dy \sim \sup_{Q \ni x} \inf_c \frac{1}{|Q|} \int_Q |f(y) - c| dy,$$

where, as usual, f_Q denotes the average of f over the cube Q containing the point x . The operator $M_\delta^\#$ is defined as $M_\delta^\#(f)(x) := [M^\#(|f|^\delta)(x)]^{\frac{1}{\delta}}$.

Definition 2.2. For $0 < \alpha < n$ and $f \in L_{loc}^1(\mathbb{R}^n)$, the standard fractional maximal operator \mathcal{M}_α is defined by

$$\mathcal{M}_\alpha(f)(x) := \sup_{Q \ni x} \frac{1}{|Q|^{1-\frac{\alpha}{n}}} \int_Q |f(y)| dy.$$

For any $r > 1$, $\mathcal{M}_{\alpha,r}$ is defined by

$$\mathcal{M}_{\alpha,r}(f)(x) := \sup_{Q \ni x} |Q|^{\frac{\alpha}{n}} \left(\frac{1}{|Q|} \int_Q |f(y)|^r dy \right)^{\frac{1}{r}}.$$

Definition 2.3. (see [37]) Let $0 < \alpha < mn$ and the multilinear fractional maximal operator \mathcal{M}_α is defined by setting. For $\vec{f} = (f_1, \dots, f_m)$ with $f_i \in L_{loc}^1(\mathbb{R}^n)$, $i = 1, \dots, m$, and $x \in \mathbb{R}^n$,

$$\mathcal{M}_\alpha(\vec{f})(x) := \sup_{Q \ni x} |Q|^{\frac{\alpha}{n}} \prod_{j=1}^m \frac{1}{|Q|} \int_Q |f_j(y_j)| dy_j = \sup_{Q \ni x} \prod_{j=1}^m \frac{1}{|Q|^{1-\frac{\alpha}{mn}}} \int_Q |f_j(y_j)| dy_j.$$

For any $r > 1$, $\mathcal{M}_{\alpha,r}$ is defined by

$$\mathcal{M}_{\alpha,r}(\vec{f})(x) := \sup_{Q \ni x} |Q|^{\frac{\alpha}{n}} \prod_{j=1}^m \left(\frac{1}{|Q|} \int_Q |f_j(y_j)|^r dy_j \right)^{\frac{1}{r}}.$$

Definition 2.4. (see [32]) Let $\vec{P} := (p_1, \dots, p_m)$ and $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ with $1 \leq p_j < \infty$ ($j = 1, \dots, m$). Given $\vec{\omega} := (\omega_1, \dots, \omega_m)$ with nonnegative functions $\omega_1, \dots, \omega_m$ on \mathbb{R}^n and $p \leq q < \infty$, we say that $\vec{\omega}$ satisfies the $A_{\vec{P},q}$ condition if

$$\sup_Q \left(\frac{1}{|Q|} \int_Q v_{\vec{\omega}}(x)^q dx \right)^{\frac{1}{q}} \prod_{j=1}^m \left(\frac{1}{|Q|} \int_Q \omega_j(x)^{-p'_j} dx \right)^{\frac{1}{p'_j}} < \infty,$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$, $v_{\vec{\omega}} = \prod_{j=1}^m \omega_j$, and $\left(\frac{1}{|Q|} \int_Q \omega_j(x)^{-p'_j} dx \right)^{\frac{1}{p'_j}}$

in the case $p_j = 1$ is understood as $\left(\inf_Q \omega_j \right)^{-1}$ when $p_j = 1$.

Definition 2.5. (see [36]) Suppose $\vec{b} := (b_1, \dots, b_m)$ with $b_j \in L_{loc}^1(\mathbb{R}^n)$ ($j = 1, \dots, m$). We define the m -linear commutator $T_{\vec{b},\alpha}^\tau$ to be

$$T_{\vec{b},\alpha}^\tau(\vec{f})(x) := \sum_{i=1}^m T_{\vec{b},\alpha}^i(\vec{f})(x).$$

Each term represents the commutator of T_α in the i -th entry with b_i , which is defined by

$$T_{\vec{b},\alpha}^i(\vec{f})(x) := b_i(x) T_\alpha(f_1, \dots, f_i, \dots, f_m)(x) - T_\alpha(f_1, \dots, b_i f_i, \dots, f_m)(x).$$

Definition 2.6. (see [30]) Suppose that $b \in L_{loc}^1(\mathbb{R}^n)$. Let

$$\|b\|_{\text{BMO}} := \sup_Q \frac{1}{|Q|} \int_Q |b(x) - b_Q| dx,$$

where the supreme is taken over all cube $Q \subset \mathbb{R}^n$, and

$$f_Q := \frac{1}{|Q|} \int_Q f(y) dy.$$

Define

$$\text{BMO}(\mathbb{R}^n) := \{f : \|f\|_{\text{BMO}} < \infty\}.$$

Let $\vec{b} = (b_1, \dots, b_m)$, if $b_j \in \text{BMO}$ for $1 \leq j \leq m$, then $\vec{b} \in \text{BMO}^m$.

Lemma 2.1. (see [15]) Let $0 < p, \delta < \infty$ and ω be any A_∞ -weight ($A_\infty := \cup_{p \geq 1} A_p$). Then there exists a constant $C > 0$ (depending on the A_∞ constant of ω), such that the inequalities

$$\int (M_\delta f(x))^p \omega(x) dx \leq C \int (M_\delta^\# f(x))^p \omega(x) dx,$$

and

$$\|M_\delta(f)\|_{L^{p,\infty}(\omega)} \leq C \|M_\delta^\#(f)\|_{L^{p,\infty}(\omega)},$$

hold for any function f for which the left hand side is finite.

Lemma 2.2. (see [29]) Let $0 < p < u < \infty$. There exists a positive constant $C = C_{p,u}$ such that the following inequality holds

$$|Q|^{-\frac{1}{p}} \|f\|_{L^p(Q)} \leq C |Q|^{-\frac{1}{u}} \|f\|_{L^{u,\infty}(Q)}.$$

Lemma 2.3. (see [32]) Let $0 < \alpha < mn$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ with $1 \leq p_j < \infty$ ($j = 1, \dots, m$), and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n} > 0$.

- (i) If $1 < p_j < \infty$ for all $j = 1, \dots, m$, then for $\vec{\omega} \in A_{\vec{P},q}$, there exists a constant $C > 0$ independent of \vec{f} such that

$$\|\mathcal{M}_\alpha(\vec{f})\|_{L^q(v_\omega^q)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(\omega_j^{p_j})}.$$

- (ii) If $1 \leq p_j < \infty$ for all $j = 1, \dots, m$ and at least one $p_j = 1$ for some $j = 1, \dots, m$, then for $\vec{\omega} \in A_{\vec{P},q}$, there exists a constant $C > 0$ independent of \vec{f} such that

$$\|\mathcal{M}_\alpha(\vec{f})\|_{L^{q,\infty}(v_\omega^q)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(\omega_j^{p_j})}.$$

Lemma 2.4. (see [32]) Let $0 < \alpha < mn$, $\vec{\omega} = (\omega_1, \dots, \omega_m)$, $v_{\vec{\omega}} = \prod_{j=1}^m \omega_j$, $\vec{P} = (p_1, \dots, p_m)$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ with $1 \leq p_j < \infty$ ($j = 1, \dots, m$), and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n} > 0$. Suppose $\vec{\omega} \in A_{\vec{P},q}$, then

$$\begin{cases} \omega_j^{-p'_j} \in A_{mp'_j} (j = 1, \dots, m), \\ v_{\vec{\omega}}^q \in A_{mq}. \end{cases}$$

Lemma 2.5. (see [2]) Let $a \geq 1$. If $b \in \text{BMO}$, then for all cubes Q , we have

- (i) $\left(\frac{1}{|Q|} \int_Q |b(y) - b_Q| dy \right)^{\frac{1}{a}} \leq C \|b\|_{\text{BMO}}.$
- (ii) $\left(\frac{1}{|2^k Q|} \int_{2^k Q} |b(y) - b_Q| dy \right)^{\frac{1}{a}} \leq C k \|b\|_{\text{BMO}}, \text{ for } k \in \mathbb{N}^+.$

2.2 Sharp Maximal Pointwise Estimates

In this subsection, we initially obtain the sharp maximal estimates for multilinear fractional integral operators with generalized kernels and multilinear commutators generated by BMO functions and these operators.

Theorem 2.1. Let $m \geq 2$, $0 < \alpha p'_0 < mn$, T_α be an m -linear fractional integral operator with generalized kernel as defined in Definition 1.2 and $\sum_{k=1}^{\infty} C_k < \infty$. Assuming $0 < \delta < \min \left\{ 1, \frac{sn}{n - s\alpha} \right\}$, there exists a constant $C > 0$ such that for all m -tuples of bounded measurable functions $\vec{f} = (f_1, \dots, f_m)$ with compact support, the following holds

$$M_\delta^\#(T_\alpha(\vec{f}))(x) \leq C \mathcal{M}_{\alpha, p'_0}(\vec{f})(x).$$

Proof. Since $||a|^e - |b|^e| \leq |a - b|^e$ for $0 < e < 1$, for any $Q \ni x$, we can estimate

$$\left(\frac{1}{|Q|} \int_Q \left| |T_\alpha(\vec{f})(z)|^\delta - |c|^\delta \right| dz \right)^{\frac{1}{\delta}} \leq C \left(\frac{1}{|Q|} \int_Q |T_\alpha(\vec{f})(z) - c|^\delta dz \right)^{\frac{1}{\delta}}.$$

Let $Q^* := 14n\sqrt{mn}Q$, we decompose $f_j := f_j^0 + f_j^\infty$ with $f_j^0 := f_j \chi_{Q^*}$, then

$$\begin{aligned} \prod_{j=1}^m f_j(y_j) &= \prod_{j=1}^m (f_j^0(y_j) + f_j^\infty(y_j)) \\ &= \sum_{(\rho_1, \dots, \rho_m) \in \{0, \infty\}} f_1^{\rho_1}(y_1) \times \dots \times f_m^{\rho_m}(y_m) \\ &= \prod_{j=1}^m f_j^0(y_j) + \sum_{(\rho_1, \dots, \rho_m) \in \rho} f_1^{\rho_1}(y_1) \times \dots \times f_m^{\rho_m}(y_m), \end{aligned}$$

where $\rho = \{(\rho_1, \dots, \rho_m) : \text{there is at least one } \rho_j = \infty\}$. It is easy to see that

$$T_\alpha(\vec{f})(z) = T_\alpha(f_1^0, \dots, f_m^0)(z) + \sum_{(\rho_1, \dots, \rho_m) \in \rho} T_\alpha(f_1^{\rho_1}, \dots, f_m^{\rho_m})(z).$$

Let $c = \sum_{(\rho_1, \dots, \rho_m) \in \rho} c_{\rho_1, \dots, \rho_m}$, then we can get

$$\begin{aligned} &\left(\frac{1}{|Q|} \int_Q |T_\alpha(\vec{f})(z) - c|^\delta dz \right)^{\frac{1}{\delta}} \\ &\leq C \left(\frac{1}{|Q|} \int_Q |T_\alpha(f_1^0, \dots, f_m^0)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\ &\quad + C \sum_{(\rho_1, \dots, \rho_m) \in \rho} \left(\frac{1}{|Q|} \int_Q |T_\alpha(f_1^{\rho_1}, \dots, f_m^{\rho_m})(z) - c_{\rho_1, \dots, \rho_m}|^\delta dz \right)^{\frac{1}{\delta}} \end{aligned}$$

$$:= I + \sum_{(\rho_1, \dots, \rho_m) \in \rho} II_{\rho_1, \dots, \rho_m}.$$

From Definition 1.2 (2), Hölder's inequality and Lemma 2.2, it follows that

$$\begin{aligned} I &\leq C|Q|^{-\frac{n-s\alpha}{sn}} \|T_\alpha(f_1^0, \dots, f_m^0)\|_{L^{\frac{sn}{n-s\alpha}, \infty}(Q)} \\ &\leq C|Q|^{\frac{\alpha}{n}} |Q|^{-\frac{1}{s}} \left(\int_{Q^*} |f_1(y_1)|^{s_1} dy_1 \right)^{\frac{1}{s_1}} \times \dots \times \left(\int_{Q^*} |f_m(y_m)|^{s_m} dy_m \right)^{\frac{1}{s_m}} \\ &\leq C|Q^*|^{\frac{\alpha}{n}} \left(\frac{1}{|Q^*|} \int_{Q^*} |f_1(y_1)|^{s_1} dy_1 \right)^{\frac{1}{s_1}} \times \dots \times \left(\frac{1}{|Q^*|} \int_{Q^*} |f_m(y_m)|^{s_m} dy_m \right)^{\frac{1}{s_m}} \\ &\leq C|Q^*|^{\frac{\alpha}{n}} \prod_{j=1}^m \left(\frac{1}{|Q^*|} \int_{Q^*} |f_j(y_j)|^{p'_0} dy_j \right)^{\frac{1}{p'_0}} \\ &\leq CM_{\alpha, p'_0}(\vec{f})(x). \end{aligned}$$

To estimate $II_{\rho_1, \dots, \rho_m}$, we choose $c_{\rho_1, \dots, \rho_m} := T_\alpha(f_1^{\rho_1}, \dots, f_m^{\rho_m})(z_0)$ for a fixed $z_0 \in 4Q \setminus 3Q$. Let $\Omega_k := Q(z_0, 2^k \sqrt{mn}|z - z_0|)$, $k \in \mathbb{N}^+$. Note that $\Omega_2 \subset Q^*$, so $(\mathbb{R}^n)^m \setminus (Q^*)^m \subset (\mathbb{R}^n)^m \setminus (\Omega_2)^m$.

When $z \in Q$, we obtain $l(Q) \leq |z - z_0| \leq \frac{5}{2} \sqrt{nl}(Q)$. By Hölder's inequality, we conclude that

$$\begin{aligned} II_{\rho_1, \dots, \rho_m} &\leq \frac{C}{|Q|} \int_Q |T_\alpha(f_1^{\rho_1}, \dots, f_m^{\rho_m})(z) - T_\alpha(f_1^{\rho_1}, \dots, f_m^{\rho_m})(z_0)| dz \\ &\leq \frac{C}{|Q|} \int_Q \int_{(\mathbb{R}^n)^m \setminus (Q^*)^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})| \prod_{j=1}^m |f_j(y_j)| d\vec{y} dz \\ &\leq \frac{C}{|Q|} \int_Q \int_{(\mathbb{R}^n)^m \setminus (\Omega_2)^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})| \prod_{j=1}^m |f_j(y_j)| d\vec{y} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} \int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})| \prod_{j=1}^m |f_j(y_j)| d\vec{y} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} \left(\int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})|^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\ &\quad \times \left(\int_{(\Omega_{k+2})^m} \prod_{j=1}^m |f_j(y_j)|^{p'_0} d\vec{y} \right)^{\frac{1}{p'_0}} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} C_k |z - z_0|^{\alpha - \frac{mn}{p'_0}} 2^{k(\alpha - \frac{mn}{p'_0})} |\Omega_{k+2}|^{\frac{m}{p'_0} - \frac{\alpha}{n}} \\ &\quad \times |\Omega_{k+2}|^{\frac{\alpha}{n}} \prod_{j=1}^m \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_j(y_j)|^{p'_0} dy_j \right)^{\frac{1}{p'_0}} dz \\ &\leq CM_{\alpha, p'_0}(\vec{f})(x). \end{aligned}$$

Combining the above estimates we get the desired result. \square

Theorem 2.2. Let $m \geq 2$, $0 < \alpha p'_0 < mn$, T_α be an m -linear fractional integral operator with generalized kernel as defined in Definition 1.2 and $\sum_{k=1}^{\infty} kC_k < \infty$. Assuming $0 < \delta < \min \left\{ 1, \frac{sn}{n-s\alpha} \right\}$,

$p'_0 < t < \infty$, $\delta < \varepsilon < \infty$ and $\vec{b} \in \text{BMO}^m$, there exists a constant $C > 0$ such that for all bounded measurable functions with compact support m -tuples $\vec{f} = (f_1, \dots, f_m)$, we have

$$M_\delta^\#(T_{\vec{b}, \alpha}(\vec{f}))(x) \leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\mathcal{M}_{\alpha, t}(\vec{f})(x) + M_\varepsilon(T_\alpha(\vec{f}))(x) \right),$$

where $\|\vec{b}\|_{(\text{BMO})^m} := \max_{1 \leq i \leq m} \|b_i\|_{\text{BMO}}$.

Proof. Let $Q^* := 14n\sqrt{mn}Q$, we decompose $f_j := f_j^0 + f_j^\infty$ with $f_j^0 := f_j \chi_{Q^*}$, then

$$\begin{aligned} \prod_{j=1}^m f_j(y_j) &= \prod_{j=1}^m (f_j^0(y_j) + f_j^\infty(y_j)) \\ &= \sum_{(\rho_1, \dots, \rho_m) \in \{0, \infty\}} f_1^{\rho_1}(y_1) \times \dots \times f_m^{\rho_m}(y_m) \\ &= \prod_{j=1}^m f_j^0(y_j) + \sum_{(\rho_1, \dots, \rho_m) \in \rho} f_1^{\rho_1}(y_1) \times \dots \times f_m^{\rho_m}(y_m), \end{aligned}$$

where $\rho = \{(\rho_1, \dots, \rho_m) : \text{there is at least one } \rho_j = \infty\}$. It is easy to see that

$$\begin{aligned} &T_\alpha(f_1, \dots, (b_j - \lambda)f_j, \dots, f_m)(z) \\ &= T_\alpha(f_1^0, \dots, (b_j - \lambda)f_j^0, \dots, f_m^0)(z) \\ &+ \sum_{(\rho_1, \dots, \rho_m) \in \rho} T_\alpha(f_1^{\rho_1}, \dots, (b_j - \lambda)f_j^{\rho_j}, \dots, f_m^{\rho_m})(z). \end{aligned}$$

Fix $x \in \mathbb{R}^n$ and let Q be a cube containing x . Denote $\lambda = b_j|_{Q^*}$. Let $c_j = - \sum_{(\rho_1, \dots, \rho_m) \in \rho} c_{j, \rho_1, \dots, \rho_m}$,

then

$$\begin{aligned} &\left(\frac{1}{|Q|} \int_Q |T_{\vec{b}, \alpha}^j(\vec{f})(z) - c_j|^\delta dz \right)^{\frac{1}{\delta}} \\ &\leq C \left(\frac{1}{|Q|} \int_Q |(b_j(z) - \lambda)T_\alpha(\vec{f})(z)|^\delta dz \right)^{\frac{1}{\delta}} + C \left(\frac{1}{|Q|} \int_Q |T_\alpha(f_1, \dots, (b_j - \lambda)f_j, \dots, f_m)(z) + c_j|^\delta dz \right)^{\frac{1}{\delta}} \\ &\leq C \left(\frac{1}{|Q|} \int_Q |(b_j(z) - \lambda)T_\alpha(\vec{f})(z)|^\delta dz \right)^{\frac{1}{\delta}} + C \left(\frac{1}{|Q|} \int_Q |T_\alpha(f_1^0, \dots, (b_j - \lambda)f_j^0, \dots, f_m^0)(z)|^\delta dz \right)^{\frac{1}{\delta}} \\ &\quad + C \sum_{(\rho_1, \dots, \rho_m) \in \rho} \left(\frac{1}{|Q|} \int_Q |T_\alpha(f_1^{\rho_1}, \dots, (b_j - \lambda)f_j^{\rho_j}, \dots, f_m^{\rho_m})(z) - c_{j, \rho_1, \dots, \rho_m}|^\delta dz \right)^{\frac{1}{\delta}} \\ &:= III + IV + \sum_{(\rho_1, \dots, \rho_m) \in \rho} V_{\rho_1, \dots, \rho_m} \end{aligned}$$

Choosing $1 < h'_1 < \min\{\frac{1}{1-\delta}, \frac{\varepsilon}{\delta}\}$ with $\frac{1}{h'_1} + \frac{1}{h_1} = 1$, by Hölder's inequality, we conclude that

$$\begin{aligned} III &\leq C \left(\frac{1}{|Q|} \int_Q |b_j(z) - \lambda|^{\delta h_1} dz \right)^{\frac{1}{\delta h_1}} \left(\frac{1}{|Q|} \int_Q |T_\alpha(\vec{f})|^{\delta h'_1} dz \right)^{\frac{1}{\delta h'_1}} \\ &\leq C \left(\frac{1}{|Q^*|} \int_{Q^*} |b_j(z) - \lambda|^{\delta h_1} dz \right)^{\frac{1}{\delta h_1}} \left(\frac{1}{|Q|} \int_Q |T_\alpha(\vec{f})|^{\delta h'_1} dz \right)^{\frac{1}{\delta h'_1}} \end{aligned}$$

$$\leq C \|b_j\|_{\text{BMO}} M_\varepsilon(T_\alpha(\vec{f}))(x).$$

Let $h_2 = \frac{t}{s_j}$ with $\frac{1}{h_2} + \frac{1}{h'_2} = 1$. From Definition 1.2 (2), Hölder's inequality, Lemma 2.2 and $s_i \leq p'_0 < t$ ($i = 1, \dots, m$), it follows that

$$\begin{aligned} IV &\leq C |Q|^{-\frac{n-s\alpha}{sn}} \|T_\alpha(f_1^0, \dots, (b_j - \lambda) f_j^0, \dots, f_m^0)\|_{L^{\frac{sn}{n-s\alpha}, \infty}(Q)} \\ &\leq C |Q^*|^{\frac{\alpha}{n}} |Q^*|^{-\frac{1}{s}} \prod_{i=1, i \neq j}^m \|f_i\|_{L^{s_i}(Q^*)} \|(b_j - \lambda) f_j\|_{L^{s_j}(Q^*)} \\ &= C |Q^*|^{\frac{\alpha}{n}} \prod_{i=1, i \neq j}^m \left(\frac{1}{|Q^*|} \int_{Q^*} |f_i(y_i)|^{s_i} dy_i \right)^{\frac{1}{s_i}} \left(\frac{1}{|Q^*|} \int_{Q^*} |b_j(y_j) - \lambda|^{s_j} |f_j(y_j)|^{s_j} dy_j \right)^{\frac{1}{s_j}} \\ &\leq C |Q^*|^{\frac{\alpha}{n}} \prod_{i=1, i \neq j}^m \left(\frac{1}{|Q^*|} \int_{Q^*} |f_i(y_i)|^{s_i} dy_i \right)^{\frac{1}{s_i}} \left(\frac{1}{|Q^*|} \int_{Q^*} |b_j(y_j) - \lambda|^{s_j h'_2} dy_j \right)^{\frac{1}{s_j h'_2}} \\ &\quad \times \left(\frac{1}{|Q^*|} \int_{Q^*} |f_j(y_j)|^{s_j h_2} dy_j \right)^{\frac{1}{s_j h_2}} \\ &\leq C \|b_j\|_{\text{BMO}} |Q^*|^{\frac{\alpha}{n}} \prod_{i=1}^m \left(\frac{1}{|Q^*|} \int_{Q^*} |f_i(y_i)|^t dy_i \right)^{\frac{1}{t}} \\ &\leq C \|b_j\|_{\text{BMO}} \mathcal{M}_{\alpha, t}(\vec{f})(x). \end{aligned}$$

To estimate $V_{\rho_1, \dots, \rho_m}$, we choose $c_{j, \rho_1, \dots, \rho_m} = T_\alpha(f_1^{\rho_1}, \dots, (b_j - \lambda) f_j^{\rho_j}, \dots, f_m^{\rho_m})(z_0)$ for a fixed $z_0 \in 4Q \setminus 3Q$. Since $\Omega_2 \subset Q^*$, we can get $\Omega_{k+2} \subset 2^k Q^*$. Since $|z - z_0| \sim l(Q)$ when $z \in Q$, then $|\Omega_{k+2}| \sim |2^k Q^*|$. Let $h_3 = \frac{t}{p'_0}$ with $\frac{1}{h_3} + \frac{1}{h'_3} = 1$. By Hölder's inequality, Lemma 2.5 and $s_i \leq p'_0 < t$ ($i = 1, \dots, m$), we conclude that

$$\begin{aligned} V_{\rho_1, \dots, \rho_m} &\leq \frac{C}{|Q|} \int_Q |T_\alpha(f_1^{\rho_1}, \dots, (b_j - \lambda) f_j^{\rho_j}, \dots, f_m^{\rho_m})(z) - T_\alpha(f_1^{\rho_1}, \dots, (b_j - \lambda) f_j^{\rho_j}, \dots, f_m^{\rho_m})(z_0)| dz \\ &\leq \frac{C}{|Q|} \int_Q \int_{(\mathbb{R}^n)^m \setminus (Q^*)^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})| \prod_{i=1}^m |f_i(y_i)| |b_j(y_j) - \lambda| d\vec{y} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} \int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})| \prod_{i=1}^m |f_i(y_i)| |b_j(y_j) - \lambda| d\vec{y} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} \left(\int_{(\Omega_{k+2})^m \setminus (\Omega_{k+1})^m} |K_\alpha(z, \vec{y}) - K_\alpha(z_0, \vec{y})|^{p_0} d\vec{y} \right)^{\frac{1}{p_0}} \\ &\quad \times \left(\int_{(\Omega_{k+2})^m} \left(\prod_{i=1}^m |f_i(y_i)| |b_j(y_j) - \lambda| \right)^{p'_0} d\vec{y} \right)^{\frac{1}{p'_0}} dz \\ &\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} C_k |z - z_0|^{\alpha - \frac{m\alpha}{p'_0}} 2^{k(\alpha - \frac{m\alpha}{p'_0})} |\Omega_{k+2}|^{\frac{m}{p'_0} - \frac{\alpha}{n}} |\Omega_{k+2}|^{\frac{\alpha}{n}} \\ &\quad \times \prod_{i=1, i \neq j}^m \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_i(y_i)|^{p'_0} dy_i \right)^{\frac{1}{p'_0}} \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |b_j(y_j) - \lambda|^{p'_0} |f_j(y_j)|^{p'_0} dy_j \right)^{\frac{1}{p'_0}} dz \end{aligned}$$

$$\begin{aligned}
&\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} C_k |\Omega_{k+2}|^{\frac{\alpha}{n}} \prod_{i=1, i \neq j}^m \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_i(y_i)|^t dy_i \right)^{\frac{1}{t}} \\
&\quad \times \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_j(y_j)|^{p'_0 h_3} dy_j \right)^{\frac{1}{p'_0 h_3}} \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |b_j(y_j) - \lambda|^{p'_0 h'_3} dy_j \right)^{\frac{1}{p'_0 h'_3}} dz \\
&\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} C_k |\Omega_{k+2}|^{\frac{\alpha}{n}} \prod_{i=1, i \neq j}^m \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_i(y_i)|^t dy_i \right)^{\frac{1}{t}} \\
&\quad \times \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_j(y_j)|^t dy_j \right)^{\frac{1}{t}} \left(\frac{1}{|2^k Q^*|} \int_{2^k Q^*} |b_j(y_j) - \lambda|^{p'_0 h'_3} dy_j \right)^{\frac{1}{p'_0 h'_3}} dz \\
&\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} C_k |\Omega_{k+2}|^{\frac{\alpha}{n}} \prod_{i=1}^m \left(\frac{1}{|\Omega_{k+2}|} \int_{\Omega_{k+2}} |f_i(y_i)|^t dy_i \right)^{\frac{1}{t}} k \|b_j\|_{\text{BMO}} dz \\
&\leq C \|b_j\|_{\text{BMO}} \mathcal{M}_{\alpha, t}(\vec{f})(x).
\end{aligned}$$

Then, we have

$$\begin{aligned}
&\inf_c \left(\frac{1}{|Q|} \int_Q \left| |T_{\vec{b}, \alpha}(\vec{f})(z)|^\delta - c| dz \right)^{\frac{1}{\delta}} \leq \left(\frac{1}{|Q|} \int_Q \left| \left| \sum_{i=1}^m T_{\vec{b}, \alpha}^i(\vec{f})(z) \right|^\delta - \left| \sum_{i=1}^m c_i \right|^\delta \right| dz \right)^{\frac{1}{\delta}} \\
&\leq C \sum_{i=1}^m \left(\frac{1}{|Q|} \int_Q \left| T_{\vec{b}, \alpha}^i(\vec{f})(z) - c_i \right|^\delta dz \right)^{\frac{1}{\delta}} \leq C \|\vec{b}\|_{(\text{BMO})^m} \left(M_\varepsilon(T_\alpha(\vec{f}))(x) + \mathcal{M}_{\alpha, t}(\vec{f})(x) \right).
\end{aligned}$$

Then, we have

$$\begin{aligned}
M_\delta^\#(T_{\vec{b}, \alpha}(\vec{f}))(x) &\sim \sup_{Q \ni x} \inf_c \left(\frac{1}{|Q|} \int_Q \left| |T_{\vec{b}, \alpha}(\vec{f})(z)|^\delta - c| dz \right)^{\frac{1}{\delta}} \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(M_\varepsilon(T_\alpha(\vec{f}))(x) + \mathcal{M}_{\alpha, t}(\vec{f})(x) \right).
\end{aligned}$$

Thus, Theorem 2.2 is proved. \square

2.3 The Boundedness on Weighted Lebesgue Spaces

In this subsection, we use Theorem 2.1 and Theorem 2.2 to derive boundedness results for multilinear fractional integral operators with generalized kernels and multilinear commutators on weighted Lebesgue spaces. The main theorem of this subsection is presented below.

Theorem 2.3. Let $m \geq 2$, $0 < \alpha p'_0 < mn$, T_α be an m -linear fractional integral operator with generalized kernel as in Definition 1.2 and $\sum_{k=1}^{\infty} C_k < \infty$. Suppose that $\vec{P} = (p_1, \dots, p_m)$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ with $p'_0 \leq p_j < \infty$ ($j = 1, \dots, m$), $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n} > 0$ and $\vec{\omega} \in A_{\vec{P}, \frac{q}{p'_0}}$.

(i) If $p'_0 < p_j < \infty$ for all $j = 1, \dots, m$, then

$$\left\| T_\alpha(\vec{f}) \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}.$$

(ii) If $p'_0 \leq p_j < \infty$ for all $j = 1, \dots, m$ and at least one of $p_j = p'_0$, then

$$\left\| T_\alpha(\vec{f}) \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}.$$

Proof. Firstly, we choose δ such that $0 < \delta < \min \left\{ 1, \frac{sn}{n - s\alpha} \right\}$ and let $\beta = \alpha p'_0$.

(i) By Lemma 2.4, for $\vec{\omega} \in A_{\frac{\vec{p}}{p'_0}, \frac{q}{p'_0}}$ there has $v_{\vec{\omega}}^{\frac{q}{p'_0}} \in A_\infty$. Then, by Lemma 2.1, Theorem 2.1 and Lemma 2.3 (i), we conclude that

$$\begin{aligned} \left\| T_\alpha(\vec{f}) \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} &\leq \left\| M_\delta(T_\alpha(\vec{f})) \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \leq C \left\| M_\delta^\#(T_\alpha(\vec{f})) \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \\ &\leq C \left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} = C \left\| [\mathcal{M}_\beta(\vec{f}^{p'_0})]^{\frac{1}{p'_0}} \right\|_{L^q \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \\ &= C \left\| \mathcal{M}_\beta(\vec{f}^{p'_0}) \right\|_{L^{\frac{q}{p'_0}} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)}^{\frac{1}{p'_0}} \leq C \prod_{j=1}^m \left\| |f_j|^{p'_0} \right\|_{L^{\frac{p_j}{p'_0}} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}^{\frac{1}{p'_0}} \\ &= C \prod_{j=1}^m \|f_j\|_{L^{p_j} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}. \end{aligned}$$

Thus, Theorem 2.3 (i) is proved.

(ii) By Lemma 2.1, Theorem 2.1 and Lemma 2.3 (ii), we obtain

$$\begin{aligned} \left\| T_\alpha(\vec{f}) \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} &\leq \left\| M_\delta(T_\alpha(\vec{f})) \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \leq C \left\| M_\delta^\#(T_\alpha(\vec{f})) \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \\ &\leq C \left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} = C \left\| [\mathcal{M}_\beta(\vec{f}^{p'_0})]^{\frac{1}{p'_0}} \right\|_{L^{q, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)} \\ &= C \sup_{\lambda > 0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{p'_0}} \left(\left\{ x \in \mathbb{R}^n : [\mathcal{M}_\beta(\vec{f}^{p'_0})]^{\frac{1}{p'_0}} > \lambda \right\} \right) \right|^{\frac{1}{q}} \\ &= C \sup_{\lambda > 0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{p'_0}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_\beta(\vec{f}^{p'_0}) > \lambda^{p'_0} \right\} \right) \right|^{\frac{1}{q}} \\ &= C \sup_{\lambda > 0} \lambda^{\frac{1}{p'_0}} \left| v_{\vec{\omega}}^{\frac{q}{p'_0}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_\beta(\vec{f}^{p'_0}) > \lambda \right\} \right) \right|^{\frac{1}{q}} \\ &= C \left[\sup_{\lambda > 0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{p'_0}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_\beta(\vec{f}^{p'_0}) > \lambda \right\} \right) \right|^{\frac{p'_0}{q}} \right]^{\frac{1}{p'_0}} \\ &= C \left\| \mathcal{M}_\beta(\vec{f}^{p'_0}) \right\|_{L^{\frac{q}{p'_0}, \infty} \left(v_{\vec{\omega}}^{\frac{q}{p'_0}} \right)}^{\frac{1}{p'_0}} \leq C \prod_{j=1}^m \left\| |f_j|^{p'_0} \right\|_{L^{\frac{p_j}{p'_0}} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}^{\frac{1}{p'_0}} \\ &= C \prod_{j=1}^m \|f_j\|_{L^{p_j} \left(\omega_j^{\frac{p_j}{p'_0}} \right)}. \end{aligned}$$

Thus, Theorem 2.3 (ii) is proved. \square

Theorem 2.4. Let $m \geq 2$, $p'_0 < t < \infty$, $0 < \alpha t < mn$, T_α be an m -linear fractional integral operator with generalized kernel as defined in Definition 1.2 and $\sum_{k=1}^{\infty} kC_k < \infty$. Suppose that $\vec{b} \in \text{BMO}^m$, $\vec{P} = (p_1, \dots, p_m)$, $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$ with $t \leq p_j < \infty$ ($j = 1, \dots, m$), $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n} > 0$ and $\vec{\omega} \in A_{\vec{P}, \frac{q}{t}}$.

(i) If $t < p_j < \infty$ for all $j = 1, \dots, m$, then

$$\left\| T_{\vec{b}, \alpha}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \leq C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{p_j}\left(\omega_j^{\frac{p_j}{t}}\right)}.$$

(ii) If $t \leq p_j < \infty$ for all $j = 1, \dots, m$ and at least one of $p_j = t$, then

$$\left\| T_{\vec{b}, \alpha}(\vec{f}) \right\|_{L^{q, \infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \leq C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{p_j}\left(\omega_j^{\frac{p_j}{t}}\right)}.$$

Proof. Firstly, we choose δ, ε such that $0 < \delta < \varepsilon < \min\left\{1, \frac{sn}{n - s\alpha}\right\}$ and let $\beta = \alpha t$.

(i) By $\vec{\omega} \in A_{\vec{P}, \frac{q}{t}}$, Lemma 2.1, Theorem 2.2, Theorem 2.1 and Lemma 2.3 (i), we conclude that

$$\begin{aligned} \left\| T_{\vec{b}, \alpha}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} &\leq \left\| M_\delta(T_{\vec{b}, \alpha}(\vec{f})) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \leq \left\| M_\delta^\#(T_{\vec{b}, \alpha}(\vec{f})) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\ &\leq C \|\vec{b}\|_{(\text{BMO})^m} \left\| M_\varepsilon(T_\alpha(\vec{f})) + \mathcal{M}_{\alpha, t}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\ &\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\left\| M_\varepsilon(T_\alpha(\vec{f})) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \left\| \mathcal{M}_{\alpha, t}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\ &\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\left\| M_\varepsilon^\#(T_\alpha(\vec{f})) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \left\| \mathcal{M}_{\alpha, t}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\ &\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \left\| \mathcal{M}_{\alpha, t}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\ &\leq C \|\vec{b}\|_{(\text{BMO})^m} \left\| \mathcal{M}_{\alpha, t}(\vec{f}) \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} = C \|\vec{b}\|_{(\text{BMO})^m} \left\| \left[\mathcal{M}_\beta(\vec{f}^t) \right]^{\frac{1}{t}} \right\|_{L^q\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\ &= C \|\vec{b}\|_{(\text{BMO})^m} \left\| \mathcal{M}_\beta(\vec{f}^t) \right\|_{L^{\frac{q}{t}}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)}^{\frac{1}{t}} \leq C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \| |f_j|^t \|_{L^{\frac{p_j}{t}}\left(\omega_j^{\frac{p_j}{t}}\right)}^{\frac{1}{t}} \\ &= C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{p_j}\left(\omega_j^{\frac{p_j}{t}}\right)}. \end{aligned}$$

Thus, Theorem 2.4 (i) is proved.

(ii) By Lemma 2.1, Theorem 2.2, Theorem 2.1 and Lemma 2.3 (ii), we conclude that

$$\begin{aligned}
\|T_{\vec{b},\alpha}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} &\leq \|M_{\delta}(T_{\vec{b},\alpha}(\vec{f}))\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \leq C \|M_{\delta}^{\#}(T_{\vec{b},\alpha}(\vec{f}))\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \|M_{\varepsilon}(T_{\alpha}(\vec{f})) + M_{\alpha,t}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\|M_{\varepsilon}(T_{\alpha}(\vec{f}))\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \|M_{\alpha,t}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\|M_{\varepsilon}^{\#}(T_{\alpha}(\vec{f}))\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \|M_{\alpha,t}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \left(\|M_{\alpha,p'_0}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} + \|M_{\alpha,t}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \right) \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \|M_{\alpha,t}(\vec{f})\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \left\| \left[\mathcal{M}_{\beta}(\vec{f}^t) \right]^{\frac{1}{t}} \right\|_{L^{q,\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \sup_{\lambda>0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{t}} \left(\left\{ x \in \mathbb{R}^n : \left[\mathcal{M}_{\beta}(\vec{f}^t) \right]^{\frac{1}{t}}(x) > \lambda \right\} \right) \right|^{\frac{1}{q}} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \sup_{\lambda>0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{t}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_{\beta}(\vec{f}^t) > \lambda^t \right\} \right) \right|^{\frac{1}{q}} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \sup_{\lambda>0} \lambda^{\frac{1}{t}} \left| v_{\vec{\omega}}^{\frac{q}{t}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_{\beta}(\vec{f}^t) > \lambda \right\} \right) \right|^{\frac{1}{q}} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \left[\sup_{\lambda>0} \lambda \left| v_{\vec{\omega}}^{\frac{q}{t}} \left(\left\{ x \in \mathbb{R}^n : \mathcal{M}_{\beta}(\vec{f}^t) > \lambda \right\} \right) \right|^{\frac{t}{q}} \right]^{\frac{1}{t}} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \left\| \mathcal{M}_{\beta}(\vec{f}^t) \right\|_{L^{\frac{q}{t},\infty}\left(v_{\vec{\omega}}^{\frac{q}{t}}\right)}^{\frac{1}{t}} \\
&\leq C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{\frac{p_j}{t}}\left(\omega_j^{\frac{p_j}{t}}\right)}^{\frac{1}{t}} \\
&= C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{p_j}\left(\omega_j^{\frac{p_j}{t}}\right)}.
\end{aligned}$$

Thus, Theorem 2.4 (ii) is proved. \square

Remark 2.1. When the generalized kernel condition goes back to the Dini kernel condition, the boundedness of m -linear fractional integral operators with Dini's type kernels on Lebesgue spaces has been demonstrated by Wu and Zhang in [37].

Corollary 2.1. When the generalized kernel condition goes back to the Dini kernel condition, the boundedness of multilinear commutators of m -linear fractional integral operators with Dini's type kernels on Lebesgue spaces remains new.

3 The Boundedness on Variable Exponent Lebesgue Spaces

Firstly, we introduce several definitions and notations that will be utilized subsequently. Then we establish the boundedness of multilinear fractional integral operators with generalized kernels and multilinear commutators on variable exponent Lebesgue spaces, respectively.

Definition 3.1. (see [9]) Let $q(\cdot) : \mathbb{R}^n \rightarrow [1, \infty)$ be a measurable function. The variable exponent Lebesgue space $L^{q(\cdot)}(\mathbb{R}^n)$ is defined by

$$L^{q(\cdot)}(\mathbb{R}^n) := \{f \text{ is measurable function} : F_q(f/\eta) < \infty \text{ for some constant } \eta > 0\},$$

where $F_q(f) := \int_{\mathbb{R}^n} |f(x)|^{q(x)} dx$ is a convex functional modular.

The space $L^{q(\cdot)}(\mathbb{R}^n)$ is a Banach function space with respect to the Luxemburg type norm.

$$\|f\|_{L^{q(\cdot)}(\mathbb{R}^n)} := \inf\{\eta > 0 : \int_{\mathbb{R}^n} \left(\frac{|f(x)|}{\eta} \right)^{q(x)} dx \leq 1\}.$$

Definition 3.2. (see [9]) Given a measurable function $q(\cdot)$ defined on \mathbb{R}^n . For $E \subset \mathbb{R}^n$, we write

$$q_-(E) := \operatorname{ess\,inf}_{x \in E} q(x), \quad q_+(E) := \operatorname{ess\,inf}_{x \in E} q(x),$$

and write $q_-(\mathbb{R}^n) = q_-$ and $q_+(\mathbb{R}^n) = q_+$ simply.

$$(i) \quad q'_- = \operatorname{ess\,inf}_{x \in \mathbb{R}^n} q'(x) = \frac{q_+}{q_+ - 1}, \quad q'_+ = \operatorname{ess\,inf}_{x \in \mathbb{R}^n} q'(x) = \frac{q_-}{q_- + 1}.$$

(ii) Denote by $\mathcal{P}(\mathbb{R}^n)$ the set of all measurable functions $q(\cdot) : \mathbb{R}^n \rightarrow (1, \infty)$ such that

$$1 < q_- \leq q(x) \leq q_+ < \infty, x \in \mathbb{R}^n.$$

(iii) The set $\mathcal{B}(\mathbb{R}^n)$ consists of all measurable functions $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ such that the Hardy-Littlewood maximal operator M is bounded on $L^{q(\cdot)}(\mathbb{R}^n)$.

$$\|Mf\|_{L^{q(\cdot)}} \leq \|f\|_{L^{q(\cdot)}}.$$

Definition 3.3. (see [10]) Let $q(\cdot)$ be a real-valued function on \mathbb{R}^n .

(i) Denote by $\mathcal{L}_{loc}^{log}(\mathbb{R}^n)$ the set of all local log-Hölder continuous functions $q(\cdot)$ which satisfy

$$|q(x) - q(y)| \leq \frac{-C}{\ln(|x - y|)}, |x - y| \leq 1/2, x, y \in \mathbb{R}^n.$$

Here, the positive constant C does not depend on x or y .

(ii) The set $\mathcal{L}_{\infty}^{log}(\mathbb{R}^n)$ consists of all log-Hölder continuous functions $q(\cdot)$ at infinity, satisfying

$$|q(x) - q_{\infty}| \leq \frac{C_{\infty}}{\ln(e + |x|)}, x \in \mathbb{R}^n,$$

$$\text{where } q_{\infty} := \lim_{|x| \rightarrow \infty} q(x).$$

(iii) Denote by $\mathcal{L}^{log}(\mathbb{R}^n) := \mathcal{L}_{loc}^{log}(\mathbb{R}^n) \cap \mathcal{L}_{\infty}^{log}(\mathbb{R}^n)$ the set of all global log-Hölder continuous functions $q(\cdot)$.

Remark 3.1. (see [37]) The condition $\mathcal{L}_\infty^{\log}(\mathbb{R}^n)$ is equivalent to the uniform continuity condition

$$|q(x) - q(y)| \leq \frac{C}{\ln(e + |x|)}, |y| \geq |x|, x, y \in \mathbb{R}^n.$$

In what follows, we denote

$$\mathcal{P}^{\log}(\mathbb{R}^n) := \mathcal{L}^{\log}(\mathbb{R}^n) \cap \mathcal{P}(\mathbb{R}^n).$$

Lemma 3.1. (see [9]) Let $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$.

(1) If $p(\cdot) \in \mathcal{L}^{\log}(\mathbb{R}^n)$, then we have $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$.

(2) The following condition are equivalent:

- (i) $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$.
- (ii) $(p(\cdot)/p_0)' \in \mathcal{B}(\mathbb{R}^n)$ for some $1 < p_0 < p_-$.

Lemma 3.2. (see [29]) Let $q(\cdot), q_1(\cdot), \dots, q_m(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfy the condition

$$\frac{1}{q(x)} = \frac{1}{q_1(x)} + \dots + \frac{1}{q_m(x)}, \text{ for a.e } x \in \mathbb{R}^n.$$

Then, for any $f_j \in L^{q_j(\cdot)}(\mathbb{R}^n)$, $j = 1, \dots, m$, one has

$$\|f_1 \cdots f_m\|_{q(\cdot)} \leq C \|f_1\|_{q_1(\cdot)} \cdots \|f_m\|_{q_m(\cdot)}.$$

Lemma 3.3. (see [14]) Let \mathcal{F} denote a family of ordered pairs of measurable functions (f, g) . suppose that for some fixed q_0 with $0 < q_0 < \infty$ and every weight $\omega \in A_1$ such that

$$\int_{\mathbb{R}^n} |f(x)|^{q_0} \omega(x) dx \leq C_0 \int_{\mathbb{R}^n} |g(x)|^{q_0} \omega(x) dx.$$

Let $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ with $q_0 \leq q_-$. If $(q(\cdot)/q_0)' \in \mathcal{B}(\mathbb{R}^n)$, then there exists a positive constant C , such that for all $(f, g) \in \mathcal{F}$,

$$\|f\|_{q(\cdot)} \leq C \|g\|_{q(\cdot)}.$$

Lemma 3.4. (see [3]) Let $0 < \alpha < n$ and $p(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $p_+ < \frac{n}{\alpha}$ and $\frac{1}{q(\cdot)} = \frac{1}{p(\cdot)} - \frac{\alpha}{n}$, then

$$\|\mathcal{M}_\alpha f\|_{q(\cdot)} \leq C \|f\|_{p(\cdot)}.$$

Theorem 3.1. Let $m \geq 2$, T_α be an m -linear fractional integral operator with generalized kernel as Definition 1.2 and $\sum_{k=1}^{\infty} C_k < \infty$. Given $\frac{p_i}{p_0}(\cdot) \in \mathcal{P}^{\log}(\mathbb{R}^n)$, $0 < \alpha_i < \frac{n}{p_{i,+}}$, $i = 1, 2, \dots, m$, $\alpha = \alpha_1 + \dots + \alpha_m$, $\frac{1}{p(\cdot)} = \frac{1}{p_1(\cdot)} + \dots + \frac{1}{p_m(\cdot)}$, $\frac{1}{q(\cdot)} = \frac{1}{p(\cdot)} - \frac{\alpha}{n}$ and $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, then there exists a positive constant C such that

$$\left\| T_\alpha(\vec{f}) \right\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j(\cdot)}(\mathbb{R}^n)}.$$

Proof. Since $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, by Lemma 3.1, there exists a q_0 with $1 < q_0 < q_-$ such that $(q(\cdot)/q_0)' \in \mathcal{B}(\mathbb{R}^n)$. For this q_0 and any $\omega \in A_1$, choosing $0 < \delta < \min \left\{ 1, \frac{sn}{n-s\alpha} \right\}$, by Theorem 2.1, we obtain

$$\left\| T_\alpha(\vec{f}) \right\|_{L^{q_0}(\omega)} \leq \left\| M_\delta(T_\alpha(\vec{f})) \right\|_{L^{q_0}(\omega)} \leq \left\| M_\delta^\#(T_\alpha(\vec{f})) \right\|_{L^{q_0}(\omega)} \leq C \left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{L^{q_0}(\omega)},$$

then we can get

$$\int_{\mathbb{R}^n} \left| T_\alpha(\vec{f})(x) \right|^{q_0} \omega(x) dx \leq C \int_{\mathbb{R}^n} \left| \mathcal{M}_{\alpha, p'_0}(\vec{f})(x) \right|^{q_0} \omega(x) dx$$

holds for all m -tuples $\vec{f} = (f_1, \dots, f_m)$ of bounded functions with compact support.

Apply Lemma 3.3 to the pair $(T_\alpha(\vec{f}), \mathcal{M}_{\alpha, p'_0}(\vec{f}))$ and obtain

$$\left\| T_\alpha(\vec{f}) \right\|_{q(\cdot)} \leq C \left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{q(\cdot)}. \quad (3.1)$$

We can get $0 < \alpha_i < \frac{n}{p_{i,+}} < \frac{n}{p'_0}$. Denote by $\frac{1}{q_i(\cdot)} = \frac{1}{p_i(\cdot)} - \frac{\alpha_i}{n}$ ($i = 1, \dots, m$), then $\frac{1}{q(\cdot)} = \frac{1}{q_1(\cdot)} + \dots + \frac{1}{q_m(\cdot)}$. By Definition 2.3, it is easy to see that

$$\mathcal{M}_{\alpha, p'_0}(\vec{f})(x) \leq \prod_{i=1}^m \mathcal{M}_{\alpha_i, p'_0}(f_i)(x) \text{ for } x \in \mathbb{R}^n. \quad (3.2)$$

Let $\beta_i = \alpha_i p'_0$. From (3.1), (3.2), Lemma 3.2, and Lemma 3.4, it follows that

$$\begin{aligned} \left\| T_\alpha(\vec{f})(x) \right\|_{q(\cdot)} &\leq C \left\| \mathcal{M}_{\alpha, p'_0}(\vec{f}) \right\|_{q(\cdot)} \leq C \prod_{i=1}^m \left\| \mathcal{M}_{\alpha_i, p'_0}(f_i) \right\|_{q_i(\cdot)} \\ &= C \prod_{i=1}^m \left\| \left[\mathcal{M}_{\beta_i}(f_i^{p'_0}) \right]^{\frac{1}{p'_0}} \right\|_{q_i(\cdot)} = C \prod_{i=1}^m \left\| \mathcal{M}_{\beta_i}(f_i^{p'_0}) \right\|_{\frac{q_i(\cdot)}{p'_0}}^{\frac{1}{p'_0}} \\ &\leq C \prod_{i=1}^m \left\| |f_i|^{p'_0} \right\|_{\frac{p_i(\cdot)}{p'_0}}^{\frac{1}{p'_0}} = C \prod_{i=1}^m \|f_i\|_{p_i(\cdot)}. \end{aligned}$$

Thus, Theorem 3.1 is proved. \square

Remark 3.2. When the generalized kernel condition goes back to the Dini kernel condition, the boundedness of m -linear fractional integral operators with Dini's type kernels on variable exponent Lebesgue spaces has been demonstrated by Wu and Zhang in [37].

Theorem 3.2. Let $m \geq 2$, T_α be an m -linear fractional integral operator with generalized kernel as Definition 1.2 and $\sum_{k=1}^{\infty} kC_k < \infty$. Given $\frac{p_i}{p'_0}(\cdot) \in \mathcal{P}^{log}(\mathbb{R}^n)$, $0 < \alpha_i < \frac{n}{p_{i,+}}$, $i = 1, 2, \dots, m$, $\alpha = \alpha_1 + \dots + \alpha_m$, $\frac{1}{p(\cdot)} = \frac{1}{p_1(\cdot)} + \dots + \frac{1}{p_m(\cdot)}$, $\frac{1}{q(\cdot)} = \frac{1}{p(\cdot)} - \frac{\alpha}{n}$, $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$ and $\vec{b} \in \text{BMO}^m$, then there exists a positive constant C such that

$$\left\| T_{\vec{b}, \alpha}(\vec{f}) \right\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C \|\vec{b}\|_{(\text{BMO})^m} \prod_{j=1}^m \|f_j\|_{L^{p_j(\cdot)}(\mathbb{R}^n)}.$$

Proof. Since $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, by Lemma 3.1 there exists a q_0 with $1 < q_0 < q_-$ such that $(q(\cdot)/q_0)^\prime \in \mathcal{B}(\mathbb{R}^n)$. For this q_0 and any $\omega \in A_1$, choosing $0 < \delta < \varepsilon < \min\left\{1, \frac{sn}{n-s\alpha}\right\}$ and $p'_0 < t < \min_{1 \leq i \leq m} p_{i,-}$, by Theorem 2.2, we obtain

$$\begin{aligned} \|T_{\vec{b},\alpha}(\vec{f})\|_{L^{q_0}(\omega)} &\leq \|M_\delta(T_{\vec{b},\alpha}(\vec{f}))\|_{L^{q_0}(\omega)} \leq \|M_\delta^\#(T_{\vec{b},\alpha}(\vec{f}))\|_{L^{q_0}(\omega)} \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \|M_\varepsilon(T_\alpha(\vec{f})) + \mathcal{M}_{\alpha,t}(\vec{f})\|_{L^{q_0}(\omega)} \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \left(\|M_\varepsilon(T_\alpha(\vec{f}))\|_{L^{q_0}(\omega)} + \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{L^{q_0}(\omega)} \right) \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \left(\|M_\varepsilon^\#(T_\alpha(\vec{f}))\|_{L^{q_0}(\omega)} + \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{L^{q_0}(\omega)} \right) \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \left(\|\mathcal{M}_{\alpha,p'_0}(\vec{f})\|_{L^{q_0}(\omega)} + \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{L^{q_0}(\omega)} \right) \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{L^{q_0}(\omega)}, \end{aligned}$$

then we can get

$$\int_{\mathbb{R}^n} |T_{\vec{b},\alpha}(\vec{f})(x)|^{q_0} \omega(x) dx \leq C\|\vec{b}\|_{(\text{BMO})^m}^{q_0} \int_{\mathbb{R}^n} |\mathcal{M}_{\alpha,t}(\vec{f})(x)|^{q_0} \omega(x) dx$$

holds for all m -tuples $\vec{f} = (f_1, \dots, f_m)$ of bounded functions with compact support.

Apply Lemma 3.3 to the pair $(T_{\vec{b},\alpha}(\vec{f}), \mathcal{M}_{\alpha,t}(\vec{f}))$ and obtain

$$\|T_{\vec{b},\alpha}(\vec{f})\|_{q(\cdot)} \leq C\|\vec{b}\|_{(\text{BMO})^m} \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{q(\cdot)}. \quad (3.3)$$

We can get $0 < \alpha_i < \frac{n}{p_{i,+}} < \frac{n}{t}$. Denote by $\frac{1}{q_i(\cdot)} = \frac{1}{p_i(\cdot)} - \frac{\alpha_i}{n}$ ($i = 1, \dots, m$), then $\frac{1}{q(\cdot)} = \frac{1}{q_1(\cdot)} + \dots + \frac{1}{q_m(\cdot)}$. By Definition 2.3, it is easy to see that

$$\mathcal{M}_{\alpha,t}(\vec{f})(x) \leq \prod_{i=1}^m \mathcal{M}_{\alpha_i,t}(f_i)(x) \text{ for } x \in \mathbb{R}^n. \quad (3.4)$$

Let $\beta_i = \alpha_i t$. The fact $\frac{p_i}{p'_0}(\cdot) \in \mathcal{P}^{log}(\mathbb{R}^n)$ implies that $\frac{p_i}{t}(\cdot) \in \mathcal{P}^{log}(\mathbb{R}^n)$, $i = 1, \dots, m$. From (3.3), (3.4), Lemma 3.2 and Lemma 3.4, it follows that

$$\begin{aligned} \|T_{\vec{b},\alpha}(\vec{f})(x)\|_{q(\cdot)} &\leq C\|\vec{b}\|_{(\text{BMO})^m} \|\mathcal{M}_{\alpha,t}(\vec{f})\|_{q(\cdot)} \leq C\|\vec{b}\|_{(\text{BMO})^m} \prod_{i=1}^m \|\mathcal{M}_{\alpha_i,t}(f_i)\|_{q_i(\cdot)} \\ &= C\|\vec{b}\|_{(\text{BMO})^m} \prod_{i=1}^m \left\| [\mathcal{M}_{\beta_i}(f_i^t)]^{\frac{1}{t}} \right\|_{q_i(\cdot)} = C\|\vec{b}\|_{(\text{BMO})^m} \prod_{i=1}^m \|\mathcal{M}_{\beta_i}(f_i^t)\|_{\frac{q_i(\cdot)}{t}}^{\frac{1}{t}} \\ &\leq C\|\vec{b}\|_{(\text{BMO})^m} \prod_{i=1}^m \| |f_i|^t \|_{\frac{p_i(\cdot)}{t}}^{\frac{1}{t}} = C\|\vec{b}\|_{(\text{BMO})^m} \prod_{i=1}^m \|f_i\|_{p_i(\cdot)}. \end{aligned}$$

Thus, Theorem 3.2 is proved. \square

Corollary 3.1. When the generalized kernel condition goes back to the Dini kernel condition, the boundedness of multilinear commutators generalized by m -linear fractional integral operators with Dini's type kernels and BMO functions on variable exponent Lebesgue spaces remains new.

References

- [1] A. Bernardis, S. Hartzstein and G. Pradolini, Weighted inequalities for commutators of fractional integrals on spaces of homogeneous type, *J. Math. Anal. Appl.*, 322 (2006), 825–846.
- [2] B. Bongioanni, E. Haboure and O. Salinas, Classes of weights related to Schrodinger operators, *J. Math. Anal. Appl.*, 373 (2011), 563–579.
- [3] C. Capone, D. Cruz-Urbe and A. Fiorenza, The fractional maximal operator and fractional integrals on variable L^p spaces, *Rev. Mat. Iberoam.*, 23 (2007), 743–770.
- [4] S. Chen and H. Wu, Multiple weighted estimates for commutators of multilinear fractional integral operators, *Sci. China Math.*, 56 (2013), 1879–1894.
- [5] X. Chen and Q. Xue, Weighted estimates for a class of multilinear fractional type operators, *J. Math. Anal. Appl.*, 362 (2010), 355–373.
- [6] R. R. Coifman and Y. Meyer, On commutators of singular integrals and bilinear singular integrals, *Trans. Amer. Math. Soc.*, 212 (1975), 315–331.
- [7] R. R. Coifman and Y. Meyer, Commutateurs d'intégrales singulières et opérateurs multilinéaires, *Ann. Inst. Fourier (Grenoble)*, 28 (1978), 177–202.
- [8] D. Cruz-Urbe, A. Fiorenza and C. J. Neugebauer, The maximal function on variable L^p spaces. *Ann. Acad. Sci. Fenn. Math.*, 28 (2003), 223–238.
- [9] D. Cruz-Urbe, A. Fiorenza, J. M. Martell and C. Pérez, The boundedness of classical operators on variable L^p spaces, *Ann. Acad. Sci. Fenn. Math.*, 31 (2006), 239–264.
- [10] D. Cruz-Urbe and A. Fiorenza, Variable Lebesgue spaces, foundations and harmonic analysis, *Appl. Numer. Harmon. Anal.*, Birkhäuser/Springer, Heidelberg, 2013, x+312 pp.
- [11] D. Cruz-Urbe, K. Moen and H. V. Nguyen, Multilinear fractional Calderón-Zygmund operators on weighted Hardy spaces, *Houston J. Math.*, 45 (2019), 853–871.
- [12] E. Dalmasso, G. Pradolini and W. Ramos, The effect of the smoothness of fractional type operators over their commutators with Lipschitz symbols on weighted spaces, *Fract. Calc. Appl. Anal.*, 21 (2018), 628–653.
- [13] L. Diening, Maximal function on Musielak-Orlicz spaces and generalized Lebesgue spaces, *Bull. Sci. math.*, 129 (2005), 657–700.
- [14] L. Diening, P. Harjulehto, P. Hästö and M. Růžička, Lebesgue and Sobolev spaces with variable exponents, Springer, Heidelberg, 2011, x+509 pp.
- [15] C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.*, 129 (1972), 137–193.
- [16] Z. Fu, L. Grafakos, Y. Lin, Y. Wu and S. Yang, Riesz transform associated with the fractional Fourier transform and applications in image edge detection, *Appl. Comput. Harmon. Anal.*, 66 (2023), 211–235.
- [17] Z. Fu, Y. Lin, D. Yang and S. Yang, Fractional Fourier transforms meet Riesz potentials and image processing, *SIAM J. Imaging Sci.*, 17 (2024), 476–500.
- [18] L. Gao, Y. Lin and S. Yang, Multiple weighted estimates for multilinear commutators of multilinear singular integrals with generalized kernels, *J. Korean Math. Soc.*, 61 (2024), 207–226.
- [19] L. Grafakos, On multilinear fractional integrals, *Studia Math.*, 102 (1992), 49–56.
- [20] L. Grafakos and R. H. Torres, Multilinear Calderón-Zygmund theory, *Adv. Math.*, 165 (2002), 124–164.
- [21] L. Grafakos and D. He, Multilinear Calderón-Zygmund operators on Hardy spaces, *J. Math. Anal. Appl.*, 416 (2014), 511–521.
- [22] C. E. Kenig and E. M. Stein, Multilinear estimates and fractional integration, *Math. Res. Lett.*, 6 (1999), 1–15.
- [23] O. Kováčik, J. Rákosník, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, *Czechoslovak Math. J.*, 41 (1991), 592–618.

- [24] K. Langley and S. J. Anderson, The Riesz transform and simultaneous representations of phase energy and orientation in spatial vision, *Vision Research*, 50 (2010), 1748–1765.
- [25] A. K. Lerner, S. Ombrosi, C. Pérez, R. H. Torres and R. González, New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory, *Adv. Math.*, 220 (2009), 1222–1264.
- [26] Y. Lin and S. Lu, Boundedness of multilinear singular integral operators on Hardy and Herz-type spaces, *Hokkaido Math. J.*, 36 (2007), 585–613.
- [27] Y. Lin and Y. Xiao, Multilinear singular integral operators with generalized kernel and their multilinear commutators, *Acta Math. Sin. (Engl. Ser.)*, 33 (2017), 1443–1462.
- [28] Y. Lin and N. Zhang, Sharp maximal and weighted estimates for multilinear iterated commutators of multilinear integrals with generalized kernels, *J. Inequal. Appl.*, 276 (2017), 1–15.
- [29] G. Lu and P. Zhang, Multilinear Calderón-Zygmund operators with kernels of Dini's type and applications, *Nonlinear Anal.*, 107 (2014), 92–117.
- [30] S. Lu, *Four Lectures on Real H^p Spaces*, World Scientific Publishing Co., Inc., River Edge, NJ, 1995, viii+217 pp.
- [31] S. Lu, Y. Ding and D. Yan, *Singular integrals and related topics*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007, viii+272 pp.
- [32] K. Moen, Weighted inequalities for multilinear fractional integral operators, *Collect. Math.*, 60 (2009), 213–238.
- [33] G. Pradolini, Weighted inequalities and pointwise estimates for the multilinear fractional integral and maximal operators, *J. Math. Anal. Appl.*, 367 (2010), 640–656.
- [34] Z. Si and S. Lu, Weighted estimates for iterated commutators of multilinear fractional operators, *Acta Math. Sin. (Engl. Ser.)*, 28 (2012), 1769–1778.
- [35] H. Wang and J. Xu, Multilinear fractional integral operators on central Morrey spaces with variable exponent, *J. Inequal. Appl.*, 1 (2019), 1–23.
- [36] B. Wei, S. Yang and Y. Lin, Multiple weighted norm inequalities for multilinear strongly singular integral operators with generalized kernels, *Bull. Iran. Math. Soc.*, 49 (2023), 1–30.
- [37] J. Wu and P. Zhang, Multilinear fractional Calderón-Zygmund operators with Dini's type kernel, *arxiv: 2307.01402*.
- [38] Q. Xue, Weighted estimates for the iterated commutators of multilinear maximal and fractional type operators, *Studia Math.*, 217 (2013), 97–122.
- [39] S. Yang, P. Li and Y. Lin, Multiple weight inequalities for multilinear singular integral operator with generalized kernels, *Adv. Math. (China)*, 53 (2024), 162–176.
- [40] K. Yabuta, Generalizations of Calderón-Zygmund operators, *Stud. Math.*, 82 (1985), 17–31.
- [41] L. Zhang and H. Li, Local image patterns using Riesz transforms: with applications to palmprint and finger-knuckle-print recognition, *Image and Vision Computing*, 30 (2012).
- [42] L. Zhang, L. Zhang and X. Mou, RFSIM: A feature based image quality assessment metric using Riesz transforms, *IEEE International Conference on Image Processing. IEEE*, 2010.
- [43] P. Zhang and J. Sun, Commutators of multilinear Calderón-Zygmund operators with kernels of Dini's type and applications, *J. Math. Inequal.*, 13 (2019), 1071–1093.

School of Science, China University of Mining and Technology, Beijing 100083, People's Republic of China

E-mails: linyan@cumtb.edu.cn (Y. Lin)

zhaoyuhang@student.cumtb.edu.cn (Y. Zhao)

yangshuhui@student.cumtb.edu.cn (S. Yang)