

# Local linear smoothing for regression surfaces on the simplex using Dirichlet kernels

Christian Genest<sup>1\*</sup> and Frédéric Ouimet<sup>1</sup>

<sup>1\*</sup>Department of Mathematics and Statistics, McGill University,  
805, rue Sherbrooke ouest, Montréal (Québec) Canada H3A 0B9.

\*Corresponding author(s). E-mail(s): [christian.genest@mcgill.ca](mailto:christian.genest@mcgill.ca);  
Contributing authors: [frederic.ouimet2@mcgill.ca](mailto:frederic.ouimet2@mcgill.ca);

## Abstract

This paper introduces a local linear smoother for regression surfaces on the simplex. The estimator solves a least-squares regression problem weighted by a locally adaptive Dirichlet kernel, ensuring excellent boundary properties. Asymptotic results for the bias, variance, mean squared error, and mean integrated squared error are derived, generalizing the univariate results of [Chen \(2002\)](#). A simulation study shows that the proposed local linear estimator with Dirichlet kernel outperforms its only direct competitor in the literature, the Nadaraya–Watson estimator with Dirichlet kernel due to [Bouzebda, Nezzal, and Elhattab \(2024\)](#).

**Keywords:** Adaptive estimator, asymmetric kernel, beta kernel, boundary bias, Dirichlet kernel, local linear smoother, mean integrated squared error, Nadaraya–Watson estimator, nonparametric regression, regression surface, simplex.

**MSC Classification:** Primary: 62G08; Secondary: 62G05 , 62H12

## 1 Introduction

Although Dirichlet kernels were proposed almost 40 years ago by [Aitchison and Lauder \(1985\)](#) for density estimation on the simplex, their application to nonparametric regression has only been considered recently. [Bouzebda et al. \(2024\)](#) introduced, in the larger framework of conditional  $U$ -statistics, a version of the Nadaraya–Watson (NW) estimator weighted by a locally adaptive Dirichlet kernel and studied some of its asymptotic properties. This development follows earlier theoretical work on Dirichlet

kernel density estimators by [Ouimet and Tolosana-Delgado \(2022\)](#) and [Bertin, Genest, Klutchnikoff, and Ouimet \(2023\)](#). [Bouzebda et al. \(2024\)](#) also investigated a NW estimator weighted by a locally adaptive multinomial kernel, building on the Bernstein approach to density estimation on the simplex proposed by [Tenbusch \(1994\)](#) in two dimensions and expanded upon by [Ouimet \(2021, 2022\)](#) in arbitrary dimension.

As an alternative to the estimator proposed by [Bouzebda et al. \(2024\)](#), the present paper introduces a local linear (LL) smoother for regression surfaces using Dirichlet kernels. This approach, which is introduced in Section 2 along with other preliminary definitions and notations, leverages the strengths of asymmetric kernels and LL smoothing to provide an efficient estimation method on the simplex with excellent boundary properties, addressing the practical limitations of implementing the multivariate LL smoother of [Ruppert and Wand \(1994\)](#) with a variable bandwidth.

Asymptotic results for the bias, variance, mean squared error (MSE), and mean integrated squared error (MISE) of the proposed smoother are stated in Section 3 and proved in Section 4, based on technical lemmas which are relegated to the Appendix. These results, which generalize those established by [Chen \(2002\)](#) in one dimension, are complemented in Section 5 with a simulation study which demonstrates the superior performance of the LL estimator with Dirichlet kernel compared to its only direct competitor, the NW estimator with Dirichlet kernel recently proposed by [Bouzebda et al. \(2024\)](#). A numerical example is then presented in Section 6 based on data from [Coakley and Rust \(1968\)](#) popularized by [Aitchison \(1986\)](#).

The rest of this section provides context. First, LL smoothers with variable bandwidth are not new; they were introduced by [Fan and Gijbels \(1992\)](#) as a tool to estimate the functional relationship between a predictor and a response. Such estimators solve a locally weighted least-squares regression problem, combining the strengths of LL smoothing, as developed by [Stone \(1977\)](#) and [Cleveland \(1979\)](#), with the local adaptivity resulting from the optimization of the variable bandwidth. A distinct advantage of LL estimation is that it adapts to both random and fixed designs, and exhibits no boundary effects, i.e., requires no modification near the boundary to reduce bias. As shown by [Fan \(1992\)](#), LL estimators also offer higher efficiency than traditional NW and Gasser–Müller (GM) regression estimators; their minimax properties were investigated by [Fan \(1993\)](#). Although LL estimation was later extended by [Ruppert and Wand \(1994\)](#) to the multiple predictor setting considered herein, LL smoothers are unfortunately currently not a viable option for compositional data, due to lack of efficient methods to compute the variable bandwidth on such supports.

An alternative approach to addressing the boundary issues of traditional regression estimators was proposed by [Chen \(2000\)](#), who replaced the fixed kernel in GM estimators with a beta kernel. The beta kernel is locally adaptive through its parameters rather than through a variable bandwidth, i.e., the kernel parameters change with every estimation point  $\mathbf{s}$  on the simplex. This local adaptivity allows the beta kernel to adjust its shape to fit the data within  $[0, 1]$ , and, if appropriately transformed, within any finite interval. [Brown and Chen \(1999\)](#) proposed a similar approach where kernel weights are convoluted with a binomial distribution instead of partitioned, offering another effective mitigation strategy for boundary bias.

The approach adopted here was inspired by the work of [Chen \(2002\)](#), who combined the ideas of LL smoothing and asymmetric kernels by replacing Fan’s fixed kernel with variable bandwidth with a beta or gamma kernel. The former is used for design variables supported on  $[0, 1]$  and the latter for  $[0, \infty)$ . This combination ensures an intrinsic local adaptivity of the kernel and the asymptotic negligibility of boundary bias. As for Fan’s LL smoother, it is suitable for both fixed and random designs.

Chen’s approach to integrating locally adaptive beta/gamma kernels with LL smoothers has been influential in the literature on asymmetric kernel estimators. The results contained in the present paper generalize the results of [Chen \(2002\)](#) to the multivariate case and documents their value in the context of nonparametric regression for density estimation on the simplex.

Many other authors, including [Fé \(2014\)](#), [Funke and Hirukawa \(2021, 2024\)](#), and [Hirukawa, Murtazashvili, and Prokhorov \(2022, 2023\)](#), have expanded on Chen’s ideas in different settings. When the slope in the locally weighted least-squares problem is zero, the regression estimator simplifies to a NW estimator weighted by an asymmetric kernel. This estimator has been studied by [Shi and Song \(2016\)](#) for gamma kernels and by [Somé and Kokonendji \(2016\)](#) within the broader theory of associated kernels.

## 2 A local linear smoother based on Dirichlet kernels

Consider data  $(\mathbf{X}_1, Y_1), \dots, (\mathbf{X}_n, Y_n)$ , where for each  $i \in \{1, \dots, n\}$ ,  $Y_i$  is a response variable related to the random design point  $\mathbf{X}_i$  by the functional model

$$Y_i = m(\mathbf{X}_i) + \varepsilon_i, \quad (1)$$

where  $\varepsilon_i$  is a random error term and  $m(\mathbf{x}) = \mathbb{E}(Y \mid \mathbf{X} = \mathbf{x})$  is an unknown regression function supported on the unit simplex

$$\mathcal{S}_d = \{\mathbf{s} \in [0, 1]^d : \|\mathbf{s}\|_1 \leq 1\},$$

where  $\|\cdot\|_1$  denotes the  $\ell^1$  norm in  $\mathbb{R}^d$ .

Assume that the design points have a common density  $f$  and that the error terms  $\varepsilon_1, \dots, \varepsilon_n$  are uncorrelated with zero mean and variance  $\sigma^2(\mathbf{X}_1), \dots, \sigma^2(\mathbf{X}_n)$ , respectively. Further let  $\sigma^2(\mathbf{x}) = \text{Var}(Y \mid \mathbf{X} = \mathbf{x})$  denote the conditional variance.

A natural local linear (LL) estimator  $\hat{m}$  of  $m$  can be obtained by minimizing the locally weighted loss function

$$L(\alpha, \beta) = \sum_{i=1}^n \{Y_i - \alpha - \beta^\top (\mathbf{X}_i - \mathbf{s})\}^2 K_{\mathbf{s}/b+1, (1-\|\mathbf{s}\|_1)/b+1}(\mathbf{X}_i),$$

where for arbitrary  $\mathbf{u} = (u_1, \dots, u_d) \in (0, \infty)^d$ ,  $v \in (0, \infty)$ , and  $\mathbf{x} \in \mathcal{S}_d$ ,

$$K_{\mathbf{u}, v}(\mathbf{x}) = \frac{\Gamma(\|\mathbf{u}\|_1 + v)}{\Gamma(v) \prod_{i=1}^d \Gamma(u_i)} (1 - \|\mathbf{x}\|_1)^{v-1} \prod_{i=1}^d x_i^{u_i-1}$$

is the density of the Dirichlet( $\mathbf{u}, v$ ) distribution.

Precisely, if  $\mathbf{e}_1 = (1, 0, \dots, 0)$  is a  $(d+1) \times 1$  vector, then one can set

$$\hat{m}_{n,b}^{(LL)}(\mathbf{s}) = \hat{\alpha}_{\mathbf{s}} = \mathbf{e}_1^\top (\mathcal{X}_{\mathbf{s}}^\top W_{\mathbf{s}} \mathcal{X}_{\mathbf{s}})^{-1} \mathcal{X}_{\mathbf{s}}^\top W_{\mathbf{s}} \mathbf{Y}, \quad (2)$$

where

$$\mathcal{X}_{\mathbf{s}} = \begin{bmatrix} 1 & (\mathbf{X}_1 - \mathbf{s})^\top \\ \vdots & \vdots \\ 1 & (\mathbf{X}_n - \mathbf{s})^\top \end{bmatrix}_{n \times (d+1)}, \quad W_{\mathbf{s}} = \text{diag} \begin{bmatrix} \kappa_{\mathbf{s},b}(\mathbf{X}_1) \\ \vdots \\ \kappa_{\mathbf{s},b}(\mathbf{X}_n) \end{bmatrix}_{n \times 1}, \quad \mathbf{Y} = \begin{bmatrix} Y_1 \\ \vdots \\ Y_n \end{bmatrix}_{n \times 1},$$

and, for simplicity, the following notation is used henceforth:

$$\kappa_{\mathbf{s},b} = K_{\mathbf{s}/b+1, (1-\|\mathbf{s}\|_1)/b+1}.$$

Estimator (2), which is new, is a natural  $d$ -variate extension of the beta kernel estimator due to [Chen \(2002\)](#). As will be seen in [Section 3](#) and proved in [Section 4](#), it enjoys excellent sampling properties. As reported in [Section 5](#), its finite-sample behavior also makes it preferable to the alternative estimator recently proposed by [Bouzebda et al. \(2024\)](#).

In what follows, the notation  $u = \mathcal{O}(v)$  means that  $\limsup |u/v| \leq C < \infty$  as  $n \rightarrow \infty$  or  $b \rightarrow \infty$ , depending on the context, where the positive constant  $C \in (0, \infty)$  may depend on the target regression function  $m$ , the design density  $f$ , the conditional variance function  $\sigma^2$ , or the dimension  $d$ , but no other variable unless explicitly written as a subscript. Similarly, the notation  $u = o(v)$  means that  $|u/v| \rightarrow 0$  as  $n \rightarrow \infty$  or  $b \rightarrow \infty$ . Subscripts indicate on which parameters the convergence rate can depend. Except for [Lemma 2](#) in the [Appendix](#), it is always assumed that the bandwidth parameter  $b = b(n)$  depends on the sample size,  $n$ , in such a way that  $b \rightarrow 0$  as  $n \rightarrow \infty$ . The shorthand  $[d] = \{1, \dots, d\}$  is often used.

### 3 Results

For any subset of indices  $\mathcal{J} \subseteq [d]$ , define

$$\psi_{\mathcal{J}}(\mathbf{s}) = \left\{ (4\pi)^{d-|\mathcal{J}|} (1 - \|\mathbf{s}\|_1) \prod_{i \in [d] \setminus \mathcal{J}} s_i \right\}^{-1/2} \quad (3)$$

and

$$h_{\mathcal{J}}(\mathbf{s}) = \begin{cases} \sum_{k, \ell \in [d] \setminus \mathcal{J}} \frac{1}{2} (s_k \mathbb{1}_{\{k=\ell\}} - s_k s_\ell) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) & \text{if } \mathcal{J} \subsetneq [d], \\ \sum_{k, \ell \in [d]} \frac{1}{2} \{(\lambda_k + 1) \mathbb{1}_{\{k=\ell\}} + 1\} \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) & \text{if } \mathcal{J} = [d], \end{cases}$$

with the convention that  $\prod_{\emptyset} = 1$  and  $\sum_{\emptyset} = 0$ . For any design point  $\mathbf{X}$ , define

$$A_b(\mathbf{s}) = \mathbb{E}\{\kappa_{\mathbf{s},b}(\mathbf{X})^2\}.$$

Further assume the following conditions:

- (A1)  $m \in C^2(\mathcal{S}_d)$ ;  $f$  and  $\sigma^2$  are Lipschitz continuous on  $\mathcal{S}_d$ .
- (A2)  $f(\mathbf{x}) \geq f_c > 0$  and  $\sigma^2(\mathbf{x}) \leq \sigma_c^2 < \infty$  for all  $\mathbf{x} \in \mathcal{S}_d$ .
- (A3)  $b \rightarrow 0$  and  $nb^{d/2} \rightarrow \infty$  as  $n \rightarrow \infty$ .

Under Assumptions (A1)–(A3), Proposition 1 below presents asymptotic expressions for the bias and variance of estimator (2) at  $\mathbf{s} \in \text{Int}(\mathcal{S}_d)$ , for the cases in which  $\mathbf{s}$  is well inside the simplex and near the boundary. This result extends similar derivations by Chen (2002) in the univariate case  $\mathcal{S}_1 \equiv [0, 1]$ . The index set  $\mathcal{J} \subseteq [d]$  corresponds to the dimensions for which  $\mathbf{s}$  is close to the boundary of the simplex.

**Proposition 1.** *Suppose that Assumptions (A1)–(A3) hold and let  $\mathcal{J} \subseteq [d]$  and  $\boldsymbol{\lambda} \in (0, \infty)^d$  be given. If a sequence  $\mathbf{s} = \mathbf{s}(b)$  is selected inside  $\text{Int}(\mathcal{S}_d)$  such that  $s_i/b \rightarrow \lambda_i$  for all  $i \in \mathcal{J}$ ,  $s_i$  is fixed for all  $i \in [d] \setminus \mathcal{J}$ , and  $(1 - \|\mathbf{s}\|_1)$  is fixed, then*

$$\text{Bias}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\} = b^{1+\mathbb{1}_{\{\mathcal{J}=[d]\}}}\{h_{\mathcal{J}}(\mathbf{s}) + \mathcal{O}\{b^{3/2(1+\mathbb{1}_{\{\mathcal{J}=[d]\}})}\}\} + \mathcal{O}\{n^{-1}A_b(\mathbf{s})\}$$

and

$$\text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\} = n^{-1}b^{-(d+|\mathcal{J}|)/2} \times \left\{ \frac{\psi_{\mathcal{J}}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} \prod_{i \in \mathcal{J}} \frac{\Gamma(2\lambda_i + 1)}{2^{2\lambda_i+1}\Gamma^2(\lambda_i + 1)} + o_{\boldsymbol{\lambda},\mathbf{s}}(1) \right\},$$

where the term  $A_b(\mathbf{s})$  is bounded and estimated in Lemma 2.

An expression for the MSE follows immediately from the above proposition. An asymptotic expression is also available for the locally optimal bandwidth under the MSE criterion. Specifically,  $\text{MSE}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\}$  is given, for any  $\mathbf{s} \in \text{Int}(\mathcal{S}_d)$ , by

$$\begin{aligned} n^{-1}b^{-(d+|\mathcal{J}|)/2} \frac{\psi_{\mathcal{J}}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} \prod_{i \in \mathcal{J}} \frac{\Gamma(2\lambda_i + 1)}{2^{2\lambda_i+1}\Gamma^2(\lambda_i + 1)} &+ b^{2+2\mathbb{1}_{\{\mathcal{J}=[d]\}}}\{h_{\mathcal{J}}(\mathbf{s})\}^2 \\ &+ \mathcal{O}_{\boldsymbol{\lambda},\mathbf{s}}(n^{-1}b^{-(d+|\mathcal{J}|-1)/2}) + \mathcal{O}\{b^{5/2(1+\mathbb{1}_{\{\mathcal{J}=[d]\}})}\} \\ &+ \mathcal{O}\{b^{1+\mathbb{1}_{\{\mathcal{J}=[d]\}}}\{n^{-1}A_b(\mathbf{s})\}\} + \mathcal{O}\{n^{-2}A_b(\mathbf{s})^2\}. \end{aligned}$$

In particular, if  $h_{\mathcal{J}}(\mathbf{s}) \neq 0$  and  $R_{\mathcal{J}} \equiv d + |\mathcal{J}| + 4 + 4\mathbb{1}_{\{\mathcal{J}=[d]\}}$ , the asymptotically optimal choice of  $b$  with respect to MSE is

$$b_{\text{opt}}(\mathbf{s}) = \left[ n^{-1} \times \frac{d + |\mathcal{J}|}{4 + 4\mathbb{1}_{\{\mathcal{J}=[d]\}}} \times \frac{\psi_{\mathcal{J}}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})\{h_{\mathcal{J}}(\mathbf{s})\}^2} \prod_{i \in \mathcal{J}} \frac{\Gamma(2\lambda_i + 1)}{2^{2\lambda_i+1}\Gamma^2(\lambda_i + 1)} \right]^{2/R_{\mathcal{J}}},$$

and the corresponding value of the MSE is

$$\begin{aligned} \text{MSE}\{\hat{m}_{n,b_{\text{opt}}}^{(\text{LL})}(\mathbf{s})\} &= n^{-(4+4\mathbb{1}_{\{\mathcal{J}=[d]\}})/R_{\mathcal{J}}} \left(1 + \frac{d + |\mathcal{J}|}{4 + 4\mathbb{1}_{\{\mathcal{J}=[d]\}}}\right) \left(\frac{d + |\mathcal{J}|}{4 + 4\mathbb{1}_{\{\mathcal{J}=[d]\}}}\right)^{-(d+|\mathcal{J}|)/R_{\mathcal{J}}} \\ &\quad \times \left\{ \frac{\psi_{\mathcal{J}}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} \prod_{i \in \mathcal{J}} \frac{\Gamma(2\lambda_i + 1)}{2^{2\lambda_i+1}\Gamma^2(\lambda_i + 1)} \right\}^{(4+4\mathbb{1}_{\{\mathcal{J}=[d]\}})/R_{\mathcal{J}}} \\ &\quad \times [\{h_{\mathcal{J}}(\mathbf{s})\}^2]^{(d+|\mathcal{J}|)/R_{\mathcal{J}}} + o_{\mathbf{s}}\left(n^{-(4+4\mathbb{1}_{\{\mathcal{J}=[d]\}})/R_{\mathcal{J}}}\right). \end{aligned}$$

By integrating the MSE and showing that the contribution from points near the boundary is negligible, one obtains an expression for the MISE, viz.

$$\text{MISE}\{\hat{m}_{n,b}^{(\text{LL})}\} = \int_{S_d} \mathbb{E}\{|\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) - m(\mathbf{s})|^2\} d\mathbf{s}.$$

This expression, provided below, could be used to implement a plug-in method; see, e.g., Section 3.6 of [Chacón and Duong \(2018\)](#).

**Proposition 2.** *Under Assumptions (A1)–(A3), one has*

$$\begin{aligned} \text{MISE}\{\hat{m}_{n,b}^{(\text{LL})}\} &= n^{-1}b^{-d/2} \int_{S_d} \frac{\psi_{\emptyset}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} d\mathbf{s} \\ &\quad + b^2 \int_{S_d} \{h_{\emptyset}(\mathbf{s})\}^2 d\mathbf{s} + o(n^{-1}b^{-d/2}) + o(b^2). \end{aligned}$$

In particular, if  $\int_{S_d} \{h_{\emptyset}(\mathbf{s})\}^2 d\mathbf{s} > 0$ , the asymptotically optimal choice of  $b$  with respect to MISE, denoted  $b_{\text{opt}}(\mathbf{s})$ , is given by

$$n^{-2/(d+4)}(d/4)^{2/(d+4)} \left\{ \int_{S_d} \frac{\psi_{\emptyset}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} d\mathbf{s} \right\}^{2/(d+4)} \left[ \int_{S_d} \{h_{\emptyset}(\mathbf{s})\}^2 d\mathbf{s} \right]^{-2/(d+4)},$$

and the corresponding value of the MISE is

$$\begin{aligned} \text{MISE}\{\hat{m}_{n,b_{\text{opt}}}^{(\text{LL})}\} &= n^{-4/(d+4)} \left\{ \frac{1 + d/4}{(d/4)^{d/(d+4)}} \right\} \left\{ \int_{S_d} \frac{\psi_{\emptyset}(\mathbf{s})\sigma^2(\mathbf{s})}{f(\mathbf{s})} d\mathbf{s} \right\}^{4/(d+4)} \\ &\quad \times \left[ \int_{S_d} \{h_{\emptyset}(\mathbf{s})\}^2 d\mathbf{s} \right]^{d/(d+4)} + o(n^{-4/(d+4)}). \end{aligned}$$

## 4 Mathematical arguments

This section contains detailed proofs of Propositions 1 and 2.

## 4.1 Proof of Proposition 1

To prove Proposition 1, first note that Assumption (A1) and a stochastic Taylor expansion (see Theorem 18.18 of Aliprantis and Border (2006)) together imply that

$$\mathbb{E}(Y | \mathbf{X}_1, \dots, \mathbf{X}_n) = \begin{bmatrix} m(\mathbf{X}_1) \\ \vdots \\ m(\mathbf{X}_n) \end{bmatrix}_{n \times 1} = \mathcal{X}_s \begin{bmatrix} m(\mathbf{s}) \\ \frac{\partial}{\partial s_1} m(\mathbf{s}) \\ \vdots \\ \frac{\partial}{\partial s_d} m(\mathbf{s}) \end{bmatrix}_{(d+1) \times 1} + \frac{1}{2} \Lambda_m(\mathbf{s}),$$

where

$$\Lambda_m(\mathbf{s}) = \begin{bmatrix} \Lambda_{m,1}(\mathbf{s}) \\ \vdots \\ \Lambda_{m,n}(\mathbf{s}) \end{bmatrix}_{n \times 1}$$

and

$$\Lambda_{m,i}(\mathbf{s}) = \sum_{k,\ell=1}^d (X_{i,k} - s_k)(X_{i,\ell} - s_\ell) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) \{1 + o_{\mathbb{P}}(1)\}.$$

Using the definition of the estimator in (2), one can write

$$\mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) - m(\mathbf{s}) | \mathbf{X}_1, \dots, \mathbf{X}_n\} = \frac{1}{2} \mathbf{e}_1^\top (n^{-1} \mathcal{X}_s^\top W_s \mathcal{X}_s)^{-1} n^{-1} \mathcal{X}_s^\top W_s \Lambda_m(\mathbf{s}). \quad (4)$$

If  $\boldsymbol{\xi}_s \sim \text{Dirichlet}[\mathbf{s}/b + \mathbf{1}, (1 - \|\mathbf{s}\|_1)/b + 1]$ , then Eq. (4) in Ouimet and Tolosana-Delgado (2022) shows that

$$\mathbb{E}(\boldsymbol{\xi}_s) = \mathbf{s} + \mathcal{O}(b), \quad \mathbb{E}\{(\boldsymbol{\xi}_s - \mathbf{s})(\boldsymbol{\xi}_s - \mathbf{s})^\top\} = b\{\text{diag}(\mathbf{s}) - \mathbf{s}\mathbf{s}^\top\} + \mathcal{O}(b^2).$$

By Assumption (A1), the weak law of large numbers and Chebyshev's inequality to control the probabilistic errors, it follows that

$$(n^{-1} \mathcal{X}_s^\top W_s \mathcal{X}_s)^{-1} = \begin{bmatrix} A_n & B_n \\ C_n & D_n \end{bmatrix}^{-1},$$

where

$$A_n = \frac{1}{n} \sum_{i=1}^n \kappa_{s,b}(\mathbf{X}_i) = f(\mathbf{s}) + \mathcal{O}_{\mathbb{P}}\{n^{-1} A_b(\mathbf{s})\} + \sum_{k=1}^d \mathcal{O}\{\mathbb{E}(|\xi_{s,k} - s_k|)\},$$

$$\begin{aligned} B_n &= \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s})^\top \kappa_{s,b}(\mathbf{X}_i) \\ &= \mathbb{E}(\boldsymbol{\xi}_s - \mathbf{s})^\top f(\mathbf{s}) + \mathcal{O}_{\mathbb{P}}\{n^{-1} A_b(\mathbf{s})\} + \sum_{k=1}^d \mathcal{O}\{\mathbb{E}(|\boldsymbol{\xi}_s - \mathbf{s}|^\top |\xi_{s,k} - s_k|)\}, \end{aligned}$$

$$\begin{aligned}
C_n &= \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s}) \kappa_{\mathbf{s},b}(\mathbf{X}_i) \\
&= \mathbb{E}(\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s}) f(\mathbf{s}) + \text{o}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\} + \sum_{k=1}^d \mathcal{O}\{\mathbb{E}(|\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s}| |\xi_{\mathbf{s},k} - s_k|)\},
\end{aligned}$$

and

$$\begin{aligned}
D_n &= \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s})(\mathbf{X}_i - \mathbf{s})^\top \kappa_{\mathbf{s},b}(\mathbf{X}_i) \\
&= \mathbb{E}\{(\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s})(\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s})^\top\} f(\mathbf{s}) + \text{o}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\} \\
&\quad + \sum_{k=1}^d \mathcal{O}\{\mathbb{E}(|\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s}| |\boldsymbol{\xi}_{\mathbf{s}} - \mathbf{s}|^\top |\xi_{\mathbf{s},k} - s_k|)\}.
\end{aligned}$$

Therefore, using Lemma 1 to invert the block matrix, one obtains

$$\begin{aligned}
&(n^{-1}\mathcal{X}_{\mathbf{s}}^\top W_{\mathbf{s}} \mathcal{X}_{\mathbf{s}})^{-1} \\
&= \left[ \begin{array}{c|c} f(\mathbf{s})^{-1} + \mathcal{O}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\} & \mathcal{O}(b\mathbf{1}^\top) + \text{o}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\mathbf{1}^\top\} \\ \hline \mathcal{O}(b\mathbf{1}) + \text{o}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\mathbf{1}\} & \star \end{array} \right]_{(d+1) \times (d+1)}, \quad (5)
\end{aligned}$$

where the expression replacing the symbol  $\star$  is irrelevant. It should be noted in these calculations that the inverse covariance matrix,  $\{\text{diag}(\mathbf{s}) - \mathbf{s}\mathbf{s}^\top\}^{-1}$ , is equal to  $\text{diag}(s_1^{-1}, \dots, s_d^{-1}) + s_{d+1}^{-1}\mathbf{1}\mathbf{1}^\top$  by Eq. (21) of [Tanabe and Sagae \(1992\)](#), and that it is bounded away from zero in each component regardless of the choice of sequence  $\mathbf{s} = \mathbf{s}(b)$  in the statement of the proposition.

Also, one has

$$n^{-1}\mathcal{X}_{\mathbf{s}}^\top W_{\mathbf{s}} \Lambda_m(\mathbf{s}) = \begin{bmatrix} E_n \\ F_n \end{bmatrix}_{(d+1) \times 1}, \quad (6)$$

where

$$E_n = \frac{1}{n} \sum_{i=1}^n \sum_{k,\ell=1}^d (X_{i,k} - s_k)(X_{i,\ell} - s_\ell) \kappa_{\mathbf{s},b}(\mathbf{X}_i) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) \{1 + \text{o}_{\mathbb{P}}(1)\},$$

and

$$F_n = \frac{1}{n} \sum_{i=1}^n \sum_{k,\ell=1}^d (X_{i,k} - s_k)(X_{i,\ell} - s_\ell) (\mathbf{X}_i - \mathbf{s}) \kappa_{\mathbf{s},b}(\mathbf{X}_i) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) \{1 + \text{o}_{\mathbb{P}}(1)\}.$$

In view of (5) and (6), one can then re-express (4) as

$$\begin{aligned} & \frac{f(\mathbf{s})^{-1}}{2n} \sum_{k,\ell=1}^d \sum_{i=1}^n (X_{i,k} - s_k)(X_{i,\ell} - s_\ell) \kappa_{\mathbf{s},b}(\mathbf{X}_i) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) \{1 + o_{\mathbb{P}}(1)\} \\ & + b \sum_{j,k,\ell=1}^d \mathcal{O} \left\{ \frac{1}{n} \sum_{i=1}^n |X_{i,j} - s_j| |X_{i,k} - s_k| |X_{i,\ell} - s_\ell| \kappa_{\mathbf{s},b}(\mathbf{X}_i) \right\} + \mathcal{O}_{\mathbb{P}}\{n^{-1}A_b(\mathbf{s})\}. \end{aligned}$$

Now, Hölder's inequality implies that, for all  $j, k, \ell \in [d]$ ,

$$\mathbb{E}(|\xi_{\mathbf{s},j} - s_j| |\xi_{\mathbf{s},k} - s_k| |\xi_{\mathbf{s},\ell} - s_\ell|) \leq \max_{1 \leq j \leq d} \{\mathbb{E}(|\xi_{\mathbf{s},j} - s_j|^6)\}^{3/6} = \mathcal{O}\{(b^3)^{3/6}\} = \mathcal{O}(b^{3/2}).$$

Therefore, by Assumption (A1), the weak law of large numbers and Chebyshev's inequality to control the probabilistic errors, taking the expectation in (4) yields

$$\begin{aligned} \mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) - m(\mathbf{s})\} &= \mathbb{E}[\mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) - m(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\}] \\ &= \frac{1}{2} \sum_{k,\ell=1}^d \mathbb{E}\{(\xi_{\mathbf{s},k} - s_k)(\xi_{\mathbf{s},\ell} - s_\ell)\} \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) \\ &\quad + \sum_{j,k,\ell=1}^d \mathcal{O}\{\mathbb{E}(|\xi_{\mathbf{s},j} - s_j| |\xi_{\mathbf{s},k} - s_k| |\xi_{\mathbf{s},\ell} - s_\ell|)\} + \mathcal{O}\{n^{-1}A_b(\mathbf{s})\}, \end{aligned}$$

which reduces to

$$\frac{b}{2} \sum_{k,\ell=1}^d (s_k \mathbb{1}_{\{k=\ell\}} - s_k s_\ell) \frac{\partial^2}{\partial s_k \partial s_\ell} m(\mathbf{s}) + \mathcal{O}(b^{3/2}) + \mathcal{O}\{n^{-1}A_b(\mathbf{s})\}. \quad (7)$$

This establishes the bias expression when  $\mathcal{J} = \emptyset$ . More generally, note that

$$\mathbb{E}\{(\xi_{\mathbf{s},k} - s_k)(\xi_{\mathbf{s},\ell} - s_\ell)\} = \mathbb{E}(\xi_{\mathbf{s},k} \xi_{\mathbf{s},\ell}) - s_k \mathbb{E}(\xi_{\mathbf{s},\ell}) - s_\ell \mathbb{E}(\xi_{\mathbf{s},k}) + s_k s_\ell$$

can be expressed as

$$\begin{aligned} & \frac{(s_k/b + 1) \mathbb{1}_{\{k=\ell\}} - \frac{(s_k/b+1)(s_\ell/b+1)}{(1/b+d+1)}}{(1/b+d+1)(1/b+d+2)} + \frac{s_k/b+1}{1/b+d+1} \times \frac{s_\ell/b+1}{1/b+d+1} \\ & \quad - s_k \frac{s_\ell/b+1}{1/b+d+1} - s_\ell \frac{s_k/b+1}{1/b+d+1} + s_k s_\ell. \end{aligned}$$

The latter expression, say  $C_b$ , is such that

$$C_b = \begin{cases} b(s_k \mathbf{1}_{\{k=\ell\}} - s_i s_\ell) + \mathcal{O}(b^2) & \text{if } s_k \text{ is fixed and } s_\ell \text{ is fixed,} \\ b^2\{1 - s_\ell(\lambda_k + d + 2)\} + \mathcal{O}(b^3) & \text{if } s_k/b \rightarrow \lambda_k \text{ and } s_\ell \text{ is fixed,} \\ b^2\{1 - s_k(\lambda_\ell + d + 2)\} + \mathcal{O}(b^3) & \text{if } s_k \text{ is fixed and } s_\ell/b \rightarrow \lambda_\ell, \\ b^2\{(\lambda_k + 1)\mathbf{1}_{\{k=\ell\}} + 1\} + \mathcal{O}(b^3) & \text{if } s_k/b \rightarrow \lambda_k \text{ and } s_\ell/b \rightarrow \lambda_\ell. \end{cases}$$

Also, if  $\mathcal{J} = [d]$ , then Hölder's inequality implies that, for all  $j, k, \ell \in [d]$ ,

$$\mathbb{E}(|\xi_{s,j} - s_j| |\xi_{s,k} - s_k| |\xi_{s,\ell} - s_\ell|) \leq \max_{1 \leq j \leq d} \{\mathbb{E}(|\xi_{s,j} - s_j|^6)\}^{3/6} = \mathcal{O}\{(b^6)^{3/6}\} = \mathcal{O}(b^3).$$

Hence, for any  $\mathcal{J} \subseteq [d]$ , one can conclude from (7) that

$$\mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) - m(\mathbf{s})\} = b^{1+\mathbf{1}_{\{\mathcal{J}=[d]\}}}\mathbf{h}_{\mathcal{J}}(\mathbf{s}) + \mathcal{O}\{b^{3/2(1+\mathbf{1}_{\{\mathcal{J}=[d]\}})}\} + \mathcal{O}\{n^{-1}A_b(\mathbf{s})\}.$$

To estimate the conditional variance of  $\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})$ , let

$$V = \text{diag}\{\sigma^2(\mathbf{X}_1), \dots, \sigma^2(\mathbf{X}_n)\}.$$

By (2), one has

$$\begin{aligned} \text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\} \\ = n^{-1} \mathbf{e}_1^\top (n^{-1} \mathcal{X}_s^\top W_s \mathcal{X}_s)^{-1} n^{-1} \mathcal{X}_s^\top W_s V W_s \mathcal{X}_s (n^{-1} \mathcal{X}_s^\top W_s \mathcal{X}_s)^{-1} \mathbf{e}_1. \end{aligned} \quad (8)$$

Furthermore, one can write

$$n^{-1} \mathcal{X}_s^\top W_s V W_s \mathcal{X}_s = \begin{bmatrix} G_n & H_n \\ I_n & J_n \end{bmatrix},$$

where

$$G_n = \frac{1}{n} \sum_{i=1}^n \kappa_{s,b}(\mathbf{X}_i)^2 \sigma^2(\mathbf{X}_i), \quad H_n = \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s})^\top \kappa_{s,b}(\mathbf{X}_i)^2 \sigma^2(\mathbf{X}_i),$$

$$I_n = \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s}) \kappa_{s,b}(\mathbf{X}_i)^2 \sigma^2(\mathbf{X}_i), \quad J_n = \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \mathbf{s})(\mathbf{X}_i - \mathbf{s})^\top \kappa_{s,b}(\mathbf{X}_i)^2 \sigma^2(\mathbf{X}_i).$$

Consequently,

$$n^{-1} \mathcal{X}_s^\top W_s V W_s \mathcal{X}_s = \left[ \begin{array}{c|c} n^{-1} A_b(\mathbf{s}) \sigma^2(\mathbf{s}) f(\mathbf{s}) + \text{op}\{n^{-1} A_b(\mathbf{s})\} & \text{op}\{n^{-1} A_b(\mathbf{s})\} \\ \text{op}\{n^{-1} A_b(\mathbf{s})\} & \star \end{array} \right],$$

where the expression replacing the symbol  $\star$  is again irrelevant. Under Assumption (A3), note that, for any given  $\mathbf{s} \in \text{Int}(\mathcal{S}_d)$ , one has  $A_b(\mathbf{s})/n \rightarrow 0$  as  $n \rightarrow \infty$ . Together with (5) and (8), one gets

$$\text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\} = n^{-1} A_b(\mathbf{s}) \sigma^2(\mathbf{s}) / f(\mathbf{s}) + o_{\mathbb{P}}\{n^{-1} A_b(\mathbf{s})\}.$$

It is also possible to show that

$$\text{Var}[\mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\}] = o\{n^{-1} A_b(\mathbf{s})\}.$$

Given that

$$\text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\} = \text{Var}[\mathbb{E}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\}] + \mathbb{E}[\text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s}) \mid \mathbf{X}_1, \dots, \mathbf{X}_n\}],$$

it follows that

$$\text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\} = n^{-1} A_b(\mathbf{s}) \sigma^2(\mathbf{s}) / f(\mathbf{s}) + o\{n^{-1} A_b(\mathbf{s})\}. \quad (9)$$

The claimed expression for the variance follows from Lemma 2 in the Appendix. This concludes the proof of Proposition 1.

## 4.2 Proof of Proposition 2

Turning to the proof of Proposition 2, one can apply the bound and convergence of  $A_b$  in Lemma 2, together with boundedness assumptions on  $f$  and  $\sigma^2$  from Assumption (A2), and the dominated convergence theorem, to deduce from (9) that

$$\begin{aligned} n b^{d/2} \int_{\mathcal{S}_d} \text{Var}\{\hat{m}_{n,b}^{(\text{LL})}(\mathbf{s})\} d\mathbf{s} &= b^{d/2} \int_{\mathcal{S}_d} \frac{A_b(\mathbf{s}) \sigma^2(\mathbf{s})}{f(\mathbf{s})} \{1 + o(1)\} d\mathbf{s} \\ &= \int_{\mathcal{S}_d} \frac{\psi_{\emptyset}(\mathbf{s}) \sigma^2(\mathbf{s})}{f(\mathbf{s})} d\mathbf{s} + o(1). \end{aligned}$$

Similarly, noting that  $m \in C^2(\mathcal{S}_d)$  by Assumption (A1), and  $\mathcal{S}_d$  is compact, one can use the boundedness of the second order partial derivatives of  $m$ , together with the dominated convergence theorem, to deduce from Proposition 1 that

$$b^{-2} \int_{\mathcal{S}_d} \text{Bias}\{\hat{m}_{n,b}(\mathbf{s})\}^2 d\mathbf{s} = \int_{\mathcal{S}_d} h_{\emptyset}^2(\mathbf{s}) d\mathbf{s} + o(1) + \mathcal{O}\{b^{-2}(n^{-1} b^{-d/2})^2\}.$$

By combining the last two equations, one finds that

$$\begin{aligned} \text{MISE}\{\hat{m}_{n,b}^{(\text{LL})}\} &= \int_{\mathcal{S}_d} \text{Var}\{\hat{m}_{n,b}(\mathbf{s})\} d\mathbf{s} + \int_{\mathcal{S}_d} \text{Bias}\{\hat{m}_{n,b}(\mathbf{s})\}^2 d\mathbf{s} \\ &= n^{-1} b^{-d/2} \int_{\mathcal{S}_d} \frac{\psi_{\emptyset}(\mathbf{s}) \sigma^2(\mathbf{s})}{f(\mathbf{s})} d\mathbf{s} + b^2 \int_{\mathcal{S}_d} h_{\emptyset}^2(\mathbf{s}) d\mathbf{s} + o(b^2) + o(n^{-1} b^{-d/2}), \end{aligned}$$

where the last error term results from the fact that  $(n^{-1}b^{-d/2})^2 = o(n^{-1}b^{-d/2})$  by Assumption (A3). This concludes the proof of Proposition 2.

## 5 Simulation study

This section reports the results of a modest simulation study comparing the performance of the LL estimator with Dirichlet kernel,  $\hat{m}_{n,b}^{(LL)}$ , against its only direct competitor in the literature, the Nadaraya–Watson (NW) estimator with Dirichlet kernel introduced recently by Bouzebda et al. (2024), viz.

$$\hat{m}_{n,b}^{(NW)}(\mathbf{s}) = \frac{\sum_{i=1}^n Y_i \kappa_{\mathbf{s},b}(\mathbf{X}_i)}{\sum_{j=1}^n \kappa_{\mathbf{s},b}(\mathbf{X}_j)}.$$

The bandwidth selection procedure considered is least-squares cross validation (LSCV). For a given method in the set {LL, NW}, and a given target regression function  $m$ , the bandwidth  $\hat{b}_n \in (0, \infty)$  is chosen to minimize the criterion

$$\text{LSCV}(b) = \frac{1}{1000} \sum_{i=1}^{1000} |\hat{m}_{n,b}^{(\text{method})}(\mathbf{U}_i) - m(\mathbf{U}_i)|^2/d!,$$

where  $\mathbf{U}_1, \dots, \mathbf{U}_{1000}$  form a random sample from the uniform distribution  $\mathcal{U}(\mathcal{S}_d)$  on the unit simplex,  $\mathcal{S}_d$ . The factor  $1/d!$  in the numerator is the normalization constant for this distribution. For each  $b \in (0, \infty)$ ,  $\text{LSCV}(b)$  is an unbiased estimator of

$$\text{MISE}\{\hat{m}_{n,b}^{(\text{method})}\} \equiv \int_{\mathcal{S}_d} \mathbb{E}\{|\hat{m}_{n,b}^{(\text{method})}(\mathbf{s}) - m(\mathbf{s})|^2\} d\mathbf{s}.$$

Three target regression functions were tested, respectively defined, for all  $\mathbf{s} = (s_1, s_2) \in \mathcal{S}_2$ , by

- (i)  $m_1(\mathbf{s}) = s_1 s_2$ ,
- (ii)  $m_2(\mathbf{s}) = \ln(1 + s_1 + s_2)$ ,
- (iii)  $m_3(\mathbf{s}) = \sin(s_1) + \cos(s_2)$ .

Let  $R = 200$  be the number of Monte Carlo replications, and suppose that the sequence of design points  $\mathbf{x}_1, \dots, \mathbf{x}_n$  consists of the fixed mesh

$$\{(w_k(i-1) + 1/2, w_k(k-j) + 1/2)/(k+1) : 1 \leq i \leq j \leq k\}$$

inside the simplex, for  $w_k = (k - 1/\sqrt{2})/(k - 1)$  and some positive integer  $k$ . For each method in {LL, NW}, each target regression function  $m_j$ ,  $j \in \{1, 2, 3\}$ , and each sample size  $n = k(k+1)/2$  with  $k \in \{7, 10, 14, 20\}$ , the mean, median, standard deviation, and interquartile range, are reported in Table 1 for the sequence of Monte

**Table 1** Comparison of the LL and NW methods in terms of the mean, median, standard deviation (SD), and interquartile range (IR), of 200  $\widehat{\text{ISE}}$  values for regression functions  $m_1, m_2, m_3$  and sample size  $n \in \{28, 55, 105, 210\}$ . The LL method leads to smaller mean, median, SD, and IR in all cases.

Function	$n$	Local Linear (LL)			
		Mean	Median	SD	IR
$m_1$	28	2.80E-05	2.80E-05	1.09E-06	1.61E-06
$m_1$	55	2.62E-05	2.63E-05	7.98E-07	1.10E-06
$m_1$	105	2.84E-05	2.84E-05	6.57E-07	9.24E-07
$m_1$	210	3.48E-05	3.47E-05	8.40E-07	1.18E-06
$m_2$	28	0.000135	0.000135	1.70E-06	2.09E-06
$m_2$	55	0.000226	0.000226	2.01E-06	2.66E-06
$m_2$	105	0.000261	0.000261	2.24E-06	2.93E-06
$m_2$	210	0.000233	0.000233	2.03E-06	2.88E-06
$m_3$	28	0.000579	0.000579	9.13E-06	1.19E-05
$m_3$	55	0.000556	0.000556	8.28E-06	1.01E-05
$m_3$	105	0.000590	0.000590	7.85E-06	1.08E-05
$m_3$	210	0.000616	0.000616	7.79E-06	1.09E-05

Function	$n$	Nadaraya–Watson (NW)			
		Mean	Median	SD	IR
$m_1$	28	0.000130	0.000130	6.60E-06	9.45E-06
$m_1$	55	8.46E-05	8.45E-05	3.50E-06	4.34E-06
$m_1$	105	6.82E-05	6.84E-05	2.44E-06	3.55E-06
$m_1$	210	7.00E-05	7.01E-05	1.64E-06	1.99E-06
$m_2$	28	0.000281	0.000281	1.39E-05	1.98E-05
$m_2$	55	0.000309	0.000308	8.93E-06	1.31E-05
$m_2$	105	0.000323	0.000323	6.53E-06	8.03E-06
$m_2$	210	0.000286	0.000286	4.43E-06	5.83E-06
$m_3$	28	0.001199	0.001193	5.18E-05	7.44E-05
$m_3$	55	0.000948	0.000945	3.11E-05	4.15E-05
$m_3$	105	0.000893	0.000892	1.85E-05	2.31E-05
$m_3$	210	0.000894	0.000895	1.34E-05	1.70E-05

Carlo estimates of the integrated squared errors:

$$\widehat{\text{ISE}}_{\text{method}}^{j,r} = \frac{1}{1000} \sum_{i=1}^{1000} |\hat{m}_{n,\hat{b}_{n,r}}^{(\text{method})}(\mathbf{U}_i) - m_j(\mathbf{U}_i)|^2/2,$$

where  $r \in \{1, \dots, R\}$ . In each case  $(m_j, n)$ , Table 1 shows that the new LL estimator with Dirichlet kernel has the smallest mean, median, standard deviation, and

interquartile range, clearly demonstrating its dominance over the NW estimator with Dirichlet kernel of Bouzebda et al. (2024).

In their paper, Bouzebda et al. (2024) also considered a NW regression estimator where the Dirichlet kernel is replaced by a discrete multinomial kernel, offering a Bernstein smoothing alternative. No advantages were found over the NW estimator with Dirichlet kernel (4), so it was not included in the present study for the sake of concision. The LL method with Dirichlet kernel outperforms both NW estimators by a significant margin. This is unsurprising, however, given that NW estimators can be viewed as local *constant* estimators, where the slope of the corresponding locally weighted least-squares problem is set to zero.

## 6 Data example

As an illustration, the proposed local linear (LL) smoother is used to take yet another look at a classical dataset reported by Aitchison (1986, p. 359) and displayed in Table 2 for convenience. These data, which are also available in the R package `DirichletReg`, are originally due to Coakley and Rust (1968), who provide the composition in terms of sand, silt, and clay of 39 samples of sediment in Stanwell-Fletcher Lake, Somerset Island (Inuktitut Kuuganajuk, Nunavut, Canada) as a function of depth (in meters). One way to assess the relation between water depth and compositional pattern is to determine the extent to which depth can be predicted by composition, which calls for a model of the form (1).

The dataset comprises 39 design points  $\mathbf{x}_i = (x_{i,1}, x_{i,2})$ , which represent the proportions of sand and silt in each sediment sample, respectively. The proportion of clay is determined by the complement,  $1 - x_{i,1} - x_{i,2}$ . Water depth (in meters) of each sediment sample, denoted  $y_i$  is an explanatory variable.

The local linear smoother  $\hat{m}_{n,b}^{(LL)}$ , as defined in Section 2, is employed to estimate the water depth based of sediment composition. The bandwidth  $b$  is selected using the leave-one-out cross-validation method which minimizes the residual sum of squares

$$\text{RSS}(b) = \frac{1}{n} \sum_{i=1}^n \{y_i - \hat{m}_{n,b,(-i)}^{(LL)}(\mathbf{x}_i)\}^2,$$

where  $\hat{m}_{n,b,(-i)}^{(LL)}$  represents the leave-one-out estimator for each  $i \in \{1, \dots, n\}$ , i.e., the local linear smoother defined without the  $i$ th pair  $(\mathbf{x}_i, y_i)$ . Fig. 1 shows the graph of RSS as a function of the bandwidth  $b$ .


Numerical computations reveal that the optimal bandwidth under this criterion is

$$\hat{b} = \operatorname{argmin}_{b \in (0, \infty)} \text{RSS}(b) \approx 0.2195.$$


The 3d plot and density plot of  $\hat{m}_{39, \hat{b}}^{(LL)}$  are illustrated in Fig. 2 and Fig. 3, respectively. These graphs make it clear that sand and silt are present in smaller proportions than clay in sediment composition in deeper areas of the lakebed. In the deepest region (in red), sediments are predominantly clay, with the remaining composition skewing heavily toward sand.

**Table 2** Sand, silt, clay compositions of 39 sediment samples at different water depths in Stanwell-Fletcher Lake, Somerset Island (Nunavut, Canada).


Sediment no.	Percentages			Water depth (meters)	Sediment no.	Percentages			Water depth (meters)
	Sand	Silt	Clay			Sand	Silt	Clay	
1	77.5	19.5	3.0	10.4	21	9.5	53.5	37.0	47.1
2	71.9	24.9	3.2	11.7	22	17.1	48.0	34.9	48.4
3	50.7	36.1	13.2	12.8	23	10.5	55.4	34.1	49.4
4	52.2	40.9	6.6	13.0	24	4.8	54.7	41.0	49.5
5	70.0	26.5	3.5	15.7	25	2.6	45.2	52.2	59.2
6	66.5	32.2	1.3	16.3	26	11.4	52.7	35.9	60.1
7	43.1	55.3	1.6	18.0	27	6.7	46.9	46.4	61.7
8	53.4	36.8	9.8	18.7	28	6.9	49.7	43.4	62.4
9	15.5	54.4	30.1	20.7	29	4.0	44.9	51.1	69.3
10	31.7	41.5	26.8	22.1	30	7.4	51.6	40.9	73.6
11	65.7	27.8	6.5	22.4	31	4.8	49.5	45.7	74.4
12	70.4	29.0	0.6	24.4	32	4.5	48.5	47.0	78.5
13	17.4	53.6	29.0	25.8	33	6.6	52.1	41.3	82.9
14	10.6	69.8	19.6	32.5	34	6.7	47.3	45.9	87.7
15	38.2	43.1	18.7	33.6	35	7.4	45.6	46.9	88.1
16	10.8	52.7	36.5	36.8	36	6.0	48.9	45.1	90.4
17	18.4	50.7	30.9	37.8	37	6.3	53.8	39.9	90.6
18	4.6	47.4	48.0	36.9	38	2.5	48.0	49.5	97.7
19	15.6	50.4	34.0	42.2	39	2.0	47.8	50.2	103.7
20	31.9	45.1	23.0	47.0					



**Fig. 1** Plot of the residual sum of squares as a function of the bandwidth  $b$ .



**Fig. 2** 3d plot of the estimated water depth as a function of the proportion of sand and silt.



**Fig. 3** Density plot of the estimated water depth as a function of the proportion of sand and silt.

## Appendix: Technical results

The first lemma recalls a well-known formula from linear algebra for the inverse of a  $2 \times 2$  block matrix, as detailed, for example, in Section 0.7 of [Horn and Johnson \(2013\)](#).

**Lemma 1.** *Let  $M$  be a matrix of width at least 2 which is partitioned into a  $2 \times 2$  block matrix. If the upper left block  $M_{11}$  is invertible and the Schur complement  $M_\star = M_{22} - M_{21}M_{11}^{-1}M_{12}$  is invertible, then*

$$M = \begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix} \Rightarrow M^{-1} = \begin{bmatrix} M_{11}^{-1} + M_{11}^{-1}M_{12}M_\star^{-1}M_{21}M_{11}^{-1} & -M_{11}^{-1}M_{12}M_\star^{-1} \\ -M_\star^{-1}M_{21}M_{11}^{-1} & M_\star^{-1} \end{bmatrix}.$$

The second lemma, which is Lemma 1 of [Ouimet and Tolosana-Delgado \(2022\)](#), provides a uniform bound on  $A_b(\mathbf{s}) = \mathbf{E}\{\kappa_{\mathbf{s},b}(\mathbf{X})^2\}$  and describes its asymptotic behavior as  $b \rightarrow 0$ , whether the sequence of points  $\mathbf{s} = \mathbf{s}(b)$  is well inside or near the boundary of the simplex  $\mathcal{S}_d$ .

**Lemma 2.** *Uniformly for  $\mathbf{s} \in \mathcal{S}_d$ , one has, as  $b \rightarrow 0$ ,*

$$0 < A_b(\mathbf{s}) \leq \frac{b^{(d+1)/2} (1/b + d)^{d+1/2}}{(4\pi)^{d/2} \sqrt{(1 - \|\mathbf{s}\|_1) \prod_{i \in [d]} s_i}} \{1 + \mathcal{O}(b)\}.$$

Also, for any vector  $\boldsymbol{\lambda} \in (0, \infty)^d$  and any subset  $\mathcal{J} \subseteq [d]$  of indices and corresponding map  $\psi_{\mathcal{J}}$  defined in (3), one has, as  $b \rightarrow 0$ ,

$$A_b(\mathbf{s}) = b^{-d/2} \psi_{\emptyset}(\mathbf{s}) \times \{1 + \mathcal{O}_{\mathbf{s}}(b)\},$$

if  $s_i$  is fixed for all  $i \in [d]$  and  $(1 - \|\mathbf{s}\|_1)$  is fixed, while

$$A_b(\mathbf{s}) = b^{-(d+|\mathcal{J}|)/2} \psi_{\mathcal{J}}(\mathbf{s}) \prod_{i \in \mathcal{J}} \frac{\Gamma(2\lambda_i + 1)}{2^{2\lambda_i + 1} \Gamma^2(\lambda_i + 1)} \times \{1 + \mathcal{O}_{\boldsymbol{\lambda}, \mathbf{s}}(b)\}$$

if  $s_i/b \rightarrow \lambda_i$  for all  $i \in \mathcal{J}$ ,  $s_i$  is fixed for all  $i \in [d] \setminus \mathcal{J}$  and  $(1 - \|\mathbf{s}\|_1)$  is fixed.

## Acknowledgment

Genest's research was funded in part by the Canada Research Chairs Program (Grant no. 950-231937) and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2024-04088). Ouimet's funding was made possible through a contribution to Genest's work from the Trottier Institute for Science and Public Policy.

## References

- Aitchison, J. (1986). *The Statistical Analysis of Compositional Data*. Chapman & Hall, London. ([MR865647](#))
- Aitchison, J., & Lauder, I.J. (1985). Kernel density estimation for compositional data. *J. Roy. Statist. Soc. Ser. C*, *34*(2), 129–137, ([doi:10.2307/2347365](#))
- Aliprantis, C.D., & Border, K.C. (2006). *Infinite Dimensional Analysis, A Hitchhiker's Guide* (Third ed.). Springer, Berlin. ([MR2378491](#))
- Bertin, K., Genest, C., Klutchnikoff, N., Ouimet, F. (2023). Minimax properties of Dirichlet kernel density estimators. *J. Multivariate Anal.*, *195*, Paper No. 105158, 16 pp., ([MR4544604](#))
- Bouzebda, S., Nezzal, A., Elhattab, I. (2024). Asymptotic properties of nonparametric conditional  $U$ -statistics smoothed by asymmetric kernels. *To appear in AIMS Mathematics*, 89 pp.,
- Brown, B.M., & Chen, S.X. (1999). Beta-Bernstein smoothing for regression curves with compact support. *Scand. J. Statist.*, *26*(1), 47–59, ([MR1685301](#))
- Chacón, J.E., & Duong, T. (2018). *Multivariate kernel smoothing and its applications*. CRC Press, Boca Raton, FL. ([MR3822372](#))
- Chen, S.X. (2000). Beta kernel smoothers for regression curves. *Statist. Sinica*, *10*(1), 73–91, ([MR1742101](#))
- Chen, S.X. (2002). Local linear smoothers using asymmetric kernels. *Ann. Inst. Statist. Math.*, *54*(2), 312–323, ([MR1910175](#))
- Cleveland, W.S. (1979). Robust locally weighted regression and smoothing scatterplots. *J. Amer. Statist. Assoc.*, *74*(368), 829–836, ([MR556476](#))
- Coakley, J.P., & Rust, B.R. (1968). Sedimentation in an Arctic lake. *J. Sed. Petrology*, *38*(4), 1290–1300, ([doi:10.1306/74D71B59-2B21-11D7-8648000102C1865D](#))
- Fan, J. (1992). Design-adaptive nonparametric regression. *J. Amer. Statist. Assoc.*, *87*(420), 998–1004, ([MR1209561](#))

- Fan, J. (1993). Local linear regression smoothers and their minimax efficiencies. *Ann. Statist.*, 21(1), 196–216, ([MR1212173](#))
- Fan, J., & Gijbels, I. (1992). Variable bandwidth and local linear regression smoothers. *Ann. Statist.*, 20(4), 2008–2036, ([MR1193323](#))
- Fé, E. (2014). Estimation and inference in regression discontinuity designs with asymmetric kernels. *J. Appl. Stat.*, 41(11), 2406–2417, ([MR3256394](#))
- Funke, B., & Hirukawa, M. (2021). Bias correction for local linear regression estimation using asymmetric kernels via the skewing method. *Econom. Stat.*, 20(C), 109–130, ([MR4302589](#))
- Funke, B., & Hirukawa, M. (2024). On uniform consistency of nonparametric estimators smoothed by the gamma kernel. *Preprint*, 29 pp.,
- Hirukawa, M., Murtazashvili, I., Prokhorov, A. (2022). Uniform convergence rates for nonparametric estimators smoothed by the beta kernel. *Scand. J. Stat.*, 49(3), 1353–1382, ([MR4471289](#))
- Hirukawa, M., Murtazashvili, I., Prokhorov, A. (2023). Yet another look at the omitted variable bias. *Econometric Rev.*, 42(1), 1–27, ([MR4556820](#))
- Horn, R.A., & Johnson, C.R. (2013). *Matrix Analysis* (Second ed.). Cambridge University Press, Cambridge. ([MR2978290](#))
- Ouimet, F. (2021). Asymptotic properties of Bernstein estimators on the simplex. *J. Multivariate Anal.*, 185, Paper No. 104784, 20 pp., ([MR4287788](#))
- Ouimet, F. (2022). On the boundary properties of Bernstein estimators on the simplex. *Open Stat.*, 3, 48–62, ([doi:10.1515/stat-2022-0111](#))
- Ouimet, F., & Tolosana-Delgado, R. (2022). Asymptotic properties of Dirichlet kernel density estimators. *J. Multivariate Anal.*, 187, Paper No. 104832, 25 pp., ([MR4319409](#))

- Ruppert, D., & Wand, M.P. (1994). Multivariate locally weighted least squares regression. *Ann. Statist.*, 22(3), 1346–1370, ([MR1311979](#))
- Shi, J., & Song, W. (2016). Asymptotic results in gamma kernel regression. *Comm. Statist. Theory Methods*, 45(12), 3489–3509, ([MR3494026](#))
- Somé, S.M., & Kokonendji, C.C. (2016). Effects of associated kernels in nonparametric multiple regressions. *J. Stat. Theory Pract.*, 10(2), 456–471, ([MR3499725](#))
- Stone, C.J. (1977). Consistent nonparametric regression. *Ann. Statist.*, 5(4), 595–645, ([MR443204](#))
- Tanabe, K., & Sagae, M. (1992). An exact Cholesky decomposition and the generalized inverse of the variance-covariance matrix of the multinomial distribution, with applications. *J. Roy. Statist. Soc. Ser. B*, 54(1), 211–219, ([MR1157720](#))
- Tenbusch, A. (1994). Two-dimensional Bernstein polynomial density estimators. *Metrika*, 41(3-4), 233–253, ([MR1293514](#))