

# On the Hamiltonian for the Hilbert-Pólya Conjecture

Enderalp Yakaboylu\*  
(Dated: August 28, 2024)

In a recent manuscript [J. Phys. A: Math. Theor. 57 235204 (2024)], a candidate Hamiltonian for the Hilbert-Pólya Conjecture was introduced. Here, we elegantly demonstrate that the corresponding eigenfunctions are square-integrable and, crucially, that the eigenvalues are real. The latter represents a significant step toward proving the Riemann Hypothesis.

The Hilbert-Pólya Conjecture, a compelling idea in mathematics, offers a potential pathway to solving one of the most profound mysteries in number theory: the Riemann Hypothesis (RH). This conjecture proposes that the nontrivial zeros of the Riemann zeta function, which are central to understanding the distribution of prime numbers—the fundamental “quanta” of arithmetic—could be the eigenvalues of a self-adjoint operator. Specifically, if there exists a self-adjoint Hamiltonian  $\hat{H}$  such that the eigenvalues of the operator  $(1/2 + i\hat{H})$  correspond to all nontrivial zeros of the Riemann zeta function, this would provide the elusive proof of the RH by confirming that all these nontrivial zeros lie on the critical line, as the RH suggests. Thus, the Hilbert-Pólya Conjecture not only offers a potential approach to proving the RH through spectral theory but also creates a bridge between number theory and quantum physics.

In a recent article [1], the following Hamiltonian was proposed to address the Hilbert-Pólya Conjecture:

$$\hat{H} = i\hat{N} - i\hat{N}_- - i \sum_{m=0}^{\infty} c_m (\hat{N}_-)^m, \quad (1)$$

acting on the Hilbert space  $L^2[0, \infty)$ . Here, the number operator,

$$\hat{N} = \frac{1}{2} \left( \hat{x} \hat{p}^2 + \hat{p}^2 \hat{x} + \frac{\hat{x}}{2} \right),$$

is defined on the positive half-line, and its eigenvalue equation can be recognized as the well-established Laguerre ODE:  $\langle x | \hat{N} | n \rangle = (n + 1/2) \langle x | n \rangle$ , with the eigenfunctions:  $\langle x | n \rangle = e^{-x/2} L_n(x)$ , where  $L_n(x)$  are the Laguerre polynomials and  $n = 0, 1, \dots, \infty$ . The ladder operators, on the other hand, obey the following algebra:  $[\hat{N}, \hat{N}_{\pm}] = \pm \hat{N}_{\pm}$  and  $[\hat{N}_+, \hat{N}_-] = -2\hat{N}$ , with the actions  $\hat{N}_+ | m \rangle = (m + 1) | m + 1 \rangle$  and  $\hat{N}_- | m \rangle = m | m - 1 \rangle$ .

The coefficients in Eq. (1) correspond to those in the series expansion of  $x(1 + e^{-x})^{-1} = \sum_{m=0}^{\infty} c_m x^m$ , and are given by  $c_m = B_m(2^m - 1)/m!$ , where  $B_m$  are the Bernoulli numbers with the convention  $B_1 = +1/2$ . Note that the sum converges only for  $0 \leq |x| < \pi$ . Nevertheless, the series is Borel summable for all  $\text{Re}(x) \geq 0$  and  $|\text{Im}(x)| < \pi$ ; see the appendix of Ref. [1].

It was shown in Ref. [1] that the eigenfunctions of the proposed Hamiltonian,  $\tilde{\Psi}_s(x)$ , associated with the eigenvalues  $i(1/2 - s)$ , vanish at the origin under the imposed

boundary condition, which is dictated by the vanishing of the Riemann zeta function:

$$\tilde{\Psi}_s(0) = (1 - 2^{1-s})\Gamma(s)\zeta(s) = 0, \quad (2)$$

where  $\Gamma(s)$  is the gamma function, and

$$\zeta(s) = \frac{1}{(1 - 2^{1-s})\Gamma(s)} \int_0^{\infty} du \frac{u^{s-1}}{1 + e^u},$$

is the Riemann zeta function defined for  $\text{Re}(s) > 0$ . As a result, the eigenvalues of the Hamiltonian (1) are given by the nontrivial Riemann zeros.

In this Letter, we present an elegant approach to demonstrating that the eigenvalues of the Hamiltonian (1) are real. To achieve this, we introduce the following similarity transformation of the Hamiltonian (1):

$$\hat{H} = e^{\hat{x}/2} \hat{H} e^{-\hat{x}/2} = -\hat{D} - i \sum_{m=0}^{\infty} c_m (\hat{T})^m, \quad (3)$$

where we identify  $\hat{x} = 2\hat{N} - \hat{N}_+ - \hat{N}_-$ , as discussed in Ref. [1], to utilize the algebra of the ladder operators. We note that the boundary condition (2) remains unchanged after applying the similarity transformation. In the Hamiltonian (3),  $\hat{D} = i(\hat{N}_- - \hat{N}_+)/2 = (\hat{x} \hat{p} + \hat{p} \hat{x})/2$  is the well-known Berry-Keating Hamiltonian [2], and the operator  $\hat{T} = \hat{N} - \hat{x}/4 = (\hat{x} \hat{p}^2 + \hat{p}^2 \hat{x})/2$  can be referred to as the Bessel operator. The eigenvalue equation of the latter corresponds to Bessel’s differential equation:

$$\langle x | \hat{T} | t \rangle = \left( -x \frac{d^2}{dx^2} - \frac{d}{dx} \right) J_0(2\sqrt{xt}) = t J_0(2\sqrt{xt}),$$

where  $\langle x | t \rangle = J_0(2\sqrt{xt})$  is the Bessel function of the first kind with  $t \in \mathbb{R}^+$ . Both operators,  $\hat{D}$  and  $\hat{T}$ , are self-adjoint on the Hilbert space  $L^2[0, \infty)$ .

The eigenvalue equation for the Hamiltonian (3),

$$\hat{H} |\Psi_s\rangle = i(1/2 - s) |\Psi_s\rangle, \quad (4)$$

can be solved by defining  $|\Psi_s\rangle = \int_0^{\infty} dt F_s(t) |t\rangle$ , which corresponds to the Hankel transformation. Consequently, the eigenvalue equation becomes

$$\int_0^{\infty} dt J_0(2\sqrt{xt}) \times \left( -it \frac{d}{dt} - \frac{i}{2} - \frac{it}{1 + e^{-t}} - i \left( \frac{1}{2} - s \right) \right) F_s(t) + it F_s(t) J_0(2\sqrt{xt}) \Big|_{t=0}^{\infty} = 0, \quad (5)$$

where the relation  $xd J_0(2\sqrt{xt})/dx = td J_0(2\sqrt{xt})/dt$  is used. The solution to the differential equation for  $F_s(t)$  in Eq. (5), which ensures the vanishing of the boundary term, is given by

$$F_s(t) = f \frac{t^{s-1}}{1+e^t},$$

where  $f$  is an integration constant. Therefore, the eigenfunctions can be expressed as

$$\langle x|\Psi_s\rangle = \Psi_s(x) = f \int_0^\infty dt \frac{t^{s-1}}{1+e^t} J_0(2\sqrt{xt}). \quad (6)$$

This result is consistent with the eigenfunctions  $\tilde{\Psi}_s(x) = e^{-x/2}\Psi_s(x)$  derived in Ref. [1]. The boundary condition,  $\Psi_s(0) = 0$ , then, identifies the eigenvalues as  $i(1/2 - \rho)$ , where  $\rho$  denotes the nontrivial Riemann zeros. Furthermore, the eigenfunctions are square-integrable:

$$\begin{aligned} \int_0^\infty dx \Psi_\rho^*(x)\Psi_\rho(x) &= |f|^2 \int_0^\infty dt \frac{t^{\rho^*+\rho-2}}{(1+e^t)^2} \\ &= |f|^2 2^{-\rho-\rho^*} \Gamma(1+\rho+\rho^*) \left( (2^{\rho+\rho^*} - 1)\zeta(1+\rho+\rho^*) \right. \\ &\quad \left. - (2^{\rho+\rho^*} - 2)\zeta(\rho+\rho^*) \right), \end{aligned}$$

where we use the identity  $\int_0^\infty dx J_0(2\sqrt{xt})J_0(2\sqrt{xt'}) = \delta(t-t')$ . This important result suggests that there may exist a weight function such that the eigenfunctions are orthonormal and the eigenvalues are real, which supports the validity of the RH.

Rather than searching for the weight function, we aim to demonstrate the reality of the eigenvalues in the following manner. We begin by considering the Hermitian conjugate of the Hamiltonian (3):

$$\hat{H}^\dagger = -\hat{D} + i \sum_{m=0}^\infty c_m \left( \hat{T} \right)^m, \quad (7)$$

which is assumed to satisfy the eigenvalue equation

$$\hat{H}^\dagger |\Phi_s\rangle = i(1/2 - s) |\Phi_s\rangle. \quad (8)$$

Similar to our previous calculation for  $|\Psi_s\rangle$ , we expand the eigenstates in terms of the eigensates of the Bessel operator  $\hat{T}$  as:  $|\Phi_s\rangle = \int_0^\infty dt G_s(t) |t\rangle$ . The corresponding eigenvalue equation can then be written as

$$\begin{aligned} &\int_0^\infty dt J_0(2\sqrt{xt}) \\ &\times \left( -it \frac{d}{dt} - \frac{i}{2} + \frac{it}{1+e^{-t}} - i \left( \frac{1}{2} - s \right) \right) G_s(t) \quad (9) \\ &+ it G_s(t) J_0(2\sqrt{xt}) \Big|_{t=0}^\infty = 0. \end{aligned}$$

It can be shown that this equation can be solved by

$$G_s(t) = g t^{s-1} (1+e^t) \int_0^t d\tau \frac{\tau^{-s}}{1+e^\tau}.$$

To derive this solution, we first set the differential equation for  $G_s(t)$  in Eq. (9) to the constant value  $(-ig)$  and omit the integration constant. We then use the vanishing of the Riemann zeta function at the boundary term and apply the identity  $\int_0^\infty dt J_0(2\sqrt{xt}) = \delta(x)$ . The latter, due to continuity, also implies that  $\Phi_s(0) = 0$  when  $s = \rho$ , thereby identifying the eigenvalues in terms of the nontrivial Riemann zeros.

This analysis reveals that the operators,  $\hat{H}$  and  $\hat{H}^\dagger$  are isospectral, meaning that there exists a well-defined similarity transformation such that  $\hat{H}^\dagger = \hat{W}\hat{H}\hat{W}^{-1}$ . Consequently, there is an associated self-adjoint operator of the form  $\hat{h} = \hat{S}\hat{H}\hat{S}^{-1}$ , where  $\hat{W} = \hat{S}^\dagger\hat{S}$ , corresponding to the weight operator in the generalized eigenvalue problem. More explicitly, the eigenkets of the self-adjoint operator,  $\hat{h}$ , are defined as  $|\psi_\rho\rangle = \hat{S}|\Psi_\rho\rangle$ , and the eigenbras as  $\langle\psi_\rho| = \langle\Phi_\rho|\hat{S}^{-1}$ , implying that  $\langle\psi_\rho|\psi_{\rho'}\rangle = \langle\Phi_\rho|\Psi_{\rho'}\rangle$  should be orthogonal and the eigenvalues should be real. This can be demonstrated explicitly as follows:

$$\langle\Phi_\rho|\left(\hat{H}|\Psi_{\rho'}\right)\rangle = i(1/2 - \rho') \langle\Phi_\rho|\Psi_{\rho'}\rangle, \quad (10a)$$

$$\left(\langle\Phi_\rho|\hat{H}\right)|\Psi_{\rho'}\rangle = -i(1/2 - \rho^*) \langle\Phi_\rho|\Psi_{\rho'}\rangle, \quad (10b)$$

where Eq. (4) is used for the first line, and Eq. (8) for the second. It can be shown that the inner product

$$\langle\Phi_\rho|\Psi_{\rho'}\rangle = (g^*f) \int_0^\infty dt t^{\rho^*+\rho'-2} \int_0^t d\tau \frac{\tau^{-\rho^*}}{1+e^\tau}$$

vanishes when  $\rho \neq \rho'$ . However, for  $\rho = \rho'$ , it is given by

$$\langle\Phi_\rho|\Psi_\rho\rangle = (g^*f) \Re \left( (2^{1-\rho} - 1)\Gamma(\rho) \frac{d\zeta(\rho)}{d\rho} \right),$$

which matches exactly with that derived in Ref. [1]. Finally, by subtracting the two equations in Eq. (10), we show that

$$\Re(\rho) = 1/2. \quad (11)$$

The author acknowledges Michael Berry for valuable discussions.

---

\* yakaboylu@proton.me

- [1] E. Yakaboylu, Journal of Physics A: Mathematical and Theoretical **57**, 235204 (2024).  
 [2] M. V. Berry and J. P. Keating, in *Supersymmetry and Trace Formulae* (Springer, 1999) pp. 355–367.