A MORE UNIFIED THEORY OF TRANSFER LEARNING

Steve Hanneke Purdue University steve.hanneke@gmail.com Samory Kpotufe Columbia University samory@columbia.edu

August 30, 2024

ABSTRACT

We show that some basic *moduli of continuity* δ —which measure how fast target risk decreases as source risk decreases—appear to be at the root of many of the classical *relatedness* measures in transfer learning and related literature. Namely, bounds in terms of δ recover many of the existing bounds in terms of other measures of relatedness—both in regression and classification—and can at times be tighter.

We are particularly interested in general situations where the learner has access to both source data and some or no target data. The unified perspective allowed by the moduli δ allow us to extend many existing notions of relatedness at once to these scenarios involving target data: interestingly, while δ itself might not be efficiently estimated, adaptive procedures exist—based on reductions to *confidence sets*—which can get nearly tight rates in terms of δ with no prior distributional knowledge. Such adaptivity to unknown δ immediately implies adaptivity to many classical relatedness notions, in terms of combined source and target samples' sizes.

Keywords Transfer Learning, Domain adaptation.

1 Introduction

Domain Adaptation or Transfer Learning refer generally to the problem of harnessing data from a source distribution P to improve prediction performance w.r.t. to a *target* distribution Q for which some or no data is available. This problem has been researched over the last few decades with a recent resurgence in interest driven by modern applications that are often characterized by a scarcity of perfect target data.

A fundamental question in the theory of domain adaptation (and variant problems on distribution shifts) is how to measure the relatedness between source P and target Q distributions. Importantly, desired measures of relatedness should not only tightly capture the predictive information P has on Q, but have to be practically useful: that is, either the measure can be estimated from data to facilitate algorithmic design, or more generally, it should somehow admit *adaptive* procedures, i.e., procedures whose performance is adaptive to the a priori unknown level of relatedness between Pand Q. Many notions have been proposed over the last few decades, starting with the seminal works of Mansour et al. [2009], Ben-David et al. [2010] on refinements of *total-variation* for domain adaptation in classification, to more recent proposals for domain adaptation in regression, e.g., Wasserstein distances Redko et al. [2017], Shen et al. [2018], or measures relating covariance structures across P and Q as in Mousavi Kalan et al. [2020], Zhang et al. [2022b], Ge et al. [2023]. These various notions of relatedness appear hard to compare at first glance, leading to a disparate theory of domain adaptation at present with no unified set of principles.

Interestingly as we show, upon closer look at the existing literature—whether in classification or regression—it turns out that in fact, many seemingly distinct measures of relatedness proposed in domain adaptation actually implicitly bound the same fundamental quantities: we refer to these quantities as *weak and strong moduli of transfer*, and they roughly measure how fast the Q-risk of predictors decrease as their P-risk decreases. These moduli always yield as tight or tighter rates of transfer than many existing notions, while also admitting adaptive procedures in general settings, as shown via a reduction to the existence of certain confidence sets for the prediction problem at hand. These reductions, while of a theoretical nature, yield insights on general adaptive transfer approaches that are less tied to specific measures of relatedness between source P and target Q. We now give a more formal description of results. We consider a general prediction setting with joint distributions Pand Q on $\mathcal{X} \times \mathcal{Y}$, and a fixed hypothesis class \mathcal{H} of functions $h : \mathcal{X} \mapsto \mathcal{Y}$. We consider general risks under a given measure μ (here P or Q) of the form $R_{\mu}(h) = \mathbb{E}_{\mu} \ell(h(X), Y)$ for a generic loss ℓ ; of special interest is the *excess risk* $\mathcal{E}_{\mu}(h) \doteq R_{\mu}(h) - \inf_{h' \in \mathcal{H}} R_{\mu}(h')$. Risk minimizers for P and Q need not be the same, and in fact need not exist. We allow a general learning setting with access to labeled samples from P and some or no labeled samples from Q. We remark in particular that many existing results on domain adaptation concern situations with no target data, while here we aim for a general theory.

• The *weak modulus of transfer* is easiest to describe, and happens to be the essential quantity at the root of many existing notions of relatedness. It is given by the following function of $\epsilon > 0$:

$$\delta(\epsilon) = \delta_{P,Q}(\epsilon) \doteq \sup \left\{ \mathcal{E}_Q(h) : h \in \mathcal{H}, \ \mathcal{E}_P(h) \le \epsilon \right\}$$

Roughly, $\delta(\epsilon)$ captures whether good predictors under P remain good under Q; intuitively, if this is the case, data from the source P should be highly informative for prediction under the target Q, and it is therefore not surprising that it is implicitly tied to many existing notions of relatedness (see Section 3.1 for examples such as \mathcal{Y} -discrepancy, transfer-exponent, Wasserstein, covariance ratios, etc.). While $\delta(\epsilon)$ is not easy to estimate from data (as it involves the unknown infimum risks), we show that it nevertheless admits adaptive procedures \hat{h} which, given data from P and some or no data from Q, and with no knowledge of δ can achieve transfer rates of the form (Theorem 1)

$$\mathcal{E}_Q(\hat{h}) \lesssim \min\left\{\epsilon_Q, \delta(\epsilon_P)\right\},\tag{1}$$

where ϵ_{μ} , for μ denoting Q or P, is roughly the best rate achievable under μ in vanilla prediction with the n_{μ} samples from μ alone; for example in classification, typically $\epsilon_u \approx n_{\mu}^{-1/2}$, while in linear regression with squared loss, $\epsilon_u \approx n_{\mu}^{-1}$, ignoring complexity terms such as dimension.

Thus, suppose for instance that the learner only has access to source samples and no target samples, as is commonly studied in the literature. Then the bound of (1) reduces to $\mathcal{E}_Q(\hat{h}) \leq \delta(\epsilon_P)$, which is trivially obtained for instance by ERM \hat{h} on the *P* sample; upper-bounds on $\delta(\epsilon)$ then easily recover many existing results in terms of various notions of relatedness. The weak modulus δ therefore appears as a good starting point towards a more general theory.

We are particularly interested in more general settings allowing some amount of target data. The rate of (1) in this general case may be viewed as interpolating between biasing towards the source data vs biasing towards the target data, whichever is more predictive under Q; this is a priori unknown, since δ is unknown. We show that this is achievable by a general adaptive procedure that reduces to the existence of *weak* confidence sets. These are empirically-induced sets $\hat{\mathcal{H}} \subset \mathcal{H}$ that contain only predictors with $\mathcal{E}_{\mu}(h) \leq \epsilon_{\mu}$, while retaining all near-optimal predictors, i.e., those satisfying $R_{\mu}(h) \approx \inf_{h'} R_{\mu}(h')$. By instantiating these rates (and confidence sets) for traditional regression and classification settings (Section 3.3), we obtain unified rates in terms of both source and target samples that automatically extend to many existing relatedness measures simultaneously.

We show in Theorem 2 via a classification lower-bound that the adaptive transfer rates of (1) cannot be improved without additional structural assumptions. This confirms that the weak modulus tightly captures, at least in a worst-case sense, the limits of transfer with both source and target data.

• The strong modulus of transfer follows as a natural refinement on the weak modulus, and aims to better capture the additional information inherent in having some target data. For intuition, suppose for instance that the source distribution P admits two distinct risk minimizers $h_P^*, h_P^{*'}$, the first with low Q risk, i.e., $\mathcal{E}_Q(h_P^*) \approx 0$, while the second has large $\mathcal{E}_Q(h_P^*') \geq \frac{1}{4}$. For example, h_P^* might focus on a feature (e.g. background color in object classification) that is predictive for data from P, but is irrelevant for prediction under Q. Then, with enough Q data, the distinction in transferability between h_P^* and $h_P^{*'}$ becomes apparent, while the weak-modulus is limited by $h_P^{*'}$. With this intuition in mind, a simplified version of the strong modulus roughly takes the form

$$\delta(\epsilon_1, \epsilon_2) \doteq \sup \{ \mathcal{E}_Q(h) : h \in \mathcal{H}, \mathcal{E}_Q(h) \le \epsilon_1 \text{ and } \mathcal{E}_P(h) \le \epsilon_2 \}$$

for a range of values of ϵ_1, ϵ_2 (see Section 2.2 for an exact definition). Importantly, the principles underlying in the strong modulus, namely that target data can improve the rate of transfer beyond source alone, are new, and not reflected in the weak modulus nor in any of the existing measures from the transfer learning literature.

Adaptation to the strong modulus is more tricky and relies on the existence of *strong confidence sets* (defined in Section 4). This yields adaptive transfer rates of the form $\delta(\epsilon_Q, \epsilon_P)$ always as tight or tighter than the rates of min $\{\epsilon_Q, \delta(\epsilon_P)\}$ achievable under the weak modulus (Corollary 2).

While the analysis presented here remains of a theoretical nature, the resulting design principles have many desirable practical implications: for one, they allow for adaptivity to many measures of relatedness at once, when the source is

informative, while also avoiding *negative transfer* [Zhang et al., 2022a], by automatically biasing towards the target data when the source is uninformative, as evidenced e.g., by the rates of (1). We hope that these design principles may yield insights into more practical procedures, e.g., via efficient approximations of confidence sets—which should be viewed as standing for the learner's ability to *identify and retain good prediction candidates*, given limited information about the strength of the relation between source and target distributions. For instance, it is easily seen that in linear regression with squared loss, our generic approaches result in efficiently solvable convex programs (see Remark 15). This is potentially the case with other tractable losses on favorable domains, which is an interesting direction for future investigation.

2 Preliminaries

2.1 General Setup

Basic Definitions. Let X, Y be jointly distributed according to some measure μ (later P or Q), where X is in some domain \mathcal{X} and $Y \in \mathcal{Y}$. A hypothesis class is a set \mathcal{H} of measurable functions $\mathcal{X} \mapsto \mathcal{Y}$. Given a loss function $\ell : \mathcal{Y}^2 \mapsto \mathbb{R}_+$, we consider risks $R_{\mu}(h) \doteq \mathbb{E}_{\mu} \ell(h(X), Y)$, as measured under a given μ .

Assumption 1 (Finiteness). We assume throughout that $R_{\mu}(h) < \infty, \forall h \in \mathcal{H}$, and for measures μ considered.

Note that the above is a mild assumption, as it does not require the loss to be uniformly bounded. For instance this admits the squared loss in linear regression.

For example, the case of binary classification corresponds to $\mathcal{Y} = \{\pm 1\}$, where we often choose $\ell(y, y') = \mathbb{1}\{y \neq y'\}$, and results below will depend on the VC dimension or Rademacher complexity of the class \mathcal{H} (see, e.g., Vapnik and Chervonenkis [1971], Koltchinskii [2006]). As another example, the case of regression corresponds to $\mathcal{Y} \subset \mathbb{R}$, and we may choose $\ell(y, y') = (y - y')^2$, and results below may depend, for instance, on the covering numbers or pseudo-dimension of the class \mathcal{H} (see, e.g. Anthony and Bartlett [1999]).

Remark 1. Our general results will in fact capture quite abstract dependences on \mathcal{H} , via an abstractly-defined notion of confidence sets, and any notion of complexity that allows for such confidence sets are therefore admissible.

Definition 1. The excess risk w.r.t. a (non-empty) subclass $\mathcal{H}_0 \subset \mathcal{H}$, and a joint distribution μ , is defined as

$$\mathcal{E}_{\mu}(h;\mathcal{H}_0) \doteq R_{\mu}(h) - \inf_{h' \in \mathcal{H}_0} R_{\mu}(h').$$

I particular, we will just refer to $\mathcal{E}_{\mu}(h;\mathcal{H})$ as excess risk, and often write $\mathcal{E}_{\mu}(h)$ in this case for simplicity.

We will need the following useful definition. Let $\overline{\mathbb{R}}_+ = (0, \infty]$ denote the extended positive reals.

Definition 2 (Constraint Set). For any distribution μ , and $\epsilon \in \mathbb{R}_+$, define $\mathcal{H}_{\mu}(\epsilon) \doteq \{h \in \mathcal{H} : \mathcal{E}_{\mu}(h) \le \epsilon\}$.

Clearly, $\mathcal{H}_{\mu}(\infty)$ is just \mathcal{H} , so the inclusion of ∞ is just for convenience as we will see later.

Transfer Setting. We consider *source* and *target* distributions P and Q on (X, Y), where we let $\mathcal{E}_P, \mathcal{E}_Q$ denote excess-risks under P and Q. We are interested in excess risk $\mathcal{E}_Q(\hat{h})$ of classifiers trained jointly on n_P i.i.d samples S_P from P, and n_Q i.i.d. samples S_Q from Q. Which \mathcal{E}_Q is achievable necessarily depends on the *discrepancy* $P \to Q$ appropriately formalized.

2.2 Moduli of Transfer

As we will argue in Section 3, the above simple definition already captures the bulk of notions of discrepancies proposed in the literature on transfer learning.

The first notion considered below serves to capture the reduction in target Q-risk induced by small source P-risk, i.e., by a potentially large amount of P data. In particular, as we will see, ϵ is to stand for the best risk achievable under P given a fixed amount of data from P.

Definition 3 (Weak Modulus). Let $\epsilon > 0$. Define the modulus

$$\delta(\epsilon) \doteq \sup \left\{ \mathcal{E}_Q(h) : h \in \mathcal{H}_P(\epsilon) \right\},\$$

or written as $\delta_{P,Q}(\epsilon)$ when the dependence on distributions is to be made explicit.

The next notion aims to refine the above notion: it serves to capture the reduction in Q-risk, induced by the combined effect of target Q and source P data. As such, in our Q risk bounds, ϵ_1 and ϵ_2 will be instantiated as functions of the amount of Q data and P data, respectively.

Definition 4 (Strong Modulus). For $\epsilon_1, \epsilon_2 > 0$, define the following bivariate modulus:

$$\delta(\epsilon_1, \epsilon_2) \doteq \sup \left\{ \mathcal{E}_Q(h) : h \in \mathcal{H}_Q(\epsilon_1), \quad \mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_1)) \le \epsilon_2 \right\},\$$

or written as $\delta_{P,Q}(\epsilon_1, \epsilon_2)$ when the dependence on distributions is to be made explicit.

We will aim to understand situations in which the strong modulus is a strict refinement over the weak modulus. The following pivotal quantity will turn out useful to our discussions as the above definition of strong modulus can be simplified for values of ϵ_1 above this pivot.

Definition 5 (Pivotal Value). The following quantity captures the infimum excess Q-risk of good classifiers under P:

$$\boldsymbol{\varepsilon}_{\mathbf{P},\mathbf{Q}} \doteq \lim_{\epsilon \to 0} \inf \left\{ \mathcal{E}_{Q}(h) : h \in \mathcal{H}_{P}(\epsilon) \right\}.$$

Remark 2 (Intuition). Consider the simplest case of a finite \mathcal{H} , and let $\mathcal{H}_P^* \doteq \arg\min_{h \in \mathcal{H}} R_P(h)$. We easily see that

$$\boldsymbol{\varepsilon}_{\mathrm{P,Q}} = \min_{h_P^* \in \mathcal{H}_P^*} R_Q(h_P^*) \text{ while } \lim_{\epsilon \to 0} \delta(\epsilon) = \max_{h_P^* \in \mathcal{H}_P^*} R_Q(h_P^*).$$

We note that $\varepsilon_{P,Q}^{\sharp} \doteq \lim_{\epsilon \to 0} \delta(\epsilon)$ is an interesting quantity on its own and will be discussed later in Section 3.1. The above general definitions simply allow us to capture a larger variety of situations including generic losses, and in particular, avoid requiring of the existence of risk minimizers.

Next, we consider an alternative expression of $\varepsilon_{P,Q}$, which will be useful in the analysis. For intuition, again consider the simplest case of a finite \mathcal{H} ; we see that $\varepsilon_{P,Q}$ is the smallest ϵ_1 for which $\mathcal{H}_Q(\epsilon_1)$ contains an h_P^* (whereas $\varepsilon_{P,Q}^{\sharp}$ represents the smallest ϵ_1 for which $\mathcal{H}_Q(\epsilon_1)$ contains every h_P^*). The proposition below generalizes this intuition.

Proposition 1 (Equivalent Form of the Pivotal Value). We have $\varepsilon_{P,Q} = \inf \{ \epsilon_1 \in \mathbb{R}_+ : \forall \epsilon > 0, \mathcal{H}_Q(\epsilon_1) \cap \mathcal{H}_P(\epsilon) \neq \emptyset \}$.

Proof. For ease of notation, let E^{\sharp} denote the set $\{\epsilon_1 \in \mathbb{R}_+ : \forall \epsilon > 0, \mathcal{H}_Q(\epsilon_1) \cap \mathcal{H}_P(\epsilon) \neq \emptyset\}$.

For any $\epsilon_1 > \epsilon_{P,Q}$, we have that $\epsilon_1 > \inf\{\mathcal{E}_Q(h) : h \in \mathcal{H}_P(\epsilon)\}$ for all $\epsilon > 0$, since this last infimum only increases as $\epsilon \to 0$; in other words, for all $\epsilon > 0$, $\mathcal{H}_Q(\epsilon_1) \cap \mathcal{H}_P(\epsilon) \neq \emptyset$, i.e., $\epsilon_1 \in E^{\sharp}$. We therefore have that $\epsilon_{P,Q} \ge \inf E^{\sharp}$. Notice that, this last inequality also holds when $\epsilon_{P,Q} = \infty$.

Now for any $\epsilon_1 \in E^{\sharp}$, by definition, we have that for all $\epsilon > 0$, $\inf_{h \in \mathcal{H}_P(\epsilon)} \mathcal{E}_Q(h) \le \epsilon_1$, in other words, $\epsilon_1 \ge \varepsilon_{PQ}$. We therefore also have that $\varepsilon_{PQ} \le \inf E^{\sharp}$.

Remark 3. We note that we may have $\varepsilon_{P,Q}$ very large, in fact matching $\sup_{a,b\in\mathcal{Y}} \ell(a,b)$ (admitting ∞ for unbounded losses). This describes *P* having little information on *Q*, i.e., where *h*'s with low *P*-risk have large *Q*-risk.

We have the following useful implications.

Corollary 1. Let $\epsilon_1, \epsilon_2 > 0$.

- If $\epsilon_1 > \epsilon_{P,Q}$, every $h \in \mathcal{H}$ has $\mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_1)) = \mathcal{E}_P(h)$. It follows that, for $\epsilon_1 > \epsilon_{P,Q}$ $\delta(\epsilon_1, \epsilon_2) = \sup \{\mathcal{E}_Q(h) : h \in \mathcal{H}_Q(\epsilon_1) \cap \mathcal{H}_P(\epsilon_2)\}.$
- If $\epsilon_1 \leq \varepsilon_{P,Q}$, then $\epsilon_1 \leq \delta(\epsilon_2)$. Equivalently, we always have $\varepsilon_{P,Q} \leq \delta(\epsilon_2)$.

Proof. For the first claim, for $\epsilon_1 > \varepsilon_{P,Q}$, we have $\epsilon_1 \in E^{\sharp}$ (where E^{\sharp} is defined in the proof of Proposition 1). Now by definition, E^{\sharp} is the set of ϵ_1 such that for all $\epsilon, \exists h \in \mathcal{H}_Q(\epsilon_1)$ satisfying $\mathcal{E}_P(h) \leq \epsilon$. Therefore, $\inf_{h \in \mathcal{H}_Q(\epsilon_1)} R_P(h) = \inf_{h \in \mathcal{H}} R_P(h)$, and the claim follows.

For the second claim, we note that $\delta(\epsilon_2) \in E^{\sharp}$, since any $\epsilon > 0$ has $\mathcal{H}_P(\epsilon) \cap \mathcal{H}_P(\epsilon_2) = \mathcal{H}_P(\min\{\epsilon, \epsilon_2\}) \neq \emptyset$, and is a subset of $\mathcal{H}_P(\epsilon_2) \subset \mathcal{H}_Q(\delta(\epsilon_2))$, where the last inclusion is from the definition of $\delta(\epsilon_2)$. Hence $\varepsilon_{PQ} \leq \delta(\epsilon_2)$.

We have the following proposition which simply states that the strong modulus is a refinement of the weak modulus.

(Make clear that Prop 1 serves to arrive at Prop2).

Corollary 2 (Relation Between Strong and Weak Moduli). For $\epsilon_1, \epsilon_2 > 0$,

$$\delta(\epsilon_1, \epsilon_2) \le \min\left\{\epsilon_1, \delta(\epsilon_2)\right\}$$

Proof. If $\epsilon_1 > \epsilon_{P,Q}$, then Corollary 1 implies $\delta(\epsilon_1, \epsilon_2) = \sup \{ \mathcal{E}_Q(h) : h \in \mathcal{H}_Q(\epsilon_1) \cap \mathcal{H}_P(\epsilon_2) \}$, which is bounded by $\min \{ \epsilon_1, \delta(\epsilon_2) \}$ by definition of $\mathcal{H}_Q(\epsilon_1)$ and $\delta(\epsilon_2)$.

If $\epsilon_1 \leq \epsilon_{P,Q}$, then since Corollary 1 implies $\epsilon_1 \leq \delta(\epsilon_2)$ in this case, we have $\min \{\epsilon_1, \delta(\epsilon_2)\} = \epsilon_1 \geq \delta(\epsilon_1, \epsilon_2)$, where the final inequality is immediate from the definition of $\delta(\epsilon_1, \epsilon_2)$.

Finally, we verify that both of these moduli are non-decreasing in either of their arguments. This is immediate by definition for the weak modulus $\delta(\cdot)$, but a bit more involved for the strong modulus $\delta(\cdot, \cdot)$.

Proposition 2. – The weak modulus $\delta(\epsilon)$ is non-decreasing in ϵ , i.e., for all $\epsilon \leq \epsilon'$ it holds that $\delta(\epsilon) \leq \delta(\epsilon')$.

- Also, the strong modulus $\delta(\epsilon_1, \epsilon_2)$ is non-decreasing, i.e., $\forall \epsilon_1 \leq \epsilon'_1, \forall \epsilon_2 \leq \epsilon'_2$ it holds that $\delta(\epsilon_1, \epsilon_2) \leq \delta(\epsilon'_1, \epsilon'_2)$.

Proof. The statement holds trivially for the weak modulus by definition. We therefore focus on the strong modulus.

The fact that $\delta(\epsilon'_1, \epsilon_2) \leq \delta(\epsilon'_1, \epsilon'_2)$ is immediate from the definition. It remains to show $\delta(\epsilon_1, \epsilon_2) \leq \delta(\epsilon'_1, \epsilon_2)$. We first note the main difficulty in this argument arises when $\epsilon_1 \leq \epsilon_{P,Q}$. In particular, the conditions on $\mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_1))$ and $\mathcal{E}_P(h; \mathcal{H}_Q(\epsilon'_1))$ may involve different *P*-infimum errors, hence the resulting sets become hard to compare.

So, first, assume $\inf_{h \in \mathcal{H}_Q(\epsilon_1)} R_P(h) = \inf_{h \in \mathcal{H}_Q(\epsilon_1')} R_P(h)$ (this is the case, e.g., when $\epsilon_1 > \varepsilon_{PQ}$, see Corollary 1). It then follows that the set $\{h \in \mathcal{H}_Q(\epsilon_1) : \mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_1)) \le \epsilon_2\}$ is contained in $\{h \in \mathcal{H}_Q(\epsilon_1') : \mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_1')) \le \epsilon_2\}$, hence $\delta(\epsilon_1, \epsilon_2) \le \delta(\epsilon_1', \epsilon_2)$.

Now, suppose the contrary. Then $\exists h \in \mathcal{H}_Q(\epsilon'_1)$ such that

$$\inf_{h'\in\mathcal{H}_Q(\epsilon_1')} R_P(h') \le R_P(h) < \min\left\{\inf_{h'\in\mathcal{H}_Q(\epsilon_1)} R_P(h'), \inf_{h'\in\mathcal{H}_Q(\epsilon_1')} R_P(h') + \epsilon_2\right\}.$$

By the second inequality, we have that $h \notin \mathcal{H}_Q(\epsilon_1)$, hence

$$\delta(\epsilon_1, \epsilon_2) \le \epsilon_1 \le \mathcal{E}_Q(h) \le \delta(\epsilon_1', \epsilon_2).$$

-		

3 Weak Modulus of Transfer

3.1 Some Existing Discrepancies vs Weak Modulus

Here we consider a few examples of existing notions of discrepancy between source and target P, Q, and illustrate the types of bounds they imply on $\delta(\epsilon)$. These bounds were already implicit in past work, even while the weak modulus $\delta(\epsilon)$ was never explicitly defined as the main object of study, or as the implied notion of discrepancy.

Some Discrepancies in Classification. We first remark that some of the notions below can be stated generally beyond classification, e.g., the \mathcal{Y} -discrepancy and the transfer-exponent, however they usually appear in works on classification. In what follows assume $\ell(a, b) = \mathbb{1}\{a \neq b\}$ and for the simplest case suppose $\mathcal{Y} = \{-1, 1\}$ (though the claims are valid for any \mathcal{Y}).

Example 1 (\mathcal{Y} -discrepancy; Mohri and Medina, 2012). Let $\operatorname{disc}_{\mathcal{Y}}(P,Q) \doteq \sup_{h \in \mathcal{H}} |R_P(h) - R_Q(h)|$. Similar to [Mohri and Medina, 2012]:

$$\begin{aligned} \mathcal{E}_Q(h) &= R_Q(h) - \inf_{h'} R_P(h') + \inf_{h'} R_P(h') - \inf_{h'} R_Q(h') \\ &\leq \mathcal{E}_P(h) + \operatorname{disc}_{\mathcal{Y}}(P,Q) + \inf_{h' \in \mathcal{H}} R_P(h') - \inf_{h' \in \mathcal{H}} R_Q(h'). \end{aligned}$$

In other words, for any $\epsilon \in (0, 1]$,

$$\delta(\epsilon) \le \epsilon + \operatorname{disc}_{\mathcal{Y}}(P, Q) + \left(\inf_{h \in \mathcal{H}} R_P(h) - \inf_{h \in \mathcal{H}} R_Q(h)\right).$$
(2)

In particular, note that the difference in risks above is at most $\operatorname{disc}(P,Q)$, i.e. we also have $\delta(\epsilon) \leq \epsilon + 2\operatorname{disc}_{\mathcal{Y}}(P,Q)$.

As noted by [Mohri and Medina, 2012, Cortes, Mohri, and Medina, 2019], the \mathcal{Y} -discrepancy in fact works for essentially any \mathcal{Y} and loss function ℓ (in particular, they focused on the cases of bounded losses and convex losses).

Remark 4. A variant of the *Y*-discrepancy discussed by Hanneke and Kpotufe [2019] instead defines

$$\operatorname{disc}_{\mathcal{Y}}'(P,Q) = \sup_{h \in \mathcal{H}} |\mathcal{E}_P(h) - \mathcal{E}_Q(h)|.$$

In this case, we directly have: $\mathcal{E}_Q(h) \leq \mathcal{E}_P(h) + \operatorname{disc}'_{\mathcal{Y}}(P,Q)$, so that $\delta(\epsilon) \leq \epsilon + \operatorname{disc}'_{\mathcal{Y}}(P,Q)$.

Example 2 (*A*-discrepancy; Ben-David et al., 2010, Mansour et al., 2009). Consider $\operatorname{disc}_{\mathcal{A}}(P,Q) \doteq \sup_{h,h'\in\mathcal{H}} |P_X(h\neq h') - Q_X(h\neq h')|$. For simplicity, as in [Mansour et al., 2009], we suppose there exist $h_Q^*, h_P^* \in \mathcal{H}$ that minimize the respective risks, and also let $h^* = \operatorname{argmin}_{h\in\mathcal{H}} R_Q(h) + R_P(h)$.

$$R_Q(h) \le Q_X(h \ne h^*) + R_Q(h^*) \le P_X(h \ne h^*) + \operatorname{disc}_{\mathcal{A}}(P,Q) + R_Q(h^*) \\ \le R_P(h) + \operatorname{disc}_{\mathcal{A}}(P,Q) + R_Q(h^*) + R_P(h^*)$$

so that

$$\mathcal{E}_Q(h) \le \mathcal{E}_P(h) + \operatorname{disc}_{\mathcal{A}}(P,Q) + (R_P(h_P^*) - R_Q(h_Q^*)) + R_Q(h^*) + R_P(h^*).$$

In particular, we have for any $\epsilon \in (0, 1]$,

 $\delta(\epsilon) \le \epsilon + \operatorname{disc}_{\mathcal{A}}(P,Q) + (R_P(h_P^*) - R_Q(h_Q^*)) + R_Q(h^*) + R_P(h^*).$ (3)

Example 3 (Transfer Exponent; Hanneke and Kpotufe, 2019, Hanneke, Kpotufe, and Mahdaviyeh, 2023). Supposing there exist $h_Q^*, h_P^* \in \mathcal{H}$ with $\mathcal{E}_Q(h_Q^*) = 0$ and $\mathcal{E}_P(h_P^*) = 0$, a value $\rho \ge 0$ is called a *transfer exponent* if there exists $h_P^* \in \operatorname{argmin}_h R_P(h)$ such that $R_Q(h) - R_Q(h_P^*) \le C_\rho \cdot \mathcal{E}_P^{1/\rho}(h)$, for some C_ρ , and for all $h \in \mathcal{H}$ (equivalently $\forall h \in \mathcal{H}_P(\epsilon_0)$, for some $\epsilon_0 > 0$). It's then immediate that

$$\mathcal{E}_Q(h) = R_Q(h) - R_Q(h_P^*) + \mathcal{E}_Q(h_P^*) \le C_\rho \cdot \mathcal{E}_P^{1/\rho}(h) + \mathcal{E}_Q(h_P^*),$$

implying that, for all $\epsilon \in (0, 1], \delta(\epsilon) \leq C_{\rho} \cdot \epsilon^{1/\rho} + \mathcal{E}_Q(h_P^*).$

A more general definition: we can extend the above definition to the general case with or without *P*-risk minimizers. Define $\varepsilon_{PQ}^{\sharp} \doteq \lim_{\epsilon \to 0} \delta(\epsilon)$. A transfer exponent (C_{ρ}, ρ) is defined as satisfying the condition: $\forall h \in \mathcal{H}, \mathcal{E}_Q(h) \leq C_{\rho} \cdot \mathcal{E}_{P}^{1/\rho}(h) + \varepsilon_{PQ}^{\sharp}$. This immediately implies the following bound:

$$\delta(\epsilon) \le C_{\rho} \cdot \epsilon^{1/\rho} + \varepsilon_{\rm PO}^{\sharp}.$$

Remark 5. These various notions appearing in the literature on classification are not directly comparable, but as we see here, many offer upper-bounds on $\delta(\epsilon)$ with varying degrees of tightness in various situations. *Thus, any transfer rate in terms of* $\delta(\epsilon)$ *immediately yields a bound in terms of these existing notions.* As it turns out, our bounds in terms of $\delta(\epsilon)$ recover many proved so far in terms of these various quantities, and can often be tighter (Theorem 1).

The transfer exponent in particular attempts to capture some desired properties of $\delta(\epsilon)$. For one, it is *asymmetric* just as δ is, and thus does not deteriorate in situations where P has much information on Q, but Q has little information on P—for instance P_X covers the decision boundary under Q but not the other way around. Second, the resulting upperbound yields the same limit $\varepsilon_{PQ}^{\sharp}$ as $\delta(\epsilon)$, i.e., it more adequately captures expected performance under large source samples. However, the polynomial form of the transfer-exponent is arbitrary, and simply allows for easier analysis.

Given the arbitrariness of the polynomial form, $\delta(\epsilon)$ can sometimes be smaller than its upper bound based on the transfer exponent. Similarly, the \mathcal{Y} -discrepancy and \mathcal{A} -discrepancy can be loose: suppose e.g., that $\varepsilon_{P,Q}^{\sharp} = 0$ (see e.g., Remark 2 for intuition), while these discrepancies are not 0, implying that the relevant upper-bounds asymptote at non-zero constants and are therefore loose for small ϵ .

Some Discrepancies in Regression. We note that, while we now focus on regression, i.e., h has continous output, the discussion below may be extended to classification in the usual way by viewing $\ell(a, b)$ as a *surrogate loss* (and the classifier as sign(h) for example).

We start with Wasserstein distance which has recently been popular in Machine Learning, and in particular for domain adaptation Redko et al. [2017], Shen et al. [2018].

Example 4 (Wasserstein 1). Let \mathcal{L} denote the set of 1-Lipschitz functions on \mathcal{X} w.r.t a metric ρ . Then consider the *integral probability metric* $W_1(P,Q) \doteq \sup_{f \in \mathcal{L}} |\mathbb{E}_{P_X}(f) - \mathbb{E}_{Q_X}(f)|$.

Suppose $\mathcal{H} \subset \lambda \cdot \mathcal{L}, \lambda > 0$, and is a set of bounded functions $\mathcal{X} \to [-M, M]$, and let $\ell(a, b) = (a - b)^2$ be the squared loss. We consider a weak covariate shift scenario, where $\mathbb{E}_P[Y|X] = \mathbb{E}_Q[Y|X] = h^*(X)$ for some $h^* \in \mathcal{H}$. Notice

that $u \mapsto u^2$ is 4M-Lipshitz on [-2M, 2M], while for any $h, h' \in \mathcal{H}$, (h - h') is 2λ -Lipschitz on \mathcal{X} , and has range within [-2M, 2M]. Then for any $h \in \mathcal{H}$, we have that $x \mapsto (h(x) - h^*(x))^2$ is $(8M\lambda)$ -Lipschitz. Hence,

$$\mathcal{E}_Q(h) = \mathop{\mathbb{E}}_Q(h(X) - h^*(X))^2 \le \mathop{\mathbb{E}}_P(h(X) - h^*(X))^2 + 8M\lambda \cdot W_1(P,Q) = \mathcal{E}_P(h) + 8M\lambda \cdot W_1(P,Q).$$

Thus, $\forall \epsilon > 0$,

$$\delta(\epsilon) \le \epsilon + 8M\lambda \cdot W_1(P,Q). \tag{4}$$

We note that Shen et al. [2018] show a similar bound as above¹, but rather than excess risk, are instead interested in relating Q-risk to P-risk (for L_1 loss). These can be transformed into excess risk bounds, however with additional terms of the form of (3), which is unsurprising as W_1 may be viewed as extending A-discrepancy to any class H of Lipschitz functions). In fact, notice that the derivations above are similar to those for the Y-discrepancy and A-discrepancy.

Similarly, certain variants of the MMD distance [Huang et al., 2007], another so-called *integral probability metric*, may be viewed as special cases of \mathcal{Y} -discrepancy or extensions of the \mathcal{A} -discrepancy to the regression setting; see e.g., Redko et al. [2017]. The implied bounds on $\delta(\epsilon)$ are analogous to those given in Examples 1 and 2 above.

Remark 6. We emphasize that $\delta(\epsilon)$ may be strictly smaller than the above bounds since (4) remains of a worst-case nature over all Lipchitz functions rather than localized to the optimal functions in the class.

Example 5 (Linear Regression). Consider $\mathcal{X} \subset \mathbb{R}^d$ and $\mathcal{Y} \subset \mathbb{R}$, and write $\Sigma_P \doteq \mathbb{E}_P X X^\top$ and $\Sigma_Q \doteq \mathbb{E}_Q X X^\top$. Assume that Σ_P is full rank. Consider a weak covariate-shift scenario where $\mathbb{E}_P[Y|X] = \mathbb{E}_Q[Y|X] = w_*^\top X$ (this would be the case under covariate-shift where $P_{Y|X} = Q_{Y|X}$). Again we consider the squared loss $\ell(a, b) = (a - b)^2$. We then have that, for any $h(x) \doteq w^\top x$:

$$\frac{\mathcal{E}_Q(h)}{\mathcal{E}_P(h)} = \frac{(w - w_*)^\top \Sigma_Q(w - w_*)}{(w - w_*)^\top \Sigma_P(w - w_*)} \le \sup_v \frac{v^\top \Sigma_Q v}{v^\top \Sigma_P v} = \sup_{v = \Sigma_P^{-\frac{1}{2}} u} \frac{u^\top \Sigma_P^{-\frac{1}{2}} \cdot \Sigma_Q \cdot \Sigma_P^{-\frac{1}{2}} u}{u^\top u} = \lambda_{\max} \left(\Sigma_P^{-1} \Sigma_Q \right),$$

where λ_{max} denotes the largest eigenvalue, and the last equality used the fact that the matrices are *similar* (see Theorem 1.3.20 of Horn and Johnson [1985]). Thus, for any $\epsilon > 0$,

$$\delta(\epsilon) \le \lambda_{\max} \left(\Sigma_P^{-1} \Sigma_Q \right) \cdot \epsilon. \tag{5}$$

Now, outside of covariate-shift, when $w_* = w_P^* \neq w_Q^*$, by the same argument as above, we have that

$$\mathcal{E}_{Q}(h) \leq 2 \|w - w_{P}^{*}\|_{\Sigma_{Q}}^{2} + 2 \|w_{P}^{*} - w_{Q}^{*}\|_{\Sigma_{Q}}^{2} \leq 2\lambda_{\max} \left(\Sigma_{P}^{-1}\Sigma_{Q}\right) \cdot \|w - w_{P}^{*}\|_{\Sigma_{P}}^{2} + 2\mathcal{E}_{Q}(w_{P}^{*})$$

(where $||v||_{\Sigma}^2 \doteq v^{\top} \Sigma v$) and we therefore have

$$\delta(\epsilon) \le 2\lambda_{\max} \left(\Sigma_P^{-1} \Sigma_Q \right) \cdot \epsilon + 2\mathcal{E}_Q(w_P^*).$$
(6)

Upper-bounds similar to the above have been shown in various recent results of Mousavi Kalan et al. [2020], Zhang et al. [2022b], Ge et al. [2023]. We note that Zhang et al. [2022b] consider more general cases with different risk minimizers under P and Q, along with refinements on $\lambda_{\max} (\Sigma_P^{-1} \Sigma_Q)$ in their upper-bounds.

3.2 Adaptive Transfer Upper-Bounds

We start with the following short notation for empirical risk minimizers over source or target samples S_P or S_Q .

Definition 6. Let μ denote either P or Q in what follows, and let $\hat{R}_{\mu}(h) \doteq \frac{1}{n_{\mu}} \sum_{(x,y) \in S_{\mu}} \ell(h(x), y)$ denote empirical risk over S_{μ} . For any $\mathcal{H}' \subset \mathcal{H}$, we let $\hat{h}_{\mu}(\mathcal{H}')$ denote the function returned by an **empirical risk minimization** function ERM_{μ}(\mathcal{H}'): namely, a function mapping the data set S_{μ} to an $h \in \mathcal{H}'$ satisfying $\hat{R}_{\mu}(h) < \inf_{h' \in \mathcal{H}'} \hat{R}_{\mu}(h') + e^{-n_{\mu}}$. In particular, when $\mathcal{H}' = \mathcal{H}$, we simply write ERM_{μ} = ERM_{μ}(\mathcal{H}), and $\hat{h}_{\mu} \doteq \hat{h}_{\mu}(\mathcal{H})$.

Remark 7. The above definition of ERM_{μ} is to allow for a generality of losses where the minimizer of \hat{R}_{μ} may not be achieved in \mathcal{H} . However, in commonly discussed settings such as classification with 0-1 loss, or linear regression with squared loss, we can let $\text{ERM}_{\mu}(\mathcal{H})$ denote any $h \in \mathcal{H}$ achieving $\inf_{h' \in \mathcal{H}} \hat{R}_{\mu}(h')$. Furthermore, for 0-1 loss, such a minimizer exists for any $\mathcal{H}' \subset \mathcal{H}$ since \hat{R}_{μ} can only take a finite number of values.

¹Their bound is on L_1 loss but the arguments are essentially the same

Definition 7. Let μ denote either P or Q. Let $0 < \tau \leq 1$ and $\epsilon > 0$. We call a random set $\hat{\mathcal{H}}_{\mu} \doteq \hat{\mathcal{H}}_{\mu}(S_{\mu})$ an (ϵ, τ) -weak confidence set (under μ) if the following conditions are met with probability at least $1 - \tau$:

(i) $\exists \hat{\epsilon}_{\mu} > \mathcal{E}_{\mu}(\hat{h}_{\mu})$ such that $\mathcal{H}_{\mu}(\hat{\epsilon}_{\mu}) \subset \hat{\mathcal{H}}_{\mu}$, and (ii) $\hat{\mathcal{H}}_{\mu} \subset \mathcal{H}_{\mu}(\epsilon)$.

Notice that, by definition, if an (ϵ, τ) -weak confidence set exists, then it necessarily follows that with probability at least $1 - \tau$, $\mathcal{E}_{\mu}(\hat{h}_{\mu}) \leq \epsilon$: that is, ϵ is a confidence upper bound on the excess risk of ERM.

We give a well-known construction of such weak confidence sets in Section 3.3

In the following algorithm, the intent is that the input sets $\hat{\mathcal{H}}_Q$, $\hat{\mathcal{H}}_P$ are weak confidence sets based on the Q-data S_Q and P-data S_P , respectively. Relevant confidence parameters are stated in the result below.

Algorithm 1. Input: $\hat{\mathcal{H}}_Q$, $\hat{\mathcal{H}}_P \subset \mathcal{H}$, and $\hat{h}_Q = \text{ERM}_Q$

If
$$\hat{\mathcal{H}}_Q \cap \hat{\mathcal{H}}_P \neq \emptyset$$
, return any $h \in \hat{\mathcal{H}}_Q \cap \hat{\mathcal{H}}_P$, otherwise return \hat{h}_Q .

The following result easily follows.

Theorem 1. Let $0 < \tau \leq 1$, and suppose $\hat{\mathcal{H}}_Q$ is an (ϵ_Q, τ) -weak confidence set under Q, and $\hat{\mathcal{H}}_P$ is an (ϵ_P, τ) -weak confidence set under P, for some $\epsilon_Q, \epsilon_P > 0$ depending respectively on n_Q and n_P . Let \hat{h} denote the hypothesis returned by Algorithm 1. We then have that, with probability at least $1 - 2\tau$:

$$\mathcal{E}_Q(h) \le \min\left\{\epsilon_Q, \delta(\epsilon_P)\right\}$$

Proof. Assume the events of Definition 7 hold for $\hat{\mathcal{H}}_Q$ and $\hat{\mathcal{H}}_P$ (with probability at least $1 - 2\tau$). By Definition 7, condition (ii), we have that any $h \in \hat{\mathcal{H}}_Q \cap \hat{\mathcal{H}}_P$ is in $\mathcal{H}_Q(\epsilon_Q) \cap \mathcal{H}_P(\epsilon_P)$ implying $\mathcal{E}_Q(h) \leq \min \{\epsilon_Q, \delta(\epsilon_P)\}$.

So suppose $\hat{\mathcal{H}}_Q \cap \hat{\mathcal{H}}_P = \emptyset$. By both conditions of Definition 7, we have that $\hat{h}_Q \in \hat{\mathcal{H}}_Q$ and satisfies $\mathcal{E}_Q(\hat{h}_Q) \leq \epsilon_Q$. Now, let $\hat{\epsilon}_P$ and $\hat{\epsilon}_Q$ as in Definition 7. Consider any $h \in \mathcal{H}_P(\hat{\epsilon}_P)$. By condition (i) we have that $h \in \hat{\mathcal{H}}_P$ so is therefore not in $\hat{\mathcal{H}}_Q$, which contains $\mathcal{H}_Q(\hat{\epsilon}_Q)$. Consequently, we have that $\mathcal{E}_Q(\hat{h}_Q) < \hat{\epsilon}_Q < \mathcal{E}_Q(h) \leq \delta(\epsilon_P)$ where the last inequality follows from that $h \in \mathcal{H}_P(\hat{\epsilon}_P) \subset \mathcal{H}_P(\epsilon_P)$.

Remark 8 (Implications and Relation to Existing Bounds). We will see in Section 3.1 below, upon instantiations of weak-confidence sets, that the above Theorem 1 recovers many of the bounds in the literature in terms of relatedness measures, and further extend them to settings with both source and target data. In particular, for the basic setting with no target data (i.e., where $n_Q = 0$), taking $\hat{\mathcal{H}}_Q = \mathcal{H}$, the bound effectively becomes $\delta(\epsilon_P)$ and holds for any choice of \hat{h} in $\hat{\mathcal{H}}_P$: in particular, simply taking $\hat{h} = \hat{h}_P$ (the ERM under P) would suffice.

For instance in classification, classical VC bounds give us $\epsilon_P \approx \sqrt{(d_H + \log(1/\tau))/n_P}$. Thus, combining with the upper-bounds of Section 3.1 on $\delta(\epsilon_P)$ we recover many classical results: for instance, (2) yields the *Q*-risk bound which can be derived from the analysis of ERM by Mohri and Medina [2012] (specifically, from equation 7 therein),² while (3) recovers the *Q*-risk results of Ben-David, Blitzer, Crammer, Kulesza, Pereira, and Vaughan [2010].

Furthermore, as discussed above for regression, integral probability metrics such as W_1 and MMD can be used to extend \mathcal{A} -divergence to regression (under e.g. L_1 and L_2 loss) for specific classes \mathcal{H} (see e.g. Redko et al. [2017], Shen et al. [2018]) resulting in bounds on $\delta(\epsilon_P)$ of similar form as (3). The upper-bound of (4) for W_1 holds under the additional assumptions we stated here.

In the case of linear regression, the bounds of (5) and (6) recover similar bounds as in the literature (see, e.g. Zhang et al. [2022b], Ge et al. [2023]). In particular, under covariate-shift, the bound of (5) on $\delta(\epsilon_P)$ matches that of Ge et al. [2023] stated under additional assumptions³.

Finally, we note that the bound $\delta(\epsilon_P)$ derived from Theorem 1 itself can be faster whenever the relevant upper-bounds of Section 3.1 are loose (see Remarks 5 and 6).

²The original bound of Mohri and Medina [2012] is expressed for a drift scenario. It can be specialized to our setting by taking all but the final distribution to be P and the final distribution to be Q. They also express their result in terms of the Rademacher complexity, which is upper bounded by the square-root bound we have presented here. We could instead use the empirical Rademacher complexity to define the weak confidence set, and recover an essentially similar result.

³Their result for linear regression is stated under specific distributional assumptions but specializes a more general bound on ratios of Fisher-Information matrices.

3.3 Examples of Weak Confidence Sets

Throughout this section, we let μ generically denote P or Q, and recall the ERM \hat{h}_{μ} . Also, we define *excess empirical* risk as $\hat{\mathcal{E}}_{\mu}(h) \doteq \hat{R}_{\mu}(h) - \inf_{h' \in \mathcal{H}} \hat{R}_{\mu}(h')$.

3.3.1 Classification with 0-1 loss.

We first remark that (ϵ, τ) -weak confidence, for $\epsilon = O(n_{\mu}^{-1/2})$ are easy to obtain for VC classes \mathcal{H} . In fact, from Proposition 6 of the later Section 4.1, a simple set of the form $\hat{\mathcal{H}} \doteq \left\{ h \in \mathcal{H} : \hat{\mathcal{E}}_{\mu}(h) \lesssim n_{\mu}^{-1/2} \right\}$ is a *strong-confidence* set (see Definition 10).

This section is dedicated to obtaining as tight a weak-confidence set as possible, while we remark that

Assume here that $\mathcal{Y} = \{-1, 1\}$ and $\ell(y, y') = \mathbb{1}\{y \neq y'\}$. In what follows, we use the short-hand notation $\mu(h \neq h') \doteq \mu_X (\{x \in \mathcal{X} : h(x) \neq h'(x)\})$, and for $\mathcal{H}' \subset \mathcal{H}$, define

$$\operatorname{diam}_{\mu}(\mathcal{H}') \doteq \sup_{h,h' \in \mathcal{H}'} \mu(h \neq h').$$

We adopt the following classical noise conditions (see e.g. [Mammen and Tsybakov, 1999, Audibert and Tsybakov, 2007, Massart and Nédélec, 2006, Koltchinskii, 2006, Bartlett et al., 2006]).

Definition 8 (BCC). We say that \mathcal{H} satisfies a **Bernstein Class Condition** (BCC), as measured under μ , with parameters $(C_{\mu}, \beta_{\mu}), C_{\mu} > 0$ and $\beta_{\mu} \in [0, 1]$, if $\forall 0 < \epsilon < 1/2$,

$$\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(\epsilon)) \leq C_{\mu} \cdot \epsilon^{\beta_{\mu}}.$$

Note that the condition trivially holds for $\beta_{\mu} = 0$, $C_{\mu} = 1$. The condition captures the hardness of the learning problem: when $\beta_{\mu} = 1$, which formalizes *low noise* regimes, we expect fast rates of the form n^{-1} , in terms of sample size *n*, while for $\beta_{\mu} = 0$, rates are of the more common form $n^{-1/2}$.

When h^* is not unique, BCC remains well defined (i.e., the definition is invariant to the choice of h^*), as it imposes (when $\beta_{\mu} > 0$) that all h^* 's differ on a set of measure 0 under the data distribution.

The following is a well-known construction of such a weak confidence set, designed to be adaptive to the noise parameters (C_{μ}, β_{μ}) .

Proposition 3. For μ either P or Q, consider the following definition of $\hat{\mathcal{H}}_{\mu}$. Define

$$\alpha_{\mu}(\tau) \doteq \frac{d_{\mathcal{H}} \log(n_{\mu}/d_{\mathcal{H}}) + \log(1/\tau)}{n_{\mu}},$$

where $d_{\mathcal{H}}$ denotes the VC dimension of \mathcal{H} [Vapnik and Chervonenkis, 1971]. For any $h, h' \in \mathcal{H}$, define the empirical distance $\hat{\mu}(h \neq h') = \frac{1}{n_{\mu}} \sum_{(x,y) \in S_{\mu}} \mathbb{1}\{h(x) \neq h'(x)\}$. There exist universal constants C, C' such that the following

$$\hat{\mathcal{H}}_{\mu} \doteq \left\{ h \in \mathcal{H} : \hat{\mathcal{E}}_{\mu}(h) \le C \sqrt{\hat{\mu}(h \neq \hat{h}_{\mu}) \cdot \alpha_{\mu}(\tau)} + C \cdot \alpha_{\mu}(\tau) \right\}$$

is an (ϵ_{μ}, τ) -weak confidence set, for $\epsilon_{\mu} = C' \left(C_{\mu} \cdot \alpha_{\mu}(\tau) \right)^{\frac{1}{2-\beta_{\mu}}}$.

The proof is based on the following lemma, which is a corollary of results of [Vapnik and Chervonenkis, 1974] (see [Hanneke and Kpotufe, 2022] for a formal proof).

Lemma 1 (Uniform Bernstein inequality). For μ either P or Q, there exists a universal constant $C_0 > 1$ such that, with probability at least $1 - \tau$, every $h, h' \in \mathcal{H}$ satisfy

$$\left| (R_{\mu}(h) - R_{\mu}(h')) - (\hat{R}_{\mu}(h) - \hat{R}_{\mu}(h')) \right| \le C_0 \sqrt{\hat{\mu}(h \neq h') \cdot \alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau)$$
(7)

and

$$\frac{1}{2}\mu(h \neq h') - C_0 \cdot \alpha_\mu(\tau) \le \hat{\mu}(h \neq h') \le 2\mu(h \neq h') + C_0 \cdot \alpha_\mu(\tau).$$
(8)

We now present the proof of Proposition 3.

Proof. (Proposition 3) In Lemma 2 of Hanneke and Kpotufe [2022], property (ii) of Definition 7 is established, on the event of probability at least $1 - \tau$ from Lemma 1. Moreover, note that on this same event from Lemma 1, for $\hat{\epsilon}_{\mu} = \mathcal{E}_{\mu}(\hat{h}_{\mu}) + \alpha_{\mu}(\tau)$, any $h \in \mathcal{H}_{\mu}(\hat{\epsilon}_{\mu})$ has

$$\hat{\mathcal{E}}_{\mu}(h) \le R_{\mu}(h) - R_{\mu}(\hat{h}_{\mu}) + C_0 \sqrt{\hat{\mu}(h \neq \hat{h}_{\mu}) \cdot \alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau) \le C_0 \sqrt{\hat{\mu}(h \neq \hat{h}_{\mu}) \cdot \alpha_{\mu}(\tau)} + (C_0 + 1) \cdot \alpha_{\mu}(\tau),$$

where the second inequality is due to $h \in \mathcal{H}_{\mu}(\hat{\epsilon}_{\mu})$, implying $R_{\mu}(h) \leq R_{\mu}(\hat{h}_{\mu})$. Therefore, any such h is also in $\hat{\mathcal{H}}_{\mu}$ (for $C \geq C_0 + 1$ in the definition of $\hat{\mathcal{H}}_{\mu}$), and hence property (i) of Definition 7 is also satisfied on this event.

Remark 9 (New Implications for Existing Relatedness Measures). Theorem 1, together with weak confidence sets instantiated as in Proposition 3 above, imply a transfer rate (hiding logarithmic terms for simplicity)

$$\mathcal{E}_Q(\hat{h}) \lesssim \min\left\{ \left(\frac{d_{\mathcal{H}}}{n_Q}\right)^{\frac{1}{2-\beta_Q}}, \delta\left(\left(\frac{d_{\mathcal{H}}}{n_P}\right)^{\frac{1}{2-\beta_P}}\right) \right\}.$$

In particular, from this we can recover (and slightly strengthen) the results of Hanneke and Kpotufe [2019] in terms of the *transfer exponent*, when combined with the bound on $\delta(\epsilon_P)$ expressed in Example 3: that is, we obtain

$$\mathcal{E}_{Q}(\hat{h}) \lesssim \min\left\{ \left(\frac{d_{\mathcal{H}}}{n_{Q}}\right)^{\frac{1}{2-\beta_{Q}}}, \left(\frac{d_{\mathcal{H}}}{n_{P}}\right)^{\frac{1}{p(2-\beta_{P})}} + \varepsilon_{P,Q}^{\sharp} \right\}$$

where the term $\varepsilon_{P,Q}^{\sharp}$ in fact has a multiplicative factor exactly 1. As discussed above (see Remark 5), the bound in terms of δ can in fact be much tighter whenever the transfer-exponent upper-bound on $\delta\left((d_{\mathcal{H}}/n_P)^{\frac{1}{2-\beta_P}}\right)$ is loose.

The weak confidence set instantiation from Proposition 3 moreover allows us to refine the results of \mathcal{Y} -discrepancy and \mathcal{A} -discrepancy discussed in Remark 8 to account for more favorable BCC parameters β_P , β_Q , while extending some of these existing results to the setting with both source and target data.

3.3.2 Regression with Squared Loss.

We consider linear regression where $\mathcal{H} \doteq \{x \mapsto h_w(x) \doteq w^\top x : w \in \mathbb{R}^d\}$ and the squared loss $\ell(y, y') \doteq (y - y')^2$. Define $\Sigma \doteq \mathbb{E}_{\mu} X X^\top$, assumed invertible. Also define $\hat{\Sigma} \doteq \mathbb{E}_{\hat{\mu}} X X^\top$, where $\hat{\mu}$ is the empirical version of μ .

It shown in Section 4.1.2 that, under common regularity conditions, a set of the form

$$\hat{\mathcal{H}}_{\mu} \doteq \left\{ h_w : (w - \hat{w})^\top \hat{\Sigma} (w - \hat{w}) \lesssim \frac{d + \log(1/\tau)}{n_{\mu}} \right\}$$

is a strong confidence set (see Definition 10), implying it is also a (ϵ, τ) -weak confidence set, for $\epsilon \approx \frac{d + \log(1/\tau)}{n_u}$.

Remark 10. Combining the above with Theorem 1, and the upper-bound of (6) on δ , yields a transfer rate of the form

$$\mathcal{E}_Q(\hat{h}) \lesssim \min\left\{\frac{d}{n_Q}, \lambda_{\max}\left(\Sigma_P^{-1}\Sigma_Q\right) \cdot \frac{d}{n_P} + 2\mathcal{E}_Q(w_P^*)\right\}$$

which extends the results of Mousavi Kalan et al. [2020], Zhang et al. [2022b], Ge et al. [2023] to the setting with both source and target samples.

3.4 Lower-Bounds

Let $\ell(a, b) = \mathbb{1}\{a \neq b\}$ and $\mathcal{Y} = \{-1, 1\}$.

Definition 9 (Weak Modulus Class in Classification). Let \mathcal{H} denote a class. Consider any non-decreasing function f on $(0,1] \mapsto [0,1]$. For any such f, and $0 \leq \beta_P, \beta_Q \leq 1$, let $\Sigma_{\mathcal{H}}(f, \beta_P, \beta_Q)$ denote all pairs of distributions P, Q such that (i) $\delta_{P,Q}(\epsilon) \leq f(\epsilon), \forall 0 < \epsilon \leq 1/4$, and (ii) P and Q satisfy BCC (Definition 8) with parameters $(1, \beta_P)$ and $(1, \beta_Q)$ respectively.

Theorem 2 (Classification Lower-Bound). In what follows, let \hat{h} denote any learner having access to independent datasets S_P and S_Q of size n_P , n_Q respectively. The following holds for any \mathcal{H} with $|\mathcal{H}| \geq 3$.

(i) Suppose the VC dimension of \mathcal{H} satisfies $d_{\mathcal{H}} \leq \max\{n_Q, n_P\}$, and that for some $\kappa \geq 1$, f satisfies $\alpha f(\epsilon) \leq \kappa f(\alpha \epsilon)$ for all $0 < \epsilon, \alpha \leq 1$. We have that, for universal constants c, c_0 :

$$\inf_{\hat{h}} \sup_{(P,Q)\in\Sigma_{\mathcal{H}}(f,\beta_{P},\beta_{Q})} \mathbb{E} \ \mathcal{E}_{Q}(\hat{h}) \ge c \cdot \frac{1}{\kappa} \cdot \min\left\{ \left(\frac{d_{\mathcal{H}}}{n_{Q}}\right)^{\frac{1}{2-\beta_{Q}}}, f\left(c_{0}\left(\frac{d_{\mathcal{H}}}{n_{P}}\right)^{\frac{1}{2-\beta_{P}}}\right) \right\}.$$
(9)

(ii) For general nondecreasing f we have that, for universal constants c, c_0 :

$$\inf_{\hat{h}} \sup_{(P,Q)\in\Sigma_{\mathcal{H}}(f,\beta_{P},\beta_{Q})} \mathbb{E} \ \mathcal{E}_{Q}(\hat{h}) \geq c \cdot \min\left\{ \left(\frac{1}{n_{Q}}\right)^{\frac{1}{2-\beta_{Q}}}, f\left(c_{0}\left(\frac{1}{n_{P}}\right)^{\frac{1}{2-\beta_{P}}}\right) \right\}$$

Furthermore, for general f, for any $2 \le d \le \max\{n_Q, n_P\}$ there exists \mathcal{H} with $d_{\mathcal{H}} = \Theta(d)$ such that (9) holds.

Remark 11. Notice that condition on f admits any concave function f with $\kappa = 1$ (in addition, nondecreasing), and allows some convex f for $\kappa \ge 1$. As it turns out, the structure is needed in our construction in order to satisfy $\delta(\epsilon) \le \epsilon$ for relevant values of ϵ .

Supporting Results.

Proposition 4 (Thm 2.5 of Tsybakov [2009]). Let $\{\Pi_h\}_{h \in \mathcal{H}}$ be a family of distributions indexed over a subset \mathcal{H} of a semi-metric (\mathcal{F} , dist). Suppose $\exists h_0, \ldots, h_M \in \mathcal{H}$, where $M \ge 2$, such that:

(i) dist
$$(h_i, h_j) \ge 2s > 0$$
, $\forall 0 \le i < j \le M$,
(ii) $\Pi_{h_i} \ll \Pi_{h_0} \quad \forall i \in [M]$, and the average KL-divergence to Π_{h_0} satisfies
 $\frac{1}{M} \sum_{i=1}^M \mathcal{D}_{kl}(\Pi_{h_i} | \Pi_{h_0}) \le \alpha \log M$, where $0 < \alpha < 1/8$.

Let $Z \sim \Pi_h$, and let $\hat{h} : Z \mapsto \mathcal{F}$ denote any *improper* learner of $h \in \mathcal{H}$. We have for any \hat{h} :

$$\sup_{h \in \mathcal{H}} \Pi_h \left(\operatorname{dist}(\hat{h}(Z), h) \ge s \right) \ge \frac{\sqrt{M}}{1 + \sqrt{M}} \left(1 - 2\alpha - \sqrt{\frac{2\alpha}{\log(M)}} \right) \ge \frac{3 - 2\sqrt{2}}{8}$$

The following proposition would be needed to construct packings (of spaces of distributions) of the appropriate size. **Proposition 5** (Varshamov-Gilbert bound). Let $d \ge 8$. Then there exists a subset $\{\sigma_0, \ldots, \sigma_M\}$ of $\{-1, 1\}^d$ such that $\sigma_0 = (1, \ldots, 1)$,

$$\operatorname{dist}(\sigma_i, \sigma_j) \ge \frac{d}{8}, \quad \forall \, 0 \le i < j \le M, \quad \text{and} \quad M \ge 2^{d/8},$$

where dist $(\sigma, \sigma') \doteq \operatorname{card}(\{i \in [m] : \sigma(i) \neq \sigma'(i)\})$ is the Hamming distance.

Results similar to the following lemma are known.

Lemma 2 (A basic KL upper-bound). For any 0 < p, q < 1, we let $\mathcal{D}_{kl}(p|q)$ denote $\mathcal{D}_{kl}(Ber(p)|Ber(q))$. Now let $0 < \epsilon < 1/2$ and let $z \in \{-1, 1\}$. We have

$$\mathcal{D}_{kl}(1/2 + (z/2) \cdot \epsilon | 1/2 - (z/2) \cdot \epsilon) \leq C_0 \cdot \epsilon^2$$
, for some C_0 independent of ϵ .

Proof of Theorem 2. For claim (i), the case $d_{\mathcal{H}} = 1$ is implied by claim (ii), so we will focus our proof of claim (i) on the case $d_{\mathcal{H}} \geq 2$. We establish both claims at once by introducing appropriate notation, as follows. For \mathcal{H} and f as in claim (i), let $d = d_{\mathcal{H}} - 1$, $\kappa_0 = \kappa$, and let x_0, x_1, \ldots, x_d be a shatterable subset of \mathcal{X} under \mathcal{H} . For \mathcal{H} and f as in claim (ii), instead let d = 1 and $\kappa_0 = 1$, and let x_0, x_1 be such that there exist $h_{-1}, h_1 \in \mathcal{H}$ with $h_y(x_1) = y$, and $h_{-1}(x_0) = h_1(x_1)$ (such x_0, x_1, h_{-1}, h_1 must exist since $|\mathcal{H}| \geq 3$), and without loss of generality suppose $h_{-1}(x_0) = h_1(x_0) = 1$. The points x_0, \ldots, x_d will form the support of marginals P_X, Q_X . Furthermore, let $\tilde{\mathcal{H}}$ denote the projection of \mathcal{H} onto $\{x_i\}_{i=0}^d$ (i.e., the quotient space of equivalences $h \equiv h'$ on $\{x_i\}$), with the additional constraint that all $h \in \tilde{\mathcal{H}}$ classify x_0 as 1. We can now restrict attention to $\tilde{\mathcal{H}}$ as the effective class.

Let $\sigma \in \{-1, 1\}^d$. We will construct a family of distribution pairs (P_{σ}, Q_{σ}) indexed by σ to which we then apply Proposition 4 above. For any P_{σ}, Q_{σ} , we let $\eta_{P,\sigma}, \eta_{Q,\sigma}$ denote the corresponding regression functions (i.e., $\mathbb{E}_{P_{\sigma}}[Y|x]$, and $\mathbb{E}_{Q_{\sigma}}[Y|x]$). To proceed, fix

$$\epsilon_P = c_0 \cdot \left(\frac{d}{n_P}\right)^{\frac{1}{2-\beta_P}}, \ \epsilon_Q = \left(\frac{d}{n_Q}\right)^{\frac{1}{2-\beta_Q}}, \ \text{and} \ \epsilon = c_1 \cdot \min\left\{\epsilon_Q, f(\epsilon_P)\right\},$$

for some c_0, c_1 to be defined so that $\epsilon_P, \epsilon < 1/2$.

- Distribution Q_{σ} . We have that $Q_{\sigma} = Q_X \times Q_{Y|X}^{\sigma}$, where $Q_X(x_0) = 1 - \epsilon^{\beta_Q}$, while $Q_X(x_i) = \frac{1}{d\kappa_0} \epsilon^{\beta_Q}$ for all $i \ge 1$. Now, the conditional $Q_{Y|X}^{\sigma}$ is fully determined by $\eta_{Q,\sigma_i}(x_0) = 1$, and $\eta_{Q,\sigma}(x_i) = 1/2 + (\sigma_i/2) \cdot \epsilon^{1-\beta_Q}$, $i \ge 1$.

- Distribution P_{σ} . We have that $P_{\sigma} = P_X \times P_{Y|X}^{\sigma}$, $P_X(x_0) = 1 - \epsilon_P^{\beta_P}$, while $P_X(x_i) = \frac{1}{d} \epsilon_P^{\beta_P}$, $i \ge 1$. Now, the conditional $P_{Y|X}^{\sigma}$ is fully determined by $\eta_{P,\sigma}(x_0) = 1$, and $\eta_{P,\sigma}(x_i) = 1/2 + (\sigma_i/2) \cdot \epsilon_P^{1-\beta_P}$, $i \ge 1$.

- Verifying that $(P_{\sigma}, Q_{\sigma}) \in \Sigma(f, \beta_P, \beta_Q)$. For any $\sigma \in \{-1, 1\}^d$, let $h_{\sigma} \in \tilde{\mathcal{H}}$ denote the corresponding Bayes classifier (remark that the Bayes is the same for both P_{σ} and Q_{σ}). Now, pick any other $h_{\sigma'} \in \tilde{\mathcal{H}}$, and let dist (σ, σ') denote the Hamming distance between σ, σ' (as in Proposition 5). We then have that

$$\mathcal{E}_{Q_{\sigma}}(h_{\sigma'}) = \operatorname{dist}(\sigma, \sigma') \cdot \frac{1}{d\kappa_0} \epsilon^{\beta_Q} \cdot \epsilon^{1-\beta_Q} = \frac{\operatorname{dist}(\sigma, \sigma')}{d\kappa_0} \cdot \epsilon, \text{ while } Q_X(h_{\sigma'} \neq h_{\sigma}) = \frac{\operatorname{dist}(\sigma, \sigma')}{d\kappa_0} \cdot \epsilon^{\beta_Q}, \quad (10)$$

and similarly,
$$\mathcal{E}_{P_{\sigma}}(h_{\sigma'}) = \frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot \epsilon_P$$
, while $P_X(h_{\sigma'} \neq h_{\sigma}) = \frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot \epsilon_P^{\beta_P}$. (11)

Since $(\operatorname{dist}(\sigma, \sigma')/d) \leq 1$ and $\kappa_0 \geq 1$, it follows that BCC (Definition 8) holds with parameters $(1, \beta_P)$ and $(1, \beta_Q)$ respectively for any P_{σ} and Q_{σ} (noting that $\mathcal{H}_P(1/2)$ and $\mathcal{H}_Q(1/2)$ both project to $\tilde{\mathcal{H}}$).

Moreover, notice that for every such $\mathcal{E}_{P_{\sigma}}(h_{\sigma'})$, we also have that (in both cases (i) and (ii))

$$\mathcal{E}_{Q_{\sigma}}(h_{\sigma'}) \leq \frac{c_1}{\kappa_0} \cdot \frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot f(\epsilon_P) \leq \frac{c_1}{\kappa_0} \cdot \kappa_0 \cdot f\left(\frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot \epsilon_P\right) \leq f\left(\mathcal{E}_{P_{\sigma}}(h_{\sigma'})\right).$$

Notice that, by construction, the mapping $\mathcal{E}_{P_{\sigma}}(h) \mapsto \mathcal{E}_{Q_{\sigma}}(h)$ is nondecreasing, hence the above implies that, for any $\epsilon' = \frac{k}{d} \cdot \epsilon_P, k \in \{0, \dots, d\}$, we do have $\delta(\epsilon') \leq f(\epsilon')$. Now, for values $\frac{k}{d} \cdot \epsilon_P < \epsilon' < \frac{k+1}{d} \cdot \epsilon_P, k \in \{0, \dots, d-1\}$, note that $\mathcal{H}_P(\epsilon') = \mathcal{H}_P(\frac{k}{d} \cdot \epsilon_P)$ so that $\delta(\epsilon') = \delta(\frac{k}{d} \cdot \epsilon_P) \leq f(\epsilon')$ (since f is non-decreasing). Noting that the requirement of $\Sigma_{\mathcal{H}}(f, \beta_P, \beta_Q)$ only concerns $\delta(\epsilon') \leq f(\epsilon')$ for $\epsilon' \leq 1/4$, and recalling that $\epsilon_P \leq 1/2$, this completely verifies that $(P_{\sigma}, Q_{\sigma}) \in \Sigma_{\mathcal{H}}(f, \beta_P, \beta_Q)$ (for both case (i) and case (ii)).

- *Reduction to a packing.* Now apply Proposition 5 to identify a subset Σ of $\{-1,1\}^d$, where $|\Sigma| = M \ge 2^{d/8}$, and $\forall \sigma, \sigma' \in \Sigma$, we have dist $(\sigma, \sigma') \ge d/8$. It should be clear then that for any $\sigma, \sigma' \in \Sigma$,

$$\mathcal{E}_{Q_{\sigma}}(h_{\sigma'}) \geq \frac{d}{8} \cdot \frac{1}{d\kappa_0} \epsilon^{\beta_Q} \cdot \epsilon^{1-\beta_Q} = \epsilon/(8\kappa_0).$$

Furthermore, by construction, any classifier $\hat{h} : \{x_i\} \mapsto \{0, 1\}$ can be reduced to a decision on σ , and we henceforth view dist (σ, σ') as the semi-metric referenced in Proposition 4, with effective indexing set Σ .

- *KL* bounds in terms of n_P and n_Q . Define $\Pi_{\sigma} = P_{\sigma}^{n_P} \times Q_{\sigma}^{n_Q}$. We can now verify that all $\Pi_{\sigma}, \Pi_{\sigma'}$ are close in KL-divergence. First notice that, for any $\sigma, \sigma' \in \Sigma$ (in fact in $\{-1, 1\}^d$)

$$\mathcal{D}_{kl}(\Pi_{\sigma}|\Pi_{\sigma'}) = n_P \cdot \mathcal{D}_{kl}(P_{\sigma}|P_{\sigma'}) + n_Q \cdot \mathcal{D}_{kl}(Q_{\sigma}|Q_{\sigma'})$$

$$= n_P \cdot \mathop{\mathbb{E}}_{P_X} \mathcal{D}_{kl}\left(P_{Y|X}^{\sigma}|P_{Y|X}^{\sigma'}\right) + n_Q \cdot \mathop{\mathbb{E}}_{Q_X} \mathcal{D}_{kl}\left(Q_{Y|X}^{\sigma}|Q_{Y|X}^{\sigma'}\right)$$

$$= n_P \cdot \sum_{i=1}^d \frac{\epsilon_P^{\beta_P}}{d} \mathcal{D}_{kl}\left(P_{Y|x_i}^{\sigma}|P_{Y|x_i}^{\sigma'}\right) + n_Q \cdot \sum_{i=1}^d \frac{\epsilon^{\beta_Q}}{d\kappa_0} \mathcal{D}_{kl}\left(Q_{Y|x_i}^{\sigma}|Q_{Y|x_i}^{\sigma'}\right)$$

$$\leq C_0\left(n_P \cdot \epsilon_P^{(2-\beta_P)} + (n_Q/\kappa_0) \cdot \epsilon^{(2-\beta_Q)}\right)$$
(12)

$$\leq C_0 d(c_0^{(2-\beta_p)} + c_1^{2-\beta_Q}) \leq 2C_0(c_0 \vee c_1)d.$$
(13)

where, for inequality (12), we used Lemma 2 to upper-bound the divergence terms. It follows that, for c_0, c_1 sufficiently small so that $2C_0(c_0 \vee c_1) \leq 1/16$, we get that (13) is upper bounded by $(1/8) \log M$. Now apply Proposition 4 and conclude with the first two claims (i) and (ii) of the theorem.

For the final claim in the theorem, consider $\mathcal{H} \doteq \{h_{\sigma}\}_{\sigma \in \Sigma}$, where Σ is supplied by Proposition 5 as just described above. In particular, since all functions in \mathcal{H} agree on x_0 , and $2^{d/8} \leq |\mathcal{H}| \leq 2^d$, the VC dimension $d_{\mathcal{H}}$ satisfies $d_{\mathcal{H}} \leq d$. Moreover, Sauer's lemma [Sauer, 1972, Vapnik and Chervonenkis, 1974] implies $2^{d/8} \leq \left(\frac{ed}{d_{\mathcal{H}}}\right)^{d_{\mathcal{H}}}$, that is, $d \leq 8d_{\mathcal{H}} \log_2\left(\frac{ed}{d_{\mathcal{H}}}\right)$, which further implies $d \leq 16d_{\mathcal{H}} \log_2(8e) \leq 72d_{\mathcal{H}}$, so that $d_{\mathcal{H}} \geq d/72$. Thus, $d_{\mathcal{H}} = \Theta(d)$.

For this class, we modify the construction slightly as follows: now set $\epsilon = \min \{\epsilon_Q, \frac{1}{8}\epsilon_P\}$, and follow the same construction as is case (i), but with κ set to 1.

It follows from (10) and (11) that the BCC conditions hold, while we can verify the weak moduli is appropriately bounded. Indeed, we have for any $\mathcal{E}_{P_{\sigma}}(h_{\sigma'}) = \frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot \epsilon_P$ that

$$\mathcal{E}_{Q_{\sigma}}(h_{\sigma'}) \leq \frac{\operatorname{dist}(\sigma, \sigma')}{d} f\left(\frac{1}{8} \cdot \epsilon_{P}\right) \leq f\left(\frac{1}{8} \cdot \epsilon_{P}\right) \leq f\left(\frac{\operatorname{dist}(\sigma, \sigma')}{d} \cdot \epsilon_{P}\right),$$

since dist $(\sigma, \sigma') \ge \frac{d}{8}$ for $\sigma \neq \sigma'$.

The rest of the argument is identical to the above cases (i) and (ii).

4 Strong Modulus of Transfer

Definition 10. Let μ denote either P or Q. Let $0 < \tau \leq 1$, $\epsilon > 0$, and C > 1. We call a random set $\hat{\mathcal{H}}_{\mu} \doteq \hat{\mathcal{H}}_{\mu}(S_{\mu})$ an (ϵ, τ, C) -strong confidence set (under μ) if the following conditions are met with probability at least $1 - \tau$:

$$\mathcal{H}_{\mu}(\epsilon/C) \subset \mathcal{H}_{\mu} \subset \mathcal{H}_{\mu}(\epsilon).$$

Notice that if we pick ϵ such that, with probability at least $1 - \tau$, $\mathcal{E}_{\mu}(\hat{h}_{\mu}) < \epsilon/C$, then an (ϵ, τ, C) -strong confidence set is also an $(\epsilon, 2\tau)$ -weak confidence set (taking $\hat{\epsilon}_{\mu} = \epsilon/C$ in Definition 7). In fact, for most typical ways of obtaining such strong confidence sets, the events referred to in the failure probabilities are nested, so that the (ϵ, τ, C) -strong confidence set is in fact an (ϵ, τ) -weak confidence set.

Algorithm 2. Input: $\hat{\mathcal{H}}_Q^{\sharp} \supset \hat{\mathcal{H}}_Q^{\flat}$, and $\hat{\mathcal{H}}_P$, all subsets of \mathcal{H} .

If
$$\hat{\mathcal{H}}_{O}^{\sharp} \cap \hat{\mathcal{H}}_{P} \neq \emptyset$$
, return any $h \in \hat{\mathcal{H}}_{O}^{\sharp} \cap \hat{\mathcal{H}}_{P}$, otherwise return $\hat{h}_{P,Q} \doteq \operatorname{argmin}_{h \in \hat{\mathcal{H}}_{O}^{\flat}} \hat{R}_{P}(h)$.

We require the following local definition of achievable rate (under P) over a given subset \mathcal{H}' of \mathcal{H} . This is because, as we consider general scenarios, it is possible that $\hat{h}_P(\mathcal{H}')$ may admit a different generalization rate than $\hat{h}_P(\mathcal{H})$ if population infimums differ; for instance in classification, the Bernstein Class Condition on \mathcal{H} (Definition 8) may allow for fast rates $\ll n_P^{-1/2}$ for $\hat{h}_P(\mathcal{H})$, but if the same BCC parameters do not hold for \mathcal{H}' we may have $\tilde{\varepsilon}_P(\mathcal{H}') \approx n_P^{-1/2}$. **Definition 11.** Fix $0 < \tau < 1$. Given a subset $\mathcal{H}' \subset \mathcal{H}$, we let $\tilde{\varepsilon}_P(\mathcal{H}')$, depending on n_P , denote the following:

$$\min\left\{\epsilon > 0: \mathbb{P}_{S_P}\left(R_P(\hat{h}_P(\mathcal{H}')) - \inf_{h \in \mathcal{H}'} R_P(h) \le \epsilon\right) \ge 1 - \tau\right\}.$$

Note that the infimum is achieved in the set by continuity of probability measures.

Theorem 3. Let $0 < \tau \leq 1$, $C^{\sharp}, C^{\flat} > 1$. Suppose $\hat{\mathcal{H}}_Q^{\sharp}, \hat{\mathcal{H}}_Q^{\flat}$ are respectively $(\epsilon_Q, \tau, C^{\sharp})$ and $(\epsilon_Q/C^{\sharp}, \tau, C^{\flat})$ strong confidence sets under Q, and $\hat{\mathcal{H}}_P$ is an (ϵ_P, τ) -weak confidence set for P, for some $\epsilon_Q, \epsilon_P > 0$ depending
respectively on n_Q and n_P . Let \hat{h} denote the hypothesis returned by Algorithm 2. Letting $\tilde{\varepsilon}_P = \tilde{\varepsilon}_P(\hat{\mathcal{H}}_Q^{\flat})$, we then
have with probability at least $1 - 4\tau$:

$$\mathcal{E}_Q(\hat{h}) \leq \begin{cases} \delta(\epsilon_Q, \epsilon_P) & \text{if } \epsilon_Q > \varepsilon_{P,Q} \text{ or } \hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P \neq \emptyset, \\ C^{\flat} \cdot \delta(\epsilon_Q/C^{\sharp}, \tilde{\varepsilon}_P) & \text{otherwise.} \end{cases}$$

Proof. Assume the events of Definition 10 (for $\hat{\mathcal{H}}_Q^{\sharp}, \hat{\mathcal{H}}_Q^{\flat}$) and Definition 7 (for $\hat{\mathcal{H}}_P$) hold (this occurs with probability at least $1 - 3\tau$ by the union bound). By Definition 10, every $h \in \hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P$ is in $\mathcal{H}_Q(\epsilon_Q) \cap \mathcal{H}_P(\epsilon_P)$. Now notice that

for any such h in this last intersection, we have $\mathcal{E}_P(h; \mathcal{H}_Q(\epsilon_Q)) \leq \mathcal{E}_P(h) \leq \epsilon_P$. Thus, by definition, we have that any $h \in \hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P$ satisfies $\mathcal{E}_Q(h) \leq \delta(\epsilon_Q, \epsilon_P)$.

Now suppose $\epsilon_Q > \epsilon_{P,Q}$, and $\hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P = \emptyset$. Recall the equivalent definition of $\delta(\epsilon_Q, \epsilon_P)$ of Corollary 1 for the case $\epsilon_Q > \epsilon_{P,Q}$. By Proposition 1, for any $\epsilon > 0$, $\mathcal{H}_Q(\epsilon_Q) \cap \mathcal{H}_P(\epsilon) \neq \emptyset$; therefore pick any $h \in \mathcal{H}_Q(\epsilon_Q) \cap \mathcal{H}_P(\hat{\epsilon}_P)$ (recalling $\hat{\epsilon}_P > 0$ from Definition 7). Since $\mathcal{H}_P(\hat{\epsilon}_P) \subset \hat{\mathcal{H}}_P \subset \mathcal{H}_P(\epsilon_P)$, it follows that $h \in \mathcal{H}_Q(\epsilon_Q) \cap \mathcal{H}_P(\epsilon_P)$, hence, $\mathcal{E}_Q(h) \leq \delta(\epsilon_Q, \epsilon_P)$. Furthermore, since $\hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P = \emptyset$, we know $h \notin \mathcal{H}_Q(\epsilon_Q/C^{\sharp}) \subset \hat{\mathcal{H}}_Q^{\sharp}$, i.e., $\mathcal{E}_Q(h) > \epsilon_Q/C^{\sharp}$. In other words, for the choice $\hat{h}_{P,Q} \in \hat{\mathcal{H}}_Q^{\flat} \subset \mathcal{H}_Q(\epsilon_Q/C^{\sharp})$, we must have that

$$\mathcal{E}_Q(\hat{h}_{P,Q}) \le \epsilon_Q/C^{\sharp} < \mathcal{E}_Q(h) \le \delta(\epsilon_Q, \epsilon_P)$$

For the final case, suppose $\epsilon_Q \leq \varepsilon_{P,Q}$, while $\hat{\mathcal{H}}_Q^{\sharp} \cap \hat{\mathcal{H}}_P = \emptyset$. We have the following two subcases.

– First, suppose that $\inf_{h \in \mathcal{H}_Q(\epsilon_Q/C^{\sharp})} R_P(h) = \inf_{h \in \mathcal{H}_Q(\epsilon_Q/(C^{\flat}C^{\sharp}))} R_P(h)$. Then it follows that

$$\mathcal{E}_P(\hat{h}_{P,Q}; \mathcal{H}_Q(\epsilon_Q/C^{\sharp})) = \mathcal{E}_P(\hat{h}_{P,Q}; \hat{\mathcal{H}}_Q^{\flat}) \leq \tilde{\varepsilon}_P \text{ (with probability at least } 1 - \tau)$$
.

Since we also have that $\hat{h}_{P,Q} \in \mathcal{H}_Q(\epsilon_Q/C^{\sharp})$, it follows by definition that it satisfies $\mathcal{E}_Q(\hat{h}_{P,Q}) \leq \delta(\epsilon_Q/C^{\sharp}, \tilde{\epsilon}_P)$. - Otherwise, let $0 < \epsilon \leq \tilde{\epsilon}_P$, and pick $\tilde{h} \in \mathcal{H}_Q(\epsilon_Q/C^{\sharp})$ such that

$$\inf_{h \in \mathcal{H}_Q(\epsilon_Q/C^{\sharp})} R_P(h) \le R_P(\tilde{h}) < \min\left\{\inf_{h \in \mathcal{H}_Q(\epsilon_Q/(C^{\flat}C^{\sharp}))} R_P(h), \inf_{h \in \mathcal{H}_Q(\epsilon_Q/C^{\sharp})} R_P(h) + \epsilon\right\}.$$

By the second inequality, namely, the first term in the min, such an $\tilde{h} \notin \mathcal{H}_Q(\epsilon_Q/(C^{\flat}C^{\sharp}))$. Using the second term in the min, we have that $\mathcal{E}_Q(\tilde{h}) \leq \delta(\epsilon_Q/C^{\sharp}, \epsilon)$. We therefore have that

$$\mathcal{E}_Q(\hat{h}_{P,Q}) \le \epsilon_Q/C^{\sharp} < C^{\flat} \mathcal{E}_Q(\tilde{h}) \le C^{\flat} \delta(\epsilon_Q/C^{\sharp}, \epsilon) \le C^{\flat} \delta(\epsilon_Q/C^{\sharp}, \tilde{\varepsilon}_P).$$

We remark that each of these final two subcases yields a bound slightly better than the one stated in the theorem, which is given as such for simplicity.

4.1 Examples of Strong Confidence Sets

We present some examples of strong confidence sets for classical problems in classification and regression. We note in particular that, while Theorem 3 requires two such confidence sets for different values of ϵ , the constructions below are for a single ϵ but are stated in a way to make it clear that different values of ϵ are admissible by varying constant factors in their definitions.

4.1.1 Classification Example

We consider classification with a VC class \mathcal{H} . In what follows assume $\ell(a, b) = \mathbb{1}\{a \neq b\}$ and for the simplest case suppose $\mathcal{Y} = \{-1, 1\}$. We first note that a simple strong confidence set follows easily from usual \sqrt{n} results in this setting as stated in the firt proposition below. However, this corresponds to assuming $\beta_{\mu} = 0$ (noise condition of Definition8); thus, for unknown β_{μ} , more sophisticated construction is required to achieve the corresponding fast rates. Such a construction is subsequently described.

Proposition 6 (\sqrt{n} -confidence sets). Let $\alpha'_{\mu}(\tau) \doteq \sqrt{n^{-1}(d_{\mathcal{H}} + \log(1/\tau))}$. For a universal constant C, the set

$$\hat{\mathcal{H}}_{\mu} \doteq \left\{ h \in \mathcal{H} : \hat{\mathcal{E}}_{\mu}(h) \le C \sqrt{\alpha'_{\mu}(\tau)} \right\}$$

is an $(\epsilon, \tau, 2)$ -strong confidence set (under μ), for $\epsilon = (4/3)C\sqrt{\alpha'_{\mu}(\tau)}$.

Proof. The classic result of Talagrand [1994] provides that, for a universal constant C, with probability at least $1 - \tau$,

$$\forall h \in \mathcal{H}, \left| \hat{R}_{\mu}(h) - R_{\mu}(h) \right| \le \frac{C}{6} \sqrt{\alpha'_{\mu}(\tau)}.$$
(14)

On this event, every $h \in \hat{\mathcal{H}}_{\mu}$ satisfies

$$\mathcal{E}_{\mu}(h) = R_{\mu}(h) - \inf_{h' \in \mathcal{H}} R_{\mu}(h') \leq \hat{R}_{\mu}(h) - \inf_{h' \in \mathcal{H}} \hat{R}_{\mu}(h') + \frac{C}{3}\sqrt{\alpha'_{\mu}(\tau)} = \hat{\mathcal{E}}_{\mu}(h) + \frac{C}{3}\sqrt{\alpha'_{\mu}(\tau)} \leq \frac{4}{3}C\sqrt{\alpha'_{\mu}(\tau)},$$

where the first inequality is by (14) and the last inequality follows from the definition of $\hat{\mathcal{H}}_{\mu}$. Moreover, on this same event, any $h \in \mathcal{H}$ with $\mathcal{E}_{\mu}(h) \leq \frac{2C}{3} \sqrt{\alpha'_{\mu}(\tau)}$ has

$$\hat{\mathcal{E}}_{\mu}(h) \leq \mathcal{E}_{\mu}(h) + \frac{C}{3}\sqrt{\alpha'_{\mu}(\tau)} \leq C\sqrt{\alpha'_{\mu}(\tau)},$$

and hence $h \in \hat{\mathcal{H}}_{\mu}$. Thus, for $\epsilon = \frac{4}{3}C\sqrt{\alpha'_{\mu}(\tau)}$, we have $\mathcal{H}_{\mu}(\epsilon/2) \subset \hat{\mathcal{H}}_{\mu} \subset \mathcal{H}_{\mu}(\epsilon)$.

Remark 12. Notice that, by the above arguments, we can also obtain a $(\epsilon', \tau, 2)$ -strong confidence set, for $\epsilon' = 2\epsilon$ simply by plugging in ϵ' wherever ϵ appears in the definition of $\hat{\mathcal{H}}_{\mu}$.

To allow for fast rates, and bounds for general β in the Bernstein Class Condition (Definition 8), we refine the above strong confidence set definition based on localization arguments involving the *uniform Bernstein inequality* (Lemma 1). The result, and underlying principle, is similar to results of Koltchinskii [2006] stated there in a more general Rademacher-based formulation (namely, a combination of Lemma 2 and Theorem 3 therein).

Proposition 7. Let C_0 and $\alpha_{\mu}(\tau)$ be as in Lemma 1. Let $2^{-\mathbb{N}_0} \doteq \{2^{-i} : i \in \mathbb{N} \cup \{0\}\}$. Let $C_1, \dots, C_7 \in \{2^i : i \in \mathbb{N}\}$. Let $\bar{\epsilon}_{\mu}^{\text{loc}}$ denote the minimal element of $2^{-\mathbb{N}_0}$ such that $\forall \epsilon' \in 2^{-\mathbb{N}_0}$ with $\epsilon' \geq \bar{\epsilon}_{\mu}^{\text{loc}}$,

$$\min\left\{C_0\sqrt{\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_1\epsilon'))\cdot\alpha_{\mu}(\tau)}+C_0\cdot\alpha_{\mu}(\tau),1\right\}\leq\epsilon'/C_2.$$
(15)

For any $\epsilon' > 0$, define $\mathcal{H}_{\hat{\mu}}(\epsilon') \doteq \left\{h \in \mathcal{H} : \hat{\mathcal{E}}_{\mu}(h) \leq \epsilon'\right\}$ and for any $\mathcal{H}' \subset \mathcal{H}$, define the empirical diameter $\operatorname{diam}_{\hat{\mu}}(\mathcal{H}') \doteq \sup_{h,h' \in \mathcal{H}'} \hat{\mu}(h \neq h')$. Let $\hat{\epsilon}_{\mu}^{\operatorname{loc}}$ denote the minimal element of $2^{-\mathbb{N}_0}$ such that $\forall \epsilon' \in 2^{-\mathbb{N}_0}$ with $\epsilon' \geq \hat{\epsilon}_{\mu}^{\operatorname{loc}}$,

$$\min\left\{C_0\sqrt{\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_3\epsilon'))\cdot\alpha_{\mu}(\tau)}+C_0\cdot\alpha_{\mu}(\tau),1\right\}\leq\epsilon'/C_4.$$
(16)

If $2C_6 \ge C_1$, $(1/C_5) + (2C_6/(C_1C_2)) \le 1$, $3/2 + (2C_6/(C_1C_2)) \le C_6$, $C_1 + 2/C_4 \le C_3$, $2C_2 \le C_4$, $(C_6C_3/C_1)C_7 \ge 1$, and $2(C_6C_3/(C_1C_2)) \le 1/C_4$, then with probability at least $1 - \tau$, for any $\epsilon \in 2^{-\mathbb{N}_0}$ with $\epsilon \ge \overline{\epsilon}_{\mu}^{\mathrm{loc}}$,

$$\mathcal{H}_{\mu}(\epsilon/C_5) \subset \mathcal{H}_{\hat{\mu}}(\epsilon) \subset \mathcal{H}_{\mu}(C_6\epsilon) \tag{17}$$

and moreover,

$$\bar{\epsilon}_{\mu}^{\rm loc} \le \hat{\epsilon}_{\mu}^{\rm loc} \le C_7 \bar{\epsilon}_{\mu}^{\rm loc}. \tag{18}$$

In particular, this implies that for any $C \in \{2^i : i \in \mathbb{N} \cup \{0\}\}, \hat{\mathcal{H}}_{\mu} \doteq \mathcal{H}_{\hat{\mu}}(C\hat{\epsilon}_{\mu}^{\text{loc}})$ is a $(C_6C_7C\bar{\epsilon}_{\mu}^{\text{loc}}, \tau, C_7C_5)$ -strong confidence set.

Moreover, for (C_{μ}, β_{μ}) as in the Bernstein Class Condition (Definition 8), it holds that

$$\bar{\epsilon}_{\mu}^{\rm loc} \le C'(C_{\mu} \cdot \alpha_{\mu}(\tau))^{\frac{1}{2-\beta_{\mu}}}$$

for a constant $C' = C(C_1, C_2)$.

Proof. Throughout the proof, suppose the event of probability at least $1 - \tau$ from Lemma 1 holds. We proceed to prove (17) by induction. As a base case, considering $\epsilon = 1$, we trivially have $\mathcal{H}_{\mu}(1/C_5) \subset \mathcal{H}_{\hat{\mu}}(1) \subset \mathcal{H}_{\mu}(C_6)$. Now take as an inductive hypothesis that, for some $\epsilon \in 2^{-\mathbb{N}_0}$ with $1 > \epsilon \ge \bar{\epsilon}_{\mu}^{\mathrm{loc}}$, it holds that $\mathcal{H}_{\mu}(2\epsilon/C_5) \subset \mathcal{H}_{\hat{\mu}}(2\epsilon) \subset \mathcal{H}_{\mu}(C_62\epsilon)$. We will extend the inclusion (17) to hold for ϵ . Toward this end, consider any $h \in \mathcal{H}_{\mu}(\epsilon/C_5)$. We trivially have $h \in \mathcal{H}_{\mu}(C_62\epsilon)$, and by the inductive hypothesis we also have $\hat{h}_{\mu} \in \mathcal{H}_{\mu}(C_62\epsilon)$ as well. Therefore $\mu(h \neq \hat{h}_{\mu}) \le \mathrm{diam}_{\mu}(\mathcal{H}_{\mu}(C_62\epsilon))$. Thus, by (7) from Lemma 1, we have

$$\hat{\mathcal{E}}_{\mu}(h) \leq \epsilon/C_5 + \min\left\{C_0 \cdot \sqrt{\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_6 2 \epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1\right\} \leq \epsilon/C_5 + (2C_6/(C_1 C_2))\epsilon,$$

where the last inequality follows from the fact that $\epsilon \geq \bar{\epsilon}_{\mu}^{\text{loc}}$ and $2C_6 \geq C_1$. Since $(1/C_5) + (2C_6/(C_1C_2)) \leq 1$, we conclude that $\hat{\mathcal{E}}_{\mu}(h) \leq \epsilon$. Hence we have verified that $\mathcal{H}_{\mu}(\epsilon/C_5) \subset \mathcal{H}_{\hat{\mu}}(\epsilon)$.

Continuing to the second inclusion, consider any $h \in \mathcal{H}_{\hat{\mu}}(\epsilon)$ and any $h' \in \mathcal{H}_{\mu}(\epsilon/2)$. By the inductive hypothesis, $h \in \mathcal{H}_{\mu}(C_62\epsilon)$, and h' is trivially also in this set. Therefore, $\mu(h \neq h') \leq \text{diam}_{\mu}(\mathcal{H}_{\mu}(C_62\epsilon))$. Thus, by (7) of Lemma 1, we have

$$R_{\mu}(h) - R_{\mu}(h') \le \epsilon + \min\left\{C_0 \cdot \sqrt{\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_6 2\epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1\right\} \le \epsilon + (2C_6/(C_1 C_2))\epsilon_{\mu}(\tau)$$

where the last inequality is again due to the fact that $\epsilon \geq \bar{\epsilon}_{\mu}^{\text{loc}}$ and $2C_6 \geq C_1$. Since $h' \in \mathcal{H}_{\mu}(\epsilon/2)$, we also have that $\mathcal{E}_{\mu}(h) \leq R_{\mu}(h) - R_{\mu}(h') + \epsilon/2$. Since $1 + (1/2) + (2C_6/(C_1C_2)) \leq C_6$, we conclude that $\mathcal{E}_{\mu}(h) \leq C_6\epsilon$. Hence we have verified that $\mathcal{H}_{\hat{\mu}}(\epsilon) \subset \mathcal{H}_{\mu}(C_6\epsilon)$. By the principle of induction, this completes the proof of (17).

Next we establish (18), again on the same event from Lemma 1. We begin with the left inequality. We again proceed by induction, to establish that any $\epsilon \in 2^{-\mathbb{N}_0}$ with $\epsilon \geq \hat{\epsilon}_{\mu}^{\text{loc}}$ also has $\epsilon \geq \bar{\epsilon}_{\mu}^{\text{loc}}$, and moreover that $\mathcal{H}_{\mu}(C_1\epsilon) \subset \mathcal{H}_{\hat{\mu}}(C_3\epsilon)$ As a base case, we trivially have $1 \geq \bar{\epsilon}_{\mu}^{\text{loc}}$ and $\mathcal{H}_{\mu}(C_1) = \mathcal{H} = \mathcal{H}_{\hat{\mu}}(C_3)$ (since $C_1, C_3 \geq 1$). If $\hat{\epsilon}_{\mu}^{\text{loc}} = 1$, this concludes the proof. Otherwise, take as an inductive hypothesis that some $\epsilon \in 2^{-\mathbb{N}_0}$ with $1 > \epsilon \geq \hat{\epsilon}_{\mu}^{\text{loc}}$ has $2\epsilon \geq \bar{\epsilon}_{\mu}^{\text{loc}}$, and moreover $\mathcal{H}_{\mu}(C_12\epsilon) \subset \mathcal{H}_{\hat{\mu}}(C_32\epsilon)$. We will first argue that $\mathcal{H}_{\mu}(C_1\epsilon) \subset \mathcal{H}_{\hat{\mu}}(C_3\epsilon)$. Consider any $h \in \mathcal{H}_{\mu}(C_1\epsilon)$. By the inductive hypothesis, $h \in \mathcal{H}_{\hat{\mu}}(C_32\epsilon)$, and we trivially have $\hat{h} \in \mathcal{H}_{\hat{\mu}}(C_32\epsilon)$ as well, so that $\hat{\mu}(h \neq \hat{h}) \leq \text{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_32\epsilon))$. By (7), we have that

$$\hat{\mathcal{E}}_{\mu}(h) \le C_1 \epsilon + \min\left\{C_0 \cdot \sqrt{\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_3 2 \epsilon) \alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1\right\} \le C_1 \epsilon + 2\epsilon/C_4,$$

where the last inequality is based on the fact that $2\epsilon \ge \hat{\epsilon}_{\mu}^{\text{loc}}$. Since $C_1 + 2/C_4 \le C_3$, we conclude that $h \in \mathcal{H}_{\hat{\mu}}(C_3\epsilon)$. Therefore $\mathcal{H}_{\mu}(C_1\epsilon) \subset \mathcal{H}_{\hat{\mu}}(C_3\epsilon)$.

We therefore have that $\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_{1}\epsilon)) \leq \operatorname{diam}_{\mu}(\mathcal{H}_{\hat{\mu}}(C_{3}\epsilon))$. By (8) from Lemma 1, we have $\operatorname{diam}_{\mu}(\mathcal{H}_{\hat{\mu}}(C_{3}\epsilon)) \leq 2\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_{3}\epsilon)) + C_{0} \cdot \alpha_{\mu}(\tau)$. Therefore,

$$\begin{split} \min \bigg\{ C_0 \cdot \sqrt{\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_1 \epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1 \bigg\} &\leq 2 \cdot \min \bigg\{ C_0 \cdot \sqrt{\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_3 \epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1 \bigg\} \\ &\leq 2\epsilon/C_4, \end{split}$$

where this last inequality is based on the fact that $\epsilon \ge \hat{\epsilon}_{\mu}^{\text{loc}}$, and hence satisfies (16). Therefore, since $2/C_4 \le 1/C_2$, we conclude that every $\epsilon \in 2^{-\mathbb{N}_0}$ with $\epsilon \ge \hat{\epsilon}_{\mu}^{\text{loc}}$ also satisfies (15). Therefore, $\hat{\epsilon}_{\mu}^{\text{loc}} \ge \bar{\epsilon}_{\mu}^{\text{loc}}$.

For the second inequality, namely $C_7 \bar{\epsilon}_{\mu}^{\text{loc}} \geq \hat{\epsilon}_{\mu}^{\text{loc}}$, it suffices to argue that for any $\epsilon \in 2^{-\mathbb{N}_0}$ with $\epsilon \geq C_7 \bar{\epsilon}_{\mu}^{\text{loc}}$, (16) holds. Fix any such ϵ . By (17), we have $\mathcal{H}_{\hat{\mu}}(C_3\epsilon) \subset \mathcal{H}_{\mu}(C_6C_3\epsilon)$. Moreover, by (8), we have $\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\mu}(C_6C_3\epsilon)) \leq 2\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_6C_3\epsilon)) + C_0 \cdot \alpha_{\mu}(\tau)$. Therefore,

$$\min\left\{C_0 \cdot \sqrt{\operatorname{diam}_{\hat{\mu}}(\mathcal{H}_{\hat{\mu}}(C_3\epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1\right\} \leq 2\min\left\{C_0 \cdot \sqrt{\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}(C_6C_3\epsilon))\alpha_{\mu}(\tau)} + C_0 \cdot \alpha_{\mu}(\tau), 1\right\} \leq 2(C_6C_3/(C_1C_2))\epsilon,$$

where this last inequality follows from $(C_6C_3/C_1)C_7 \ge 1$. Since $2(C_6C_3/(C_1C_2)) \le 1/C_4$, we conclude that this ϵ indeed satisfies (16), which completes the proof of this inequality.

For the last claim in the proposition, note that by the Bernstein Class Condition, $\operatorname{diam}_{\mu}(\mathcal{H}_{\mu}((C_1/2)\bar{\epsilon}_{\mu}^{\operatorname{loc}})) \leq C_{\mu}((C_1/2)\bar{\epsilon}_{\mu}^{\operatorname{loc}})^{\beta_{\mu}}$. Minimality of $\bar{\epsilon}_{\mu}^{\operatorname{loc}}$ implies

$$(1/2)\overline{\epsilon}_{\mu}^{\mathrm{loc}}/C_{2} < C_{0}\sqrt{\mathrm{diam}_{\mu}\left(\mathcal{H}_{\mu}((C_{1}/2)\overline{\epsilon}_{\mu}^{\mathrm{loc}})\right) \cdot \alpha_{\mu}(\tau)} + C_{0} \cdot \alpha_{\mu}(\tau)$$

$$\leq C_{0}\sqrt{C_{\mu}((C_{1}/2)\overline{\epsilon}_{\mu}^{\mathrm{loc}})^{\beta_{\mu}} \cdot \alpha_{\mu}(\tau)} + C_{0} \cdot \alpha_{\mu}(\tau) \leq 2C_{0}\max\left\{\sqrt{C_{\mu}((C_{1}/2)\overline{\epsilon}_{\mu}^{\mathrm{loc}})^{\beta_{\mu}} \cdot \alpha_{\mu}(\tau)}, \alpha_{\mu}(\tau)\right\}.$$

Thus, either the $\alpha_{\mu}(\tau)$ term achieves the max, in which case $\bar{\epsilon}_{\mu}^{\text{loc}} \leq 4C_2C_0 \cdot \alpha_{\mu}(\tau) \leq 4C_2C_0 \cdot (C_{\mu} \cdot \alpha_{\mu}(\tau))^{\frac{1}{2-\beta_{\mu}}}$, or else the first term achieves the max, in which case solving for $\bar{\epsilon}_{\mu}^{\text{loc}}$ yields

$$\bar{\epsilon}_{\mu}^{\text{loc}} < \left(4C_2C_0\right)^{\frac{2}{2-\beta\mu}} \left(C_1/2\right)^{\frac{\beta\mu}{2-\beta\mu}} \left(C_{\mu} \cdot \alpha_{\mu}(\tau)\right)^{\frac{1}{2-\beta\mu}} \le \left(4C_2C_0\right)^2 \left(1 \lor C_1/2\right) \left(C_{\mu} \cdot \alpha_{\mu}(\tau)\right)^{\frac{1}{2-\beta\mu}}.$$

Remark 13. We remark that, using the above proposition to specify an (ϵ, τ, C) -strong confidence set, one can easily adjust it to be a $(C'\epsilon, \tau, C'' \cdot C)$ -strong confidence set, for any choices of $C', C'' \ge 1$, by scaling the constants C_3, C_4 appropriately in the definition.

4.1.2 Regression Example

We consider a linear regression setting with $X \in \mathbb{R}^d$ and $Y \in \mathbb{R}$, jointly distributed under μ . Assume throughout this section that $||X|| \leq 1$, and $Y - \mathbb{E}[Y|x]$ is uniformly subGaussian with parameter $\sigma_Y^2 \geq 1$; see Condition 3 of Hsu et al. [2012]. We consider the squared loss $\ell(y, y') \doteq (y - y')^2$, and $\mathcal{H} \doteq \{x \mapsto h_w(x) \doteq w^\top x : w \in \mathbb{R}^d\}$.

Define $\Sigma \doteq \mathbb{E}_{\mu} X X^{\top}$, assumed invertible, and $\hat{\Sigma} \doteq \mathbb{E}_{\hat{\mu}} X X^{\top}$, where $\hat{\mu}$ is the emprirical version of μ on $S_{\mu} \sim \mu^{n_{\mu}}$. We make use of the following corollary to classical concentration results.

Lemma 3 (Matrix concentration). For any $0 < \tau < 1$, the following holds with probability at least $1 - \tau$. There exists $C = C(\tau, d, \text{eigs}(\Sigma))$ such that, for $n_{\mu} \ge C$, we have (writing $A \preceq B$ for A - B is negative semidefinite):

$$\frac{1}{2}\Sigma \preceq \hat{\Sigma} \preceq \frac{3}{2}\Sigma. \quad \text{Equivalently } \forall v \in \mathbb{R}^d, \quad \frac{1}{2} \|v\|_{\Sigma}^2 \le \|v\|_{\hat{\Sigma}}^2 \le \frac{3}{2} \|v\|_{\Sigma}^2$$

Proof. Let $\hat{A} \doteq \Sigma^{-\frac{1}{2}} \hat{\Sigma} \Sigma^{-\frac{1}{2}}$ so that $\mathbb{E}\hat{A} = I_d$. Then standard matrix Bernstein bounds (see e.g., [Tropp et al., 2015, Theorem 1.6.2]) imply that, with probability at least $1 - \tau$,

$$\left\|\hat{A} - I_d\right\|_{\text{op}} \le \alpha_{\mu}, \text{ for } \alpha_{\mu} \doteq \sqrt{c \frac{\log(d/\tau)}{n_{\mu}} + c \frac{\log(d/\tau)}{n_{\mu}}} \text{ where } c = c \left(\sup_{X} \left\| \Sigma^{-\frac{1}{2}} X \right\|^2 \right) = c \left(\lambda_{\min}(\Sigma) \right).$$

In other words, we have $(1 - \alpha_{\mu})I_d \preceq \hat{A} \preceq (1 + \alpha_{\mu})I_d$. We obtain the result by multiplying either side of each term by $\Sigma^{\frac{1}{2}}$, and setting n_{μ} sufficiently large so that $\alpha_u \leq \frac{1}{2}$.

We have the following proposition. Notice that, the proposition assumes knowledge of upper-bounds on distributional parameters, namely the subGaussian parameter σ_Y , and a parameter c_{μ} which we assume to be an upper bound on $R_{\mu}(h_{w_*}) - R^*_{\mu}$, where we let R^*_{μ} denotes Bayes risk, and w_* minimizes $R(h_w)$ over the linear class. Thus, if the regression function $\mathbb{E}[Y|x]$ is linear as often assumed, we have that $c_{\mu} = 0$; more generally, if Y itself is bounded, then c_{μ} can be taken as the square of this bound.

Proposition 8. Let \hat{w} minimize $\hat{R}(h_w)$ over $w \in \mathbb{R}^d$, and for any such w, and symmetric and positive $A \in \mathbb{R}^{d \times d}$, define $\|w\|_A^2 = w^\top A w$. Assume $n_\mu \ge C(\tau, d, \operatorname{eigs}(\Sigma))$ as in Lemma 3.

In what follows, c_0 is a universal constant, and c_{μ} depends on μ as explained above. We have that, the following set is a $(26\epsilon, 2\tau, 26)$ -strong confidence set, for $\epsilon = c_0 \cdot \sigma_Y^2 \frac{d+c_{\mu}+\log(1/\tau)}{n_{\mu}}$:

$$\hat{\mathcal{H}}_{\mu} \doteq \left\{ h_w : \|w - \hat{w}\|_{\hat{\Sigma}}^2 \le 6\epsilon \right\}.$$

Proof. We rely on the following two facts (see e.g. Hsu et al. [2012]). First, with probability at least $1 - \tau$, we have that $\mathcal{E}_{\mu}(h_{\hat{w}}) \leq \epsilon$, for some universal c_0 . Second, it is a classifical fact that, for any $w \in \mathbb{R}^d$, $\mathcal{E}(h_{\hat{w}}) = \|w - w_*\|_{\Sigma}^2$, where w_* minimizes $R(h_w)$. Using Lemma 3, we then have that, with probability at least $1 - 2\tau$, the following holds.

- For any $h_w \in \hat{\mathcal{H}}_{\mu}$, we have that

$$\mathcal{E}(h_w) = \|w - w_*\|_{\Sigma}^2 \le 2 \|w - \hat{w}\|_{\Sigma}^2 + 2 \|\hat{w} - w_*\|_{\Sigma}^2 \le 4 \|w - \hat{w}\|_{\hat{\Sigma}}^2 + 2\epsilon \le 26\epsilon.$$

- For any $h_w \in \mathcal{H}_{\mu}(\epsilon)$, we have that

$$\|w - \hat{w}\|_{\hat{\Sigma}}^{2} \leq \frac{3}{2} \|w - \hat{w}\|_{\Sigma}^{2} \leq 3 \left(\|w - w_{*}\|_{\Sigma}^{2} + \|\hat{w} - w_{*}\|_{\Sigma}^{2}\right) \leq 6\epsilon.$$

Remark 14. Notice that, by the above arguments, we can also obtain a $(26\epsilon', 2\tau, 26)$ -strong confidence set, for $\epsilon' = 26\epsilon$ simply by plugging in ϵ' wherever ϵ appears in the definition of $\hat{\mathcal{H}}_{\mu}$.

Remark 15 (Efficient Implementation). It is easily seen that, with the instantiation of strong confidence sets expressed in Proposition 8, our methods used in Theorems 1 and 3 can be implemented efficiently via formulation as convex programs with quadratic constraints.

References

- M. Anthony and P. Bartlett. Neural Networks Learning: Theoretical Foundations. Cambridge University Press, 1999.
- Jean-Yves Audibert and Alexandre B Tsybakov. Fast learning rates for plug-in classifiers. *The Annals of Statistics*, 35 (2):608–633, 2007.
- P. Bartlett, M. I. Jordan, and J. McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006.
- Shai Ben-David, John Blitzer, Koby Crammer, Alex Kulesza, Fernando Pereira, and Jennifer Wortman Vaughan. A theory of learning from different domains. *Machine Learning*, 79(1-2):151–175, 2010.
- Corinna Cortes, Mehryar Mohri, and Andrés Munoz Medina. Adaptation based on generalized discrepancy. *Journal of Machine Learning Research*, 20:1–30, 2019.
- Jiawei Ge, Shange Tang, Jianqing Fan, Cong Ma, and Chi Jin. Maximum likelihood estimation is all you need for well-specified covariate shift. *arXiv preprint arXiv:2311.15961*, 2023.
- S. Hanneke, S. Kpotufe, and Y. Mahdaviyeh. Limits of model selection under transfer learning. In *Proceedings of the* 36th Annual Conference on Learning Theory, 2023.
- Steve Hanneke and Samory Kpotufe. On the value of target data in transfer learning. In Advances in Neural Information Processing Systems, 2019.
- Steve Hanneke and Samory Kpotufe. A no-free-lunch theorem for multitask learning. *The Annals of Statistics*, 50(6): 3119–3143, 2022.
- Roger A. Horn and Charles R. Johnson. Matrix Analysis. Cambridge University Press, 1985.
- Daniel Hsu, Sham M. Kakade, and Tong Zhang. Random design analysis of ridge regression. In *Proceedings of the* 25th Annual Conference on Learning Theory, 2012.
- Jiayuan Huang, Arthur Gretton, Karsten M Borgwardt, Bernhard Schölkopf, and Alex J Smola. Correcting sample selection bias by unlabeled data. In *Advances in neural information processing systems*, pages 601–608, 2007.
- V. Koltchinskii. Local Rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6):2593–2656, 2006.
- E. Mammen and A. B. Tsybakov. Smooth discrimination analysis. The Annals of Statistics, 27(6):1808–1829, 1999.
- Yishay Mansour, Mehryar Mohri, and Afshin Rostamizadeh. Domain adaptation: Learning bounds and algorithms. In *Proceedings of the 22nd Conference on Learning Theory*, 2009.
- P. Massart and É. Nédélec. Risk bounds for statistical learning. The Annals of Statistics, 34(5):2326–2366, 2006.
- Mehryar Mohri and Andres Munoz Medina. New analysis and algorithm for learning with drifting distributions. In *International Conference on Algorithmic Learning Theory*, 2012.
- Mohammadreza Mousavi Kalan, Zalan Fabian, Salman Avestimehr, and Mahdi Soltanolkotabi. Minimax lower bounds for transfer learning with linear and one-hidden layer neural networks. *Advances in Neural Information Processing Systems*, 33:1959–1969, 2020.
- Ievgen Redko, Amaury Habrard, and Marc Sebban. Theoretical analysis of domain adaptation with optimal transport. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, 2017.
- Norbert Sauer. On the density of families of sets. Journal of Combinatorial Theory, Series A, 13(1):145–147, 1972.
- Jian Shen, Yanru Qu, Weinan Zhang, and Yong Yu. Wasserstein distance guided representation learning for domain adaptation. In 32nd AAAI Conference on Artificial Intelligence, 2018.
- M. Talagrand. Sharper bounds for Gaussian and empirical processes. The Annals of Probability, 22(1):28–76, 1994.
- Joel A Tropp et al. An introduction to matrix concentration inequalities. *Foundations and Trends*® *in Machine Learning*, 8(1-2):1–230, 2015.
- Alexandre B Tsybakov. Introduction to Nonparametric Estimation. Springer, 2009.
- V. Vapnik and A. Chervonenkis. On the uniform convergence of relative frequencies of events to their expectation. *Theory of Probability and its Applications*, 16:264–280, 1971.
- Vladimir Vapnik and Alexey Chervonenkis. Theory of Pattern Recognition. Nauka, 1974.
- Wen Zhang, Lingfei Deng, Lei Zhang, and Dongrui Wu. A survey on negative transfer. IEEE/CAA Journal of Automatica Sinica, 10(2):305–329, 2022a.
- Xuhui Zhang, Jose Blanchet, Soumyadip Ghosh, and Mark S Squillante. A class of geometric structures in transfer learning: Minimax bounds and optimality. In *International Conference on Artificial Intelligence and Statistics*, pages 3794–3820. PMLR, 2022b.