# Foundations of ghost stability

Verónica Errasti Díez \*1,2,3, Jordi Gaset Rifà <sup>†3</sup>, and Georgina Staudt <sup>‡4,5</sup>

<sup>1</sup> Universitäts-Sternwarte, Fakultät für Physik, Ludwig-Maximilians-Universität München, Scheinerstraße 1, 81679

München, Germany

<sup>2</sup> Excellence Cluster ORIGINS, Boltzmannstraße 2, 85748 Garching, Germany

<sup>3</sup> Department of Quantitative Methods, CUNEF Universidad, Calle Pirineos 55, 28040 Madrid, Spain

<sup>4</sup> Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Boltzmannstraße 8, 85748 Garching, Germany

<sup>5</sup> Arnold-Sommerfeld Center for Theoretical Physics, Theresienstraße 37, 80333 München, Germany

#### Abstract

We present a new method to analytically prove global stability in ghost-ridden dynamical systems. Our proposal encompasses all prior results and consequentially extends them. In particular, we show that stability can follow from a conserved quantity that is unbounded from below, contrary to expectation. Novel examples illustrate all our results. Our findings take root on a careful examination of the literature, here comprehensively reviewed for the first time. This work lays the mathematical basis for ulterior extensions to field theory and quantization, and it constitutes a gateway for inter-disciplinary research in dynamics and integrability.

### 1 Motivation

Higher-order theories enjoy a long, fruitful history. Archetypal examples include the Pais-Uhlenbeck oscillator [1], Podolsky electrodynamics [2–4] and the gravitational models by Weyl [5,6] and Lovelock [7]. Interest in higher-order settings is heterogeneous, to say the least. For illustration purposes, we mention geometry [8–12], ultraviolet regularization [13–17], cosmology [18–22] and string theory [23–27]. Recent constructions of novel higher-order settings [28–34] evince topicality and augur continued zeal.

The so-called Ostrogradsky theorem [35, 36] states that higher-order Lagrangians generically correspond to Hamiltonians that are not bounded from below. Such unbounded scenarios lie beyond the scope of the Lagrange-Dirichlet theorem [37, 38], wherein strict minima of the energy correspond to stable points of the dynamics, in the Lyapunov sense [39] — definition L2 later on. For a field theoretical treatment, see [40].

In pursuit of stability within the higher-order arena, the most heavily transited way around the hurdle consists in constraining the equations of motion so as to restore a lower bound on the Hamiltonian. Hefty efforts have been made in this regard [41–47] with remarkable outcomes, particularly in Modified Gravity. Influential representatives encompass f(R) theories [48,49] and de Rham-Gabadadze-Tolley (dRGT) massive

<sup>\*</sup>veronica.errasti@cunef.edu

<sup>&</sup>lt;sup>†</sup>jordi.gaset@cunef.edu

<sup>&</sup>lt;sup>‡</sup>georgina@mpp.mpg.de

gravity [50]. Nonetheless, recall that caution must be exercised in such serpentine scenarios [51, 52]. We refer to [53] for a pedagogical presentation.

A prevalent misconception hampers alternative progress. The fallacious Ostrogradsky instability affirms that the absence of a lower bound for the Hamiltonian entails the instability of the dynamical system. This is in direct contradiction with a substantial amount of counterexamples, e.g. [54-60]. For their clarity and vehemence against the faulty identification of Ostrogradsky theorem with instability, we advert to [60-62]. In this work, we shall exploit the breach and investigate the classical stability of systems whose Hamiltonian is manifestly not bounded from below. For concision, we shall employ the widespread nickname ghost systems.

Before proceeding ahead and for completeness, we register the analogy between the higher-order and nonlocal frameworks. To a lesser but far from negligible extent, nonlocal theories share the features of being well-established [63–65], possessing multi-disciplinary scope [66–69] and deserving persistent attention [70]. A prominent exemplar is the Wess-Zumino-Witten model [71–73]. Nonlocal theories tend to be associated with a Hamiltonian that is not bounded from below, which again does not imply their dynamical instability. The scrutiny of nonlocal theories calls for tailored techniques beyond their loose understanding as infinite-order settings, see [74,75] and references therein. Hence, we shall not expressly consider them. Notwithstanding, the essence of our account and conclusions apply therein.

**Organization.** The paper is arranged as follows. In section 2, we classify the notions of classical stability that have been put forward for ghost systems. Additionally, we provide a succinct yet all-encompassing overview of the literature. In section 3, we formalize the global subset of the said classical stability notions. With a focus on one such formalization, in section 4 we expose the mathematical foundation of known stable ghost systems, and develop a framework for novel settings, all of which are exemplified in the subsequent section 5. We summarize and draw our conclusions in the final section 6. Tangential technicalities are consigned to appendices A-D.

# 2 State of the art: multifacetedness of ghost stability

Granted that ghost systems can be stable, the question arises as to what is meant by their stability. Indeed, beyond the misreckoning of unbounded Hamiltonians as necessarily yielding unstable dynamics, we identify a secondary hindrance in the multiplicity of stability notions currently in use for such systems. Accordingly, we proceed to their categorization. We restrict attention to the classical realm.

Global notions of stability. These apply to all solutions, irrespective of initial conditions.

- G1. Lagrange stability [37,76]. Solutions are bounded.
- G2. Benign ghost [55,77]. Solutions are either bounded or diverge only as time tends to infinity (runaway). This excludes solutions that diverge within finite time (blow ups).

Local notions of stability. These apply to some solutions, characterized by initial conditions.

L1. Island of stability [78, 79]. Local counterpart to either of the above global notions.

L2. Lyapunov stability [39] and generalizations thereof. Notions of stability in the immediate vicinity of a critical point of the system.

By definition,  $G1 \subset G2$ . We remark that, given a dynamical system with free parameters and/or functions, any fixed notion of stability is typically attained only within a certain parametric and/or functional region.

### 2.1 Main classical results

For intelligibility and contextualization, we here provide a necessarily abridged recount of the cardinal outcomes within each stability notion. As a preliminary observation, we note that all notions originate and are employed beyond our targeted ghost systems.

G1 studies flourished motivated by engineering applications wherein Lyapunov stability L2 is overrestrictive [80]. In particular, G1 stability is a befitting property of dynamical systems intended to model situations where variables need not be kept within a stringent tolerance for an acceptable performance. As regards ghost systems, G1 is a prolific entry in our stability dictionary. Kaparulin and coauthors have developed a vast bodywork, with onset [81] and culmination [57], encompassing [56,82–93]. These works prove G1 stability in numerous mechanical and field theoretical settings of patent physical relevance and complexity. Central to the clearly prescribed yet not fully systematized construction of such theories is their bi-Hamiltonian character. (Bi-Hamiltonian systems were pioneered by Magri [94]. A cogent introduction can be found in chapter 7.3 of [95].) In broad strokes, the expounded frameworks possess a ghost-like Hamiltonian, but are also associated with a non-canonical Hamiltonian that is bounded from below. The latter warrants G1 stability. The method requires such a compendium of attributes from a root system and proves its preservation in non-trivially extended scenarios. For an articulate summary, we recommend the introduction in [96]. On the other hand, Deffayet et al. [59,97] have shown G1 stability for a broad class of mechanical integrable systems. Along these lines, Heredia and Llosa have incorporated further examples in their second version of [60].

In certain settings, G2 stability can be assimilated to the completeness of the Hamiltonian vector field — see the discussion below definition 2 in the subsequent section 3. In this sense, G2 is a classic subject of study in differential geometry, e.g. [98,99]. Applied to ghost systems, G2 stability has been the goal of significant sundry efforts by Smilga and collaborators. Prefatory studies [78,79,100–103] ultimately led to G2 stable supersymmetric models: mechanical at the outset [77] and field theoretical later [55]. A lucid chronicle is [61], while forefront results are conveyed in [58]. Therein, G2 stable systems are systematically obtained from non-ghost G1 stable systems, without appeal to supersymmetry. The construction incorporates [55,77] as subcases.

(The kind of) Global stability in some systems awaits future advancements. In pursuit of G2 stability, Smilga suggested a proliferation of examples [104], including the renowned Toda chain [105] and the Korteweg-de Vries (KdV) system [106], with time and space interchanged. Both Toda and especially KdV are epitomic instances of bi-Hamiltonian systems. In spite of ulterior progress [29, 58, 107] in relation to time-space flipped KdV and certain modifications thereof, G1 and/or G2 stability remain inconclusive.

As local correlative to G1, ghostly L1 stability counts two apogees. First, in an ambitious generalization of their frame, Kaparulin-Lyakhovic-Nosyrev [96] undertake the stabilization of fully unbounded from below bi-Hamiltonian root systems via interacting augmentations. As a result, they prove L1 stability starting with two such root systems: the resonant Pais-Uhlenbeck oscillator and zero mass Podolsky electrodynamics.

This scheme does not generalize to G1 stability, at least not straightforwardly. Second, following the thread of [58, 104, 107], Fring-Taira-Turner [29] have recently proven the oscillatory behavior of a number of modified and time-space swapped KdV systems, for specific initial conditions. We already indicated that G1 stability persists an enticing open question. Complementarily, as local incarnation of G2 stability, L1 stability has made an appearance in a variety of cosmological contexts benefitting from ghost states, e.g. [108–112]. We single out [78, 79] as decisive for our present discussion, since they triggered the very notion of G2 stability.

Lyapunov stability L2 and refinements thereof, particularly its asymptotic variant, are vertebral to dynamical systems. Within scope, we have been able to trace back L2 studies in ghost systems to [54]. The latest contribution [60] includes proofs of L2 stability for nonlocal mechanical systems.

Numerical studies [113–117] have been instrumental to the development of the above reviewed analytic results and continue to motivate enticing questions [118].

#### 2.2 Comments on quantization

Although beyond scope, for completeness we here provide a lightning recap on quantum ghost settings.

Sustained consensus identifies unitarity as the paramount feature for quantum viability and the existence of a well-defined ground state as the embodiment of quantum stability. A priori in clash, the canonical quantization of generic ghost settings involves a dichotomy: negative norm states or a spectrum that is not bounded from below [29,55,61,77,119–124]. Early works, such as Stelle's influential proof of renormalizability for fourth-order quantum gravity [13] or its reconsideration in [119], tend to focus on the reconciliation of unitarity and negative norm states. Recently, the study of negative eigenvalues has gathered attention, as emphatically defended in [123].

In this regard, Smilga [61, 121], also with Robert [77], surmounts the paradox in certain cases, proving unitarity of the evolution operator while minutely characterizing the unorthodox, unbounded from below spectrum. Their analyses heavily rely on solvability, direct or by means of Liouville integrability via a non-trivial change of coordinates to action-angle variables. With a solid basis on the Pais-Uhlenbeck oscillator [121] and their propounded G2 stable mechanical system [77], a remarkable conjecture is put forward: global stability of a ghost system ensures its quantum viability. More precisely, both G1 and G2 stability warrant quantum unitarity, with pure point and band-wise continuous dense spectra, respectively.

The exception to the rule are [124] and the very recent [125], where negative norm states are shown to be compatible with unitarity. In [124], the approach is mechanical in nature. It calls for coordinate operators that are odd under time reflection and also for a self-adjoint yet not Hermitian Hamiltonian. See also [126]. In [125], a new inner product is proposed which, used in the Born rule, provides a sensible probability interpretation — in particular, respecting unitarity. This is worked out in detail in mechanical scenarios and toy-models that support its validity in four dimensional field theory. The focus is on classically unstable systems, which aligns well with Smilga's conjecture. In this regard, see [127] and references therein.

Notwithstanding, most efforts center around the outright cancellation of the dichotomy. We highlight three such possibilities. By construction, the bi-Hamiltonian theories considered by Kaparulin and friends admit canonical quantization of a non-canonical Hamiltonian bounded from below, with respect to appropriate Poisson brackets. In this manner, positive norm states and a ground state are obtained. The technique was developed in [128–130].

Besides, it has been suggested that the Pais-Uhlenbeck oscillator be viewed as invariant under Parity and Time reflection (PT) and the apposite non-canonical quantization scheme applied [131–133]. (An instructive recapitulation of PT quantum mechanics appeared not long ago [134].) Yet again, the result is positive norm states and a ground state. On the not so bright side, PT symmetry is obtained through a rather unnatural analytic continuation and the proposal does not straightforwardly extend to the non-trivial interacting scenario [121]. To the best of our knowledge, such criticisms have never been addressed.

Lastly, a popular circumvention consists in the development of ghost confinement mechanisms, e.g. [135–140]. These techniques remove the modes responsible for the bottom unboundedness of the Hamiltonian from the physical spectrum, including the classical regime.

# 3 Definitions: the big picture of global stability

Additional progress in the above reviewed subject of stable ghost systems necessitates, or at least greatly benefits from, a rigorous comprehension of the various stability notions at play. Therefore, this section is devoted to the formalization of the comparatively under-examined global stability notions G1 and G2. Local stability L1 is then recovered for particular sets of solutions, characterized by initial conditions. Our work thus complements the long-standing formalization of Lyapunov stability L2 and plentiful generalizations thereof. We underscore the generality of the forthcoming discussion, which shall be explicitly applied to (certain) ghost systems in section 5. The domain and interrelation of our proposed definitions are succinctly conveyed in figure 1.

We set our scope to dynamical systems. We require their generic characterization, independent of a Lagrangian and/or Hamiltonian. The rationale is three-fold. First, this entails the same level of abstraction employed by Lyapunov. Second, numerous dynamical systems exist for which a standard Lagrangian and/or Hamiltonian cannot be defined. Well-known examples include the Lorenz system [141] and nonlocal actions [60]. Third, we thus remove potential ambiguities that may arise in bi-Hamiltonian systems [121], such as the extensively researched Pais-Uhlenbeck oscillator [1]. Accordingly, we consider the set of solutions:

$$Sol = \{\gamma : I_{\gamma} \subset \mathbb{R} \to \mathcal{P} \text{ continuous}\},\tag{3.1}$$

for some topological space  $\mathcal{P}$ ; henceforth taken to be at least Hausdorff. Here,  $I_{\gamma}$  is a fixed (non-empty) interval of time where the curve  $\gamma$  is defined. For example, given a Lagrangian system,  $\mathcal{P}$  is the tangent space of the configuration space, Sol are the solutions of the Euler-Lagrange equations and  $I_{\gamma}$  is their maximal interval of definition.

In this broad setting, the notion G1 admits two formalizations, depending on  $\mathcal{P}$ . The definition closest to the understanding employed in the ghost literature is:

**Definition 1.a.** For  $\mathcal{P}$  metric, a system is **G1** in the bounded sense if, for all curves  $\gamma \in Sol$ , their image  $\gamma(I_{\gamma})$  is bounded.

Notice that the metric nature of  $\mathcal{P}$  allows for a notion of distance, which in turn renders boundedness meaningful. We here propose the broader notion:

**Definition 1.b.** For  $\mathcal{P}$  Hausdorff, a system is **G1** in the compact sense if, for all curves  $\gamma \in Sol$ , their image  $\gamma(I_{\gamma})$  is relatively compact with respect to  $\mathcal{P}$ .

Recall that a subset U of a topological space  $\mathcal{P}$  is relatively compact with respect to  $\mathcal{P}$  if its closure  $\overline{U}$  is compact. Moreover, for  $\mathcal{P}$  Hausdorff, U is relatively compact with respect to  $\mathcal{P}$  iff it is inside a compact



Figure 1: Visualization of the scope of global stability definitions 1.a, 1.b and 2, together with relations between them. These formalize the preexisting notions G1 and G2.

set. (The definition of Hausdorff, relatively compact and allied concepts and properties can be found in e.g. [142], see 17.15.)

Two important observations follow. First, all metric spaces are Hausdorff, but the converse is not true, see e.g. 11.5 in [143]. This implies that the scope of definition 1.b is larger than that of definition 1.a. Second, for  $\mathcal{P}$  metric (and hence Hausdorff), definitions 1.a and 1.b are not equivalent. In this case, definition 1.b implies definition 1.a, but the converse is not true. Equivalence is achieved by requiring the Heine-Borel property for  $\mathcal{P}$  [144], which states that any closed and bounded subset of  $\mathcal{P}$  is compact. For a counterexample, the renowned Riesz's lemma shows that the closed unit ball of infinite-dimensional normed vector spaces is not compact — see e.g. 5.2 in [145].

Beyond its greater scope, section 4 (particularly our main result theorem 3) unequivocally leads to our prevailing of definition 1.b over definition 1.a. More generally, it is essential to keep watch on metric  $\mathcal{P}$ 's without the Heine-Borel property, since they are physically relevant. Conspicuous instances include infinite-dimensional Banach and Hilbert spaces (as particular exemplars of infinite-dimensional normed vector spaces) and geodesically incomplete Riemannian manifolds; for example,  $\mathbb{R}^n \setminus \{0\}$  with the Euclidean metric. In all such scenarios, it becomes indispensable to pick out between definitions 1.a and 1.b.

Moving on to the stability notion G2, we observe that this is a weaker version of G1. Specifically, G2 accepts solutions that diverge at infinite time. For any such  $\gamma \in Sol$ , the image  $\gamma(I_{\gamma})$  is not inside a compact. Nevertheless, G2 excludes solutions that diverge at a finite time. In order to detect these  $\gamma$ 's and banish them from the formal definition of G2, we follow the work of Palais [99] and probe spans of time that are relatively compact with respect to  $\mathbb{R}$ , instead of probing the whole time range. Accordingly, we put forward:

**Definition 2.** For  $\mathcal{P}$  Hausdorff, a system is **G2** if, for all curves  $\gamma \in \text{Sol}$ , and for all subsets  $S \subset I_{\gamma}$  relatively compact with respect to  $\mathbb{R}$ , the image  $\gamma(S)$  is relatively compact with respect to  $\mathcal{P}$ .

If Sol is the set of maximal integral curves of a smooth<sup>1</sup> vector field X, then by negation of the Escape Lemma (see, e.g. [146]), the system is G2 iff X is complete. This is the case of regular Lagrangian and/or

<sup>&</sup>lt;sup>1</sup>Classical texts consider X smooth. For the purists, we note it is enough to require that X is  $C^1$ .

Hamiltonian systems. Bear in mind that a vector field is complete if its integral curves can be defined for all times:  $I_{\gamma} = \mathbb{R}$  in (3.1).

It is obvious that definition 1.b implies definition 2, but the converse is not true. Moreover, for  $\mathcal{P}$  metric, definition 1.a does not imply 2.

All our proposed definitions 1.a, 1.b and 2 are manifestly intrinsic: they do not appeal to coordinates.

## 4 Proving G1: the pivotal role of conserved quantities

Henceforth, we concentrate on global stability G1, in the compact sense of definition 1.b. In contraposition to G2 global stability in the formal sense of definition 2, our prevailed conceptualization a priori prevents the divergence of wave functions and probability distributions upon quantization, which in principle span the whole configuration space. This argument was also presented in [59, 122] and is reinforced by calculations [61,77,121] that signal a comparatively less unconventional quantum structure. In contraposition to G1 global stability in the bounded sense of definition 1.a, our prevailed conceptualization enjoys a broader scope in its definition (all Hausdorff spaces) and always implies G2 global stability.

In this section, we investigate a suitable tool kit for the implementation of definition 1.b in systems of the form (3.1). We begin by briefly expounding the mathematical results that support such G1 stability proofs and then proceed to their unprecedented generalization. To this aim, the following comprehension of a conserved quantity is employed hereafter.

**Definition 3.** Let N be a topological space. A continuous function  $Q : \mathcal{P} \to N$  is a **conserved quantity** of Sol if, for any fixed curve  $\gamma \in$  Sol, the value of Q is unchanged; that is,  $Q(\gamma(t_1)) = Q(\gamma(t_2))$  for all points  $t_1, t_2 \in I_{\gamma}$ .

Observe that our reference definition 1.b is topological in nature. Befittingly, definition 3 brings the concept of conserved quantity to the same level of abstraction.

### 4.1 When does energy work?

A well-known resolution to the interrogation is as follows.

**Theorem 1.** Given a system Sol with  $\mathcal{P} = \mathbb{R}^n$ , if the energy  $E : \mathbb{R}^n \to \mathbb{R}$  is a twice differentiable conserved quantity that has a minimum and whose Hessian is positive definite everywhere, then the system is G1.

Since  $\mathbb{R}^n$  is metric and has the Heine-Borel property, the system is G1 in the sense of both equivalent definitions 1.a and 1.b. The converse of theorem 1 is not true, as abundantly exemplified in section 2.

For completeness, a recount of the proof is provided in appendix A. (Alternatively, theorem 1 can be viewed as a corollary of theorem 2, also proved in appendix A.) It is beneficial to bear in mind that the proof relies on two facts. First, solutions  $\gamma \in Sol$  are inside level sets of the energy E because the latter is a conserved quantity. Second, the level sets of E are bounded, as a direct consequence of E having both a minimum and a positive definite Hessian everywhere — the very definition of which necessitates the auxiliary property that E is twice differentiable. To avoid confusion stemming from polysemy, we note that we abide by the understanding of level set of a function f as the set of points in its domain for which ftakes a fixed constant value. Expressly, for  $f: M \to N$  and  $c \in N$ , the level set is

$$f^{\neg}(c) := \{ x \in M \mid f(x) = c \}.$$
(4.1)

Notice that theorem 1 does not ensure G1 stability of the system Sol if either of the two requirements on the conserved, twice differentiable energy E is not met. Namely, if either E has a minimum but its Hessian is not positive definite somewhere or if E does not attain a minimum in spite of having a positive definite Hessian everywhere, then the axioms of theorem 1 do not apply. Respective counterexamples are the Hamiltonian systems on  $\mathbb{R}^2$ 

$$H = \frac{1}{2}(p-x)^2, \qquad H = \exp(x^2 + p), \tag{4.2}$$

which can be easily verified to each admit a solution that is not relatively compact with respect to  $\mathbb{R}^2$ :

$$\begin{cases} x = t, \\ p = t + 1, \end{cases} \qquad \begin{cases} x = t, \\ p = -t^2. \end{cases}$$
(4.3)

We remark that (4.2) illustrate the fact that (in)stability is a feature of generic dynamical systems, including but transcending our targeted ghost systems. (As quantum correlative, it is interesting to mention that unitarity loss can occur in ghost-free settings [147].)

### 4.2 The case of ghost systems

An enticing overspill of the hypotheses of theorem 1 is provided by (mechanical) ghost systems, whose imprint is an energy that is not bounded from below. Close inspection of the apposite proof in appendix A readily reveals that any conserved quantity Q, not necessarily the energy E, can be employed in the theorem. This cognizance underlies the great majority of G1 stable ghost systems known to date, developed by Kaparulin et al.  $[56, 57, 81-93, 96]^2$ .

The remaining results, those of Deffayet and friends [59,97] and more recently by Heredia and Llosa [60], consider a more general kind of conserved quantity: coercive instead of positive definite. For self-containment, we recall:

**Definition 4.** A function  $f : \mathbb{R}^n \to \mathbb{R}$  is coercive if the absolute value of f(x) diverges as the modulus of x diverges:

$$\lim_{\|x\|\to+\infty} |f(x)| \to +\infty.$$
(4.4)

(En passant, we point out that coercive functions can in fact be defined for arbitrary metric spaces, see lemma 2 in appendix A. Upon demanding the Heine-Borel property, coercive functions are identified with proper functions. The double implication is established in appendix B, and its relevance is discussed.) Using the above, we proceed to formalize the theorem implicitly in force in [59,97], which is a restatement of theorem 5 in the second version of [60].

**Theorem 2.** Given a system Sol with  $\mathcal{P} = \mathbb{R}^n$ , if there exists a conserved quantity  $Q : \mathbb{R}^n \to \mathbb{R}$  that is coercive, then the system is G1.

 $<sup>^{2}</sup>$ Specifically and as already heeded, these works avail themselves of a Q that is another, inequivalent Hamiltonian of the system.

As in theorem 1 before, G1 refers to both equivalent definitions 1.a and 1.b. The converse of theorem 2 is not true; a counterexample being the renowned Kronecker foliation of the torus with irrational slope<sup>3</sup>, see e.g. 19.18 in [146].

The proof is relegated to appendix A. It is worth highlighting that theorem 2 is a corollary of theorem 3 later on, whose proof is deferred to section 4.3.

A weighty remark is due. Given a fixed dynamical system, the unveiling of conserved quantities is widely recognized to be a non-trivial task. This recalcitrant open problem has recently attracted the efforts of the machine learning community, with noteworthy success in specific settings [148, 149], including the symbolic (i.e. analytic) characterization of some of the thusly disclosed conserved quantities [150]. Furthermore, given a set of (non-trivial, functionally independent) conserved quantities for a system, it is arbitrarily difficult to prove their exhaustiveness. Arnold-Kozlov-Neishtadt insightfully revisit six approaches to tackle such an ambitious goal [151].

In the ghost system literature, there exist four proposals to surmount the hardships of conserved quantity revelation. Smilga has put forward a systematic variational approach to generate multi-variable conserved ghost Hamiltonians starting from a single-variable conserved Hamiltonian, see [58] and (self-)references therein. Additionally, he suggested [107] appropriate modifications on the Korteweg-de Vries (KdV) system [106] for its ghostly conversion, later widened in [29]. Kaparulin and coauthors have developed a methodology to extend their targeted bi-Hamiltonian theories in a manner that the original conserved quantities are preserved [56, 81, 87]. Lastly, Deffayet et al. consider integrable ghost systems resulting from the application of complex canonical transformations on standard integrable Hamiltonian systems [59]. Certainly, one may always attempt to pull through direct computation, especially in propitious settings and equipped with a simplifying ansatz [60], as we do to obtain (5.3).

#### 4.3 Main result: generalization of the method

In light of definitions 1.b and 3, critical examination of theorems 1 and 2 leads to the following fertile conclusion. Consider a system (3.1) possessing a conserved quantity Q. For such a system to be G1 in the compact sense, it is sufficient that each path-connected component of a level set of Q is relatively compact with respect to  $\mathcal{P}$ .

To grasp the meaning and value of our inference, let us take a moment to examine a purposeful example. Let  $\mathcal{P} = \mathbb{R}^2$  and let Sol be a system of the form (3.1) with the conserved quantity

$$Q = \sin(u^2 + v^2). \tag{4.5}$$

The level set

$$Q^{\neg}(0) = \left\{ (u, v) \in \mathbb{R}^2 \,|\, u^2 + v^2 = k\pi, \, k \in \mathbb{N} \cup \{0\} \right\}$$
(4.6)

consists of concentric circles in  $\mathbb{R}^2$ . Each of these circles is a path-connected component of  $Q^{\neg}(0)$ . In fact, for each  $c \in [-1,1]$ , the level set  $Q^{\neg}(c)$  is made up of concentric circles. Considered together, the set of level sets  $\{Q^{\neg}(c)\}$  covers  $\mathbb{R}^2$ . Now, since all curves  $\gamma \in$  Sol are continuous, the image of each such curve

<sup>&</sup>lt;sup>3</sup>This consists of curves winding on the torus, the image of each curve being dense on the torus. As such, the closure of each curve is the whole torus, which is compact. Thus, the system is G1 in the compact sense of definition 1.b. However, the system does not have a non-trivial conserved quantity in the sense of definition 3. To see this, recall that a conserved quantity has constant value on the image of each fixed curve. As noted, this is a dense subset on the torus. For a continuous conserved quantity, this entails a constant value on the whole torus, rendering the conserved quantity trivial.

 $\gamma(I_{\gamma})$  can only be contained within one of the circles (i.e. path-connected components) of the appropriate level set  $Q^{\neg}(c)$ . Even though  $Q^{\neg}(c)$  is not relatively compact with respect to  $\mathbb{R}^2$  (it contains points with arbitrarily large modulus), the contemplated system is G1 in the compact sense because each path-connected component of  $Q^{\neg}(c)$  is compact — thus, relatively compact — with respect to  $\mathbb{R}^2$ .

The above indicated and illustrated sufficient criterion for the enforcement of definition 1.b on a system Sol motivates our putting forward the subsequent definition.

**Definition 5.** Let  $\mathcal{P}$  be a Hausdorff space and let  $\mathcal{S}$  be a set. A function  $f : \mathcal{P} \to \mathcal{S}$  is confining if the path-connected components of its level sets are relatively compact with respect to  $\mathcal{P}$ .

To the best of our knowledge, the above is a novel concept. It generalizes coercive (proper) functions, in the appropriate setting, as shown in proposition 1 (3) of appendix A (B). Its interest is justified by our main result:

**Theorem 3.** Given a system Sol with  $\mathcal{P}$  Hausdorff, if there exists a conserved quantity  $Q : \mathcal{P} \to N$  that is confining, then the system is G1 in the compact sense.

Recall that G1 in the compact sense was defined in 1.b. Notice that definition 3 entails that N in theorem 3 must be some topological space. The converse of theorem 3 is not true. A counterexample is again provided by the Kronecker foliation of the torus with irrational slope, see e.g. 19.18 in [146] and footnote 3.

Our theorem 3 can be demonstrated as follows.

Proof. Consider an arbitrary but fixed curve  $\gamma \in \text{Sol}$  and a point  $t_0 \in I_\gamma$ . Let  $q_0 \equiv Q(\gamma(t_0))$ . Since Q is a conserved quantity, for all  $t \in I_\gamma$ , the equality  $Q(\gamma(t)) = q_0$  holds true. Therefore, the image of the curve  $\gamma$  is contained in a level set of  $q_0: \gamma(I_\gamma) \subset Q^{\neg}(q_0)$ . Moreover, because the curve  $\gamma$  is continuous, it must be inside a path-connected component of  $Q^{\neg}(q_0)$ . Let A denote such a component. Since Q is confining, A is relatively compact with respect to  $\mathcal{P}$ . Hence, its closure  $\overline{A}$  is compact. Consequently,  $\gamma(I_\gamma)$  is inside a compact. As  $\mathcal{P}$  is Hausdorff,  $\gamma(I_\gamma)$  is relatively compact.

We stress that our theorem 3 provides a generalization to theorem 2, which in turn extends theorem 1. The concatenation is vertebrated by the following doublet of implications. First, a function  $f : \mathbb{R}^n \to \mathbb{R}$  that is twice differentiable satisfies

f has a minimum and positive definite Hessian everywhere  $\Rightarrow$  f has a minimum and is coercive. (4.7)

Second, a function  $f : \mathbb{R}^n \to \mathbb{R}$  fulfills

$$f \text{ is coercive } \Rightarrow f \text{ is confining.}$$
(4.8)

The first (second) implication is proven in lemma 1 (proposition 1) of the appendix. The converses are not true. Ghostly counterexamples can be found in the next section 5: see (5.26) and (5.31), respectively. Also, see figure 2.

Before advancing to concrete families of ghost systems that manifest the non-triviality of the stepwise progression (4.7)-(4.8), we comment on an often overlooked yet physically germane relation: that between the protagonist conserved quantity Q and the symmetries of the system Sol.

#### 4.4 Relation to symmetries

All methods known to date that prove G1 stability (in the sense of definition 1.b) in a given ghost system use a conserved quantity. The symmetries associated with such G1-certifying conserved quantities are generically not natural. This is true for the examples in [59, 60, 97], as well as our examples in the subsequent section 5. Given that unnatural symmetries come about sporadically in the physics literature, we here provide a lightning introduction, based on the all-encompassing recent book [152] on the geometry of mechanical systems.

A dynamical symmetry of a system Sol is defined as a diffeomorphism  $\Psi : \mathcal{P} \to \mathcal{P}$  mapping solutions to solutions. The subset of dynamical symmetries related to an (at least  $C^2$ ) conserved quantity via Noether's theorem are called Noether symmetries. For regular Lagrangian systems, such as all aforementioned ones, a Noether symmetry leaves both the Lagrangian energy and the symplectic form invariant, but not necessarily the Lagrangian.

On the other hand, let  $\mathcal{P}$  be the tangent space of the configuration space  $\mathcal{P} = T\mathcal{C}$ . A diffeomorphism  $\Psi: T\mathcal{C} \to T\mathcal{C}$  is said to be natural if it consists of a transformation in positions that is canonically lifted to velocities:

$$\Psi(\dot{x}^{i}, x^{i}) = (\dot{x}^{\prime i}(\dot{x}^{i}, x^{i}), x^{\prime i}(\dot{x}^{i}, x^{i})), \qquad \text{such that} \qquad x^{\prime i}(\dot{x}^{i}, x^{i}) = x^{\prime i}(x^{i}) \quad \text{and} \quad \dot{x}^{\prime i} = \frac{\partial x^{\prime i}}{\partial x^{j}} \dot{x}^{j}. \tag{4.9}$$

~ *li* 

A Noether symmetry with the naturalness property leaves the Lagrangian invariant (up to a divergence). The converse is also true: a natural symmetry that leaves the Lagrangian invariant (up to a divergence) is a natural Noether symmetry.

It follows that the Noether symmetries of the G1-enforcing conserved quantity in ghost systems typically present two unconventionalities. First, they entail a transformation of positions involving velocities. Second, they may, or may not, leave the Lagrangian invariant (up to a divergence).

The quintessential example of unnatural symmetries is given by the Laplace-Runge-Lenz vector in the Kepler problem. (Re)Discovered several times, it acquired sizeable notoriety in the early days of quantum mechanics, when Pauli employed it to derive the energy levels of the hydrogen atom [153]. Ulterior works showed that the symmetries associated to the Laplace-Runge-Lenz vector, together with the angular momentum, form a Lie group acting on the phase space taken as a whole — i.e. considering both positions and momenta. The interested reader is referred to the enjoyable text [154] and references therein.

**Relation to G2 and Liouville-Arnold theorem.** Two reminders and a fruitful application follow. First, proving that a system Sol is G1 in the sense of definition 1.b proves that the system is G2 in the sense of definition 2 — recall figure 1. Second, if the system Sol can be described by an (at least  $C^1$ ) vector field X, then G2 stability is equivalent to the completeness of X — summon back the comments below definition 2 and observe this applies to all examples in [59,60,97] and section 5. In this case, G1 implies the completeness of X. This direct consequence is constructive in the context of integrable systems.

We already stressed that finding conserved quantities is generically difficult. Consider nonetheless the auspicious scenario wherein a dynamical system Sol possesses at least  $n = \dim(\mathcal{C})$  functionally independent conserved quantities  $\{Q\}$  in involution. Such a system is said to be *integrable*. The study of integrable systems is a decidedly active area of reasearch on its own [155], often invoking supplementary non-trivial hypotheses from the onset. This has lead to a notable ramification within the field — for a first impression on the bustle, see the start of chapter 2.2.3 in the wonderful book [156]. In our fleeting incursion into the

subject, we opt for following [157] in our understanding of the vertebral Liouville-Arnold theorem [158, 159] and thus require that the infinitesimal symmetries associated to the conserved quantities be complete vector fields.

In an integrable mechanical system proven to be G1 by means of our theorem 3, at least one of the conserved quantities in  $\{Q\}$  is confining. As a result, the hypotheses of the Liouville-Arnold theorem are satisfied. Moreover, the system can be described as free particles moving *exclusively* around torii. Thusly, theorem 3 connects stable ghost systems with the forefront of the field of integrable systems, see [155] and references therein. The particulars and proof of this relation are allocated to appendix C.

### 5 Examples

This section is devoted to illustrating the impactful implications of our previous results, especially definition 1.b and theorem 3, in the particular arena of ghost systems. To achieve this goal, we begin by presenting a two free parameter and one free function family of integrable ghost systems (5.1). The subsequent, staggered demand that the axioms of theorems 1, 2 and 3 be fulfilled by (5.1) yields increasingly vast ghost systems proven to be G1 in the compact sense of definition 1.b. Most remarkably, the G1 ghost systems obtained via the enforcement of theorem 3 are not only new, but also could not have been derived with the techniques in the literature.

Consider the regular Lagrangian system

$$L = \frac{1}{2}(\dot{x}^2 - \dot{y}^2) - \frac{\omega^2}{2}(x^2 - y^2) - f(x + \alpha y),$$
(5.1)

where dot denotes derivation with respect to time, the function  $f : \mathbb{R} \to \mathbb{R}$  is taken to be at least twice differentiable and  $\omega^2, \alpha \in \mathbb{R}$ . We highlight that (5.1) is not an instance of a self-interacting extension of the abundantly studied Pais-Uhlenbeck oscillator [83, 114, 122, 132, 160], owing to the degeneracy in the frequencies of the independent variables (x, y).

Because (5.1) is autonomous, the (Lagrangian) energy

$$E_L = \frac{1}{2}(\dot{x}^2 - \dot{y}^2) + \frac{\omega^2}{2}(x^2 - y^2) + f(x + \alpha y)$$
(5.2)

is an obvious conserved quantity of the system. Furthermore, it is apparent that (5.1) is ghostly: the energy (5.2) does not have a global minimum.

It can be readily verified that the system (5.1) has the functionally independent conserved quantity

$$Q = \frac{1}{2} \left( \dot{x}^2 + \dot{y}^2 + 2c_1 \dot{x} \dot{y} \right) + \frac{\omega^2}{2} \left( x^2 + y^2 + 2c_1 xy \right) + c_2 f(x + \alpha y), \tag{5.3}$$

with<sup>4</sup>

$$c_1 = \frac{2\alpha}{1+\alpha^2}, \qquad c_2 = \frac{1-\alpha^2}{1+\alpha^2}.$$
 (5.4)

We have unveiled (5.3) by direct computation, in a befitting adaptation of the classical results by Darboux [161].

<sup>&</sup>lt;sup>4</sup>The shrewd reader may prefer to let  $\alpha = \tan \theta$ , so that  $c_1 = \sin(2\theta)$  and  $c_2 = \cos(2\theta)$ .

To the best of our knowledge, (5.1) with (5.3) is a novel family of integrable ghost systems. In particular, it cannot be obtained via complex canonical transformations on previously known systems, as suggested in [59]. Besides, it is neither part of the family in equation (34) of [81] nor contained in any of the families I-IV in section 3.4 of [60].

Further properties of (5.1) are discussed in appendix D, including symmetries and its bi-Hamiltonian nature.

As announced, we proceed to carving out various cases from (5.1) that are G1 in the compact sense of definition 1.b. To this aim, we shall employ the results in section 4.

#### 5.1 An application of theorem 1

In order to determine the instances in which the system (5.1) is G1 in the equivalent sense of definitions 1.a and 1.b, we here employ theorem 1. We apply it, not on the energy (5.2) that always fails to meet the hypotheses, but rather on the conserved quantity (5.3). Specifically, we require that (5.3) has a positive definite Hessian everywhere, as well as a minimum.

For starters, we compute the Hessian of (5.3), which is of block-diagonal form

$$\operatorname{Hess}(Q) = \begin{pmatrix} H_1 & 0\\ 0 & H_2 \end{pmatrix}, \quad \text{where} \quad H_1 = \begin{pmatrix} 1 & c_1\\ c_1 & 1 \end{pmatrix}, \quad H_2 = \omega^2 H_1 + c_2 f'' \begin{pmatrix} 1 & \alpha\\ \alpha & \alpha^2 \end{pmatrix}. \quad (5.5)$$

Here, the prime stands for total derivative:  $f'(z) \equiv df/dz$  and so on. As per Sylvester's criterion, (5.5) is positive definite iff all its principal minors have a positive determinant. The first principal minor is evidently positive. For the second principal minor  $H_1$  to have positive determinant, we require that

$$1 - c_1^2 > 0 \implies (\alpha^2 - 1)^2 > 0 \implies \alpha \neq \pm 1.$$
 (5.6)

Provided that (5.6) is fulfilled, the positiveness of the determinant of the third principal minor entails

$$(1+\alpha^2)\omega^2 + (1-\alpha^2)f'' > 0.$$
(5.7)

Besides, the determinant of (5.5) is positive when

$$(1 - c_1^2)\omega^4 + c_2\omega^2(\alpha^2 - 2c_1\alpha + 1)f'' > 0 \implies \omega^2\left(\omega^2 + (1 - \alpha^2)f''\right) > 0.$$
(5.8)

Minor algebraic effort is involved in exhaustively choosing the tightest bound among (5.7) and (5.8). We thus arrive at the necessary and sufficient conditions for the positive definiteness of (5.5):

$$\omega^2 > 0, \qquad f'' \gtrless \frac{\omega^2}{\alpha^2 - 1} \quad \text{for } |\alpha| \le 1,$$
(5.9)

with  $|\alpha| = 1$  forbidden by (5.6). We call attention to the fact that we have derived  $\omega^2 > 0$ , as opposed to taking it as a priori datum.

On the other hand, for (5.3) to have a minimum, we demand

$$\frac{\partial Q}{\partial \dot{x}} = 0, \qquad \frac{\partial Q}{\partial \dot{y}} = 0, \qquad \frac{\partial Q}{\partial x} = 0, \qquad \frac{\partial Q}{\partial y} = 0.$$
 (5.10)

The first two requirements immediately yield  $\dot{x} = 0 = \dot{y}$ . Operating through, the latter two requirements give rise to an additional, necessary and sufficient condition for the existence of a minimum: that there exists a point  $z_0 \in \mathbb{R}$  such that

$$\omega^2 z_0 + (1 - \alpha^2) f'(z_0) = 0.$$
(5.11)

If extant, such  $z_0$  can be readily employed to determine the point at which (5.3) achieves its minimum:

$$x = -\frac{1}{\omega^2} f'(z_0), \qquad y = \frac{\alpha}{\omega^2} f'(z_0).$$
 (5.12)

All in all, (5.9) and (5.11) comprise the necessary and sufficient conditions for Q in (5.3) to have a positive definite Hessian everywhere and a minimum.

**Cases in point.** Let  $n \in \mathbb{N}$  and let  $a, b \in \mathbb{R}$ . Exemplars from the family (5.1) satisfying both (5.9) and (5.11) include

$$f(z) = z^{2n}$$
 for  $|\alpha| < 1$ ,  $f(z) = az + b$ ,  $f(z) = (z^2 + 1)^{1/n}$ , (5.13)

As around (4.2) before, we again call attention to the fact that the simultaneous fulfillment of (5.9) and (5.11) is needed in order to prove G1 stability of a system of the form (5.1). Counterexamples are

$$f(z) = z^{2n}$$
 for  $|\alpha| > 1$ ,  $f(z) = \frac{\omega^2}{1 - \alpha^2} (\exp(z) - z^2/2)$   $\forall \alpha \neq \pm 1.$  (5.14)

These fail to meet (5.9) and (5.11), respectively.

An altogether distinct example of a ghost system that is proven to be G1 as per theorem 1 appeared not long ago, in section 3.3 of [60].

#### 5.2 An application of theorem 2

Next, we ascertain the subset of (5.1) that is G1 in the equivalent sense of definitions 1.a and 1.b with a basis on theorem 2. Consequently, we move forward to imposing that the conserved quantity Q in (5.3) is coercive.

In fact, this is easier said than done. Definition 4 of a coercive function is a usual suspect for finding counterexamples, as opposed to a powerful hammer for forging a proof. Accordingly, we start by restricting the free parameter space of Q, spotting escape directions that are incompatible with its desired coerciveness. First, we note that, for  $\alpha = \pm 1$ , points of the form  $(\dot{x}, \dot{y}, x, y) = (\lambda, \mp \lambda, 0, 0) \equiv \mathbf{p}_{\lambda}$ , with  $\lambda \in \mathbb{R}$  satisfy

$$\lim_{\lambda \to +\infty} |Q(\mathbf{p}_{\lambda})| = |c_2 f(0)| \neq +\infty, \tag{5.15}$$

The above contradicts definition 4 of a coercive function. Thence, in analogy to the preceding section 5.2, we hereafter restrict attention to systems where  $\alpha \neq \pm 1$ . Second, consider

$$\mathbf{q}_{\lambda} \equiv \lambda(-\alpha \sqrt{|\omega^2|}, \sqrt{|\omega^2|}, -\alpha, 1), \qquad Q(\mathbf{q}_{\lambda}) = c_2(1-\alpha^2) \left(\frac{|\omega^2|+\omega^2}{2}\right) \lambda^2 + c_2 f(0). \tag{5.16}$$

If  $\omega^2 \leq 0$ , it is clear that

$$\lim_{\lambda \to \pm \infty} Q(\mathbf{q}_{\lambda}) = c_2 f(0) \neq +\infty.$$
(5.17)

We point out the restrictive nature of (5.17) as compared to definition 4. In detail, we have dropped out the absolute value, thereby forcing  $Q(\mathbf{q}_{\lambda})$  to tend to *plus* infinity as  $\lambda$  diverges. Indeed, this is the only possibility for Q in (5.3) to be coercive, since its kinetic piece is invariably positive definite with a minimum for all  $\alpha \neq \pm 1$  — recall (5.6) and around. (This concept is formally introduced in appendix A: see definition 6 of a *positively coercive* function.) Thus, (5.17) contradicts definition 4. As a result, once more paralleling the preceding section 5.1, we derive the requirement that frequencies be real for Q to be coercive:  $\omega^2 > 0$ .

In the same spirit, we move forward to restricting the free functional space of Q. Let the velocities  $(\dot{x}, \dot{y})$  take a fixed value. In this case, the enforcement of definition 6 on Q stipulates that

$$U(x,y) = \frac{\omega^2}{2} \left( x^2 + y^2 + 2c_1 xy \right) + c_2 f(x + \alpha y)$$
(5.18)

be positively coercive. It is beneficial to perform the change of variables

$$\begin{cases} \zeta = x + \alpha y, \\ \eta = \alpha x + y, \end{cases}$$
(5.19)

so that

$$U(\zeta,\eta) = \frac{\omega^2}{2(1+\alpha^2)}(\zeta^2+\eta^2) + c_2f(\zeta) = P(\zeta,\eta) + c_2f(\zeta).$$
(5.20)

Let  $\eta$  take a fixed value. For later convenience, we single out  $\eta = 0$ . Then, for Q to be positively coercive, it must hold true that

$$\lim_{\zeta \to \pm \infty} P(\zeta, 0) + c_2 f(\zeta) = +\infty, \tag{5.21}$$

which readily leads to  $f(\zeta)$ 's of the form

$$c_2 f(\zeta) = h(\zeta) - P(\zeta, 0),$$
 (5.22)

for any  $C^2$  positively coercive function  $h : \mathbb{R} \to \mathbb{R}$ . (That h be  $C^2$ , as opposed to simply continuous, follows from the Lagrangian nature of the system (5.1).)

The structure (5.22) is not only necessary, but also sufficient for Q to be coercive. To see this, we need only to prove that the potential piece  $P + c_2 f$  of Q is positively coercive — the kinetic piece always being positive definite with minimum. With this goal in mind and as per definition 6, we must ensure that the multivariate limit to infinity of  $P + c_2 f$  diverges. Namely, we must find, for every k > 0, a radius R > 0such that  $P + c_2 f$  takes a value larger than k at any point with modulus larger than R.

Because h is positively coercive, for any k > 0 there exist a  $\zeta_0 \in \mathbb{R}$  such that, for all  $|\zeta| > \zeta_0$ ,

$$P(\zeta, 0) + c_2 f(\zeta) > k.$$
(5.23)

Since  $[-\zeta_0, \zeta_0]$  is compact with respect to  $\mathbb{R}$  and f is continuous,  $c_2 f$  reaches a global minimum when evaluated in this interval. Let m denote such minimum. Notice that, by analogy to the kinetic piece, P has positive definite Hessian everywhere and a minimum at  $\zeta = 0 = \eta$ . This implies that there exists R > 0such that, for all  $\zeta, \eta$  satisfying  $\|(\zeta, \eta)\| > R$ , it holds true that  $P(\zeta, \eta) > k - m$ . This completes the proof, owing to the fact that, for  $\|(\zeta, \eta)\| > R$ , either  $|\zeta| > \zeta_0$  and

$$P(\zeta, \eta) + c_2 f(\zeta) > P(\zeta, 0) + c_2 f(\zeta) > k,$$
(5.24)

or  $\zeta \in [-\zeta_0, \zeta_0]$ , in which case

$$P(\zeta, \eta) + c_2 f(\zeta) > k - m + m = k.$$
(5.25)

Summing up, for Q to be coercive, it is necessary and sufficient that  $|\alpha| \neq 1$ ,  $\omega^2 > 0$  and (5.22) is fulfilled. We draw attention to the fact that these requirements encompass the previously derived necessary and sufficient conditions (5.9) and (5.11) for Q to have a positive definite Hessian everywhere and a minimum. In other words, the systems of the form (5.1) that are G1 on the grounds of theorem 2 include and extend those that are G1 on the grounds of theorem 1.

**Cases in point.** Let  $a, b \in \mathbb{R}$ . Concrete cases from the family (5.1) with a conserved quantity Q of the form (5.3) that is coercive include: all previous examples with a Q that has positive definite Hessian everywhere and minimum (5.13) and instances beyond, such as

$$f(z) = a(z^2 - 1)(z^2 - 4)$$
 for  $a \ge 0$  and  $|\alpha| \le 1$ ,  $f(z) = b\sin(z)$ . (5.26)

These proposals' Qs only have a Hessian that is positive definite everywhere when

$$0 \leq a \leq \frac{\omega^2}{10(1-\alpha^2)} \text{ and } |\alpha| \leq 1, \qquad |b| < \frac{\omega^2}{|1-\alpha^2|},$$
 (5.27)

respectively.

Independent ghost systems formerly proven to be G1 on the (implicit) grounds of theorem 2 can be found in [59, 97] and in section 3.4 of [60].

### 5.3 An application of theorem 3

At last, we establish the subset of (5.1) that is G1 in the equivalent sense of definitions 1.a and 1.b as per theorem 3. Namely, we now require that the conserved quantity (5.3) is confining — recall our proposed definition 5.

The same reasoning as that between equations (5.15) and (5.17) earlier on applies in this case. Indeed, our discussion shows the existence of a line inside a level set of Q. In particular, this line is inside a path-connected component of a level set of Q. This contradicts definition 5. Thus, for Q to be confining, it must hold true that  $\alpha \neq \pm 1$  and  $\omega^2 > 0$ .

Having restricted the free parameter space in (5.1) so as to make it compatible with a confining Q in (5.3), we replicate the strategy in the previous section 5.2 and turn to restricting the free functional space. To this aim, we reconsider (5.22). Specifically, we ponder over relaxations of the (sufficiently smooth) function  $h(\zeta)$ . In seeking to exceed the positively coercive requirement, we envisage an oscillatory function whose maxima ultimately increase while minima remain below some threshold. For instance, see plot D in figure 2.

Rigorously, let  $f(\zeta)$  be of the form

$$c_2 f(\zeta) = h(\zeta) - P(\zeta, 0),$$
 (5.28)

where  $h : \mathbb{R} \to \mathbb{R}$  is any  $C^2$  function such that, for all  $k \in \mathbb{R}$  and for all  $\zeta \in \mathbb{R}$ , there exists  $\zeta_1, \zeta_2 \in \mathbb{R}$  with  $\zeta_1 < \zeta < \zeta_2$  such that  $f(\zeta_1) > k$  and  $f(\zeta_2) > k$ . Then, (5.3) can be conveniently rewritten as

$$Q = \frac{1}{2} \left( \dot{x}^2 + \dot{y}^2 + 2c_1 \dot{x} \dot{y} \right) + P(\zeta, \eta) + h(\zeta) - P(\zeta, 0) = Q_1 + h(\zeta) = Q_2 + h(\zeta) - P(\zeta, 0).$$
(5.29)

Subsequently, we prove that (5.29) is confining iff (5.28) is satisfied.

Consider the level set  $Q^{\neg}(c)$  for some arbitrary  $c \in \mathbb{R}$ . Let  $A \subset \mathbb{R}^4$  denote a path-connected component thereof. Further, let  $J \subset \mathbb{R}$  denote the projection of A to the  $\zeta$  variable. First, we establish that J is relatively compact with respect to  $\mathbb{R}$ . Afterwards, we demonstrate that A is relatively compact with respect to  $\mathbb{R}^4$ . This entails the sufficiency of (5.28) for rendering (5.29) confining.

By virtue of (5.28), for any point  $\zeta_0 \in J$  and choosing k = c, there exist  $\zeta_1 < \zeta_0 < \zeta_2$  such that  $h(\zeta_1) > c$ and  $h(\zeta_2) > c$ . Within  $Q^-(c)$ , the values of  $h(\zeta)$  are contained in  $(-\infty, c]$ , since zero is the absolute minimum value of  $Q_1$ . Consequently, such  $\zeta_1$  and  $\zeta_2$  do not belong in J. Furthermore, J is path-connected because it is the image of the path-connected set A by the projection, which is a continuous map. Altogether, J is a subset of  $[\zeta_1, \zeta_2]$ ; thus, J is relatively compact with respect to  $\mathbb{R}$ .

The continuity of  $h(\zeta) - P(\zeta, 0)$  warrants its attainment of a global minimum value m on the closure of J. Consistently, the maximum value of  $Q_2$  on A is c - m. In fact,  $Q_2$  is a positive elliptic paraboloid and, as such, positively coercive as a function of its variables — which becomes apparent in terms of  $(\dot{x}, \dot{y}, x, y)$ . Definition 4, in the restrictive sense wherein the absolute value is removed, assures us of the existence of a radius R such that  $Q_2(\dot{x}, \dot{y}, x, y) > c - m$  whenever  $\|(\dot{x}, \dot{y}, x, y)\| > R$ . It follows that  $A \subset B(0, R)$ ; thereby, A is relatively compact with respect to  $\mathbb{R}^4$ .

To wrap up the proof, we show by contradiction that (5.28) is not only sufficient but also necessary a condition for (5.29) to be confining. Assume  $h(\zeta)$  does not have the property (5.28). I.e. there exist  $k \in \mathbb{R}$  and  $\zeta_0 \in \mathbb{R}$  such that  $h(\zeta) \leq k$  for all  $\zeta_1 < \zeta_0$  (and similarly for  $\zeta_2 > \zeta_0$ ). Then, the level set  $Q^-(k)$  contains the subset

$$\{(\dot{x}, \dot{y}, \eta, \zeta) \in \mathbb{R}^4 \,|\, \dot{x} = \sqrt{2(k - h(\zeta))}, \, \dot{y} = 0, \, \eta = 0, \, \zeta \leqslant \zeta_0\},\tag{5.30}$$

which is path-connected yet not relatively compact; so Q is not confining.

Retrospectively, we can enhance our understanding of the successful ansatz (5.28). The free function f is decomposed into two pieces: the free function h possessing the crucial property that guarantees the system is G1, plus the fixed function  $P(\zeta, 0)$  that balances the relevant term of Q — recall (5.20). (This intuition also holds for the coercive instance (5.22), where  $P(\zeta, 0)$  surfaced naturally.) Moreover, the decomposition unlocks the exhaustiveness of the result. The term  $P(\zeta, 0)$  provides the exact cancellation of the quadratic terms  $\zeta^2$  inside the square root in (5.30), rendering it well-defined for all values of  $\zeta \leq \zeta_0$ .

In short, for Q to be confining, it is necessary and sufficient that  $|\alpha| \neq 1$ ,  $\omega^2 > 0$  and (5.28) holds. We highlight that (5.28) generalizes (5.22), so that the systems of the form (5.1) that are G1 on the grounds of theorem 3 broaden those that are G1 on the grounds of theorem 2.

**Cases in point.** Let  $a, b \in \mathbb{R}$ . Exemplars of (5.1) with a conserved quantity Q of the form (5.3) that is confining include: all previous examples with a Q that has minimum and positive definite Hessian everywhere (5.13), those associated with a coercive Q (5.26) and instances beyond, such as

$$f(z) = ae^{z}\cos(z) \text{ for } a \neq 0 \qquad f(z) = bz^{2}\sin(z) \text{ for } |b| \ge \frac{\omega^{2}}{2|1 - \alpha^{2}|} \equiv b_{c}.$$
 (5.31)

The left-most example is neither bounded from below nor from above, regardless of the sign of  $a^5$ . The right-most example neatly showcases the doublet of single-sided implications (4.7)-(4.8).

<sup>&</sup>lt;sup>5</sup>This example refutes footnote 5 in [59]: the corresponding conserved quantity Q is bounded neither from below nor from above, yet it is confining and thus guarantees that the system is G1 as per both definitions 1.a and 1.b.



Figure 2: Archetypal instances of the rich phenomenology of the right-most example in (5.31). The function (5.32) is shown, with its variable  $\zeta$  labelling the abscissa. The free parameters have been set to  $\omega^2 = 1$ ,  $\alpha = 0.3$ ,  $\nu = 7$  and b = 0 (plot A), b = 1/8 (plot B),  $b = b_c \approx 0.55$  (plot C), b = 2 (plot D). Thereby, the itemization below (5.31) is illustrated.

All these are exemplary potentials of conserved quantities that can be used to analytically prove G1 stability in the compact sense of definition 1.b. The lower half portrays radically new proposals, including totally unbounded scenarios.

- For b = 0, the system decouples. The conserved quantity Q has positive definite Hessian everywhere and minimum. Accordingly, the system can be proven to be G1 through either theorems 1, 2 or 3.
- For  $|b| \in (0, b_c)$ , the conserved quantity Q is coercive, but the Hessian is not positive definite at some points. Thence, the system can be proven to be G1 via either theorems 2 or 3, but not via theorem 1.
- For  $|b| \in [b_c, \infty)$ , the conserved quantity Q is confining, but it is neither coercive nor does it possess a positive definite Hessian everywhere. In this case, the system can only be proven to be G1 by means of theorem 3.
  - If  $|b| = b_c$ , Q has (infinitely many) global minima.
  - $\circ\,$  Else, Q is unbounded from above and below.

Figure 2 exhibits the aforementioned progression in plots A to D, respectively. In more detail, it represents the G1-triggering piece h of the conserved quantity Q, for the choice of f in the right-most example of (5.31) and in terms of the  $\zeta$  variable introduced in (5.19):

$$h(\zeta) = P(\zeta, 0) + c_2 f(\zeta) = \frac{\omega^2}{2(1+\alpha^2)} \zeta^2 + c_2 b \zeta^2 \sin(\nu \zeta).$$
(5.32)

(At this point, it is beneficial to bring the discussion below (5.30) to mind.) The parameter  $\nu \in \mathbb{R} - \{0\}$  has been introduced by hand, for an easier appreciation of the qualitative behavior of the plotted functions.

We reiterate that examples of (ghost) systems provable to be G1 exclusively on the basis of theorem 3 are novel in a deep sense. Indeed, they could not have been obtained without an increase in both the understanding and techniques employed before.

# 6 Summary and conclusions

On the basis of the profuse literature in stable ghost theories, we have put forward a cartography for classical notions of stability currently in force. Crucially, we have differentiated between global and local stability notions, depending on their application to *all* or *some* solutions of the system: see G1-G2 versus L1-L2. We stress that only Lyapunov-like proposals L2 enjoy a long and fruitful tradition in the mathematical literature of dynamical systems. From this perspective, alternative classical notions of stability enforced in ghost physics open the door to expansive studies in an altogether distinct discipline.

Subsequently, we have formalized the global notions G1 and G2: see 1.a-1.b and 2, respectively. It is worth emphasizing that our propounded definitions apply to generic dynamical systems (3.1), independent of their possible Lagrangian/Hamiltonian and/or ghostly nature. The fact that G1 admits two formalizations 1.a-1.b is consequential for dynamical systems whose solutions are defined in metric spaces without the Heine-Borel property, as shown in figure 1. We call attention to the relevance of these scenarios, which encompass infinite-dimensional Banach and Hilbert spaces, as well as geodesically incomplete Riemannian manifolds. In all such cases, there is no equivalence between 1.a and 1.b, and only 1.b implies 2. The dichotomy is to be resolved on physical grounds, according to the desired phenomenology of the systems under study. For ghost systems, we regard this as an important open question.

Taking definition 1.b as bedrock throughout, we have (re)explicated the mathematical foundations sustaining the analytic proofs of global stability G1 for ghost systems: see theorems 1 and 2. The crux of the proofs relies on two insights. First, the characterization of ghost systems as dynamical Lagrangian/Hamiltonian systems for which the energy does not have a global minimum. Second, the identification of a functionally independent conserved quantity Q possessing a key property as the means to ascertain the desired G1 feature in the system. The key property in (a befitting adaptation of) theorem 1 is two-fold: Q has a minimum and its Hessian is positive definite everywhere. A more sophisticated key property is employed in theorem 2: Q is coercive, in the sense of definition 4. Theorems 1 and 2 formalize and classify the examples in [56, 57, 60, 81–93] and [59, 60, 97], respectively.

We have extended the aforementioned known frameworks in an unprecedented manner: see our main result theorem 3. In more detail, within the same paradigm, we have generalized the key property to: Q is confining, in the sense of our proposed novel definition 5. The architecture conformed by theorem 3 and definition 5 applies to generic dynamical systems in Hausdorff spaces. The supervened augmentation in ghost systems provable to be G1 as a direct consequence of our work is both quantitative and qualitative. In the latter regard, we have explicitly proved that Q need *not* be bounded from below for the system to be G1, thereby rigorously contravening the status quo of ghost physics in a counter-intuitive way. For an instinctive impression, compare the upper and lower plots in figure 2, remarkably the totally unbounded scenario in plot D.

We have introduced confining functions tailored to the stability definition 1.b for a dynamical system. Confining functions englobe coercive functions in metric spaces with the Heine-Borel property, see proposition 1. Our definition 5 is intrinsic (i.e. coordinate-independent) and it is practicable. Indeed, its direct implementation has allowed us to put forward novel examples of G1 stable ghost systems that could not have been obtained with the artillery in the literature, see (5.31).

Moreover, we have explicitly related confining functions to the Liouville-Arnold theorem [157]. In lemma 3, we have proved that, given an integrable mechanical system, the presence of one confining conserved quantity guarantees the fulfillment of all hypotheses and ensures that the phase space is foliated exlusively by compact leaves. This result connects our work with open questions in the mathematical field of integrable systems [155]. Its exploitation conforms to yet another appealing direction for inter-disciplinary research.

**Outlook.** More ambitiously, one may envisage a change of paradigm in the analytic proof of global stability: without appeal to conserved quantities. This tantalizing possibility has been considered in [59]. Deffayet and coauthors set forth and illustrate a conjecture in such an unexplored arena. We deem the search for analytic proofs of global stability without calling for a conserved quantity pivotal to the development of realistic ghost models. This physically motivated search seems particularly enticing for the community studying non-integrable systems.

The application of our results to field theory and their quantization are obvious follow ups of noteworthy physical interest. In both contexts, the inequivalence between global stability definitions 1.a and 1.b is consequential. This disjunction and other functional analytical subtleties are essential aspects to bear in mind, even in non-ghostly basic settings such as a free massless scalar field [162, 163]. We defer the investigation of such compelling topics to future works.

Acknowledgements. We would like to show our high appreciation to Xavier Gràcia, for his suggestion to include the relation to proper functions, for perspicacious feedback and for pointing out [99,143,164]. We are indebted to Jose Beltrán Jiménez, Saskia Demulder, Manuel Lainz Valcázar and Alexander Vikman for insightful conversations at the onset of investigations. We have benefitted from Ivano Basile's constructive comments on an earlier version of this text. VED and JGR greatly thank Shinji Mukohyama for enlightening

discussions during a visit to the Yukawa Institute for Theoretical Physics (YITP). Additionally, VED and JGR are deeply grateful to all members of the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU) for their extended hospitality during the development of this work; in particular, host Elisa G. M. Ferreira and vibrant interlocutors Andrés Nahuel Briones, Thomas Edward Melia and Tsutomu Yanagida.

The work of VED, as well as an invited visit of JGR to the Excellence Cluster ORIGINS, are funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), under Germany's Excellence Strategy – EXC-2094 – 390783311. JGR acknowledges financial support from Ministerio de Ciencia, Innovación y Universidades in Spain, project D2021-125515NB-21 and from Agencia Estatal de Investigación, project RED2022-134301-T.

# A Supplementary proofs

This appendix proves theorems 1 and 2, as well as the one-sided implications (4.7) and (4.8). The reader may find useful the books in topology by Schechter [142] and especially by Lee [164].

With the aim of proving implication (4.7), we introduce the notion of a positively coercive function.

**Definition 6.** A function  $f : \mathbb{R}^n \to \mathbb{R}$  is **positively coercive** if the value of f(x) tends to plus infinity as the modulus of x diverges:

$$\lim_{\|x\|\to+\infty} f(x)\to +\infty. \tag{A.1}$$

Notice that a positively coercive function is an instance of a coercive function — recall definition 4. In particular, coercive functions with a global minimum are positively coercive. Implication (4.7) follows readily.

**Lemma 1.** [Entails implication (4.7).] If a twice-differentiable function  $f : \mathbb{R}^n \to \mathbb{R}$  has a minimum and a positive definite Hessian everywhere, then it is positively coercive.

*Proof.* Let  $x_0 \in \mathbb{R}^n$  be the point where f has a minimum. The Taylor expansion up to second order of f around  $x_0$  satisfies

$$f(x) = f(x_0) + \nabla f(x_0)(x - x_0) + (x - x_0)^T \operatorname{Hess} f(x_0 + a(x - x_0))(x - x_0),$$
(A.2)

for some  $a \in (0, 1)$ . Because f has a positive definite Hessian everywhere,  $f(x) \ge f(x_0)$ . The equality only holds for  $x = x_0$ . Namely,  $x_0$  is the global minimum.

Because  $\mathbb{R}^n$  has the Heine-Borel property, any ball therein is a compact set. The corresponding boundary is a closed set inside the ball and thus itself is a compact set. Any continuous function in a compact set achieves both an absolute maximum and an absolute minimum in the set. In particular, consider the boundary of the unit ball centered on  $x_0$ . Let  $f_*$  denote the absolute minimum of f in  $\partial B(x_0, 1)$ . Notice that  $f_* > f(x_0)$ .

By definition, f is positively coercive if, for every  $d \in \mathbb{R}^+$ , there exists a radius  $R_d$  such that, for any point  $x \in \mathbb{R}^n$  with bigger modulus  $||x|| > R_d$ , the inequality f(x) > d holds true. Without loss of generality, we take  $d > f(x_0)$ . In the following, we prove that the ansatz

$$R_d = \frac{d - f(x_0)}{f_* - f(x_0)} + \|x_0\| + 1$$
(A.3)

realizes the requirement. Observe that  $d > f(x_0)$  ensures  $R_d > 0$ .

Let  $r = ||x - x_0||$  and consider the auxiliary point

$$y = \frac{x - x_0}{r} + x_0 \in \partial B(x_0, 1).$$
 (A.4)

By virtue of the mean value theorem,

$$f(y) - f(x_0) = \nabla f(z)(y - x_0), \qquad z \equiv y + b(x_0 - y), \tag{A.5}$$

for some  $b \in (0, 1)$ . The Taylor expansion up to second order of f around z fulfills

$$f(x) = f(z) + \nabla f(z)(x-z) + (x-z)^T \operatorname{Hess} f(z+c(x-z))(x-z),$$
(A.6)

for some  $c \in (0,1)$ . Because f has a positive definite Hessian everywhere,  $f(x) \ge f(z) + \nabla f(z)(x-z)$ . Since  $x - z = (y - x_0)(r - 1 + b)$  and making use of (A.5), it follows that

$$f(x) \ge f(z) + \nabla f(z)(y - x_0)(r - 1 + b) = f(z) + [f(y) - f(x_0)](r - 1 + b).$$
(A.7)

Provided that  $||x|| > R_d$  in (A.3) and employing the triangle inequality,

$$r - 1 + b = \|x - x_0\| - 1 + b > \|x\| - \|x_0\| - 1 + b > \frac{d - f(x_0)}{f_* - f(x_0)}.$$
(A.8)

holds true. Substitution of (A.8) in (A.7), in light of  $x_0$  being a global minimum and the very definitions of y and  $f_*$ , finally yields

$$f(x) > f(z) + \left[f(y) - f(x_0)\right] \frac{d - f(x_0)}{f_* - f(x_0)} > f(x_0) + \left[f_* - f(x_0)\right] \frac{d - f(x_0)}{f_* - f(x_0)} = d.$$
(A.9)

Definition 4 of a coercive function can be straightforwardly extended to generic metric spaces.

**Lemma 2.** Let M and N be metric spaces. A coercive function  $f : M \to N$  has bounded level sets. Moreover, if M has the Heine-Borel property, the level sets are relatively compact with respect to M.

*Proof.* Consider an arbitrary but fixed  $c \in N$ . Since f is coercive, for  $||c|| \in \mathbb{R}$  there exists a radius  $R_c \in \mathbb{R}^+$  such that, for any point  $x \in M$  with bigger modulus  $||x|| > R_c$ , the inequality ||f(x)|| > ||c|| holds true. In particular, at any point outside the ball  $B(0, R_c) \subset M$ , f takes a value different from c. Consequently, the level set  $f^{\neg}(c) \subset B(0, R_c)$  and, therefore, it is bounded.

If M has the Heine-Borel property, then balls are compact sets. Since balls are closed in metric spaces, then the closure of  $f^{\neg}(c)$  is also inside  $B(0, R_c)$ . The closure of  $f^{\neg}(c)$  is compact as a result of it being a closed set inside a compact.

The above generalization allows us to succinctly prove theorems 1 and 2, which are hereafter restated for the convenience of the reader.

**Theorem 1.** Given a system Sol with  $\mathcal{P} = \mathbb{R}^n$ , if the energy  $E : \mathbb{R}^n \to \mathbb{R}$  is a twice differentiable conserved quantity that has a minimum and whose Hessian is positive definite everywhere, then the system is G1.

*Proof.* Let e denote the constant value of E along a fixed solution  $\gamma$ . It follows that  $\gamma(I_{\gamma}) \subset E^{\neg}(e)$ .

Because E is twice differentiable, has a minimum and has a positive definite Hessian everywhere, by virtue of lemma 1, it is coercive. By lemma 2, its level sets are both bounded and relatively compact with respect to  $\mathbb{R}^n$ , which has the (crucial) Heine-Borel property. Ergo the system is G1 in the equivalent sense of definitions 1.a and 1.b.

**Theorem 2.** Given a system Sol with  $\mathcal{P} = \mathbb{R}^n$ , if there exists a conserved quantity  $Q : \mathbb{R}^n \to \mathbb{R}$  that is coercive, then the system is G1.

*Proof.* Let q denote the constant value of Q along a fixed solution  $\gamma$ . It follows that  $\gamma(I_{\gamma}) \subset Q^{\neg}(q)$ .

Because Q is coercive, by virtue of lemma 2, its level sets are both bounded and relatively compact with respect to  $\mathbb{R}^n$ , which has the (crucial) Heine-Borel property. Ergo the system is G1 in the equivalent sense of definitions 1.a and 1.b.

Lastly, we prove implication (4.8), which holds true for metric spaces, as long as the source space has the Heine-Borel property.

**Proposition 1.** [Entails implication (4.8).] Let M be a metric space with the Heine-Borel property and let N be a metric space. A coercive function  $f: M \to N$  is confining.

*Proof.* Consider an arbitrary but fixed  $c \in N$ . Let A denote a path-connected component of the level set  $f^{\neg}(c)$ . The latter satisfies  $A \subset f^{\neg}(c)$ , which is bounded as per lemma 2. As such,  $f^{\neg}(c) \subset M$  is contained in a certain ball. Since M has the Heine-Borel property, balls are compact. Moreover, as a metric space, M is Hausdorff. Thus, A is relatively compact with respect to M because it is inside a compact.  $\Box$ 

# **B** Relation to proper functions

In topology and in differential geometry, there exists the appurtenant concept of a *proper function*. This is closely related to the notion of coercivity, see proposition 2. Crucially for us, proper functions have compact level sets. What is more, there are useful results in the literature providing sufficient conditions for a function to be proper, see e.g. proposition 4.93 in [164]. Altogether, we find proper functions to be a promising addition to the study of G1 global stability, in the sense of definition 1.b. In the following, we relate our work with proper functions.

Proper functions suffer from an occasional polysemy in the literature. We employ the meaning in [164]:

**Definition 7.** Let M and N be topological spaces. A function  $f: M \to N$  is **proper** if the anti-image of each compact set of N is compact.

Understood in this precise sense and as announced, proper functions are closely related to coercive functions.

**Proposition 2.** Let M and N be metric spaces with the Heine-Borel property. A continuous function  $f: M \to N$  is coercive iff it is proper.

*Proof.* Coercive  $\Rightarrow$  proper. Consider an arbitrary compact set  $K \subset N$ . As such, it is both closed and bounded. Because f is continuous, the first property implies that  $f^{\neg}(K)$  is also closed. The second property entails the existence of a distance  $d_K \in \mathbb{R}^+$  such that  $K \subset B(0, d_K)$ . Since f is coercive, there exists a

corresponding radius  $R_{d_K}$  such that, for all  $x \in M$  with  $||x|| > R_{d_K}$ , the inequality  $||f(x)|| > d_K$  holds true. In particular,

$$f^{\neg}(K) \subset f^{\neg}(B(0, d_K)) \subset B(0, R_{d_K}).$$
(B.1)

The ball  $B(0, R_{d_K})$  is compact because M has the Heine-Borel property. Consequently,  $f^{-}(K)$  is a compact set: it is a closed set inside a compact.

**Proper**  $\Rightarrow$  coercive. Given an arbitrary distance  $d \in \mathbb{R}^+$ , the ball  $B(0,d) \subset N$  is a compact set because N has the Heine-Borel property. Since f is proper,  $f^{\neg}(B(0,d))$  is a compact set in M. As such, it is bounded. Namely, there exists a radius  $R_d$  such that  $f^{\neg}(B(0,d)) \subset B(0,R_d)$ . Equivalently, for all  $x \in M$ with  $||x|| > R_d$ , the inequality ||f(x)|| > d holds true.

Proper functions are relevant to our work owing to the following two results. First, following straightforwardly from definition 7: the level sets of a proper function are compact. Second, holding true provided that the initial space is Hausdorff:

**Proposition 3.** Let  $\mathcal{P}$  be a Hausdorff space and let N be a topological space. A proper function  $f : \mathcal{P} \to N$  is confining.

*Proof.* Consider an arbitrary but fixed point  $c \in N$ . Let A denote a path-connected component of the level set  $f^{\neg}(c)$ . The latter satisfies  $A \subset f^{\neg}(c)$ , which is compact by definition of a proper function. Since M is Hausdorff and A is inside a compact, A is relatively compact with respect to M.

Counterexamples of the converse are (5.31).

## C Integrability in light of a confining conserved quantity

For completeness, we begin by introducing the relevant geometric structures of a mechanical system. A more general and detailed presentation can be found in chapter 2 of [152]. Alternatively, see the last chapter in [146].

Let  $\mathcal{P}$  denote a smooth manifold, which is Hausdorff by definition [146]. Further, let  $\omega$  be a symplectic form on  $\mathcal{P}$ . Any (at least  $C^2$ ) function  $F : \mathcal{P} \to \mathbb{R}$  has an associated vector field  $X_F$ , called the Hamiltonian vector field of F, defined as

$$\iota_{X_F}\omega = \mathrm{d}F.\tag{C.1}$$

When F is a conserved quantity F = Q,  $X_Q$  is called an infinitesimal symmetry [152]. When F is the Hamiltonian (or Lagrangian energy) of the dynamical system F = H, the integral curves of  $X_H$  are the solutions of the system.

The assignation of  $X_F$  to F induces a canonical Poisson bracket. For two functions  $F, G : \mathcal{P} \to \mathbb{R}$ ,

$$\{F,G\} := \iota_{X_G} \iota_{X_F} \omega = \iota_{X_G} dF = X_G(F).$$
(C.2)

When (C.2) vanishes, F and G are said to be in involution.

Recount definition 5 of a confining function and the definition of a complete vector field — restated below definition 2. The following lemma shows that a confining function can only be in involution with functions whose Hamiltonian vector field generates G1 dynamics, in the compact sense of definition 1.b. This entails the completeness of such Hamiltonian vector fields. **Lemma 3.** Let  $\omega$  be a symplectic form on a smooth manifold  $\mathcal{P}$ . Consider two  $C^2$  functions  $F, G : \mathcal{P} \to \mathbb{R}$  such that F is confining. If F and G are in involution, then the integral curves of  $X_G$  are relatively compact with respect to  $\mathcal{P}$  and, hence,  $X_G$  is complete.

*Proof.* If F and G are in involution

$$0 = \{F, G\} = X_G(F), \tag{C.3}$$

then F is constant along the integral curves  $\{\gamma\}$  of  $X_G$ . In particular, each  $\gamma$  is inside a path-connected component of a level set of F, which is relatively compact with respect to  $\mathcal{P}$  because F is confining. Since  $\mathcal{P}$  is Hausdorff, each  $\gamma$  is inside a compact set and thus its closure  $\overline{\gamma}$  is compact. By negation of the Escape Lemma (see, e.g. [146]),  $X_G$  is complete.  $\Box$ 

This result is relevant for integrable systems, in the sense of section 4.4.

The celebrated Liouville-Arnold theorem ensures the existence of action-angle coordinates for integrable systems that trivialize the dynamics, provided that the infinitesimal symmetries are complete vector fields. In more detail, the theorem ascertains that the phase space of the system is foliated by (the path-connected components of) the intersection of the level sets of the conserved quantities. Furthermore, these leaves are diffeomorphic to  $\mathbb{T}^k \times \mathbb{R}^{n-k}$ , and there exist action-angle coordinates in terms of which the dynamics correspond to that of *n* free particles. (For a rigorous formulation, see [157].)

Assume an integrable system defined on  $\mathcal{P}$  where one of the conserved quantities Q is confining. By virtue of lemma 3, the infinitesimal symmetries of the conserved quantities in involution are guaranteed to be complete. Thus, the hypotheses of the Liouville-Arnold theorem are satisfied. Moreover, since the path-connected components of the level sets of Q are relatively compact with respect to  $\mathcal{P}$ , the leaves provided by the Liouville-Arnold theorem are *compact* (i.e. k = n). It follows that the dynamics, described in terms of action-angle variables, is that of free particles moving *exclusively* around torii.

In brief, theorem 3 — and its corollaries theorems 1 and 2 — accept an interpretation as methods to enforce the completeness of vector fields associated with symmetries and the compactness of the leaves in integrable systems.

### D Supplementary properties of the system (5.1)

In this appendix, we discuss properties of the system (5.1) against the backdrop of section 4.4.

Unnatural yet Lagrangian-preserving symmetry of Q. The symplectic form induced by the Lagrangian 5.1 is

$$\omega_L = \mathrm{d}x \wedge \mathrm{d}\dot{x} - \mathrm{d}y \wedge \mathrm{d}\dot{y},\tag{D.1}$$

and the Hamiltonian function of the system is the Lagrangian energy (5.2).

The infinitesimal symmetry (C.1) associated to the conserved quantity (5.3) via Noether's theorem is the vector field

$$X_Q = (\dot{x} + c_1 \dot{y}) \frac{\partial}{\partial x} - (\dot{y} + c_1 \dot{x}) \frac{\partial}{\partial y} - \left[ \omega^2 (x + c_1 y) + c_2 f' \right] \frac{\partial}{\partial \dot{x}} + \left[ \omega^2 (y + c_1 x) + \alpha c_2 f' \right] \frac{\partial}{\partial \dot{y}}.$$
 (D.2)

The associated infinitesimal changes for positions and velocities are

$$\begin{cases} x \to x' = x + (\dot{x} + c_1 \dot{y})\varepsilon, \\ y \to y' = y - (\dot{y} + c_1 \dot{x})\varepsilon, \end{cases} \qquad \begin{cases} \dot{x} \to \dot{x}' = \dot{x} - \left[\omega^2 (x + c_1 y) + c_2 f'\right]\varepsilon, \\ \dot{y} \to \dot{y}' = \dot{y} + \left[\omega^2 (y + c_1 x) + \alpha c_2 f'\right]\varepsilon, \end{cases}$$
(D.3)

respectively. Here,  $\varepsilon$  is an infinitesimal parameter. Notice that the changes in positions depend on velocities: the signature of an unnatural symmetry.

Being a Noether symmetry, (D.2) leaves invariant the Lagrangian energy and the symplectic form. However, it need not be a symmetry of the Lagrangian. In this case, it does turn out to be one:

$$\mathcal{L}_{X_Q}L = -2\frac{\mathrm{d}}{\mathrm{d}t} \left[ \frac{\omega^2}{2} \left( x^2 + y^2 + 2c_1 xy \right) + c_2 f(x + \alpha y) \right].$$
(D.4)

**Bi-Hamiltonian nature of the system.** A mechanical system is (quasi-)bi-Hamiltonian (in the sense of [165]) if there exists a compatible symplectic form  $\hat{\omega}$  and a function  $\hat{H}$  with the same dynamics as the original pair ( $\omega_L, E_L$ ).

That is the case for the system (5.1), whenever  $\alpha \neq \pm 1^6$ . Then, the compatible symplectic form is

$$\hat{\omega} = \mathrm{d}x \wedge \mathrm{d}\dot{x} + c_1 \mathrm{d}x \wedge \mathrm{d}\dot{y} + \mathrm{d}y \wedge \mathrm{d}\dot{y} + c_1 \mathrm{d}y \wedge \mathrm{d}\dot{x} \tag{D.5}$$

and  $\hat{H} = Q$  given by (5.3). Explicitly, the vector fields that encode the dynamics in both settings,

$$\iota_{X_{E_L}}\omega_L = \mathrm{d}E_L \quad \text{and} \quad \iota_{\hat{X}_Q}\hat{\omega} = \mathrm{d}Q,$$
(D.6)

are equal:  $X_{E_L} = \hat{X}_Q$ . Besides, both structures are compatible because the technical requirement that the Nijenhuis torsion vanishes is met.

### References

- A. Pais and G. E. Uhlenbeck, "On Field Theories with Non-Localized Action," Phys. Rev. 79, 145 (1950) doi:10.1103/PhysRev.79.145
- [2] B. Podolsky, "A Generalized Electrodynamics Part I-Non-Quantum," Phys. Rev. 62, 68 (1942) doi:10.1103/PhysRev.62.68
- B. Podolsky and C. Kikuchi, "A Generalized Electrodynamics Part II-Quantum," Phys. Rev. 65, 228 (1944) doi:10.1103/PhysRev.65.228
- [4] B. Podolsky and C. Kikuchi, "Auxiliary Conditions and Electrostatic Interaction in Generalized Quantum Electrodynamics," Phys. Rev. 67, 184 (1945) doi:10.1103/PhysRev.67.184
- [5] H. Weyl, "A New Extension of Relativity Theory," Annalen Phys. 59, 101 (1919) doi:10.1002/andp.19193641002
- [6] H. Weyl, "Electricity and Gravitation," Nature **106**, 800 (1921) doi:10.1038/106800a0

<sup>&</sup>lt;sup>6</sup>For  $\alpha = \pm 1$ , the two-form  $\hat{\omega}$  is not regular, i.e. it is presymplectic. This case is not further developed because the G1 stable examples in sections 5.1-5.3 require  $\alpha \neq \pm 1$ .

- [7] D. Lovelock, "The Einstein tensor and its generalizations," J. Math. Phys. 12, 498 (1971) doi:10.1063/1.1665613
- [8] E. Cartan, "Sur les variétés de courbure constante d'un espace euclidien ou non-euclidien," Bull. Soc. Math. Fr. 47, 125 (1919); 48, 132 (1920) doi:10.24033/bsmf.1007
- X. Gràcia, J. M. Pons and N. Román-Roy, "Higher-order Lagrangian systems: Geometric structures, dynamics, and constraints," J. Math. Phys. 32, 2744 (1991) doi:10.1063/1.529066
- [10] F. Etayo Gordejuela, P. L. García Pérez, J. Muñoz Masqué and J. Pérez Álvarez, "Higher-order Utiyama-Yang-Mills Lagrangians," J. Geom. Phys. 57, 1089 (2007) doi:10.1016/j.geomphys.2006.09.004
- [11] A. J. Bruce, K. Grabowska and J. Grabowski, "Higher order mechanics on graded bundles," J. Phys. A 48 205203 (2015) doi:10.1088/1751-8113/48/20/205203 [arXiv:1412.2719 [math-ph]]
- [12] M. Castrillón and J. Gaset Rifà, "Higher-order Utiyama theorem," Manuscript under preparation
- K. S. Stelle, "Renormalization of Higher Derivative Quantum Gravity," Phys. Rev. D 16, 953 (1977) doi:10.1103/PhysRevD.16.953
- [14] B. Grinstein, D. O'Connell and M. B. Wise, "The Lee-Wick standard model," Phys. Rev. D 77, 025012 (2008) doi:10.1103/PhysRevD.77.025012 [arXiv:0704.1845 [hep-ph]]
- [15] E. T. Tomboulis, "Renormalization and unitarity in higher derivative and nonlocal gravity theories," Mod. Phys. Lett. A 30, 1540005 (2015) doi:10.1142/S0217732315400052
- [16] P. D. Mannheim, "Normalization of the vacuum and the ultraviolet completion of Einstein gravity," Int. J. Mod. Phys. D 32, 2350096 (2023) doi:10.1142/S0218271823500967 [arXiv:2303.10827 [hep-th]]
- [17] P. D. Mannheim, "Pauli-Villars and the ultraviolet completion of Einstein gravity," [arXiv:2404.16148 [hep-th]]
- [18] R. R. Caldwell, "A Phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state," Phys. Lett. B 545, 23 (2002) doi:10.1016/S0370-2693(02)02589-3 [arXiv:astro-ph/9908168 [astro-ph]]
- [19] S. Nojiri and S. D. Odintsov, "Modified gravity with negative and positive powers of the curvature: Unification of the inflation and of the cosmic acceleration," Phys. Rev. D 68, 123512 (2003) doi:10.1103/PhysRevD.68.123512 [arXiv:hep-th/0307288 [hep-th]]
- [20] C. Charmousis, "Higher order gravity theories and their black hole solutions," Lect. Notes Phys. 769, 299 (2009) doi:10.1007/978-3-540-88460-6\_8 [arXiv:0805.0568 [gr-qc]]
- [21] R. Brandenberger and P. Peter, "Bouncing Cosmologies: Progress and Problems," Found. Phys. 47, 797 (2017) doi:10.1007/s10701-016-0057-0 [arXiv:1603.05834 [hep-th]]
- [22] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess and J. Silk, "In the realm of the Hubble tension—a review of solutions," Class. Quant. Grav. 38, 153001 (2021) doi:10.1088/1361-6382/ac086d [arXiv:2103.01183 [astro-ph.CO]]

- [23] M. D. Freeman, C. N. Pope, M. F. Sohnius and K. S. Stelle, "Higher Order σ Model Counterterms and the Effective Action for Superstrings," Phys. Lett. B 178, 199 (1986) doi:10.1016/0370-2693(86)91495-4
- [24] A. M. Polyakov, "Fine Structure of Strings," Nucl. Phys. B 268, 406 (1986) doi:10.1016/0550-3213(86)90162-8
- [25] J. McOrist and S. Sethi, "M-theory and Type IIA Flux Compactifications," JHEP 12, 122 (2012) doi:10.1007/JHEP12(2012)122 [arXiv:1208.0261 [hep-th]]
- [26] J. T. Liu and R. Minasian, "Higher-derivative couplings in string theory: dualities and the B-field," Nucl. Phys. B 874, 413 (2013) doi:10.1016/j.nuclphysb.2013.06.002 [arXiv:1304.3137 [hep-th]]
- [27] J. Calderón-Infante, M. Delgado and A. M. Uranga, "Emergence of species scale black hole horizons," JHEP 01, 003 (2024) doi:10.1007/JHEP01(2024)003 [arXiv:2310.04488 [hep-th]]
- [28] J. Kubo and T. Kugo, "Anti-Instability of Complex Ghost," PTEP 2024, 053B01 (2024) doi:10.1093/ptep/ptae053 [arXiv:2402.15956 [hep-th]]
- [29] A. Fring, T. Taira and B. Turner, "Higher time-derivative theories from space-time interchanged integrable field theories," Universe 10, 198 (2024) doi:10.3390/universe10050198 [arXiv:2403.11949 [hep-th]]
- [30] S. M. Kuzenko and E. S. N. Raptakis, "Higher-derivative deformations of the ModMax theory," JHEP 06, 162 (2024) doi:10.1007/JHEP06(2024)162 [arXiv:2404.09108 [hep-th]]
- [31] A. Colléaux, D. Langlois and K. Noui, "Degenerate Higher-Order Maxwell Theories in Flat Space-Time," [arXiv:2404.18715 [gr-qc]]
- [32] S. Mironov and M. Valencia-Villegas, "Healthy Horndeski gravities with torsion," JCAP 07, 030 (2024) doi:10.1088/1475-7516/2024/07/030 [arXiv:2405.08673 [gr-qc]]
- [33] S. Mandal and S. Shankaranarayanan, "Origin of effective graviton mass from spontaneous symmetry breaking of scalar field," [arXiv:2407.06572 [hep-th]]
- [34] S. Mironov, A. Shtennikova and M. Valencia-Villegas, "Higher derivative SVT theories from Kaluza-Klein reductions of Horndeski theory," [arXiv:2408.04626 [hep-th]]
- [35] M. Ostrogradsky, "Mémoires sur les équations différentielles, relatives au problème des isopérimètres," Mem. Acad. St. Petersbourg 6, 385 (1850) https://hdl.handle.net/2027/mdp.39015038710128
- [36] D. M. Gitman, S. L. Lyakhovich and I. V. Tyutin, "Hamilton formulation of a theory with high derivatives," Sov. Phys. J. 26, 730 (1983) doi:10.1007/BF00898884
- [37] J. L. Lagrange, "Mécanique Analytique," Veuve Desaint, Paris (1788); Cambridge University Press (2009) doi:10.1017/CBO9780511701788
- [38] G. L. Dirichlet, "Über die Stabilität des Gleichgewichts," J. Reine Angew. Math. 32, 85 (1846) doi:10.1515/crll.1846.32.85
- [39] A. M. Lyapunov, "Problème général de la stabilité du mouvement," Ann. Fac. Sci. Toulouse Math. 9, 203 (1947); Princeton University Press (1948) doi:10.1515/9781400882311

- [40] T. S. Biró, S. G. Matinyan and B. Müller, "Chaos and gauge field theory," World Sci. Lect. Notes Phys. 56, 1 (1994) doi:10.1142/2584
- [41] T. J. Chen, M. Fasiello, E. A. Lim and A. J. Tolley, "Higher derivative theories with constraints: Exorcising Ostrogradski's Ghost," JCAP 02, 042 (2013) doi:10.1088/1475-7516/2013/02/042 [arXiv:1209.0583 [hep-th]]
- [42] B. Díaz, D. Higuita and M. Montesinos, "Lagrangian approach to the physical degree of freedom count," J. Math. Phys. 55, 122901 (2014) doi:10.1063/1.4903183 [arXiv:1406.1156 [hep-th]]
- [43] R. Klein and D. Roest, "Exorcising the Ostrogradsky ghost in coupled systems," JHEP 07, 130 (2016) doi:10.1007/JHEP07(2016)130 [arXiv:1604.01719 [hep-th]]
- [44] M. Crisostomi, R. Klein and D. Roest, "Higher Derivative Field Theories: Degeneracy Conditions and Classes," JHEP 06, 124 (2017) doi:10.1007/JHEP06(2017)124 [arXiv:1703.01623 [hep-th]]
- [45] B. Díaz and M. Montesinos, "Geometric Lagrangian approach to the physical degree of freedom count in field theory," J. Math. Phys. 59, 052901 (2018) doi:10.1063/1.5008740 [arXiv:1710.01371 [gr-qc]]
- [46] A. Ganz and K. Noui, "Reconsidering the Ostrogradsky theorem: Higher-derivatives Lagrangians, Ghosts and Degeneracy," Class. Quant. Grav. 38, 075005 (2021) doi:10.1088/1361-6382/abe31d [arXiv:2007.01063 [hep-th]]
- [47] V. Errasti Díez, M. Maier and J. A. Méndez-Zavaleta, "Constraint characterization and degree of freedom counting in Lagrangian field theory," Phys. Rev. D 109, 025010 (2024) doi:10.1103/PhysRevD.109.025010 [arXiv:2310.12218 [hep-th]]
- [48] T. P. Sotiriou and V. Faraoni, "f(R) Theories Of Gravity," Rev. Mod. Phys. 82, 451 (2010) doi:10.1103/RevModPhys.82.451 [arXiv:0805.1726 [gr-qc]]
- [49] A. De Felice and S. Tsujikawa, "f(R) theories," Living Rev. Rel. 13, 3 (2010) doi:10.12942/lrr-2010-3 [arXiv:1002.4928 [gr-qc]]
- [50] C. de Rham, G. Gabadadze and A. J. Tolley, "Resummation of Massive Gravity," Phys. Rev. Lett. 106, 231101 (2011) doi:10.1103/PhysRevLett.106.231101 [arXiv:1011.1232 [hep-th]]
- [51] D. Gorbunov, K. Koyama and S. Sibiryakov, "More on ghosts in DGP model," Phys. Rev. D 73, 044016 (2006) doi:10.1103/PhysRevD.73.044016 [arXiv:hep-th/0512097 [hep-th]]
- [52] J. Beltrán Jiménez and A. Delhom, "Ghosts in metric-affine higher order curvature gravity," Eur. Phys. J. C 79, 656 (2019) doi:10.1140/epjc/s10052-019-7149-x [arXiv:1901.08988 [gr-qc]]
- [53] A. Delhom, A. Jiménez-Cano and F. J. Maldonado Torralba, "Instabilities in Field Theory: A Primer with Applications in Modified Gravity," Springer (2023) doi:10.1007/978-3-031-40433-7 [arXiv:2207.13431 [gr-qc]]
- [54] E. Pagani, G. Tecchiolli and S. Zerbini, "On the Problem of Stability for Higher Order Derivatives: Lagrangian Systems," Lett. Math. Phys. 14, 311 (1987) doi:10.1007/BF00402140

- [55] A. V. Smilga, "Supersymmetric field theory with benign ghosts," J. Phys. A 47, 052001 (2014) doi:10.1088/1751-8113/47/5/052001 [arXiv:1306.6066 [hep-th]]
- [56] D. S. Kaparulin, S. L. Lyakhovich and A. A. Sharapov, "Stable interactions via proper deformations," J. Phys. A 49, 155204 (2016) doi:10.1088/1751-8113/49/15/155204 [arXiv:1510.08365 [hep-th]]
- [57] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Stable interactions in higher derivative field theories of derived type," Phys. Rev. D 99, 045020 (2019) doi:10.1103/PhysRevD.99.045020 [arXiv:1811.10019 [hep-th]]
- [58] T. Damour and A. Smilga, "Dynamical systems with benign ghosts," Phys. Rev. D 105, 045018 (2022) doi:10.1103/PhysRevD.105.045018 [arXiv:2110.11175 [hep-th]]
- [59] C. Deffayet, A. Held, S. Mukohyama and A. Vikman, "Global and local stability for ghosts coupled to positive energy degrees of freedom," JCAP 11, 031 (2023) doi:10.1088/1475-7516/2023/11/031 [arXiv:2305.09631 [gr-qc]]
- [60] C. Heredia and J. Llosa, "Are nonlocal Lagrangian systems fatally unstable?," [arXiv:2403.19777 [hep-th]]
- [61] A. Smilga, "Classical and quantum dynamics of higher-derivative systems," Int. J. Mod. Phys. A 32, 1730025 (2017) doi:10.1142/S0217751X17300253 [arXiv:1710.11538 [hep-th]]
- [62] M. Pavšič, "On negative energies, strings, branes, and braneworlds: A review of novel approaches," Int. J. Mod. Phys. A 35, 2030020 (2020) doi:10.1142/S0217751X20300203 [arXiv:2012.04976 [hep-th]]
- [63] N. Barnaby and N. Kamran, "Dynamics with infinitely many derivatives: The Initial value problem," JHEP 02, 008 (2008) doi:10.1088/1126-6708/2008/02/008 [arXiv:0709.3968 [hep-th]]
- [64] C. Heredia and J. Llosa, "Non-local Lagrangian mechanics: Noether's theorem and Hamiltonian formalism," J. Phys. A 54, 425202 (2021) doi:10.1088/1751-8121/ac265c [arXiv:2105.10442 [hep-th]]
- [65] C. Heredia and J. Llosa, "Nonlocal Lagrangian fields: Noether's theorem and Hamiltonian formalism," Phys. Rev. D 105, 126002 (2022) doi:10.1103/PhysRevD.105.126002 [arXiv:2203.02206 [hep-th]]
- [66] S. Deser and R. P. Woodard, "Nonlocal Cosmology," Phys. Rev. Lett. 99, 111301 (2007) doi:10.1103/PhysRevLett.99.111301 [arXiv:0706.2151 [astro-ph]]
- [67] T. Biswas, T. Koivisto and A. Mazumdar, "Towards a resolution of the cosmological singularity in nonlocal higher derivative theories of gravity," JCAP 11, 008 (2010) doi:10.1088/1475-7516/2010/11/008 [arXiv:1005.0590 [hep-th]]
- [68] G. Calcagni and L. Modesto, "Nonlocality in string theory," J. Phys. A 47, 355402 (2014) doi:10.1088/1751-8113/47/35/355402 [arXiv:1310.4957 [hep-th]]
- [69] G. Calcagni, B. L. Giacchini, L. Modesto, T. de Paula Netto and L. Rachwal, "Renormalizability of nonlocal quantum gravity coupled to matter," JHEP 09, 034 (2023) doi:10.1007/JHEP09(2023)034 [arXiv:2306.09416 [hep-th]]

- [70] C. Deffayet and R. P. Woodard, "The Price of Abandoning Dark Matter Is Nonlocality," JCAP 05, 042 (2024) doi:10.1088/1475-7516/2024/05/042 [arXiv:2402.11716 [gr-qc]]
- [71] J. Wess and B. Zumino, "Consequences of anomalous Ward identities," Phys. Lett. B 37, 95 (1971) doi:10.1016/0370-2693(71)90582-X
- [72] S. P. Novikov, "The Hamiltonian formalism and a many valued analog of Morse theory," Usp. Mat. Nauk 37N5, 3 (1982) doi:10.1070/RM1982v037n05ABEH004020
- [73] E. Witten, "Nonabelian Bosonization in Two-Dimensions," Commun. Math. Phys. 92, 455 (1984) doi:10.1007/BF01215276
- [74] J. Gomis, K. Kamimura and T. Mateos, "Gauge and BRST generators for space-time noncommutative U(1) theory," JHEP 03, 010 (2001) doi:10.1088/1126-6708/2001/03/010 [arXiv:hep-th/0009158 [hep-th]]
- [75] C. Heredia and J. Llosa, "Infinite Derivatives vs Integral Operators. The Moeller-Zwiebach paradox," J. Phys. A 57, 235202 (2024) doi:10.1088/1751-8121/ad4aa5 [arXiv:2302.00684 [hep-th]]
- [76] J. LaSalle and S. Lefschetz, "Stability by Liapunov's Direct Method with Applications" Academic Press (1961) https://archive.org/details/stabilitybyliapu0004jose
- [77] D. Robert and A. V. Smilga, "Supersymmetry vs ghosts," J. Math. Phys. 49, 042104 (2008) doi:10.1063/1.2904474 [arXiv:math-ph/0611023 [math-ph]]
- [78] A. V. Smilga, "Benign versus malicious ghosts in higher-derivative theories," Nucl. Phys. B 706, 598 (2005) doi:10.1016/j.nuclphysb.2004.10.037 [arXiv:hep-th/0407231 [hep-th]]
- [79] A. V. Smilga, "Ghost-free higher-derivative theory," Phys. Lett. B 632, 433 (2006) doi:10.1016/j.physletb.2005.10.014 [arXiv:hep-th/0503213 [hep-th]]
- [80] E. P. Gyftopoulos, "Lagrange Stability by Liapunov's Direct Method," Proc. Sym. Rea. Kin. Con. TID 7662, 227 (1963) https://elias-gyftopoulos-memorial-collection.unibs.it/otherpublications.htm
- [81] D. S. Kaparulin, S. L. Lyakhovich and A. A. Sharapov, "Classical and quantum stability of higher-derivative dynamics," Eur. Phys. J. C 74, 3072 (2014) doi:10.1140/epjc/s10052-014-3072-3 [arXiv:1407.8481 [hep-th]]
- [82] D. S. Kaparulin and S. L. Lyakhovich, "On the Stability of a Nonlinear Oscillator with Higher Derivatives," Russ. Phys. J. 57, 1561 (2015) doi:10.1007/s11182-015-0419-7
- [83] D. S. Kaparulin and S. L. Lyakhovich, "Energy and Stability of the Pais-Uhlenbeck Oscillator," Geom. Meth. Phys. Birkhäuser, Cham (2015) doi:10.1007/978-3-319-18212-4\_8 [arXiv:1506.07422 [hep-th]]
- [84] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Bounded Hamiltonian in the Fourth-Order Extension of the Chern–Simons Theory," Izv. Vuz. Fiz. 60, 40 (2017) doi:10.1007/s11182-018-1331-8
- [85] D. S. Kaparulin, I. Y. Karataeva and S. L. Lyakhovich, "Extension of the Chern-Simons Theory: Conservation Laws, Lagrange Structures, and Stability," Russ. Phys. J. 59, 1930 (2017) doi:10.1007/s11182-017-0997-7

- [86] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Multi-Hamiltonian formulations and stability of higher-derivative extensions of 3d Chern–Simons," Eur. Phys. J. C 78, 115 (2018) doi:10.1140/epjc/s10052-018-5601-y [arXiv:1711.07897 [hep-th]]
- [87] D. S. Kaparulin and S. L. Lyakhovich, "On the Equivalence of Two Approaches to the Construction of Interactions in Higher-Derivative Theories," Russ. Phys. J. 59, 2041 (2017) doi:10.1007/s11182-017-1012-z
- [88] D. S. Kaparulin, I. Y. Karataeva and S. L. Lyakhovich, "Third order extensions of 3d Chern-Simons interacting to gravity: Hamiltonian formalism and stability," Nucl. Phys. B 934, 634 (2018) doi:10.1016/j.nuclphysb.2018.08.001 [arXiv:1806.00936 [hep-th]]
- [89] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Stable interactions between the extended Chern-Simons theory and a charged scalar field with higher derivatives: Hamiltonian formalism," Russ. Phys. J. 62, 12 (2019) doi:10.1007/s11182-019-01677-0 [arXiv:1905.00263 [hep-th]]
- [90] D. S. Kaparulin, "Conservation Laws and Stability of Field Theories of Derived Type," Symmetry 11, 642 (2019) doi:10.3390/sym11050642 [arXiv:1907.03068 [hep-th]]
- [91] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Conservation laws and stability of higher derivative extended Chern-Simons," J. Phys. Conf. Ser. 1337, 012001 (2019) doi:10.1088/1742-6596/1337/1/012001 [arXiv:1907.02267 [hep-th]]
- [92] V. A. Abakumova, D. S. Kaparulin and S. L. Lyakhovich, "Stable Interactions between Higher Derivative Extended Chern-Simons and Charged Scalar Field," AIP Conf. Proc. 2163, 090001 (2019) doi:10.1063/1.5130123 [arXiv:1907.08075 [hep-th]]
- [93] D. S. Kaparulin, "A stable higher-derivative theory with the Yang-Mills gauge symmetry," [arXiv:2011.12928 [hep-th]]
- [94] F. Magri, "A Simple model of the integrable Hamiltonian equation," J. Math. Phys. 19, 1156 (1978) doi:10.1063/1.523777
- [95] P. J. Olver, "Applications of Lie Groups to Differential Equations," Springer-Verlag (1993) doi:10.1007/978-1-4612-4350-2
- [96] D. S. Kaparulin, S. L. Lyakhovich and O. D. Nosyrev, "Resonance and stability of higher derivative theories of a derived type," Phys. Rev. D 101, 125004 (2020) doi:10.1103/PhysRevD.101.125004 [arXiv:2003.10860 [hep-th]]
- [97] C. Deffayet, S. Mukohyama and A. Vikman, "Ghosts without Runaway Instabilities," Phys. Rev. Lett. 128, 041301 (2022) doi:10.1103/PhysRevLett.128.041301 [arXiv:2108.06294 [gr-qc]]
- [98] H. Hopf and W. Rinow, "Ueber den Begriff der vollstÀndigen differentialgeometrischen FlÀche," Commentarii Mathematici Helvetic 3, 209 (1931) doi:10.1007/BF01601813
- [99] R. S. Palais, "A global formulation of the Lie theory of transformation groups," Memoirs of the American Mathematical Society 22, 123 (1957) doi.org/10.1090/memo/0022

- [100] E. A. Ivanov, A. V. Smilga and B. M. Zupnik, "Renormalizable supersymmetric gauge theory in six dimensions," Nucl. Phys. B 726, 131 (2005) doi:10.1016/j.nuclphysb.2005.08.014 [arXiv:hep-th/0505082 [hep-th]]
- [101] A. V. Smilga, "6D superconformal theory as the theory of everything," Gribov Memorial Volume, 443 (2006) doi:10.1142/9789812773784\_0038 [arXiv:hep-th/0509022 [hep-th]]
- [102] A. V. Smilga, "On dynamics of 5-D superconformal theories," Phys. Atom. Nucl. 70, 960 (2007) doi:10.1134/S1063778807050183 [arXiv:hep-th/0602201 [hep-th]]
- [103] A. V. Smilga, "Chiral anomalies in higher-derivative supersymmetric 6D theories," Phys. Lett. B 647, 298 (2007) doi:10.1016/j.physletb.2007.02.002 [arXiv:hep-th/0606139 [hep-th]]
- [104] A. V. Smilga, "On exactly solvable ghost-ridden systems," Phys. Lett. A 389, 127104 (2021) doi:10.1016/j.physleta.2020.127104 [arXiv:2008.12966 [hep-th]]
- [105] M. Toda, "Vibration of a Chain with Nonlinear Interaction," J. Phys. Soc. Jpn. 22, 431 (1967) doi:10.1143/JPSJ.22.431
- [106] D. J. Korteweg and G. de Vries, "On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves," Lon. Edi. Dub. Phil. M. J. Sci. 39, 422 (1895) doi:10.1080/14786449508620739
- [107] A. Smilga, "Modified Korteweg-de Vries equation as a system with benign ghosts," Acta Polytech.
  62, 190 (2022) doi:10.14311/AP.2022.62.0190 [arXiv:2112.14120 [hep-th]]
- [108] A. D. Linde, "Universe multiplication and the cosmological constant problem," Phys. Lett. B 200, 272 (1988) doi:10.1016/0370-2693(88)90770-8
- [109] S. M. Carroll, M. Hoffman and M. Trodden, "Can the dark energy equation-of-state parameter w be less than -1?," Phys. Rev. D 68, 023509 (2003) doi:10.1103/PhysRevD.68.023509 [arXiv:astro-ph/0301273 [astro-ph]]
- [110] D. E. Kaplan and R. Sundrum, "A Symmetry for the cosmological constant," JHEP 07, 042 (2006) doi:10.1088/1126-6708/2006/07/042 [arXiv:hep-th/0505265 [hep-th]]
- [111] A. Salvio, "Metastability in Quadratic Gravity," Phys. Rev. D 99, 103507 (2019) doi:10.1103/PhysRevD.99.103507 [arXiv:1902.09557 [gr-qc]]
- [112] C. Gross, A. Strumia, D. Teresi and M. Zirilli, "Is negative kinetic energy metastable?," Phys. Rev. D 103, 115025 (2021) doi:10.1103/PhysRevD.103.115025 [arXiv:2007.05541 [hep-th]]
- [113] M. Pavšič, "Quantum Field Theories in Spaces with Neutral Signatures," J. Phys. Conf. Ser. 437, 012006 (2013) doi:10.1088/1742-6596/437/1/012006 [arXiv:1210.6820 [hep-th]]
- [114] M. Pavšič, "Stable Self-Interacting Pais-Uhlenbeck Oscillator," Mod. Phys. Lett. A 28, 1350165 (2013) doi:10.1142/S0217732313501654 [arXiv:1302.5257 [gr-qc]]
- [115] M. Pavšič, "Pais-Uhlenbeck Oscillator with a Benign Friction Force," Phys. Rev. D 87, 107502 (2013) doi:10.1103/PhysRevD.87.107502 [arXiv:1304.1325 [gr-qc]]

- [116] M. Pavšič, "Pais–Uhlenbeck oscillator and negative energies," Int. J. Geom. Meth. Mod. Phys. 13, 1630015 (2016) doi:10.1142/S0219887816300154 [arXiv:1607.06589 [gr-qc]]
- [117] A. Fring and B. Turner, "Higher derivative Hamiltonians with benign ghosts from affine Toda lattices," J. Phys. A 56, 295203 (2023) doi:10.1088/1751-8121/ace0e6 [arXiv:2301.11317 [hep-th]]
- [118] A. Fring, T. Taira and B. Turner, "Nonlinear evolution of disturbances in higher time-derivative theories," [arXiv:2406.18255 [nlin.SI]]
- [119] N. H. Barth and S. M. Christensen, "Quantizing fourth-order gravity theories: The functional integral," Phys. Rev. D 28, 1876 (1983) doi:10.1103/PhysRevD.28.1876
- [120] H. A. Weldon, "Quantization of higher-derivative field theories," Ann. Phys. 305, 137 (2003) doi:10.1016/S0003-4916(03)00070-8
- [121] A. V. Smilga, "Comments on the dynamics of the Pais-Uhlenbeck oscillator," SIGMA 5, 017 (2009) doi:10.3842/Sigma.2009.017 [arXiv:0808.0139 [quant-ph]]
- [122] I. B. Ilhan and A. Kovner, "Some Comments on Ghosts and Unitarity: The Pais-Uhlenbeck Oscillator Revisited," Phys. Rev. D 88, 044045 (2013) doi:10.1103/PhysRevD.88.044045 [arXiv:1301.4879 [hep-th]]
- [123] R. P. Woodard, "Ostrogradsky's theorem on Hamiltonian instability," Scholarpedia 10, 32243 (2015) doi:10.4249/scholarpedia.32243 [arXiv:1506.02210 [hep-th]]
- [124] A. Salvio and A. Strumia, "Quantum mechanics of 4-derivative theories," Eur. Phys. J. C 76, 227 (2016) doi:10.1140/epjc/s10052-016-4079-8 [arXiv:1512.01237 [hep-th]]
- [125] B. Holdom, "Making sense of ghosts," [arXiv:2408.04089 [hep-th]]
- [126] A. Strumia, "Interpretation of quantum mechanics with indefinite norm," MDPI Physics 1, 17 (2019) doi:10.3390/physics1010003 [arXiv:1709.04925 [quant-ph]]
- [127] P. Romatschke, "Out-Of-Time-Ordered-Correlators for the Pure Inverted Quartic Oscillator: Classical Chaos meets Quantum Stability," [arXiv:2408.12643 [hep-th]]
- [128] P. O. Kazinski, S. L. Lyakhovich and A. A. Sharapov, "Lagrange structure and quantization," JHEP 07, 076 (2005) doi:10.1088/1126-6708/2005/07/076 [arXiv:hep-th/0506093 [hep-th]]
- [129] S. L. Lyakhovich and A. A. Sharapov, "Schwinger-Dyson equation for non-Lagrangian field theory," JHEP 02, 007 (2006) doi:10.1088/1126-6708/2006/02/007 [arXiv:hep-th/0512119 [hep-th]]
- [130] S. L. Lyakhovich and A. A. Sharapov, "Quantizing non-Lagrangian gauge theories: An Augmentation method," JHEP 01, 047 (2007) doi:10.1088/1126-6708/2007/01/047 [arXiv:hep-th/0612086 [hep-th]]
- [131] C. M. Bender and P. D. Mannheim, "No-ghost theorem for the fourth-order derivative Pais-Uhlenbeck oscillator model," Phys. Rev. Lett. 100, 110402 (2008) doi:10.1103/PhysRevLett.100.110402 [arXiv:0706.0207 [hep-th]]
- [132] C. M. Bender and P. D. Mannheim, "Exactly solvable PT-symmetric Hamiltonian having no Hermitian counterpart," Phys. Rev. D 78, 025022 (2008) doi:10.1103/PhysRevD.78.025022 [arXiv:0804.4190 [hep-th]]

- [133] C. M. Bender and P. D. Mannheim, "Giving up the ghost," J. Phys. A 41, 304018 (2008) doi:10.1088/1751-8113/41/30/304018 [arXiv:0807.2607 [hep-th]]
- [134] C. M. Bender and D. W. Hook, "PT-symmetric quantum mechanics," [arXiv:2312.17386 [quant-ph]]
- [135] S. Kawasaki and T. Kimura, "A Possible Mechanism of Ghost Confinement in a Renormalizable Quantum Gravity," Prog. Theor. Phys. 65, 1767 (1981) doi:10.1143/PTP.65.1767
- [136] N. Arkani-Hamed, H. C. Cheng, M. A. Luty and S. Mukohyama, "Ghost condensation and a consistent infrared modification of gravity," JHEP 05, 074 (2004) doi:10.1088/1126-6708/2004/05/074 [arXiv:hep-th/0312099 [hep-th]]
- [137] S. Mukohyama, "Ghost condensate and generalized second law," JHEP 09, 070 (2009) doi:10.1088/1126-6708/2009/09/070 [arXiv:0901.3595 [hep-th]]
- [138] J. F. Donoghue, "Quartic propagators, negative norms and the physical spectrum," Phys. Rev. D 96, 044007 (2017) doi:10.1103/PhysRevD.96.044007 [arXiv:1704.01533 [hep-th]]
- [139] M. Frasca, A. Ghoshal and A. S. Koshelev, "Confining complex ghost degrees of freedom," Phys. Lett. B 841, 137924 (2023) doi:10.1016/j.physletb.2023.137924 [arXiv:2207.06394 [hep-th]]
- [140] J. Liu, L. Modesto and G. Calcagni, "Quantum field theory with ghost pairs," JHEP 02, 140 (2023) doi:10.1007/JHEP02(2023)140 [arXiv:2208.13536 [hep-th]]
- [141] E. N. Lorenz, "Deterministic nonperiodic flow," J. Atmos. Sci. 20, 130 (1963) doi:10.1175/1520-0469(1963)020<0130:DNF>2.0.CO;2
- [142] E. Schechter, "Handbook of Analysis and Its Foundations," Elsevier Academic (1997) doi:10.1016/B978-0-12-622760-4.X5000-6 (Available online at https://staffwww.dcs.shef.ac.uk/people/R.Chisholm/docs/)
- [143] L. A. Steen and J. A. Seebach, "Counterexamples in Topology," Springer New York (1978) doi:10.1007/978-1-4612-6290-9
- [144] R. Williamson and L. Janos, "Constructing metrics with the Heine-Borel property," Proc. Amer. Math. Soc. 100, 567 (1987) doi:10.1090/S0002-9939-1987-0891165-X
- [145] P. D. Lax, "Functional analysis," Wiley (2002) ISBN 978-0-471-55604-6 (Available online at http://users.math.uoc.gr/ frantzikinakis/FunctionalGrad2015/Lax.pdf)
- [146] J. M. Lee, "Introduction to Smooth Manifolds," Springer (2012) doi.org/10.1007/978-1-4419-9982-5
- [147] M. Asorey, L. Rachwal and I. L. Shapiro, "Unitary Issues in Some Higher Derivative Field Theories," Galaxies 6, 23 (2018) doi:10.3390/galaxies6010023 [arXiv:1802.01036 [hep-th]]
- [148] Z. Liu and M. Tegmark, "Machine Learning Conservation Laws from Trajectories," Phys. Rev. Lett. 126, 180604 (2021) doi:10.1103/PhysRevLett.126.180604 [arXiv:2011.04698 [cs.LG]]
- [149] S. Ha and H. Jeong, "Discovering invariants via machine learning," Phys. Rev. Research 3, L042035 (2021) doi:10.1103/PhysRevResearch.3.L042035 [arXiv:2102.04008 [cs.LG]]

- [150] Z. Liu, V. Madhavan and M. Tegmark, "Machine learning conservation laws from differential equations," Phys. Rev. E 106, 045307 (2022) doi:10.1103/PhysRevE.106.045307 [arXiv:2203.12610 [cs.LG]]
- [151] V. I. Arnold, V. V. Kozlov and A. I. Neishtadt, "Mathematical Aspects of Classical and Celestial Mechanics," Springer (2006) doi:10.1007/978-3-540-48926-9
- [152] M. C. Muñoz-Lecanda and N. Román-Roy, "Geometry of Mechanics," [arXiv:2401.12650 [math-ph]]
- [153] W. Pauli, "Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik," Z. Physik
   36, 336 (1926) doi:10.1007/BF01450175
- [154] J. Baez, "Mysteries of the gravitational 2-body problem," Personal webpage: https://math.ucr.edu/home/baez/gravitational.html
- [155] A. Bolsinov, VS. Matveev, E. Miranda and S. Tabachnikov "Open problems, questions and challenges in finite-dimensional integrable systems," Phil. Trans. R. Soc. A 376, 20170430 (2018) doi:10.1098/rsta.2017.0430 [arXiv:1804.03737 [math.DS]]
- [156] H. S. Dumas, "The KAM story: a friendly introduction to the content, history, and significance of classical Kolmogorov-Arnold-Moser theory," World Scientific Publishing (2014) doi:10.1142/8955
- [157] E. Fioranil, G. Giachetta and G. Sardanashvily, "The Liouville-Arnold-Nekhoroshev theorem for noncompact invariant manifolds," J. Phys. A: Math. Gen. 36, L101 (2003) doi:10.1088/0305-4470/36/7/102 [arXiv:math/0210346 [math.DS]]
- [158] J. Liouville, "Note sur l'intégration des équations différentielles de la dynamique," J. Mat. Pur. App. 20, 137 (1855) http://www.numdam.org/item/JMPA\_1855\_1\_20\_137\_0/
- [159] V. I. Arnold, "Mathematical Methods of Classical Mechanics," Springer-Verlag (1978) doi:10.1007/978-1-4757-1693-1
- [160] A. Mostafazadeh, "A Hamiltonian Formulation of the Pais-Uhlenbeck Oscillator that Yields a Stable and Unitary Quantum System," Phys. Lett. A 375, 93 (2010) doi:10.1016/j.physleta.2010.10.050 [arXiv:1008.4678 [hep-th]]
- [161] G. Darboux, "Sur un problème de mécanique," Archives Néerlandaises des Sciences Exactes et Naturelles (1901) https://www.biodiversitylibrary.org/item/159022
- [162] J. F. Barbero G., M. Basquens, B. Díaz and E. J. S. Villaseñor, "Poisson brackets in Sobolev spaces: a mock holonomy-flux algebra," Phys. Scripta 97, 125202 (2022) doi:10.1088/1402-4896/ac99a9 [arXiv:2207.00342 [math-ph]]
- [163] J. F. Barbero G., "Free scalar field theory on a Sobolev space over a bounded interval," Gen. Rel. Grav. 56, 69 (2024) doi:10.1007/s10714-024-03256-2 [arXiv:2405.17996 [hep-th]]
- [164] J. M. Lee, "Introduction to Topological Manifolds," Springer (2011) doi:10.1007/978-1-4419-7940-7
- [165] R. Brouzet, R. Caboz, J. Rabenivo and V. Ravoson, "Two degrees of freedom quasi-bi-Hamiltonian systems," J. Phys. A: Math. Gen. 29, 2069 (1996) doi:10.1088/0305-4470/29/9/019