

# 3-manifold spine cyclic presentations with seldom seen Whitehead graphs

Gerald Williams

September 2, 2024

## Abstract

We consider two families of cyclic presentations and show that, subject to certain conditions on the defining parameters, they are spines of closed 3-manifolds. For the first family, the Whitehead graphs have not previously been observed in this context, and the corresponding manifolds are lens spaces. The second family provides new examples where the reduced Whitehead graphs are those of the Fractional Fibonacci presentations; here the corresponding manifolds are often (but not always) hyperbolic.

**Keywords:** 3-manifold spine, Whitehead graph, cyclically presented group, Fractional Fibonacci group.

**MSC:** 20F05, 57M05, 57M50.

## 1 Introduction

The *cyclically presented group*  $G_n(w)$  is the group defined by the *cyclic presentation*

$$\mathcal{G}_n(w) = \langle x_0, \dots, x_{n-1} \mid w, \theta(w), \dots, \theta^{n-1}(w) \rangle$$

where  $w(x_0, \dots, x_{n-1})$  is a cyclically reduced word in the free group  $F_n$  with generators  $x_0, \dots, x_{n-1}$  and  $\theta : F_n \rightarrow F_n$  is the *shift automorphism* of  $F_n$  given by  $\theta(x_i) = x_{i+1}$  for each  $0 \leq i < n$  (subscripts mod  $n$ ,  $n > 0$ ). Cyclic presentations that are spines of 3-manifolds have been widely researched, with notable early studies by Dunwoody [18], Sieradski [50], Helling, Kim, Mennicke [22], and Cavicchioli and Spaggiari [16].

A necessary condition for a presentation to be a spine of a closed 3-manifold is that its Whitehead graph is planar (see, for example, [25, page 33], [4, Documentation, Section 11]). The planar, reduced, Whitehead graphs of cyclically reduced cyclic presentations were classified in [27], whose labelling (I.j), (II.j), (III.j) we now adopt. In the overwhelming majority of studies of cyclic presentations as spines of 3-manifolds (for example [1, 2, 7, 8, 9, 10, 11, 12, 13, 15, 18, 20, 21, 26, 33, 34, 35, 36, 37, 38, 42, 44, 48, 50, 51, 52, 53]), the reduced Whitehead graph is of one of the forms (I.1) or (I.3) (which correspond to the graphs given by Dunwoody in [18, Figure 1]) and, as such, provide examples of so-called *Dunwoody manifolds*. Cyclic presentations that are spines of 3-manifolds where the reduced Whitehead graph is of a different form to these are few and far between. We are only aware of the following families of such presentations. The non-cyclically reduced cyclic presentation  $\mathcal{G}_n(x_0x_1x_1^{-1})$  is a spine of  $S^3$  for all  $n \geq 1$  ([26, Lemma 9]). Its Whitehead graph contains loops, but once these are removed the graph is of type (I.4). Helling, Kim and Mennicke [22] and Cavicchioli and Spaggiari [16, Theorem 3] showed that, for even  $n$ , the Fibonacci presentations  $\mathcal{F}(n) = \mathcal{G}_n(x_0x_1x_2^{-1})$  are spines of closed, connected, oriented 3-manifolds; here the Whitehead graph is of type (II.11). A special case of

[47, Theorem 6] gives that, for coprime integers  $k, l \geq 1$  and even  $n$ , the cyclic presentations  $\mathcal{G}_n(x_0^l x_1^k x_2^{-l})$  are spines of closed, connected, oriented 3-manifolds. These presentations are the *Fractional Fibonacci presentations*  $\mathcal{F}^{k/l}(n)$  of [55, 56] (or of [40, 41] in the case  $l = 1$ ) and if  $(k, l) \neq (1, 1)$  their reduced Whitehead graph is of type (II.7). Jeong and Wang [28, 29] showed that, for  $l \geq 2$  and even  $n$  the cyclic presentations  $\mathcal{G}_n(x_0(x_2^{-1}x_1)^l)$  are spines of closed, connected, oriented 3-manifolds; here the reduced Whitehead graph is of type (II.14).

In Theorem A we present a family of cyclic presentations that are spines of closed 3-manifolds where the Whitehead graphs are of type (I.5) and show that the manifolds are cyclic branched covers of a lens space. This construction is essentially well known (see, for example [49, page 217], [43, page 4], or [39, Section 3]), but the connection to the planar Whitehead graph classification of [27] has not previously been observed. In Theorem B we present a new family of cyclic presentations that are spines of closed 3-manifolds where the reduced Whitehead graphs are of type (II.7) (i.e. the same type as those of the Fractional Fibonacci presentations) and show that many, but not all, of the manifolds are hyperbolic. Experiments in Heegaard [4] were used in formulating these results.

## 2 Presentation complexes as spines of closed 3-manifolds

The *presentation complex* (or *cellular model*)  $K = K_{\mathcal{P}}$  of a group presentation  $\mathcal{P} = \langle X \mid R \rangle$  is the 2-complex with one 0-cell  $O$ , a loop at  $O$  for each generator  $x \in X$  and a 2-cell for each relator (the boundary of that 2-cell spelling the relator). If  $N$  is a regular neighbourhood of  $O$  then  $K \cap \partial N$  is a 1-dimensional cell complex called the *Whitehead graph* or *link graph* of  $\mathcal{P}$ . Thus the Whitehead graph of  $\mathcal{P}$  is the graph with  $2|X|$  vertices  $v_x, v'_x$  ( $x \in X$ ) and an edge  $(v_x, v_y)$  (resp.  $(v'_x, v'_y), (v_x, v'_y)$ ) for each occurrence of a cyclic subword  $xy^{-1}$  (resp.  $x^{-1}y, (xy)^{\pm 1}$ ) in a relator  $r \in R$ . The *reduced Whitehead graph* is the graph obtained from the Whitehead graph by replacing all multiedges between two vertices by a single edge. We say that a group presentation  $\mathcal{P}$  is the *spine* of a closed, connected, 3-manifold  $M$  if there exists a 3-ball  $B^3 \subset M$  such that  $M - \mathring{B}^3$  collapses onto  $K_{\mathcal{P}}$  (where  $\mathring{B}^3$  denotes the interior of  $B^3$ ).

Suppose that a group presentation  $\mathcal{P} = \langle X \mid R \rangle$  with an equal number of generators and relators is the spine of a closed, connected, oriented 3-manifold. Then (see [49, Chapter 9], [45], [50, page 125]) there is a 3-complex  $C$  whose set of faces consists of precisely one pair of oppositely oriented faces  $F_r^+, F_r^-$  for each relator  $r \in R$ , whose boundaries spell  $r$ . Let  $M_0$  denote the 3-complex obtained from  $C$  by identifying the faces  $F_r^+, F_r^-$  ( $r \in R$ ). Then  $M_0$  is a closed, connected, oriented pseudo-manifold. The *Seifert-Threlfall condition* for  $M_0$  to be a manifold  $M$  is that its Euler characteristic is zero [49, Theorem I, Section 60]. In this case the cell structure on  $M = M_0$  has one vertex, one 3-cell, and 2-skeleton homeomorphic to the presentation complex  $K_{\mathcal{P}}$  of  $\mathcal{P}$ . The Whitehead graph  $\Gamma$  of  $\mathcal{P}$  is the link of the single vertex of  $K_{\mathcal{P}}$ , and so embeds in the link of the single vertex of  $M$ . The link of the vertex of  $M$  is the 2-sphere, and so  $\Gamma$  has a planar embedding on this sphere. When  $\Gamma$  is connected the 3-complex  $C$  is a polyhedron  $\pi$  bounding a 3-ball. Given a planar embedding of  $\Gamma$  there is a one-to-one correspondence between the faces  $F$  of this embedding and the vertices  $u_F$  of  $\pi$ . This correspondence maps a face of degree  $d$  to a vertex of degree  $d$  of  $\pi$ . Moreover, if the vertices in the face read cyclically around the face are  $w_1, \dots, w_d$  where  $w_i \in \{v_x, v'_x \mid x \in X\}$  then the arcs  $e_1, \dots, e_d$  incident to  $u_F$ , read cyclically, are directed towards  $u_F$  if  $w_i = v_x$  and away from  $u_F$  if  $w_i = v'_x$ , and the arc  $e_j$ , corresponding to vertex  $w_j = v_x$  or  $v'_x$ , is labelled  $x$ .




Figure 1: Whitehead graph for  $\mathcal{H}(r, n)$  (where  $(r, n) = 1$ )

### 3 Cyclic presentations $\mathcal{H}(r, n)$ as spines of type (I.5)

For  $n > 1, r \geq 1$  let

$$\mathcal{H}(r, n) = \mathcal{G}_n(x_0 x_1 \dots x_{r-1})$$

and let  $H(r, n)$  be the group it defines. By [54, Theorems 2,3],  $H(r, n)$  is finite if and only if  $(n, r) = 1$ , in which case  $H(r, n) \cong \mathbb{Z}_r$ . The Whitehead graph of  $\mathcal{H}(r, n)$  is connected if and only if  $(n, r) = 1$ , and in this case it is of type (I.5), as shown in Figure 1, where (as also in Figure 5) the edge labels denote their multiplicities, a vertex label  $i$  denotes  $v_{x_i}$ , and a vertex label  $\bar{i}$  denotes  $v'_{x_i}$ .

**Theorem A.** *Let  $n \geq 2, r \geq 2$  and suppose  $(n, r) = 1$ . Then the cyclic presentation  $\mathcal{H}(r, n)$  is a spine of a closed, connected, oriented 3-manifold  $M$ , which is an  $n$ -fold cyclic branched cover of the lens space  $L(r, 1)$ .*

*Proof.* Consider the polyhedron in Figure 2 with face pairing given by identifying the faces  $F_i^+, F_i^-$  ( $0 \leq i < n$ ). We shall show that the identification of faces results in a 3-complex  $M$  whose 2-skeleton is the presentation complex of  $\mathcal{H}(r, n)$ , so  $M$  has one 0-cell,  $n$  1-cells,  $n$  2-cells, and one 3-cell, so is a manifold by [49, Theorem I, Section 60].

Let  $[S, N]_i$  denote the arc of the face  $F_i^+$  with initial vertex  $S$  and terminal vertex  $N$ . All the arcs labelled  $x_0$  are contained in the following cycle (of length  $r$ ):

$$\begin{aligned} [S, N]_0 &\xrightarrow{F_{n-(r-1)}} [w_{n-(r-1)}^2, w_{n-(r-1)}^1] \xrightarrow{F_{n-(r-2)}} [w_{n-(r-2)}^3, w_{n-(r-2)}^2] \\ &\xrightarrow{F_{n-(r-3)}} [w_{n-(r-3)}^4, w_{n-(r-3)}^3] \longrightarrow \dots \xrightarrow{F_{n-2}} [w_{n-2}^{r-1}, w_{n-2}^{r-2}] \\ &\xrightarrow{F_{n-1}} [N, w_{n-1}^{r-1}] \xrightarrow{F_0} [S, N]_0. \end{aligned}$$

All the vertices that are a vertex (either initial or terminal) of an arc labelled  $x_0$  are contained in the (induced) cycle of the initial vertices in the arcs above, so in the resulting complex  $M$  these vertices are identified. In particular, vertices  $N, S$  are identified, and (by comparing initial vertices) for  $1 < j < r$  vertices  $w_{n-j}^{r-j+1}$  are identified with  $S$ . Therefore, for each  $0 \leq i < n$ ,  $1 < j < r$  the vertex  $w_i^j = \theta^{i+r-j+1}(w_{n-(r-j+1)}^j)$  is identified with  $S$ . Therefore all the vertices of the polyhedron are identified. Thus the quotient  $M$  has one 3-cell,  $n$  2-cells,  $n$  1-cells, and one 0-cell, and since the boundaries of the 2-cells spell the relators of  $\mathcal{H}(r, n)$ , it follows that  $M$  is a closed, connected, oriented 3-manifold, and  $\mathcal{H}(r, n)$  is the spine of  $M$ .

As in (for example) [57, proof of Proposition 2], a Heegaard diagram for  $M$  arising from the face pairing polyhedron is given in Figure 3. This diagram has a rotational symmetry  $\rho$  of order  $n$  cyclically permuting the faces  $F_i^+ \rightarrow F_{i+r}^+$  and  $F_i^- \rightarrow F_{i+r}^-$ . The quotient of  $M$  by  $\rho$  is




Figure 2: Face pairing polyhedron for  $\mathcal{H}(r, n)$




Figure 3: Heegaard diagram for the manifold  $M(r, n)$




Figure 4: Heegaard diagram for the manifold  $M(r, n)/\rho$




Figure 5: Whitehead graph for  $\mathcal{G}^{k/l}(n, f)$  (where  $\lambda = 2l + k - 3$ ,  $n$  even and  $fk \equiv 0$  or  $2 \pmod n$ )

a 3-orbifold in which the image of the rotation axis is a singular set. This yields the Heegaard diagram for  $M/\rho$  in Figure 4, which is the canonical diagram of the lens space  $L(r, 1)$ . Hence the manifold  $M$  is the branched cyclic  $n$ -fold cover of  $L(r, 1)$ , as required.  $\square$

#### 4 Cyclic presentations $\mathcal{G}^{k/l}(n, f)$ as spines of type (II.7)

For  $n \geq 2$ ,  $k, l \geq 1$ ,  $0 \leq f < n$ , let  $\mathcal{G}^{k/l}(n, f)$  be the cyclic presentation

$$\mathcal{G}_n((y_0 y_f \cdots y_{(l-1)f})(y_{lf+1} y_{(l+1)f+1} \cdots y_{(l+(k-1)f+1}))(y_{2f} y_{2f+f} \cdots y_{2+(l-1)f})^{-1})$$

and let  $G^{k/l}(n, f)$  be the group it defines. When  $n \geq 4$ , the Whitehead graph of  $\mathcal{G}^{k/l}(n, f)$  is planar if and only if  $n$  is even and either  $fk \equiv 0 \pmod n$  or  $fk \equiv 2 \pmod n$ , in which case it is of type (II.7) if  $(k, l) \neq (1, 1)$  and is of type (II.11) if  $k = l = 1$ ; see Figure 5. Our main result for this section is the following:

**Theorem B.** *Let  $n \geq 4$ ,  $fk \not\equiv 2 \pmod n$ , and  $(k, l) \in \{(k, 1), (1, l), (5, 2), (2, 5)\}$ . Then  $\mathcal{G}^{k/l}(n, f)$  is the spine of a closed, connected, oriented 3-manifold  $M^{k/l}(n, f)$  if and only if  $n$  and  $f$  are even and  $fk \equiv 0 \pmod n$ .*

Our motivation for studying these presentations (and, in particular, under the condition  $fk \equiv 0 \pmod n$  but not under the condition  $fk \equiv 2 \pmod n$ ) stems from the following connection to the Fractional Fibonacci groups. The presentations  $\mathcal{G}^{k/l}(n, 0)$  are the Fractional Fibonacci group presentations

$$\mathcal{F}^{k/l}(n) = \mathcal{G}_n(x_0^l x_1^k x_2^{-l})$$

introduced in [40, 41, 55, 56], generalizing the Fibonacci group presentations  $\mathcal{F}^{1/1}(n)$ . For even  $n$  and coprime integers  $k, l \geq 1$ , the Fractional Fibonacci group  $F^{k/l}(n)$  is a 3-manifold group [40, Theorem 4.1], [41, Section 4], [55, Section 2], [56, Section 2]. If  $n \geq 2$  is even then  $\mathcal{F}^{1/1}(n)$  was shown to be the spine of a closed, connected, oriented, 3-manifold in [16],[22],[23],[24]. More generally, if  $k, l \geq 1$  are coprime and  $n$  is even, then setting  $s_i = q_i = l$ ,  $p_i = -r_i = k$  in [47, Theorem 6] gives that  $\mathcal{F}^{k/l}(n)$  is the spine of a closed, connected, oriented, 3-manifold. If  $n$  is odd then  $\mathcal{F}^{k/l}(n)$  is not a spine, since its Whitehead graph is non-planar. The question as to when, for odd  $n$ ,  $F^{k/l}(n)$  is a 3-manifold group is considered in [26, Theorem 3] for the case  $k = l = 1$ , and in [17, Theorem 6.2] for the case  $l = 1$  and the case  $n = 3$ . The abelianisation  $F^{k/l}(n)^{\text{ab}}$  is obtained in [41, Lemma 1], [46, Corollary 4.3].

The *shift extension*  $E^{k/l}(n) = F^{k/l}(n) \rtimes_{\theta} \mathbb{Z}_n$  of  $F^{k/l}(n)$  by  $\mathbb{Z}_n = \langle t \mid t^n \rangle$  has the presentation

$$E^{k/l}(n) = \langle x, t \mid t^n, x^l t x^k t x^{-l} t^{-2} \rangle, \quad (1)$$

where the second relator is obtained by rewriting the relators of  $F^{k/l}(n)$  in terms of the substitutions  $x_i = t^i x t^{-i}$  (see, for example, [31, Theorem 4]). As set out in [5, Section 2], for  $0 \leq f < n$  such that  $fk \equiv 0 \pmod n$ , there is a retraction  $\nu^f : E^{k/l}(n) \rightarrow \mathbb{Z}_n = \langle t \mid t^n \rangle$  given by  $\nu^f(t) = t$ ,  $\nu^f(x) = t^f$ . The kernel,  $\ker \nu^f$ , has a presentation with generators  $y_i = t^i x t^{-(i+f)}$  ( $0 \leq i < n$ ) and relators that are rewrites of conjugates of the second relator of  $E^{k/l}(n)$  by powers of  $t$ , and so has the cyclic presentation  $\mathcal{G}^{k/l}(n, f)$ . In particular (under the condition  $fk \equiv 0 \pmod n$ ),

$$\begin{aligned}\mathcal{G}^{k/1}(n, 0) &= \mathcal{G}_n(y_0 y_1^k y_2^{-1}) = \mathcal{F}^{k/1}(n), \\ \mathcal{G}^{1/l}(n, f) &= \mathcal{G}^{1/l}(n, 0) = \mathcal{G}_n(y_0^l y_1 y_2^{-l}) = \mathcal{F}^{1/l}(n), \\ \mathcal{G}^{5/2}(n, f) &= \mathcal{G}_n((y_0 y_f)(y_{2f+1} y_{3f+1} y_{4f+1} y_{1f+1})(y_2 y_{2+f})^{-1}), \\ \mathcal{G}^{2/5}(n, f) &= \mathcal{G}_n((y_0 y_f y_0 y_f y_0)(y_{f+1} y_1)(y_2 y_{2+f} y_2 y_{2+f} y_2)^{-1}).\end{aligned}$$

The shift extension

$$\begin{aligned}G^{k/l}(n, f) \rtimes_{\theta} \mathbb{Z}_n &= \langle y, t \mid t^n, (y t^f)^l t (y t^f)^k t^{-fk+1} (y t^f)^{-l} t^{-2} \rangle \\ &= \langle x, t \mid t^n, x^l t x^k t^{-fk+1} x^{-l} t^{-2} \rangle\end{aligned}$$

(by setting  $x = y t^f$  and eliminating  $y$ ). In the case  $fk \equiv 0 \pmod n$ , this coincides with  $E^{k/l}(n)$ , which is independent of the value of  $f$ . This implies, for example, that the order of  $G^{k/l}(n, f)$  is independent of  $f$ , the shift dynamics of  $G^{k/l}(n, f)$  are identical for all values of  $f$  [5, Lemma 2.2], and that  $G^{k/l}(n, f)$  is (non-elementary) word hyperbolic if and only if  $F^{k/l}(n)$  is (non-elementary) word hyperbolic. In Theorem 4.4 we similarly show that, for fixed  $k, l, n$ , if two groups  $G^{k/l}(n, f_1), G^{k/l}(n, f_2)$  sharing a shift extension are fundamental groups of closed, connected, orientable 3-manifolds then either both manifolds are hyperbolic, or neither are. On the other hand, for fixed  $k, l, n$ , two groups  $G^{k/l}(n, f_1), G^{k/l}(n, f_2)$  sharing a shift extension will typically be non-isomorphic. We give examples of this in Example 4.1 and Lemma 4.2. Further, in the case  $fk \equiv 0 \pmod n$ , we also have that the presentation  $\mathcal{G}^{k/l}(n, f)$  has a planar Whitehead graph. Since the group  $G^{k/l}(n, f)$  shares its shift extension with that of  $F^{k/l}(n)$ , these properties combined suggest that the geometric properties of  $F^{k/l}(n)$  and  $\mathcal{F}^{k/l}(n)$  may be inherited by  $G^{k/l}(n, f)$  and  $\mathcal{G}^{k/l}(n, f)$ . However, while [47, Theorem 6] implies that  $\mathcal{F}^{k/l}(n)$  is a spine for all coprime  $k, l$  and even  $n \geq 2$ , Lemma 4.3 will show that, additionally,  $f$  must be even for  $\mathcal{G}^{k/l}(n, f)$  to be a spine. In the other planar case,  $fk \equiv 2 \pmod n$ , the groups  $G^{k/l}(n, f)$  do not share a shift extension with  $F^{k/l}(n)$ , so there is no a priori reason to expect similar geometric properties to hold.

**Example 4.1** (Non-cyclic finite groups.). The groups  $G^{3/1}(3, 0), G^{3/1}(3, 1)$ , are solvable of order 3528, and derived lengths 3, and 4, respectively ([41, Remark 1], [30, page 282]) and hence are non-isomorphic. The groups  $G^{3/2}(3, 0), G^{3/2}(3, 1)$ , are solvable of order 504, and derived lengths 2, and 3 respectively (using [19]) and hence are non-isomorphic.

In Lemma 4.2 we use the fact that the order of the abelianisation  $G^{k/l}(n, f)^{\text{ab}}$  is given by  $|\text{Res}(p_f(t), t^n - 1)|$ , if this is non-zero, and is infinite otherwise, where

$$p_f(t) = (1 - t^2)(1 + t^f + \dots + t^{(l-1)f}) + t^{lf+1}(1 + t^f + \dots + t^{(k-1)f})$$

is the *representer polynomial* of  $G^{k/l}(n, f)$ , and  $\text{Res}(\cdot, \cdot)$  denotes the resultant [32, page 82]. As we will only be interested in the absolute values of resultants (and not the sign), to avoid repetitive use of modulus signs we will take  $\text{Res}(\cdot, \cdot)$  to mean  $|\text{Res}(\cdot, \cdot)|$ .

**Lemma 4.2.** *Let  $n, k, l \geq 1$  where  $n, k$  are even and  $\gcd(k, l) = 1$  then  $F^{k/l}(n) = G^{k/l}(n, 0) \not\cong G^{k/l}(n, n/2)$ .*

*Proof.* Note that  $l$  is odd, since  $\gcd(k, l) = 1$  and  $k$  is even. For any  $0 \leq f < n$

$$\operatorname{Res}(p_f(t), t^n - 1) = \operatorname{Res}(p_f(t), t^{n/2} - 1) \cdot \operatorname{Res}(p_f(t), t^{n/2} + 1). \quad (2)$$

We have

$$\begin{aligned} \operatorname{Res}(p_{n/2}(t), t^{n/2} - 1) &= \operatorname{Res}(l(1 - t^2) + kt, t^{n/2} - 1) \\ &= \operatorname{Res}(p_0(t), t^{n/2} - 1), \end{aligned}$$

and, setting  $\alpha, \bar{\alpha} = (k \pm \sqrt{k^2 + 4l^2})/(2l)$ ,

$$\begin{aligned} \operatorname{Res}(p_0(t), t^{n/2} + 1) &= \operatorname{Res}\left(l(t - \alpha)(t - \bar{\alpha}), t^{n/2} + 1\right) \\ &= l^{n/2}(\alpha^{n/2} + 1)(\bar{\alpha}^{n/2} + 1) \\ &= l^{n/2} \left( (-1)^{n/2} + 1 + (\alpha^{n/2} + \bar{\alpha}^{n/2}) \right). \end{aligned}$$

On the other hand,

$$\begin{aligned} \operatorname{Res}(p_{n/2}(t), t^{n/2} + 1) &= \operatorname{Res}\left(l(1 - t^2) + kt(1 + t^{n/2})/2, t^{n/2} + 1\right) \\ &= \operatorname{Res}\left(l(1 - t^2), t^{n/2} + 1\right) \\ &= l^{n/2} \cdot 2(1 + (-1)^{n/2}). \end{aligned}$$

Therefore  $\operatorname{Res}(p_0(t), t^{n/2} + 1) \neq \operatorname{Res}(p_{n/2}(t), t^{n/2} + 1)$  so, by equation (2),  $\operatorname{Res}(p_0(t), t^n - 1) \neq \operatorname{Res}(p_{n/2}(t), t^n - 1)$ . Hence  $|G^{k/l}(n, 0)^{\text{ab}}| \neq |G^{k/l}(n, n/2)^{\text{ab}}|$ , and the result follows.  $\square$

**Lemma 4.3.** *Suppose  $n \geq 4$ ,  $k, l \geq 1$ ,  $0 \leq f < n$  where  $n$  is even,  $fk \equiv 0 \pmod{n}$ , and  $\gcd(k, l) = 1$ . If  $f$  is odd then  $\mathcal{G}^{k/l}(n, f)$  is not the spine of a closed, connected, oriented 3-manifold.*

*Proof.* First note that if  $f$  is odd, then the hypotheses imply that  $l$  is odd. The Whitehead graph of  $\mathcal{G}^{k/l}(n, f)$  has a planar embedding, as shown in Figure 5, which is unique up to self homeomorphism of  $S^2$ . It has two  $n/2$ -gons  $v'_{y_0} - v'_{y_2} - \dots - v'_{y_{n-2}} - v'_{y_0}$  and  $v'_{y_1} - v'_{y_3} - \dots - v'_{y_{n-1}} - v'_{y_1}$ ,  $n$  3-gons  $v_{y_i} - v_{y_{i+1}} - v'_{y_{i+f+1}} - v_{y_i}$ ,  $n$  4-gons  $v_{y_i} - v_{y_{i+1}} - v'_{y_{i+f+2}} - v'_{y_{i+f}} - v_{y_i}$ , and  $\lambda n$  2-gons  $v_{y_i} - v'_{y_{i+f+1}} - v_{y_i}$ , where  $\lambda = 2l + k - 3$ .

If  $\mathcal{G}^{k/l}(n, f)$  is a 3-manifold spine then in a putative corresponding face-pairing polyhedron, there are two degree  $n/2$  source vertices,  $N, S$ , say, where  $N$  has outgoing arcs in cyclic order  $y_0, y_2, y_4, \dots, y_{n-2}$  and  $S$  has outgoing arcs in cyclic order  $y_1, y_3, y_5, \dots, y_{n-1}$ . The 2-cells incident to  $N$  are therefore, in cyclic order,  $F_0^-, F_2^-, F_4^-, \dots, F_{n-2}^-$ , where the boundary of  $F_i^-$  reads, anticlockwise, the  $i$ 'th relator of  $\mathcal{G}^{k/l}(n, f)$ . The arc labelled  $y_2$ , with initial vertex  $N$ , is the first of a path  $y_2 - y_4 - \dots - y_{2+(l-1)f}$ . Let  $v$  denote the terminal vertex of the last arc in this path. Then  $v$  has two incoming arcs labelled  $y_{(l-1)f+1}, y_{(l-1)f+2}$ , and an outgoing arc labelled  $y_{lf+3}$ . Therefore  $v$  is a degree 4 vertex, and its remaining arc is outgoing, and labelled  $y_{lf+1}$ . The outgoing arcs  $y_{lf+1}, y_{lf+3}$  then bound the  $F_{lf+1}^-$  face. Since  $l, f$  are odd,  $lf + 1$  is even, so the face pairing contains two  $F_{lf+1}^-$  faces, a contradiction.  $\square$

In Figures 6,8,7,9 we present face-pairing polyhedra with  $n$  pairs of faces  $F_i^+, F_i^-$  ( $0 \leq i < n$ ) of opposite orientation whose boundaries spell the defining relators of the relevant presentation  $\mathcal{G}^{k/l}(n, f)$ . In the proof of Theorem B we show that the identification of faces  $F_i^+, F_i^-$  results in a 3-complex that satisfies the Seifert-Threlfall condition, and so is a manifold whose spine

is the presentation complex of  $\mathcal{G}^{k/l}(n, f)$ . (Note that, in these figures, the condition that  $f$  is even is necessary for each relator to appear as the label of a pair of oppositely oriented faces.) The diagrams have different forms depending on whether  $l > k$  or  $k > l$ . The figures should provide the enthusiastic reader with sufficient information to construct face-pairing polyhedra in the general cases. The values  $\{k, l\} = \{2, 5\}$  were selected so that  $k, l, |k - l|$  are distinct, to make it straightforward to infer paths of lengths  $k, l$ , and  $|k - l|$ . (For example, in Figure 7 the path from  $v_{2f}$  to  $u_2$  has length  $3 = k - l$ , and in Figure 9 the path from  $w_f$  to  $v_{f-1}$  has length  $3 = l - k$ .) We expect the Seifert-Threlfall condition to hold in the general case for coprime  $k, l$ ; however, the corresponding analysis to check this would be highly technical. Since our goal is to exhibit new cyclic presentations that arise as spines of closed 3-manifolds, whose Whitehead graphs are of type (II.7), rather than to be exhaustive, we have not sought to do this.

*Proof of Theorem B.* We may assume  $n$  is even and  $fk \equiv 0$ , for otherwise the Whitehead graph of  $\mathcal{G}^{k/l}(n, f)$  is not planar [27] (noting that  $fk \not\equiv 2 \pmod n$ , by hypothesis). Moreover, by Lemma 4.3 we may assume  $f$  is even.

*Case 1:  $(k, l) = (k, 1)$ .* Consider the face pairing polyhedron depicted in Figure 6 (where the arc labels can be deduced from the position of the unique source in the boundary of each face) with face pairing given by identifying the faces  $F_i^+, F_i^-$  ( $0 \leq i < n$ ). We shall show that the identification of faces results in a 3-complex  $M$  whose 2-skeleton is the presentation complex of  $\mathcal{G}^{k/l}(n, f)$ , so  $M$  has one 0-cell,  $n$  1-cells,  $n$  2-cells, and one 3-cell, so is a manifold by [49, Theorem I, Section 60].

All the arcs labelled  $x_0$  are contained in the following cycle:

$$\begin{aligned} [N, u_0] &\xrightarrow{F_0} [u_{1-f}, v_0] \xrightarrow{F_{-1}} [w_f^{k-2}, u_1] \xrightarrow{F_{f-1}} [w_{2f}^{k-3}, w_{2f}^{k-2}] \\ &\xrightarrow{F_{2f-1}} [w_{3f}^{k-4}, w_{3f}^{k-3}] \xrightarrow{F_{3f-1}} \dots \xrightarrow{F_{(k-3)f-1}} [w_{(k-2)f}^1, w_{(k-2)f}^2] \\ &\xrightarrow{F_{(k-2)f-1}} [v_{(k-1)f-2}, w_{(k-1)f}^1] \xrightarrow{F_{(k-1)f-1}} [v_{(k-1)f-3}, w_0^1] \xrightarrow{F_{-2}} [N, u_0]. \end{aligned}$$

Therefore, in the resulting complex  $M$  all the arcs labelled  $x_0$  are identified. Moreover, all the vertices that are a vertex (either initial or terminal) of an arc labelled  $x_0$  are contained in the (induced) cycle of the initial vertices in the arcs above, so these vertices are identified in  $M$ . By applying the shift  $\theta^{2j}$  ( $0 \leq j < n/2$ ) to the above, all the arcs labelled  $x_{2j}$  are identified and all the vertices of such arcs are identified. That is, the vertices  $N, u_{2j+(1-f)}, w_{2j+f}^{k-2}, w_{2j+2f}^{k-3}, \dots, w_{2j+(k-2)f}^1, v_{2j+(k-1)f-2}$  are identified; equivalently,  $N, u_{2j+1}, w_{2j}^1, \dots, w_{2j}^{k-2}, v_{2j}$  ( $0 \leq j < n/2$ ) are identified.

All the arcs labelled  $x_1$  are contained in the following cycle:

$$\begin{aligned} [S, u_1] &\xrightarrow{F_1} [u_{2-f}, v_1] \xrightarrow{F_0} [w_{f+1}^{k-2}, u_2] \xrightarrow{F_f} [w_{2f+1}^{k-3}, w_{2f+1}^{k-2}] \\ &\xrightarrow{F_{2f}} [w_{3f+1}^{k-4}, w_{3f+1}^{k-3}] \xrightarrow{F_{3f}} \dots \xrightarrow{F_{(k-3)f}} [w_{(k-2)f+1}^1, w_{(k-2)f+1}^2] \\ &\xrightarrow{F_{(k-2)f}} [v_{(k-1)f-1}, w_{(k-1)f+1}^1] \xrightarrow{F_{(k-1)f}} [v_{(k-1)f-2}, w_1^1] \xrightarrow{F_{-2}} [S, u_1]. \end{aligned}$$

Therefore, in  $M$  all the arcs labelled  $x_1$  are identified. Moreover, all the vertices that are a vertex (either initial or terminal) of an arc labelled  $x_1$  are contained in the (induced) cycle of the initial vertices in the arcs above, so these vertices are identified in  $M$ . By applying the shift  $\theta^{2j}$  ( $0 \leq j < n/2$ ) to the above, all the arcs labelled  $x_{2j+1}$  are identified and all the vertices of such arcs are identified. That is, the vertices  $S, u_{2j+(2-f)}, w_{2j+f+1}^{k-2}, w_{2j+2f+1}^{k-3}, \dots, w_{2j+(k-2)f+1}^1, v_{2j+(k-1)f-1}$  are identified; that is  $S, u_{2j}, w_{2j+1}^1, \dots, w_{2j+1}^{k-2}, v_{2j+1}$  ( $0 \leq j < n/2$ ) are identified. Moreover, examining the terminal vertices of the first cycle above we see that  $u_0$  and  $u_1$  are identified.




Figure 6: Face pairing polyhedron for  $\mathcal{G}^{k/l}(n, f)$

Therefore the vertices induced from the first cycle are identified with the vertices from the second cycle, so all vertices of the polyhedron are identified. Thus  $M$  has one 3-cell,  $n$  2-cells,  $n$  1-cells, and one 0-cell, and since the boundaries of the 2-cells spell the relators of  $\mathcal{G}^{1/l}(n, 0)$ , it follows that  $\mathcal{G}^{1/l}(n, 0)$  is the spine of a closed 3-manifold, as required.

*Case 2:*  $(k, l) = (5, 2)$ . Consider the face pairing polyhedron depicted in Figure 7 (where the arc labels can be deduced from the position of the unique source in the boundary of each face) with face pairing given by identifying the faces  $F_i^+, F_i^-$  ( $0 \leq i < n$ ). All the arcs labelled  $x_0$  are contained in the following cycle:

$$\begin{aligned}
[N, s_0] &\xrightarrow{F_0} [u_{1-2f}, t_0] \xrightarrow{F_{-1}} [w_{2f-1}^1, w_{2f-1}^2] \xrightarrow{F_{2f-1}^-} [r_{2f-1}, v_{4f-1}] \\
&\xrightarrow{F_{4f-2}^+} [s_{4f}, u_{4f}] \xrightarrow{F_{4f}} [t_{4f}, v_{4f}] \xrightarrow{F_{4f-1}^+} [w_{f-1}^2, u_{-f+1}] \\
&\xrightarrow{F_{f-1}^-} [v_{3f-1}, w_{3f-1}^1] \xrightarrow{F_{3f-1}^-} [u_{3f-1}, r_{3f-1}] \xrightarrow{F_{-2}^-} [N, s_0],
\end{aligned}$$

and all the arcs labelled  $x_1$  are contained in the following cycle:

$$\begin{aligned}
[S, s_1] &\xrightarrow{F_1} [u_{2-2f}, t_1] \xrightarrow{F_0} [w_{2f}^1, w_{2f}^2] \xrightarrow{F_{2f}} [r_{2f}, v_{4f}] \xrightarrow{F_{4f-1}^+} [s_{4f+1}, u_{4f+1}] \\
&\xrightarrow{F_{4f+1}^+} [t_{4f+1}, v_{4f+1}] \xrightarrow{F_{4f}^+} [w_f^2, u_{-f+2}] \xrightarrow{F_f} [v_{3f}, w_{3f}^1] \xrightarrow{F_{3f}^+} [u_{3f}, r_{3f}] \xrightarrow{F_{-1}^-} [S, s_1].
\end{aligned}$$

The proof then proceeds as in Case 1.

*Case 3:*  $(k, l) = (1, l)$ . Consider the face pairing polyhedron depicted in Figure 8. All the arcs labelled  $x_0$  are contained in the following cycle:




Figure 7: Face pairing polyhedron for  $\mathcal{G}^{5/2}(n, f)$

$$\begin{aligned}
[N, u_0^1] &\xrightarrow{F_0} [w_0, t_0^1] \xrightarrow{F_{-2}} [u_0^1, u_0^2] \xrightarrow{F_0} [t_0^1, t_0^2] \xrightarrow{F_{-2}} [u_0^2, u_0^3] \\
&\xrightarrow{F_0} [t_0^2, t_0^3] \longrightarrow \cdots \xrightarrow{F_{-2}} [u_0^{l-2}, u_0^{l-1}] \xrightarrow{F_0} [t_0^{l-2}, v_{-1}] \\
&\xrightarrow{F_{-2}} [u_0^{l-1}, w_{-1}] \xrightarrow{F_0} [v_{-1}, v_0] \xrightarrow{F_{-1}} [w_{-2}, w_0] \xrightarrow{F_{-2}} [N, u_0^1],
\end{aligned}$$

and all the arcs labelled  $x_1$  are contained in the following cycle:

$$\begin{aligned}
[S, u_1^1] &\xrightarrow{F_1} [w_1, t_1^1] \xrightarrow{F_{-1}} [u_1^1, u_1^2] \xrightarrow{F_1} [t_1^1, t_1^2] \xrightarrow{F_{-1}} [u_1^2, u_1^3] \\
&\xrightarrow{F_1} [t_1^2, t_1^3] \longrightarrow \cdots \xrightarrow{F_{-1}} [u_1^{l-2}, u_1^{l-1}] \xrightarrow{F_1} [t_1^{l-2}, v_0] \\
&\xrightarrow{F_{-1}} [u_1^{l-1}, w_0] \xrightarrow{F_1} [v_0, v_1] \xrightarrow{F_0} [w_{-1}, w_1] \xrightarrow{F_{-1}} [S, u_1^1].
\end{aligned}$$

The proof then proceeds as in Case 1.

*Case 4:*  $(k, l) = (2, 5)$ . Consider the face pairing polyhedron depicted in Figure 9. All the arcs labelled  $x_0$  are contained in the following cycle:

$$\begin{aligned}
[N, u_0^1] &\xrightarrow{F_0} [w_0, t_0^1] \xrightarrow{F_{-2}} [u_0^2, u_0^3] \xrightarrow{F_0} [t_0^2, v_{-1}] \xrightarrow{F_{-2}} [u_0^4, w_{f-1}] \\
&\xrightarrow{F_0} [s_{-1}, v_0] \xrightarrow{F_{-1}} [r_{f-2}, w_f] \xrightarrow{F_{f-2}} [u_f^1, u_f^2] \xrightarrow{F_f} [t_f^1, t_f^2] \\
&\xrightarrow{F_{f-2}} [u_f^3, u_f^4] \xrightarrow{F_f} [v_{f-1}, s_{f-1}] \xrightarrow{F_{f-1}} [w_{2f-2}, r_{2f-2}] \xrightarrow{F_{2f-2}} [N, u_0^1],
\end{aligned}$$

and all the arcs labelled  $x_1$  are contained in the following cycle:

$$\begin{aligned}
[S, u_1^1] &\xrightarrow{F_1} [w_1, t_1^1] \xrightarrow{F_{-1}} [u_1^2, u_1^3] \xrightarrow{F_1} [t_1^2, v_0] \xrightarrow{F_{-1}} [u_1^4, w_f] \\
&\xrightarrow{F_1} [s_0, v_1] \xrightarrow{F_0} [r_{f-1}, w_{f+1}] \xrightarrow{F_{f-1}} [u_{f+1}^1, u_{f+1}^2] \xrightarrow{F_{f+1}} [t_{f+1}^1, t_{f+1}^2] \\
&\xrightarrow{F_{f-1}} [u_{f+1}^3, u_{f+1}^4] \xrightarrow{F_{f+1}} [v_f, s_f] \xrightarrow{F_f} [w_{2f-1}, r_{2f-1}] \xrightarrow{F_{2f-1}} [S, u_1^1].
\end{aligned}$$

The proof then proceeds as in Case 1. □

We now consider the structures of the manifolds  $M^{k/l}(n, f)$  of Theorem B.

**Theorem 4.4.** *Let  $n \geq 2$ ,  $k, l \geq 1$ ,  $0 \leq f < n$ , and  $fk \equiv 0 \pmod n$ , where  $n, f$  are even, and suppose  $(k, l) \in \{(k, 1), (1, l), (5, 2), (2, 5)\}$ . Then  $M^{k/l}(n, f)$  is hyperbolic if and only if  $M^{k/l}(n, 0)$  is hyperbolic.*

*Proof.* By [17, Corollary 3.3] the shift  $\theta_{G^{k/l}(n,0)}$  has order  $n$ , and hence the shift  $\theta_{G^{k/l}(n,f)}$  also has order  $n$ . Suppose  $M^{k/l}(n, 0)$  is hyperbolic. Then since by Theorem B,  $M^{k/l}(n, 0)$  is a closed, connected, orientable 3-manifold,  $G^{k/l}(n, 0)$  is a subgroup of  $\text{Isom}^+(H^3) \cong PSL(2, \mathbb{C})$ . As in the proof of [40, Theorem 3.1] (see also [12, Theorem 3.1], [14, Theorem 3.1], [3, Theorem 3.1]) Mostow rigidity implies that  $E^{k/l}(n)$  is a subgroup of  $PSL(2, \mathbb{C})$ . Hence  $G^{k/l}(n, f)$  is a subgroup of  $PSL(2, \mathbb{C})$ , and so  $M^{k/l}(n, f)$  is hyperbolic. Repeating the argument with the roles of  $G^{k/l}(n, 0)$  and  $G^{k/l}(n, f)$  interchanged and the roles of  $M^{k/l}(n, 0)$  and  $M^{k/l}(n, f)$  interchanged proves the converse. □

*Remark 4.5.* The argument of the proof of Theorem 4.4 holds in the more general setting of [5, Theorem 2.3]. Namely, let  $E = \langle x, t \mid t^n, W(x, t) \rangle$  ( $n \geq 2$ ), and suppose there are retractions  $\nu^0, \nu^f : E \rightarrow \langle t \mid t^n \rangle$  ( $0 \leq f < n$ ) given by  $\nu^0(t) = \nu^f(t) = t$  and  $\nu^0(x) = t^0$ ,  $\nu^f(x) = t^f$ . Then  $K_0 = \ker(\nu_0)$ ,  $K_f = \ker(\nu_f)$  have cyclic presentations  $\mathcal{G}_n(w_0), \mathcal{G}_n(w_f)$ , respectively (for some words  $w_0, w_f$  in generators  $x_0, \dots, x_{n-1}$ ). Suppose that the shift automorphism has order  $n$  for




Figure 8: Face pairing polyhedron for  $\mathcal{G}^{1/l}(n, 0) = \mathcal{F}^{1/l}(n)$




Figure 9: Face pairing polyhedron for  $\mathcal{G}^{2/5}(n, f)$

either (and hence both) of these groups, and that  $K_0, K_f$  are fundamental groups of closed, connected, orientable 3-manifolds  $M_0, M_f$ , respectively. Then  $M_0$  is hyperbolic if and only if  $M_f$  is hyperbolic. (Theorem 4.4 corresponds to the case  $W(x, t) = x^l t x^k t x^{-l} t^{-2}$ , as in (1), where  $(k, l) \in \{(k, 1), (1, l), (5, 2), (2, 5)\}$ .)

The significance of Theorem 4.4 is that the manifolds  $M^{k/l}(n, 0)$  are known to be hyperbolic in many cases, as we now describe. By [56, Corollary 2.1], if  $n \geq 6$  is even then  $M^{k/l}(n, 0)$  is hyperbolic for all but finitely many pairs of coprime integers  $k, l$ , and it is conjectured ([56, page 658]) that  $n = 6$  and  $k = l = 1$  is the only non-hyperbolic case (which is an affine Riemannian manifold by [22, Proposition 6]). Moreover,  $M^{k/l}(n, 0)$  is hyperbolic in each of the following cases:  $k \geq 2, l = 1, n \geq 6$  and even [41, Theorem 3];  $k = 1, l \geq 2, n \geq 6$  and even [56, Corollary 3.5];  $k = l = 1, n \geq 8$  and even [22, Theorem C]. If  $n = 4$  and  $k = 1$  then  $M^{k/l}(n, 0) = M^{1/l}(4, 0)$  is the lens space  $L(4l^2 + 1, 2l)$  and  $G^{1/l}(4, 0) \cong \mathbb{Z}_{4l^2+1}$  by [56, Corollary 3.4]. For  $k \geq 2$  the manifolds  $M^{k/1}(4, 0)$  are not hyperbolic and are described in [41, pages 170–171]. In the next theorem we consider the manifolds  $M^{k/l}(4, f)$  where  $k \geq 2$ .

**Theorem 4.6.** *Let  $k \geq 2, l \geq 1$ , where  $\gcd(k, l) = 1$  and suppose  $fk \equiv 0 \pmod{4}$ . Then  $G^{k/l}(4, f)$  contains a  $\mathbb{Z} \oplus \mathbb{Z}$  subgroup and hence  $M^{k/l}(4, f)$  is not hyperbolic.*

*Proof.* We show that the subgroup  $A$  of  $E = E^{k/l}(4)$  generated by  $x^k, tx^k t^{-1}$  is free abelian of rank 2. Then, since  $\nu_f(x^k) = 1$  and  $\nu_f(tx^k t^{-1}) = 1$ ,  $A = \mathbb{Z} \oplus \mathbb{Z}$  is a subgroup of  $\ker(\nu_f) = G^{k/l}(4, f)$ , and hence  $M^{k/l}(4, f)$  is not hyperbolic by [6, Theorem 3.3].

Let  $G = \ker(\nu_0)$ , so  $G = F^{k/l}(4)$  and is generated by  $x_i = t^i x t^{-i}$ , and let  $G_1 = \langle x_0, x_2 \mid x_0^k x_2^k \rangle$ ,  $G_2 = \langle x_1, x_3 \mid x_1^k x_3^k \rangle$ ,  $H = \langle a, b \mid ab = ba \rangle \cong \mathbb{Z}^2$ . Let  $\phi_1 : H \rightarrow G_1$ ,  $\phi_2 : H \rightarrow G_2$ , be given by  $\phi_1(a) = x_0^k$ ,  $\phi_1(b) = x_0^{-l} x_2^l$ ,  $\phi_2(a) = x_3^{-l} x_1^l$ ,  $\phi_2(b) = x_1^k$ . We shall show that  $\phi_1, \phi_2$  are injections. This implies that  $G$  can be expressed as an amalgamated free product  $G = G_1 *_H G_2$ , and  $\phi_1(a) = x_0^k$ ,  $\phi_1(b) = x_0^{-l} x_2^l$  generate a free abelian subgroup of  $G$  (compare [41, page 171], which deals with the case  $l = 1$ ). That is,  $x_0^k$  and  $x_1^k$  generate a free abelian subgroup of  $G$  and so  $x^k$  and  $tx^k t^{-1}$  generate a free abelian subgroup of  $E$ , as required.

We show that  $\phi_1$  is an injection, the argument for  $\phi_2$  being similar. Let  $\alpha : G_1 \rightarrow \mathbb{Z}_k$  be given by  $\alpha(x_0) = 1, \alpha(x_2) = 1 \in \mathbb{Z}_k$ . Then  $K = \ker(\alpha)$  is generated by  $z = x_0^k$ ,  $g_i = x_0^i x_2 x_0^{-(i+1)}$  ( $0 \leq i \leq k-2$ ),  $g_{k-1} = x_0^{k-1} x_2$ , and has a presentation

$$\langle z, g_0, \dots, g_{k-1} \mid z g_0 g_1 \dots g_{k-2} g_{k-1}, z g_1 g_2 \dots g_{k-1} g_0, \dots, z g_{k-1} g_0 \dots g_{k-3} g_{k-2} \rangle.$$

Now let  $\theta : K \rightarrow \mathbb{Z}^k$  be an epimorphism given by  $\theta(g_1) = (1, 0, \dots, 0)$ ,  $\theta(g_2) = (0, 1, \dots, 0), \dots$ ,  $\theta(g_{k-1}) = (0, 0, \dots, 1)$ , and  $\theta(z) = (-1, -1, \dots, -1)$ . Let  $l = qk + r$  where  $q \geq 0$ ,  $0 \leq r < k$  and note that  $r \neq 0$ , since  $\gcd(k, l) = 1$ . Then  $x_0^{-l} x_2^l = (x_0^k)^{-q} x_0^{-r} (x_2^k)^q x_2^r = (x_0^k)^{-q} x_0^{-r} (x_0^{-k})^q x_2^r = z^{-2q-1} g_{k-r} g_{k-(r-1)} \dots g_{k-1}$  and  $x_2^k = z$ . Therefore

$$\theta(x_0^{-l} x_2^l) = (2q+1)(1, 1, \dots, 1) + (0, \dots, 0, 1, \dots, 1) = (2q+1, \dots, 2q+1, 2q+2, \dots, 2q+2)$$

and  $\theta(x_2^k) = (-1, -1, \dots, -1)$  so  $\theta(x_0^{-l} x_2^l), \theta(x_2^k)$  generate a free abelian subgroup of rank 2 in  $\mathbb{Z}^k$ . Moreover, in  $G_1$ ,  $x_0^{-l} x_2^l$  and  $x_2^k$  commute and so generate a free abelian subgroup of  $G_1$ , and hence  $\phi_1$  is injective.  $\square$

## Acknowledgements

I thank Jim Howie for insightful comments on a draft of this article and Alessia Cattabriga for interesting discussions relating to this work, and for bringing pertinent results from [43] and [47] to my attention.

## References

- [1] Nurullah Ankaralioglu and Huseyin Aydin. Some Dunwoody parameters and cyclic presentations. *Gen. Math.*, 16(2):85–93, 2008.
- [2] H. Aydin, I. Gultekyn, and M. Mulazzani. Torus knots and Dunwoody manifolds. *Sibirsk. Mat. Zh.*, 45(1):3–10, 2004.
- [3] V.G. Bardakov and A.Yu. Vesnin. A generalization of Fibonacci groups. *Alg. and Logic*, 42:131–160, 2003.
- [4] John Berge. Heegaard. <https://github.com/3-manifolds/heegaard>.
- [5] William A. Bogley. On shift dynamics for cyclically presented groups. *J. Algebra*, 418:154–173, 2014.
- [6] P.J. Callahan and A.W. Reid. Hyperbolic structures on knot complements. *Chaos, Solitons & Fractals*, 9(4):705–738, 1998. Knot Theory and Its Applications: Expository Articles on Current Research.
- [7] Alessia Cattabriga and Michele Mulazzani. Strongly-cyclic branched coverings of  $(1, 1)$ -knots and cyclic presentations of groups. *Math. Proc. Cambridge Philos. Soc.*, 135(1):137–146, 2003.
- [8] Alessia Cattabriga and Michele Mulazzani. All strongly-cyclic branched coverings of  $(1, 1)$ -knots are Dunwoody manifolds. *J. London Math. Soc. (2)*, 70(2):512–528, 2004.
- [9] Alessia Cattabriga and Michele Mulazzani. Representations of  $(1, 1)$ -knots. *Fund. Math.*, 188:45–57, 2005.
- [10] Alessia Cattabriga, Michele Mulazzani, and Andrei Vesnin. Complexity, Heegaard diagrams and generalized Dunwoody manifolds. *J. Korean Math. Soc.*, 47(3):585–599, 2010.
- [11] A. Cavicchioli, F. Hegenbarth, and A.C. Kim. A geometric study of Sieradski groups. *Algebra Colloq.*, 5(2):203–217, 1998.
- [12] A. Cavicchioli, D. Repovš, and F. Spaggiari. Topological properties of cyclically presented groups. *J. Knot Theory Ramifications*, 12(2):243–268, 2003.
- [13] Alberto Cavicchioli, Friedrich Hegenbarth, and Ann Chi Kim. On cyclic branched coverings of torus knots. *J. Geom.*, 64(1-2):55–66, 1999.
- [14] Alberto Cavicchioli, Dušan Repovš, and Fulvia Spaggiari. Families of group presentations related to topology. *J. Algebra*, 286(1):41–56, 2005.
- [15] Alberto Cavicchioli, Beatrice Ruini, and Fulvia Spaggiari. On a conjecture of M. J. Dunwoody. *Algebra Colloq.*, 8(2):169–218, 2001.
- [16] Alberto Cavicchioli and Fulvia Spaggiari. The classification of 3-manifolds with spines related to Fibonacci groups. Algebraic topology, Proc. Conf., S. Feliu de Guíxols/Spain 1990, Lect. Notes Math. 1509, 50-78, 1992.
- [17] Ihechukwu Chinyere and Gerald Williams. Fractional Fibonacci groups with an odd number of generators. *Topology Appl.*, 312:Paper No. 108083, 16, 2022.

- [18] M.J. Dunwoody. Cyclic presentations and 3-manifolds. In *Groups - Korea '94. Proceedings of the international conference, Pusan, Korea, August 18-25, 1994*. Berlin: Walter de Gruyter, 1995.
- [19] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.13.1*, 2024.
- [20] L. Grasselli and M. Mulazzani. Seifert manifolds and  $(1,1)$ -knots. *Sibirsk. Mat. Zh.*, 50(1):28–39, 2009.
- [21] Luigi Grasselli and Michele Mulazzani. Genus one 1-bridge knots and Dunwoody manifolds. *Forum Math.*, 13(3):379–397, 2001.
- [22] H. Helling, A.C. Kim, and J.L. Mennicke. A geometric study of Fibonacci groups. *J. Lie Theory*, 8(1):1–23, 1998.
- [23] Hugh Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On a remarkable polyhedron geometrizing the figure eight knot cone manifolds. *J. Math. Sci., Tokyo*, 2(3):501–561, 1995.
- [24] Hugh M. Hilden, María Teresa Lozano, and Jose María Montesinos-Amilibia. The arithmeticity of the figure eight knot orbifolds. In *Topology '90. Contributions from a research semester in low dimensional topology, held at Ohio State University, Columbus, OH (USA), from Feb. through June 1990*, pages 169–183. Berlin: Walter de Gruyter, 1992.
- [25] Cynthia Hog-Angeloni and Wolfgang Metzler. Geometric aspects of two-dimensional complexes. In Hog-Angeloni et al., editor, *Two-dimensional homotopy and combinatorial group theory*, number 197 in Lond. Math. Soc. Lect. Note Ser., pages 1–50. Cambridge: Cambridge University Press, 1993.
- [26] James Howie and Gerald Williams. Fibonacci type presentations and 3-manifolds. *Topology Appl.*, 215:24–34, 2017.
- [27] James Howie and Gerald Williams. Planar Whitehead graphs with cyclic symmetry arising from the study of Dunwoody manifolds. *Discrete Math.*, 343(12):112096, 18, 2020.
- [28] Kwang-Woo Jeong. On the cyclic branched coverings of the 2-bridge knot  $b(17,4)$ . *Algebra Colloq.*, 13(1):173–180, 2006.
- [29] Kwang-Woo Jeong and Moon-Ok Wang. Notes on more Fibonacci groups. *Algebra Colloq.*, 15(4):699–706, 2008.
- [30] D. L. Johnson and E. F. Robertson. Finite groups of deficiency zero. In *Homological group theory (Proc. Sympos., Durham, 1977)*, volume 36 of *London Math. Soc. Lecture Note Ser.*, pages 275–289. Cambridge Univ. Press, Cambridge-New York, 1979.
- [31] D. L. Johnson, J. W. Wamsley, and D. Wright. The Fibonacci groups. *Proc. London Math. Soc. (3)*, 29:577–592, 1974.
- [32] D.L. Johnson. *Topics in the Theory of Group Presentations*, volume 42 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1980.
- [33] Goansu Kim, Yangkok Kim, and Andrei Vesnin. The knot  $5_2$  and cyclically presented groups. *J. Korean Math. Soc.*, 35(4):961–980, 1998.

- [34] Soo Hwan Kim and Yangkok Kim. Torus knots and 3-manifolds. *J. Knot Theory Ramifications*, 13(8):1103–1119, 2004.
- [35] Soo Hwan Kim and Yangkok Kim. On the 2-bridge knots of Dunwoody  $(1, 1)$ -knots. *Bull. Korean Math. Soc.*, 48(1):197–211, 2011.
- [36] Soo Hwan Kim and Yangkok Kim. On the polynomial of the Dunwoody  $(1, 1)$ -knots. *Kyungpook Math. J.*, 52(2):223–243, 2012.
- [37] Soo Hwan Kim and Yangkok Kim. The dual and mirror images of the Dunwoody 3-manifolds. *Int. J. Math. Math. Sci.*, 7 pages, 2013. Art. ID 103209.
- [38] Soo Hwan Kim and Yangkok Kim. On the generalized Dunwoody 3-manifolds. *Osaka J. Math.*, 50(2):457–476, 2013.
- [39] Tatyana Kozlovskaya. Heegaard splittings of branched cyclic coverings of connected sums of lens spaces. *Bull. Korean Math. Soc.*, 54(5):1851–1857, 2017.
- [40] C. Maclachlan. Generalisations of Fibonacci numbers, groups and manifolds. Duncan, A.J. et al., editor, Combinatorial and geometric group theory. Proceedings of a workshop held at Heriot-Watt University, Edinburgh, 1993. Cambridge University Press. Lond. Math. Soc. Lect. Note Ser. 204, 233-238, 1995.
- [41] C. Maclachlan and A. W. Reid. Generalised Fibonacci manifolds. *Transform. Groups*, 2(2):165–182, 1997.
- [42] Kirk McDermott. On the topology of the groups of type  $Z$ . *Topology and its Applications*, 307:107767, 2022.
- [43] Jerome Minkus. The branched cyclic coverings of 2 bridge knots and links. *Mem. Amer. Math. Soc.*, 35(255):iv+68, 1982.
- [44] Michele Mulazzani.  $(1, 1)$ -knots and Dunwoody manifolds. In *Recent advances in geometry and topology*, pages 273–277. Cluj Univ. Press, Cluj-Napoca, 2004.
- [45] L. Neuwirth. An algorithm for the construction of 3-manifolds from 2-complexes. *Proc. Camb. Philos. Soc.*, 64:603–613, 1968.
- [46] Vanni Noferini and Gerald Williams. Smith forms of matrices in companion rings, with group theoretic and topological applications, 2024. arXiv: 2408.08662.
- [47] Beatrice Ruini and Fulvia Spaggiari. On the structure of Takahashi manifolds. *Tsukuba J. Math.*, 22(3):723–739, 1998.
- [48] Beatrice Ruini, Fulvia Spaggiari, and Andrei Vesnin. On spines of Seifert fibered manifolds. *Aequationes Math.*, 65(1-2):40–60, 2003.
- [49] Herbert Seifert and William Threlfall. *A textbook of topology*, volume 89 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, German edition, 1980. With a preface by Joan S. Birman, With “Topology of 3-dimensional fibered spaces” by Seifert, Translated from the German by Wolfgang Heil.
- [50] Allan J. Sieradski. Combinatorial squashings, 3-manifolds, and the third homology of groups. *Invent. Math.*, 84:121–139, 1986.

- [51] F. Spaggiari and A. I. Telloni. Isometry groups of some Dunwoody manifolds. *Algebra Colloq.*, 23(1):117–128, 2016.
- [52] Fulvia Spaggiari. A geometric study of generalized Neuwirth groups. *Forum Math.*, 18(5):803–827, 2006.
- [53] Agnese Ilaria Telloni. Combinatorics of a class of groups with cyclic presentation. *Discrete Math.*, 310(22):3072–3079, 2010.
- [54] Abdullahi Umar. Some remarks about Fibonacci groups and semigroups. *Communications in Algebra*, 25(12):3973–3977, 1997.
- [55] A. Yu. Vesnin and A. Ch. Kim. A topological study of the Fractional Fibonacci groups. *Universität Bielefeld, Preprint 96-049*, 1996.
- [56] A. Yu. Vesnin and A. Ch. Kim. Fractional Fibonacci groups and manifolds. *Sibirsk. Mat. Zh.*, 39(4):765–775, i, 1998.
- [57] A. Yu. Vesnin and T. A. Kozlovskaya. Brieskorn manifolds, generalized Sieradski groups and coverings of lens spaces. *Tr. Inst. Mat. Mekh.*, 23(4):85–97, 2017.

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF ESSEX, WIVENHOE PARK, COLCHESTER, ESSEX CO4 3SQ, UK.

*E-mail address*, `Gerald.Williams@essex.ac.uk`