Polynomial bounds for monochromatic tight cycle partition in *r*-edge-coloured $K_n^{(k)}$

Debmalya Bandyopadhyay * Allan Lo[†]

September 2, 2024

Abstract

Let $K_n^{(k)}$ be the complete k-graph on n vertices. A k-uniform tight cycle is a k-graph with its vertices cyclically ordered so that every k consecutive vertices form an edge and any two consecutive edges share exactly k-1 vertices. A result of Bustamante, Corsten, Frankl, Pokrovskiy and Skokan shows that all r-edge coloured $K_n^{(k)}$ can be partitioned into $c_{r,k}$ vertex disjoint monochromatic tight cycles. However, the constant $c_{r,k}$ is of tower-type. In this work, we show that $c_{r,k}$ is a polynomial in r.

1 Introduction

An *r*-edge-colouring of a graph or a *k*-uniform hypergraph is a colouring of its edges with *r* colours. The set of colours is usually identified with the set $\{1, 2, ..., r\}$. A monochromatic subgraph of an edge-coloured graph is a subgraph where all the edges are assigned the same colour. On the other hand, a rainbow subgraph is a subgraph all of whose edges have a different colour.

Lehel conjectured that for any 2-edge-colouring of a complete graph, there exist two vertex-disjoint monochromatic cycles (one of each colour) covering all vertices. Isolated vertices and single-edges are considered to be degenerate cycles. For large n, this conjecture was proved by Luczak, Rödl and Szemerédi [18] using Szemerédi's Regularity Lemma. The bound on n was improved later by Allen [1]. In 2010, Bessy and Thomassé [3] finally resolved this conjecture for all $n \in \mathbb{N}$.

When $r \geq 3$, Erdős, Gyárfás and Pyber [7] proved that any *r*-edge-coloured complete graph can be partitioned into $O(r^2 \log r)$ monochromatic cycles and conjectured that *r* monochromatic cycles would be enough. This was one of the first instances of using the absorbing method. Gyárfás, Ruszinkó, Sárközy and Szemerédi [11] improved their result and proved that $O(r \log r)$ monochromatic cycles suffice. However, Pokrovskiy [21] found a counter-example disproving the conjecture. A weaker conjecture was proposed stating that any *r*-edge-coloured K_n contains *r* vertex-disjoint monochromatic cycles covering all but c(r) of the vertices, where c(r) is a constant depending only on *r*. Pokrovskiy [20] also proved that $c(3) \leq 43000$. Korándi, Lang, Letzter and Pokrovskiy [14] determined a tight minimum degree threshold for monochromatic cycle partition of edge-coloured graphs, namely they showed that there exists a constant c > 0 such that any *r*-edge-coloured graph *G* on *n* vertices with $\delta(G) \geq n/2 + cr \log n$ has a partition into $O(r^2)$ monochromatic cycles.

^{*}School of Mathematics, University of Birmingham, *E-mail*: dxb209@student.bham.ac.uk

[†]School of Mathematics, University of Birmingham, *E-mail*: s.a.lo@bham.ac.uk. The research leading to these results was supported by EPSRC, grant no. EP/V002279/1 and EP/V048287/1. There is no additional data beyond that contained within the main manuscript.

A local r-colouring of a graph is an edge-colouring such that every vertex is incident with at most r edges of distinct colours. Sárközy [23] showed that any large locally r-coloured K_n can be partitioned into $O(r \log r)$ monochromatic cycles.

A k-uniform hypergraph (or k-graph) is a pair H = (V(H), E(H)) where $E(H) \subseteq \binom{V(H)}{k}$. ¹ Let $K_n^{(k)}$ denote the complete graph on n vertices, where all $\binom{n}{k}$ edges are present. For positive integers $1 \leq \ell < k \leq n$, a k-uniform ℓ -cycle is a k-graph with its vertices cyclically ordered so that every edge contains k consecutive vertices and any two consecutive edges share exactly ℓ vertices. Note that 1-cycles are called *loose cycles* and (k-1)-cycles are called *tight cycles*.

Lehel's problem has been generalised for hypergraphs and studied for both tight and loose cycles. As in the case of graphs, any set of at most k vertices is considered as a degenerate cycle. Gyárfás and Sárközy [12] showed that for loose cycles, every r-edge-coloured $K_n^{(k)}$ can be partitioned into C(k,r) vertex-disjoint monochromatic loose cycles. For sufficiently large n, Sárközy [22] proved that $50kr\log(kr)$ loose cycles suffice. For an overview of results related to monochromatic partitions of (hyper)graphs we refer the reader to the surveys [8] and [9].

In this paper, we focus on monochromatic tight cycle partition. For k = 3, Bustamante, Hàn and Stein [5] proved that any 2-edge-coloured $K_n^{(3)}$ contains two vertex-disjoint monochromatic cycles of distinct colours covering all but at most o(n) vertices. Lo and Pfenninger [16] proved the corresponding result when k = 4. Recently, Pfenninger [19] generalised the result to all k improving a previous result [15].

For any $r, k \ge 3$, Bustamante, Corsten, Frankl, Pokrovskiy and Skokan [4] proved that any r-edgecoloured $K_n^{(k)}$ can be partitioned into at most C'(k,r) monochromatic tight cycles. They achieved this bound by using the regularity method for hypergraphs, as a result of which the constant C'(k,r) is a tower bound.

In this paper we show that C'(k, r) is a polynomial of r (for fixed k).

Theorem 1.1. For all $r \in \mathbb{N}$ and $k \geq 3$, there exists an integer $n_0 = n_0(r, k)$ such that for all r-edgecoloured $K_n^{(k)}$ with $n \geq n_0$, there exists a monochromatic tight cycle partition of $V\left(K_n^{(k)}\right)$ into at most $(2r)^{2^{k+4}} + 2^{k+8}r\log(2r)$ tight cycles.

We make no attempt to improve the coefficients.

To prove Theorem 1.1, we use the absorption method motivated by the works of Erdős, Gyárfás and Pyber [7] and Gyárfás, Ruszinkó, Sárközy and Szemerédi [11], as well as the connected matching method that is often credited to Luczak [17]. We first reserve a special vertex subset. Then we use a result by Allen, Böttcher, Cooley and Mycroft [2] (Theorem 2.1) to greedily remove monochromatic tight cycles from the rest of the hypergraph, until there is a small leftover set of vertices. We then use properties of the reserved structures to 'absorb' the leftover set with a few monochromatic tight cycles.

We now outline the layout of the paper. We prove Theorem 1.1 through a series of reductions. In Section 2, we reduce Theorem 1.1 to the absorbing lemma, see Lemma 2.3. Roughly speaking, Lemma 2.3 finds a vertex set R such that, given any small vertex subset $B, R \cup B$ can be partitioned into few monochromatic tight cycles. The rest of this paper is focused on proving Lemma 2.3. In Sections 3 and 4, we introduce some basic notations and the hypergraph regularity lemma. In Section 5, we reduce the size of the small vertex subset B needed in Lemma 2.3. In Section 6, we translate the problem into finding rainbow cycle partitions in edge-coloured multigraphs, see Lemma 6.2. We prove this lemma in Sections 7 and 8. In Section 9, we end the paper with some concluding remarks and further directions. There will be further motivation and discussion in each section.

¹For a set V and $k \in \mathbb{N}$, $\binom{V}{k}$ denotes the set of all subsets of V of size k.

2 Proof of Theorem 1.1

We typically assume n to be a large integer. Let $[n] = \{1, \ldots, n\}$ and for integers $a \leq b$, let $[a, b] = \{a, a+1, \ldots, b\}$. We will use hierarchies in our statements. The phrase " $a \ll b$ " means "for every b > 0, there exists $a_0 > 0$, such that for all $0 < a \leq a_0$ the following statements hold". We implicitly assume all constants in such hierarchies are positive and if 1/m appears we assume m is an integer. For the rest of this paper, r and k will denote the number of colours and uniformity of the hypergraph, respectively.

We first show that one can cover most vertices of an *r*-edge-coloured $K_n^{(k)}$ with few vertex-disjoint monochromatic tight cycles. We need the following theorem on the Turán density of tight cycles.

Theorem 2.1. (Allen, Böttcher, Cooley, Mycroft [2]) Let $1/n \ll \delta, 1/k \leq 1/3$ and $\alpha \in [0, 1]$. Let G be a k-graph on n vertices with $e(G) \geq (\alpha + \delta) {n \choose k}$. Then G contains a tight cycle of length ℓ for every $\ell \leq \alpha n$ that is divisible by k.

Proposition 2.2. Let $1/n \ll \varepsilon, 1/k, 1/r$. Let $K_n^{(k)}$ be r-edge-coloured. Then all but at most εn vertices of $K_n^{(k)}$ can be covered by $2r \log(1/\varepsilon)$ vertex-disjoint monochromatic tight cycles.

Proof. Since $K_n^{(k)}$ is r-edge-coloured, there exists a monochromatic subgraph G of $K_n^{(k)}$ such that $e(G) \ge {n \choose k}/r$ edges. Theorem 2.1 with 1/3r, 2/3r playing the roles of δ, α , respectively, implies that there is a monochromatic tight cycle of length at least $2n/3r - k \ge n/2r$. Remove this cycle and repeat the argument. After removing *i* tight cycles, there are at most $(1 - \frac{1}{2r})^i n \le ne^{-i/2r}$ many vertices left. By setting $i = 2r \log(1/\varepsilon)$, we have at most εn vertices uncovered.

We need the following absorbing lemma which will be proved later.

Lemma 2.3. Let $k \ge 3$ and $1/N \ll 1/r, 1/k$. Let H be an r-edge-coloured $K_N^{(k)}$. Then there exists a vertex set $R \subseteq V(H)$ such that for any $B \subseteq V(H) \setminus R$ with $|B| \le (2r)^{-2^{k+7}} k^{-1}N$, $H[R \cup B]$ can be partitioned into at most $2^{19}r(2r)^{5 \cdot 2^{k+1}} + 3$ monochromatic tight cycles.

We now prove Theorem 1.1 assuming Lemma 2.3.

Proof of Theorem 1.1. Let $\varepsilon^* = (2r)^{-2^{k+7}}k^{-1}$ and $m = 2^{19}r(2r)^{5\cdot 2^{k+1}} + 3$. Choose constants t_0, ε_k, ψ such that $1/n \ll 1/t_0 \ll \varepsilon_k \ll \psi \ll 1/r, 1/k$. By Lemma 2.3 with n playing the role of N, there exists a vertex set $R \subseteq V(H)$ such that, for any $B \subseteq V(H) \setminus R$ with $|B| \leq \varepsilon^* n$, $H[R \cup B]$ can be partitioned into at most m monochromatic tight cycles. Let $H' = H \setminus R$ and n' = |V(H')|. By Proposition 2.2 with ε^*, n' playing the roles of ε, n , respectively, all but at most $\varepsilon^* n' \leq \varepsilon^* n$ vertices of H' can be covered by at most

$$2r\log(1/\varepsilon^*) = 2^{k+8}r\log(2r) + 2r\log(k)$$

vertex-disjoint monochromatic tight cycles. Let B be the set of uncovered vertices, so $|B| \leq \varepsilon^* n$. Thus $H[R \cup B]$ can be partitioned into at most m monochromatic tight cycles. This covers up H with at most

$$2^{k+8}r\log(2r) + 2r\log(k) + m \le 2^{20+5\cdot 2^{k+1}}r^{5\cdot 2^{k+1}+1} + 2^{k+8}r\log(2r) \le (2r)^{2^{k+4}} + 2^{k+8}r\log(2r)$$

vertex-disjoint monochromatic tight cycles. This concludes the proof of the theorem.

3 Notation

We omit floors and ceilings if they do not affect the calculations. For two sets A and B, $A\Delta B$ denotes their set difference. We often write $v_1v_2\ldots v_k$ for $\{v_1,\ldots,v_k\}$.

Let G be a graph. The neighbourhood of a vertex u, denoted $N_G(u)$, is the set of vertices $\{v \in V(G) \setminus \{u\} : uv \in E(G)\}$. The closed neighbourhood of a vertex u is defined as $\{u\} \cup N_G(u)$ and denoted as $N_G[u]$. For a path $P = v_1 \cdots v_\ell$, the internal vertices of P, denoted by int(P), are $v_2, \ldots, v_{\ell-1}$. For a vertex set $U \subseteq V(G)$, we denote the compliment of U as \overline{U} . When G is a digraph, its minimum out-degree is denoted as $\delta^+(G)$.

Let H be a k-graph. We write e(H) = |E(H)|. For a set of vertices $U \subseteq V(H)$, H[U] denotes the subgraph induced on U. For k-graphs G and H, $H \setminus G$ denotes the subgraph obtained by deleting V(G) from H and H - G denotes the subgraph after deleting E(G) from E(H). For a vertex set $U \subseteq V(H)$, $H \setminus U = H[V(H) \setminus U]$. The link graph of a vertex z, denoted by H(z), is the (k-1)-graph on V(H) so that $S \in E(H(z))$ if and only if $S \cup z \in E(H)$. For $S \subseteq V(H)$, $N_H(S) = \{T \subseteq V(H) \setminus S : S \cup T \in E(H)\}$ and $d_H(S) = |N_H(S)|$. Denote

$$\delta_{\ell}(H) = \min_{S \in \binom{V(H)}{\ell}} d(S) \text{ and } \Delta_{\ell}(H) = \max_{S \in \binom{V(H)}{\ell}} d(S)$$

We write δ and Δ respectively for the above when $\ell = 1$. For vertex sets V and W, we define $N_H(V, W)$ to be the set $\{e \in \binom{W}{k-|V|}: e \cup V \in E(H)\}$. In particular when $V = \{v\}$ and W = V(H), we have $|N_H(V, W)| = d(v)$.

A set of edges \mathcal{E} in a k-graph H is called *tightly connected* if for any pair of distinct edges $e, f \in \mathcal{E}$, there exists a sequence of edges $e_1, \ldots, e_t \in E(H)$ such that $e_1 = e, e_t = f$ and for $i \in [t-1]$, $|e_i \cap e_{i+1}| = k - 1$. A *tight component* of a k-graph is a set of edges that is maximal with respect to this property. Note that we treat tight components as k-graphs. A k-partite k-graph H has a partition of its vertex set into distinct and disjoint vertex classes V_1, \ldots, V_k such that for any edge $e \in E(H)$, $|e \cap V_i| = 1$ for each $i \in [k]$. For vertex sets $X_1, \ldots, X_k \subseteq V(H)$, an $X_1X_2 \ldots X_k$ -edge is an edge $x_1 \ldots x_k$ where for each $i \in [k]$, we have $x_i \in X_i$.

We work with edge-coloured multigraphs and use the following notation frequently. We denote the edge-colouring by ϕ . Let G be an edge-coloured (multi-)k-graph. The colour set $\phi(G)$ of G is the set of colours that appear in G. When G is a multi-k-graph, for an edge $e, \phi(e)$ is known from the context. For a colour subset $C \subseteq \phi(G)$, G_C is the induced subgraph of G with edges of colours in C after removing any isolated vertices. Let $c \in \phi(G)$ and $S \subseteq V(G)$. We write $N_{c,G}(S)$ for $N_{G_c}(S)$ and $d_{c,G}(S) = |N_{c,G}(S)|$. We say S sees a colour c in a graph G, if $d_{c,G}(S) > 0$. The set of colours that are seen by S is denoted by $\phi_G(S)$. We write $\delta_{\text{mon}}(G) = \min_{c \in \phi(G)} \delta(G_c)$.² Let $V^*(G) = \{v \in V(G) : |\phi_G(v)| \ge 2\}$. We also define

$$\delta_{\mathrm{mon}}^*(G) = \min_{v \in V^*(G)} \min_{c \in \phi_G(v)} d_c(v).$$

An edge-coloured k-graph H is locally r-edge-coloured if any set of k-1 vertices see at most r colours in H. It is easy to check that an r-edge-coloured k-graph is also locally r-edge-coloured. A monochromatic tight component is a tight component that is monochromatic, i.e. between any two edges in the monochromatic tight component there is a monochromatic tight path in a colour fixed for that component. Note that a monochromatic tight component T has colour $\phi_H(T)$.

We drop the subscript when the underlying k-graph is clear from context. Additional notation will be introduced within a section if only needed there.

We use the following three standard concentration inequalities in this work.

²Note that this is different from the monochromatic colour degree in literature, as G_c has no isolated vertices.

Lemma 3.1 (Chernoff Bound c.f. [13, Remark 2.5]). Let $0 < \delta \leq 3/2$ and $X \sim Bin(n,p)$. Then $\mathbb{P}(|X - \mathbb{E}(X)| \geq \delta \mathbb{E}(X)) \leq 2e^{-\frac{\delta^2 \mathbb{E}(X)}{3}}$.

Let N, n, m be positive integers such that $\max\{n, m\} \leq N$. The hypergeometric distribution Hyp(N, n, m) is the distribution of the random variable X obtained by choosing a set σ of m elements from a set τ of N elements, with $X = |\sigma \cap \tau_n|$ where τ_n is a random subset of τ with n elements. We need the following concentration inequality for hypergeometric random variables.

Lemma 3.2 (Hoeffding's inequality c.f. [13, Theorem 2.10]). Let $0 < \varepsilon \leq 3/2$ and $X \sim \text{Hyp}(N, n, m)$. Then $\mathbb{P}[|X - \mathbb{E}[X]| \geq \varepsilon \mathbb{E}[X]] \leq 2e^{-\frac{\varepsilon^2}{3}\mathbb{E}[X]}$.

Lemma 3.3 (Azuma's inequality c.f. [13, Theorem 2.25]). Let $Z_0, ..., Z_n$ be a martingale with $|Z_i - Z_{i-1}| \le c_i$ for all $i \in [n]$. Then for all a > 0,

$$\mathbb{P}[|Z_n - Z_0| \ge a] \le 2e^{-\frac{a^2}{2\sum_{i=1}^n c_i^2}}$$

4 Hypergraph regularity

In this section, we formulate the notion of hypergraph regularity that we use, closely following the formulation from Allen, Böttcher, Cooley and Mycroft [2]. Recall that a hypergraph \mathcal{H} is an ordered pair $(V(\mathcal{H}), E(\mathcal{H}))$, where $E(\mathcal{H}) \subseteq 2^{V(\mathcal{H})}$. We identify the hypergraph \mathcal{H} with its edge set $E(\mathcal{H})$. A subgraph \mathcal{H}' of \mathcal{H} is a hypergraph with $V(\mathcal{H}') \subseteq V(\mathcal{H})$ and $E(\mathcal{H}') \subseteq E(\mathcal{H})$. It is spanning if $V(\mathcal{H}') = V(\mathcal{H})$. For $U \subseteq V(\mathcal{H})$, we define $\mathcal{H}[U]$ to be the subgraph of \mathcal{H} with $V(\mathcal{H}[U]) = U$ and $E(\mathcal{H}[U]) = \{e \in E(\mathcal{H}) : e \subseteq U\}$.

A hypergraph \mathcal{H} is called a *complex* if \mathcal{H} is down-closed, that is if for an edge $e \in \mathcal{H}$ and $f \subseteq e$, then $f \in \mathcal{H}$. A *k*-complex is a complex having only edges of size at most *k*. We denote by $\mathcal{H}^{(i)}$ the spanning subgraph of \mathcal{H} containing only the edges of size *i*. Let \mathcal{P} be a partition of $V(\mathcal{H})$ into vertex classes V_1, \ldots, V_s . Then we say that a set $S \subseteq V(\mathcal{H})$ is \mathcal{P} -partite if $|S \cap V_i| \leq 1$ for all $i \in [s]$. For $\mathcal{P}' = \{V_{i_1}, \ldots, V_{i_r}\} \subseteq \mathcal{P}$, we define the subgraph of \mathcal{H} induced by \mathcal{P}' , denoted by $\mathcal{H}[\mathcal{P}']$ or $\mathcal{H}[V_{i_1}, \ldots, V_{i_r}]$, to be the subgraph of $\mathcal{H}[\bigcup \mathcal{P}']$ containing only the edges that are \mathcal{P}' -partite. The hypergraph \mathcal{H} is said to be \mathcal{P} -partite if all of its edges are \mathcal{P} -partite. We say that \mathcal{H} is *s*-partite if it is \mathcal{P} -partite for some partition \mathcal{P} of $V(\mathcal{H})$ into *s* parts. Let \mathcal{H} be a \mathcal{P} -partite hypergraph. If *X* is a *k*-set of vertex classes of \mathcal{H} , then we write \mathcal{H}_X for the *k*-partite subgraph of $\mathcal{H}^{(k)}$ induced by $\bigcup X$, whose vertex classes are the elements of *X*. Moreover, we denote by $\mathcal{H}_{X^<}$ the *k*-partite hypergraph with $V(\mathcal{H}_{X^<}) = \bigcup X$ and $E(\mathcal{H}_{X^<}) = \bigcup_{X' \subseteq X} \mathcal{H}_{X'}$. In particular, if \mathcal{H} is a complex, then $\mathcal{H}_{X^<}$ is a (k-1)-complex because *X* is a set of size *k*.

Let $i \geq 2$ and let \mathcal{P}_i be a partition of a vertex set V into i parts. Let H_i and H_{i-1} be a \mathcal{P}_i -partite i-graph and a \mathcal{P}_i -partite (i-1)-graph on a common vertex set V, respectively. We say that a \mathcal{P}_i -partite i-set in V is supported on H_{i-1} if it induces a copy of the complete (i-1)-graph $K_i^{(i-1)}$ on i vertices in H_{i-1} . We denote by $K_i(H_{i-1})$ the \mathcal{P}_i -partite i-graph on V whose edges are all \mathcal{P}_i -partite i-sets contained in V which are supported on H_{i-1} . Now we define the density of H_i with respect to H_{i-1} to be

$$d(H_i \mid H_{i-1}) = \frac{|K_i(H_{i-1}) \cap H_i|}{|K_i(H_{i-1})|}$$

if $|K_i(H_{i-1})| > 0$ and $d(H_i | H_{i-1}) = 0$ if $|K_i(H_{i-1})| = 0$. So $d(H_i | H_{i-1})$ is the proportion of \mathcal{P}_i -partite copies of K_i^{i-1} in H_{i-1} which are also edges of H_i . More generally, if $\mathbf{Q} = (Q_1, Q_2, \dots, Q_r)$ is a collection

of r (not necessarily disjoint) subgraphs of H_{i-1} , we define $K_i(\mathbf{Q}) = \bigcup_{j=1}^r K_i(Q_j)$ and

$$d(H_i \mid \mathbf{Q}) = \frac{|K_i(\mathbf{Q}) \cap H_i|}{|K_i(\mathbf{Q})|}$$

if $|K_i(\mathbf{Q})| > 0$ and $d(H_i | \mathbf{Q}) = 0$ if $|K_i(\mathbf{Q})| = 0$. We say that H_i is (d_i, ε, r) -regular with respect to H_{i-1} , if we have $d(H_i | \mathbf{Q}) = d_i \pm \varepsilon$ for every r-set \mathbf{Q} of subgraphs of H_{i-1} with $|K_i(\mathbf{Q})| > \varepsilon |K_i(H_{i-1})|$. We say that H_i is (ε, r) -regular with respect to H_{i-1} if there exists some d_i for which H_i is (d_i, ε, r) -regular with respect to H_{i-1} . Finally, given an *i*-graph G whose vertex set contains that of H_{i-1} , we say that Gis (d_i, ε, r) -regular with respect to H_{i-1} if the *i*-partite subgraph of G induced by the vertex classes of H_{i-1} is (d_i, ε, r) -regular with respect to H_{i-1} . We refer to the density of this *i*-partite subgraph of Gwith respect to H_{i-1} as the relative density of G with respect to H_{i-1} .

Now let $s \geq k \geq 3$ and let \mathcal{H} be an s-partite k-complex on vertex classes V_1, \ldots, V_s . For any set $A \subseteq [s]$, we write V_A for $\bigcup_{i \in A} V_i$. Note that, if $e \in \mathcal{H}^{(i)}$ for some $2 \leq i \leq k$, then the vertices of e induce a copy of K_i^{i-1} in $\mathcal{H}^{(i-1)}$. Therefore, for any set $A \in \binom{[s]}{i}$, the density $d(\mathcal{H}^{(i)}[V_A] \mid \mathcal{H}^{(i-1)}[V_A])$ is the proportion of 'possible edges' of $\mathcal{H}^{(i)}[V_A]$, which are indeed edges. We say that \mathcal{H} is $(d_k, \ldots, d_2, \varepsilon_k, \varepsilon, r)$ -regular if

(a) for any $2 \leq i \leq k-1$ and any $A \in {[s] \choose i}$, the induced subgraph $\mathcal{H}^{(i)}[V_A]$ is $(d_i, \varepsilon, 1)$ -regular with respect to $\mathcal{H}^{(i-1)}[V_A]$ and

(b) for any $A \in {\binom{[s]}{k}}$, the induced subgraph $\mathcal{H}^{(k)}[V_A]$ is (d_k, ε_k, r) -regular with respect to $\mathcal{H}^{(k-1)}[V_A]$.

For $\mathbf{d} = (d_k, \dots, d_2)$, we write $(\mathbf{d}, \varepsilon_k, \varepsilon, r)$ -regular to mean $(d_k, \dots, d_2, \varepsilon_k, \varepsilon, r)$ -regular. We say that a (k-1)-complex \mathcal{J} is (t_0, t_1, ε) -equitable if it has the following properties.

- (a) \mathcal{J} is \mathcal{P} -partite for some \mathcal{P} which partitions $V(\mathcal{J})$ into t parts, where $t_0 \leq t \leq t_1$, of equal size. We refer to \mathcal{P} as the ground partition of \mathcal{J} and to the parts of \mathcal{P} as the clusters of \mathcal{J} .
- (b) There exists a *density vector* $\mathbf{d} = (d_{k-1}, \ldots, d_2)$ such that, for each $2 \leq i \leq k-1$, we have $d_i \geq 1/t_1$ and $1/d_i \in \mathbb{N}$ and \mathcal{J} is $(\mathbf{d}, \varepsilon, \varepsilon, 1)$ -regular.

For any k-set X of clusters of \mathcal{J} , we denote by $\hat{\mathcal{J}}_X$ the k-partite (k-1)-graph $(\mathcal{J}_{X^{<}})^{(k-1)}$ and call $\hat{\mathcal{J}}_X$ a polyad. Given a (t_0, t_1, ε) -equitable (k-1)-complex \mathcal{J} and a k-graph G on $V(\mathcal{J})$, we say that G is (ε_k, r) -regular with respect to a k-set X of clusters of \mathcal{J} if there exists some d such that G is (d, ε_k, r) regular with respect to the polyad $\hat{\mathcal{J}}_X$. Moreover, we write $d^*_{G,\mathcal{J}}(X)$ for the relative density of G with respect to $\hat{\mathcal{J}}_X$; we may drop either subscript if it is clear from context.

We can now give the crucial definition of a regular slice.

Definition 4.1 (Regular slice). Given ε , $\varepsilon_k > 0, r, t_0, t_1 \in \mathbb{N}$, a k-graph G, a (k-1)-complex \mathcal{J} on V(G), is a $(t_0, t_1, \varepsilon, \varepsilon_k, r)$ -regular slice for G if \mathcal{J} is (t_0, t_1, ε) -equitable and G is (ε_k, r) -regular with respect to all but at most $\varepsilon_k {t \choose k}$ of the k-sets of clusters of \mathcal{J} , where t is the number of clusters of \mathcal{J} .

If we specify the density vector **d** and the number of clusters t of an equitable complex or a regular slice, then it is not necessary to specify t_0 and t_1 (since the only role of these is to bound **d** and t). In this situation we write that \mathcal{J} is $(\cdot, \cdot, \varepsilon)$ -equitable, or is a $(\cdot, \cdot, \varepsilon, \varepsilon_k, r)$ -regular slice for G.

Given a regular slice \mathcal{J} for a k-graph G, we define the d-reduced k-graph $\mathcal{R}^{\mathcal{J}}_{d}(G)$ as follows.

Definition 4.2 (The d-reduced k-graph). Let $k \geq 3$. Let G be a k-graph and let \mathcal{J} be a $(t_0, t_1, \varepsilon, \varepsilon_k, r)$ regular slice for G. Then, for d > 0, we define the d-reduced k-graph $\mathcal{R}^{\mathcal{J}}_d(G)$ to be the k-graph whose
vertices are the clusters of \mathcal{J} and whose edges are all k-sets X of clusters of \mathcal{J} such that G is (ε_k, r) regular with respect to X and $d^*(X) \geq d$.

We now state the statement of the Regular Slice Lemma that we need, that is a straightforward consequence of [2, Lemma 10].

Lemma 4.3 (Regular Slice Lemma [2, Lemma 10]). Let $k \ge 3$. For all $t_0, r, s \in \mathbb{N}$, $\varepsilon_k > 0$ and all functions $r' \colon \mathbb{N} \to \mathbb{N}$ and $\varepsilon \colon \mathbb{N} \to (0, 1]$, there are integers t_1 and n_0 such that the following holds for all $n \ge n_0$ which are divisible by t_1 !. Let H be an r-edge-coloured $K_n^{(k)}$ with colour set [r]. For $i \in [r]$, let H_i denote the monochromatic subgraph of H in colour i. Then there exists a (k-1)-complex \mathcal{J} on V(H) which is a $(t_0, t_1, \varepsilon(t_1), \varepsilon_k, r'(t_1))$ -regular slice for each H_i .

The following lemma which is a direct generalisation of [16, Lemma 12] shows that the union of the corresponding reduced graphs $\bigcup_{i \in [r]} \mathcal{R}_d^{\mathcal{J}}(H_i)$ is almost complete.

Lemma 4.4. Let $k \geq 3$, $\varepsilon, \varepsilon_k > 0, r \in \mathbb{N}$ and $r' : \mathbb{N} \to \mathbb{N}$. Let H be an r-edge-coloured $K_n^{(k)}$ and for $i \in [r]$ let \mathcal{J} be a $(\cdot, \cdot, \varepsilon, \varepsilon_k, r')$ -regular slice for each H_i . Let t be the number of clusters of \mathcal{J} . Then, provided that $d \leq 1/r$, we have $\left| \bigcup_{i \in [r]} \mathcal{R}_d^{\mathcal{J}}(H_i) \right| \geq (1 - r\varepsilon_k) {t \choose k}$.

Proof. Since \mathcal{J} is a $(\cdot, \cdot, \varepsilon, \varepsilon_k, r)$ -regular slice for each H_i there are at least $(1 - r\varepsilon_k) {t \choose k}$ k-sets X of clusters of \mathcal{J} such that each H_i is (ε_k, r) -regular with respect to X. Let X be such a k-set. Since the H_i are edge-disjoint and $\bigcup_{i \in [r]} H_i = H$, we have $\sum_{i \in [r]} d^*_{H_i}(X) = 1$. Hence for some $i \in [r], d^*_{H_i}(X) \ge 1/r$ and thus, since $d \le 1/r$, we have $X \in \bigcup_{i \in [r]} \mathcal{R}^{\mathcal{J}}_d(H_i)$.

Let $r \in \mathbb{N}, 0 < d \leq 1/r, \varepsilon > 0$ and $0 < \varepsilon_k < 1/2r$. Let H be an r-edge-coloured $K_n^{(k)}$ on colour set [r]. Let \mathcal{J} be a $(\cdot, \cdot, \varepsilon, \varepsilon_k, r')$ -regular slice for each H_i with partition \mathcal{P} of V(H). By Lemma 4.4, we have $|\mathcal{R}_d^{\mathcal{J}}(H)| \geq (1 - r\varepsilon_k) {t \choose k} \geq {t \choose k}/2$. Given the following natural edge-colouring, we will be working with an edge-coloured reduced graph. For an edge X in $\mathcal{R}_d^{\mathcal{J}}(H)$, let $e_{H,i}(X)$ denote the number of edges in colour i among the vertex clusters in H that are induced by vertices in X. Assign each such k-set X the colour i if $e_{H,i}(X) \geq e_H(X)/r$ (if multiple colours satisfy this, then choose one of them arbitrarily). Note that $\mathcal{R}_d^{\mathcal{J}}(H)$ is an r-edge-coloured k-graph with at least ${t \choose k}/2$ edges. We write \mathcal{R} for $\mathcal{R}_d^{\mathcal{J}}(H)$. For each $v \in V(\mathcal{R})$, we denote V_v to be the cluster of \mathcal{P} corresponding to v.

Let *H* be a *k*-graph. A fractional matching in *H* is a function $\omega : E(H) \to [0,1]$ such that for all $v \in V(H), \omega(v) \coloneqq \sum_{e \in H: v \in e} \omega(e) \leq 1$. The weight of the fractional matching is defined to be $\sum_{e \in H} \omega(e)$. A fractional matching is tightly connected if the subgraph induced by the edges *e* with $\omega(e) > 0$ is tightly connected in *H*. For a fractional matching ω , we write the size of ω as $|\omega| = \sum_{e \in E(G)} \omega(e)$.

We would use the following lemma to lift a tightly connected fractional matching in the reduced graph to a monochromatic tight cycle covering almost all vertices in the corresponding clusters.

Lemma 4.5 ([16, Lemma 18]). Let $1/n \ll 1/r'$, $\varepsilon \ll \varepsilon_k$, d_{k-1}, \ldots, d_2 and $\varepsilon_k \ll \varepsilon' \ll \psi$, d_k , β , $1/k \le 1/3$ and $1/n \ll 1/t$ such that t divides n and $1/d_i \in \mathbb{N}$ for all $2 \le i \le k-1$. Let G be a k-graph on n vertices and \mathcal{J} be a $(\cdot, \cdot, \varepsilon, \varepsilon_k, r')$ -regular slice for G. Further, let \mathcal{J} have t clusters V_1, \ldots, V_t all of size n/t and density vector $\mathbf{d} = (d_{k-1}, \ldots, d_2)$. Suppose that the reduced graph $\mathcal{R}_{d_k}^{\mathcal{J}}(G)$ contains a tightly connected fractional matching φ with weight μ . Assume that all edges with non-zero weight have weight at least β . For each $i \in [t]$, let $W_i \subseteq V_i$ be such that $|W_i| \ge ((1 - 3\varepsilon')\varphi(V_i) + \varepsilon')n/t$. Then $G\left[\bigcup_{i \in [t]} W_i\right]$ contains a tight cycle of length ℓ for each $\ell \le (1 - \psi)k\mu n/t$ that is divisible by k.

Another tool we use in the next section is the notion of a hypergraph being (μ, α) -dense. For constants $\mu, \alpha > 0$, we say that a k-graph H on n vertices is (μ, α) -dense if, for each $i \in [k-1]$, we have $d_H(S) \ge \mu \binom{n}{k-i}$ for all but at most $\alpha \binom{n}{i}$ sets $S \in \binom{V(H)}{i}$ and $d_H(S) = 0$ for all other $S \in \binom{V(H)}{i}$. We use the following result from [16].

Proposition 4.6 ([16, Proposition 5]). Let $1/n \ll \alpha \ll 1/k \leq 1/2$. Let H be a k-graph on n vertices with $|H| \geq (1 - \alpha) \binom{n}{k}$. Then there exists a subgraph H' of H such that V(H') = V(H) and H' is $(1 - 2\alpha^{1/4k^2}, 2\alpha^{1/4k})$ -dense.

5 The reduction of Lemma 2.3

We now generalise the triangle cycle from Erdős, Gyárfás and Pyber [7]. Let t, k, m be positive integers such that m = (k - 1)t. The *k*-uniform triangle cycle $T_m^{(k)}$ consists of a *k*-uniform tight cycle $a_1 \cdots a_m$ and a vertex set $B = \{b_1, \ldots, b_t\}$ such that

$$a_{(k-1)i-(k-2)}a_{(k-1)i-(k-3)}\dots a_{(k-1)i}b_ia_{(k-1)i+1}\dots a_{(k-1)i+(k-1)}$$

is a k-uniform tight path for all $i \in [t]$ (when subscripts are taken modulo m). Moreover, there is a tight cycle on $V\left(T_m^{(k)}\right) \setminus B'$ for all $B' \subseteq B$. Note that $\Delta\left(T_m^{(k)}\right) = 2k$. For a k-graph H and $r \in \mathbb{N}$, the multicolour Ramsey number $R_r(H)$ is the minimum positive integer

For a k-graph H and $r \in \mathbb{N}$, the multicolour Ramsey number $R_r(H)$ is the minimum positive integer N such that any r-edge-coloured $K_N^{(k)}$ contains a monochromatic copy of H. We use the following result of Conlon, Fox and Sudakov on the Ramsey number of hypergraphs with bounded maximum degree, to obtain a monochromatic triangle cycle $T_m^{(k)}$.

Theorem 5.1 (Conlon, Fox and Sudakov [6, Theorem 5]). Let $\Delta, k, r \in \mathbb{N}$. There exists a constant $c(\Delta, k, r)$ such that if H is a k-graph on n vertices with maximum degree Δ , then $R_r(H) \leq c(\Delta, k, r)n$.

Note that $c(\Delta, k, r)$ is a tower-type function in r and k. Since $T_m^{(k)}$ has bounded degree, we get the following corollary.

Corollary 5.2. Let $m, n, r, k \in \mathbb{N}$ and $1/n \ll \alpha \ll 1/r, 1/k$ be such that $m = (k-1)\alpha n$. Then a monochromatic triangle cycle $T_m^{(k)}$ exists in every r-edge-coloured $K_n^{(k)}$.

The main aim of this section is to reduce Lemma 2.3 to the following lemma.

Lemma 5.3. Let $k, r \in \mathbb{N}$, $\delta_0 = (2r)^{-2^{k+1}}$ and $\alpha = 2^5 r k (2r)^{2^{k+1}}$. Let H be an r-edge-coloured $K_n^{(k)}$ and X, Z be disjoint subsets of V(H) such that $|X| \ge \alpha |Z|$. Then there exists a set C of vertex-disjoint monochromatic tight cycles covering Z with $|\mathcal{C}| \le 2^{18} r \delta_0^{-5}$ and $|V(\mathcal{C}) \cap X| = (k-1)|Z|$.

Suppose that Lemma 5.3 holds. A naive approach of proving Lemma 2.3 is to first reserve a monochromatic $T_m^{(k)}$ using Corollary 5.2. We then apply Lemma 5.3 with B (from $T_m^{(k)}$) playing the role of X to cover the remaining vertices. However, $T_m^{(k)}$ obtained is too short (unless the number of cycles we remove using Proposition 2.2 in the proof of Theorem 1.1 is a super polynomial of r). In order to achieve this, we use the following lemma, which is motivated by Gyárfás, Ruszinkó, Sárközy and Szemerédi [11].

Lemma 5.4. Let $1/n \ll \psi \ll 1/r$, 1/k. Let $\delta_0 = (2r)^{-2^{k+1}}$ and $\varepsilon^* = (2r)^{-2^{k+7}}/k$. Let H be an r-edgecoloured $K_n^{(k)}$. Then there exists a vertex set U^* with $|U^*| \leq 3n/4$ such that, for any vertex subset B^* of $V(H) \setminus U^*$ with $|B^*| = 4\varepsilon^* n$, $H[B^* \cup U^*]$ can be covered by at most $2^{18}r\delta_0^{-5} + 1$ monochromatic tight cycles and at most ψn isolated vertices.

This lemma will be proved in Section 5.2. Observe that this lemma significantly reduces the size of the leftover vertex set. However, we would require precisely $4\varepsilon^*n$ leftover vertices, which may not be guaranteed by Proposition 2.2. We deal with it using the following lemma.

Lemma 5.5. Let $1/n \ll \psi \ll 1/r, 1/k$. Let H be an r-edge-coloured $K_n^{(k)}$. Then there exists a vertex subset $W \subseteq V(H)$ such that $n/5r \leq |W| \leq n/4r$ and H[W] contains a monochromatic tight cycle of length ℓ for all $\ell \leq (1 - \psi)|W|$ with $\ell \equiv 0 \pmod{k}$.

Proof. Let $d_k = 1/r$,

$$1/n \ll 1/t_1 \ll 1/t_0,$$

$$1/n \ll 1/r', \tilde{\varepsilon} \ll \varepsilon_k, d_{k-1}, \dots, d_2,$$

$$\varepsilon_k \ll \varepsilon' \ll \psi \ll 1/r, 1/k.$$

Let H' be an induced subgraph of H such that $|V(H')| \equiv 0 \pmod{t_1!}$ and $|V(H \setminus H')| < t_1!$. By Lemma 4.3 with H' playing the role of H, there exists a $(\cdot, \cdot, \tilde{\varepsilon}, \varepsilon_k, r')$ -regular slice $\tilde{\mathcal{J}}$ with partition \mathcal{P} of V(H'). Let m = |V| for $V \in \mathcal{P}$ and $t = |\mathcal{P}|$. Note that $mt \geq n - t_1!$.

Let $\mathcal{R} = \mathcal{R}_{d_k}^{\tilde{\mathcal{J}}}(H')$. Recall that for each $x \in V(\mathcal{R})$, V_x denotes the corresponding cluster in \mathcal{P} . By Lemma 4.4 (with $d = d_k$), \mathcal{R} contains at least $(1 - r\varepsilon_k) {t \choose k} \ge {t \choose k}/2$ edges. There exists a colour $j \in [r]$ such that $|E(\mathcal{R}_j)| \ge {t \choose k}/2r$. By Theorem 2.1 with $\delta = \alpha = 1/4r$, \mathcal{R}_j contains a monochromatic tight cycle $C = v_1 v_2 \dots v_\ell$ with $t/5r \le \ell \le t/4r$ and $\ell \equiv 0 \pmod{k}$. Set $W = \bigcup_{i \in [\ell]} V_{v_i}$. Let φ be a fractional matching on C such that for any edge $e \in E(C)$, we have $\varphi(e) = 1/k$. For $i \in [\ell]$, note that $\varphi(v_i) = 1$ and $|V_{v_i}| \ge ((1 - 3\varepsilon')\varphi(v_i) + \varepsilon')m$. Therefore, Lemma 4.5 implies our result.

5.1 Proof of Lemma 2.3 assuming Lemmas 5.3 and 5.4

We now prove Lemma 2.3 assuming Lemmas 5.3 and 5.4.

Proof of Lemma 2.3. Set

$$\delta_0 = (2r)^{-2^{k+1}}, \ \varepsilon^* = (2r)^{-2^{k+7}}/k, \ \alpha = 2^5 r k (2r)^{2^{k+1}}, \ \alpha_\Delta = 2\alpha\psi \text{ and } \tilde{m} = (k-1)\alpha_\Delta N.$$
(5.1)

By Corollary 5.2 with α_{Δ} , \tilde{m} playing the roles of α, m , respectively, H contains a monochromatic kuniform triangle cycle $T_{\tilde{m}}^{(k)}$. Note that $T_{\tilde{m}}^{(k)}$ contains a vertex-set \tilde{B} with

$$|\tilde{B}| = \tilde{m}/(k-1) = \alpha_{\Delta}N \tag{5.2}$$

such that for any $\tilde{B}' \subseteq \tilde{B}$, there is a monochromatic tight cycle on $V\left(T_{\tilde{m}}^{(k)} \setminus \tilde{B}'\right)$. Note that $\left|V\left(T_{\tilde{m}}^{(k)}\right)\right| = \tilde{m} + \alpha_{\Delta}N = k\alpha_{\Delta}N$. Let $H_1 = H \setminus T_{\tilde{m}}^{(k)}$ and

$$n = |V(H_1)| = N - k\alpha_{\Delta}N \ge N/2. \tag{5.3}$$

Lemma 5.4 with H_1 playing the role of H implies that there exists a vertex set $U^* \subseteq V(H_1)$ with $|U^*| \leq 3n/4$ such that, for any $B^* \subseteq V(H_1) \setminus U^*$ with $|B^*| = 4\varepsilon^* n$, $H[B^* \cup U^*]$ can be covered by at most $2^{18}r\delta_0^{-5} + 1$ vertex-disjoint monochromatic tight cycles and at most ψn isolated vertices. Let $H_2 = H_1 \setminus U^*$ and $n' = |V(H_2)| = n - |U^*| \geq n - 3n/4 = n/4$.

By Lemma 5.5 with H_2, n' playing the role of H, n, respectively, there exists a vertex subset $W \subseteq V(H_2)$ with

$$n/20r \le n'/5r \le |W| \le n'/4r \le n/4r$$

such that H[W] contains a monochromatic tight cycle of length ℓ for all $\ell \leq (1 - \psi)|W|$ with $\ell \equiv 0 \pmod{k}$.

Set $R = V\left(T_{\tilde{m}}^{(k)}\right) \cup W \cup U^*$. We now show that R has the desired properties. Let B be a subset of $V(H) \setminus R$ with

$$|B| \le \varepsilon^* N \stackrel{(5.3)}{\le} 2\varepsilon^* n.$$

Let ℓ be the largest integer such that $\ell \equiv 0 \pmod{k}$ and $|B| + |W| - \ell \geq 2\varepsilon^* n$. Since $\psi \ll \varepsilon^*$, $\ell \leq (1 - \psi)|W|$ and therefore there exists a monochromatic tight cycle C_1 in H[W] of length ℓ . Let B^+ and L_1 be disjoint vertex sets such that

$$B^+ \cup L_1 = B \cup (W \setminus V(C_1)), \ |B^+| = 4\varepsilon^* n \text{ and } |L_1| < k.$$
 (5.4)

Thus $H[U^* \cup B^+]$ can be covered by a set C of monochromatic tight cycles such that $|C| \leq 2^{18} r \delta_0^{-5} + 1$ and

$$|U^* \setminus V(\mathcal{C})| \le \psi n \le \psi N. \tag{5.5}$$

Let $L = L_1 \cup (U^* \setminus V(\mathcal{C}))$. We deduce that

$$|L| = |L_1| + |U^* \setminus V(\mathcal{C})| \stackrel{(5.4), (5.5)}{\leq} k + \psi N \stackrel{(5.1)}{\leq} \alpha_{\Delta} N / \alpha \stackrel{(5.2)}{=} |\tilde{B}| / \alpha.$$

By Lemma 5.3 with \tilde{B}, L playing the roles of X, Z, respectively, $H[\tilde{B} \cup L]$ contains a set \mathcal{C}' of vertexdisjoint monochromatic tight cycles such that $L \subseteq V(\mathcal{C}')$ and $|\mathcal{C}'| \leq 2^{18} r \delta_0^{-5}$. By the property of $T_{\tilde{m}}^{(k)}$, there exists a monochromatic tight cycle C_2 with vertex set $V\left(T_{\tilde{m}}^{(k)}\right) \setminus (\tilde{B} \cap V(\mathcal{C}'))$. Thus, we have partitioned $R \cup B$ into monochromatic cycles, namely $\mathcal{C} \cup \mathcal{C}' \cup \{C_1, C_2\}$. The total number of cycles created is at most $2^{19} r \delta_0^{-5} + 3 = 2^{19} r (2r)^{5 \cdot 2^{k+1}} + 3$. This completes the proof of the lemma.

5.2 Proof of Lemma 5.4

The proof of Lemma 5.4 is motivated by Gyárfás, Ruszinkó, Sárközy and Szemerédi [11]. Let $t \in \mathbb{N}$. A *t*-half dense matching in a *k*-partite *k*-graph *H* with vertex classes X_1, \ldots, X_k is a matching $M = \{x_{i,1}x_{i,2}\ldots x_{i,k}: i \in [\ell]\}$ where for all $i \in [\ell], x_{i,j} \in X_j$ and for any vertex $x_1 \in V(M) \cap X_1$, we have

$$|\{j \in [\ell] : x_1 x_{j,2} \dots x_{j,k} \in E(H)\}| \ge t.$$

We say a matching M is *t-semi-dense* if for all $i \in [\ell]$, we have

$$|\{j \in [\ell] : x_{i,1}x_{i,2} \dots x_{i,k-1}x_{j,k} \in E(H)\}| \ge t.$$

Note that the order of $X_1, \ldots X_k$ matters, but it will be clear from context. We shall need the following simple fact.

Fact 5.6. A matching in a 2-graph is t-semi-dense if and only if it is t-half-dense.

We need the following result by Gyárfás, Ruszinkó, Sárközy and Szemerédi [11].

Lemma 5.7 ([11, Lemma 4]). Every graph of average degree 8k has a connected k-half dense matching.

Recall that a k-graph H is locally r-edge-coloured if any set of k-1 vertices sees at most r colours in H. We use the following result by Gyárfás, Lehel, Nešetřil, Rödl, Schelp and Tuza [10].

Lemma 5.8 ([10, Corollary 3]). Let G be a graph with average degree d that is r-locally edge-coloured. Then there is a monochromatic subgraph G' such that $|E(G')| \ge d^2/2r^2$. For a k-graph H, we denote by ∂H the (k-1)-graph on V(H) whose edges are all (k-1)-tuples of vertices contained in some edge in H. Therefore, $|E(\partial H)| \leq k|E(H)|$. Recall that a monochromatic tight component is a tight component that is monochromatic, i.e. there is a sequence of edges of the same colour joining any two edges in the monochromatic tight component, such that any two consecutive edges in the sequence share k-1 vertices.

The following lemma shows that one can find a linear-sized semi-dense matching in a monochromatic tight component of a locally r-edge-coloured almost complete k-graph.

Lemma 5.9. Let $0 < 1/t \ll \varepsilon_k \ll \varepsilon_{k-1} \ll \cdots \ll \varepsilon_2 \ll 1/r, 1/k$ and $\delta(r,k) = \left(2^{2^k+3k-5}r^{2^k-2}\right)^{-1}$. Let R be a locally r-edge-coloured k-graph on t vertices with $|E(R)| \ge (1-\varepsilon_k)\binom{t}{k}$. Then there exists a monochromatic tight component of R containing a $\delta(r,k)t$ -semi-dense matching.

Proof. We prove the lemma by induction on k. Note that

$$\delta(r,2) = \left(2^5 r^2\right)^{-1} \text{ and } \delta(r,k+1) = \delta(2r^2,k)/2^5 r^2.$$
(5.6)

Suppose that k = 2. The average degree of R is

$$\frac{2|E(R)|}{t} \ge \frac{2(1-\varepsilon_2)\binom{t}{2}}{t} \ge \frac{t}{2}.$$

By Lemma 5.8, there is a monochromatic subgraph R' such that $|E(R')| \ge t^2/8r^2$. The average degree of R' is

$$\frac{2|E(R')|}{t} \ge \frac{t}{4r^2} = 8\delta(r,2)t.$$

By Lemma 5.7, R' contains a connected $\delta(r, 2)t$ -half-dense matching. By Fact 5.6, we are done.

Thus, we may assume $k \geq 3$. By Proposition 4.6 with ε_k , t playing the role of α , n, respectively, there exists a spanning subgraph R' of R that is $(1-2\varepsilon_k^{1/4k^2}, 2\varepsilon_k^{1/4k})$ -dense. Let \mathcal{T} be the set of monochromatic tight components of R'. For $T \in \mathcal{T}$, let $\phi_{R'}(T)$ be the colour of T. Let

$$n = \frac{t}{8r}.\tag{5.7}$$

Partition V(R') into V, W with |V| = n. Let $G = (\partial R')[V]$, so G is a (k-1)-graph. Note that

$$|E(G)| \ge \binom{n}{k-1} - 2\varepsilon_k^{1/4k^2} \binom{t}{k-1} \ge (1-\varepsilon_{k-1})\binom{n}{k-1}$$

For every edge $e \in E(G)$, we have

$$d_{R'}(e, W) \ge |W| - 2\varepsilon_k^{1/4k^2} t \ge |W|/2.$$

Define an edge-colouring ϕ_G of G with colour set \mathcal{T} so that, for an edge $e \in E(G)$, we have

$$\phi_G(e) = T$$
 if $d_{T,G}(e, W) \ge |W|/2r$.

If multiple $T \in \mathcal{T}$ satisfy this, pick one such T arbitrarily.

We now show that G is locally $2r^2$ -edge-coloured. Suppose for a contradiction, there exists a set S of k-2 vertices x_1, \ldots, x_{k-2} in G such that $|\phi_G(S)| \ge 2r^2 + 1$. Since R' is locally r-edge-coloured, there exist a colour $c \in \phi(R')$, vertices y_1, \ldots, y_{2r+1} and distinct tight components T_1, \ldots, T_{2r+1} in R' for which $\phi_{R'}(T_i) = c$ and $\phi_G(S \cup y_i) = T_i$ for all $i \in [2r+1]$. Then there exist distinct $i, j \in [2r+1]$ such that $N_{T_i}(S \cup y_i, W) \cap N_{T_j}(S \cup y_j, W) \neq \emptyset$. Let $w \in N_{T_i}(S \cup y_i, W) \cap N_{T_j}(S \cup y_j, W)$. Note that $y_i x_1 \dots x_{k-2} w y_j$ is a tight path in R' where both edges are coloured c implying $T_i = T_j$, a contradiction.

By our induction hypothesis, G contains a monochromatic (k-1)-uniform $\delta(2r^2, k-1)n$ -semi-dense matching $M^{(k-1)}$ of size ℓ . Note that

$$\delta(2r^2, k-1)n \le \ell \le n/2 = t/16r.$$

Let $T_0 \in \mathcal{T}$ be the colour of $M^{(k-1)}$ in G. Let $M^{(k-1)} = \{x_{i,1} \dots x_{i,k-1} : i \in [\ell]\}$ be such that for each $i \in [\ell]$, we have

$$|\{j \in [\ell] : x_{i,1} \dots x_{i,k-2} x_{j,k-1} \in E(T_0)\}| \ge \delta(2r^2, k-1)n.$$

We now extend $M^{(k-1)}$ to a k-uniform $\delta(r, k)t$ -semi-dense matching $M^{(k)}$ in $T \subseteq R' \subseteq R$ using vertices from W.

Let $\eta = \delta(2r^2, k-1)$ and m = |W|. For each $i \in [\ell]$, let $X^i = \{x_{i,1}, x_{i,2}, \ldots, x_{i,k-2}\}$ and let $X_{k-1} = \{x_{i,k-1} : i \in [\ell]\}$. Delete some edges in $T_0[V(M^{(k-1)})]$ if necessary so that for each $i \in [\ell]$, we have $d_{T_0}[V(M^{(k-1)})](X^i, X_{k-1}) = \eta n$. We now choose vertices w_1, \ldots, w_ℓ from W in turns such that $d_{T_0}(X^i \cup w_i, X_{k-1}) \geq \delta(r, k)t$. Fix $i \in [\ell]$. For each $x' \in N_{G[V(M^{(k-1)})]}(X^i, X_{k-1})$, we have $d_{T_0}(X^i \cup x', W) \geq m/2r$. Therefore, T_0 contains at least $\eta nm/2r$ many $x_{i,1} \ldots x_{i,k-2}X_{k-1}W$ -edges. At most $\ell\eta n \leq \eta n^2/2$ such edges contain a vertex in w_1, \ldots, w_{i-1} . Thus there exists a vertex $w_i \in W \setminus \{w_j : j \in [i-1]\}$ such that

$$d_{T_0}(X^i \cup w_i, X_{k-1}) \ge \frac{\eta n m / (2r) - \eta n^2 / 2}{m} \ge \left(\frac{\eta n}{2r}\right) \left(1 - \frac{rn}{m}\right) \\\ge \frac{\eta n}{4r} = \frac{\delta(2r^2, k-1)n}{4r} \stackrel{(5.6), (5.7)}{\ge} \delta(r, k)t.$$

Let $M^{(k)} = \{w_i x_{i,1} \dots x_{i,k-2} x_{i,k-1} : i \in [\ell]\}$. Note that $M^{(k)}$ is the desired k-uniform semi-dense matching.

We now convert the semi-dense matching into a half-dense matching. The following lemma finds a half-dense matching in a bipartite graph where we control the vertex set with a large minimum degree. Its proof is based on [11, Lemma 3]. For a matching M in a graph G, an M-augmented path is a path in G where every alternate edge is in M.

Lemma 5.10. Let $1/n \ll \delta \leq 1$ and G be a bipartite graph with vertex classes X and Y such that $|X|, |Y| \leq n$ and $|E(G)| \geq \delta n^2$. Then G contains a matching M such that for all $x \in X \cap V(M)$, we have $d_{G[V(M)]}(x) \geq \delta^2 n/8$.

Proof. Let $X' \subseteq X$ and $Y' \subseteq Y$ be such that $\delta(G[X' \cup Y']) \geq \delta n/2$. Let $m = \min\{|X'|, |Y'|\}$, so $m \geq \delta n/2$.

Claim 5.11. There exist subsets $X^* \subseteq X'$ and $Y^* \subseteq Y'$ such that $|X^*| = |Y^*| = m$ and $\delta(G[X^* \cup Y^*]) \ge \delta^2 n/8$.

Proof of claim. Suppose that |X'| = m (and the case |Y'| = m is proved similarly). Let $X^* = X'$. Pick a subset Y^* of Y' of size m uniformly at random. Note that for any $y \in Y^*$, $d(y, X^*) \ge \delta n/2$. Let $x \in X^*$. Note that $d(x, Y^*) \sim \text{Hyp}(|Y'|, m, d(x, Y'))$ and

$$\mathbb{E}d(x, Y^*) = \frac{|Y^*|d(x, Y')|}{|Y'|} \ge \frac{m(\delta n/2)}{n} \ge \frac{\delta^2 n}{4}.$$

Applying Lemma 3.2, we deduce that for $x \in X^*$

$$\mathbb{P}\left(d(x,Y^*) < \frac{\delta^2 n}{8}\right) \le 2e^{-\frac{\delta^2 n}{12}}.$$

Taking a union bound over all $x \in X^*$, we have that with high probability, $\delta(G[X^* \cup Y^*]) \ge \delta^2 n/8$. Fix such X^* and Y^* .

Let $G^* = G[X^* \cup Y^*]$. Pick a largest matching M^* in G^* . If M^* is spanning on $V(G^*)$, then we are done by setting $M = M^*$. Hence we may assume $Y^* \setminus V(M^*) \neq \emptyset$. Let X_1^* be the set of vertices in $V(M^*) \cap X^*$ that can be reached by an M^* -augmented path from $X^* \setminus V(M^*)$. Observe that $E(G^*[X_1^* \cup (Y^* \setminus V(M^*))]) = \emptyset$. Otherwise for a vertex $x \in X_1^*$, let xy be an edge in $E(G^*[X_1^* \cup (Y^* \setminus V(M^*))])$. There exists an M^* -augmented path P between x and $X^* \setminus V(M^*)$. But $(M^* \Delta E(P)) \cup \{xy\}$ is a larger matching, contradicting that M^* is the largest matching.

Let $Y_1^* = N_{G^*}(X_1^*) \subseteq V(M^*) \cap Y^*$, so $E(G[X_1^* \cup (Y^* \cap V(M^*)) \setminus Y_1^*]) = \emptyset$. Since M^* is maximal, $E(G^* \setminus V(M^*)) = \emptyset$. On the other hand, $E(G[X^* \setminus V(M^*), Y^*]) \ge \delta(G^*) > 0$. Hence $E(G[X^* \setminus V(M^*), Y^* \cap V(M^*)]) \neq \emptyset$ implying that $X_1^* \neq \emptyset$.

Let
$$M = M^*[X_1^* \cup Y_1^*]$$
. We have $\delta_M(x) \ge \delta^2 n/8$ for all $x \in X_1^*$.

Proposition 5.12. Let $1/t \ll \delta \ll 1/r$, 1/k and R be a k-graph on t vertices with a δt -semi-dense matching. Then R contains a $\delta^3 t/2$ -half-dense matching.

Proof. Let $M = \{v_{i,1}v_{i,2} \dots v_{i,k} : i \in [\ell]\}$ be a δt -semi-dense matching and let $V_k = \{v_{i,k} : i \in [\ell]\}$. Note that

$$\delta t \le \ell \le t/2. \tag{5.8}$$

Let H be the auxiliary bipartite graph with vertex classes $[\ell]$ and V_k such that, for each $i \in [\ell]$ and $v_{j,k} \in V_k$, we have $iv_{j,k} \in E(H)$ if and only if $v_{i,1}v_{i,2} \dots v_{i,k-1}v_{j,k} \in E(R)$. We deduce that

$$e(H) \ge \delta t \ell \stackrel{(5.8)}{\ge} 2\delta \ell^2.$$

By Lemma 5.10 with $H, \ell, (\delta/2), V_k, [\ell]$ playing the roles of G, n, δ, X, Y , respectively, H contains a matching M' such that, for all $v \in V_k \cap V(M')$, we have

$$d_{H[V(M')]}(v) \ge \delta^2 \ell/2 \stackrel{(5.8)}{\ge} \delta^3 t/2.$$
 (5.9)

Without loss of generality, $M' = \{iv_{i,k}\}_{i \in [\ell]}$. Let $M'' = \{v_{i,k}v_{i,k-1} \dots v_{i,1} : i \in [\ell]\}$. By (5.9), M'' is the required half-dense matching.

Corollary 5.13. Let $1/t \ll \varepsilon_k \ll 1/r, 1/k$. Let R be an r-edge-coloured k-graph on t vertices with $|E(R)| \ge (1-\varepsilon_k) {t \choose k}$. Then there exists a monochromatic tight component of R containing a $\left(2^{9k}(2r)^{3\cdot 2^k}\right)^{-1}$ t-half-dense matching.

Proof. Let $1/t \ll \varepsilon_k \ll \varepsilon_{k-1} \ll \cdots \ll \varepsilon_2 \ll 1/r$, 1/k and let $\delta = \left(2^{2^k+3k-5}r^{2^k-2}\right)^{-1}$. By Lemma 5.9 there exists a monochromatic tight component of R containing a δt -semi-dense matching. By Proposition 5.12, there is a $\delta^3 t/2$ -half-dense matching in R. Note that

$$\frac{\delta^3 t}{2} = \frac{t}{2^{3 \cdot 2^k + 9k - 14} r^{3 \cdot 2^k - 6}} \ge \frac{t}{2^{9k} (2r)^{3 \cdot 2^k}}$$

This completes the proof of the corollary.

We need the following lemma on bipartite graphs (Lemma 5.15) for the main result in this subsection. The lemma will enable us to find a suitable fractional matching. To prove it we need the following result by Gyárfás, Ruszinkó, Sárközy and Szemerédi [11].

Lemma 5.14 ([11, Lemma 5]). Let $1/n \ll c \leq 0.001$ and G be a directed graph on n vertices with minimum out-degree $\delta^+(G) \geq cn$. Then there are subsets $Y \subseteq X \subseteq V(G)$ such that $|Y| \geq cn/2$ and for all $x \in X$ and $y \in Y$, there are at least c^6n internally vertex-disjoint paths from x to y of length at most c^{-3} .

Lemma 5.15. Let $1/n \ll c, \mu < 1$ with $c + \mu \leq 1/8$. Let G be a bipartite graph with vertex classes $X = \{x_i : i \in [n]\}, Y = \{y_i : i \in [n]\}$ and a perfect matching $M = \{x_iy_i : i \in [n]\}$. Suppose that for all $x \in X, d(x, Y) \geq \delta n$. Then there exist disjoint subsets I^+ and I^- of [n] with $|I^+| = |I^-| = \delta n/4$ such that the following hold. Let ω be a vertex weighting of G such that we have

$$\begin{aligned}
\omega(y_i) &= 1/2 & \text{for all } i \in [n], \\
\omega(x_i) &\in [1/2 - c, 1/2] & \text{for all } i \in [n] \setminus I^+, \\
\omega(x_i) &\in [1/2, 1/2 + c] & \text{for all } i \in [n] \setminus I^- \text{ and} \end{aligned} (5.10)$$

$$\sum_{i \in I^+} (\omega(x_i) - \omega(y_i)) = \sum_{j \in I^-} (\omega(y_j) - \omega(x_j)) < \frac{\delta^9 n}{8}.$$
(5.11)

Then G has a fractional matching ω^* of weight at least $\sum_{i \in [n]} \omega(x_i) - \mu n$ such that, for each $v \in V(G)$, we have $1/2 - c \leq \omega^*(v) \leq \omega(v)$ and each non-zero weighted edge has weight at least 1/8.

Proof. We start with the following claim identifying I^+ and I^- .

Claim 5.16. There exists a subset I of [n] with $|I| = \delta n/2$ such that for all distinct $i, i' \in I$, there are at least $\delta^6 n$ internally vertex-disjoint M-augmented paths from x_i to $x_{i'}$ of the form $x_i y_{i_1} x_{i_1} \dots y_{i'} x_{i'}$.

Proof of claim. Define an auxiliary digraph H on X such that $x_i x_j \in E(H)$ is directed from x_i to x_j if $x_i y_j \in E(G)$. Note that $\delta^+(H) \geq \delta n$. By Lemma 5.14, there exists $X_2 \subseteq X_1 \subseteq X$ such that $|X_2| \geq \delta n/2$ and every $x_1 \in X_1$ has at least $\delta^6 n$ internally vertex-disjoint paths of length at most δ^{-3} to every $x_2 \in X_2$. Let $I \subseteq [n]$ with $|I| = \delta n/2$ be such that $\{x_i : i \in I\} \subseteq X_2$. Note that a path $x_{i_1} \dots x_{i_\ell}$ in H corresponds to an M-augmented path in G from x_{i_1} to x_{i_ℓ} , namely $x_{i_1} y_{i_2} x_{i_2} \dots y_{i_\ell} x_{i_\ell}$.

Let I be given by Claim 5.16. Partition I into I^+ and I^- so that $|I^+| = |I^-| = \delta n/4$. We write $X^+ = \{x_i : i \in I^+\}$ and write X^-, Y^+, Y^-, M^+ and M^- similarly. Let ω be a vertex weighting satisfying the assumptions of the lemma.

Define a fractional matching ω_0^* on G such that, for all $xy \in E(G)$ with $x \in X$ and $y \in Y$,

$$\omega_0^*(xy) = \begin{cases} \min\{\omega(x), \omega(y)\} = 1/2 & \text{if } xy \in M \setminus M^-, \\ \min\{\omega(x), \omega(y)\} = \omega(x) & \text{if } xy \in M^-, \\ 0 & \text{otherwise.} \end{cases}$$

For a fractional matching ω^* on G, let

$$||\omega^* - \omega_0^*|| = \sum_{e \in E(G)} |\omega^*(e) - \omega_0^*(e)|,$$

which denotes the sum of the edge-weight differences between ω^* and ω_0^* . Let ω^* be a fractional matching on G such that

- (a) $1/2 c \le \omega^*(v) \le \omega(v)$ for all $v \in V(G)$;
- (b) $\omega^*(x_i y_i) \ge 1/8$ for all $i \in [\ell]$;
- (c) $||\omega^* \omega_0^*|| \le 2\delta^{-3}(|\omega^*| |\omega_0^*|);$
- (d) $\omega^*(v) = \omega(v)$ for all $v \in V(G) \setminus (X^+ \cup Y^-)$.

Such an ω^* exists by taking $\omega^* = \omega_0^*$. We further assume that $|\omega^*|$ is maximal. We may also assume that $|\omega^*| < \sum_{i \in [n]} \omega(x_i) - \mu n$ or else we are done. By (d), there exist $x^+ \in X^+$ and $y^- \in Y^-$ such that

$$\omega(x^{+}) - \omega^{*}(x^{+}), \ \omega(y^{-}) - \omega^{*}(y^{-}) \ge \mu.$$
(5.12)

We aim to add weight to x^+ and y^- . Let $x^- = N_M(y^-)$. Let $E_0 = \{e \in E(G) : |\omega^*(e) - \omega_0^*(e)| \ge 1/4\}$. Note that

$$|E_0| \le \frac{||\omega^* - \omega_0^*||}{1/4} \stackrel{(c)}{\le} 8\delta^{-3} \left(|\omega^*| - |\omega_0^*|\right) \stackrel{(d)}{\le} 8\delta^{-3} \left(\sum_{i \in I^+} \omega(x_i) - \omega(y_i)\right) \stackrel{(5.11)}{<} \delta^6 n$$

Thus, by the definitions of X^+ and X^- , there exists an *M*-augmented path *P* in *G* from x^+ to x^- of length at most δ^{-3} with $E(P) \cap E_0 = \emptyset$. Without loss of generality, let $P = x_1 y_2 x_2 y_3 \dots x_{\ell-1} y_{\ell} x_{\ell}$ where $x_1 = x^+, x_{\ell} = x^-, y_{\ell} = y^-$ and $\ell \leq \delta^{-3}$. Define the fractional matching $\omega_1^* : E(G) \to [0, 1]$ such that, for any edge $e \in E(G)$, we have

$$\omega_{1}^{*}(e) = \begin{cases} \omega^{*}(e) - \mu & \text{if } e = y_{i}x_{i} \text{ with } i \in [\ell - 1] \setminus \{1\}, \\ \omega^{*}(e) + \mu & \text{if } e = x_{i}y_{i+1} \text{ with } i \in [\ell - 1], \\ \omega^{*}(e) & \text{otherwise.} \end{cases}$$
(5.13)

Note that

$$|\omega_1^*| = |\omega^*| + \mu \tag{5.14}$$

and for $v \in V(G)$,

$$\omega_1^*(v) = \begin{cases} \omega^*(v) + \mu & \text{if } v \in \{x^+, y^-\}, \\ \omega^*(v) & \text{otherwise.} \end{cases}$$

Therefore by (5.12), we have that ω_1^* satisfies (a) and (d).

To see ω_1^* satisfies (b), if $e \in E(G) \setminus (E(P) \setminus \{x_\ell y_\ell\})$, then $\omega_1^*(e) = \omega^*(e)$. If $e \in E(P) \setminus \{x_\ell y_\ell\} \subseteq E(G) \setminus E_0$, then $|\omega_1^*(e) - \omega_0^*(e)| = \mu$ and so

$$\omega_1^*(e) \begin{cases} \ge \omega_0^*(e) - 1/4 - \mu \ge 1/8 & \text{if } e \in E(M) \\ \le \omega_0^*(e) + 1/4 + \mu \le 1/2 & \text{if } e \notin E(M). \end{cases}$$

Thus, ω_1^* indeed satisfies (b) and moreover is a fractional matching on G.

To verify that ω_1^* satisfies (c), recall that P has $2\ell - 1 \leq 2\delta^{-3}$ edges. Therefore, we deduce that

$$\begin{aligned} ||\omega_1^* - \omega_0^*|| &\leq ||\omega_1^* - \omega^*|| + ||\omega^* - \omega_0^*|| \leq \mu(2\ell - 1) + 2\delta^{-3}(|\omega^*| - |\omega_0^*|) \\ &\leq 2\delta^{-3}\mu + 2\delta^{-3}(|\omega^*| - |\omega_0^*|) \stackrel{(5.14)}{=} 2\delta^{-3}(|\omega_1^*| - |\omega_0^*|). \end{aligned}$$

Thus, ω_1^* satisfies (a) to (d) and has larger weight than ω^* , a contradiction.

We now prove Lemma 5.4 assuming Lemma 5.3. Firstly, we find a monochromatic tightly connected half-dense matching in the reduced k-graph and reserve some unbalanced vertex clusters corresponding to the vertices of the matching, as the vertex set U^* . We then use Lemma 5.3 to cover B^* with a set C of few monochromatic tight cycles using some vertices from U^* . Finally, we use the half-dense property of the matching and Lemma 5.15 to find one long tight cycle covering almost all the vertices in $U^* \setminus V(C)$.

Proof of Lemma 5.4. Let
$$d_k = 1/r$$
, $\delta = \left(2^{9k}(2r)^{3 \cdot 2^k}\right)^{-1}$,
 $1/n \ll 1/t_1 \ll 1/t_0 \ll \varepsilon_k \ll \varepsilon', \mu \ll \psi \ll 1/r, 1/k,$
 $1/n \ll 1/r', \tilde{\varepsilon} \ll \varepsilon_k, d_2, \dots, d_{k-1}.$

Step 1: Defining U^* .

Let H' be an induced subgraph of H such that $|V(H')| \equiv 0 \pmod{t_1!}$ and $|V(H \setminus H')| < t_1!$. By Lemma 4.3 with H' playing the role of H, there exists a $(\cdot, \cdot, \tilde{\varepsilon}, \varepsilon_k, r')$ -regular slice $\tilde{\mathcal{J}}$ with partition \mathcal{P} of V(H'). Let m = |V| for $V \in \mathcal{P}$, $t = |\mathcal{P}|$ and $t_0 \leq t \leq t_1$. Note that

$$n \ge mt \ge n - t_1! \ge n/2.$$
 (5.15)

Let $\mathcal{R} = \mathcal{R}_{d_k}^{\widetilde{\mathcal{J}}}(H')$. Note that $|E(\mathcal{R})| \geq (1 - r\varepsilon_k) {t \choose k}$ by Lemma 4.4. By Corollary 5.13 with $\mathcal{R}, r\varepsilon_k$ playing the roles of R, ε , respectively, there exists a monochromatic tight component T of \mathcal{R} and a δt -half dense matching M of size ℓ , where

$$\delta t \le \ell \le t/k \le t/2. \tag{5.16}$$

Let $M = \{x_i y_{i,2} \dots y_{i,k} : i \in [\ell]\}$ and $X = \{x_i : i \in [\ell]\}$ be such that, for all $x \in X$, we have $|\{i \in [\ell] : xy_{i,2} \dots y_{i,k} \in E(T)\}| \ge \delta t.$

Define an auxiliary bipartite graph G on X and $Y = \{y_1, \ldots, y_\ell\}$ such that $x_i y_j \in E(G)$ if and only if $x_i y_{j,2} \ldots y_{j,k} \in E(T)$. Note that M corresponds to a half-dense matching $M' = \{x_i y_i : i \in [\ell]\}$ in G and for each $x \in X$,

$$d_G(x,Y) \ge \delta t \stackrel{(5.16)}{\ge} 2\delta \ell.$$

By Lemma 5.15 with $\ell, 2\delta$ playing the roles of n, δ , respectively, there exist disjoint $I^+, I^- \subseteq [\ell]$ such that

$$|I^+| = |I^-| = \delta\ell/2 \stackrel{(5.16)}{\geq} \delta^2 t/2 \stackrel{(5.15)}{\geq} \delta^2 (n/2m) - \delta^2 t_1!/(2m) \ge \delta^2 n/4m.$$

Define $X^+ = \{x_i \in X : i \in I^+\}$ and define X^-, Y^+, Y^- similarly. Let

$$\alpha = 2^5 r k (2r)^{2^{k+1}}, \gamma = 4\varepsilon^* \alpha n / (m|I^+|) \text{ and } \gamma_a = (k-1)\gamma / \alpha = 4\varepsilon^* (k-1)n / (m|I^+|).$$
(5.17)

Note that Lemma 5.15 also implies that if ω is a vertex-weighting satisfying (5.10) and (5.11) with $\gamma - \gamma_a, \ell$ playing the roles of c, n, respectively, then G has a fractional matching ω^* with $|\omega^*| \geq \sum_{i \in [\ell]} \omega(x_i) - \mu \ell$, where each vertex v has weight between $1/2 - \gamma + \gamma_a$ and $\omega(v)$ and furthermore each non-zero weighted edge has weight at least 1/8.

For each $v \in V(\mathcal{R})$, recall that V_v is the cluster of \mathcal{P} corresponding to v. We pick disjoint $U_v, A_v \subseteq V_v$ such that

$$|A_v| = \begin{cases} \gamma m & \text{if } v \in X^+, \\ 0 & \text{otherwise,} \end{cases}$$
$$|U_v| = \begin{cases} \left(\frac{1}{2} - \gamma + \gamma_a\right)m & \text{if } v \in X^-, \\ \frac{m}{2} & \text{if } v \in V(M') \setminus X^-, \\ \varepsilon'm & \text{otherwise.} \end{cases}$$

Let $A = \bigcup_{v \in V(\mathcal{R})} A_v$, $U = \bigcup_{v \in V(\mathcal{R})} U_v$ and $U^* = U \cup A$. Therefore, we have $|A| = |I^+|\gamma m = 4\alpha \varepsilon^* n$ by (5.17) and

$$\begin{aligned} |U^*| &= |U| + |A| = km\ell/2 - (\gamma - \gamma_a)m|X^-| + \varepsilon'm(t - k\ell) + \gamma m|X^+| \\ &= km\ell/2 + \gamma_a m|X^-| + \varepsilon'm(t - k\ell) = km\ell/2 + 4(k - 1)\varepsilon^*n + \varepsilon'm(t - k\ell) \\ &\stackrel{(5.16)}{\leq} mt/2 + 4(k - 1)\varepsilon^*n + \varepsilon'm(t - k\ell) \stackrel{(5.15)}{\leq} 3n/4. \end{aligned}$$

Step 2: Verifying properties of U^* . Let $B^* \subseteq V(H) \setminus U^*$ be such that $|B^*| = 4\varepsilon^* n$. **Step 2A:** Covering B^* .

By Lemma 5.3 with A, B^* playing the roles of X, Z, respectively, there exists a set C of vertex-disjoint monochromatic tight cycles covering B^* with

$$|\mathcal{C}| \le 2^{18} r \delta_0^{-5}$$
 and $|V(\mathcal{C}) \cap A| = 4(k-1)\varepsilon^* n = \gamma_a m |I^+|.$

Step 2B: Covering $U^* \setminus V(\mathcal{C})$.

Let $A' = A \setminus V(\mathcal{C})$, so $|A'| = (\gamma - \gamma_a)|I^+|m$. For each $x \in V(\mathcal{R})$, let $A'_x = A' \cap V_x$, so $\bigcup_{x \in X^+} A'_x = A'$. Note that

$$km\ell/2 \le |U \cup A'| \le km\ell/2 + \varepsilon'mt.$$
(5.18)

We define a vertex-weighting $\omega: V(M') \to [0,1]$ to be such that for $v \in V(M')$,

$$\omega(v) = \frac{|V_v \cap (U \cup A')|}{m} = \frac{|U_v| + |A'_v|}{m} \begin{cases} \in [\frac{1}{2}, \frac{1}{2} + \gamma] & \text{if } v \in X^+, \\ = \frac{1}{2} - \gamma + \gamma_a & \text{if } v \in X^-, \\ = \frac{1}{2} & \text{if } v \in V(M') \setminus (X^+ \cup X^-). \end{cases}$$

Firstly, note that

$$\sum_{i \in I^+} (\omega(x_i) - \omega(y_i)) = \sum_{i \in I^+} \left(\left(\frac{1}{2} + \frac{|A'_{x_i}|}{m} \right) - \frac{1}{2} \right) = \frac{|A'|}{m}$$
$$= (\gamma - \gamma_a)|I^+| = \sum_{i \in I^-} (\omega(y_i) - \omega(x_i)).$$

Also

$$\sum_{i \in I^+} (\omega(x_i) - \omega(y_i)) = (\gamma - \gamma_a) |I^+| \le \gamma |I^+|$$

$$\stackrel{(5.17)}{=} 4\varepsilon^* \alpha n/m \stackrel{(5.15)}{\le} 8\varepsilon^* \alpha t \stackrel{(5.16)}{\le} 8\varepsilon^* \alpha \ell/\delta < (2\delta)^9 \ell/8.$$

Therefore, all the conditions for Lemma 5.15 (with 2δ playing the role of δ) are satisfied. Furthermore, $\sum_{i \in [\ell]} \omega(x_i) = \sum_{i \in [\ell]} \omega(y_i)$. We deduce that G has a fractional matching ω^* with

$$|\omega^*| \ge \sum_{i \in [\ell]} \omega(x_i) - \mu \ell = \sum_{i \in [\ell]} \omega(y_i) - \mu \ell = \left(\frac{1}{2} - \mu\right) \ell, \tag{5.19}$$

where each vertex v has weight at least $1/2 - \gamma + \gamma_a$ and at most $\omega(v)$ and each non-zero weighted edge has weight at least 1/8. Define a corresponding fractional matching ω_H^* on H such that for an edge $e \in E(H)$, $\omega_H^*(e) = \omega^*(x_i y_j)$ if $e = x_i y_{j,2} \dots y_{j,k}$ and $\omega_H^*(e) = 0$ otherwise. Note that $|\omega_H^*| = |\omega^*|$. For any vertex $u \notin V(M)$ we have $\omega_H^*(u) = 0$, so $|V_u \cap (U \cup A')| = \varepsilon' m = ((1 - 3\varepsilon')\omega_H^*(u) + \varepsilon')m$. For each $u \in V(M)$, we have $\omega_H^*(u) \ge 1/3$ and so

$$|V_u \cap (U \cup A')| = |U_u \cup A'_u| = \omega(u)m \ge \omega_H^*(u)m \ge ((1 - 3\varepsilon')\omega_H^*(u) + \varepsilon')m.$$

By Lemma 4.5, $H[A' \cup U]$ contains a tight cycle C of length ℓ_0 with

$$\ell_0 \ge (1-\psi)k|\omega_H^*|m-k \ge (1-\psi)(1/2-\mu)k\ell m-k \ge (1-\psi)km\ell/2-\mu km\ell.$$

Recall that $\mu, \varepsilon', 1/t \ll \psi$ and $1/m \ll 1/k$. Hence,

$$|(A' \cup U) \setminus V(C)| \stackrel{(5.18)}{\leq} \psi km\ell/2 + \mu km\ell + \varepsilon'mt \leq \psi km\ell \stackrel{(5.16)}{\leq} \psi mt \stackrel{(5.15)}{\leq} \psi mt$$

as required.

6 Edge-coloured Multigraphs

Let *H* be a *k*-partite *k*-graph with vertex classes X_1, \ldots, X_k with $X_i = \{x_{i,j} : j \in [n]\}$. We denote the 2-blowup of a *k*-edge by $K_k^{(k)}(2)$. Note that $K_k^{(k)}(2)$ is the complete *k*-partite *k*-graph with each vertex class of order 2. We say that an edge-coloured multigraph *G* respects *H* if the following hold:

- the colour set of G is a subset of X_k ;
- $V(G) \subseteq \{v_i : i \in [n]\};$
- if $\phi(v_i v_{i'}) = x_{k,t}$ for some $t \in [n]$, then $\{x_{i,j} x_{i',j} : j \in [k-1]\}$ forms a $K_{k-1}^{(k-1)}(2)$ in $H(x_{k,t})$.

There is a natural correspondence between a tight cycle in H and a rainbow cycle in G.

Fact 6.1. Let H be a k-partite k-graph with vertex sets X_1, \ldots, X_k such that for $i \in [k-1]$, $X_i = \{x_{i,j} : j \in [n]\}$. Let G be an edge-coloured multigraph respecting H. Suppose that G contains a rainbow cycle $v_1v_2 \cdots v_\ell$ with $\phi(v_iv_{i+1}) = x_{k,i}$ for $i \in [\ell - 1]$ and $\phi(v_\ell v_1) = x_{1,\ell}$. Then H contains a tight cycle with vertex set $\{x_{i,j} : i \in [k], j \in [\ell]\}$.

Let G be an edge-coloured multigraph. Recall that $\phi(G)$ is the set of colours in G and for a colour $c \in \phi(G)$, G_c is the subgraph with edges only of the colour c and no isolated vertices. The monochromatic colour degree of G to be $\delta_{\text{mon}}(G) = \min_{c \in \phi(G)} \delta(G_c)$. A path system is a collection of vertex-disjoint paths. A rainbow path system is a path system such that its union (viewed as a graph) is rainbow. A (rainbow) cycle system is defined similarly. The main goal of this section is to reduce Lemma 5.3 into the following lemma, which will be proved in the next section.

Lemma 6.2. Let $1/n \ll \delta_0 \leq 1/8$. Suppose G is an edge-coloured multigraph on n vertices with $\delta_{mon}(G) \geq \delta_0 n$ and $|\phi(G)| \leq \delta_0 n/16$. Then there exists a rainbow cycle system C with $\phi(C) = \phi(G)$ and $|\mathcal{C}| \leq 2^{18} \delta_0^{-5}$.

To reduce Lemma 5.3 to Lemma 6.2, we need the help of the following lemma, which will be proved in the next subsection.

Lemma 6.3. Let H be an r-edge-coloured $K_n^{(k)}$ with disjoint vertex sets X and Z with $|X| \ge (k-1)|Z|$. Then for each $i \in [r]$, there exist an edge-coloured multigraph G^i and a subgraph H^i of H such that

- (A1) H^i is a monochromatic k-partite k-graph on vertex classes X_1^i, \ldots, X_k^i of colour i;
- (A2) $\phi(G^i) = X_k^i, Z = \bigcup_{i \in [r]} X_k^i \text{ and } \bigcup_{j \in [k-1]} X_j^i \subseteq X;$
- (A3) if H^i is not empty, then $|V(G^i)| = \left(1 \frac{1}{4kr^k}\right) \frac{|X|}{r(k-1)}$, $\delta_{mon}(G^i) \ge (2r)^{-2^k} |V(G^i)|/8$ and G^i respects H^i ;
- (A4) for distinct $i, j \in [r]$, we have $V(H^i) \cap V(H^j) = \emptyset$.

We now prove Lemma 5.3 assuming Lemmas 6.2 and 6.3.

Proof of Lemma 5.3. By Lemma 6.3, for each $i \in [r]$, there exist an edge-coloured multigraph G^i and a k-partite k-graph H^i such that G^i respects H^i . Fix $i \in [r]$. Observe that if H^i is not empty, then $\delta_{\text{mon}}(G^i) \geq \delta_0 |V(G^i)|$ by (A3). Moreover

$$\left|\phi(G^{i})\right| \stackrel{(\mathbf{A2})}{\leq} |Z| \leq \frac{|X|}{\alpha} \leq \frac{\delta_{0}|X|}{32r(k-1)} \leq \frac{\delta_{0}}{16} \left(1 - \frac{1}{4kr^{k}}\right) \frac{|X|}{r(k-1)} \stackrel{(\mathbf{A3})}{=} \frac{\delta_{0}|V(G^{i})|}{16}.$$

By Lemma 6.2 with G^i playing the role of G, we deduce that G^i contains a rainbow cycle system \mathcal{C}^i with $\phi(\mathcal{C}^i) = \phi(G^i) = X_k^i$ and $|\mathcal{C}^i| \leq 2^{18} \delta_0^{-5}$. By Fact 6.1 and (A1), \mathcal{C}^i corresponds to a set $\hat{\mathcal{C}}^i$ of vertex-disjoint monochromatic tight cycles in H^i covering X_k^i . Therefore by (A4), $\bigcup_{i \in [r]} \hat{\mathcal{C}}^i$ is a set of vertex-disjoint monochromatic tight cycles covering $\bigcup_{i \in [r]} X_k^i = Z$. Note that $\left|\bigcup_{i \in [r]} \hat{\mathcal{C}}^i\right| \leq 2^{18} r \delta_0^{-5}$. This completes the proof of the lemma.

6.1 Proof of Lemma 6.3

We start with the following result, which allows us to partition $X \cup Z$ into the vertex-disjoint kpartite k-graphs H^i for $i \in [r]$. Let H be an r-edge-coloured k-graph. For vertex sets X_1, \ldots, X_k , let $e_j(X_1, \ldots, X_k)$ be the number of $X_1 \ldots X_k$ -edges of colour j in H.

Lemma 6.4. Let $1/N \ll 1/k, 1/r \leq 1/2$. Let H be an r-edge-coloured complete k-partite k-graph with vertex classes X_1, X_2, \ldots, X_k where for all $i \in [k-1], |X_i| = N$ and $|X_k| \leq N$. Then there exists a partition of X_1, \ldots, X_k into $\{X_1^j : j \in [r] \cup \{0\}\}, \ldots, \{X_{k-1}^j : j \in [r] \cup \{0\}\}, \{X_k^j : j \in [r]\}$ such that, for each $i \in [k-1], j \in [r]$ and $x \in X_k^j$, we have

$$|X_i^j| = \left(1 - \frac{1}{4kr^k}\right) \left(\frac{N}{r}\right) \text{ and } e_j\left(X_1^j, \dots, X_{k-1}^j, x\right) \ge \frac{\left|X_1^j\right| \cdots \left|X_{k-1}^j\right|}{2r}.$$

Proof. Partition X_k into X_k^1, \ldots, X_k^r such that for all $j \in [r]$ and $x_k \in X_k^j$,

$$e_j(X_1, \dots, X_{k-1}, x_k) \ge \frac{|X_1| \dots |X_{k-1}|}{r}.$$

Let $\varepsilon = (4kr^k)^{-1}$.

Claim 6.5. There exists a partition of X_i for each $i \in [k-1]$ into X_i^1, \ldots, X_i^r such that for all $j \in [r]$, $\left| \left| X_i^j \right| - \frac{N}{r} \right| \le \varepsilon \frac{N}{r}$ and

$$\left| e_j \left(X_1^j, \dots, X_i^j, X_{i+1}, \dots, X_{k-1}, x_k \right) - \frac{e_j \left(X_1^j, \dots, X_{i-1}^j, X_i, \dots, X_k \right)}{r} \right| < \varepsilon N^{k-1}.$$

Proof of claim. We proceed by induction on i. We only prove the base case i = 1 as the rest can be proved analogously.

Let $X_1 = X_1^1 \cup \cdots \cup X_1^r$ be a random partition of X_1 , where for all $x_1 \in X_1$ and $i \in [r]$, x_1 is assigned to X_1^j independently with probability 1/r. Consider $x_i \in X_i$ for $i \in [2, k]$ and $j \in [r]$. Note that $\mathbb{E}\left(e_j(X_1^j, x_2, \ldots, x_k)\right) = \frac{e_j(X_1, x_2, \ldots, x_k)}{r}$. If $e_j(X_1, x_2, \ldots, x_k) \leq \varepsilon N$, then clearly $e_j(X_1^i, x_2, \ldots, x_k) \leq \varepsilon N$. If $e_j(X_1^j, x_2, \ldots, x_k) \geq \varepsilon N$, then by Lemma 3.1, we have

$$\mathbb{P}\left(\left|e_j(X_1^j, x_2, \dots, x_k) - \frac{e_j(X_1, x_2, \dots, x_k)}{r}\right| > \varepsilon \frac{e_j(X_1, x_2, \dots, x_k)}{r}\right) \le 2e^{-\frac{\varepsilon^3 N}{3r}}.$$

Note that $\mathbb{E}|X_1^j| = N/r$. By Lemma 3.1, we have

$$\mathbb{P}\left(\left||X_1^j| - \frac{N}{r}\right| > \varepsilon \frac{N}{r}\right) < 2e^{-\frac{\varepsilon^2 N}{3r}}.$$

Hence by the union bound, with positive probability for all $j \in [r]$, $i \in [2, k]$ and $x_i \in X_i$, we have $\left| |X_1^j| - \frac{N}{r} \right| < \varepsilon \frac{N}{r}$ and $\left| e_j(X_1^j, x_2, \dots, x_k) - \frac{e_j(X_1, x_2, \dots, x_k)}{r} \right| \le \varepsilon N$. Fix such a partition. We have

$$\left| e_j(X_1^j, X_2, \dots, X_{k-1}, x_k) - \frac{e_j(X_1, X_2, \dots, X_{k-1}, x_k)}{r} \right|$$
$$= \left| \sum_{\substack{x_i \in X_i, \\ i \in [2, k-1]}} \left(e_j(X_1^j, x_2, \dots, x_k) - \frac{e_j(X_1, x_2, \dots, x_{k-1}, x_k)}{r} \right) \right| \le \sum_{\substack{x_i \in X_i, \\ i \in [2, k-1]}} \varepsilon N \le \varepsilon N^{k-1}.$$

Fix a partition of X_1, \ldots, X_{k-1} given by Claim 6.5. For each $i \in [k-1]$ and $j \in [r]$, remove additional vertices from X_i^j if necessary so that, for all $i \in [k-1]$ and $j \in [r]$, $\left|X_i^j\right| = (1-\varepsilon)\frac{N}{r}$. Let $X_i^0 = X_i \setminus \bigcup_{j \in [r]} X_i^j$ for each $i \in [k-1]$. Each X_i^j has at most $2\varepsilon N/r$ vertices removed. Thus for all $j \in [r]$ and $x_k \in X_k^j$, we have

$$e_{j}(X_{1}^{j}, \dots, X_{k-1}^{j}, x_{k}) \geq \frac{e_{j}(X_{1}, \dots, X_{k-1}, x_{k})}{r^{k-1}} - k\varepsilon N^{k-1} - N^{k-2} \left(\frac{2\varepsilon N}{r}\right)$$
$$\geq \frac{N^{k-1}}{r^{k}} - 2k\varepsilon N^{k-1} \geq \frac{N^{k-1}}{2r^{k}} \geq \frac{|X_{1}^{j}| \dots |X_{k-1}^{j}|}{2r}.$$

We need a lower bound on the number of $K_k^{(k)}(2)$ in a dense k-partite k-graph H.

Lemma 6.6. Let $1/n \ll 1/k, d \leq 1$. Let H be a k-partite k-graph with n vertices in each vertex class and $e(H) \geq dn^k$. Then there are at least $d^{2^k}n^{2k}/2$ many $K_k^{(k)}(2)$ in H.

Proof. Let the vertex classes of H be X_1, \ldots, X_k . For vertices $x_1, \ldots, x_k \in V(H)$, we write $\mathbb{1}(x_1 \ldots x_k)$ for the indicator function $\mathbb{1}(x_1 \ldots x_k \in E(H))$. For $j \in [k] \cup \{0\}$, let

$$f(j) = \sum_{\substack{x_i \in X_i, \\ i \in [k-j]}} \sum_{\substack{x_s, y_s \in X_s, \\ s \in [k-j+1,k]}} \prod_{\substack{v_t \in \{x_t, y_t\}, \\ t \in [k-j+1,k]}} \mathbb{1}(x_1 \dots x_{k-j} v_{k-j+1} \dots v_k).$$

Note that

$$dn^{k} \leq |E(H)| = \sum_{x_{i} \in X_{i}, i \in [k]} \mathbb{1}(x_{1} \dots x_{k}) = f(0).$$
(6.1)

Claim 6.7. For $t \in [k]$, we have $n^{2^{t}k-k-t}f(t) \ge f(0)^{2^{t}}$.

Proof of claim. For any $j \in [k-1] \cup \{0\}$, we apply the Cauchy-Schwarz inequality to obtain

$$f(j)^{2} = \left(\sum_{\substack{x_{i} \in X_{i}, \\ i \in [k-j-1]}} \sum_{\substack{x_{s}, y_{s} \in X_{s}, \\ i \in [k-j+1,k]}} \left(\sum_{\substack{x_{k-j} \in X_{k-j} \\ t \in [k-j+1,k]}} \prod_{\substack{v_{t} \in \{x_{t}, y_{t}\}, \\ i \in [k-j-1]}} \mathbb{1}(x_{1} \dots x_{k-j} v_{k-j+1} \dots v_{k})\right)\right)^{2}$$

$$\leq n^{k+j-1} \sum_{\substack{x_{i} \in X_{i}, \\ i \in [k-j-1]}} \sum_{\substack{x_{s}, y_{s} \in X_{s}, \\ s \in [k-j+1,k]}} \left(\sum_{\substack{x_{k-j} \in X_{k-j} \\ t \in [k-j+1,k]}} \prod_{\substack{v_{t} \in \{x_{t}, y_{t}\}, \\ t \in [k-j+1]}} \mathbb{1}(x_{1} \dots x_{k-j} v_{k-j+1} \dots v_{k})\right)^{2}$$

$$= n^{k+j-1} \sum_{\substack{x_{i} \in X_{i}, \\ i \in [k-j,k]}} \sum_{\substack{x_{s}, y_{s} \in X_{s}, \\ t \in [k-j,k]}} \prod_{\substack{v_{t} \in \{x_{t}, y_{t}\}, \\ t \in [k-j,k]}} \mathbb{1}(x_{1} \dots x_{k-j-1} v_{k-j} \dots v_{k})$$

$$= n^{k+j-1} f(j+1). \tag{6.2}$$

We now prove the claim by induction on t. Note that $f(0)^2 \leq n^{k-1}f(1)$ by (6.2). Suppose that t > 1. Thus

$$n^{2^{t}k-k-t}f(t) \stackrel{(6.2)}{\geq} n^{2^{t}k-k-t}n^{-(k+t-2)}f(t-1)^{2} = \left(n^{2^{t-1}k-k-(t+1)}f(t-1)\right)^{2} \ge f(0)^{2^{t}}, \tag{6.3}$$

where the last inequality is due to our induction hypotheses.

The number of $K_k^{(k)}(2)$ in H is

$$\sum_{\substack{x_i, y_i \in X_i, \\ x_i \neq y_i, \\ i \in [k]}} \prod_{\substack{v_t \in \{x_t, y_t\}, \\ t \in [k]}} \mathbb{1}(v_1 \dots v_k) \ge f(k) - kn^{2k-1} \stackrel{\text{Claim}}{\ge} \frac{6.7}{n^{2k-k2^k}} n^{2k-k2^k} f(0)^{2^k} - kn^{2k-1}$$

We now show that there exists a (non-empty) subgraph H_0 of H such that each edge is in many $K_k^{(k)}(2)$.

Lemma 6.8. Let $1/n \ll d \leq 1$, $k \geq 2$ and $\gamma = d^{2^k}/2$. Let H be a k-partite k-graph with n vertices in each vertex class and $e(H) \geq dn^k$. Then there exists a subgraph H_0 of H such that $|E(H_0)| \geq 2^{k-1}\gamma n^k$ and every edge of H_0 is contained in at least $\gamma n^k/2$ many $K_k^{(k)}(2)$ in H_0 .

Proof. Define the auxiliary 2k-graph \mathcal{A} to be such that $V(\mathcal{A}) = V(H)$ and each $K_k^{(k)}(2)$ of H is an edge of \mathcal{A} . By Lemma 6.6, $|E(\mathcal{A})| \geq \gamma n^{2k}$. If there exists $S \in \binom{V(\mathcal{A})}{k}$ such that $|S \cap V_i| = 1$ for $i \in [k]$ and $d_{\mathcal{A}}(S) < (\gamma/2)n^k$, then we delete all edges in \mathcal{A} containing S. Repeat this process and call the resulting 2k-graph \mathcal{A}' . Note that S can be chosen in at most n^k ways, so

$$|E(\mathcal{A}')| \ge \gamma n^{2k} - n^k \left(\frac{\gamma n^k}{2}\right) = \frac{\gamma n^{2k}}{2}.$$

Let H_0 be the subgraph of H induced by $e \in E(H)$ with $d_{\mathcal{A}'}(e) \geq 1$. For each $e \in E(H_0)$, we have $d_{\mathcal{A}'}(e) \geq \gamma n^k/2$ and therefore e is contained in at least $\gamma n^k/2$ many $K_k^{(k)}(2)$ in H_0 . Since each edge can be contained in at most n^k many $K_k^{(k)}(2)$ in H_0 ,

$$|E(H_0)| \ge \frac{2^k |E(\mathcal{A}')|}{n^k} = 2^{k-1} \gamma n^k$$

as required.

Let S_n denote the set of permutations on [n]. The following lemma shows that if the vertices within each class of a k-partite k-graph are permuted randomly, then the number of 'horizontal' edges is concentrated around its expectation.

Lemma 6.9. Let $1/n \ll 1/k \leq 1/2$ and H be a k-partite k-graph with vertex classes X_1, \ldots, X_k each of size n. Suppose $\sigma_1, \ldots, \sigma_{k-1} \in S_n$ are chosen independently and uniformly at random. Let $H_{\sigma_1,\ldots,\sigma_{k-1}} = \{x_{1,\sigma_1(i)} \cdots x_{k-1,\sigma_{k-1}(i)} x_{k,i} : i \in [n]\} \cap E(H)$. Then

$$\mathbb{P}\left(\left|\mathbb{E}|H_{\sigma_1,\dots,\sigma_{k-1}}| - \frac{|E(H)|}{n^{k-1}}\right| > \varepsilon n\right) \le 2e^{-\frac{\varepsilon^2 n}{32}}.$$

Proof. We consider the martingale obtained by exposing each position of $\sigma_1, \ldots, \sigma_{k-1}$ in turns as follows. For $i \in [k-1]$ and $j \in [n]$, let $Z_{i,j}$ be the j^{th} exposed position of σ_i . Let

$$B_{0} = \mathbb{E} \left| H_{\sigma_{1},\dots,\sigma_{k-1}} \right| = \sum_{x_{1,i_{1}}\dots,x_{k-1,i_{k-1}}x_{k,i}\in E(H)} \mathbb{P}(i_{1} = \sigma_{1}(i),\dots,i_{k-1} = \sigma_{k-1}(i))$$
$$= \sum_{x_{1,i_{1}}\dots,x_{k-1,i_{k-1}}x_{k,i}\in E(H)} \prod_{j\in[k-1]} \mathbb{P}(i_{j} = \sigma_{j}(i)) = \frac{|E(H)|}{n^{k-1}}.$$

Define

$$B_{(i-1)n+j} = \mathbb{E}\left(\left|H_{\sigma_1,\dots,\sigma_{k-1}}\right| | Z_{1,1},\dots,Z_{1,n}, Z_{2,1},\dots,Z_{2,n},\dots,Z_{i,1},\dots,Z_{i,j}\right).$$

Note that $B_0, \ldots, B_{(k-1)n}$ forms a martingale.

Claim 6.10. For $t \in [(k-1)n-1]$, we have $|B_t - B_{t+1}| \le 4$.

Proof of claim. For simplicity, we only consider the case when $t \le n-1$ (and the other cases are proved similarly). Thus

$$B_t = \mathbb{E}(|H_{\sigma_1,\dots,\sigma_{k-1}}||Z_{1,1},\dots,Z_{1,t}) \text{ and } B_{t+1} = \mathbb{E}(|H_{\sigma_1,\dots,\sigma_{k-1}}||Z_{1,1},\dots,Z_{1,t+1}))$$

If t = n - 1, then $B_t = B_{t+1}$. Thus we may assume $t \le n - 2$. For $t_1, t_2 \in [n]$, define $\pi_{t_1,t_2} : S_n \to S_n$ to be such that for all $\sigma' \in S_n$ and $s \in [n]$,

$$\pi_{t_1,t_2}(\sigma')(s) = \begin{cases} \sigma'(t_2) & \text{if } s = t_1, \\ \sigma'(t_1) & \text{if } s = t_2, \\ \sigma'(s) & \text{otherwise.} \end{cases}$$

Equivalently $\pi_{t_1,t_2}(\sigma')$ swaps the t_1^{th} and t_2^{th} position of σ' . For all $\sigma'_1,\ldots,\sigma'_{k-1}\in S_n$,

$$\left| H_{\sigma_{1}',\ldots,\sigma_{k-1}'} \Delta H_{\pi_{t_{1},t_{2}}(\sigma_{1}')\sigma_{2}',\ldots,\sigma_{k-1}'} \right| \leq \left| \left\{ x_{1,\sigma_{1}'(i)} \ldots x_{k-1,\sigma_{k-1}'(i)} x_{k,i}, \ x_{1,\pi_{t_{1},t_{2}}(\sigma_{1}'(i))} \ldots x_{k-1,\pi_{t_{1},t_{2}}(\sigma_{k-1}'(i))} x_{k,i} : i \in \{t_{1},t_{2}\} \right\} \right| \leq 4.$$
(6.4)

Let $A = \{\sigma \in S_n : \sigma(j) = Z_{1,j} \text{ for } j \in [t]\}$. Given $Z_{1,1}, \ldots, Z_{1,t}$, the probability space is now reduced to $A \times (S_n)^{k-2}$. Pick $j_0 \in [n] \setminus \{Z_{1,1} \ldots Z_{1,t}\}$. Let $A' = \{\sigma \in A : \sigma(t+1) = j_0\}$. Note that A can be partitioned into $\{\pi_{t,j}(A') : j \in [t+1,n]\}$. Hence the probability space can be partitioned into $\{\pi_{t,j}(A') \times (S_n)^{k-2} : j \in [t+1,n]\}$. Hence the claim follows by (6.4).

By Azuma's inequality (Lemma 3.3), we have

$$\mathbb{P}\left(\left|\mathbb{E}\left|H_{\sigma_{1},\ldots,\sigma_{k-1}}\right| - \frac{|E(H)|}{n^{k-1}}\right| > \varepsilon n\right) \le 2e^{-\frac{\varepsilon^{2}n}{32}}$$

as required.

We are now ready to prove Lemma 6.3.

Proof of Lemma 6.3. Let

$$n = \left(1 - \frac{1}{4kr^k}\right) \left(\frac{|X|}{r(k-1)}\right).$$

Partition X into equally sized subsets X_1, \ldots, X_{k-1} so that for $j \in [k-1]$, we have $|X_j| = |X|/(k-1) \ge |Z|$. By Lemma 6.4, there exists a partition of X_1, \ldots, X_{k-1}, Z into $\{X_1^j : j \in [r] \cup \{0\}\}, \ldots, \{X_{k-1}^j : j \in [r] \cup \{0\}\}, \{Z^j : j \in [r]\}$ such that for each $i \in [k-1], j \in [r]$ and $z \in Z^j$, we have

$$\left|X_{i}^{j}\right| = n \text{ and } e_{j}\left(X_{1}^{j}, \dots, X_{k-1}^{j}, z\right) \ge \frac{\left|X_{1}^{j}\right| \cdots \left|X_{k-1}^{j}\right|}{2r} = \frac{n^{k-1}}{2r}$$

For each $j \in [r]$, let H^j be the k-partite k-graph with vertex classes $X_1^j, \ldots, X_{k-1}^j, Z^j$ and $E(H^j) = E\left(H_j\left[X_1^j, \ldots, X_{k-1}^j, Z^j\right]\right)$.

Let $d = (2r)^{-1}$ and $\gamma = d^{2^k}/2$. Fix $j \in [r]$. Consider $z \in Z^j$ and let $H^j(z)$ be the link graph of z on vertex set X_1^j, \ldots, X_{k-1}^j . Note that $|E(H^j(z))| \ge dn^{k-1}$. By Lemma 6.8 with $H^j(z)$ playing the role of H, there exists a (k-1)-partite (k-1)-uniform subhypergraph J^z of $H^j(z)$ such that $|E(J^z)| \ge \gamma 2^{k-2}n^{k-1}$ and every edge in J^z is contained in at least $\gamma n^{k-1}/2$ many $K_{k-1}^{(k-1)}(2)$ in J^z . For each edge e in J^z , let $J^{z,e}$ be the (k-1)-partite (k-1)-graph on vertex classes X_1^j, \ldots, X_{k-1}^j such that fis an edge in $J^{z,e}$ if and only if $e \cup f$ forms a $K_{k-1}^{(k-1)}(2)$ in J^z . For all $z \in Z^j$ and $e \in E(J^z)$, we have

$$|E(J^z)| \ge 2^{k-2} \gamma n^{k-1}$$
 and $|E(J^{z,e})| \ge \gamma n^{k-1}/2.$

Let $X_i^j = \{x_{i,1}, \ldots, x_{i,n}\}$. Choose $\sigma_1, \ldots, \sigma_{k-2} \in S_n$ independently and uniformly at random. For a (k-1)-partite (k-1)-graph J with vertex classes X_1^j, \ldots, X_{k-1}^j , let

$$J_{\sigma_1,\dots,\sigma_{k-2}} = \{x_{1,\sigma_1(i)}\dots x_{k-2,\sigma_{k-2}(i)}x_{k-1,i} : i \in [n]\} \cap E(J).$$

By Lemma 6.9 we have

$$\mathbb{P}\left(\left|J_{\sigma_1,\dots,\sigma_{k-2}}^z\right| < \gamma 2^{k-3}n\right) \le 2e^{-\gamma^2 2^{2k-11}n} \text{ and } \mathbb{P}\left(\left|J_{\sigma_1,\dots,\sigma_{k-2}}^{z,e}\right| < \frac{\gamma n}{4}\right) \le 2e^{-\frac{\gamma^2 n}{512}}$$

By the union bound, there exist some $\sigma_1, \ldots, \sigma_{k-2} \in S_n$ such that for all $z \in Z^j$ and all $e \in E(J^z)$,

$$\left|J_{\sigma_1,\dots,\sigma_{k-2}}^z\right| \ge \gamma 2^{k-3} n = d^{2^k} 2^{k-4} n \text{ and } \left|J_{\sigma_1,\dots,\sigma_{k-2}}^{z,e}\right| \ge \frac{\gamma n}{4} = \frac{d^{2^k} n}{8} = \frac{(2r)^{-2^k} n}{8}.$$
 (6.5)

Without loss of generality (by relabelling vertices of X_1^j, \ldots, X_{k-2}^j) we may assume $\sigma_1 = \cdots = \sigma_{k-2} = id$. For $z \in Z^j$, define the edge-coloured graph G_z on $V = \{v_i : i \in [n]\}$ with colour z to be such that $v_i v_j \in E(G)$ if and only if $x_{1,i} \ldots x_{k-1,i} x_{1,j} \ldots x_{k-1,j}$ forms a $K_{k-1}^{(k-1)}(2)$ in J^z . Hence after removing isolated vertices from G_z , $\delta(G_z) \geq (2r)^{-2^k} n/8$ by (6.5). Let $G^j = \bigcup_{z \in Z^j} G_z$. It is easy to check that (A1) to (A4) are satisfied. This completes the proof of the lemma.

7 Rainbow Path Systems

From now on, we will work with edge-coloured multigraph G. Note that the colours of G are different to those in $K_n^{(k)}$. We often assume that $\delta_{\text{mon}}(G)$ is linear in |V(G)|. Recall that a rainbow path system is a path system (collection of vertex-disjoint paths) such that its union (viewed as a graph) is rainbow. Our first goal is to prove the following result, which shows that there is a rainbow path system in Gwith few paths.

Lemma 7.1. Let $n, d \in \mathbb{N}$ and G be an edge-coloured multigraph on n vertices with $\delta_{mon}(G) \geq d \geq 4|\phi(G)|$. Then there exists a rainbow path system \mathcal{P} such that $\phi(\mathcal{P}) = \phi(G)$ and $|\mathcal{P}| \leq 2n/d$.

To show this, we need the help of the following proposition which lets us start building \mathcal{P} with a rainbow matching.

Proposition 7.2. Let G be an edge-coloured multigraph with $\delta_{mon}(G) \ge 2|\phi(G)| - 1$. Then there exists a rainbow matching M such that $\phi(M) = \phi(G)$.

Proof. Without loss of generality let $\phi(G) = [d]$. Suppose for some $i \in [d]$, we have already found a rainbow matching M_{i-1} such that $\phi(M_{i-1}) = [i-1]$. We now construct M_i as follows. Since

$$\delta_{\mathrm{mon}}(G) \ge 2|\phi(G)| - 1 > 2|\phi(M_{i-1})| = |V(M_{i-1})|,$$

there exists a vertex $v \in V(G) \setminus V(M_{i-1})$ such that $N_i(v, \overline{V(M_{i-1})}) \neq \emptyset$. Pick a vertex $v' \in N_i(v, \overline{V(M_{i-1})})$ and let $M_i = M_{i-1} \cup \{vv'\}$. Let $M = M_d$.

We now present the proof of Lemma 7.1.

Proof of Lemma 7.1. Let \mathcal{P} be a rainbow path system of G with $\phi(\mathcal{P}) = \phi(G)$, which exists by Proposition 7.2. Suppose that $|\mathcal{P}|$ is minimal. If $|\mathcal{P}| \leq 2n/d$, then we are done. So we may assume $|\mathcal{P}| > 2n/d$. Let $\mathcal{P} = \{P_1, P_2, \ldots, P_\ell\}$ be such that $P_i = v_1^i \ldots v_{q_i}^i$ and without loss of generality $\phi(v_1^i v_2^i) = i$. Note that

$$d_i(v_2^i, \overline{V(\mathcal{P})}) \ge d - |V(\mathcal{P})| \ge d - 2|\phi(G)| \ge d/2.$$

Hence there exist distinct $i, j \in [\ell]$ such that $(N_i(v_2^i) \cap N_j(v_2^j)) \setminus V(\mathcal{P}) \neq \emptyset$. Let $w \in (N_i(v_2^i) \cap N_j(v_2^j)) \setminus V(\mathcal{P})$. Let $\mathcal{P}' = (\mathcal{P} \setminus \{P_i, P_j\}) \cup \{v_q^j ... v_3^j v_2^j w v_2^j v_3^i ... v_q^i\}$. Note that \mathcal{P}' is also a rainbow path system with $\phi(\mathcal{P}') = \phi(G)$ and $|\mathcal{P}'| < |\mathcal{P}|$. This contradicts the minimality of $|\mathcal{P}|$.

The following lemma is our absorption result, the proof of which will be presented in the next section.

Lemma 7.3. Let $1/n \ll \delta$ and G be an edge-coloured multigraph satisfying |V(G)| = n and $\delta_{mon}^*(G) \ge \delta n$. Then there exists a subgraph G^* of G such that $|V(G) \setminus V(G^*)|, |\phi(G) \setminus \phi(G^*)| < 2000\delta^{-4}$. Moreover, for all rainbow paths P^* in G^* with $|V(P^*)| \ge 3$ and vertex sets $S \subseteq V(G^*) \setminus V(P^*)$ such that $|V(P^*) \cup S| \le \delta n/4, G \setminus S$ contains a rainbow cycle C^* with $\phi(P^*) \subseteq \phi(C^*) \subseteq \phi(P^*) \cup (\phi(G) \setminus \phi(G^*))$.

We now prove Lemma 6.2 using Lemma 7.3. We iteratively reserve a set of vertices and colours, which shall be used to close the rainbow path system constructed at the end, one path at a time.

Proof of Lemma 6.2. If $|\phi(G)| \leq 2^{18} \delta_0^{-5}$ then we have $\delta_{\text{mon}}(G) \geq 2|\phi(G)|$. By Proposition 7.2, G contains a rainbow matching M such that $\phi(M) = \phi(G)$. Since each edge is a degenerate cycle, we are done setting $\mathcal{C} = M$. Therefore, we may assume $|\phi(G)| > 2^{18} \delta_0^{-5}$.

done setting C = M. Therefore, we may assume $|\phi(G)| > 2^{18} \delta_0^{-5}$. Let $G^1 = G$. For each $j \in [4\delta_0^{-1}]$ in turns, we apply Lemma 7.3 with $G^j, \delta_0/2$ playing the roles of G, δ , respectively, to obtain a subgraph G^{j+1} of G such that

$$|V(G^j) \setminus V(G^{j+1})|, \ |\phi(G^j) \setminus \phi(G^{j+1})| < 2000(\delta_0/2)^{-4} \le 2^{15}\delta_0^{-4}.$$

Moreover, for all rainbow paths P in G^{j+1} with $|V(P)| \ge 3$ and a vertex set $S \subseteq V(G^{j+1}) \setminus V(P)$ such that $|V(P) \cup S| \le \delta_0 n/8$, $G^{j+1} \setminus S$ contains a rainbow cycle C with

$$\phi(P) \subseteq \phi(C) \subseteq \phi(P) \cup (\phi(G^{j}) \setminus \phi(G^{j+1})).$$

Note that

$$\delta_{\text{mon}}(G^{j+1}) \ge \delta_{\text{mon}}(G^j) - |V(G^j) \setminus V(G^{j+1})| \ge \delta_{\text{mon}}(G^j) - 2^{15}\delta_0^{-4} \ge \delta_{\text{mon}}(G) - 2^{15}\delta_0^{-4}j \ge \delta_0 n/2.$$
(7.1)

Let $G^* = G^{4\delta_0^{-1}}$. Note that

$$\delta_{\mathrm{mon}}(G^*) = \delta_{\mathrm{mon}}(G^{4\delta_0^{-1}}) \ge \delta_0 n/2 \ge 8\phi(G^*)$$

and

$$|\phi(G) \setminus \phi(G^*)|, |V(G) \setminus V(G^*)| \le 2^{15} \delta_0^{-4} (4\delta_0^{-1}) = 2^{17} \delta_0^{-5}.$$
(7.2)

By Lemma 7.1 with G^* , $\delta_0 n/2$ playing the roles of G, d, respectively, there exists a rainbow path system \mathcal{P} such that $\phi(\mathcal{P}) = \phi(G^*)$ and

$$|\mathcal{P}| \le \frac{2|V(G^*)|}{\delta_0 n/2} \le 4\delta_0^{-1}.$$
(7.3)

Note that

$$|V(\mathcal{P})| = |\phi(G^*)| + |\mathcal{P}| \le \frac{\delta_0 n}{16} + 4\delta_0^{-1} \le \frac{\delta_0 n}{8}.$$
(7.4)

Moreover

$$\delta_{\mathrm{mon}}(G \setminus V(\mathcal{P})) \stackrel{(7.4)}{\geq} \delta_0 n - \frac{\delta_0 n}{8} \ge \frac{\delta_0 n}{2} \ge 2\phi(G)$$

and $\phi(G \setminus V(\mathcal{P})) \supseteq \phi(G) \setminus \phi(G^*)$. By Proposition 7.2, we deduce that there exists a rainbow matching M in $(G \setminus V(\mathcal{P}))_{\phi(G) \setminus \phi(G^*)}$ with $\phi(M) = \phi(G) \setminus \phi(G^*)$. Let $Q_0 = M \cup \mathcal{P}$, so $\phi(Q_0) = \phi(G)$ and

$$|Q_0| \le |\mathcal{P}| + |\phi(G) \setminus \phi(G^*)| \stackrel{(7.3), (7.2)}{\le} 4\delta_0^{-1} + 2^{17}\delta_0^{-5} \le 2^{18}\delta_0^{-5}.$$

Let P_1, \ldots, P_ℓ be paths in Q_0 such that for $j \in [\ell], |V(P_j)| \ge 3$. Suppose for some $j \in [\ell - 1] \cup \{0\}$, we have constructed Q_j such that

- (i) Q_j is a union of vertex-disjoint paths and cycles with $|Q_j| \le |Q_0|$;
- (ii) Q_j is rainbow with $\phi(Q_j) = \phi(G)$;
- (iii) the paths in Q_j of order at least 3 are precisely $P_{j+1}, \ldots, P_{\ell}$;
- (iv) all edges in Q_j of colours in $\phi(G^{j+1}) \setminus \phi(G^*)$ form a rainbow matching.

We now construct Q_{j+1} as follows. Note that $P_{j+1} \subseteq G^* \subseteq G^{j+1}$. Let $S = V(Q_j \setminus P_{j+1}) \cap V(G^{j+1})$. Note that

$$|S \cup V(P_{j+1})| \le |V(Q_j)| \le |\phi(Q_j)| + |Q_j| \le |\phi(G)| + |Q_0| \le \frac{\delta_0 n}{16} + 2^{18} \delta_0^{-5} \le \frac{\delta_0 n}{8} \le \frac{\delta_0 n}{4}.$$

By the property of G^{j+1} , there is a rainbow cycle C^{j+1} in $G^{j+1} \setminus S$ such that

$$\phi(P_{j+1}) \subseteq \phi(C^{j+1}) \subseteq \phi(P_j) \cup \left(\phi(G^j) \setminus \phi(G^{j+1})\right).$$

Since $\mathcal{P} \subseteq G^*$, the subgraph $G_{\phi(C^{j+1})} \cap Q_j$ of Q_j induced by the colours in $\phi(C^{j+1})$ is precisely the path P_{j+1} and single edges of colour in $\phi(C^{j+1}) \setminus \phi(P_{j+1}) \subseteq \phi(G^j) \setminus \phi(G^{j+1})$. Let $Q_{j+1} = (Q_j - G_{\phi(C^{j+1})}) \cup C^{j+1}$. It is easy to check that (i) to (iv) hold. Finally, note that Q_ℓ consists of cycles and single edges. The proof of the lemma is complete.

8 Closing rainbow paths

The aim of this section is to prove Lemma 7.3. To prove Lemma 7.3, we would need to reserve some vertices and colours so that we can close any rainbow path. Ideally our aim is to find a vertex v, small disjoint vertex sets W_1, W_2 and small disjoint colour sets C_1, C_2 such that for every $x \in V(G)$ and $i \in [2]$, there exists a rainbow path $P_{i,x}$ from x to v such that $V(P_{i,x}) \subseteq \{x, v\} \cup W_i$ and $\phi(P_{i,x}) \subseteq C_i$. We now reserve the vertex set $v \cup W_1 \cup W_2$ and colour set $C_1 \cup C_2$. Given any rainbow path P with end vertices x and $y, P \cup P_{1,x} \cup P_{1,y}$ forms a rainbow cycle. This will motivate our definition of a bowtie (see later). First, we need the following definition which describes how to reach a vertex v through a vertex set W using a colour set C.

Recall that $V^*(G) = \{v \in V(G) : |\phi_G(v)| \ge 2\}$ and for a path $P = v_1 \dots v_\ell$, we denote $\operatorname{int}(P) = \{v_i : 2 \le i \le \ell - 1\}$. Let G be an edge-coloured multigraph with vertex set V. For $v \in V, C \subseteq \phi(G)$ and $W \subseteq V$, define $U_G(v, C', W)$ to be the subset $U \subseteq V$ such that for all vertices $u \in U$, there is a rainbow path P from v to u with $\phi(P) \subseteq C$ and $\operatorname{int}(P) \subseteq W$. We will always assume that $\{v\} \cup W \subseteq V^*(G) \cap U_G(v, C, W)$. Let $U = U_G(v, C, W)$ and $g \in \mathbb{N}$. We say that U is g-maximal in G if for all $u \in U, w_1, w_2 \in V$ and distinct $c, c_1, c_2 \notin C$ with $\phi(uw_1) = c_1$ and $\phi(w_1w_2) = c_2$,

$$d_c(u,\overline{U}), d_{c_1}(w_1,\overline{U}), d_{c_2}(w_2,\overline{U}) < g.$$

If U is not g-maximal, then by adding at most two vertices to W and three colours to C, we can enlarge U by at least g. This leads to the following proposition.

Proposition 8.1. Let G be an edge-coloured multigraph. Let $v \in V^*(G)$ and $c \in \phi_G(v)$. Then there exists a colour set $C \subseteq \phi(G)$ containing c and a vertex-set $W \subseteq V^*(G) \setminus v$ such that $U_G(v, C, W)$ is g-maximal, $|U_G(v, C, W)| \ge d_c(v)$ and $|C|, |W| \le 3|U_G(v, C, W)|/g$.

Proof. Initially, we set $W = \emptyset$ and $C = \{c\}$. Note that $U_G(v, C, W) = N_c(v)$, so $|U_G(v, C, W)| \ge d_c(v)$ and $|W|, |C| \le 3|U_G(v, C, W)|/g$.

If $U_G(v, C, W)$ is g-maximal, then we are done. Suppose that $U_G(v, C, W)$ is not g-maximal. If there exists $u \in U_G(v, C, W)$ and a colour $c' \notin C$ such that $d_{c'}(u, \overline{U_G(v, C, W)}) \geq g$, then add c' to C and u to W. If there exists $u \in U_G(v, C, W)$, $w \in V^*(G)$ and a colour $c' \notin C$ such that $d_{c'}(w, \overline{U_G(v, C, W)}) \geq g$ and $\phi(uw) \notin \{c'\} \cup C$, then add $c', \phi(uw)$ to C and u, w to W. If there exists $u \in U_G(v, C, W)$, $w, w' \in V^*(G)$ and a colour c' such that $d_{c'}(w', \overline{U_G(v, C, W)}) \geq g$ and $\phi(uw), \phi(ww'), c'$ are distinct colours not in C, then add $c', \phi(uw), \phi(ww')$ to C and u, w, w' to W.

Note that we still have $|W|, |C| \leq 3|U_G(v, C, W)|/g$ and $|U_G(v, C, W)| \geq d_c(v)$. Note that $W \subseteq V^*(G)$. We repeat this process until $U_G(v, C, W)$ is g-maximal.

Recall that $\delta_{\text{mon}}^*(G) = \min_{v \in V^*(G)} \min_{c \in \phi_G(v)} d_c(v)$. In other words, $\delta_{\text{mon}}^*(G)$ is the (non-zero) minimum number of edges seen by a vertex v in a particular colour, minimised over all the vertices in G that see at least two colours. We are mainly concerned about vertices in $V^*(G)$ because if a vertex $v \notin V^*(G)$, then it cannot be contained in a non-degenerate rainbow cycle. Also recall that the *closed* neighborhood of a vertex w in a graph G is defined as $\{w\} \cup N_G(w)$ and denoted by $N_G[w]$.

Let U = U(v, C, W) be g-maximal in G. The following crucial lemma says that if a rainbow path P satisfies $int(P) \cap N_{G-G_C}[U] \neq \phi$, then $int(P) \subseteq N_{G-G_C}[U]$.

Lemma 8.2. Let $g \in \mathbb{N}$ and G be an edge-coloured multigraph on n vertices with $\delta^*_{mon}(G) > g$. Let $v \in V(G)$, $W \subseteq V(G)$ and $C \subseteq \phi(G)$. Suppose that $U = U_G(v, C, W)$ is g-maximal. Furthermore let $G' = G - G_C$ and $U^* = N_{G'}[U]$. Then the following hold:

- (i) for all $x \in U^* \cap V^*(G')$ and $c \in \phi(G')$, we have $d_c(x, \overline{U}) < g$;
- (ii) if P is a rainbow path in G' such that $int(P) \cap U^* \neq \emptyset$, then $int(P) \subseteq U^*$.

Proof. Consider $x \in U^* \cap V^*(G')$ and $c \in \phi(G')$. If $x \in U$, then since $c \notin C$, by the g-maximality of U, we have $d_c(x,\overline{U}) < g$. If $x \in U^* \setminus U$, then there exists a vertex $u \in U$ and a colour $c' \notin C$ such that $\phi(ux) = c'$. If $c' \neq c$, then $d_c(x,\overline{U}) < g$ by the g-maximality of U. If c = c', then there exists a colour $c'' \notin \phi_{G'}(v) \setminus \{c\}$ as $v \in V^*(G')$. By the argument above, we have $d_{c''}(x,\overline{U}) < g$. Since $\delta^*_{\mathrm{mon}}(G) > g$, we deduce that $N_{c''}(x,U) \neq \emptyset$. Pick $u' \in N_{c''}(x,U)$. Now, $\phi(u'x) = c'' \neq c$. Again by the previous argument we have $d_c(x,\overline{U}) < g$. Hence (i) holds.

We now prove (ii). Let $P = x_1 \dots x_\ell$ be a rainbow path in G' with $\operatorname{int}(P) \cap U^* \neq \emptyset$. Suppose that $\phi(x_{j-1}x_j) = j$ for $j \in [\ell] \setminus \{1\}$. Furthermore assume that $x_2 \in U^*$ and $\ell \geq 4$. (Indeed, if $x_i \in U^*$ with $i \in [3, \ell - 2]$, then consider the two rainbow paths $x_{i-1}x_i \dots x_\ell$ and $x_{i+1}x_i \dots x_1$ separately.) Thus it is enough to show that $x_3 \in U^*$, as we can then consider the rainbow path $x_2x_3 \dots x_\ell$. If $x_2 \in U$, then $x_3 \in U^*$. If $x_2 \in U^* \setminus U$, then (i) implies $d_2(x_2, \overline{U}) < g$ and so $N_2(x_2, U) \neq \emptyset$. Then by g-maximality of $U, d_4(x_3, \overline{U}) < g$ and so $N_4(x_3, U) \neq \emptyset$. Thus $x_3 \in U^*$ as required.

Let G be an edge-coloured multigraph. We say that $B = (v, C_1, W_1, C_2, W_2)$ is a bowtie in G if $v \in V(G)$, $W_1, W_2 \subseteq V(G) \setminus \{v\}$ are disjoint, C_1 and C_2 are disjoint nonempty colour sets. Denote $\phi(B) = C_1 \cup C_2$, $W(B) = \{v\} \cup W_1 \cup W_2$ and for $i \in [2]$, $U_i(B|G) = U_{G \setminus W_{3-i}-G_{C_{3-i}}}(v, C_i, W_i)$. A bowtie B is g-maximal in G if for $i \in [2]$, $U_i(B|G)$ is g-maximal in $G \setminus W_{3-i} - G_{C_{3-i}}$.

The following corollary shows that a g-maximal bowtie exists, which follows from Proposition 8.1.

Corollary 8.3. Let $g \in \mathbb{N}$ and G be an edge-coloured multigraph on n vertices. Let v be a vertex such that $d_{c_1}(v), d_{c_2}(v) \geq g + 3n/g$ for distinct colours c_1, c_2 . Then there exists a g-maximal bowtie $B(v, C_1, W_1, C_2, W_2)$ such that for $i \in [2]$, $c_i \in C_i$, $W(B) \subseteq V^*(G)$, $|U_i(B|G)| \geq d_{c_i}(v) - 3n/g$ and $|\phi(B)|, |W(B)| \leq 6n/g$.

Proof. Apply Proposition 8.1 with $v, c_1, G - G_{c_2}$ playing the roles of v, c, G, respectively and obtain a colour set C_1 and a vertex set W_1 such that $c_1 \in C_1$ and $U_{G-G_{c_2}}(v, C_1, W_1)$ is g-maximal in $G - G_{c_2}$ and $|U_{G-G_{c_2}}(v, C_1, W_1)| \ge d_{c_1}(v)$. Note that

$$|C_1|, |W_1| \le 3|U_{G-G_{c_2}}(v, C_1, W_1)|/g \le 3n/g.$$

Let $G' = G \setminus W_1 - G_{C_1}$. Apply Proposition 8.1 again with v, c_2, G' , playing the roles of v, c, G, respectively and obtain a colour set C_2 and a vertex set W_2 such that $c_2 \in C_2$ and $U_{G'}(v, C_2, W_2)$ is g-maximal in G'and

$$|U_{G'}(v, C_2, W_2)| \ge d_{c_2, G'}(v) \ge d_{c_2, G}(v) - |W_1| \ge d_{c_2, G}(v) - 3n/g.$$

Set $B = (v, C_1, W_1, C_2, W_2)$. Note that B is g-maximal and for $i \in [2]$, $|U_i(B|G)| \ge d_{c_i}(v) - 3n/g$. Furthermore,

$$|\phi(B)|, |W(B)| \le \frac{3}{g} \left(\sum_{i \in [2]} |U_i(B|G)| \right) \le \frac{6n}{g}.$$

We now continue our motivation for the proof of Lemma 7.3. Recall that our aim is to use a bowtie to close a rainbow path. First, we show that there exist a small set of bowties B_1, \ldots, B_t such that $\{U_2(B_i|G) : i \in [t]\}$ partitions $V^*(G)$. This will ensure that any rainbow path can be extended via one of these bowties (but we may not be able to close it).

Let $d, g \in \mathbb{N}$ and G be an edge-coloured multigraph. Let \mathcal{B} be a family of bowties B_1, \ldots, B_t in G. Let $\phi(\mathcal{B}) = \bigcup_{i \in [t]} \phi(B_i)$ and $W(\mathcal{B}) = \bigcup_{i \in [t]} W(B_i)$. We write $G - \mathcal{B}$ for $G \setminus W(\mathcal{B}) - G_{\phi(\mathcal{B})}$. For a bowtie $B \in \mathcal{B}$ and $i \in [2]$, we denote $U_i(B|G, \mathcal{B}) = U_i(B|G - (\mathcal{B} \setminus B))$ and $U_i^*(B|G, \mathcal{B}) = N_{G-G_{\phi(\mathcal{B})}}[U_i(B|G, \mathcal{B})] \cap V^*(G - \mathcal{B})$. We say \mathcal{B} is a (d, g)-partition of G if the following hold

(P1) for all $i \in [2]$ and $j \in [t]$, we have $|U_i(B_j|G, \mathcal{B})| \ge d$;

- (**P2**) $W(B_1), \ldots, W(B_t)$ are all disjoint and $W(\mathcal{B}) \subseteq V^*(G)$;
- (**P3**) $\phi(B_1), \ldots, \phi(B_t)$ are all disjoint;

(**P4**) for all $j \in [t]$, B_j is g-maximal in $G - (\mathcal{B} \setminus B_j)$;

(**P5**) for all distinct $j, j' \in [t], U_2^*(B_j|G, \mathcal{B}) \cap U_2^*(B_{j'}|G, \mathcal{B}) = \emptyset;$

(P6) $U_2^*(B_1|G, \mathcal{B}), U_2^*(B_2|G, \mathcal{B}), \ldots, U_2^*(B_t|G, \mathcal{B})$ partition $V^*(G - \mathcal{B})$.

We say that \mathcal{B} is a weak (d, g)-partition of G if only (P1) to (P5) hold.

Fact 8.4. Let $d, g, n \in \mathbb{N}$ and G be a graph on n vertices. Let \mathcal{B} be a weak (d, g)-partition of G. Then $|\mathcal{B}| \leq n/d$.

Proof. Consider $B \in \mathcal{B}$. By (P1), $|U_2^*(B|G, \mathcal{B}) \cup W(B)| \ge |U_2(B|G, \mathcal{B})| \ge d$. Note that $U_2^*(B|G, \mathcal{B}) \subseteq V^*(G - \mathcal{B}) \subseteq V(G) \setminus W(\mathcal{B})$. By (P2) and (P3), for any distinct $B, B' \in \mathcal{B}$,

$$(U_2^*(B|G,\mathcal{B})\cup W(B))\cap (U_2^*(B'|G,\mathcal{B})\cup W(B'))=\emptyset.$$

Therefore, $|\mathcal{B}| \leq n/d$.

The next lemma shows that one can extend a weak (d, g)-partition into a (d, g)-partition.

Lemma 8.5. Let $d, g, n \in \mathbb{N}$ with $d \geq 4g$ and $g \geq \max\{3n/g, 12n^2/gd\}$. Let G be an edge-coloured multigraph on n vertices such that $\delta^*_{mon}(G) \geq d$. Suppose \mathcal{B}_0 is a weak (d/2, g)-partition of G with $|\phi(\mathcal{B}_0)|, |W(\mathcal{B}_0)| \leq 6n|\mathcal{B}_0|/g$. Then there exists a (d/2, g)-partition \mathcal{B}^* so that $\mathcal{B}_0 \subseteq \mathcal{B}^*$ and $|\phi(\mathcal{B}^*)|, |W(\mathcal{B}^*)| \leq 6n|\mathcal{B}^*|/g \leq 12n^2/gd$.

Proof. Suppose we have already constructed a weak (d/2, g)-partition $\mathcal{B} = \{B_1, \ldots, B_t\}$ with $\mathcal{B}_0 \subseteq \mathcal{B}$ and $|\phi(\mathcal{B})|, |W(\mathcal{B})| \leq t(6n/g)$. By Fact 8.4,

$$t \le 2n/d. \tag{8.1}$$

Hence

$$|\phi(\mathcal{B}^*)|, |W(\mathcal{B}^*)| \le \left(\frac{2n}{d}\right) \left(\frac{6n}{g}\right) = \frac{12n^2}{gd}.$$

We further assume that t is maximal. If \mathcal{B} satisfies (P6) then we are done by setting $\mathcal{B}^* = \mathcal{B}$.

Thus we may assume that (**P6**) does not hold. We now construct a bowtie B_{t+1} as follows. Let $G' = G - \mathcal{B}$. Pick $v^{t+1} \in V^*(G') \setminus \bigcup_{i \in [t]} U_2^*(B_i | G, \mathcal{B})$ and $c_1, c_2 \in \phi_{G'}(v^{t+1})$, which exist as (**P6**) does not hold for \mathcal{B} . Note that

$$\delta_{\text{mon}}^*(G') \ge \delta_{\text{mon}}^*(G) - |W(\mathcal{B})| \ge d - (12n^2/gd) \ge d - g \ge d/2 \ge 2g \ge g + 3n/g.$$

By Corollary 8.3, G' contains a g-maximal bowtie B_{t+1} such that $|\phi(B_{t+1})|, |W(B_{t+1})| \leq 6n/g$ and $W(B_{t+1}) \subseteq V^*(G')$.

Let $\mathcal{B}' = \mathcal{B} \cup B_{t+1}$. Clearly, $|\phi(\mathcal{B}')|, |W(\mathcal{B}')| \leq (t+1)(6n/g)$. We now show that \mathcal{B}' is a weak (d/2, g)-partition, contradicting the maximality of t. Fix $i \in [2]$ and $j \in [t+1]$. Note that

$$\delta_{\mathrm{mon}}^*(G - (\mathcal{B}' \setminus B_j)) \ge d - |W(\mathcal{B}' \setminus B_j)| \ge d - (12n^2/gd) \ge d/2$$

Thus, $|U_i(B_j|G, \mathcal{B}')| \ge \delta^*_{\text{mon}}(G - (\mathcal{B}' \setminus B_j)) \ge d/2$ and so (**P1**) holds. Note that (**P2**) and (**P3**) hold by our construction. For $j \in [t]$, B_j is g-maximal in $G - (\mathcal{B} \setminus B_j)$ so it is g-maximal in $G - (\mathcal{B}' \setminus B_j)$. Recall B_{t+1} is g-maximal in $G' = G - \mathcal{B} = G - (\mathcal{B}' \setminus B_{t+1})$. Thus (**P4**) holds for \mathcal{B}' .

It remains to show (**P5**) holds for \mathcal{B}' . For $i \in [t]$, $U_2^*(B_i|G, \mathcal{B}') \subseteq U_2^*(B_i|G, \mathcal{B})$. Since \mathcal{B} is a weak (d/2, g)-partition of G, (**P5**) of \mathcal{B} implies that $U_2^*(B_1|G, \mathcal{B}'), \ldots, U_2^*(B_t|G, \mathcal{B}')$ are disjoint. Suppose that $U_2^*(B_{t+1}|G, \mathcal{B}') \cap U_2^*(B_j|G, \mathcal{B}') \neq \emptyset$ for some $j \in [t]$. Let $u \in U_2^*(B_{t+1}|G, \mathcal{B}') \cap U_2^*(B_j|G, \mathcal{B}') \subseteq V^*(G-\mathcal{B})$ and $c \in \phi_{G-\mathcal{B}'}(u)$. Note that

$$d_{c,G-\mathcal{B}'}(u) \ge \delta^*_{\mathrm{mon}}(G-\mathcal{B}') \ge d - (12n^2/gd) > g.$$

Lemma 8.2(i) and (P4) imply that $d_{c,G-\mathcal{B}'}(u, \overline{U_2(B_j|G,\mathcal{B}')}) < g$. Therefore,

$$d_{c,G}(u, U_2(B_j|G, \mathcal{B}')) \ge d_{c,G}(u) - |W(\mathcal{B}')| - d_{c,G-\mathcal{B}'}(u, \overline{U_2(B_j|G, \mathcal{B}')}) > d_{c,G}(u) - \frac{12n^2}{qd} - g \ge \frac{d_{c,G}(u)}{2}.$$

Similarly, we have $d_{c,G}(u, U_2(B_{t+1}|G, \mathcal{B}')) > d_{c,G}(u)/2$. Hence,

$$\begin{aligned} |U_2(B_j|G,\mathcal{B}') \cap U_2(B_{t+1}|G,\mathcal{B}')| &\geq |U_2(B_j|G,\mathcal{B}') \cap U_2(B_{t+1}|G,\mathcal{B}') \cap N_{c,G}(u)| \\ &\geq d_{c,G}(u,U_2(B_j|G,\mathcal{B}')) + d_{c,G}(u,U_2(B_{t+1}|G,\mathcal{B}')) - d_{c,G}(u) \\ &> 0. \end{aligned}$$

Let $w \in U_2(B_j|G, \mathcal{B}') \cap U_2(B_{t+1}|G, \mathcal{B}')$. Let $B_{t+1} = (v^{t+1}, C_1^{t+1}, W_1^{t+1}, C_2^{t+1}, W_2^{t+1})$. Since $w \in U_2(B_{t+1}|G, \mathcal{B}')$, there exists a rainbow path $P = wv_1 \dots v_\ell v^{t+1}$ with $\phi(P) \subseteq C_2^{t+1}$ and $int(P) \subseteq W_2^{t+1}$. By $(\mathbf{P3}), \phi(P) \cap \phi(B_j) = \emptyset$. We deduce that $v_1 \in U_2^*(B_j|G, \mathcal{B}')$. Let $c_1 \in C_1^{t+1} \cap \phi_G(v^{t+1})$. Note that

$$d_{c_1,G'}(v^{t+1}) \ge \delta_{\text{mon}}^*(G') \ge d/2 > |C_2^{t+1}| \ge |V(P)|.$$

Pick $v' \in N_{c_1,G'}(v^{t+1}) \setminus V(P)$. Then the path $P' = wv_1 \dots v_\ell v^{t+1} v'$ is a rainbow path with $\phi(P') \cap \phi(B_j) = \emptyset$. By Lemma 8.2(ii), $v^{t+1} \in int(P') \subseteq U_2^*(B_j|G,\mathcal{B})$, contradicting the fact that v^{t+1} was chosen from $V^*(G') \setminus \bigcup_{i \in [t]} U_2^*(B_i|G,\mathcal{B})$.

8.1 Proof of Lemma 7.3

We now present the ideas in the proof of Lemma 7.3. Let $d = \delta n$.

We apply Lemma 8.5 to obtain a (d/2, g)-partition \mathcal{B} of G. Let P be a rainbow path in $G - \mathcal{B}$. Since $int(P) \subseteq V^*(G - \mathcal{B})$, we deduce (by (P4), (P6) and Lemma 8.2(ii)) that there exists $B \in \mathcal{B}$ such that $int(P) \subseteq U_2^*(B|G, \mathcal{B})$. We now discuss how to augment P into a rainbow cycle using B.

Let $P = x_1 x_2 \dots x_{\ell}$ with $\phi(x_1 x_2) = 2$ and $\phi(x_{\ell-1} x_{\ell}) = \ell$.

Case 1: $|U_2(B|G, \mathcal{B})| \leq 3d/4$. By Lemma 8.2(i), we can find $x \in U_2(B|G, \mathcal{B})$ to replace both x_1 and x_ℓ in P. This transforms P into a rainbow cycle.

Case 2: $U_2^*(B|G, \mathcal{B}) = U_1^*(B|G, \mathcal{B})$. By Lemma 8.2(i), we may assume that $x_1 \in U_1(B|G, \mathcal{B})$ and $x_\ell \in U_2(B|G, \mathcal{B})$. Let $B = (v, C_1, W_1, C_2, W_2)$. There exists a rainbow path P' from x_1 to x_ℓ through v such that $\phi(P') \subseteq \phi(B)$ and $\operatorname{int}(P') \subseteq W(B)$. Hence PP' is a rainbow cycle.

Therefore we would like to ensure Case 1 or Case 2 hold. Our aim is to refine \mathcal{B} by replacing bowties with smaller ones so that Case 1 will hold eventually. In particular, we will increase the number of bowties in each step. For simplicity, suppose $V^*(G - \mathcal{B}) = V(G)$. Note that $B \in \mathcal{B}$ is gmaximal in $G - (\mathcal{B} \setminus B)$. Lemma 8.2(i) implies that "the subgraph H_B induced by $U_2^*(B|G, \mathcal{B})$ satisfies $\delta_{\text{mon}}^*(H_B) \geq d - g \geq d/2$ ". Hence we can apply Lemma 8.5 to obtain a (d/4, g)-partition \mathcal{B}' of H_B . Thus, we refine \mathcal{B} by replacing B with \mathcal{B}' .

Suppose that we are unable to refine \mathcal{B} further and for simplicity \mathcal{B} consists of only one bowtie $B = (v, C_1, W_1, C_2, W_2)$. If Case 2 fails, then $U_1^*(B|G, \mathcal{B})$ is smaller than $U_2^*(B|G, \mathcal{B})$. We consider the 'swapped' bowtie $B' = (v, C_2, W_2, C_1, W_1)$ instead. We extend this weak partition $\{B'\}$ into a (d/4, g)-partition of G (using Lemma 8.5). Note that we increase the number of bowties in the partition.

Proof of Lemma 7.3. Set $d = \delta n$, $\gamma = \delta^2/16$ and $g = \gamma n$. Note that

$$d \ge 4g + 2$$
 and $g \ge \max\{6n/g, 12n^2/gd\}$.

Let $G^0 = G$, $J^0 = G$ and $H^0 = G$. By Lemma 8.5 with H^0, d, g, \emptyset playing the roles of G, d, g, \mathcal{B}_0 , respectively, we obtain a (d/2, g)-partition \mathcal{B}^1 of H^0 and $|\phi(\mathcal{B}^1)|, |W(\mathcal{B}^1)| \leq 2n|\mathcal{B}^1|/g$ and $W(\mathcal{B}^1) \subseteq V^*(H^0)$.

Suppose for some $i \in \mathbb{N}$ we have already constructed families $\mathcal{B}^1, \mathcal{B}^2, \ldots, \mathcal{B}^i$ of bowties, edge-coloured multigraphs $J^0, G^0, H^0, \ldots, J^{i-1}, G^{i-1}, H^{i-1}$ whose properties will be specified later. Let

$$\begin{split} J^{i} &= G^{i-1} - \mathcal{B}^{i}, \\ G^{i} &= J^{i} - \bigcup_{\substack{B,B' \in \mathcal{B}^{i} \\ B \neq B'}} J^{i}[U_{2}^{*}(B|H^{i-1},\mathcal{B}^{i}), U_{2}^{*}(B'|H^{i-1},\mathcal{B}^{i})]. \end{split}$$

In other words, G^i is obtained from $G^{i-1} - \mathcal{B}^i$ by removing all edges between $U_2^*(B|H^{i-1}, \mathcal{B}^i)$ and $U_2^*(B'|H^{i-1}, \mathcal{B}^i)$ for distinct $B, B' \in \mathcal{B}^i$.

Let $\mathcal{B}_1^i = \{B \in \mathcal{B}^i : |U_2(B|H^{i-1}, \mathcal{B}^i)| \leq 3d/4\}$. (These will consist of bowties that satisfy Case 1.) Consider $i \in \mathbb{N}$ and $B \in \mathcal{B}^i$. For a bowtie family \mathcal{B}' in H^i , define

$$\partial_{\mathcal{B}'}(B) = \{ B' \in \mathcal{B}' : W(B') \subseteq U_2^*(B|H^{i-1}, \mathcal{B}^i) \}.$$

We say B is covered by \mathcal{B}' if $\partial_{\mathcal{B}'}(B) = \{B'\}$ and $U_2^*(B'|H^i, \mathcal{B}') = U_1^*(B'|H^i, \mathcal{B}')$. We write ∂B for $\partial_{\mathcal{B}^{i+1}}B$. Let

$$\mathcal{B}_2^i = \begin{cases} \bigcup \{ \partial B \colon B \in \mathcal{B}^{i-1} \text{ is covered by } \mathcal{B}^i \} & \text{if } i \ge 2, \\ \emptyset & \text{if } i = 1. \end{cases}$$

(These bowties in \mathcal{B}_2^i will satisfy Case 2.) Let $\mathcal{B}_3^i = \mathcal{B}^i \setminus (\mathcal{B}_1^i \cup \mathcal{B}_2^i)$ and

$$H^{i} = G^{i} \setminus \left(V^{*}(G^{i}) \setminus \bigcup_{B \in \mathcal{B}_{3}^{i}} U_{2}^{*}(B|H^{i-1}, \mathcal{B}^{i}) \right) = G^{i} \left[(V(G^{i}) \setminus V^{*}(G^{i})) \cup \bigcup_{B \in \mathcal{B}_{3}^{i}} U_{2}^{*}(B|H^{i-1}, \mathcal{B}^{i}) \right].$$

In other words, H^i is induced by $U_2^*(B|H^{i-1}, \mathcal{B}^i)$ for all $B \in \mathcal{B}_3^i$ and the vertices in G^i that see at most one colour. From now on, for $B \in \mathcal{B}^j$, we write $U_i^*(B)$ for $U_i^*(B|H^{j-1}, \mathcal{B}^j)$. We now list the desired properties of G^i and H^i . Suppose that for all $j \in [i]$,

- (i) \mathcal{B}^j is a (d/4, g)-partition of H^{j-1} with $|\phi(\mathcal{B}^j)|, |W(\mathcal{B}^j)| < 384\delta^{-3}$ and $W(\mathcal{B}^j) \subseteq V^*(H^{j-1});$
- (ii) if $B' \in \mathcal{B}^{j-1}$ and $|\partial B'| = 1$, then B' is covered by \mathcal{B}^j ;
- (iii) if $j \ge 2$, then $\mathcal{B}^j = \bigcup_{B \in \mathcal{B}_2^{j-1}} \partial B$;
- (iv) for all $B \in \mathcal{B}_{3}^{j}, |U_{2}^{*}(B)| \leq n j(d/4);$
- (v) $\delta_{\text{mon}}(G^j) \ge d j(g + 12n^2/gd) \ge d 2gj;$
- (vi) if $V^*(H^j)$ is not empty, then $\delta^*_{\text{mon}}(H^j) \ge d j(g + 12n^2/gd) \ge d 2gj;$
- (vii) $V^*(H^j) \subseteq \bigcup_{B \in \mathcal{B}_3^j} U_2^*(B);$
- (viii) $\phi(G^j) = \phi(G^{j-1}) \setminus \phi(\mathcal{B}^j);$
- (ix) G^j has no edge between $U_2^*(B)$ and $V^*(G^j) \setminus U_2^*(B)$ for all $B \in \mathcal{B}^j$;
- (x) for each $B \in \mathcal{B}^j$, B is g-maximal in $G^{j-1} (\mathcal{B}^j \setminus B)$.

It should be noted that (i) and (ii) imply (iii) to (x). Hence we technically only require $\mathcal{B}^1, \ldots, \mathcal{B}^j$ to satisfy (i) and (ii). Note that (iv) implies that

$$i \le 4n/d = 4/\delta. \tag{8.2}$$

Case A: $V^*(H^i) \neq \emptyset$. We now construct \mathcal{B}^{i+1} as follows. We have

$$\delta_{\text{mon}}^*(H^i) \ge d - 2g(4\delta^{-1}) \ge d/2.$$

By Lemma 8.5 with $H^i, d/2, g, \emptyset$ playing the roles of G, d, g, \mathcal{B}_0 , respectively, there exists a (d/4, g)-partition \mathcal{B} of H^i with $|\phi(\mathcal{B})|, |W(\mathcal{B})| \leq 6n|\mathcal{B}|/g = 192\delta^{-3}$.

Claim 8.6. Let \mathcal{B} be a (d/4, g)-partition of H^i with $|\phi(\mathcal{B})|, |W(\mathcal{B})| \leq 12n^2/gd$. Let $B^+ \in \mathcal{B}^i$ and $B \in \partial_{\mathcal{B}}B^+$. Then for $t \in [2]$, $U_t^*(B|H^i, \mathcal{B}) \subseteq U_2^*(B^+)$. Moreover, if $\partial_{\mathcal{B}}B^+ = \{B\}$, then $U_2^*(B|H^i, \mathcal{B}) = U_2^*(B^+) \cap V^*(H^i - \mathcal{B})$.

Proof of claim. We prove the case for t = 2 (and the case for t = 1 is proven analogously). Let $B = (v, C_1, W_1, C_2, W_2)$, so $v \in U_2^*(B^+)$. Suppose there exists a vertex

$$u \in U_2^*(B|H^i, \mathcal{B}) \setminus U_2^*(B^+).$$

$$(8.3)$$

Then there exists a rainbow path $P = vw_1 \dots w_\ell u$ such that $w_1, \dots, w_{\ell-1} \in W_2$, $\phi(P) \subseteq C_2 \cup \phi(w_\ell u) \subseteq \phi(H^i)$ and $C_1 \cap \phi(P) = \emptyset$. Note that

$$|P| \le |W(\mathcal{B})| + 2 \le 192\delta^{-3} + 2 < d/2 \tag{8.4}$$

and $u \in V^*(H^i - \mathcal{B})$. Thus, there exists a colour $c' \in \phi_{H^i - \mathcal{B}}(u) \setminus \phi(w_\ell v)$. By (vi),

$$d_{c',H^i}(u) \ge d - 2ig \ge d - \frac{8g}{\delta} \stackrel{(8.4)}{>} |P|.$$

Pick $u' \in N_{c',H^i}(u) \setminus V(P)$. Recall that $\phi(H^i) \supseteq \phi(\mathcal{B}) \supseteq C_1 \neq \emptyset$. Pick $c_1 \in C_1 \cap \phi_{H^i}(v) \subseteq \phi(H^i)$. By (vi),

$$d_{c_1,H^{i-1}}(v) \ge d - 2g(i-1) \ge d - \frac{8g}{\delta} > |P| + 1.$$

Pick $v' \in N_{c_1,H^{i-1}}(u) \setminus (V(P) \cup \{u'\})$. Then the path $P' = v'Pu' = v'vw_1 \dots w_\ell uu'$ is rainbow in H^{i-1} with

$$\phi(P') \cap \phi(\mathcal{B}^i) \subseteq \phi(H^i) \cap \phi(\mathcal{B}^i) \subseteq \phi(G^i) \cap \phi(\mathcal{B}^i) = \emptyset$$

by (viii). Recall that $U_2^*(B^+)$ is g-maximal in $H^{i-1} - (\mathcal{B}^i \setminus B^+)$. Lemma 8.2(ii) implies that $u \in int(\mathbf{P}') \subseteq U_2^*(\mathbf{B}^+)$ contradicting (8.3). The moreover statement follows.

For each $B^+ \in \mathcal{B}^i$ with $\partial_{\mathcal{B}}B^+ = \{B\}$ that is not covered by \mathcal{B} , we replace the bowtie $B = (v, C_1, W_1, C_2, W_2)$ with $B' = (v, C_2, W_2, C_1, W_1)$. We call the resulting family \mathcal{B}' . We now show that \mathcal{B}' is a weak (d/4, g)-partition of H^i . Note that (**P1**) to (**P4**) hold. To show (**P5**), note that if $B' \in \mathcal{B}' \setminus \mathcal{B}$, then there exist unique $B \in \mathcal{B} \setminus \mathcal{B}'$ and $B^+ \in \mathcal{B}^i$ with

$$B = (v, C_1, W_1, C_2, W_2), B' = (v, C_2, W_2, C_1, W_1) \text{ and } \partial_{\mathcal{B}} B^+ = \{B\}.$$

By Claim 8.6 and the fact $\partial_{\mathcal{B}}B^+ = \{B\}$, we deduce that

$$U_2^*(B'|H^i, \mathcal{B}') = U_1^*(B|H^i, \mathcal{B}) \subseteq U_2^*(B^+) \cap V^*(G^{i+1}) = U_2^*(B|H^i, \mathcal{B}).$$

Recall that \mathcal{B} is a (d/4, g)-partition of H^i , in particular this implies that $\{U_2^*(B|H^i, \mathcal{B}) : B \in \mathcal{B}\}$ are pairwise disjoint. Thus, \mathcal{B}' satisfies (**P5**) and is a weak (d/4, g)-partition. Apply Lemma 8.5 with $H^i, d/2, g, \mathcal{B}'$ playing the roles of G, d, g, \mathcal{B}_0 , respectively and obtain (d/4, g)-partition \mathcal{B}^{i+1} containing \mathcal{B}' and satisfying $W(\mathcal{B}^{i+1}) \subseteq V^*(H^i)$ and

$$|\phi(\mathcal{B}^{i+1})|, |W(\mathcal{B}^{i+1})| \le 2(6n|\mathcal{B}^{i+1}|/g) \le 24n^2/gd \le 384\delta^{-3}.$$

Claim 8.7. $\mathcal{B}^{i+1}, G^{i+1}, H^{i+1}$ satisfy (i) to (x).

Proof of claim. Note that (i) holds by construction.

Consider $B^+ \in \mathcal{B}^i$. If $|\partial_{\mathcal{B}}B^+| = 1$ and B^+ is not covered by \mathcal{B} , then $|\partial_{\mathcal{B}^{i+1}}B^+| > 1$. Otherwise $\partial_{\mathcal{B}}B^+ = \partial_{\mathcal{B}^{i+1}}B^+$. Suppose that $\partial B^+ = \{B\}$. By (**P6**) and Claim 8.6, $U_2^*(B|H^i, \mathcal{B}^{i+1}) = U_2^*(B^+|H^{i-1}, \mathcal{B}^i) \cap V^*(H^i - \mathcal{B}^{i+1})$. Note that B^+ is covered by \mathcal{B} , therefore covered by \mathcal{B}^{i+1} . Thus (ii) holds.

Consider $B = (v, C_1, W_1, C_2, W_2) \in \mathcal{B}^{i+1}$. For each $w \in W(B) \setminus \{v\}$, there is a path from v to w in H^i . Thus W(B) is contained in a component of $H^i[V^*(H^i)]$. Since $H^i \subseteq G^i$, (ix) for G^i implies that $W(B) \subseteq U_2^*(B^+)$ for some $B^+ \in \mathcal{B}_3^i$. Therefore (iii) holds for G^{i+1} .

We now show that (iv) holds. Let $B \in \mathcal{B}_3^{i+1}$. Then there exists $B^+ \in \mathcal{B}^i$ and $B' \in \mathcal{B}_3^{i+1}$ such that $B, B' \in \partial B^+$. By (**P1**), (**P5**) and Claim 8.6, we deduce that $U_2^*(B) \cup W(B)$ and $U_2^*(B') \cup W(B')$ each has size at least d/4, are disjoint and contained in $U_2^*(B^+)$. Hence,

$$U_2^*(B)| \le |U_2^*(B^+) - |U_2^*(B') \cup W(B')|$$

$$\stackrel{(iv)}{\le} n - i(d/4) - (d/4) = n - (i+1)(d/4).$$

To show (v), consider a vertex $u \in V(G^{i+1})$ and a colour $c \in \phi_{G^{i+1}}(u)$. It is enough to show that $d_{c,G^{i+1}}(u) \ge d - (i+1)(g+12n^2/gd)$. By (v) for G^i , we have $d_{c,G^i}(u) \ge d - i(g+12n^2/gd)$. Then

$$d_{c,J^{i+1}}(u) \ge d_{c,G^i}(u) - |W(\mathcal{B}^{i+1})| \ge d - ig - (i+1)(12n^2/gd).$$

If $u \in V(G^{i+1}) \setminus V^*(G^{i+1})$, then $d_{c,G^{i+1}}(u) = d_{c,J^{i+1}}(u)$. If $u \in V^*(G^{i+1})$, then by (**P6**), $u \in U_2^*(B)$ for some $B \in \mathcal{B}^{i+1}$. Then by g-maximality of B in $H^i - (\mathcal{B}^{i+1} \setminus B)$, Lemma 8.2(i) implies that $d_{c,G^{i+1}}(u, \overline{U_2^*(B)}) < g$. We deduce that

$$d_{c,G^{i+1}}(u) \ge d_{c,J^{i+1}}(u) - d_{c,G^{i+1}}(u, \overline{U_2^*(B))}) \ge d - (i+1)(g + 12n^2/gd).$$

This argument also implies (vi), (vii) and $\phi(J^{i+1}) = \phi(G^{i+1})$. Since $\delta_{\text{mon}}(G^i) \ge d/4 > |W(\mathcal{B}^{i+1})|$, we have $\phi(J^{i+1}) = \phi(G^i) \setminus \phi(\mathcal{B}^{i+1})$ implying (viii).

To show (ix), consider $B \in \mathcal{B}^{i+1}$. Let $B^+ \in \mathcal{B}^i$ with $B \in \partial B^+$. By Claim 8.6, $U_2(B) \subseteq U_2^*(B^+)$. Hence G^{i+1} contains no edge between $U_2^*(B)$ and $U_2^*(B^+) \setminus U_2^*(B)$. By (ix) for G^i , there are no edges in G^i (and so in G^{i+1}) between $U_2^*(B^+)$ and $V^*(G^i) \setminus U_2^*(B^+)$ in G^i and so in G^{i+1} . Thus (ix) holds for G^{i+1} .

Suppose for contradiction (x) does not hold. That is, some bowtie $B \in \mathcal{B}^{i+1}$ is not g-maximal in $G^i - (\mathcal{B}^{i+1} \setminus B)$. Note that B is g-maximal in $H^i - (\mathcal{B}^{i+1} \setminus B)$. Let $B^+ \in \mathcal{B}^i$ with $B \in \partial B^+$. By (ix), G^i has no edge between $U_2^*(B^+|H^{i-1}, \mathcal{B}^i)$ and $V^*(G^i) \setminus U_2^*(B^+|H^{i-1}, \mathcal{B}^i)$. Thus for any $v \in V^*(H^i)$, the neighbourhoods of v are the same in H^i and in G^i . Hence any vertices u, w, w' that could prevent g-maximality must also be contained in $V^*(H^i)$. Therefore B is g-maximal in $G^i - (\mathcal{B}^{i+1} \setminus B)$.

Case B: $V^*(H^i) = \emptyset$. Set $C^* = \bigcup_{j \in [i]} \phi(\mathcal{B}^j)$ and $W^* = \bigcup_{j \in [i]} W(\mathcal{B}^j)$. Let $G^* = G^i$. Clearly,

$$|W^*| \le \sum_{j \in [i]} |W(\mathcal{B}^j)| \stackrel{(\mathbf{i}), \ (\mathbf{8.2})}{\le} (384\delta^{-3})(4\delta^{-1}) < 2000\delta^{-4}.$$
(8.5)

There is a similar upper bound on $|C^*|$. Note that $\phi(G^*) = \phi(G) \setminus C^*$ by (viii). Moreover, by (v),

$$\delta_{\text{mon}}(G^*) \ge d - |W^*| \stackrel{(8.5)}{\ge} d - 2000\delta^{-4}.$$

Consider a rainbow path P in G^* with $|V(P)| \ge 3$ and $S \subseteq V^*(G) \setminus V(P)$ such that $|V(P) \cup S| \le d/4$. We shall find a rainbow cycle C^* in $G \setminus S$, which closes P with $\phi(P) \subseteq \phi(C^*) \subseteq \phi(P) \cup \phi(G) \setminus \phi(G^*)$. **Claim 8.8.** There exists $j \in [i]$ and $B \in \mathcal{B}_1^j \cup \mathcal{B}_2^j$ such that $int(P) \subseteq U_2^*(B)$.

Proof of claim. Let $v \in int(P) \subseteq V^*(G^*) \subseteq V^*(G)$. Note that $V^*(G) = V^*(H^0) \supseteq \cdots \supseteq V^*(H^i) = \emptyset$. Let $j \in [i]$ be such that $v \in V^*(H^j)$ but $v \notin V^*(H^{j+1})$. By (vii), there exists $B \in \mathcal{B}_1^j \cup \mathcal{B}_2^j$ such that $v \in U_2^*(B)$. By (x), B is g-maximal in $G^{j-1} - (\mathcal{B}^j \setminus B)$. Therefore Lemma 8.2(ii) implies that $int(P) \subseteq U_2^*(B)$.

Let B and j be as given by the claim. Let $P = x_1 \dots x_\ell$ with $\phi(x_1 x_2) = 2$ and $\phi(x_{\ell-1} x_\ell) = \ell$. Let $H' = H^{j-1} - (\mathcal{B}^j \setminus B)$.

Case B(i): $B \in \mathcal{B}_1^j$. Note that $\operatorname{int}(P) \subseteq U_2^*(B) \subseteq V^*(H^{j-1})$. Moreover, $|\operatorname{int}(P) \cup S| < d/4$. Let $U_2 = U_2(B)$, so $|U_2| \leq 3d/4$ as $B \in \mathcal{B}_1^j$. By Lemma 8.2(i), $d_{2,H'}(x_2, \overline{U_2}) < g$. Then by (vi),

$$d_{2,H'}(x_2, U_2) > \delta_{\text{mon}}^* \left(H^{j-1} \right) - g - |W^*| \stackrel{(8.5)}{\geq} d - 2g(j-1) - g - 2000\delta^{-4} \ge d/2.$$

Similarly, we have $d_{\ell,H'}(x_{\ell-1}, U_2) \ge d/2$. We deduce that

$$\begin{aligned} |N_{2,H'}(x_2, U_2) \cap N_{\ell,H'}(x_{\ell-1}, U_2) \setminus (S \cup V(P))| \\ \geq d_{2,H'}(x_2, U_2) + d_{\ell,H'}(x_{\ell-1}, U_2) - |U_2| - |S \cup V(P)| \\ > d/2 + d/2 - 3d/4 - d/4 = 0. \end{aligned}$$

Pick $x \in (N_{2,H'}(x_2, U_2) \cap N_{\ell,H'}(x_{\ell-1}, U_2)) \setminus (S \cup V(P))$. Then $C^* = xx_2 \cdots x_{\ell-1}x$ is a rainbow cycle with $\phi(C^*) = \phi(P)$.

Case B(ii): $B \in \mathcal{B}_2^j$. There exists $B' \in \mathcal{B}^{j-1}$ with $\partial B' = \{B\}$. Let $B = (v, C_1, W_1, C_2, W_2)$. Furthermore, $V^*(G^j) \cap U_2^*(B') = U_1^*(B) = U_2^*(B)$. Since P is in G^* , we have $\operatorname{int}(P) \subseteq V^*(G^*)$. Note that B is g-maximal in H'. We deduce that

$$\begin{aligned} d_{2,H'}(x_2, U_1(B|H^{j-1}, \mathcal{B}^j)) &\geq \delta^*_{\text{mon}}(H^j) - d_{2,H'}(x_2, U_1(B)) \\ (\text{vi}), \text{ Lemma } \underbrace{8.2(\mathbf{i})}_{\geq d-2gj-g} \geq d - (2i+1)g \overset{(8.2),(8.5)}{\geq} |S \cup V(P)| + |V(G) \setminus V(G^*)|. \end{aligned}$$

Pick $y_1 \in U_1(B) \cap (V(G^*) \setminus (S \cup V(P)))$ such that $\phi(y_1x_2) = 2$. Similarly, we can find a distinct vertex $y_2 \in U_2(B) \cap (V(G^*) \setminus (S \cup V(P)))$ such that $\phi(y_2x_{\ell-1}) = \ell$. Let $P_{y_1,v}$ be the rainbow path from y_1 to v with $\operatorname{int}(P_{y_1,v}) \subseteq W_1$ and $\phi(P_{y_1,v}) \subseteq C_1$; and P_{v,y_2} be a rainbow path from y_2 to v with $\operatorname{int}(P_{v,y_2}) \subseteq W_2$ and $\phi(P_{v,y_2}) \subseteq C_2$. Then $C^* = y_1 P_{y_1,v} P_{v,y_2} y_2 x_{\ell-1} x_{\ell-2} \dots x_2 y_1$ is a rainbow cycle in $G \setminus S$. Note that

$$\phi(P) \subseteq \phi(C^*) \subseteq \phi(P) \cup \phi(G) \setminus \phi(G^*).$$

This concludes the proof of the lemma.

9 Concluding Remarks

In this work, we showed that there exists a monochromatic tight cycle partition of any large r-edgecoloured $K_n^{(k)}$ with a number of cycles that is a polynomial of r. For $k \ge 3$, it is easy to construct an r-edge-coloured hypergraph which needs at least r monochromatic tight cycles to partition all the vertices. We show such a construction below for completeness.

Proposition 9.1. Let $k, r \in \mathbb{N}$ such that $k \geq 2$. There exists an r-edge-coloured complete k-graph such that at least r monochromatic cycles are required for a monochromatic tight cycle partition of its vertex set.

Proof. Let H be a complete k-graph with vertex set partitioned as $V = \bigcup_{i \in [r]} V_i$, where $V_1 \neq \emptyset$, for distinct $i, j \in [r], V_i \cap V_j = \emptyset$ and for each $i \in [r] \setminus \{1\}$,

$$|V_i| > (k-1) \sum_{j \in [i-1]} |V_j|.$$
(9.1)

We colour each $e \in E(G)$ with colour $\phi(e) = \min\{i \in [r] : V_i \cap e \neq \emptyset\}$. Let \mathcal{C} be a set of vertex-disjoint monochromatic tight cycles that partition V(H). It is enough to show that for each $i \in [r]$, there is a tight cycle in \mathcal{C} coloured *i*. Clearly, a monochromatic tight cycle *C* that contains at least one vertex from V_i satisfies $\phi(C) \leq i$. Let

$$\mathcal{C}^{$$

It suffices to show that $V_i \not\subseteq V(\mathcal{C}^{<i})$. Note that each $C \in \mathcal{C}^{<i}$ satisfies $\left|V(C) \cap \bigcup_{j < i} V_j\right| \ge |V(C)|/k$ and therefore $|V(C) \cap V_i| \le (k-1) \left|V(C) \cap \bigcup_{j < i} V_j\right|$. We deduce that

$$|V(\mathcal{C}^{< i}) \cap V_i| \le \sum_{C \in \mathcal{C}^{< i}} |V(C) \cap V_i| \le (k-1) \sum_{j \in [i-1]} |V_j| \stackrel{(9,1)}{<} |V_i|.$$

This completes the proof of the proposition.

When k = 2 and $r \ge 3$, Pokrovskiy [21] showed that there exist infinitely many r-edge-coloured K_n which require at least r + 1 monochromatic cycles. Lo and Pfenninger [16] showed that for any $k \ge 3$, there are 2-edge-coloured $K_n^{(k)}$ that cannot be partitioned into two monochromatic tight cycles of distinct colours. It is interesting to consider whether the same holds for r-edge-coloured complete k-graphs when $r, k \ge 3$.

We believe that an r-edge-coloured $K_n^{(k)}$ can be partitioned into linearly many monochromatic tight cycles.

Conjecture 9.2. Let $r, k \in \mathbb{N}$ with $r, k \geq 2$. Then there exists a constant C = C(k) such that every *r*-edge-coloured $K_n^{(k)}$ can be partitioned into at most Cr monochromatic tight cycles.

If one can strengthen Lemma 5.3 to show that f(r,k) monochromatic tight cycles suffice, then our proof method would imply $O(r \log r) + 2f(r,k) + 3$ cycles partition any *r*-edge-coloured $K_n^{(k)}$. The $O(r \log r)$ term corresponds to the number of monochromatic cycles required to cover almost all the vertices.

Conjecture 9.3. For all $r, k \ge 2$, there exists C = C(k) such that any r-edge-coloured $K_n^{(k)}$ contains at most Cr cycles covering (1 - o(1))n vertices.

Note that the case k = 2 and general r is still open.

Acknowledgements

The authors would like to thank Vincent Pfenninger, Simona Boyadzhiyska and Andrea Freschi for their comments on the draft and helpful discussions.

References

- [1] P. Allen. "Covering two-edge-coloured complete graphs with two disjoint monochromatic cycles". Combin. Probab. Comput. 17.4 (2008), 471–486. DOI: 10.1017/S0963548308009164.
- P. Allen, J. Böttcher, O. Cooley, and R. Mycroft. "Tight cycles and regular slices in dense hypergraphs". J. Combin. Theory Ser. A 149 (2017), 30–100. DOI: 10.1016/j.jcta.2017.01.003.
- S. Bessy and S. Thomassé. "Partitioning a graph into a cycle and an anticycle, a proof of Lehel's conjecture". J. Combin. Theory Ser. B 100.2 (2010), 176–180. DOI: 10.1016/j.jctb.2009.07.001.
- [4] S. Bustamante, J. Corsten, N. Frankl, A. Pokrovskiy, and J. Skokan. "Partitioning edge-colored hypergraphs into few monochromatic tight cycles". *SIAM J. Discrete Math.* 34.2 (2020), 1460–1471. DOI: 10.1137/19M1269786.
- [5] S. Bustamante, H. Hàn, and M. Stein. "Almost partitioning 2-colored complete 3-uniform hypergraphs into two monochromatic tight or loose cycles". J. Graph Theory 91.1 (2019), 5–15. DOI: 10.1002/jgt.22417.
- [6] D. Conlon, J. Fox, and B. Sudakov. "Ramsey numbers of sparse hypergraphs". Random Structures Algorithms 35.1 (2009), 1–14. DOI: 10.1002/rsa.20260.
- [7] P. Erdős, A. Gyárfás, and L. Pyber. "Vertex coverings by monochromatic cycles and trees". J. Combin. Theory Ser. B 51.1 (1991), 90–95. DOI: 10.1016/0095-8956(91)90007-7.
- S. Fujita, H. Liu, and C. Magnant. "Monochromatic structures in edge-coloured graphs and hypergraphs-a survey". *Int. J. Graph Theory Applications* 1.1 (2015), 3–56. URL: https://www.mililink.com/upload/article/813658924IJGTA%20v1i1%20
- [9] A. Gyárfás. "Vertex covers by monochromatic pieces—a survey of results and problems". Discrete Math. 339.7 (2016), 1970–1977. DOI: 10.1016/j.disc.2015.07.007.
- [10] A. Gyárfás, J. Lehel, J. Nešetřil, V. Rödl, R. H. Schelp, and Zs. Tuza. "Local k-colorings of graphs and hypergraphs". J. Combin. Theory Ser. B 43.2 (1987), 127–139. DOI: 10.1016/0095-8956(87)90017-7.
- [11] A. Gyárfás, M. Ruszinkó, G. N. Sárközy, and E. Szemerédi. "An improved bound for the monochromatic cycle partition number". J. Combin. Theory Ser. B 96.6 (2006), 855–873. DOI: 10.1016/j.jctb.2006.02.007.
- [12] A. Gyárfás and G. N. Sárközy. "Monochromatic path and cycle partitions in hypergraphs". Electron. J. Combin. 20.1 (2013), Paper 18. DOI: 10.37236/2631.
- [13] S. Janson, T. Luczak, and A. Ruciński. "Random graphs". Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, 2000. DOI: 10.1002/9781118032718.
- [14] D. Korándi, R. Lang, S. Letzter, and A. Pokrovskiy. "Minimum degree conditions for monochromatic cycle partitioning". J. Combin. Theory Ser. B 146 (2021), 96–123. DOI: 10.1016/j.jctb.2020.07.005.
- [15] A. Lo and V. Pfenninger. "Almost partitioning every 2-edge-coloured complete k-graph into k monochromatic tight cycles". Innovations in Graph Theory, to appear.
- [16] A. Lo and V. Pfenninger. "Towards Lehel's conjecture for 4-uniform tight cycles". *Electron. J. Combin.* 30.1 (2023), Paper No. 1.13. DOI: 10.37236/10604.
- [17] T. Luczak. " $R(C_n, C_n, C_n) \le (4+o(1))n$ ". J. Combin. Theory Ser. B 75.2 (1999), 174–187. DOI: 10.1006/jctb.1998.1874.
- [18] T. Luczak, V. Rödl, and E. Szemerédi. "Partitioning two-coloured complete graphs into two monochromatic cycles". Combin. Probab. Comput. 7.4 (1998), 423–436. DOI: 10.1017/S0963548398003599.
- [19] V. Pfenninger. "On k-uniform tight cycles: the Ramsey number for $C_{kn}^{(k)}$ and an approximate Lehel's conjecture". ArXiv: 2406.14468. Preprint (2024).
- [20] A. Pokrovskiy. "Partitioning a graph into a cycle and a sparse graph". Discrete Math. 346.1 (2023), Paper No. 113161, 21. DOI: 10.1016/j.disc.2022.113161.
- [21] A. Pokrovskiy. "Partitioning edge-coloured complete graphs into monochromatic cycles and paths". J. Combin. Theory Ser. B 106 (2014), 70–97. DOI: 10.1016/j.jctb.2014.01.003.
- [22] G. N. Sárközy. "Improved monochromatic loose cycle partitions in hypergraphs". *Discrete Math.* 334 (2014), 52–62. DOI: 10.1016/j.disc.2014.06.025.
- [23] G. N. Sárközy. "Monochromatic partitions in local edge colorings". Acta Math. Hungar. 161.2 (2020), 412–426. DOI: 10.1007/s10474-020-01054-1.