

IMPROVED PERFORMANCE GUARANTEES FOR TUKEY’S MEDIAN

STANISLAV MINSKER AND YINAN SHEN

ABSTRACT. Is there a natural way to order data in dimension greater than one? The approach based on the notion of data depth, often associated with the name of John Tukey, is among the most popular. Tukey’s depth has found applications in robust statistics, graph theory, and the study of elections and social choice. We present improved performance guarantees for empirical Tukey’s median, a deepest point associated with the given sample, when the data-generating distribution is elliptically symmetric and possibly anisotropic. Some of our results remain valid in the class of affine equivariant estimators. As a corollary of our bounds, we show that the diameter of the set of all empirical Tukey’s medians scales like $o(n^{-1/2})$ where n is the sample size.

1. INTRODUCTION

The fundamental notions of order statistics and ranks form a basis of many inferential procedures. They are the backbone of methods in robust statistics that studies stability of algorithms under perturbations of the data (Huber and Ronchetti, 2009). As the Euclidean space of dimension 2 and higher lacks the canonical order, the “canonical” versions of order statistics and ranks are difficult to construct. Many useful versions have been proposed over the years: examples include the componentwise ranks (Hodges, 1955), spatial ranks and quantiles (Brown, 1983; Koltchinskii, 1997), Mahalanobis ranks (Hallin and Paindaveine, 2002), and recently introduced Monge-Kantorovich ranks and quantiles (Chernozhukov et al., 2017). In this paper, we will focus on the notion of *data depth* that gives rise to the depth-based ranks and quantiles, see (Serfling, 2002) for an overview. The idea of depth and the associated ordering with respect to a probability measure P on \mathbb{R} goes back to Hotelling (1929): given $z \in \mathbb{R}$, its *depth* is defined as the minimum among $P(-\infty, z]$ and $P[z, \infty)$. Hodges (1955) and Tukey (1975) extended this idea to \mathbb{R}^2 , and finally Donoho (1982); Donoho and Gasko (1992) formalized the general notion of depth of a point $z \in \mathbb{R}^d$ with respect to a probability measure P as the infimum of all univariate depth evaluated over projections of P on the lines passing through z . Equivalently, Tukey’s (or half-space) depth of z is

$$(1) \quad D_P(z) = \inf_{u \in S^{d-1}} P(H(z, u)),$$

where $H(z, u) = \{x \in \mathbb{R}^d : x^T u \geq z^T u\}$ is the half-space passing through z in direction u , and S^{d-1} is the unit sphere in \mathbb{R}^d with respect to the Euclidean norm $\|\cdot\|_2$. Let us remark that the notion of half-space depth and its applications appear in other areas, for example in graph theory (Small, 1997; Cerdeira and Silva, 2021), in convex geometry in relation to the convex floating boudies (Nagy et al., 2019), and the theory of social choice in economics under the name of the “min-max majority” (Caplin and Nalebuff, 1988; Nehring and Puppe, 2023). We refer the reader to the habilitation thesis of S. Nagy (2022) for an excellent overview of the history and recent developments in the theoretical and algorithmic aspects of Tukey’s depth.

If P is such that $P(\partial H) = 0$ for the boundary ∂H of any closed half-space H , then the infimum in (1) is attained (Massé, 2004, Proposition 4.5). For example, this is the case for

S. Minsker acknowledges support by the National Science Foundation grant DMS CAREER-2045068.

distributions that are absolutely continuous with respect to the Lebesgue measure. Of particular interest are the points of maximal depth, that is,

$$\mu_* := \mu_*(P) \in \arg \max D_P(z).$$

The set of deepest points with respect to P will be referred to as Tukey's median set, or simply Tukey's set, while the barycenter of this set is Tukey's median; when $d = 1$, it coincides with the standard median. According to Propositions 7 and 9 in (Rousseeuw and Ruts, 1999), Tukey's median always exists, and $\alpha_* := D_P(\mu_*) \geq \frac{1}{d+1}$ for any P . For distributions possessing symmetry properties, α_* can be much larger. For example, if P is halfspace-symmetric¹ and absolutely continuous with respect to the Lebesgue measure, then its center of symmetry coincides with Tukey's median, and its depth α_* equals $\frac{1}{2}$. The multivariate normal distribution, and more generally all elliptically symmetric distributions, are known to satisfy this property. Whenever α_* is large, Tukey's median also has a high breakdown point, and therefore features strong robustness properties – for instance, its breakdown point equals $\frac{1}{3}$ for halfspace-symmetric distributions, see Proposition 1 in (Chen, 1995) for the precise statement. The upper-level sets of half-space depth, defined as

$$R_P(\alpha) := \{z \in \mathbb{R}^d : D_P(z) \geq \alpha_* - \alpha\}, \quad \alpha < \alpha_*,$$

are convex and compact. These sets are often called the *central regions* (Serfling, 2002) and their boundaries – the quantile surfaces or depth contours (Liu et al., 1999).

Let $X_1, \dots, X_n \in \mathbb{R}^d$ be a sequence of independent copies of a random vector X with distribution P . The empirical measure P_n^X is a discrete measure with atoms X_1, \dots, X_n having weight $1/n$ each. In particular, for any Borel measurable set $A \subseteq \mathbb{R}^d$,

$$P_n^X(A) := \frac{1}{n} \sum_{j=1}^n I_A(X_j), \quad \text{where } I_A(z) = \begin{cases} 1, & z \in A, \\ 0, & z \notin A. \end{cases}$$

The *empirical depth* corresponding to P_n^X will be denoted $D_n(z)$, and it yields a natural way to order the points in a sample, giving rise to the depth-based ranks. It also admits the following convenient geometric interpretation: $nD_n(z)$ equals the cardinality of a smallest subset $S \subset \{X_1, \dots, X_n\}$ such that z is not in the convex hull of $\{X_1, \dots, X_n\} \setminus S$, with the convention that the cardinality of the empty set is 0. The empirical version of Tukey's set is defined as the set of points of maximal empirical depth,

$$(2) \quad \hat{\mu}_n := \hat{\mu}_n(X_1, \dots, X_n) \in \arg \max_{z \in \mathbb{R}^d} D_n(z).$$

The barycenter of this set is known as the empirical Tukey's median, and it will also be denoted $\hat{\mu}_n$ with slight abuse of notation. The definition is illustrated in Figure 1 that displays the (empirical) depth contours, upper-level sets and the Tukey's median corresponding to a sample of size $n = 100$ from the isotropic bivariate Gaussian distribution as well as the isotropic bivariate Student's t distribution with 2.1 degrees freedom. Figure 2 shows that the empirical contours become smoother and approach their population counterparts as the sample size grows.

Even when μ_* is unique, the Tukey's set can have a non-empty interior. For instance, if $d = 3$, then the set $\arg \max_{z \in \mathbb{R}^d} D_n(z)$ can be a single point, an interval, or a convex polytope, see example 3.1 in (Pokorný et al., 2024). This raises a natural question: *what is the typical diameter of this set?* In the univariate case ($d = 1$), the sample median is unique when n is odd, and when n is even, the diameter equals $L_n = \left| X_{(\frac{n}{2}+1)} - X_{(\frac{n}{2})} \right|$. Under mild assumptions and using the classical results on order statistics (Rényi, 1953), the expected value of L_n can be shown to be of order c/n for a constant c that depends on the distribution. To the best of our

¹The distribution of a random vector X is halfspace-symmetric with respect to some $\mu \in \mathbb{R}^d$ if the random variables $u^T(X - \mu)$ and $u^T(\mu - X)$ are equidistributed for all $u \in S^{d-1}$.


(A) $n = 100$, standard Gaussian(B) $n = 100$, Student's t with $\nu = 2.1$ d.f.

FIGURE 1. Depth contours: lighter-colored regions correspond to higher depth. Empirical Tukey's median is marked with a circle and the sample mean – with a cross

knowledge, the question was open for $d \geq 2$. We make a step towards answering it and prove (see Corollary 2.3) that with high probability, diameter of empirical Tukey's set decays faster than $n^{-1/2}$.

Among the attractive properties of the half-space depth is the *affine invariance*: for any affine map $z \mapsto T(z) = Mz + b$ such that $M \in \mathbb{R}^{d \times d}$ is non-singular and $b \in \mathbb{R}^d$,

$$D_{P \circ T^{-1}}(Mz + b) = D_P(z), \quad \forall z \in \mathbb{R}^d,$$

where $P \circ T^{-1}(A) = P(T^{-1}(A))$. This implies that

$$\mu_*(P \circ T^{-1}) = M\mu_*(P) + b,$$

so that Tukey's median “respects” affine transformations. For example, if X has mean μ and the covariance matrix $\Sigma = \mathbb{E}(X - \mu)(X - \mu)^T$ is non-degenerate, then

$$\hat{\mu}_n(X_1, \dots, X_n) = \Sigma^{1/2} \hat{\mu}_n(Z_1, \dots, Z_n) + \mu,$$

where $Z_j = \Sigma^{-1/2}(X_j - \mu)$ has centered isotropic distribution (i.e. the covariance of Z_1 is the identity matrix). Due to this property, existing literature that investigates the proximity between the population median μ and its data-dependent counterpart $\hat{\mu}_n$ often focuses on the estimation error measured in the Mahalanobis distance $\|\Sigma^{-1/2}(\hat{\mu}_n(X_1, \dots, X_n) - \mu)\|_2$. Since Σ is typically unknown, one may instead prefer to measure the error with respect to the usual Euclidean distance $\|\hat{\mu}_n - \mu\|_2$.

In this paper, we establish certain optimality properties of Tukey's median. Specifically, we prove (see Theorem 2.1) that the size of the error $\|\hat{\mu}_n - \mu\|_2$ is controlled by the *effective rank* of Σ defined as the ratio of the trace $\text{tr}(\Sigma)$ and the spectral norm $\|\Sigma\|$,

$$r(\Sigma) = \frac{\text{tr}(\Sigma)}{\|\Sigma\|},$$

as opposed to the ambient dimension d . Finally, we investigate robustness properties of Tukey's median and the effects of adversarial contamination on the error. While this question has been answered for isotropic distributions (equivalently, for the error with respect to the Mahalanobis norm, see (Chen et al., 2018)), the case of Euclidean distance poses new challenges. We give a partial answer in Corollary 2.4. The remaining open problems are discussed in the section 3.


(A) $n = 50$, standard Gaussian(B) $n = 500$, standard Gaussian

FIGURE 2. Empirical depth contours approach their population limit. The blue circle denotes Tukey's median.

1.1. Notation. Given $z_1, z_2 \in \mathbb{R}^d$, $\langle z_1, z_2 \rangle$ will stand for the standard inner product and $\|\cdot\|_2$ for the associated Euclidean norm. For a matrix $A \in \mathbb{R}^{d \times d}$, $\text{tr}(A)$ represents its trace and $\|A\|$ - its spectral (operator) norm. I_d will denote the $d \times d$ identity matrix.

We will employ the standard small-o and big-O asymptotic notation throughout the paper. Given two sequences $\{a_j\}_{j \geq 1}$, $\{b_j\}_{j \geq 1} \subset \mathbb{R}_+$, we will say that $a_j \ll b_j$ if $a_j = o(b_j)$ as $j \rightarrow \infty$. Finally, we will write $o_P(1)$ to denote a sequence of random variables ξ_j , $j \geq 1$ that converge to 0 in probability as $j \rightarrow \infty$. Indicator function of an event \mathcal{E} will be denoted $I_{\mathcal{E}}$ or $I\{\mathcal{E}\}$. Additional notation will be introduced on demand.

2. MAIN RESULTS

Assume that a fraction of the data has been replaced by arbitrary values, that is, we observe Y_1, \dots, Y_n where $\frac{1}{n} \sum_{j=1}^n I\{Y_j \neq X_j\} = \varepsilon$.² In a scenario when X_1, \dots, X_n have multivariate normal distribution $N(\mu, \Sigma)$ such that Σ is non-singular, Chen et al. (2018) showed that whenever $\varepsilon < 1/5$ and $\frac{d}{n} + \frac{t}{n} \leq c$,

$$\|\Sigma^{-1/2}(\hat{\mu}_n(Y_1, \dots, Y_n) - \mu)\|_2 \leq C \left(\sqrt{\frac{d+t}{n}} + \varepsilon \right),$$

holds with probability at least $1 - e^{-t}$, where $c, C > 0$ are absolute constants. The key feature of this inequality is the fact that it implies optimal dependence of the error on the contamination proportion ε ; Tukey's median is among the first estimators known to possess this property. This robustness property of Tukey's median is illustrated in figure 3 where it is juxtaposed with the sample mean. Recently, Minasyan and Zhivotovskiy (2023) proved that there exists an estimator $\tilde{\mu}_n$ that satisfies the inequality

$$(3) \quad \|\tilde{\mu}_n(Y_1, \dots, Y_n) - \mu\|_2 \leq C \|\Sigma\|^{1/2} \left(\sqrt{\frac{r(\Sigma) + t}{n}} + \varepsilon \right)$$

with probability at least $1 - e^{-t}$, whenever $\frac{r(\Sigma) + t}{n} \leq c$. Furthermore, if $\|\Sigma\|$ and $r(\Sigma)$ are known up to some absolute multiplicative factor, then the requirement $\frac{r(\Sigma) + t}{n} \leq c$ can be dropped. This inequality exhibits optimal dependence on the ‘‘dimensional’’ characteristic expressed via the

²This is commonly referred to as the *adversarial contamination framework*, see (Diakonikolas and Kane, 2023)

effective rank as opposed to the ambient dimension that can be much larger, as well as correct dependence on the degree of contamination ε . Deviation inequalities of type (3) are often referred to as *sub-Gaussian guarantees*, since they hold, for $\varepsilon = 0$, for the sample mean of i.i.d. Gaussian random vectors due to the Borell-TIS inequality (Borell, 1975; Tsirelson et al., 1976). Whether Tukey's median admits a bound of the form (3) remained an open question: specifically, it was unknown whether d can be replaced by $r(\Sigma)$. We will give a (partial) affirmative answer below. As this question is mainly interesting in the situation when d is large, we will assume that $d \geq 2$ in the remainder of the paper.

2.1. Contamination-free framework. Our starting point is the contamination-free scenario corresponding to $\varepsilon = 0$. Recall that a random vector $X \in \mathbb{R}^d$ has elliptically symmetric distribution $\mathcal{E}(\mu, \Sigma, F)$, denoted $X \sim \mathcal{E}(\mu, \Sigma, F)$ in the sequel, if

$$(4) \quad X \stackrel{d}{=} \eta \cdot BU + \mu,$$

where $\stackrel{d}{=}$ denotes equality in distribution, η is a nonnegative random variable with cumulative distribution function F , B is a fixed $d \times d$ matrix such that $\Sigma = BB^T$, and $U \in \mathbb{R}^d$ is uniformly distributed over the unit sphere \mathbb{S}^{d-1} and independent of η . The distribution $\mathcal{E}(\mu, \Sigma, F)$ is well defined, as if $B_1 B_1^T = B_2 B_2^T$, then there exists an orthogonal matrix Q such that $B_1 = B_2 Q$, and $QU \stackrel{d}{=} U$. To avoid such ambiguity, we will allow B to be any matrix satisfying $BB^T = \Sigma$. Whenever $\mathbb{E}\eta^2 < \infty$, we will also assume that η is scaled so that $\mathbb{E}\eta^2 = 1$, in which case the covariance of X exists and is equal to Σ .

A useful property of elliptically symmetric distributions that is evident from the definition is the following: for every orthogonal matrix Q , $Q\Sigma^{-1/2}(X - \mu)$ has the same distribution as $\Sigma^{-1/2}(X - \mu)$. In other words, $\Sigma^{-1/2}(X - \mu)$ is spherically symmetric. We are ready to state our first result that essentially hinges on this fact.

Theorem 2.1. *Let X_1, \dots, X_n be i.i.d. copies of $X \sim \mathcal{E}(\mu, \Sigma, F)$ where Σ is non-singular. Let $\tilde{\mu}_n$ be any affine equivariant estimator of μ . Suppose that*

$$\|\Sigma^{-1/2}(\tilde{\mu}_n - \mu)\|_2 \leq e(n, t, d)$$

holds with probability at least $1 - p(t)$. Then

$$\|\tilde{\mu}_n - \mu\|_2 \leq e(n, t, d) \|\Sigma\|^{1/2} \left(\sqrt{\frac{r(\Sigma)}{d}} + \sqrt{\frac{2t}{d}} \right)$$

with probability at least $1 - p(t) - e^{-t}$.

In typical scenarios,

$$e(n, t, d) \asymp \sqrt{\frac{d}{n}} + \sqrt{\frac{t}{n}} \text{ and } p(t) = e^{-t},$$

whence

$$(5) \quad \|\tilde{\mu}_n - \mu\|_2 \leq C \left(\sqrt{\frac{d}{n}} + \sqrt{\frac{t}{n}} \right) \sqrt{\|\Sigma\|} \left(\sqrt{\frac{r(\Sigma)}{d}} + \sqrt{\frac{t}{d}} \right) \leq C_1 \sqrt{\|\Sigma\|} \left(\sqrt{\frac{r(\Sigma)}{n}} + \sqrt{\frac{t}{n}} \right)$$

with probability at least $1 - 2e^{-t}$ whenever $t \leq d$. The proof of the theorem is given in appendix A. Next, we state the corollaries for Tukey's median and the Stahel-Donoho estimator, another well known affine-equivariant robust estimator of location (see (Stahel, 1981; Donoho, 1982) for its definition and properties).

Corollary 2.1 (Tukey Median). *Let X_1, \dots, X_n be i.i.d. copies of $X \sim N(\mu, \Sigma)$ where Σ is non-singular, and let $\hat{\mu}_n$ be the Tukey's median, the barycenter of the set $\arg \max_{z \in \mathbb{R}^d} D_n(z)$. Then there exist absolute constants $c, C > 0$ such that*

$$\|\hat{\mu}_n - \mu\|_2 \leq C \left(\sqrt{\frac{\text{tr}(\Sigma)}{n}} + \sqrt{\|\Sigma\|} \sqrt{\frac{t}{n}} \right)$$

with probability at least $1 - 2e^{-t}$ whenever $\frac{d}{n} + \frac{t}{n} \leq c$.

Proof. Theorem 2.1 in (Chen et al., 2018) states that

$$e(n, t, d) = \sqrt{\|\Sigma\|} \left(\sqrt{\frac{d}{n}} + \sqrt{\frac{t}{n}} \right),$$

and $p(t) = e^{-t}$ whenever $\frac{d}{n} + \frac{t}{n} \leq c$. Inequality (5) thus implies the desired result whenever $t \leq d$. If $t > d$, it suffices to show that under the stated assumptions, $\|\Sigma^{-1/2}(\hat{\mu}_n - \mu)\| \leq C\sqrt{\frac{t}{n}}$ with probability at least $1 - 2e^{-t}$, which follows directly from Theorem 2.1 in (Chen et al., 2018). \square

Remark 2.1. *The bound of Corollary 2.1 remains valid whenever $d \geq n+1$: indeed, in this case with probability 1 the convex hull of X_1, \dots, X_n is a simplex with n vertices, and every point in the convex hull has (maximal) depth $1/n$, hence $\hat{\mu}_n$ coincides with the sample mean.*

Up to the values of numerical constants, the same result is valid for the Stahel-Donoho estimator.

Corollary 2.2 (Stahel-Donoho Estimators). *Let X_1, \dots, X_n be i.i.d. copies of $X \sim N(\mu, \Sigma)$ where Σ is non-singular, and let $\tilde{\mu}_n$ be the barycenter of the set of Stahel-Donoho estimators. Then there exist absolute constants $c, C > 0$ such that*

$$\|\tilde{\mu}_n - \mu\|_2 \leq C \left(\sqrt{\frac{\text{tr}(\Sigma)}{n}} + \sqrt{\|\Sigma\|} \sqrt{\frac{t}{n}} \right)$$

with probability at least $1 - 2e^{-t}$ whenever $\frac{d}{n} + \frac{t}{n} \leq c$.

Proof. The argument repeats the proof of Corollary 2.1 *mutatis mutandis*, where Theorem 1 in (Depersin and Lecué, 2023) plays the role of Theorem 2.1 in (Chen et al., 2018). \square

Remark 2.2. *Results of both corollaries hold not just for Gaussian measures but more generally for a wide class of elliptically symmetric distributions. Indeed, the arguments in (Chen et al., 2018; Depersin and Lecué, 2023) leading to the bounds for $e(n, t, d)$ remain valid if we assume that $X \sim \mathcal{E}(\mu, \Sigma, F)$ is such that the density $q(t)$ of one-dimensional projections*

$$\langle \Sigma^{-1/2}(X - \mu), v \rangle = \eta \langle U, v \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean inner product, is continuous and strictly positive at the origin (also see remark 2.2 in (Chen et al., 2018)). For example, multivariate t -distribution satisfies this assumption.

2.2. Performance guarantees in the adversarial contamination framework. Recall the adversarial contamination framework that was introduced in the beginning of section 2. All results stated below will be specific to Tukey's median as we rely on the existing rich literature on the subject. However, we expect our claims to be valid for other affine equivariant estimators such as the Stahel-Donoho estimator mentioned previously.

In what follows, we will let $\hat{\mu}_n^X$ be the empirical Tukey's median based on the uncontaminated data X_1, \dots, X_n and $\hat{\mu}_n^Y$ – its counterpart based on the sample Y_1, \dots, Y_n . We will let

$$D_n^X(z) = \inf_{\|u\|_2=1} P_n^X H(z, u)$$




FIGURE 3. Contaminated sample with contamination proportion $\varepsilon = 0.1$. Circle mark denotes Tukey median and the cross represents the mean.

stand for the empirical depth of a point $z \in \mathbb{R}^d$, where the empirical distribution P_n^X is based on X_1, \dots, X_n , and $\hat{d}_n := D_n^X(\hat{\mu}_n^X)$ be the maximal (empirical) depth. Consider the set

$$R_n(\delta) := \{z \in \mathbb{R}^d : D_n^X(z) \geq \hat{d}_n - \delta\}$$

of “ $\hat{d}_n - \delta$ -deep” points. $R_n(\delta)$ is a convex polytope whose facets belong to the hyperplanes defined by the elements of the sample. Observe that the empirical depth $D_n^Y(z)$ (based on the contaminated dataset) of any point $z \notin R_n(\delta)$ satisfies the inequality

$$D_n^Y(z) < \varepsilon + \hat{d}_n - \delta.$$

Indeed,

$$(6) \quad D_n^Y(z) = \inf_{\|u\|_2=1} P_n^Y H(z, u) \leq \underbrace{\inf_{\|u\|_2=1} P_n^X H(z, u)}_{< \hat{d}_n - \delta} + \underbrace{\sup_{\|u\|_2=1} |(P_n^Y - P_n^X) H(z, u)|}_{\leq \varepsilon}.$$

The latter does not exceed $\hat{d}_n - \varepsilon$ whenever $\delta > 2\varepsilon$. Since $D_n^Y(\hat{\mu}_n)$ is at least $(\hat{d}_n - \varepsilon)$ due to the inequality in the spirit of (6), we deduce the following lemma.

Lemma 2.1. *Let $\hat{\mu}_N^Y \in \arg \max_{z \in \mathbb{R}^d} D_n^Y(z)$. Then*

$$\hat{\mu}_N^Y \in R_n(2\varepsilon).$$

An immediate corollary of this lemma is that in the framework of Corollary 2.1,

$$(7) \quad \|\widehat{\mu}_n^Y - \mu\|_2 \leq \|\widehat{\mu}_n^X - \mu\|_2 + \text{diam}(R_n(2\varepsilon)),$$

where $\text{diam}(A)$ stands for the diameter of a set $A \subseteq \mathbb{R}^d$. Whenever ε is large, meaning that $1/5 > \varepsilon \geq K\sqrt{\frac{d}{n}}$ for some constant $K > 0$, results by Chen et al. (2018) imply that

$$\text{diam}(R_n(2\varepsilon)) \leq C(K)\sqrt{\|\Sigma\|} \left(\sqrt{\frac{t}{n}} + \varepsilon \right)$$

with probability at least $1 - 2e^{-t}$ for $t \leq c_1 n$. Therefore, for such values of ε and t ,

$$(8) \quad \|\widehat{\mu}_n^Y - \mu\|_2 \leq C(K) \left(\sqrt{\frac{\text{tr}(\Sigma)}{n}} + \sqrt{\|\Sigma\|} \left(\sqrt{\frac{t}{n}} + \varepsilon \right) \right)$$

with probability at least $1 - 2e^{-t}$.

The problem is less straightforward for $\varepsilon := \varepsilon_n \ll \sqrt{\frac{d}{n}}$. In the remainder of this section, we will assume that d and $r(\Sigma)$ are fixed, where d is potentially much larger than $r(\Sigma)$, and $n \rightarrow \infty$. The following theorem is the main result of this section.

Theorem 2.2. *Assume that X_1, \dots, X_n are i.i.d. random vectors sampled from the isotropic normal distribution $N(\mu, I_d)$. There exists an event Θ of probability at least $1 - 2e^{-t} - \frac{2}{n}$ such that for any sequence $\varepsilon_n = o(n^{-1/2})$, the diameter of the set $R_n(\varepsilon_n)$ satisfies*

$$\text{diam}(R_n(\varepsilon_n)) \cdot I_\Theta = o_P(n^{-1/2}),$$

where I_Θ is the indicator of the event Θ .

Before turning to the proof of this result, we will state its immediate corollaries.

Corollary 2.3 (Tukey's median set). *The diameter of the set $R_n(0)$ satisfies*

$$\text{diam}(R_n(0)) = o_P(n^{-1/2}).$$

Corollary 2.4 (Adversarial Contamination). *Assume that X_1, \dots, X_n are i.i.d. random vectors sampled from the multivariate normal distribution $N(\mu, \Sigma)$. In the adversarial contamination framework with $\varepsilon_n = o(n^{-1/2})$,*

$$\|\widehat{\mu}_n^Y - \mu\|_2 \leq C \left(\sqrt{\frac{\text{tr}(\Sigma)}{n}} + \sqrt{\|\Sigma\|} \sqrt{\frac{t}{n}} \right)$$

with probability at least $1 - 2e^{-t} - o(1)$, where $C > 0$ is an absolute constant.

Proof. First, note that due to affine invariance of the depth function $D_n(z)$, the isotropic and non-isotropic depth regions are related via

$$R_n(2\varepsilon_n; X_1, \dots, X_n) = \mu + \Sigma^{1/2} R_n(2\varepsilon_n; \Sigma^{-1/2}(X_1 - \mu), \dots, \Sigma^{-1/2}(X_n - \mu)).$$

Therefore,

$$\text{diam}(R_n(2\varepsilon_n; X_1, \dots, X_n)) \leq \|\Sigma\|^{1/2} \text{diam}\left(R_n(2\varepsilon_n; \Sigma^{-1/2}(X_1 - \mu), \dots, \Sigma^{-1/2}(X_n - \mu))\right).$$

The latter is $o_P(n^{-1/2})$ in view of Theorem 2.2. The result follows in view of the inequality (7) combined with Theorems 2.2 and 2.1. \square

Of course, for $\varepsilon_n = o(n^{-1/2})$, the corollary trivially yields that

$$\|\hat{\mu}_n^Y - \mu\|_2 \leq C \left(\sqrt{\frac{\text{tr}(\Sigma)}{n}} + \sqrt{\|\Sigma\|} \left(\sqrt{\frac{t}{n}} + \varepsilon_n \right) \right)$$

with high probability. In view of (8), this inequality is also valid for large values of ε_n , above $K\sqrt{\frac{d}{n}}$. Whether a bound of this form also holds whenever $\varepsilon_n \asymp \sqrt{\frac{r(\Sigma)}{n}}$ (or more generally in the regime $C_1\sqrt{\frac{r(\Sigma)}{n}} \leq \varepsilon_n \leq C_2\sqrt{\frac{d}{n}}$, where $C_1, C_2 > 0$ are some fixed constants), remains an open problem. Next, we turn our attention to the proof of Theorem 2.2.

Proof of Theorem 2.2. Everywhere below, we will assume without loss of generality that $\mu = 0$. The idea of the proof relies on the strong approximation approximation of the depth process via a P-Brownian bridge. We will need several preliminary facts. Let $C > 0$ be a sufficiently large absolute constants. Recall that in view of results by Chen et al. (2018),

$$\text{diam}(R_n(2\varepsilon_n)) \leq C_1 \sqrt{\frac{d+t}{n}},$$

with probability at least $1 - e^{-t}$, and in particular on this event $\|\hat{\mu}_n\|_2 \leq C_1 \sqrt{\frac{d+t}{n}}$. Next, define

$$W_n(z) = \sqrt{n} \inf_{\|v\|_2=1} \left(P_n^X H \left(\frac{z}{\sqrt{n}}, v \right) - \frac{1}{2} \right), \quad \|z\|_2 \leq C\sqrt{d+t}.$$

On event of probability at least $1 - e^{-t}$, we can equivalently define $R_n(2\varepsilon_n)$ via

$$R_n(2\varepsilon_n) = \left\{ \frac{z}{\sqrt{n}} : \|z\|_2 \leq C\sqrt{d+t} \text{ and } W_n(z) \geq \sup_z W_n(z) - 2\sqrt{n}\varepsilon_n \right\}.$$

Moreover, the depth process $W_n(z)$ converges weakly to

$$W(z) := W_G(z) = \inf_{\|v\|_2=1} \left\{ G(v, 0) - \frac{1}{\sqrt{2\pi}} \langle z, v \rangle \right\}, \quad \|z\| \leq C\sqrt{d+t},$$

where $G(v, y)$ is a Brownian bridge indexed by half-spaces – a centered Gaussian process with covariance function

$$\mathbb{E}G(u, z)G(v, y) = \text{Cov} (I\{\langle X - z, u \rangle \geq 0\} I\{\langle X - y, v \rangle \geq 0\}), \quad X \sim N(0, I_d).$$

This result is known and is due to Massé (2002). We will also need the following estimate on the rate of convergence of W_n to W .

Lemma 2.2. *For each $n \geq 1$, one can construct on the same probability space independent random vectors X_1, \dots, X_n with standard normal distribution and a Brownian bridge G such that*

$$\sup_{\|z\|_2 \leq C\sqrt{d+t}} |W_n(z) - W(z)| \leq K(C) \left(\tilde{\gamma}_n + \left(\frac{d+t}{n} \right)^{1/4} \sqrt{\log(n)} + \sqrt{\frac{d+t}{n}} \sqrt{\log(2\sqrt{n/d})} \right)$$

with probability at least $1 - 2/n$, where $\tilde{\gamma}_n = n^{-\frac{1}{2+5d}} (\log(n))^{\frac{4+5d}{4+10d}}$.

We include the proof of the lemma in the appendix. In what follows, we will set

$$\gamma_n := K(C) \left(\tilde{\gamma}_n + \left(\frac{d}{n} \right)^{1/4} \sqrt{\log(n)} + \sqrt{\frac{d+t}{n}} \sqrt{\log(2\sqrt{n/d})} \right).$$

Lemma 2.3. *Let $\hat{z} \in \arg \max_{z \in \mathbb{R}^d} W(z)$ and $R(\beta) = \{\|z\|_2 \in \mathbb{R}^d : W(z) \geq \sup_{z \in \mathbb{R}^d} W(z) - \beta\}$. Then there exists an event \mathcal{E} of probability at least $1 - e^{-t}$ such that on this event, $\|\hat{z}\|_2 \leq C\sqrt{d+t}$. Moreover, for any sequence $\beta_j = o(1)$ and any $\tau > 0$,*

$$\lim_{j \rightarrow \infty} \mathbb{P}(\{\text{diam}(R(\beta_j)) \geq \tau\} \cap \mathcal{E}) = 0.$$

Proof. Lemma 3.8 in (Massé, 2002) states that with probability 1, \hat{z} is unique. To establish the first claim, note that in view of Gaussian concentration inequality (Tsirelson et al., 1976),

$$\inf_{\|v\|_2=1} G(v, 0) \geq \mathbb{E} \inf_{\|v\|_2=1} G(v, 0) - \sqrt{2t} \sup_{\|v\|_2=1} \text{var}^{1/2}(G(v, 0))$$

with probability at least $1 - e^{-t}$. A standard argument³ based on Dudley's entropy integral bound (Dudley, 1967) and the fact that the Vapnik-Chervonenkis (VC) dimension of the class of half-spaces passing through the origin equals d implies that

$$\mathbb{E} \inf_{\|v\|_2=1} G(v, 0) \geq -C_1\sqrt{d}.$$

Since $\text{var}(G(v, 0)) = 1/4$ for all v , we deduce that

$$\inf_{\|v\|_2=1} G(v, 0) \geq -C_2\sqrt{d+t}$$

on event \mathcal{E} of probability at least $1 - e^{-t}$. We conclude that on \mathcal{E} ,

$$W(0) = \inf_{\|v\|_2=1} G(v, 0) \geq -C_2\sqrt{d+t}.$$

Moreover, for $\|z\|_2 > C\sqrt{d+t}$, we have

$$W(z) \leq -\frac{1}{\sqrt{2\pi}}\|z\|_2 + \sup_{\|v\|_2=1} G(v, 0) < C_2\sqrt{d+t},$$

whenever C is sufficiently large. Therefore, $\|\hat{z}\|_2 \leq C\sqrt{d+t}$ on \mathcal{E} . The proof of the second claim now proceeds as follows. Assume that for some increasing sequence $\{j_i\}_{i \geq 1}$,

$$\inf_{i \geq 1} \mathbb{P}(\{\text{diam}(R(\beta_{j_i})) \geq \tau\} \cap \mathcal{E}) \geq c > 0.$$

Since the sets $R(\beta_{j_i})$ are nested, it implies that

$$\mathbb{P}\left(\left\{\text{diam}\left(\bigcap_{i \geq 1} R(\beta_{j_i})\right) \geq \tau\right\} \cap \mathcal{E}\right) > 0.$$

But any element of the set $\bigcap_{i \geq 1} R(\beta_{j_i})$ must be the $\arg \max_z W(z)$, contradicting the a.s. uniqueness of the latter. The claim follows. \square

We will now deduce the statement of the theorem. It follows from the facts established above that there exists an event Θ of probability at least $1 - 2e^{-t} - \frac{2}{n}$ such that the claims of Lemmas 2.3 and 2.2 hold on Θ , and moreover the inequality $\text{diam}(R_n(2\varepsilon_n)) \leq C_1\sqrt{\frac{d+t}{n}}$ is satisfied.

³For example, it suffices to combine Theorem 2.37 with Theorems 4.47 or 4.52 in (Dudley, 2014).

Therefore, on Θ

$$\begin{aligned} R_n(\varepsilon_n) &= \left\{ \frac{z}{\sqrt{n}} : \|z\|_2 \leq C\sqrt{d+t} \text{ and } W_n(z) \geq \sup_z W_n(z) - 2\sqrt{n}\varepsilon_n \right\} \\ &\subseteq \left\{ \frac{z}{\sqrt{n}} : \|z\|_2 \leq C\sqrt{d+t} \text{ and } W(z) \geq \sup_z W(z) - 2 \sup_{\|z\| \leq C\sqrt{d+t}} |W_n(z) - W(z)| - 2\sqrt{n}\varepsilon_n \right\} \\ &\subseteq \left\{ \frac{z}{\sqrt{n}} : \|z\|_2 \leq C\sqrt{d+t} \text{ and } W(z) \geq \sup_z W(z) - 2\gamma_n - 2\sqrt{n}\varepsilon_n \right\} := \mathcal{A}_n. \end{aligned}$$

Since $\gamma_n + \sqrt{n}\varepsilon_n = o(1)$ as $n \rightarrow \infty$, we conclude that

$$\sqrt{n} \text{diam}(\mathcal{A}_n) \cdot I_\Theta \rightarrow 0 \text{ in probability as } n \rightarrow \infty,$$

yielding the conclusion. \square

3. DISCUSSION

We proved that Tukey's median, and more generally affine equivariant estimators of location, are sensitive to the geometry of the underlying distribution, in a sense that their performance, measured with respect to the Euclidean distance, is controlled by the effective rank of the "shape" matrix rather than the ambient dimension. In the adversarial contamination framework, our claim remains valid as long as the contamination proportion ε is either large (larger than $\sqrt{\frac{d}{n}}$) or small, namely of order $o(n^{-1/2})$. There is still a significant gap between the bounds for Tukey's median proved in this paper and best known guarantees for the alternative estimators, for example by Minasyan and Zhivotovskiy (2023). This fact motivates the following questions:

- (1) Is it possible to prove a version of Corollary 2.4 that allows the effective rank $r(\Sigma)$ and the ambient dimension d grow with the sample size n ?
- (2) What are the best guarantees for Tukey's median in the adversarial contamination framework with ε_n of order $\sqrt{\frac{r(\Sigma)}{n}}$?
- (3) More generally, does Tukey's median satisfy non-asymptotic, finite-sample guarantees of the form (3)?
- (4) Can the bound of Corollary 2.3 be sharpened? In other words, what is the typical diameter of Tukey's set when the data follow an isotropic normal distribution?

REFERENCES

- Aubrun, G., Jenkinson, J., and Szarek, S. J. (2024). Optimal constants in concentration inequalities on the sphere and in the Gauss space. *arXiv preprint arXiv:2406.13581*.
- Berthet, P. and Mason, D. M. (2006). Revisiting two strong approximation results of Dudley and Philipp. *High Dimensional Probability*, pages 155–172.
- Borell, C. (1975). The Brunn-Minkowski inequality in Gauss space. *Inventiones mathematicae*, 30(2):207–216.
- Brown, B. M. (1983). Statistical uses of the spatial median. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 45(1):25–30.
- Caplin, A. and Nalebuff, B. (1988). On 64%-majority rule. *Econometrica: Journal of the Econometric Society*, pages 787–814.
- Cerdeira, J. O. and Silva, P. C. (2021). A centrality notion for graphs based on Tukey depth. *Applied Mathematics and Computation*, 409:126409.
- Chen, M., Gao, C., and Ren, Z. (2018). Robust covariance and scatter matrix estimation under huber's contamination model. *The Annals of Statistics*, 46(5):1932–1960.

- Chen, Z. (1995). Robustness of the half-space median. *Journal of statistical planning and inference*, 46(2):175–181.
- Chernozhukov, V., Galichon, A., Hallin, M., and Henry, M. (2017). Monge-Kantorovich depth, quantiles, ranks and signs. *Annals of Statistics*, 45(1):223–256.
- Depersin, J. and Lecué, G. (2023). On the robustness to adversarial corruption and to heavy-tailed data of the Stahel–Donoho median of means. *Information and Inference: A Journal of the IMA*, 12(2):814–850.
- Diakonikolas, I. and Kane, D. M. (2023). *Algorithmic high-dimensional robust statistics*. Cambridge university press.
- Donoho, D. L. (1982). Breakdown properties of multivariate location estimators. Ph.D. qualifying paper, Harvard University. *Statistics Dept.*
- Donoho, D. L. and Gasko, M. (1992). Breakdown properties of location estimates based on halfspace depth and projected outlyingness. *The Annals of Statistics*, pages 1803–1827.
- Dudley, R. M. (1967). The sizes of compact subsets of hilbert space and continuity of Gaussian processes. *Journal of Functional Analysis*, 1(3):290–330.
- Dudley, R. M. (2014). *Uniform central limit theorems*, volume 142. Cambridge university press.
- Hallin, M. and Paindaveine, D. (2002). Optimal tests for multivariate location based on interdirections and pseudo-Mahalanobis ranks. *The Annals of Statistics*, 30(4):1103–1133.
- Hodges, J. L. (1955). A bivariate sign test. *The Annals of Mathematical Statistics*, 26(3):523–527.
- Hotelling, H. (1929). Stability in competition. *The economic journal*, 39(153):41–57.
- Huber, P. J. and Ronchetti, E. M. (2009). *Robust statistics; 2nd ed.* Wiley Series in Probability and Statistics. Wiley, Hoboken, NJ.
- Koltchinskii, V. I. (1997). M-estimation, convexity and quantiles. *The Annals of Statistics*, pages 435–477.
- Lévy, P. (1951). Problèmes concrets d’analyse fonctionnelle. *2nd ed. Gauthier-Villars, Paris.*
- Liu, R. Y., Parelius, J. M., and Singh, K. (1999). Multivariate analysis by data depth: descriptive statistics, graphics and inference. *The Annals of Statistics*, 27(3):783–858.
- Massé, J.-C. (2002). Asymptotics for the Tukey median. *Journal of Multivariate Analysis*, 81(2):286–300.
- Massé, J.-C. (2004). Asymptotics for the Tukey depth process, with an application to a multivariate trimmed mean. *Bernoulli*, 10(3):397–419.
- Minasyan, A. and Zhivotovskiy, N. (2023). Statistically optimal robust mean and covariance estimation for anisotropic Gaussians. *arXiv preprint arXiv:2301.09024*.
- Nagy, S. (2022). Halfspace depth: Theory and computation. *Habilitation Thesis*.
- Nagy, S., Schuett, C., and Werner, E. (2019). Data depth and floating body. *Statistics surveys*, 13.
- Nehring, K. and Puppe, C. (2023). Multi-dimensional social choice under frugal information: the Tukey median as Condorcet winner ex ante by. Technical report, KIT Working Paper Series in Economics.
- Pokorný, D., Laketa, P., and Nagy, S. (2024). Another look at halfspace depth: flag halfspaces with applications. *Journal of Nonparametric Statistics*, 36(1):165–181.
- Rényi, A. (1953). On the theory of order statistics. *Acta Math. Acad. Sci. Hung.* 4(2):48–89.
- Rousseeuw, P. J. and Ruts, I. (1999). The depth function of a population distribution. *Metrika*, 49(3):213–244.
- Serfling, R. (2002). Quantile functions for multivariate analysis: approaches and applications. *Statistica Neerlandica*, 56(2):214–232.
- Small, C. G. (1997). Multidimensional medians arising from geodesics on graphs. *The Annals of Statistics*, pages 478–494.

- Stahel, W. A. (1981). *Breakdown of covariance estimators*. Fachgruppe für Statistik, Eidgenössische Techn. Hochsch.
- Tsirelson, B., Ibragimov, I., and Sudakov, V. (1976). Norms of Gaussian sample functions. In *Proceedings of the Third Japan—USSR Symposium on Probability Theory*, pages 20–41. Springer.
- Tukey, J. W. (1975). Mathematics and the picturing of data. In *Proceedings of the International Congress of Mathematicians*, volume 2, pages 523–531.

APPENDIX A. PROOF OF THEOREM 2.1.

Define the random vector

$$W := \Sigma^{-1/2}(\tilde{\mu}_n(X_1, \dots, X_n) - \mu),$$

and observe that we need to estimate $\|\Sigma^{1/2}W\|_2$. Moreover, set $Z_j = \Sigma^{-1/2}(X_j - \mu)$, $j = 1, \dots, n$. Let us record the following identity:

$$\Sigma^{1/2}W = \|W\|_2 \Sigma^{1/2} \left[\frac{W}{\|W\|_2} \right],$$

where $U := \frac{W}{\|W\|_2}$ is defined as 0 if $W = 0$. We claim that conditionally on $W \neq 0$, U has uniform distribution over the sphere of radius 1. Indeed, for any orthogonal matrix Q , the vector QZ_1 has the same distribution as Z_1 in view of elliptical symmetry, which implies that $\tilde{\mu}_n(QZ_1, \dots, QZ_n)$ and $\tilde{\mu}_n(Z_1, \dots, Z_n)$ have the same distribution. But

$$\begin{aligned} \tilde{\mu}_n(QZ_1, \dots, QZ_n) &= Q\tilde{\mu}_n(Z_1, \dots, Z_n), \\ \tilde{\mu}_n(Z_1, \dots, Z_n) &= \Sigma^{-1/2}(\tilde{\mu}_n(X_1, \dots, X_n) - \mu) = W, \end{aligned}$$

due to affine equivariance, hence $W \stackrel{d}{=} QW$. It implies that the conditional distribution of $\frac{W}{\|W\|}$ given that $W \neq 0$ is indeed uniform.

The function $w \mapsto \|\Sigma^{1/2}w\|_2$ Lipschitz continuous with Lipschitz constant equal to $\|\Sigma\|^{1/2}$, therefore we can apply Lévy's lemma (Lévy, 1951), again conditionally on $W \neq 0$, to get that

$$\|\Sigma^{1/2}U\|_2 \leq \mathbb{E}\|\Sigma^{1/2}U\|_2 + \sqrt{\frac{2\|\Sigma\|^{1/2}t}{d}}$$

with probability at least $1 - e^{-t}$ (this version of the inequality with sharp constants is due to Aubrun et al. (2024)). The same result also holds unconditionally since $U = 0$ if $W = 0$. It remains to observe that

$$\mathbb{E}\|\Sigma^{1/2}U\|_2 \leq \mathbb{E}^{1/2}\|\Sigma^{1/2}U\|_2^2 = \sqrt{\frac{\text{tr}(\Sigma)}{d}},$$

hence we conclude by the union bound that

$$\|\Sigma^{1/2}W\|_2 = \|W\|_2 \|\Sigma^{1/2}U\|_2 \leq e(n, t, d) \left(\sqrt{\frac{\text{tr}(\Sigma)}{d}} + \sqrt{\frac{2\|\Sigma\|^{1/2}t}{d}} \right)$$

with probability at least $1 - p(t) - e^{-t}$.

Remark A.1. *Spherical symmetry plays a crucial role in the proof of the theorem. Due to the affine invariance of the depth function, it is not possible to capture the dependence of the error on the effective rank by simply comparing the empirical and the true depth. For example, let*

X_1, \dots, X_n be i.i.d. $N(0, \Sigma)$ random vectors where Σ is non-singular, and note that for any half-space $H(0, u)$, $P(X_1 \in H(0, u)) = \frac{1}{2}$. Therefore,

$$\sup_{\|u\|=1} \left| \frac{1}{n} \sum_{j=1}^n I\{X_j \in H(0, u)\} - \frac{1}{2} \right| = \sup_{\|u\|=1} \left| \frac{1}{n} \sum_{j=1}^n I\{\Sigma^{-1/2} X_j \in H(0, u)\} - \frac{1}{2} \right|,$$

hence the distribution of the supremum does not depend on Σ .

APPENDIX B. PROOF OF LEMMA 2.2.

The strong approximation result by Berthet and Mason (2006, Proposition 1) guarantees the existence of a sequence X_1, \dots, X_n and the Brownian bridge $G(v, z)$ defined on the same probability space such that

$$(9) \quad \sup_{z \in \mathbb{R}^d, \|v\|=1} \left| \sqrt{n}(P_n^X - P)H(z, v) - G(v, z) \right| \leq \delta_n$$

with probability at least $1 - 1/n$. Set

$$Z_n(z) = \inf_{\|v\|=1} \left(\sqrt{n}(P_n^X - P)I \left\{ \left\langle \cdot - \frac{z}{\sqrt{n}}, v \right\rangle \right\} \right)$$

for brevity. Fix z and consider two possibilities: (a) $W_n(z) - W(z) \geq 0$ and (b) $W_n(z) - W(z) < 0$. Observe that the infimum in the definitions of $W_n(z)$ and $W(z)$ is attained, almost surely, at some unit vectors $v_n(z)$ and $v(z)$. In case (a),

$$(10) \quad 0 \leq W_n(z) - W(z) \\ = \underbrace{\sqrt{n} \left(P_n^X H \left(\frac{z}{\sqrt{n}}, v_n(z) \right) - \frac{1}{2} \right) - \sqrt{n} \left(P_n^X H \left(\frac{z}{\sqrt{n}}, v(z) \right) - \frac{1}{2} \right)}_{\leq 0} \\ + \sqrt{n} \left(P_n^X H \left(\frac{z}{\sqrt{n}}, v(z) \right) - \frac{1}{2} \right) - \sqrt{n}(P_n^X - P)H \left(\frac{z}{\sqrt{n}}, v(z) \right) \\ + \sqrt{n}(P_n^X - P)H \left(\frac{z}{\sqrt{n}}, v(z) \right) - G \left(v(z), \frac{z}{\sqrt{n}} \right) + G \left(v(z), \frac{z}{\sqrt{n}} \right) - W(z).$$

Note that

$$\sup_{\|z\| \leq C\sqrt{d+t}} \left| \sqrt{n} \left(P_n^X H \left(\frac{z}{\sqrt{n}}, v(z) \right) - \frac{1}{2} \right) - \sqrt{n}(P_n^X - P)H \left(\frac{z}{\sqrt{n}}, v(z) \right) \right| \\ = \sup_{\|z\| \leq C\sqrt{d+t}} \left| PH \left(\frac{z}{\sqrt{n}}, v(z) \right) - \frac{1}{2} \right| \leq C_1 \frac{\sqrt{d+t}}{\sqrt{n}}$$

since P is the standard normal law. Moreover, inequality (9) implies that

$$\sup_{z \in \mathbb{R}^d} \left| \sqrt{n}(P_n^X - P)H \left(\frac{z}{\sqrt{n}}, v(z) \right) - G \left(v(z), \frac{z}{\sqrt{n}} \right) \right| \leq \delta_n$$

with probability at least $1 - 1/n$. Finally,

$$\left| G \left(v(z), \frac{z}{\sqrt{n}} \right) - W(z) \right| \leq \sup_{\|v\|=1} \left| G \left(v, \frac{z}{\sqrt{n}} \right) - G(v, 0) \right| + \frac{\|z\|}{\sqrt{2\pi n}}.$$

In view of the Gaussian concentration inequality (Tsirolson et al., 1976),

$$\begin{aligned} & \sup_{\|z\| \leq C\sqrt{d+t}, \|v\|=1} \left| G\left(v, \frac{z}{\sqrt{n}}\right) - G(v, 0) \right| \\ & \leq \mathbb{E} \sup_{\|z\| \leq C\sqrt{d+t}, \|v\|=1} \left| G\left(v, \frac{z}{\sqrt{n}}\right) - G(v, 0) \right| + C \left(\frac{d+t}{n}\right)^{1/4} \sqrt{s} \end{aligned}$$

with probability at least $1 - e^{-s}$. Dudley's entropy integral bound yields, via a standard argument (as explained in the proof of Lemma 2.3), that

$$\mathbb{E} \sup_{\|z\| \leq C\sqrt{d+t}, \|v\|=1} \left| G\left(v, \frac{z}{\sqrt{n}}\right) - G(v, 0) \right| \leq \sqrt{\frac{d+t}{n}} \sqrt{\log(2\sqrt{n/d})}.$$

In case (b), the reasoning is very similar. Collecting the estimates of each term in (10), we deduce the desired conclusion.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA
Email address: `minsker@usc.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA
Email address: `yinanshe@usc.edu`