Whittaker–Shintani functions for Fourier–Jacobi models on unitary groups

Paul Boisseau

November 1, 2024

Abstract

We state and prove a formula for the Whittaker–Shintani functions associated to Fourier– Jacobi models for *p*-adic unitary groups and general linear groups. These generalized spherical functions play a fundamental role in the proof of the Gan–Gross–Prasad conjecture for Fourier– Jacobi models. As an application we prove the unramified Ichino–Ikeda conjecture.

Contents

1	Introduction	1
2	Whittaker–Shintani functions	6
3	Multiplicity one for Whittaker–Shintani functions	11
4	Integral expression and analytic continuation	20
5	Formula for the normalized Whittaker–Shintani function	22
6	Unfolding of tempered periods	31
7	Unramified L-function for Fourier–Jacobi models	35

1. Introduction

1.0.1. Let V be a non degenerate n-dimensional skew-Hermitian space relative to a quadratic extension of number fields k'/k, and let $W \subset V$ be a non degenerate subspace of dimension m such that the corank n - m =: 2r is even. Set $G := U(V) \times U(W)$. Assume that the orthogonal complement W^{\perp} is split, which means that it contains an isotropic subspace X of dimension r. Gan Gross and Prasad have introduced in [GGP12] a subgroup $H \leq G$ and an automorphic representation $\overline{\nu}_{\mu,\psi} = \otimes \overline{\nu}_{\mu,\psi,v}$ of H built from the Weil representation of U(W). It depends on μ a character of the idele class group $(k')^{\times} \setminus \mathbb{A}_{k'}^{\times}$ lifting the quadratic character $\eta_{k'/k}$ attached to k'/k

by global class field theory, and on ψ a non trivial character of $k \setminus \mathbb{A}_k$. If $\sigma = \otimes \sigma_v$ is a cuspidal automorphic representation of $G(\mathbb{A}_k)$, one can define a global Fourier–Jacobi period

$$\mathcal{P}_{H}(\varphi,\phi) := \int_{H(k) \setminus H(\mathbb{A}_{k})} \varphi(h) \theta(h,\phi) dh$$

for $\varphi \in \sigma$, $\phi \in \overline{\nu}_{\mu,\psi}$, where θ is a theta series. Assume that $\operatorname{Hom}_{H(k_v)}(\sigma_v \otimes \overline{\nu}_{\mu,\psi,v}, \mathbb{C}) \neq \{0\}$ for any place v of k. The Gan–Gross–Prasad conjecture for Fourier–Jacobi models then states that \mathcal{P}_H is non-zero if and only if the central value of the complete Rankin–Selberg *L*-function $L(\frac{1}{2}, \sigma \otimes \overline{\mu})$ is non-zero. Progress has been obtained by Xue in [Xue14] and [Xue16] in the corank 0 case under local assumptions on π . With Lu and Xue, we proved the conjecture for any even corank and for every cuspidal representation of G whose base change is generic in [BLX24].

By uniqueness of local Fourier–Jacobi functionals, \mathcal{P}_H decomposes as an Eulerian product of linear forms $\mathcal{L}_{H_v} \in \operatorname{Hom}_{H_v}(\sigma_v \otimes \overline{\nu}_{\mu,\psi,v}, \mathbb{C})$. In [BPC23], Beuzart-Plessis and Chaudouard used an unfolding argument to reduce the conjecture in the Bessel case (i.e. when n-m is odd) to the corank 1 situation. Moreover, if σ is assumed to be everywhere tempered, integrating matrix coefficients yields a H_v -invariant sesquilinear form \mathcal{P}_{H_v} on $\sigma_v \otimes \overline{\nu}_{\mu,\psi,v}$. In this setting, Xue formulated in [Xue16] a variation of the Ichino–Ikeda conjecture [II10] relating $|\mathcal{P}_H|^2$ to $\prod_v \mathcal{P}_{H_v}$. For both problems, it is necessary to first check that the corresponding local statement holds in the unramified setting in order to regularize the Eulerian product. The goal of this article is to understand the unramified behaviour of the local Fourier–Jacobi functionals \mathcal{L}_{H_v} . This includes two cases: if the place v is inert, then $U(V)_v$ and $U(W)_v$ are quasi-split unitary groups, if it is split they are isomorphic to $GL_n(k_v)$ and $GL_m(k_v)$.

1.0.2. We now describe our main results. We work in the local unramified situation and we drop the place v from the notation, so that E/F is an unramified quadratic extension of p-adic fields, or $E = F \times F$. Denote by q_F and q_E the cardinals of the respective residual fields (if $E = F \times F$, take $q_E = q_F$). Assume that 2 does not divide q_F , that V and W are split in the sense of §2.1.2 and that μ and ψ are unramified. We identify any algebraic group over F with its group of Fpoints. Let (T, B^+) be a Borel pair in G such that HB^+ is open in G and T contains $N_G(H) \cap B^+$. Write $T = T_V \times T_W$. Let $K = K_V \times K_W$ be a hyperspecial open-compact subgroup of G in good position relatively to T. Let χ and η be unramified characters of T_V and T_W . Assume that σ is the unramified irreducible subquotient of the parabolic induction $I_{B^+}^G(\chi \boxtimes \eta)$. Let $\varphi^\circ \in \sigma^K$ be a non-zero K-spherical vector. The representation $\overline{\nu}_{\mu,\psi}$ of H extends to a representation of a group K_J in J, and $\overline{\nu}_{\mu,\psi}^{K_J}$ has dimension 1. Let $\phi^\circ \in \overline{\nu}_{\mu,\psi}^{K_J}$ be non-zero. Consider

$$\mathcal{W}_{\chi,\overline{\mu}\eta}(g) := \mathcal{L}_H(\sigma(g)\varphi^\circ \otimes \phi^\circ), \quad g \in \mathcal{U}(V).$$
(1.0.2.1)

This $\mathcal{W}_{\chi,\overline{\mu}\eta}$ belongs to the space $\mathrm{WS}_{\chi,\overline{\mu}\eta}$ of *Whittaker–Shintani functions* associated to $(\chi,\overline{\mu}\eta)$. It is roughly defined as follows. In §2.3 we introduce a space WS of functions $\mathcal{W} \in C^{\infty}(\mathrm{U}(V))$ which satisfy some bi-invariance properties. It is a module on the Hecke algebra $\mathcal{H}(\mathrm{U}(V), K_V) \otimes \mathcal{H}(J, K_J)$ by left and right convolutions. $\mathrm{WS}_{\chi,\overline{\mu}\eta}$ is then the eigenspace of functions that transform by the Satake character of $\sigma \otimes \overline{\nu}_{\mu,\psi}$. Our first result is multiplicity one for such functions. It is proved in §3.3.

Theorem 1.0.2.1. The space $WS_{\chi,\overline{\mu}\eta}$ has dimension one.

The space WS admits another more natural description. Let $\nu_{\mu,\psi}$ be the complex conjugate of $\overline{\nu}_{\mu,\psi}$ and consider $(\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ the space of spherical vectors in the smooth induction. This is a $\mathcal{H}(G, K)$ -module. There exists a surjective map of \mathbb{C} -algebras $\mathcal{H}(U(V), K_V) \otimes \mathcal{H}(J, K_J) \to \mathcal{H}(G, K)$ and we have an isomorphism of $\mathcal{H}(U(V), K_V) \otimes \mathcal{H}(J, K_J)$ -modules

WS
$$\simeq \left(\operatorname{Ind}_{H}^{G} \nu_{\mu,\psi} \right)^{K}$$
. (1.0.2.2)

Note that a modification is needed if r = 0 (see §3.4). Let $\Lambda \subset T$ be the group of cocharacters of T, and let Λ^- be the subset of negative cocharacters relatively to B^+ . By duality, Theorem 1.0.2.1 is a consequence of the following result which is of independent interest. It is a combination of Proposition 3.2.2.2 and Proposition 3.2.2.3.

Proposition 1.0.2.2. The spherical space in the compact induction $(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ decomposes as

$$\left(\mathrm{ind}_{H}^{G}\nu_{\mu,\psi}\right)^{K} = \bigoplus_{\lambda \in \Lambda^{-}} \mathbb{C} \Phi_{\lambda},$$

where Φ_{λ} is the unique vector in $(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ such that $\operatorname{supp}(\Phi_{\lambda}) \subset H\lambda K$ and $\Phi_{\lambda}(\lambda) = \phi^{\circ}$. Moreover, $(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ is a free $\mathcal{H}(G,K)$ -module of rank 1.

The image in $(\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ of the function $\mathcal{W}_{\chi,\overline{\mu}\eta}$ defined in (1.0.2.1) is easy to describe. By choosing an invariant inner product $(\cdot, \cdot)_{\nu}$, we have an identification $\overline{\nu}_{\mu,\psi}^{\vee} \simeq \nu_{\mu,\psi}$. By duality, the operator \mathcal{L}_{H} corresponds to $\mathcal{L}_{H}^{\vee} \in \operatorname{Hom}_{H}(\sigma, \nu_{\mu,\psi})$. Consider the map

$$\Phi_{\chi,\overline{\mu}\eta}(g) = \mathcal{L}_{H}^{\vee}(\sigma(g)\varphi^{\circ}) \in \nu_{\mu,\psi}, \quad g \in G.$$
(1.0.2.3)

Then $\Phi_{\chi,\overline{\mu}\eta} \in (\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ is the image of $\mathcal{W}_{\chi,\overline{\mu}\eta}$ by (1.0.2.2).

Our second main result is a formula for $\Phi_{\chi,\overline{\mu}\eta}$. Set $n_- = \lfloor n/2 \rfloor$, $m_- = \lfloor m/2 \rfloor$ and r' = m + r. Let δ_{B^+} be the the modular character of B^+ . Denote by Σ_{nd}^+ the set of non-divisible positive roots of T in B^+ . Then T is identified in a natural way with the canonical maximal torus T_G of G which is $(\operatorname{Res}_{E/F}\mathbb{G}_m)^{n_-+m_-}$ in the inert even case, $(\operatorname{Res}_{E/F}\mathbb{G}_m)^{n_-+m_-} \times U(1)^2$ in the inert odd case, and \mathbb{G}_m^{n+m} in the split case. We identify the character $\chi \boxtimes \eta$ with a couple $(\chi, \eta) \in$ $(\mathbb{C}^{\times})^{n_-+m_-}$ or $(\mathbb{C}^{\times})^{n+m}$. Let W_G be the Weyl group of G and let w_0 be its longest element. Define $\zeta_F(s) = (1 - q_F^{-s})^{-1}$ and $\zeta_E(s) = (1 - q_E^{-s})^{-1}$ the local zeta functions, and for $x, s \in \mathbb{C}$ set $L_F(s,x) = (1 - xq_F^{-s})^{-1}$ and $L_E(s,x) = (1 - xq_E^{-s})^{-1}$. The RHS of Theorem 1.0.2.3 is well-defined for generic (χ, η) and in fact extends to a regular function. The exact constants involved in the formula depend on whether E/F is inert or split, and n is even or odd. The theorem is proved in two steps in §5.5 and §7.2.

Theorem 1.0.2.3. We have $(\Phi_{\chi,\overline{\mu}\eta}(1),\phi^{\circ})_{\nu} = 0$ if and only if $\Phi_{\chi,\overline{\mu}\eta} = 0$. Moreover if $\Phi_{\chi,\overline{\mu}\eta} \neq 0$ then for all $\lambda \in \Lambda^-$ we have

$$\frac{(\Phi_{\chi,\overline{\mu}\eta}(\lambda),\phi^{\circ})_{\nu}}{(\Phi_{\chi,\overline{\mu}\eta}(1),\phi^{\circ})_{\nu}} = \frac{\Delta_{\mathrm{U}(W)}}{\Delta_{T_{W}}} \sum_{w \in W_{G}} \mathbf{b}(w_{V}\chi,w_{W}\overline{\mu}\eta)\mathbf{d}(w(\chi \boxtimes \eta)) \left(w(\chi \boxtimes \eta)\delta_{B^{+}}^{-\frac{1}{2}}\right)(\lambda), \quad (1.0.2.4)$$

where $w = (w_V, w_W)$,

$$\Delta_{T_W} = \begin{cases} \zeta_E(1)^{m_-} & inert \ even \ case, \\ \zeta_E(1)^{m_-} L_F(1, \eta_{E/F}) & inert \ odd \ case, \\ \zeta_F(1)^m & split \ case, \end{cases} \quad \Delta_{\mathcal{U}(W)} = \begin{cases} \prod_{i=1}^m L_F(i, \eta_{E/F}^i) & inert \ case, \\ \prod_{i=1}^m \zeta_F(i) & split \ case, \end{cases}$$

$$\mathbf{d}(\chi \boxtimes \eta) = \prod_{\alpha \in \Sigma_{\mathrm{nd}}^+} \frac{1}{1 - \langle \chi \boxtimes \eta, \alpha^{\vee} \rangle},$$

and $\mathbf{b}(\chi,\eta)$ is

$$\begin{cases} \prod_{\substack{1 \le i \le n-\\ 1 \le j \le m-\\ 1 \le j \le m-}} L_E^{-1}(\frac{1}{2}, \chi_i \eta_j) \prod_{i < r+j} L_E^{-1}(\frac{1}{2}, \chi_i \eta_j^{-1}) \prod_{i > r+j} L_E^{-1}(\frac{1}{2}, \chi_i^{-1} \eta_j) & \text{inert } n \text{ even} \\ \prod_{\substack{1 \le i \le n-\\ 1 \le j \le m-\\ 1 \le j \le m-}} L_E^{-1}(\frac{1}{2}, \chi_i \eta_j) L_E^{-1}(\frac{1}{2}, -\chi_i) L_E^{-1}(\frac{1}{2}, \eta_j) \prod_{i < r+j} L_E^{-1}(\frac{1}{2}, \chi_i \eta_j^{-1}) \prod_{i > r+j} L_E^{-1}(\frac{1}{2}, \chi_i^{-1} \eta_j) & \text{inert } n \text{ odd,} \\ \prod_{i+j < r'+1} L_F^{-1}(\frac{1}{2}, \chi_i \eta_j) \prod_{i+j > r'+1} L_F^{-1}(\frac{1}{2}, \chi_i^{-1} \eta_j^{-1}) & \text{split.} \end{cases}$$

(1.0.2.5)

Although Theorem 1.0.2.3 is the appropriate result for applications, the factors $\mathbf{b}(\chi,\eta)$ lack a satisfying interpretation. In the split case, we propose a more robust version. Let \hat{G} be the Langlands dual group of G with Borel pair (\hat{T}, \hat{B}^+) . We have the *L*-groups LG and LT . Let $S \in {}^LT$ be a representative of the Satake parameter of σ . We define in §7.1.2 a representation $\mathcal{R}_{\overline{\mu}} : {}^LG \to \mathrm{GL}(\mathbb{C}^n \otimes \mathbb{C}^m \oplus \mathbb{C}^n \otimes \mathbb{C}^m)$. It satisfies

$$L(s, \sigma \otimes \overline{\mu}) = \det(1 - q_F^{-s} \mathcal{R}_{\overline{\mu}}(S)).$$
(1.0.2.6)

The representation $\mathcal{R}_{\overline{\mu}}$ is symplectic. In §7.1.3 we choose a Lagrangian subspace \mathcal{Y} which is stable by the restriction of $\mathcal{R}_{\overline{\mu}}$ to ${}^{L}T$. Let $\mathcal{Y}_{\overline{\mu}}$ be this representation of ${}^{L}T$. Denote by $D_{\widehat{G}/\widehat{B}^{+}}(S)$ the determinant of $1 - \operatorname{Ad}(S)$ on $\operatorname{Lie}(\widehat{G})/\operatorname{Lie}(\widehat{B}^{+})$. We introduce in §3.1.3 an element $\phi^{\times} \in \overline{\nu}_{\mu,\psi}$. It is not spherical but is fixed by $I_{W} \cap w_{0}I_{W}w_{0}^{-1}$, where I_{W} is a Iwahori subgroup of GL_{m} . The vector ϕ^{\times} depends on the choice of a Schrödinger model of $\overline{\nu}_{\mu,\psi}$ which is equivalent to the choice of a Lagrangian subspace Y of the symplectic space $\operatorname{Res}_{E/F}W$. The Lagrangian subspaces \mathcal{Y} and Y are chosen to be compatible. The following proposition is proved in §7.2.5.

Proposition 1.0.2.4. Assume that we are in the split case and that $\Phi_{\chi,\overline{\mu}\eta} \neq 0$. For $\lambda \in \Lambda^-$ we have

$$\frac{(\Phi_{\chi,\overline{\mu}\eta}(\lambda),\phi^{\times})_{\nu}}{(\Phi_{\chi,\overline{\mu}\eta}(1),\phi^{\circ})_{\nu}} = \Delta_{\mathrm{U}(W)} \sum_{w \in W_{G}} \frac{\det(1-q_{F}^{-\overline{2}}\mathcal{Y}_{\overline{\mu}}(wS))}{D_{\widehat{G}/\widehat{B}^{+}}(wS)} \left(w_{0}w(\chi \boxtimes \eta)\delta_{B^{+}}^{-\frac{1}{2}}\right)(\lambda).$$

In the inert case, there is no counterpart to ϕ^{\times} as any $I_W \cap w_0 I_W w_0^{-1}$ -fixed vector is K_W -fixed (see Lemma 3.1.3.1), and moreover there is no Lagrangian subspace of $\mathcal{R}_{\overline{\mu}}$ stable by ${}^L T$. However, the representation $\mathcal{R}_{\overline{\mu}}$ makes sense and (1.0.2.6) is satisfied. The factor $\mathbf{b}(\chi,\eta)$ of Theorem 1.0.2.3 can then be interpreted as $\det(1-q_F^{-\frac{1}{2}}(\mathcal{R}_{\overline{\mu}})_{|\mathcal{V}_-}(S))$, where \mathcal{V}_- is an isotropic subspace of $\mathcal{R}_{\overline{\mu}}$ which is not Lagrangian but which is stable by ${}^L T$.

Our proof of Theorem 1.0.2.3 and Proposition 1.0.2.4 follows the strategy initiated in [Cas80] and [CS80] for Whittaker functionals, which has been used to establish formulas for Whittaker–Shintani functions for GL_n in [KMS], orthogonal groups in [KMS03], Bessel models for unitary groups in [Kho08] and GL_n in [Zha18], and for symplectic groups in [She13a]. This last case is the closest in spirit to ours. In our setting, the main difficulties come from dealing with the representation $\overline{\nu}_{\mu,\psi}$: this is done in §2.2 by interpreting $\sigma \otimes \overline{\nu}_{\mu,\psi}$ as a principal series on the bigger

Jacobi group $U(V) \times J$. This does not fit into the usual theory of induction on reductive groups and complicates the computations.

Note that our formulae lie outside of the abstract result established in [Sak13] for functions on spherical varieties. Ben-Zvi, Sakellaridis and Venkatesh have recently proposed in [BZSV24] a new framework that could be used to interpret them.

1.0.3. We give an application of Theorem 1.0.2.3 to the Ichino–Ikeda conjecture for Fourier–Jacobi periods formulated in [Xue16]. Assume that σ is tempered. We equip it with an invariant inner product (\cdot, \cdot) . We take a model over \mathcal{O}_F of H (see §2.1.8) and give H the left-invariant measure dh normalized so that $H(\mathcal{O}_F)$ has volume 1. Define the local Ichino–Ikeda period for $\varphi \in \sigma$, $\phi \in \overline{\nu}_{\mu,\psi}$

$$\mathcal{P}_{H}(\varphi,\phi) := \int_{H} (\sigma(h)\varphi,\varphi) (\overline{\nu}_{\mu,\psi}(h)\phi,\phi)_{\nu} dh.$$
(1.0.3.1)

This integral is only convergent if r = 0 and needs to be interpreted as the unique continuous extension of a linear form on $C_c^{\infty}(G)$ to the space of tempered functions $C^w(G)$ otherwise (see §6.2 for a precise statement which is not specific to the unramified setting and will be given for any local field of characteristic zero). The following theorem is the unramified Ichino–Ikeda conjecture. It had been proved in [Har12] for Bessel models on unitary groups. For Fourier–Jacobi models it was only known in corank zero by [Xue16]. It is proved in §7.3.

Theorem 1.0.3.1. Let $\varphi^{\circ} \in \sigma$ be a non-zero spherical vector. Then

$$\frac{\mathcal{P}_H(\varphi^{\circ},\phi^{\circ})}{(\varphi^{\circ},\varphi^{\circ})(\phi^{\circ},\phi^{\circ})_{\nu}} = \Delta_{\mathrm{U}(V)} \frac{L(\frac{1}{2},\sigma\otimes\overline{\mu})}{L(1,\sigma,\mathrm{Ad})}$$

1.0.4. The paper is organized as follows. In §2 we give a precise definition of Whittaker–Shintani functions for Fourier-Jacobi models using the Jacobi group J. In §3, we relate these functions to the Heisenberg-Weil representation $\overline{\nu}_{\mu,\psi}$ and prove the multiplicity one Theorem 1.0.2.1. In the process, we show Proposition 1.0.2.2. We then proceed in §4 to produce an element $\mathcal{W}_{\chi,\eta}^{I} \in WS_{\chi,\eta}$ defined by an integral expression for (χ,η) in some open subset, which we then extend to generic (χ,η) by meromorphic continuation. In §5, we compute an explicit formula for $\mathcal{W}_{\chi,\eta}^{I}$: this is done by calculating the γ factors caused by the uniqueness of Fourier–Jacobi models. To prove Theorem 1.0.2.3, it then remains to compute the normalization value $\mathcal{W}_{\chi,\eta}^{I}(1)$. In §6 we temporarily leave the unramified setting to discuss convergence and state an unfolding identity for tempered Ichino–Ikeda periods. In §7, we reformulate our results in terms of Satake parameters and representations of the Langlands dual group ${}^{L}G$. We then use the Weyl character formula to determine $\mathcal{W}_{\chi,\eta}^{I}(1)$ (thus completing the proof of Theorem 1.0.2.3), and finally prove Theorem 1.0.3.1 by reduction to the known corank zero result using §6.

1.0.5. Acknowledgement The author thanks Raphaël Beuzart-Plessis for helpful discussions and comments on many parts of this article. He is also grateful to Hang Xue for explanations on his results.

This work was partly funded by the European Union ERC Consolidator Grant, RELANTRA, project number 101044930. Views and opinions expressed are however those of the author only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

2. Whittaker–Shintani functions

2.1. General notations

2.1.1. Unless specified otherwise, F will be a non-Archimedean local field of characteristic zero, with ring of integers \mathcal{O}_F and maximal ideal \mathfrak{p}_F . We denote by q_F the cardinality of the residual field $\mathcal{O}_F/\mathfrak{p}_F$ and assume that it is odd. We fix an uniformizer ϖ and normalize the valuation v_F and the absolute value $|\cdot|_F$ on F such that $v_F(\varpi) = 1$ and $|x|_F = q_F^{-v_F(x)}$ for $x \in F$.

Let E be a quadratic unramified extension of F (referred to as the *inert* case), or $E := F \times F$ (the *split* case). In the inert case, write \mathcal{O}_E and q_E as before, and normalize v_E and $|\cdot|_E$ such that $v_E(\varpi) = 1$ and $|x|_E = q_E^{-v_E(x)}$. Let $c \in \operatorname{Gal}(E/F)$ be the non-trivial element, and set $E^- := \{x \in E \mid c(x) = -x\}$. Fix $\tau \in \mathcal{O}_E^{\times} \cap E^-$ (which exists as q_F is odd). In the split case, write $\mathcal{O}_E := \mathcal{O}_F \times \mathcal{O}_F$. We have $\operatorname{Gal}(E/F) = \{1, c\}$ where c switches the coordinates of elements in $F \times F$. Let $\operatorname{Tr}_{E/F}$ and $N_{E/F}$ be the trace and norm maps of E/F which make sense in both settings. Let $|\cdot|$ be $|\cdot|_E$ in the inert case and $|\cdot|_F$ in the split case, v be v_E in the inert case and v_F in the split case, q be q_E in the inert case and q_F in the split case.

2.1.2. Let $(V, \langle ., . \rangle_V)$ be a *n*-dimensional non-degenerate skew-Hermitian space over E/F and $W \subset V$ be a non-degenerate subspace of dimension m such that n = m + 2r with $r \geq 0$. In the inert case, set $n_- := \lfloor \frac{n}{2} \rfloor$ and $m_- := \lfloor \frac{m}{2} \rfloor$. If n is even (referred to as the *even* case), define $n_+ = n_-$ and $m_+ = m_-$, while if n is odd (the *odd* case), set $n_+ = n_- + 1$ and $m_+ = m_- + 1$. In the split case, set $n_- = n_+ = n$ and $m_- = m_+ = m$. Set r' = m + r.

In the inert case, we assume that $V = W \oplus W^{\perp}$ is split. This means that there exists an *E*-basis $\{v_i, v_i^* \mid 1 \le i \le r\}$ of W^{\perp} and an *E*-basis $\{w_i, w_i^* \mid 1 \le i \le m_+\}$ of *W* (with the convention $w_{m_+} = w_{m_+}^*$ in the odd case) such that for all $1 \le i, j \le r$, $1 \le i' \le m_+$ and $1 \le j' \le m_-$ we have

$$\begin{cases} \langle v_i, v_j \rangle_V = \langle v_i^*, v_j^* \rangle_V = \langle w_{i'}, w_{j'} \rangle_V = \langle w_{i'}^*, w_{j'}^* \rangle_V = 0, \\ \langle v_i, v_j^* \rangle_V = \delta_{i,j}, \\ \langle w_{i'}, w_{j'}^* \rangle_V = \delta_{i',j'}, \\ \langle w_{m_+}, w_{m_+} \rangle_V = \tau \text{ in the odd case.} \end{cases}$$

$$(2.1.2.1)$$

We obtain an *E*-basis $\mathfrak{B} = (v_1, \ldots, v_r, w_1, \ldots, w_{m_+}, w_{m_-}^*, \ldots, w_1^*, v_r^*, \ldots, v_1^*)$ of *V*.

In the split case, the above construction can be made explicit and it will be convenient to choose a particular \mathfrak{B} . We have $V = F^n \times F^n$ and we can and will assume that $\langle ., . \rangle_V$ is given by

$$\langle (x,y), (x',y') \rangle_V = ({}^t x y', -{}^t y x'),$$

where we identify F^n with column vectors. Let $\{\mathbf{e}_i\}$ and $\{\mathbf{e}_i^*\}$ be the canonical basis of $F^n \times \{0\}$ and $\{0\} \times F^n$ respectively. Define for $1 \le i \le n_+$ the elements $v_i = \mathbf{e}_i + \mathbf{e}_{n-i+1}^*$ and $v_i^* = -\mathbf{e}_{n-i+1} + \mathbf{e}_i^*$. The basis $\mathfrak{B} = (v_1, \ldots, v_{n_+}, v_{n_-}^*, \ldots, v_1^*)$ is an *E*-basis of *V* satisfying (2.1.2.1) with $\tau = (1, -1)$ (up to taking $w_i = v_{r+i}$ and $w_i^* = v_{r+i}^*$). Set

$$W = \operatorname{span}_E(v_i, v_i^* \mid r+1 \le i \le n_+).$$

For $1 \leq j \leq m$, set $w_j = \mathbf{e}_{j+r}$ and $w_j^* = \mathbf{e}_{j+r}^* \in W$. Note that $(w_1, \ldots, w_m, w_1^*, \ldots, w_m^*)$ is a *F*-basis of *W*.

In both cases, set

$$X = \operatorname{span}_E(v_i \mid 1 \le i \le r), \text{ and } X^* = \operatorname{span}_E(v_i^* \mid 1 \le i \le r),$$

so that $V = X \oplus W \oplus X^*$.

2.1.3. Denote by U(V) and U(W) the subgroups of GL(V) and GL(W) consisting of the *E*-linear unitary transformations of *V* and *W* respectively. Note that n_- (resp. m_-) is always the split *F*rank of U(V) (resp. U(W)). We will identify U(W) as the subgroup of U(V) of $g \in U(V)$ which act by the identity on W^{\perp} . The basis \mathfrak{B} chosen in §2.1.2 yields an isomorphism $GL(V) \cong GL_n(E)$ and two embeddings $U(W), U(V) \hookrightarrow GL_n(E)$. Denote by (T_V, B_V) and (T_W, B_W) the inverse image in U(V) and U(W) of the torus of diagonal matrices and the Borel subgroup of upper triangular matrices by this morphism. In the split case, we will furthermore identify U(V) with $GL_n(F)$ and U(W) with $GL_m(F)$ by projecting on the first coordinate. Note that through this isomorphism B_V and B_W are identified with the Borel subgroups of upper triangular matrices in $GL_n(F)$ and $GL_m(F)$ respectively.

Denote by N_V and N_W the unipotent radicals of B_V and B_W respectively. The opposite Borel subgroups will be denoted by $B_V^- = T_V N_V^-$ and $B_W^- = T_W N_W^-$. Set $T = T_V \times T_W$, $B = B_V \times B_W$ and $B^+ = B_V^- \times B_W$.

Any $t_V \in T_V$ is of the form $t_V(\mathbf{t})$ where

$$t_{V}(\mathbf{t}) = \begin{cases} \operatorname{diag}(t_{1}, \dots, t_{n_{-}}, c(t_{n_{-}})^{-1}, \dots, c(t_{1})^{-1}), \ \mathbf{t} \in (E^{\times})^{n_{-}} & \text{inert even case,} \\ \operatorname{diag}(t_{1}, \dots, t_{n_{-}}, t_{n_{+}}, c(t_{n_{-}})^{-1}, \dots, c(t_{1})^{-1}) \ \mathbf{t} \in (E^{\times})^{n_{-}} \times \mathrm{U}(1), & \text{inert odd case,} \\ \operatorname{diag}(t_{1}, \dots, t_{n}) \ \mathbf{t} \in (F^{\times})^{n}, & \text{split case.} \end{cases}$$

(2.1.3.1)

If $t \in T_V$ we will write t_i for the coordinates of **t** in (2.1.3.1). Set $|t| = \prod_{i=1}^{n_-} |t_i|$. We let Λ_V be the group of cocharacters of T_V , and Λ_V^+ be its cone of positive cocharacters with respect to B_V , i.e.

$$\Lambda_V^+ := \begin{cases} \{(a_1, \dots, a_{n_-}) \in \mathbb{Z}^{n_-} \mid a_1 \ge \dots \ge a_{n_-} \ge 0\} & \text{inert case;} \\ \{(a_1, \dots, a_n) \in \mathbb{Z}^n \mid a_1 \ge \dots \ge a_n\} & \text{split case.} \end{cases}$$

If $\lambda_V \in \Lambda_V^+$, we will simply write λ_V for $\lambda_V(\varpi)$.

The subgroup of U(V) of automorphisms preserving X and X^* and acting by identity on W is isomorphic to $\operatorname{GL}_r(E)$ and is denoted by G_r . Set $T_r = G_r \cap T_V$ and $B_r = G_r \cap B_V$. Let Λ_r be its group of cocharacters and Λ_r^+ be its the subcone of positive cocharacters with respect to B_r .

With the same definitions, we also have T_W , $t_W(\mathbf{t})$, Λ_W , Λ_W^+ , Λ_W^- and $|\cdot|$. Let $\Lambda = \Lambda_V \times \Lambda_W$ be the group of cocharacters of T, and let $\Lambda^- = \Lambda_V^+ \times \Lambda_W^-$ be the cone of negative cocharacters relatively to B^+ .

2.1.4. Let A_V (resp. A_W) be the split component of T_V (resp. T_W). Let $\langle ., . \rangle$ be the canonical pairing between the groups of characters $X^*(A_V)$ and cocharacters $X_*(A_V)$ of A_V . Denote by Σ_V the set of roots of A_V in U(V), $\Sigma_{V,nd}$ the subset of non-divisible roots, $\Sigma_{V,nd}^+$ the set of non-divisible positive roots with respect to B_V . Let Δ_V be the simple roots in $\Sigma_{V,nd}^+$. If $\alpha \in \Sigma_V$, write α^{\vee} for the associated coroot. Then we write $\alpha > 0$ if for all $\beta \in \Delta_V$ we have $\langle \alpha, \beta^{\vee} \rangle > 0$. Let W_V be the Weyl group of U(V). It has an action on Σ_V which we will write $w.\alpha$. The longest element in W_V will be denoted by w_0

For each $1 \leq i \leq n$, let e_i be the character of A_V such that $e_i(t) = t_i$ with $t \in A_V$. If $\alpha \in \Delta_V$, we will write N_α for the corresponding root subgroup. Denote by P_α the parabolic subgroup corresponding to α , with Levi component G_α whose derived group is $D(G_\alpha)$. Denote by w_α the simple reflection corresponding to α . The N_α 's are isomorphic to product of copies of F or Ethrough maps n_α . More precisely, denote by $\{E_{i,j}\}$ the canonical basis of $M_n(E)$ in the inert case, and $M_n(F)$ in the split case, where $M_n(E)$ (resp. $M_n(F)$) is the vector space of $n \times n$ -matrices over E (resp. over F). Then we can write each $\alpha \in \Delta_V$ as $\alpha = e_i - e_{i+1}$ and we have the description

	Case	n_{lpha}	$D(G_{\alpha})$
$1 \le i < n$	inert	$n_{\alpha}(x) = I_n + x E_{i,i+1} - c(x) E_{n-i,n-i+1}, \ x \in E$	$\operatorname{SL}_2(E)$
$1 \le i < n$	split	$n_{\alpha}(x) = I_n + x E_{i,i+1}, \ x \in F$	$\operatorname{SL}_2(F)$
$i = n_{-}$	inert even	$n_{\alpha}(x) = I_n + xE_{n-,n-+1}, \ x \in F$	$\operatorname{SL}_2(F)$
$i = n_{-}$	inert odd	$n_{\alpha}(x_1, x_2) = I_n - \tau c(x_1) E_{n, n_+} + x_1 E_{n_+, n_+ + 1} + x_2 E_{n, n_+ + 1}$ $x_1, x_2 \in E, \ \text{Tr}_{E/F}(\frac{x_2}{\tau}) = -N_{E/F}(x_1)$	$SU_3(F)$

Similar notations hold for U(W). To avoid any confusion we will write n'_{β} if $\beta \in \Sigma_W$.

2.1.5. Consider the *F*-vector spaces $\mathbb{V} := \operatorname{Res}_{E/F} V$ and $\mathbb{W} := \operatorname{Res}_{E/F} W$ equipped with the symplectic form $\langle ., . \rangle_{\mathbb{V}} := \operatorname{Tr}_{E/F} \circ \langle ., . \rangle_{V}$ and let $\mathbb{H}(\mathbb{W}) = \mathbb{W} \times F$ be the associated Heisenberg group, with group law $(w_1, z_1).(w_2, z_2) = (w_1 + w_2, z_1 + z_2 + \frac{1}{2} \langle w_1, w_2 \rangle_{\mathbb{V}})$. If $r \geq 1$, we have an injective morphism $h : \mathbb{H}(\mathbb{W}) \to N_V$ characterized by

$$\begin{cases}
h(w, z)(v_i) = v_i & 1 \le i \le r, \\
h(w, z)(w') = w' - \langle w', w \rangle_V v_r & w' \in W, \\
h(w, z)(v_r^*) = (-\frac{1}{2} \langle w, w \rangle_V + z) v_r + w + v_r^*, \\
h(w, z)(v_i^*) = v_i^* & 1 \le i \le r - 1.
\end{cases}$$
(2.1.5.1)

If $g_1, g_2 \in U(V)$, we will write $g_1 g_2 := g_1 g_2 g_1^{-1}$. The subgroup $\mathbb{H}(\mathbb{W})$ of U(V) is stable by conjugation by U(W) and satisfies

$${}^{g_W}h(w,z) = h(g_W w, z), \ g_W \in \mathcal{U}(W), \ w \in W, \ z \in F.$$
 (2.1.5.2)

We define

$$J := \mathcal{U}(W) \ltimes \mathbb{H}(\mathbb{W}) \tag{2.1.5.3}$$

the Jacobi group of W. If $r \ge 1$ it is a subgroup of U(V) by (2.1.5.1). Its center h(0, F) will be simply denoted by Z.

Let U be the unipotent radical of the parabolic subgroup of U(V) stabilizing the flag $Ev_1 \subset Ev_1 \oplus Ev_2 \subset \ldots \subset Ev_1 \oplus \ldots Ev_{r-1}$. Define

$$G = U(V) \times U(W), \quad H := J \ltimes U \text{ if } r \ge 1, \quad H = U(W) \text{ if } r = 0.$$
 (2.1.5.4)

Then H is a subgroup of G by the natural inclusion in the first component and the projection on U(W) in the second (this is the diagonal embedding if r = 0). Set $W_G := W_V \times W_W$.

2.1.6. For $1 \le k \le r$, set $G_k := \operatorname{GL}_k(E)$ seen as the subgroup of $\operatorname{U}(V)$ stabilizing $\operatorname{Span}_E(v_i \mid 1 \le i \le k)$ and $\operatorname{Span}_E(v_i^* \mid 1 \le i \le k)$ and acting trivially on their orthogonal complement. Set $B_k := G_k \cap B_V = T_k N_k$. Let P(X) be the parabolic subgroup of $\operatorname{U}(V)$ stabilizing X. Its Levi subgroup stabilizing X* is isomorphic to $G_r \times \operatorname{U}(W)$. Let N(X) be its unipotent radical. Set $L := G \times G_r$. It is a Levi subgroup of the parabolic $\operatorname{U}(V) \times P(X)$ of $\operatorname{U}(V) \times \operatorname{U}(V)$. If $r \ge 1$ define

$$H^L = H \times N_r \subset L.$$

Let P be the parabolic subgroup of U(V) stabilizing the full flag $0 \subset Ev_1 \subset ... \subset X$ of X, and N be its unipotent radical. Then we have the alternative description $H = U(W) \ltimes N$. Note that HB^+ is open in G.

2.1.7. In the inert case, if R is a subgroup of E set $W(R) := \operatorname{span}_R(w_i, w_i^* \mid 1 \le i \le m_+)$, $Y_-(R) := \operatorname{span}_R(w_i \mid 1 \le i \le m_-), Y_-^*(R) := \operatorname{span}_R(w_i^* \mid 1 \le i \le m_-)$ and $Y_{m_+} = \{0\}$ in the even case, $Y_{m_+}(R) = Y_{m_+}^*(R) = Rw_{m_+}$ in the odd case. Let $Y_+ := Y_- \oplus Y_{m_+}$ and $Y_+^* := Y_{m_+}^* \oplus Y_-^*$. Set

$$Y := Y_{-}(E) \oplus Y_{m_{+}}(\mathcal{O}_{E}).$$

In the split case, we use the same notations with $Y_{-}(R) = \operatorname{span}_{R}(w_{i} \mid 1 \leq i \leq m)$, $Y_{-}^{*}(R) = \operatorname{span}_{R}(w_{i}^{*} \mid 1 \leq i \leq m)$ and $Y_{m_{+}} = \{0\}$. Set $Y := Y_{-}(E)$. All these groups embed into J by h and we identify them with their image. Consider

$$N_J := N_W \ltimes YZ, \text{ and } B_J := T_W \ltimes N_J. \tag{2.1.7.1}$$

Note that we can identify $Y_{-}(E)$ and $Y_{-}^{*}(E)$ with $E^{m_{-}}$ (inert case) or F^{m} (split case) with our choice of basis. For $y \in Y_{-}(E)$ or $y^{*} \in Y_{-}^{*}(E)$, we will use (y_{i}) and (y_{i}^{*}) to write the coordinates and write $t_{W}(y)$ or $t_{W}(y^{*})$ as in (2.1.3.1) if they are all non-zero. Let $1_{Y_{+}}^{*}$ be the element of $Y_{+}^{*}(E)$ whose coordinates are all 1's. Define $\mu_{+}^{*} := h(1_{Y_{+}^{*}}, 0)$.

The subgroup of U(W) stabilizing Y_{-} and Y_{-}^{*} and acting by identity on $Y_{m_{+}}$ can be identified with $\operatorname{GL}_{m_{-}}(E)$ in the inert case and $\operatorname{GL}_{m}(F)$ in the split case by restricting to Y_{-} . Then $g \in$ $\operatorname{GL}_{m_{-}}(E)$ (resp. $\operatorname{GL}_{m}(F)$) acts by g on $Y_{-}(E)$ and $(g^{*})^{-1}$ on $Y_{-}^{*}(E)$, where $g^{*} = {}^{t}c(g)$ (resp. $g^{*} = {}^{t}g$). Note that this property holds in the split case because of our particular choice of F-basis $\{w_{j}, w_{j}^{*}\}$ of W (see §2.1.2). Combined with (2.1.5.2), we get for $w = y + y_{m_{+}} + y^{*} \in$ $Y_{-}(E) \oplus Y_{m_{+}}(E) \oplus Y_{-}^{*}(E) = W$ and $g \in \operatorname{GL}_{m_{-}}(E)$ (resp. $\operatorname{GL}_{m}(F)$ in the split case)

$${}^{g}h(w,0) = h(gy + y_{m_{+}} + (g^{*})^{-1}y^{*}, 0).$$
 (2.1.7.2)

In the odd inert case, there is also a copy of U(1) contained in T_W that acts on $Y_{m_+}(E)$.

2.1.8. Let $K_V \leq U(V)$ be the stabilizer of the lattice $\bigoplus_{v \in \mathfrak{B}} \mathcal{O}_E v$ and set $K_W = K_V \cap U(W)$. These are hyperspecial maximal open compact subgroups of U(V) and U(W) respectively. Recall that $U(V) \subset \operatorname{GL}_n(E)$ and set for $i \geq 0$

$$\mathbb{G}^{i} = \mathbb{G} \cap K_{V} \cap (I_{n} + M_{n \times n}(\varpi^{i}\mathcal{O}_{E})),$$

where $\mathbb G$ is any of the groups we have considered so far.

Set $K_J := K_W \ltimes W^0 Z^0$, an open-compact subgroup of J. Set $I_V = N_V^{-,1} T_V^0 N_V^0$ and $I_W = N_W^{-,1} T_W^0 N_W^0$ which are Iwahori subgroups of U(V) and U(W). Define $I_J := I_W \ltimes W^0 Z^0$. Set $K = K_V \times K_W \subset G$.

2.1.9. We equip equip all our groups \mathbb{G} with the left invariant Haar-measure giving \mathbb{G}^0 volume 1. In particular, $db_J := db_W dy dz$ will be our measure on $B_J = B_W \ltimes YZ$. The modular character is

$$\delta_{B_J}(t_W n_J) = \delta_{B_W}(t_W) |t_W| \ t_W \in T_W, \ n_J \in N_J.$$
(2.1.9.1)

where we recall that $|t_W| = \prod_{j=1}^{m_-} |t_{W,j}|$.

2.1.10. Denote by ζ_F and ζ_E the zeta functions associated to F and E, i.e. $\zeta_F(s) = (1 - q_F^{-s})^{-1}$, $\zeta_E(s) = (1 - q_E^{-s})^{-1}$ in the inert case and $\zeta_E(s) = (1 - q_F^{-s})^{-2}$ in the split case. For $x \in \mathbb{C}$, define

$$L_F(s,x) = (1 - xq_F^{-s})^{-1}$$
 and $L_E(s,x) = (1 - xq_E^{-s})^{-1}$.

Set $\Delta_{U(V)} = L_{U(V)}(0)$, $\Delta_{G_k} = L_{G_k}(0)$ and $\Delta_{T_W} = L_{T_W}(0)$, where $L_{\mathbb{G}}$ is the Artin–Tate *L*-function associated to \mathbb{G} defined in [Gro97]. In our case,

$$\Delta_{\mathrm{U}(V)} = \prod_{i=1}^{n} L_{F}(i, \eta_{E/F}^{i}), \ \Delta_{G_{k}} = \prod_{i=1}^{k} \zeta_{E}(i), \ \Delta_{T_{W}} = \begin{cases} \zeta_{F}(1)^{m_{-}} L(1, \eta_{E/F})^{m_{+}} & \text{inert case,} \\ \zeta_{F}(1)^{m} & \text{split case,} \end{cases}$$
(2.1.10.1)

where $\eta_{E/F}$ is the quadratic character associated to E/F. We will also use Δ'_{T_W} which is Δ_{T_W} in the inert even case and the split case, and $\Delta'_{T_W} = \zeta_F(1)^{m_-} L(1, \eta_{E/F})^{m_-}$ in the inert odd case.

2.2. Representations

Let G_2 be a locally compact totally disconnected group, let $G_1 \leq G_2$ be a closed subgroup. If π is a smooth representation of G_1 , let $\operatorname{Ind}_{G_1}^{G_2}\pi$ be the smooth induction, and $\operatorname{ind}_{G_1}^{G_2}\pi$ be the compact smooth induction (see [Ren10, Section III.2.2]). We reserve the notation I for parabolic induction.

We fix once and for all ψ an unramified character of F of conductor \mathcal{O}_F which will appear in the definition of Whittaker–Shintani functions. It is a consequence of Theorem 1.0.2.3 that the latter are independent of ψ , so that we will often drop it from the notations.

Let χ and η be unramified characters of T_V and T_W extended to B_V and B_W , that we identify with elements of $(\mathbb{C}^{\times})^{n_-}$ and $(\mathbb{C}^{\times})^{m_-}$ via the rules $\chi(t_V(\mathbf{t})) = \prod_{i=1}^{n_-} \chi_i^{v(t_i)}$ and $\eta(t_W(\mathbf{t})) = \prod_{j=1}^{m_-} \eta_j^{v(t_j)}$. We also extend $|\cdot|$ to B_V and B_W . Note that the map

$$(\eta\overline{\psi})(b_Wh(y,z)) := \eta(b_W)\overline{\psi}(z), \ b_W \in B_W, \ y \in Y, \ z \in F,$$

defines a character of B_J . We introduce the parabolic induced representations

$$I_{B_{V}}^{U(V)}(\chi) = \operatorname{Ind}_{B_{V}}^{U(V)}(\chi \delta_{B_{V}}^{\frac{1}{2}}), \quad I_{B_{W}}^{U(W)}(\eta) = \operatorname{Ind}_{B_{W}}^{U(W)}(\eta \delta_{B_{W}}^{\frac{1}{2}}), \quad I_{B_{J}}^{J}(\eta \overline{\psi}) = \operatorname{Ind}_{B_{J}}^{J}(\eta \overline{\psi} \delta_{B_{J}}^{\frac{1}{2}})$$

It is well-known that the parabolic induction $I_{B_V}^{U(V)}(\chi)$ and $I_{B_W}^{U(W)}(\eta)$ are compact inductions.

Lemma 2.2.0.1. We have $I_{B_J}^J(\eta\overline{\psi}) = \operatorname{ind}_{B_J}^J(\eta\overline{\psi}\delta_{B_J}^{\frac{1}{2}})$. Moreover, the spherical subspace $(I_{B_J}^J\eta\overline{\psi})^{K_J}$ is of dimension 1.

Proof. Let $f \in I_{B_J}^J(\eta \overline{\psi})$. By the Iwasawa decomposition we have $J = B_J Y_+^*(E) K_J$. Let $y^* \in Y_+^*(E)$. By smoothness, there exists an open neighbourhood \mathcal{U} of 0 in Y independent of y^* such that for all $k \in K_J$ and $y \in \mathcal{U}$ we have $f(h(y^*, 0)h(y, 0)k) = f(h(y^*, 0)k)$. But $f(h(y^*, 0)h(y, 0)k) = \overline{\psi}(\langle y^*, y \rangle_{\mathbb{W}})f(h(y^*, 0)k)$. If $f(h(y^*, 0)k) \neq 0$, this implies that y^* lives in a compact of $Y_+^*(E)$.

If f is K_J invariant, we can take $\mathcal{U} = Y_+(\mathcal{O}_E)$ so that $f(h(y^*, 0)) \neq 0$ implies $y^* \in Y_+^*(\mathcal{O}_E)$ as ψ is unramified.

For $\mathbb{G} \in {U(V), U(W), J}$, let $C_c^{\infty}(\mathbb{G})$ be the space of smooth compactly supported functions on \mathbb{G} . Set

$$\begin{split} F_{\chi}(f_{V})(g_{V}) &= \int_{B_{V}} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}})(b_{V}) f_{V}(b_{V} g_{V}) db_{V}, \quad f_{V} \in C_{c}^{\infty}(\mathrm{U}(V)), \ g_{V} \in \mathrm{U}(V), \\ F_{\eta}(f_{W})(g_{W}) &= \int_{B_{W}} (\eta^{-1} \delta_{B_{W}}^{\frac{1}{2}})(b_{W}) f_{W}(b_{W} g_{W}) db_{W}, \quad f_{W} \in C_{c}^{\infty}(\mathrm{U}(W)), \ g_{W} \in \mathrm{U}(W), \\ F_{\eta}^{J}(f_{J})(g_{J}) &= \int_{B_{J}} (\eta^{-1} \delta_{B_{J}}^{\frac{1}{2}} \psi)(b_{J}) f_{J}(b_{J} g_{J}) db_{J}, \quad f_{J} \in C_{c}^{\infty}(J), \ g_{J} \in J. \end{split}$$

This yields surjective intertwining maps $F_{\chi}: C_c^{\infty}(\mathrm{U}(V)) \to I_{B_V}^{\mathrm{U}(V)}(\chi), F_{\eta}: C_c^{\infty}(\mathrm{U}(W)) \to I_{B_W}^{\mathrm{U}(W)}(\eta)$ and $F_{\eta}^J: C_c^{\infty}(J) \to I_{B_J}^J(\eta\overline{\psi})$. In particular, we introduce the spherical vectors

$$\Phi_{\chi}^{\circ} := F_{\chi}(1_{K_V}), \ \Phi_{\eta}^{\circ} := F_{\eta}(1_{K_W}) \text{ and } \Phi_{\eta}^{\circ,J} := F_{\eta}^J(1_{K_J}),$$
(2.2.0.1)

and for $w_V \in W_V$ and $w_W \in W_W$ the Iwahori-fixed vectors

$$\Psi_{w_V,\chi} := F_{\chi}(1_{I_V w_V I_V}), \ \Psi_{w_W,\eta} := F_{\eta}(1_{I_W w_W I_W}) \text{ and } \Psi^J_{w_W,\eta} := F^J_{\eta}(1_{I_J w_W I_J}).$$
(2.2.0.2)

2.3. Whittaker–Shintani functions

Let $\mathcal{H}(\mathcal{U}(V), K_V)$, $\mathcal{H}(\mathcal{U}(W), K_W)$ and $\mathcal{H}(J, K_J)$ be the spherical Hecke algebras of $\mathcal{U}(V)$, $\mathcal{U}(W)$ and J. By Lemma 2.2.0.1, they act by characters ξ_{χ} , ξ_{η} and $\xi_{\eta\overline{\psi}}^{J}$ on $(I_{B_V}^{\mathcal{U}(V)}\chi)^{K_V}$, $(I_{B_W}^{\mathcal{U}(W)}\eta)^{K_W}$ and $(I_{B_J}^J\eta\overline{\psi})^{K_J}$ respectively. For any function f on $\mathcal{U}(V)$, $\mathcal{U}(W)$ or J, we have the regular actions $L(g)f = f(g^{-1})$ and R(g)f = f(.g). Set $\psi_E := \psi \circ \frac{1}{2} \operatorname{Tr}_{E/F}$ and consider the generic character of U

$$\psi_U(u) := \psi_E\left(\sum_{i=1}^{r-1} \langle u(v_{i+1}), v_i^* \rangle_V\right), \ u \in U.$$

We define the space WS of Whittaker–Shintani functions to be the \mathbb{C} -vector space of functions $\mathcal{W} \in C^{\infty}(\mathcal{U}(V))$ if $r \geq 1$, or $\mathcal{W} \in C^{\infty}(J)$ if r = 0, such that for $z \in Z$, $u \in U$, $k_J \in K_J$, $k_V \in K_V$ and $g \in \mathcal{U}(V)$ if $r \geq 1$, $g \in J$ if r = 0, we have

$$\mathcal{W}(zuk_Jgk_V) = \psi(z)\psi_U(u)W(g).$$

It has a structure of $\mathcal{H}(\mathrm{U}(V), K_V) \otimes \mathcal{H}(J, K_J)$ -module by the $R \otimes L$ action. We denote by $\mathrm{WS}_{\chi,\eta}$ the $\xi_{\chi} \otimes \xi_{\eta\overline{\psi}}^J$ -isotypic component of WS, that is the space of $\mathcal{W}_{\chi,\eta} \in \mathrm{WS}$ such that for all $f_V \in \mathcal{H}(\mathrm{U}(V), K_V)$ and $f_J \in \mathcal{H}(J, K_J)$ we have

$$R(f_V)L(f_J)\mathcal{W}_{\chi,\eta} = \xi_{\chi}(f_V)\xi_{\eta\overline{\psi}}^J(f_J)\mathcal{W}_{\chi,\eta}.$$

3. Multiplicity one for Whittaker–Shintani functions

In this section we show that for every unramified characters χ and η of T_V and T_W respectively, the space WS_{χ,η} has dimension at most one. In §4 and §5 we will build a non zero element $W^{\circ}_{\chi,\eta}$, so that it has dimension exactly one. This proves the multiplicity one result Theorem 1.0.2.1.

3.1. Heisenberg–Weil representation of the Jacobi group

By the *p*-adic Stone–von Neumann Theorem [MVW87, Théorème I.1] there exists a unique smooth irreducible representation ρ_{ψ} of $\mathbb{H}(\mathbb{W})$ with central character ψ . Consider a S^1 metaplectic cover $Mp(\mathbb{W})$ of the symplectic group $Sp(\mathbb{W})$, which is unique up to isomorphism. There is a natural map $U(W) \to Sp(\mathbb{W})$. The data of ψ determines a smooth representation ω_{ψ} of $Mp(\mathbb{W})$ ([MVW87, Section 2.II]), realised on ρ_{ψ} . Let μ be a character of E^{\times} lifting $\eta_{E/F}$. Then the data of (μ, ψ) determines a splitting of $Mp(\mathbb{W})$ over U(W) which yields the Weil representation $\omega_{\mu,\psi}$ of U(W). We will write $\overline{\omega}_{\mu,\psi} = \omega_{\overline{\mu},\overline{\psi}}$ and $\overline{\rho}_{\psi} = \rho_{\overline{\psi}}$. By definition, for any $g \in U(W)$ the automorphism $\omega_{\mu,\psi}(g)$ is an intertwining operator between

By definition, for any $g \in U(W)$ the automorphism $\omega_{\mu,\psi}(g)$ is an intertwining operator between ρ_{ψ} and ρ_{ψ}^{g} and moreover the action of J by conjugation on ψ_{U} is trivial, so that we can define the representation $\nu_{\mu,\psi}$ of $J \ltimes U$ by the rule

$$\nu_{\mu,\psi}(g_W h u) = \psi_U(u)\omega_{\mu,\psi}(g_W)\rho_{\psi}(h), \ u \in U, \ g_W \in \mathrm{U}(W), \ h \in \mathbb{H}(\mathbb{W}).$$

Let $\overline{\nu}_{\mu,\psi}$ be its complex conjugation which is defined by the same formula with $\overline{\psi}_U$, $\overline{\omega}_{\mu,\psi}$ and $\overline{\rho}_{\psi}$.

3.1.1. Mixed Schrödinger-lattice models. In the split case or the inert even case, we have a polarization $\mathbb{W} = Y_{-}(E) \oplus Y_{-}^{*}(E)$, so that $\nu_{\mu,\psi}$ is realized on $C_{c}^{\infty}(Y_{-}^{*}(E)) = C_{c}^{\infty}(Y_{+}^{*}(E))$ by the Schrödinger model ([MVW87, Section II.6]). Otherwise, set

$$C_{c}^{\infty}(Y_{m_{+}}(E),\psi) := \{ f \in C_{c}^{\infty}(Y_{m_{+}}(E)) \mid f(y+y') = \psi(\frac{1}{2}\langle y, y' \rangle_{\mathbb{V}}) f(y), \ y \in Y_{m_{+}}(E), \ y' \in Y_{m_{+}}(\mathcal{O}_{E}) \}.$$
(3.1.1)

We may realize $\nu_{\mu,\psi}$ on $C_c^{\infty}(Y_-^*(E)) \otimes C_c^{\infty}(Y_{m_+}(E),\psi) \subset C_c^{\infty}(Y_+^*(E))$ by mixing a Schrödinger and a lattice model ([MVW87, Sections II.7 and II.8]). Then $\nu_{\mu,\psi}$ is unitary for the inner product

$$(\phi, \phi')_{\nu} = \int_{Y_{+}^{*}(E)} \phi(y^{*}) \overline{\phi'(y^{*})} dy^{*}, \quad \phi, \phi' \in \nu_{\mu,\psi}.$$
 (3.1.1.2)

This identifies $\nu_{\mu,\psi}^{\vee}$ with $\overline{\nu}_{\mu,\psi}$. There is a unique $\phi^{\circ} \in \nu_{\mu,\psi}^{K_J}$ with $(\phi^{\circ},\phi^{\circ})_{\nu} = 1$, namely $\phi^{\circ} := 1_{Y_+^*(\mathcal{O}_E)}$.

Let P(Y) be the parabolic subgroup of U(W) stabilizing the flag $\{0\} \subset Y_- \subset Y_+ \subset W$. Let M(Y) be its Levi subgroup stabilizing Y_{m_+} and Y_-^* . Then M(Y) is isomorphic to $GL(Y_-)$ in the inert even case and the split case, and $GL(Y_-) \times U(1)$ in the inert odd case, where $GL(Y_-) = GL_{m_-}(E)$ in the inert case and $GL(Y_-) = GL_m(F)$ in the split case. In the mixed Schrödingerlattice model, the representation $\nu_{\mu,\psi}$ admits an explicit description as a $P(Y) \ltimes \mathbb{H}(\mathbb{W})$ module ([GI16, Section 7.4]). It implies in particular that

$$(\nu_{\mu,\psi})_{|B_W \ltimes \mathbb{H}(\mathbb{W})} \simeq \operatorname{Ind}_{B_J}^{B_W \ltimes \mathbb{H}(\mathbb{W})} |\cdot|^{\frac{1}{2}} \mu \psi \simeq \operatorname{ind}_{B_J}^{B_W \ltimes \mathbb{H}(\mathbb{W})} |\cdot|^{\frac{1}{2}} \mu \psi, \qquad (3.1.1.3)$$

where we write μ for $\mu \circ \det$. Moreover, for $\phi \in \nu_{\mu,\psi}$, $a \in \operatorname{GL}(Y_-)$, $y \in Y_-(E)$, $y^* \in Y_-^*(E)$ and $y_0^* \in Y_+^*(E)$ we have

$$\nu_{\mu,\psi}(a)\phi(y_0^*) = \mu(\det a) |\det a|^{\frac{1}{2}} \phi(a^* y_0^*), \qquad (3.1.1.4)$$

$$\nu_{\mu,\psi}(h(y+y^*,0))\phi(y_0^*) = \psi\left(\langle y_0^*, y \rangle_{\mathbb{W}} + \frac{1}{2}\langle y^*, y \rangle_{\mathbb{W}}\right)\phi(y^*+y_0^*).$$
(3.1.1.5)

and in the inert odd case, for $g \in U(1)$, $y^* \in Y^*_{-}(E)$ and $y^*_{+} \in Ew_{m_{+}}$

$$\nu_{\mu,\psi}(g)\phi(y^* + y^*_+) = \phi(y^* + g^{-1}y^*_+).$$
(3.1.1.6)

3.1.2. Lattice models. There exists another model of $\nu_{\mu,\psi}$ called the "lattice" model (see [GKT23, Section 1.2.3]). Set $\mathbb{H}(\mathbb{W}^0) = \mathbb{H}(W^0, F)$. Then $\nu_{\mu,\psi}$ can be realized on the induction $\mathrm{Ind}_{\mathbb{H}(\mathbb{W}^0)}^{\mathbb{H}(\mathbb{W})}\psi$. In this model, $\mathbb{H}(\mathbb{W})$ acts by right translations and the subgroup K_W acts by left translation : for $\phi \in \mathrm{ind}_{\mathbb{H}(\mathbb{W}^0)}^{\mathbb{H}(\mathbb{W})}\psi$, $k_W \in K_W$, $w \in W$ and $z \in Z$ we have

$$\nu_{\mu,\psi}(k_W)\phi(h(w,z)) = \phi(h(k_W^{-1}w,z)), \qquad (3.1.2.1)$$

which by (2.1.5.2) amounts to

$$(\nu_{\mu,\psi})_{|K_W \ltimes H(W)} \simeq \operatorname{Ind}_{K_W \ltimes \mathbb{H}(\mathbb{W}^0)}^{K_W \ltimes \mathbb{H}(\mathbb{W})} \psi \simeq \operatorname{ind}_{K_W \ltimes \mathbb{H}(\mathbb{W}^0)}^{K_W \ltimes \mathbb{H}(\mathbb{W})} \psi$$
(3.1.2.2)

The space $(\nu_{\mu,\psi})^{K_J}$ has dimension one and is spanned by $\phi_{\text{latt}}^\circ : h(w,z) \mapsto \psi(z) \mathbb{1}_{W(\mathcal{O}_E)}(w).$

There is an unitary isomorphism $F_{W^0,Y}$ between the two models described in [GKT23, Proposition 1.23]. For $\phi \in \operatorname{ind}_{\mathbb{H}(\mathbb{W}^0)}^{\mathbb{H}(\mathbb{W})} \psi$ and $y^* \in Y^*_+(E)$ we have

$$F_{W^0,Y}(\phi)(y_+^*) = \int_{Y_-(E)} \phi(h(y,0)h(y_+^*,0))dy, \qquad (3.1.2.3)$$

and it sends $\phi_{\text{latt}}^{\circ}$ to $\phi^{\circ} = 1_{Y_{+}^{*}(\mathcal{O}_{E})}$.

3.1.3. Spherical vectors. Set $\phi^1 = 1_{Y_-^*(\varpi \mathcal{O}_E) \otimes Y_{m_+}(\mathcal{O}_E)}$ and $\phi^{\times} = \Delta'_{T_W} 1_{Y_-^*(\mathcal{O}_E^{\times}) \otimes Y_{m_+}(\mathcal{O}_E)}$ in the Schrödinger-lattice model of $\nu_{\mu,\psi}$. By (3.1.1.4) and (3.1.1.5) we have

$$\phi^{\times} = q^{m_{-}} \nu_{\mu,\psi}(1_{T_{W}^{0}}) \nu_{\mu,\psi}(\mu_{+}^{*}) \phi^{1}, \qquad (3.1.3.1)$$

where we recall that $\operatorname{vol}(T_W^0) = 1$. Note that in the split case $\Delta_{T_W} = \Delta'_{T_W}$ and $m = m_{-}$.

Lemma 3.1.3.1. In the inert case, $\nu_{\mu,\psi}^{I_W \cap {}^{w_0}I_W} = \mathbb{C}\phi^{\circ}$. In the split case, $\phi^{\times} \in \nu_{\mu,\psi}^{I_W \cap {}^{w_0}I_W}$.

Proof. We first treat the inert case with U(W) = U(1). Let $\phi \in \nu_{\mu,\psi}^{U(1)}$ which we identify with a subspace of $C_c^{\infty}(E,\psi)$ by (3.1.1.1). Let $y = u\varpi^d \in E$ with $u \in \mathcal{O}_E^{\times}$ and d < 0. Let $s_2 \in \mathcal{O}_F^{\times}$. By Hensel's Lemma and because q_F is odd, there exists $s_1 \in \mathcal{O}_F$ such that $s := \frac{\varpi^d}{(\varpi^d + s_1 + \tau s_2)} \in U(1)$. By (3.1.2.1) we have

$$\phi(y) = \nu_{\mu,\psi}(s)\phi(y) = \psi(N_{E/F}(u)\tau^2 s_2 \varpi^d)\phi(y).$$

As s_2 is arbitrary this implies that $\phi(y) = 0$ and therefore that $\phi \in \mathbb{C} \phi^{\circ}$.

To conclude the proof in the inert case, by (3.1.1.6) it remains to show that $\nu_{\mu,\psi}^{I_W \cap w_0 I_W} = \mathbb{C}\phi^\circ$ in the even case. For $c \in \operatorname{Hom}(Y^*_-, Y_-)$, let $c^* \in \operatorname{Hom}(Y^*_-, Y_-)$ be such that $\langle cy, y' \rangle_W = \langle y, c^*y' \rangle_W$ for all $y, y' \in Y^*_-$. Set

$$\operatorname{Herm}(Y_{-}^{*}, Y_{-}) = \{ c \in \operatorname{Hom}(Y_{-}^{*}, Y_{-}) \mid c^{*} = -c \}.$$

By [GI16, Section 7.1], there is an embedding $\operatorname{Herm}(Y_{-}^{*}, Y_{-}) \subset P(Y)$. By our choice of basis, $\operatorname{Herm}(Y_{-}^{*}, Y_{-})$ has an integral structure over \mathcal{O}_{F} and $\varpi\operatorname{Herm}(Y_{-}^{*}, Y_{-})(\mathcal{O}_{F}) \subset I_{W} \cap {}^{w_{0}}I_{W}$. By [GI16, Section 7.4], the action of $\operatorname{Herm}(Y_{-}^{*}, Y_{-})$ in the Schrödinger model is given by

$$\nu_{\mu,\psi}(c)\phi(y) = \psi(\langle cy, y \rangle_W)\phi(y), \quad c \in \operatorname{Herm}(Y_-^*, Y_-), \quad y \in Y_-^*.$$

Therefore, any $\phi \in \nu_{\mu,\psi}^{w_0 I_W}$ is in fact fixed by $\operatorname{Herm}(Y_-^*, Y_-)(\mathcal{O}_F)$ and thus supported in $Y_-^*(\mathcal{O}_E)$. By [GI16, Section 7.4], w_0 acts by the Fourier transform $\hat{\cdot}$ on $\nu_{\mu,\psi}$. Thus, $\nu_{\mu,\psi}^{I_W \cap w_0 I_W}$ is contained in the space of functions ϕ such that ϕ and $\hat{\phi}$ are supported in $Y_-^*(\mathcal{O}_E)$. They are multiple of ϕ° .

In the split case, note that $\phi^1 = \mu(\varpi)q_F^{m/2}\nu_{\mu,\psi}(t_W(\varpi,\ldots,\varpi)^{-1})\phi^\circ$, so that $\phi^1 \in \nu_{\mu,\psi}^{K_W}$. Moreover, for any $g \in I_W \cap {}^{w_0}I_W$ we have ${}^g\mu_+^* \in T_W^0\mu_+^*T_W^0$. The result follows from (3.1.3.1).

3.2. Spherical vectors of $\operatorname{ind}_{H}^{G} \nu_{\mu,\psi}$

In this subsection, we always assume that $r \ge 1$. We show that the space of K-spherical vectors of the compact induction $\operatorname{ind}_{H}^{G}\nu_{\mu,\psi}$ is a free $\mathcal{H}(G,K)$ -module of rank 1. We explain in §3.4 how to adapt this result to the r = 0 case.

3.2.1. We first record some results on the homogeneous space $X = H \setminus G$. Set $\Lambda_X = \Lambda_r + \Lambda_W^+$ and $\Lambda_X^- = \Lambda_r^+ + \Lambda_W^+$, which we identify with subsets of $\Lambda_V \subset G$. By the Iwasawa decomposition on U(V) and the Cartan decomposition on U(W), we see that Λ_X is a complete set of representatives of $H \setminus G/K$. Recall that $\Lambda = \Lambda_V \times \Lambda_W$ and $\Lambda^- = \Lambda_V^+ \times \Lambda_W^-$. There is a projection $\lambda = (\lambda_V, \lambda_W) \in \Lambda^- \mapsto \lambda_X := \lambda_V - \lambda_W \in \Lambda_X^-$. Note that $\lambda_X = \lambda_X'$ is equivalent to $H \lambda K = H \lambda' K$. For $\lambda_V, \lambda_V' \in \Lambda_V$, we write $\lambda_V \leq_V \lambda_V'$ if $\lambda_V' - \lambda_V = \sum_{\alpha \in \Delta_V} c_\alpha \alpha^{\vee}$ for some $c_\alpha \in \mathbb{Z}_{\geq 0}$. For $\lambda_X, \lambda_X' \in \Lambda_X^-$, we write $\lambda_X \leq_X \lambda_X'$ if $\lambda_X \leq_V \lambda_X'$. Note that here there is a slight inconsistency of notation as any $\lambda_X \in \Lambda_X^-$ satisfies $\lambda_X \geq_X 0$. The following lemma holds for any quasi-split reductive group.

Lemma 3.2.1.1. Let $\lambda_V, \lambda'_V \in \Lambda_V^+$. Then $K_V \lambda_V K_V \lambda_V^{-1} \cap K_V (\lambda'_V)^{-1} K_V \lambda'_V \subset K_V$.

Proof. Denote by * the convolution on $\mathcal{H}(\mathcal{U}(V), K_V)$. By [Gro98], $1_{K_V \lambda'_V K_V} * 1_{K_V \lambda_V K_V} (\lambda'_V \lambda_V) = 1$, which is equivalent to $\operatorname{vol}(K_V \lambda'_V K_V \cap \lambda'_V \lambda_V K_V \lambda_V^{-1} K_V) = 1$. As we have $\lambda'_V K_V \subset K_V \lambda'_V K_V \cap \lambda'_V \lambda_V K_V \lambda_V^{-1} K_V$, this implies that

$$\lambda'_V K_V = K_V \lambda'_V K_V \cap \lambda'_V \lambda_V K_V \lambda_V^{-1} K_V.$$

and in turn that

$$K_V \supset (\lambda'_V)^{-1} K_V \lambda'_V \cap \lambda_V K_V \lambda_V^{-1} K_V,$$

which is the desired inclusion.

We now describe the combinatorics between the $K \times K$ and $H \times K$ orbits in G. Recall that we can write $H = U(W) \ltimes N$, where N is the unipotent radical of P the parabolic subgroup of U(V) stabilizing the flag $0 \subset Ev_1 \subset \ldots \subset X$ of X.

Lemma 3.2.1.2. Let $\lambda, \lambda' \in \Lambda^-$. Assume that

$$K\lambda K \cap H\lambda' K \neq \emptyset. \tag{3.2.1.1}$$

Then $\lambda'_X \leq_X \lambda_X$. Moreover, $H \cap K \lambda K \lambda^{-1} \subset K$.

Proof. Let us prove the first point. The hypothesis (3.2.1.1) implies that

$$K_W \lambda_W K_W (\lambda'_W)^{-1} \cap K_V \lambda_V K_V (\lambda'_V)^{-1} N \neq \emptyset$$

It follows that there exists $n \in N$ such that

$${}^{\lambda'_W)^{-1}}n(\lambda'_W)^{-1}\lambda'_V \in K_W \lambda_W^{-1}K_V \lambda_V K_V.$$

Choose $\lambda_r \in \Lambda_r^+$ with $\lambda_r - \lambda_W \in \Lambda_V^+$ and $\lambda_r (\lambda'_W)^{-1} n \in K_V$. As λ_r commutes with U(W), we get

$$\lambda_r(\lambda'_W)^{-1}\lambda'_V \in K_V \lambda_r \lambda_W^{-1} K_V \lambda_V K_V.$$

By [Gro98, Equation (2.9)], we have

$$K_V \lambda_r \lambda_W^{-1} K_V \lambda_V K_V \subset \bigsqcup_{\substack{\tilde{\lambda}_V \in \Lambda_V^+ \\ \tilde{\lambda}_V \leq V \lambda_r - \lambda_W + \lambda_V}} K_V \tilde{\lambda}_V K_V,$$

so that $\lambda_r - \lambda'_W + \lambda'_V \leq_V \lambda_r - \lambda_W + \lambda_V$. This is exactly $\lambda'_X \leq_X \lambda_X$.

Assume now that $g_W \in U(W)$ and $n \in N$ are such that $(g_W, g_W n)\lambda \in K\lambda K$. Choose λ_r as before. Then we get

$$g_W n \in K_V \lambda_V K_V \lambda_V^{-1} \cap K_W \lambda_W K_W \lambda_W^{-1} n = K_V \lambda_V K_V \lambda_V^{-1} \cap K_W \lambda_r^{-1} \lambda_W K_W^{\lambda_r \lambda_W^{-1}} n \lambda_r \lambda_W^{-1}$$
$$\subset K_V \lambda_V K_V \lambda_V^{-1} \cap K_V \lambda_r^{-1} \lambda_W K_V \lambda_r \lambda_W^{-1}$$

Lemma 3.2.1.1 yields $g_W n \in K_V$, which concludes the proof.

3.2.2. Proof of Proposition 1.0.2.2. In this section, we prove that $(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ is a free $\mathcal{H}(G, K)$ -module of rank 1 (Proposition 1.0.2.2).

Lemma 3.2.2.1. We have a decomposition as a direct sum

$$\nu_{\mu,\psi}^{B_W^0} = \bigoplus_{\lambda_W \in \Lambda_W^+} \mathbb{C} \ \nu_{\mu,\psi}(\lambda_W^{-1})\phi^{\circ}.$$
(3.2.2.1)

Moreover, let $\tilde{\lambda}_X \in \Lambda_X^-$. Then

$$\nu_{\mu,\psi}^{\bar{\lambda}_X K_W \cap K_W \ltimes \bar{\lambda}_X \mathbb{H}(\mathbb{W})^0} = \bigoplus_{\substack{\lambda \in \Lambda^- \\ \lambda_X = \bar{\lambda}_X}} \mathbb{C} \ \nu_{\mu,\psi}(\lambda_W^{-1}) \phi^{\circ}.$$
(3.2.2.2)

Proof. We identify $\nu_{\mu,\psi}$ with its Schrödinger-lattice model which is included in the space $C_c^{\infty}(Y_+^*(E))$. By Lemma 3.1.3.1, every $\phi \in \nu_{\mu,\psi}^{B_W^0}$ is of the form $\phi_- \otimes 1_{Y_{m_+}^*(\mathcal{O}_E)}$ for $\phi_- \in C_c^{\infty}(Y_-^*(E))$. By (3.1.1.4), for any $\lambda_W \in \Lambda_W$ we know that $\nu_{\mu,\psi}(\lambda_W^{-1})\phi^{\circ}$ is, up to a scalar, the function $1_{\lambda_W Y_+^*(\mathcal{O}_E)}$. By doing induction on any finite subset of Λ_W equipped with the lexicographic order, we see that the family $(\nu_{\mu,\psi}(\lambda_W^{-1})\phi^{\circ})_{\lambda_W\in\Lambda_W}$ is a basis of $\nu_{\mu,\psi}^{T_W^0}$ so that the RHS of (3.2.2.1) is indeed a direct sum.

We now prove (3.2.2.1). Let $\phi \in \nu_{\mu,\psi}^{B_W^0}$ be non-zero. By the previous discussion, there exist a finite set $\Lambda_{\phi} \subset \Lambda_W$ and constants $c_{\lambda_W} \in \mathbb{C}^{\times}$ for $\lambda_W \in \Lambda_{\phi}$ such that $\phi = \sum_{\lambda_W \in \Lambda_{\phi}} c_{\lambda_W} \nu_{\mu,\psi}(\lambda_W^{-1})\phi^{\circ}$. Let $\beta \in \Delta_W$, and set $l := \min\{\langle \lambda_W, \beta \rangle \mid \lambda_W \in \Lambda_{\phi}\}$. We have to show that $l \ge 0$. Assume by contradiction that l < 0. Write

$$\phi_{l} := \sum_{\substack{\lambda \in \Lambda_{\phi} \\ \langle \lambda_{W}, \beta \rangle = l}} c_{\lambda_{W}} \nu_{\mu, \psi}(\lambda_{W}^{-1}) \phi^{\circ}, \quad \phi_{>l} := \sum_{\substack{\lambda \in \Lambda_{\phi} \\ \langle \lambda_{W}, \beta \rangle > l}} c_{\lambda_{W}} \nu_{\mu, \psi}(\lambda_{W}^{-1}) \phi^{\circ}.$$

Recall that G_{β} is the Levi subgroup of the parabolic subgroup P_{β} of U(W) corresponding to β . Set $K_{\beta} = K_W \cap G_{\beta} \subset H$. For $\lambda_W \in \Lambda_W$, the subgroup $\lambda_W^{-1} K_{\beta} \lambda_W$ only depends on $\langle \lambda_W, \beta \rangle$. If $\langle \lambda_W, \beta \rangle = k$, write $K_{\beta}^k = \lambda_W^{-1} K_{\beta} \lambda_W$. Set $\Lambda_{W,\beta} = \Lambda_W \cap G_{\beta}$. By the Cartan decomposition on G_{β} ,

$$G_{\beta} = K^l_{\beta} \Lambda_{W,\beta} K^l_{\beta}.$$

Note that ϕ_l is stabilized by K^l_{β} . Moreover, as the family $(\nu_{\mu,\psi}(\lambda_W^{-1})\phi^\circ)_{\lambda_W \in \Lambda_W}$ is free the stabilizer of ϕ_l in $\Lambda_{W,\beta}$ is trivial. Therefore, the stabilizer of ϕ_l in G_β is exactly K^l_{β} .

For any k, set $B_{\beta}^{k} = K_{\beta}^{k} \cap B_{W}$ so that $B_{\beta}^{k+1} \subseteq B_{\beta}^{k}$. Then for any $\lambda_{W} \in \Lambda_{W}$ with $\langle \lambda_{W}, \beta \rangle > l$, $\nu_{\mu,\psi}(\lambda_{W}^{-1})(\phi^{\circ})$ is stable by B_{β}^{l+1} . It follows that $\phi_{>l}$ is invariant by B_{β}^{l+1} , but by the previous discussion ϕ_{l} is not as its stabilizer in $B_{W} \cap G_{\beta}$ is B_{β}^{l} . This is a contradiction as $B_{\beta}^{l+1} \subset B_{W}^{0}$ because l < 0, which concludes the proof of (3.2.2.1).

We now show (3.2.2.2). The inclusion \supset is automatic so that we prove the reverse. Write $\tilde{\lambda}_X = \tilde{\lambda}_r + \tilde{\lambda}_W$ with $\tilde{\lambda}_r \in \Lambda_r^+$ and $\tilde{\lambda}_W \in \Lambda_W^+$. As $\tilde{\lambda}_X B_W^0 \subset \tilde{\lambda}_X K_W \cap K_W$ and $w_0 B_W^0 \subset \tilde{\lambda}_X K_W \cap K_W$ (where w_0 is the longest element in W_W), we see by (3.2.2.1) that

$$\nu_{\mu,\psi}^{\tilde{\lambda}_X K_W \cap K_W \ltimes^{\tilde{\lambda}_X \mathbb{H}(\mathbb{W})^0}} \subset \bigoplus_{\substack{\lambda_W \in \Lambda_W^-\\ \tilde{\lambda}_W + \lambda_W \in \Lambda_W^+}} \mathbb{C}\nu_{\mu,\psi}(\lambda_W^{-1})\phi^{\circ}.$$
(3.2.2.3)

To conclude, we have to show that any λ_W appearing in the RHS of (3.2.2.3) satisfies $\tilde{\lambda}_X + \lambda_W \in \Lambda_V^+$, which means that $\langle \tilde{\lambda}_X + \lambda_W, \alpha \rangle \geq 0$ for $\alpha = e_r - e_{r+1} \in \Delta_V$ (or $\alpha = e_{r'} - e_{r'+1} \in \Delta_V$ in the split case). This follows from similar considerations as in the proof of (3.2.2.1), noting that by (3.1.1.5) the stabilizer of ϕ° in $N_V \cap G_\alpha \subset H$ is $N_V^0 \cap G_\alpha$.

Proposition 3.2.2.2. We have a decomposition

$$\left(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi}\right)^{K} = \bigoplus_{\lambda \in \Lambda^{-}} \mathbb{C} \Phi_{\lambda},$$

where Φ_{λ} is the unique vector in $(\operatorname{ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ such that $\operatorname{supp}(\Phi_{\lambda}) \subset H\lambda K$ and $\Phi_{\lambda}(\lambda) = \phi^{\circ}$.

Proof. As Λ_X is a complete system of representatives of $H \setminus G/K$, by evaluating we have

$$\left(\mathrm{ind}_{H}^{G}\nu_{\mu,\psi}\right)^{K} = \bigoplus_{\tilde{\lambda}_{X} \in \Lambda_{X}} \nu_{\mu,\psi}^{\tilde{\lambda}_{X}K_{W} \cap K_{W} \ltimes \tilde{\lambda}_{X}(\mathbb{H}(\mathbb{W})^{0}U^{0})}$$

As ψ is unramified, $\nu_{\mu,\psi}^{\tilde{\lambda}_X U^0}$ is zero unless $\tilde{\lambda}_X U^0 \subset U^0$ which implies that $\tilde{\lambda}_X \in \Lambda_X^-$. The lemma now follows from Lemma 3.2.2.1 by letting Φ_{λ} be the unique vector in $(\operatorname{ind}_H^G \nu_{\mu,\psi})^K$ such that $\operatorname{supp}(\Phi_{\lambda}) \subset H\lambda K$ and $\Phi_{\lambda}(\lambda_X) = \nu_{\mu,\psi}(\lambda_W^{-1})\phi^\circ$.

We now end the proof of Proposition 1.0.2.2.

Proposition 3.2.2.3. The $\mathcal{H}(G, K)$ -module $(\operatorname{ind}_{H}^{G} \nu_{\mu,\psi})^{K}$ is free of rank 1 generated by Φ_{0} .

Proof. Let $\lambda \in \Lambda^-$. By Lemma 3.2.1.2 and Proposition 3.2.2.2, for every $\lambda' \in \Lambda^-$ there exists $c(\lambda, \lambda') \in \mathbb{C}$ with

$$R(1_{K\lambda^{-1}K})\Phi_0 = \sum_{\substack{\lambda' \in \Lambda^-\\\lambda'_X \le x\lambda_X}} c(\lambda, \lambda')\Phi_{\lambda'},$$

and the $c(\lambda, \lambda')$ are almost all zero. By the second part of Lemma 3.2.1.2, we have $R(1_{K\lambda^{-1}K})\Phi_0(\lambda) = \phi^\circ$, so that by Proposition 3.2.2.2 we see that

$$R(1_{K\lambda^{-1}K})\Phi_0 = \Phi_{\lambda} + \sum_{\substack{\lambda' \in \Lambda^-\\\lambda'_X < x\lambda_X}} c(\lambda, \lambda')\Phi_{\lambda'}.$$
(3.2.2.4)

As $r \geq 1$, the map $\Lambda^- \to \Lambda_X^-$ has finite fibers, and for every $\lambda_X \in \Lambda_X^-$ the set of $\lambda'_X \in \Lambda_X^-$ such that $\lambda'_X \leq_X \lambda_X$ is finite. Moreover, by the Cartan decomposition the family $(1_{K\lambda^{-1}K})_{\lambda\in\Lambda^-}$ is a basis of $\mathcal{H}(G, K)$. We conclude by induction on (3.2.2.4) and by Proposition 3.2.2.2 that the $\mathcal{H}(G, K)$ -module $(\operatorname{ind}_H^G \nu_{\mu,\psi})^K$ is generated by the torsion-free element Φ_0 , so that it is free of rank 1.

3.3. Proof of Theorem 1.0.2.1

We prove Theorem 1.0.2.1 under the assumption that $r \ge 1$ (see §3.4 for the r = 0 case).

Lemma 3.3.0.1. Let η be an unramified character of T_W . There is an isomorphism of H-representations (which also holds for r = 0)

$$S_{\overline{\mu}\eta}: \left(I_{B_W}^{\mathrm{U}(W)}\eta\right) \otimes \overline{\nu}_{\mu,\psi} \to \overline{\psi}_U \boxtimes \left(I_{B_J}^J \overline{\mu} \eta \overline{\psi}\right) \tag{3.3.0.1}$$

which satisfies $S_{\overline{\mu}\eta}(\Phi_{\eta}^{\circ}\otimes\phi^{\circ})=\Phi_{\overline{\mu}\eta}^{\circ,J}$ and $S_{\overline{\mu}\eta}(\Psi_{w,\eta}\otimes\phi^{\circ})=\Psi_{w,\overline{\mu}\eta}^{J}$.

Proof. The existence of $S_{\overline{\mu}\eta}$ follows from (2.1.9.1), (3.1.1.3) and the isomorphisms of J-representations

$$\left(I_{B_W}^{\mathrm{U}(W)} \eta \right) \otimes \overline{\nu}_{\mu,\psi} \simeq \mathrm{Ind}_{B_W \ltimes \mathbb{H}(\mathbb{W})}^J \left(\delta_{B_W}^{\frac{1}{2}} \eta \otimes (\overline{\nu}_{\mu,\psi})_{|B_W \ltimes \mathbb{H}(\mathbb{W})} \right) \simeq \mathrm{Ind}_{B_W \ltimes \mathbb{H}(\mathbb{W})}^J \left(\delta_{B_J}^{\frac{1}{2}} \eta \otimes I_{B_J}^{B_W \ltimes \mathbb{H}(\mathbb{W})} \overline{\mu} \overline{\psi} \right) \\ \simeq I_{B_J}^J \overline{\mu} \eta \overline{\psi}.$$

Explicitly, we have $S_{\overline{\mu}\eta}(f \otimes \phi)(hg_W) = (\overline{\nu}_{\mu,\psi}(hg_W)\phi)(0)f(g_W)$ for $h \in \mathbb{H}(\mathbb{W})$ and $g_W \in U(W)$. This implies the last point.

Proposition 3.3.0.2. There exists a surjective morphism of \mathbb{C} -algebras

$$f_J \in \mathcal{H}(J, K_J) \mapsto f_J^{\psi} \in \mathcal{H}(\mathcal{U}(W), K_W)$$
 (3.3.0.2)

such that for all $f_J \in \mathcal{H}(J, K_J)$ we have $\xi^J_{\overline{\mu}\eta\overline{\psi}}(f_J) = \xi_{\eta}(f_J^{\psi})$. Moreover, there exists an isomorphism of $\mathcal{H}(\mathrm{U}(V), K_V) \otimes \mathcal{H}(J, K_J)$ -modules

$$\mathcal{W} \in \mathrm{WS} \mapsto \Phi_{\mathcal{W}} \in \left(\mathrm{Ind}_{H}^{G} \nu_{\mu,\psi}\right)^{K}.$$
 (3.3.0.3)

Remark 3.3.0.3. Let $\mathcal{L}_H \in \operatorname{Hom}_H(I_{B_V}^{\mathrm{U}(V)}\chi \otimes I_{B_W}^{\mathrm{U}(W)}\eta \otimes \overline{\nu}_{\mu,\psi},\mathbb{C})$, and let $\mathcal{L}_H^{\vee} \in \operatorname{Hom}_H(I_{B_V}^{\mathrm{U}(V)}\chi \otimes I_{B_W}^{\mathrm{U}(W)}\eta,\nu_{\mu,\psi})$ be the associated operator by duality. If $\mathcal{W}_{\chi,\overline{\mu}\eta}$ is of the form $\mathcal{W}_{\chi,\overline{\mu}\eta}(g) = \mathcal{L}_H(R(g)\Phi_{\chi}^{\circ}\otimes \Phi_{\eta}^{\circ}\otimes \phi^{\circ})$ for Φ_{χ}° and Φ_{η}° spherical, then we have

$$\Phi_{\mathcal{W}_{\chi,\overline{\mu}\eta}}(g_V,g_W) = \mathcal{L}_H^{\vee}(R(g_V)\Phi_{\chi}^{\circ} \otimes R(g_W)\Phi_{\eta}^{\circ}), \quad g_V \in \mathrm{U}(V), \quad g_W \in \mathrm{U}(W).$$
(3.3.0.4)

Proof. We first describe the map $\mathcal{W} \mapsto \Phi_{\mathcal{W}}$ and show that it is an isomorphism of \mathbb{C} -vector spaces. Let $\mathcal{W} \in WS$. For every $g_V \in U(V)$, consider the function $\phi_{\mathcal{W},g_V} : h \in \mathbb{H}(\mathbb{W}) \mapsto \mathcal{W}(hg_V)$. Then $\phi_{\mathcal{W},g_V} \in \operatorname{ind}_{\mathbb{H}(\mathbb{W}^0)}^{\mathbb{H}(\mathbb{W})} \psi$, which we identify with the lattice model of $\nu_{\mu,\psi}$ by §3.1.2. We now define

$$\Phi_{\mathcal{W}}: (g_V, g_W) \in G \mapsto \nu_{\mu, \psi}(g_W) \phi_{\mathcal{W}, g_W^{-1}g_V} \in \nu_{\mu, \psi}.$$

This yields the desired morphism WS $\rightarrow (\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$. For $\Phi \in (\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$, define

$$\mathcal{W}_{\Phi}: g_V \mapsto \Phi(g_V)(0) = (\Phi(g_V), \phi_{\text{latt}}^\circ)_{\text{latt}} \in \mathbb{C}, \qquad (3.3.0.5)$$

where we identify $\Phi(g_V)$ with an element of $\operatorname{Ind}_{\mathbb{H}(\mathbb{W}^0)}^{\mathbb{H}(\mathbb{W})}\psi$. Then $\mathcal{W}_{\Phi} \in WS$. Indeed, the key point is that by (3.1.2.1) for every $w \in W^0$, $k_W \in K_W$ and $g_V \in U(V)$ we have

$$\mathcal{W}_{\Phi}(h(w,0)k_Wg_V) = \Phi((k_W,k_W)(g_V,1)(1,k_W^{-1}))(w) = (\nu_{\mu,\psi}(k_W)\Phi(g_V,1))(0) = \mathcal{W}_{\Phi}(g_V).$$

The maps $\mathcal{W} \mapsto \Phi_{\mathcal{W}}$ and $\Phi \mapsto \mathcal{W}_{\Phi}$ are inverse of each other.

We now describe the morphism (3.3.0.2). Let $C_c^{\infty}((K_J Z, \psi) \setminus J)$ be the compact induction $\operatorname{ind}_{K_J Z}^J, \psi$. Define $\mathcal{H}(J, K_J Z, \psi)$ to be the Hecke algebra of functions J which transform by ψ under left and right translations by $K_J Z$, and which have compact support modulo the action of $K_J Z$. Note that $K_J Z$ is an open subgroup of the locally compact totally disconnected group J which is compact modulo its center. We therefore have an isomorphism of \mathbb{C} -algebras

$$\operatorname{End}_J\left(C_c^{\infty}((K_J Z, \psi) \backslash J)\right) \simeq \mathcal{H}(J, K_J Z, \psi), \qquad (3.3.0.6)$$

where $\mathcal{H}(J, K_J Z, \psi)$ acts on $C_c^{\infty}((K_J Z, \psi) \setminus J)$ by left-translations. Using (3.1.2.2) we see that

$$\operatorname{ind}_{K_J Z}^J \psi \simeq \operatorname{ind}_{K_W \ltimes \mathbb{H}(\mathbb{W})}^J \left(\operatorname{ind}_{K_W \ltimes \mathbb{H}(\mathbb{W}^0)}^{K_W \ltimes \mathbb{H}(\mathbb{W})} \psi \right) \simeq \operatorname{ind}_{K_W \ltimes \mathbb{H}(\mathbb{W})}^J (\nu_{\mu,\psi})_{|K_W \ltimes \mathbb{H}(\mathbb{W})} \simeq \operatorname{ind}_{K_W \ltimes \mathbb{H}(\mathbb{W})}^J (1) \otimes \nu_{\mu,\psi}$$

$$\simeq C_c^\infty (K_W \setminus U(W)) \otimes \nu_{\mu,\psi},$$

(3.3.0.7)

where J acts on U(W) by the composition of R with the projection $J \to U(W)$. As $\nu_{\mu,\psi}$ is irreducible (it is already as a $\mathbb{H}(\mathbb{W})$ -representation), Schur's lemma implies that we have an isomorphism of \mathbb{C} -algebras

$$\operatorname{End}_J(C_c^{\infty}((K_JZ,\psi)\backslash J)) \simeq \operatorname{End}_{U(W)}(C_c^{\infty}(K_W\backslash U(W))) \simeq \mathcal{H}(U(W),K_W).$$
(3.3.0.8)

Composing (3.3.0.8) with (3.3.0.6) and the surjective morphism $\mathcal{H}(K, K_J) \to \mathcal{H}(K, K_J Z, \psi)$ given by integrating against $\overline{\psi}$ on Z, we obtain the desired map (3.3.0.2).

Let us show that $\mathcal{W} \mapsto \Phi_{\mathcal{W}}$ is an morphism of $\mathcal{H}(\mathcal{U}(V), K_V) \otimes \mathcal{H}(J, K_J)$ -modules. For any $f_V \in \mathcal{H}(\mathcal{U}(V), K_V)$, we have $R(f_V)\Phi_{\mathcal{W}} = \Phi_{R(f_V)\mathcal{W}}$. Let $\mathcal{W} \in WS$ and $f_J \in \mathcal{H}(J, K_J)$. It is enough to show that for any $g_W \in \mathcal{U}(W)$ we have

$$R(f_J^{\psi})\Phi_{\mathcal{W}}(1,g_W) = \Phi_{L(f_J)\mathcal{W}}(1,g_W).$$

Note that $j \mapsto \mathcal{W}(j) \in \operatorname{ind}_{K_J Z}^J \psi$. We can assume that this element is sent to $f_W \otimes \phi \in C_c(K_W \setminus U(W)) \otimes \nu_{\mu,\psi}$ by (3.3.0.7). By definition, we have

$$L(f_J)\mathcal{W} = L(f_J^{\psi})f_W \otimes \phi$$
, and $\Phi_{\mathcal{W}}(1, g_W) = f_W(g_W^{-1})\phi$.

This concludes that $R(f_J^{\psi})\Phi_{\mathcal{W}} = \Phi_{L(f_J)\mathcal{W}}$ and therefore that (3.3.0.3) is indeed an isomorphism of modules.

Finally, let us prove that $\xi^J_{\overline{\mu}n\overline{\psi}}(f_J) = \xi_\eta(f^\psi_J)$ for every $f_J \in \mathcal{H}(J, K_J)$. We have

$$\int_{Z} f_J(zg_W h)\overline{\psi}(z)dz = f_J^{\psi}(g_W)\nu_{\mu,\psi}(g_W h)\phi_{\text{latt}}^{\circ}(0), \quad g_W \in \mathcal{U}(W), \ h \in \mathbb{H}(\mathbb{W}).$$
(3.3.0.9)

By Lemma 3.3.0.1, the isomorphism of *J*-representations $(I_{B_W}^{U(W)}\eta) \otimes \overline{\nu}_{\mu,\psi} \to I_{B_J}^J \overline{\mu} \eta \overline{\psi}$ sends $\Phi_{\eta}^{\circ} \otimes \phi^{\circ}$ to $\Phi_{\overline{\mu}\eta}^{\circ,J}$. Let $f_J \in \mathcal{H}(J, K_J)$. It follows from (3.1.1.5), (3.1.2.3) and (3.3.0.9) that

$$\begin{split} \xi^{J}_{\overline{\mu}\eta\overline{\psi}}(f_{J}) &= \int_{\mathrm{U}(W)} \int_{W} \int_{Z} f_{J}(h(w,z)g_{W}) \Phi^{\circ}_{\eta}(g_{W})\overline{\nu}_{\mu,\psi}(h(w,z)g_{W})\phi^{\circ}(0)dzdwdg_{W} \\ &= \int_{\mathrm{U}(W)} f^{\psi}_{J}(g_{W}) \Phi^{\circ}_{\eta}(g_{W}) \int_{W} \nu_{\mu,\psi}(h(w,0)g_{W})\phi^{\circ}_{\mathrm{latt}}(0)\overline{\nu}_{\mu,\psi}(h(w,0)g_{W})\phi^{\circ}(0)dwdg_{W} \\ &= \int_{\mathrm{U}(W)} f^{\psi}_{J}(g_{W}) \Phi^{\circ}_{\eta}(g_{W}) \int_{Y^{*}_{+}(E)} \overline{\nu}_{\mu,\psi}(g_{W})\phi^{\circ}(y^{*}) \int_{Y_{-}(E)} \nu_{\mu,\psi}(h(y^{*},0)g_{W})\phi^{\circ}_{\mathrm{latt}}(y)dydy^{*}dg_{W} \\ &= \int_{\mathrm{U}(W)} f^{\psi}_{J}(g_{W}) \Phi^{\circ}_{\eta}(g_{W})(\overline{\nu}_{\mu,\psi}(g_{W})\phi^{\circ},\overline{\nu}_{\mu,\psi}(g_{W})\phi^{\circ})_{\overline{\nu}}dg_{W}. \end{split}$$

As $\overline{\nu}_{\mu,\psi}$ is unitary, we see that this last expression reduces to $\xi_{\eta}(f_J^{\psi})$, which concludes the proof. \Box

We now end the proof of Theorem 1.0.2.1. Let χ and η be unramified characters of T_V and T_W respectively. The space $WS_{\chi,\overline{\mu}\eta}$ is the $\xi_{\chi} \otimes \xi^J_{\overline{\mu}\eta\overline{\psi}}$ -eigenspace in WS (see §2.3). By Proposition 3.3.0.2, it is therefore enough to prove than for any character ξ of $\mathcal{H}(G, K)$ the ξ -eigenspace $(\operatorname{Ind}_H^G \nu_{\mu,\psi})^K_{\xi}$ has dimension at most one. As $\nu_{\mu,\psi}$ is unitary and admissible, its contragredient $\widetilde{\nu_{\mu,\psi}}$ is $\overline{\nu}_{\mu,\psi}$. By [Ren10, Théorème III.2.7] we have the isomorphism of $\mathcal{H}(G)$ -modules

$$\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi}\simeq (\operatorname{ind}_{H}^{G}\overline{\nu}_{\mu,\psi})\widetilde{.}$$

Let $L \in \left((\operatorname{ind}_{H}^{G} \overline{\nu}_{\mu,\psi})^{\sim} \right)_{\xi}^{K}$. Then L is determined by its values on $(\operatorname{ind}_{H}^{G} \overline{\nu}_{\mu,\psi})^{K}$, and therefore by $L(\Phi_{0})$ by Proposition 3.2.2.3. This shows that $(\operatorname{Ind}_{H}^{G} \nu_{\mu,\psi})_{\xi}^{K}$ has dimension at most one. In §4 and §5 we build a non-zero element in this space. This concludes the proof of Theorem 1.0.2.1.

3.4. The r = 0 case

We now assume that r = 0 and briefly explain how to adapt the proof of Theorem 1.0.2.1. Set $\Lambda_X^- = \Lambda_V^+$ embedded in the first factor of G. Define $\Delta_J = \Delta_V \cup \{-e_1\}$ in the inert case, and $\Delta_J = \Delta_V \cup \{-e_1, e_n\}$ in the split case. Set

$$\Lambda_J^- = \{ \lambda_V \in \Lambda_V \mid \forall \alpha \in \Delta_J, \ \langle \lambda_V, \alpha \rangle \ge 0 \}.$$

For any $\lambda_V \in \Lambda_X^-$, the proof of Lemma 3.2.2.1 shows that

$$\nu_{\mu,\psi}^{\lambda_V K_V \cap K_V \ltimes^{\lambda_V} \mathbb{H}(\mathbb{W})^0} = \bigoplus_{\substack{\lambda \in \Lambda_V^- \\ \lambda_V + \lambda \in \Lambda_J^-}} \mathbb{C} \ \nu_{\mu,\psi}(\lambda^{-1}) \phi^{\circ}.$$

But $\Lambda_J^- = \{0\}$, so that $\nu_{\mu,\psi}^{\lambda_V K_V \cap K_V \ltimes^{\lambda_V} \mathbb{H}(\mathbb{W})^0} = \mathbb{C} \ \overline{\nu}_{\mu,\psi}(\lambda_V)\phi^\circ$. As all the other results of §3.2 remain valid, we prove as in Proposition 3.2.2.3 that the $\mathcal{H}(\mathcal{U}(V), K_V) \otimes \mathcal{H}(J, K_J)$ -module $(\operatorname{ind}_J^{\mathcal{U}(V) \times J} \overline{\nu}_{\mu,\psi})^{K_V \times K_J}$ is free of rank 1. An easy adaptation of Proposition 3.3.0.2 shows that we have an isomorphism WS $\simeq (\text{Ind}_J^{U(V) \times J} \nu_{\mu,\psi})^{K_V \times K_J}$ with the same properties as in the $r \ge 1$ case. Theorem 1.0.2.1 now follows from argument of §3.3. Note that (3.3.0.4) still holds in this case.

Integral expression and analytic continuation 4.

In this section we produce a non-trivial element $\mathcal{W}^{I}_{\chi,\eta}$ in WS_{χ,η}, first for (χ,η) in a non-empty open subset of $(\mathbb{C}^{\times})^{n_{-}} \times (\mathbb{C}^{\times})^{m_{-}}$ using Frobenius reciprocity and a convergent integral pairing, and then for (χ, η) generic by analytic continuation.

Integral pairing 4.1.

4.1.1. For A and B subsets of $\{1, \ldots, n\}$ and $\{1, \ldots, m\}$ respectively with |A| = |B| and $g \in U(V)$, we define $\Delta_{A,B}(g)$ to be the determinant of the $A \times B$ minor of g. We will also use this notation if $g \in J$ by identifying it with a $(m+2) \times (m+2)$ matrix.

If $r \ge 1$, define for $1 \le k \le n_+$ the set $I_k = \{n+1-k, \ldots, n\}$, and for $1 \le l \le m_+$ the sets $J_l = \{1, \dots, l\}$ and $J'_l = \{1, \dots, r-1, r+1, \dots, r+l\}$. If r = 0, we use rather $I_k = \{n+2-k, \dots, n+1\}$, $I'_l = \{1, n+3-l, \dots, n+1\}$ and $J_l = \{2, \dots, 1+l\}$. Set

$$\alpha_k(g) = \Delta_{I_k, J_k}(g) \text{ and } \beta_l(g) = \Delta_{I_{r+l-1}, J_l'}(g) \text{ if } r \ge 1 \text{ (resp. } \beta_l(g) = \Delta_{I_l', J_l}(g) \text{ if } r = 0 \text{)}.$$
(4.1.1.1)

We define $\beta_{m_{+}}$ to be $\beta_{m_{+}}$ in the inert odd case, and 0 otherwise. It is easily checked that these applications enjoy the following properties for all g. Recall that $\mu_{+}^{*} = h(1_{Y_{+}^{*}}, 0)$ (see §2.1.7).

- For $n_1, n_2 \in N_V$, $\alpha_k(n_1gn_2) = \alpha_k(g)$ $(n_2 \in N_J \text{ if } r = 0)$,
- $\alpha_k(t_V(\mathbf{t})gt_V(\mathbf{s})) = \begin{cases} \prod_{i=1}^k {}^c t_i^{-1} s_i \cdot \alpha_k(g) & \text{inert case,} \\ \prod_{i=1}^k t_{n-i+1} s_i \cdot \alpha_k(g) & \text{split case.} \end{cases}$
- If $w \in W_V$, then $\alpha_k(w) \neq 0$ for all k if and only if $w = w_0$.
- For any $n_V \in N_V$, $\beta_l(n_V q) = \beta_l(q)$.
- For any $n_J \in N_J$, $u \in U$ and $l \leq m_-$, $\beta_l(gn_J u) = \beta_l(g)$, and in the odd inert case, for $y \in Y_{-}(E), z \in F, n_W \in N_W, \tilde{\beta}_{m_+}(gh(y,z)n_W u) = \tilde{\beta}_{m_+}(g).$
- $\beta_l(t_V(\mathbf{t})gt_W(\mathbf{s})) = \begin{cases} \prod_{i=1}^r ct_i^{-1} \cdot \prod_{i=1}^{l-1} ct_{r+i}^{-1} \cdot \prod_{i=1}^l s_i \cdot \beta_l(g) \\ \prod_{i=1}^r t_{n-i+1} \cdot \prod_{i=1}^{l-1} t_{n-r-i+1} \cdot \prod_{i=1}^l s_i \cdot \beta_l(g) \end{cases}$ inert case, split case.

• For $y^* \in Y^*_{-}(E)$ and $l \le m_{-}$, $\beta_l(w_0h(y^*, 0)) = \pm y^*_l$, and in the inert odd case, for $y_{m_+} \in E$ we have $\tilde{\beta}_{m_+}(w_0h(y^* + y_{m_+}w_{m_+}, 0)) = \pm y_{m_+}$.

Proposition 4.1.1.1. We have the equality

$$B_V w_0 \mu_+^* B_J U = \left\{ g \left| \alpha_k(g) \neq 0, \ \beta_l(g) \neq 0, \ \left| \frac{\tilde{\beta}_{m_+}(g) \prod_{i=1}^{m_-} \alpha_{r+i-1}(g)}{\alpha_{n_-}(g) \prod_{i=1}^{m_-} \beta_i(g)} \right| \le 1, \ 1 \le k \le n_+, \\ 1 \le l \le m_- \right\} \right\}$$

where $g \in U(V)$ if $r \ge 1$ and $g \in J$ if r = 0.

Proof. Assume $r \ge 1$ and that we are in the inert odd case. For $\mathbf{a} = (a_i) \in \{0, 1\}^{m_-}$, write $y_{\mathbf{a}}^* = \sum a_i w_i^*$. Then the Bruhat decomposition on U(V) yields

$$U(V) = \bigcup_{w \in W_V} \bigcup_{\mathbf{a} \in \{0,1\}^{m_-}} \bigcup_{y_{m_+} \in E/U(1)} B_V w h(y_{\mathbf{a}}^* + y_{m_+} w_{m_+}, 0) B_J U,$$

where U(1) acts on E by multiplication. The result now follows from the above properties of the α_k and β_l . The proof is the same for r = 0 and for the other cases.

4.1.2. Let χ and η be unramified characters of T_V and T_W . For $g \in U(V)$ (or $g \in J$ for r = 0) set

$$Y_{\chi,\eta}(g) := \chi^{-1}(\delta_{B_V}^{\frac{1}{2}})(b_V)(\eta \overline{\psi} \delta_{B_J}^{-\frac{1}{2}})(b_J) \overline{\psi}_{\mathrm{U}}(u) \quad \text{if } g = b_V w_0 \mu_+^* b_J u \in B_V w_0 \mu_+^* B_J U, \tag{4.1.2.1}$$

and $Y_{\chi,\eta}(g) = 0$ if $g \notin B_V w_0 \mu_+^* B_J U$. From section 4.1.1 we see that for $g = t_V(\mathbf{t}) n_V w_0 \mu_+^* t_W(\mathbf{s}) n_J u$ we have, writing α_i (resp. β_j) for $\alpha_i(g)$ (resp. $\beta_j(g)$)

$$|Y_{\chi,\eta}(g)| = \prod_{i=1}^{r-1} \left| \chi_i \chi_{i+1}^{-1} \left| .\right|_E^{-1} \right| (\alpha_i) \prod_{i=r}^{n-1} \left| \chi_i \eta_{i-r+1}^{-1} \left| .\right|_E^{-\frac{1}{2}} \right| (\alpha_i) \prod_{j=1}^{m-1} \left| \eta_j \chi_{j+r}^{-1} \left| .\right|_E^{-\frac{1}{2}} \right| (\beta_j) \times \begin{cases} \left| \chi_{n-1} \left| .\right|_E^{-\frac{1}{2}} \right| (\alpha_{n-1}) \left| \chi_{n-1} \left| \chi_{n-1} \left| .\right|_E^{-\frac{1}{2}} \right| (\alpha_{n-1}) \left| \chi_{n-1} \left|$$

in the inert case, depending on whether n is even or odd, and

$$\begin{aligned} |Y_{\chi,\eta}(g)| &= \left|\chi_1^{-1} \left|.\right|_F^{\frac{n-1}{2}}\right| \left(\det(g)\right) \prod_{i=1}^r \left|\chi_i \chi_{i+1}^{-1} \left|.\right|_F^{-1}\right| \left(\alpha_{n-i}\right) \prod_{i=r+1}^{r'} \left|\chi_{i+1}^{-1} \eta_{r'-i+1}^{-1} \left|.\right|_F^{-\frac{1}{2}}\right| \left(\alpha_{n-i}\right) \\ &\times \prod_{i=r'+1}^{n-1} \left|\chi_i \chi_{i+1}^{-1} \left|.\right|_F^{-1}\right| \left(\alpha_{n-i}\right) \prod_{j=1}^m \left|\eta_j \chi_{r'-j+1} \left|.\right|_F^{-\frac{1}{2}}\right| \left(\beta_j\right). \end{aligned}$$

in the split case. We now define \mathcal{U} to be the non empty open subset of (χ, η) satisfying

$$\begin{cases} \left| \chi_{i}\chi_{i+1}^{-1} \right| < q_{E}^{-1} & 1 \leq i < r, \\ \left| \chi_{i}\eta_{i-r+1}^{-1} \right| < q_{E}^{-\frac{1}{2}} & r \leq i < n_{-}, \\ \left| \chi_{i+1}^{-1}\eta_{i-r+1} \right| < q_{E}^{-\frac{1}{2}} & r \leq i < n_{-}, \\ \left| \chi_{i+1}^{-1}\eta_{i-r+1}^{-1} \right| < q_{E}^{-\frac{1}{2}} & r \leq i < n_{-}, \\ \left| \chi_{i}\chi_{i+1}^{-1} \right| < q_{F}^{-1} & 1 \leq i \leq r, \\ \left| \chi_{i}\chi_{i+1}^{-1} \right| < q_{F}^{-1} & r < i \leq r', \\ \left| \chi_{i}\chi_{i+1}^{-1} \right| < q_{F}^{-1} & r' < i < n, \\ \left| \chi_{i}\chi_{i+1}^{-1} \right| < q_{F}^{-1} & 1 \leq j \leq m, \\ \left| \eta_{j}\chi_{r'-j+1} \right| < q_{F}^{-\frac{1}{2}} & 1 \leq j \leq m, \\ \end{cases}$$

in the inert and split case respectively. It follows from Proposition 4.1.1.1 that for any $(\chi, \eta) \in \mathcal{U}$ the function $Y_{\chi,\eta}$ is continuous on U(V).

4.1.3. For $(\chi, \eta) \in \mathcal{U}, f_V \in C_c^{\infty}(\mathcal{U}(V)), f_J \in C_c^{\infty}(J)$, consider the absolutely convergent integral

$$\mathcal{L}_{\chi,\eta}(F_{\chi}(f_V) \otimes F_{\eta}^J(f_J)) = \int_{\mathcal{U}(V)} \int_J f_V(g_V) f_J(g_J) Y_{\chi,\eta}(g_V g_J^{-1}) dg_V dg_J.$$
(4.1.3.1)

Then $\mathcal{L}_{\chi,\eta}$ defines a non zero element in $\operatorname{Hom}_H(I_{B_V}^{U(V)}\chi\otimes\overline{\psi}_U.I_{B_J}^J\eta\overline{\psi},\mathbb{C})$, and it follows that

$$\mathcal{W}^{I}_{\chi,\eta}(g) := \begin{cases} \mathcal{L}_{\chi,\eta}(R(g).\Phi^{\circ}_{\chi} \otimes \Phi^{\circ}_{\eta}), & g \in \mathrm{U}(V), & \text{if } r \ge 1; \\ \mathcal{L}_{\chi,\eta}(\Phi^{\circ}_{\chi} \otimes R(g^{-1}).\Phi^{\circ}_{\eta}), & g \in J, & \text{if } r = 0. \end{cases}$$
(4.1.3.2)

is a Whittaker–Shintani function. We will explicitly compute $\mathcal{W}_{\chi,n}^{I}(1)$ in Proposition 5.5.2.1.

4.2. Analytic section

By Lemma 3.3.0.1 we have

$$\operatorname{Hom}_{H}(I_{B_{V}}^{\mathrm{U}(V)}\chi\otimes I_{B_{W}}^{\mathrm{U}(W)}(\mu\eta)\otimes\overline{\nu}_{\mu,\psi},\mathbb{C}) = \operatorname{Hom}_{H}(I_{B_{V}}^{\mathrm{U}(V)}\chi\otimes\overline{\psi}_{U}.I_{B_{J}}^{J}(\eta\overline{\psi}),\mathbb{C}).$$
(4.2.0.1)

For χ and η generic, $I_{B_V}^{U(V)}\chi \otimes I_{B_W}^{U(W)}(\mu\eta)$ is irreducible. By the multiplicity one result [GGP12, Corollary 16.3] we have

$$\dim \operatorname{Hom}_{H}(I_{B_{V}}^{\mathrm{U}(V)}\chi \otimes I_{B_{W}}^{\mathrm{U}(W)}(\mu\eta) \otimes \overline{\nu}_{\mu,\psi}, \mathbb{C}) \leq 1.$$

$$(4.2.0.2)$$

We can now apply Bernstein's Theorem [Ban98] (see also [She13a, Lemma 5.1]).

Proposition 4.2.0.1. There exists a dense subset \mathcal{V} of $(\mathbb{C}^{\times})^{n_{-}} \times (\mathbb{C}^{\times})^{m_{-}}$ such that for every f_{V} and f_{J} the map $(\chi, \eta) \mapsto \mathcal{L}_{\chi,\eta}(F_{\chi}(f_{V}) \otimes F_{\eta}^{J}(f_{J}))$ defined on \mathcal{U} by (4.1.3.1) extends to a rational function on \mathcal{V} .

In particular, we obtain a non zero $\mathcal{W}_{\chi,\eta}^{I} \in WS_{\chi,\eta}$ for (χ,η) generic. Let $\Phi_{\chi,\eta}^{I} \in (\mathrm{Ind}_{H}^{G}\nu_{\mu,\psi})^{K}$ $((\mathrm{Ind}_{J}^{\mathrm{U}(V)\times J}\nu_{\mu,\psi})^{K_{V}\times K_{J}}$ if r=0) be the associated function by Proposition 3.3.0.2.

5. Formula for the normalized Whittaker–Shintani function

We prove the formulas for Whittaker–Shintani functions of Theorem 1.0.2.3 and Proposition 1.0.2.4.

5.1. Iwahori-invariant vectors

5.1.1. Let χ be a generic unramified character of T_V . For $\alpha \in \Sigma_{V,\mathrm{nd}}^+$, let $c_{\alpha}(\chi)$ be the constant defined in [Cas80, Section 3], i.e.

$$c_{e_{a}\pm e_{b}}(\chi) = \begin{cases} \frac{1-q_{E}^{-1}\chi_{a}\chi_{b}^{\pm 1}}{1-\chi_{a}\chi_{b}^{\pm 1}} & \text{inert case,} \\ \frac{1-q_{F}^{-1}\chi_{a}\chi_{b}^{-1}}{1-\chi_{a}\chi_{b}^{-1}} & \text{split case,} \end{cases} \text{ and } \begin{cases} c_{2e_{a}}(\chi) = \frac{1-q_{F}^{-1}\chi_{a}}{1-\chi_{a}} & \text{inert even case,} \\ c_{e_{a}}(\chi) = \frac{(1-q_{F}^{-1}\chi_{a})(1+q_{F}^{-1}\chi_{a})}{1-\chi_{a}^{2}} & \text{inert odd case,} \end{cases}$$

where we only consider $e_a - e_b$ in the split case. For $w \in W_V$, set

$$c_w^V(\chi) = \prod_{\substack{\alpha \in \Sigma_{V,\mathrm{nd}}^+ \\ w\alpha < 0}} c_\alpha(\chi).$$
(5.1.1.1)

There exists an intertwining operator $T_{w,\chi}: I_{B_V}^{\mathrm{U}(V)}\chi \to I_{B_V}^{\mathrm{U}(V)}w\chi$ introduced in [Cas80, Section 3]. Define the normalization $\overline{T}_{w,\chi}:=c_w^V(\chi)^{-1}T_{w,\chi}$. Recall that we have introduced a spherical vector Φ_{χ}° and Iwahori-fixed vectors $\Psi_{w,\chi}$ in (2.2.0.1) and (2.2.0.2) respectively. By [Cas80, Theorem 3.1] it satisfies the relations $\overline{T}_{w_1,w_2\chi} \circ \overline{T}_{w_2,\chi} = \overline{T}_{w_1w_2,\chi}$ and $\overline{T}_{w,\chi}(\Phi_{\chi}^{\circ}) = \Phi_{w\chi}^{\circ}$. Using the Bruhat decomposition and evaluating Φ_{χ}° on W_V , we have

$$\Phi_{\chi}^{\circ} = \frac{1}{\operatorname{vol}(I_V)} \sum_{w \in W_V} c_{w_0}^V(w\chi) \overline{T}_{w^{-1},w\chi} \Psi_{1,w\chi},$$

and it follows from [KMS03, Proposition 1.10] that for $\lambda_V \in \Lambda_V^-$

$$R(I_V\lambda_V I_V)\Phi_{\chi}^{\circ} = \frac{\operatorname{vol}(I_V\lambda_V I_V)}{\operatorname{vol}(I_V)} \sum_{w \in W_V} c_{w_0}^V(w\chi) \cdot (w\chi) \delta_{B_V}^{-\frac{1}{2}}(\lambda_V) \overline{T}_{w^{-1},w\chi} \Psi_{1,w\chi}.$$
(5.1.1.2)

5.1.2. Let η be a generic unramified character of T_W and let $w \in W_W$. We adapt the preceding discussion to $I_{B_J}^J \eta \overline{\psi}$. Denote by $T_{w,\mu\eta} : I_{B_W}^{\mathrm{U}(W)} \mu \eta \to I_{B_W}^{\mathrm{U}(W)} w(\mu \eta)$ the map built in §5.1.1. Then

$$T_{w,\eta}^J := S_{w\eta} \circ (T_{w,\mu\eta} \otimes \mathrm{id}) \circ S_\eta^{-1}$$

provides an intertwining operator $I_{B_I}^J \eta \overline{\psi} \to I_{B_I}^J w \eta \overline{\psi}$ (where S_η is defined in (3.3.0.1)). Define $c_w^W(\eta)$ as in (5.1.1.1), and consider the normalized version $\overline{T}_{w,\eta}^J = c_w^W(\mu\eta)^{-1}T_{w,\eta}^J$ and recall that $\phi^\circ := 1_{Y^*_+(\mathcal{O}_E)} \in \overline{\nu}_{\mu,\psi}$. By Lemma 3.3.0.1 we have $S_\eta(\Phi^\circ_{\mu\eta} \otimes \phi^\circ) = \Phi^{\circ,J}_\eta$ and $S_\eta(\Psi_{w,\mu\eta} \otimes \phi^\circ) = \Psi^J_{w,\eta}$, where $\Phi_{\mu\eta}^{\circ}$ and $\Psi_{w,\mu\eta}$ are the spherical and Iwahori fixed vectors in $I_{B_W}^{U(W)}\mu\eta$ introduced in (2.2.0.1) and (2.2.0.2) respectively. In particular, $\overline{T}_{w,\eta}^J(\Phi_{\eta}^{\circ,J}) = \Phi_{w,\eta}^{\circ,J}$. By the same proof as Lemma 2.2.0.1, we now see that any $f \in (I_{B_I}^J \eta \overline{\psi})^{I_J}$ is supported on $B_J W_W I_J$ so that we also obtain

$$\Phi_{\eta}^{\circ,J} = \frac{1}{\text{vol}(I_J)} \sum_{w \in W_W} c_{w_0}^W(w\mu\eta) \overline{T}_{w^{-1},w\eta}^J \Psi_{1,w\eta}^J, \qquad (5.1.2.1)$$

and therefore for $\lambda_W \in \Lambda_W^-$

$$R(I_W \lambda_W I_W) \Phi_{\eta}^{\circ, J} = \frac{\operatorname{vol}(I_W \lambda_W I_W)}{\operatorname{vol}(I_J)} \sum_{w \in W_W} c_{w_0}^W(w\mu\eta) . (w\mu\eta) \delta_{B_W}^{-\frac{1}{2}}(\lambda_W) \overline{T}_{w^{-1}, w\eta}^J \Psi_{1, w\eta}^J.$$
(5.1.2.2)

γ -factors 5.2.

For generic $\chi, \eta \in (\mathbb{C}^{\times})^{n_{-}} \times (\mathbb{C}^{\times})^{m_{-}}$, set $\Gamma_{1}^{V}(\chi) = \frac{c_{w_{0}}^{V}(\chi)}{\mathbf{d}_{V}(\chi)}$, $\Gamma_{1}^{W}(\eta) = \frac{c_{w_{0}}^{W}(\eta)}{\mathbf{d}_{W}(\eta)}$ and let $\Gamma_{2}(\chi, \eta)$ be $\begin{cases} \prod_{j=1}^{m_{-}} \left(\left(\prod_{i=1}^{n_{-}} L_{E}(\frac{1}{2}, \chi_{i}\eta_{j}) L_{E}(\frac{1}{2}, \chi_{i}^{-1}\eta_{j}) \right) \left(\prod_{i=1}^{r+j-1} \frac{L_{E}(\frac{1}{2}, \chi_{i}\eta_{j}^{-1})}{L_{E}(\frac{1}{2}, \chi_{i}^{-1}\eta_{j})} \right) \right), \\ \prod_{j=1}^{m_{-}} \left(\left(\prod_{i=1}^{n_{-}} L_{E}(\frac{1}{2}, \chi_{i}\eta_{j}) L_{E}(\frac{1}{2}, \chi_{i}^{-1}\eta_{j}) \right) \left(\prod_{i=1}^{r+j-1} \frac{L_{E}(\frac{1}{2}, \chi_{i}\eta_{j}^{-1})}{L_{E}(\frac{1}{2}, \chi_{i}^{-1}\eta_{j})} \right) \right) \prod_{i=1}^{n_{-}} L_{F}(1, -\chi_{i}) \prod_{i=1}^{m_{-}} L_{F}(1, \eta_{i}), \\ \prod_{i+j \leq r'+1} L_{F}(\frac{1}{2}, \chi_{i}\eta_{j}) \prod_{i+j > r'+1} L_{F}(\frac{1}{2}, \chi_{i}^{-1}\eta_{j}^{-1}), \end{cases}$ (5.2.0.1) in the inert even case, inert odd case and split case respectively. Set $\Gamma(\chi,\eta) := \Gamma_1^V(\chi)\Gamma_1^W(\mu\eta)\Gamma_2(\chi,\eta)$.

For $w = (w_V, w_W) \in W_G$, set $\overline{T}_{w,\chi,\eta} = \overline{T}_{w_V,\chi} \otimes \overline{T}_{w_W,\eta}^J$. The proof of Theorem 1.0.2.3 relies on the following result.

Theorem 5.2.0.1. For generic χ and η we have for every $w \in W_G$ the functional equation

$$\frac{\mathcal{L}_{\chi,\eta}}{\Gamma(\chi,\eta)} = \frac{\mathcal{L}_{w_V\chi,w_W\eta} \circ T_{w,\chi,\eta}}{\Gamma(w_V\chi,w_W\eta)}$$

By the multiplicity one Theorem (4.2.0.2), for generic χ and η there exists $\gamma(\chi, \eta, w_V, w_W) \in \mathbb{C}$ such that

$$\mathcal{L}_{w_V\chi,w_W\eta} \circ \overline{T}_{w,\chi,\eta} = \gamma(\chi,\eta,w_V,w_W)\mathcal{L}_{\chi,\eta}.$$

It is enough to prove that $\gamma(\chi, \eta, w_V, w_W) = \frac{\Gamma(w_V \chi, w_W \eta)}{\Gamma(\chi, \eta)}$ for $(w_V, w_W) = (w_\alpha, 1)$ or $(1, w_\beta)$ with $\alpha \in \Delta_V$ and $\beta \in \Delta_W$. By Proposition 4.2.0.1, γ is a rational function so that we now assume that $(\chi, \eta) \in \mathcal{U}$ is generic and use the integral expression (4.1.3.1). Theorem 5.2.0.1 will follow from (5.2.0.1), Propositions 5.3.1.2 and 5.4.1.1, and elementary computations.

5.3. Computation of $\gamma(\chi, \eta, w_{\alpha}, 1)$

5.3.1. Let $\alpha = e_i - e_{i+1} \in \Delta_V$. Define $I_J^1 := I_W \ltimes Y^0_+ Y^{*,1}_- Z^0$ and $\Psi^{1,J}_\eta = F^J_\eta(1_{I_J^1})$. By [Cas80, Theorem 3.4] we have

$$\overline{T}_{w_{\alpha},\chi}(\Psi_{1,\chi}+\Psi_{w_{\alpha},\chi})=\Psi_{1,w_{\alpha}\chi}+\Psi_{w_{\alpha},w_{\alpha}\chi},$$

so that

$$\gamma(\chi,\eta,w_{\alpha},1) = \frac{\mathcal{L}_{w_{\alpha}\chi,\eta}(R(\mu_{+}^{*}w_{0})(\Psi_{1,w_{\alpha}\chi}+\Psi_{w_{\alpha},w_{\alpha}\chi})\otimes\Psi_{\eta}^{1,J})}{\mathcal{L}_{\chi,\eta}(R(\mu_{+}^{*}w_{0})(\Psi_{1,\chi}+\Psi_{w_{\alpha},\chi})\otimes\Psi_{\eta}^{1,J})},$$
(5.3.1.1)

granted the denominator is non-zero, where we recall that $\mu_{+}^{*} = h(1_{Y_{+}^{*}}, 0)$.

Remark 5.3.1.1. In (5.3.1.1) we implicitly assume that $r \ge 1$. Otherwise, we compute instead

$$\gamma(\chi,\eta,w_{\alpha},1) = \frac{\mathcal{L}_{w_{\alpha}\chi,\eta}(R(w_{0})(\Psi_{1,w_{\alpha}\chi} + \Psi_{w_{\alpha},w_{\alpha}\chi}) \otimes R((\mu_{+}^{*})^{-1})\Psi_{\eta}^{1,J})}{\mathcal{L}_{\chi,\eta}(R(w_{0})(\Psi_{1,\chi} + \Psi_{w_{\alpha},\chi}) \otimes R((\mu_{+}^{*})^{-1})\Psi_{\eta}^{1,J})}$$

As the calculations are identical, we ignore this issue to simplify the proof. We will work under the assumption $r \ge 1$ throughout this section, and leave the easy modifications for r = 0 to the reader.

Theorem 5.2.0.1 in that case therefore follows from the next proposition which we prove in the rest of this subsection.

Proposition 5.3.1.2. The value of $\mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)(\Psi_{1,\chi}+\Psi_{w_\alpha,\chi})\otimes\Psi_{\eta}^{1,J})$ is

$$\operatorname{vol}(I_V)\operatorname{vol}(I_J^1) \times \begin{cases} q_E L_E(1,\chi_i\chi_{i+1}^{-1})^{-1} & 1 \leq i < r & inert \ case, \\ q_F L_F(1,\chi_i\chi_{i+1}^{-1})^{-1} & \begin{cases} 1 \leq i \leq r, \\ or \ r' < i < n \end{cases} \text{ split \ case,} \\ (q_E - 1) \frac{L_E(\frac{1}{2},\chi_i\eta_{i-r+1}^{-1})L_E(\frac{1}{2},\chi_{i+1}^{-1}\eta_{i-r+1})}{L_E(1,\chi_i\chi_{i+1}^{-1})} & r \leq i < n_- & inert \ case, \\ (q_F - 1) \frac{L_F(\frac{1}{2},\chi_i\eta_{i-r'+1})L_F(\frac{1}{2},\chi_{i+1}^{-1}\eta_{i-r+1})}{L_F(1,\chi_i\chi_{i+1}^{-1})} & r < i \leq r' & split \ case, \\ q_F L_F(1,\chi_{n_-})^{-1} & i = n_- & inert \ even \ case, \\ q_E q_F L_E(1,\chi_{n_-})^{-1} & i = n_- & inert \ odd \ case. \end{cases}$$

5.3.2. The following lemma will also be used to compute $\gamma(\chi, \eta, 1, w_{\beta})$. It is a straightforward adaptation of [She13b, Lemma 2.8.2].

Lemma 5.3.2.1. Let $\alpha \in \Delta_V$, $\beta \in \Delta_W$. Then for any values of ± 1

$$I_V w_0^{\pm 1} \mu_+^{*,\pm 1} I_J^1 U^0 = B_V^0 w_0^{\pm 1} \mu_+^{*,\pm 1} B_J^0 U^0, \qquad (5.3.2.1)$$

$$I_V w_0^{\pm 1} \mu_+^{*,\pm 1} I_W U^0 \subset B_V^0 w_0^{\pm 1} B_J^0 Y_+^{*,5} U^0,$$
(5.3.2.2)

$$I_V w_{\alpha} I_V w_0^{\pm 1} \mu_+^{*,\pm 1} I_J^1 U^0 = B_V^0 w_{\alpha} N_{\alpha}^0 w_0^{\pm 1} \mu_+^{*,\pm 1} B_J^0 U^0, \qquad (5.3.2.3)$$

$$I_V w_0^{\pm 1} \mu_+^{*,\pm 1} I_J w_\beta I_J U^0 = B_V^0 w_0^{\pm 1} N_\beta^0 w_\beta Y_+^{*,0} B_J^0 U^0.$$
(5.3.2.4)

Lemma 5.3.2.2. We have

$$\mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)\Psi_{1,\chi}\otimes\Psi_\eta^{1,J})=\mathrm{vol}(I_V)\mathrm{vol}(I_J^1).$$

Proof. Using the integral expression (4.1.3.1) we have

$$\mathcal{L}_{\chi,\eta}(R(\mu_{+}^{*}w_{0})\Psi_{1,\chi}\otimes\Psi_{\eta}^{1,J}) = \int_{I_{V}\times I_{J}^{1}}Y_{\chi,\eta}(g_{V}w_{0}^{-1}\mu_{+}^{*,-1}g_{J})dg_{V}dg_{J}$$

The result follows from the invariance properties of $Y_{\chi,\eta}$ in (4.1.2.1) and Lemma 5.3.2.1 (5.3.2.1).

5.3.3. We now compute $\mathcal{L}_{w_{\alpha}\chi,\eta}(R(\mu_{+}^{*}w_{0})\Psi_{1,w_{\alpha}\chi} \otimes \Psi_{\eta}^{1,J})$. Using (4.1.2.1) and Lemma 5.3.2.1 (5.3.2.3) this is

$$\int_{I_V w_\alpha I_V} \int_{I_J^1} Y_{\chi,\eta}(g_V w_0^{-1} \mu_+^{*,-1} g_J) dg_V dg_J = \operatorname{vol}(I_V w_\alpha I_V) \operatorname{vol}(I_J^1) \int_{N_\alpha^0} Y_{\chi,\eta}(w_\alpha n_\alpha w_0 \mu_+^*) dn_\alpha.$$
(5.3.3.1)

Denote by I_{α} this last integral.

Case SL₂ We first treat the case $D(G_{\alpha}) = \text{SL}_2(E)$ or $\text{SL}_2(F)$, so that N_{α} is isomorphic to E or F. We write α^{\vee} for the corresponding coroot of the maximal torus of $D(G_{\alpha})$ (which is a cocharacter over E in the first case). For $t \neq 0$ we have $w_{\alpha}n_{\alpha}(t) = \alpha^{\vee}(-t^{-1})n_{\alpha}(-t)n_{-\alpha}(t^{-1})$. Therefore

$$I_{\alpha} = \int_{|t| \le 1} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}}) (\alpha^{\vee}(t^{-1})) Y_{\chi,\eta}(w_{0} n_{w_{0}(-\alpha)}(t^{-1}) \mu_{+}^{*}) dt.$$

Case SL₂ **1.** In the inert case with $1 \le i < r - 1$, or the split case with $1 \le i < r$ or $r' + 1 < i \le n$, then $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = \mu_+^* n_{w_0(-\alpha)}(t^{-1})$. In the inert case with i = r - 1, or the split case with i = r' + 1, $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = \mu_+^* n_{w_0(-\alpha)}(t^{-1})u$ with $\psi_U(u) = 1$. Finally, in the split case with i = r, $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = \mu_+^* n_{w_0(-\alpha)}(t^{-1})h(0, t^{-1})$. Therefore we always have

$$I_{\alpha} = \int_{|t| \le 1} (\chi^{-1} \delta_{B_V}^{\frac{1}{2}}) (\alpha^{\vee}(t^{-1})) \overline{\psi_E}(t^{-1}) dt.$$
 (5.3.3.2)

Case SL₂ 2. In the inert case with $r \leq i < n_{-}$ set j = i - r, and in the split case with $r + 1 \leq i \leq r'$, set j = r' - i. Let $\bar{}$ be $c \in \text{Gal}(E/F)$ in the inert case, and identity in the split case. Define $\mathbf{t}_{j+1}(t) = t_W((1 - t^{-1})w_{j+1}) \in T_W$ and $y_{j+1}^* = \sum_{k=1}^{m_-} w_k^* - \bar{t}^{-1}w_{j+1}^*$. Then $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = h(y_1^*, 0)$ if j = 0, and $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = h(y_{j+1}^*, 0)n_{w_0(-\alpha)}(t^{-1})$ otherwise. But $h(y_{j+1}^*, 0) = \mathbf{t}_{j+1}(t)^{-1}\mu_+^*\mathbf{t}_{j+1}(t)$ so that

$$I_{\alpha} = \int_{|t| \le 1} (\chi^{-1} \delta_{B_V}^{\frac{1}{2}}) (\alpha^{\vee}(t^{-1})) (\chi^{-1} \delta_{B_V}^{\frac{1}{2}}) ({}^{w_0} \mathbf{t}_{j+1}(t)^{-1}) (\eta \delta_{B_J}^{-\frac{1}{2}}) (\mathbf{t}_{j+1}(t)) dt.$$
(5.3.3.3)

Case SL₂ **3.** In the inert even case with $i = n_{-}$, $\alpha = 2e_{n_{-}}$ and $D(G_{\alpha})$ is SL₂(F). Then $n_{w_0(-\alpha)}(t^{-1})\mu_+^* = \mu_+^*h(t^{-1}w_{m_{-}},t^{-1})n_{w_0(-\alpha)}(t^{-1})$ and

$$I_{\alpha} = \int_{|t| \le 1} (\chi^{-1} \delta_{B_V}^{\frac{1}{2}}) (\alpha^{\vee}(t^{-1})) \overline{\psi_E}(t^{-1}) dt.$$
 (5.3.3.4)

Case SU₃ Assume now that $D(G_{\alpha})$ is SU₃(F), which occurs in the inert odd case with $i = n_{-}$. Replace $e_{n_{-}}$ by $\alpha = 2e_{n_{-}}$. In that case, recall that

$$n_{\alpha}(x_1, x_2) = \begin{pmatrix} 1 & -\tau \overline{x_1} & x_2 \\ & 1 & x_1 \\ & & 1 \end{pmatrix}, \ x_1, x_2 \in E, \ \operatorname{Tr}_{E/F}(\frac{x_2}{\tau}) = -N_{E/F}(x_1).$$
(5.3.3.5)

Then for $x_2 \neq 0$ we have $w_{\alpha}n_{\alpha}(x_1, x_2)w_0 \in T_V(\mathcal{O}_E)n_{\alpha}(\frac{x_1}{x_2}, \frac{-1}{x_2})\alpha^{\vee}(\frac{1}{x_2})w_0n_{\alpha}(\frac{x_1}{x_2}, \frac{-1}{x_2})$ and furthermore $n_{\alpha}(x_1, x_2)\mu_+^*n_{\alpha}(x_1, x_2)^{-1} = \mu_+^*h(x_1w_{m_+} + (-\tau \overline{x_1} + x_2)w_{m_-}, \frac{1}{2}\mathrm{Tr}(x_2))$. Therefore by Proposition 4.1.1.1 we have

$$Y_{\chi,\eta}(w_{\alpha}n_{\alpha}(x_{1},x_{2})w_{0}\mu_{+}^{*}) = (\chi^{-1}\delta_{B_{V}}^{\frac{1}{2}})\left(\alpha^{\vee}(x_{2}^{-1})\right)Y_{\chi,\eta}\left(w_{0}\mu_{+}^{*}h\left(\frac{x_{1}}{x_{2}}w_{m_{+}},-\frac{1}{2}\mathrm{Tr}(\frac{1}{x_{2}})\right)\right) \quad (5.3.3.6)$$
$$= \begin{cases} 0 & \text{if } \left|\frac{x_{1}}{x_{2}}\right| > 1\\ (\chi\delta_{B_{V}}^{-\frac{1}{2}})(\alpha^{\vee}(x_{2}))\overline{\psi}\left(-\frac{1}{2}\mathrm{Tr}(\frac{1}{x_{2}})\right) & \text{if } \left|\frac{x_{1}}{x_{2}}\right| \le 1 \end{cases}$$

Let $x \in E^{\times}$ and $y \in F^{\times}$. In our coordinates, $x_1 = x$ and $x_2 = y - \tau \frac{N(x)}{2}$ and we are computing

$$I_{\alpha} = \int_{|x|_{E} \le 1} \int_{|y|_{F} \le 1} (\chi \delta_{B_{V}}^{-\frac{1}{2}}) \left(\alpha^{\vee}(x_{2}) \right) \overline{\psi} \left(-\frac{1}{2} \operatorname{Tr}(\frac{1}{x_{2}}) \right) \mathbf{1}_{|x|_{E} \le |x_{2}|_{E}} dy dx.$$

We now split the integral.

- If $|x|_E = 1$, then the integrand is 1 so that we get $\frac{q_E 1}{q_E}$.
- If $|x|_E < 1$ and $|y|_E \le |N(x)|_E$, then $|x_2|_E = |N(x)|_E < |x|_E$, and the integrand is 0.
- If $|x|_E < 1$ and $|y|_E > |N(x)|_E$, we need $|x|_E \le |x_2|_E = |y|_E$ otherwise the integrand is zero. In this range, $\overline{\psi}\left(-\frac{1}{2}\operatorname{Tr}(\frac{1}{x_2})\right) = \overline{\psi}(-y^{-1})$.

Therefore,

$$I_{\alpha} = \frac{q_E - 1}{q_E} + \int_{|x|_E < 1} \int_{\substack{y \in F^{\times} \\ |x|_E \le |y|_E \le 1}} (\chi \delta_{B_V}^{-\frac{1}{2}}) \left(\alpha^{\vee}(y)\right) \overline{\psi}(y^{-1}) dy dx.$$
(5.3.3.7)

The integrals (5.3.3.2), (5.3.3.3), (5.3.3.4) and (5.3.3.7) can now be computed by standard techniques ([She13a, Lemmas 8.5, 8.9 and 8.11]) and Proposition 5.3.1.2 follows from Lemma 5.3.2.2, (5.3.3.1) and the value of $\operatorname{vol}(I_V w_{\alpha} I_V)$. Note that $(\chi, \eta) \in \mathcal{U}$ implies that these integrals converge.

5.4. Computation of $\gamma(\chi, \eta, 1, w_{\beta})$

5.4.1. Let $\beta = e_j - e_{j+1} \in \Delta_W$. As in section 5.3.1 we have

$$\gamma(\chi,\eta,1,w_{\beta}) = \frac{\mathcal{L}_{\chi,w_{\beta}\eta}(R(\mu_{+}^{*}w_{0})\Psi_{1,\chi}\otimes(\Psi_{1,w_{\beta}\eta}^{J}+\Psi_{w_{\beta},w_{\beta}\eta}^{J}))}{\mathcal{L}_{\chi,\eta}(R(\mu_{+}^{*}w_{0})\Psi_{1,\chi}\otimes(\Psi_{1,\eta}^{J}+\Psi_{w_{\beta},\eta}^{J}))}.$$

Proposition 5.4.1.1. Recall that Δ'_{T_W} was defined in §2.1.10. Set

$$\Pi(\chi,\eta) = \Delta_{T_W}^{',-1} \times \begin{cases} \prod_{\substack{j=1 \ m}}^{m_-} L_E(\frac{1}{2},\eta_j\chi_{r+j}^{-1}) & inert \ case, \\ \prod_{j=1}^{m} L_F(\frac{1}{2},\eta_j\chi_{r'-j+1}^{-1}) & split \ case. \end{cases}$$

The value of $\mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)\Psi_{1,\chi}\otimes(\Psi_{1,\eta}^J+\Psi_{w_\beta,\eta}^J))$ is

$$\operatorname{vol}(I_V)\operatorname{vol}(I_J)\Pi(\chi,\eta) \times \begin{cases} q_E \frac{L_E(\frac{1}{2},\eta_j\chi_{r+j+1}^{-1})L_E(\frac{1}{2},\chi_{r+j}\eta_{j+1}^{-1})}{L_E(1,\eta_j\eta_{j+1}^{-1})L_E(1,\chi_{r+j}\chi_{r+j+1}^{-1})} & 1 \leq j < m_- & inert \ case, \\ q_F \frac{L_F(\frac{1}{2},\eta_j\chi_{r'-j})L_F(\frac{1}{2},\eta_{j+1}^{-1}\chi_{r'-j+1})}{L_F(1,\eta_j\eta_{j+1}^{-1})L_F(1,\chi_{r'-j}\chi_{r'-j+1}^{-1})} & 1 \leq j < m & split \ case, \\ q_F \frac{L_F(1,\eta_j\eta_{j+1}^{-1})L_F(1,\chi_{r-j}\chi_{r'-j+1})}{L_F(1,\chi_{m-1})L_F(1,\chi_{m-1})} & j = m_- & inert \ even \ case, \\ q_E q_F \frac{L_F(1,\chi_m,\eta_n)}{L_E(1,\eta_m)L_E(1,\chi_{n-1})} & j = m_- & inert \ odd \ case. \end{cases}$$

5.4.2.

Lemma 5.4.2.1.

$$\mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)\Psi_{1,\chi}\otimes\Psi_{1,\eta}^J)=\mathrm{vol}(I_V)\mathrm{vol}(I_J)\Pi(\chi,\eta).$$

Proof. By Lemma 5.3.2.1 (5.3.2.4) we have

$$\mathcal{L}_{\chi,\eta}(R(\mu_{+}^{*}w_{0})\Psi_{1,\chi}\otimes\Psi_{1,\eta}^{J}) = \operatorname{vol}(I_{V})\operatorname{vol}(I_{J})\int_{Y_{-}^{*}(\mathcal{O}_{E})}Y_{\chi,\eta}(w_{0}h(y^{*},0))dy^{*}.$$

For y^* with all coordinates being non zero, we have $h(y^*, 0) = t_W(y^*)^{-1} \mu_+^* t_W(y^*)$ so that

$$\int_{Y_{-}^{*}(\mathcal{O}_{E})} Y_{\chi,\eta}(w_{0}h(y^{*},0))dy^{*} = \int_{Y_{-}^{*}(\mathcal{O}_{E})} (\chi^{-1}\delta_{B_{V}}^{\frac{1}{2}})(^{w_{0}}t_{W}(y^{*})^{-1})(\eta\delta_{B_{J}}^{-\frac{1}{2}})(t_{W}(y^{*}))dy^{*}.$$

This can be computed by decomposing as a product (see [She13a, Lemma 8.9]).

5.4.3. By Lemma 5.3.2.1 (5.3.2.4) again we have

$$\mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)\Psi_{1,\chi}\otimes\Psi_{w_\beta,\eta}^J) = \operatorname{vol}(I_V)\operatorname{vol}(I_Jw_\beta I_J) \int_{N_\beta^{0,\prime}} \int_{Y_-^*(\mathcal{O}_E)} Y_{\chi,\eta}(w_0 n_\beta' w_\beta h(y^*,0)) dy^* dn_\beta.$$
(5.4.3.1)

Denote by I_{β} this last double integral.

Case SL₂. Assume that $D(G_{\beta})$ is SL₂(F) or SL₂(E). We write again β^{\vee} for the corresponding coroot of the maximal torus of $D(G(\beta))$, so that $n'_{\beta}(t)w_{\beta} = \beta^{\vee}(-t)n'_{-\beta}(t)n'_{\beta}(-t^{-1})$. Therefore

$$I_{\beta} = \int_{|t| \le 1} \int_{Y_{-}^{*}(\mathcal{O}_{E})} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}}) ({}^{w_{0}} \beta^{\vee}(t)) Y_{\chi,\eta}(w_{0} n_{\beta}'(t^{-1}) h(y^{*}, 0) n_{\beta}'(t^{-1})^{-1}) dy^{*} dt.$$

Case SL₂ **1.** In the inert case with $j < m_-$ or in the split case with any j, set $y^*(t) = y^* - y_j \overline{t}^{-1} w_{j+1}^*$ so that when all coordinates are non zero

$$n'_{\beta}(t^{-1})h(y^*,0)n'_{\beta}(t^{-1})^{-1} = h(y^*(t),0) = t_W(y^*(t))^{-1}\mu_+^*t_W(y^*(t))$$

Therefore

$$I_{\beta} = \int_{|t| \le 1} \int_{Y_{-}^{*}(\mathcal{O}_{E})} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}}) ({}^{w_{0}} \beta^{\vee}(t) {}^{w_{0}} t_{W}(y^{*}(t))^{-1}) (\eta \delta_{B_{J}}^{-\frac{1}{2}}) (t_{W}(y^{*}(t))) dy^{*} dt.$$
(5.4.3.2)

Case SL₂ **2.** In the inert even case with $j = m_{-}$ we have $\beta = 2e_{m_{-}}$ and

$$n'_{\beta}(t^{-1})h(y^*,0)n'_{\beta}(t^{-1})^{-1} = h(y^* + t^{-1}y^*_{m_-}w_{m_-},0) = t^{-1}_W(y^*)\mu^*_+t_W(y^*)h(t^{-1}y^*_{m_-}w_{m_-},t^{-1}N_{E/F}(y^*_{m_-})).$$

We get

$$I_{\beta} = \int_{|t| \le 1} \int_{Y_{-}^{*}(\mathcal{O}_{E})} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}}) ({}^{w_{0}} \beta^{\vee}(t) {}^{w_{0}} t_{W}(y^{*})^{-1}) (\eta \delta_{B_{J}}^{-\frac{1}{2}}) (t_{W}(y^{*})) \overline{\psi}(t^{-1} N_{E/F}(y_{m_{-}}^{*})) dy^{*} dt.$$

$$(5.4.3.3)$$

Case SU₃ Assume that $D(G_{\alpha})$ is SU₃, which occurs in the inert odd case with $j = m_{-}$. Take $\beta = e_{m_{-}}$. Using the same coordinates as in (5.3.3.6) and by a similar computation we see that $Y_{\chi,\eta}(w_0 n'_{\beta}(x_1, x_2) w_{\beta} h(y^*, 0))$ is

$$\begin{split} & (\chi^{-1}\delta_{B_{V}}^{\frac{1}{2}}) \left({}^{w_{0}}t_{W}(y^{*})^{-1}\beta^{\vee}(x_{2})^{-1}\right) (\eta\delta_{B_{J}}^{-\frac{1}{2}})(t_{W}(y^{*}))Y_{\chi,\eta} \left(w_{0}\mu_{+}^{*}h\left(y_{m_{-}}^{*}\frac{x_{1}}{x_{2}}w_{m_{+}}, -\frac{1}{2}\mathrm{Tr}(\frac{N_{E/F}(y_{m_{-}}^{*})}{x_{2}})\right)\right) \\ & = \begin{cases} 0 & \text{if } \left|y_{m_{-}}^{*}\frac{x_{1}}{x_{2}}\right| > 1, \\ (\chi^{-1}\delta_{B_{V}}^{\frac{1}{2}}) \left(t_{W}(y^{*})\beta^{\vee}(x_{2})^{-1}\right) (\eta\delta_{B_{J}}^{-\frac{1}{2}})(t_{W}(y^{*}))\overline{\psi} \left(-\frac{1}{2}\mathrm{Tr}(\frac{N_{E/F}(y_{m_{-}}^{*})}{x_{2}})\right) & \text{if } \left|y_{m_{-}}^{*}\frac{x_{1}}{x_{2}}\right| \le 1. \end{cases}$$

 I_{β} is a product of an integral over y_1^*, \ldots, y_{m-1}^* , which is computed as in Lemma 5.4.2.1, and of

$$\int_{|z|_{E} \leq 1} \int_{\substack{|x|_{E} \leq 1 \\ |y|_{F} \leq 1}} (\chi^{-1} \delta_{B_{V}}^{\frac{1}{2}}) \left(\beta^{\vee}(x_{2}^{-1}z)\right) (\eta \delta_{B_{J}}^{-\frac{1}{2}}) (\beta^{\vee}(z)) \overline{\psi} \left(-\frac{1}{2} \operatorname{Tr}(\frac{N_{E/F}(z)}{x_{2}})\right) \mathbf{1}_{|zx|_{E} \leq |x_{2}|_{E}} dy dx dz,$$

$$(5.4.3.4)$$

where we recall that $y \in F$, $x_2 = y - \tau \frac{N_{E/F}(x)}{2}$, and we have replaced $y_{m_-}^*$ by z to ease notations. We now split the integral.

• If $|y|_E \leq |N_{E/F}(x)|_E$, then $|x_2|_E = |N_{E/F}(x)|_E$ and the integrand is zero unless $|z|_E \leq |x|_E$. In that case $\overline{\psi}\left(-\frac{1}{2}\mathrm{Tr}(\frac{N_{E/F}(z)}{x_2})\right) = 1$ and we are simply integrating a product of multiplicative characters. • If $|y|_E > |N_{E/F}(x)|_E$, then the integrand is zero unless $|zx|_E \le |y|_E$. In this case we see that $\overline{\psi}\left(-\frac{1}{2}\mathrm{Tr}(\frac{N_{E/F}(z)}{x_2})\right) = \overline{\psi}\left(-\frac{N_{E/F}(z)}{y}\right).$

The integrals (5.4.3.2), (5.4.3.3) and (5.4.3.4) can now be computed by splitting further depending on valuations thanks to [She13a, Lemmas 8.5, 8.9 and 8.11], and Proposition 5.4.1.1 follows from Lemma 5.4.2.1 and (5.4.3.1). This concludes the proof of Theorem 5.2.0.1.

5.5. Proofs of Theorem 1.0.2.3 and Proposition 1.0.2.4

In this section we work under the assumption that $r \ge 1$. The case r = 0 follows from the modification explained in Remark 5.3.1.1.

Let us first remark that the conventions of Theorem 1.0.2.3 are different from the ones we are currently using as we are inducing from B rather than $B^+ = B_V^- \times B_W$, and because moreover we are working with $WS_{\chi,\eta}$ instead of $WS_{\chi,\overline{\mu}\eta}$. Set

$$\mathbf{d}_{V}(\chi) = \prod_{\alpha \in \Sigma_{V,\mathrm{nd}}^{+}} \frac{1}{1 - \langle \chi, \alpha^{\vee} \rangle}, \quad \mathbf{d}_{W}(\eta) = \prod_{\beta \in \Sigma_{W,\mathrm{nd}}^{+}} \frac{1}{1 - \langle \eta, \beta^{\vee} \rangle}.$$
 (5.5.0.1)

Then $\mathbf{d}(\chi \boxtimes \eta) = \mathbf{d}_V(w_0\chi)\mathbf{d}_W(\eta)$. Moreover, because $w_{0,V}B_V = B_V^-$ the identification of T with the canonical torus T_G coming from $T \subset B$ and $T \subset B^+$ differ by a conjugation by $w_{0,V}$. In particular, after a change of variable we see that (1.0.2.4) in Theorem 1.0.2.3 is equivalent to, for $\lambda \in \Lambda^-$,

$$\frac{(\Phi_{\chi,\eta}(\lambda),\phi^{\circ})_{\nu}}{(\Phi_{\chi,\eta}(1),\phi^{\circ})_{\nu}} = \frac{\Delta_{\mathrm{U}(W)}}{\Delta_{T_{W}}} \sum_{w \in W_{G}} \mathbf{b}(w_{V}\chi,w_{W}\eta) \mathbf{d}_{V}(w_{V}\chi) \mathbf{d}_{W}(w_{W}\mu\eta) \left((w_{0}w_{V}\chi \boxtimes w_{W}\mu\eta) \,\delta_{B^{+}}^{-\frac{1}{2}} \right)(\lambda).$$
(5.5.0.2)

5.5.1. Recall that $\Phi^I_{\chi,\eta}$ was defined in §4.2. By Remark 3.3.0.3, for any $\phi \in \overline{\nu}_{\mu,\psi}$ we have

$$(\Phi_{\chi,\eta}^{I}(g_{V},g_{W}),\phi)_{\nu} = \mathcal{L}_{\chi,\eta}\left(R(g_{V})\Phi_{\chi}^{\circ}\otimes S_{\eta}\left(R(g_{W})\Phi_{\mu\eta}^{\circ}\otimes\phi\right)\right), \quad (g_{V},g_{W})\in G.$$
(5.5.1.1)

Proposition 5.5.1.1. For every $\lambda \in \Lambda^-$ and generic (χ, η) we have

$$\frac{(\Phi_{\chi,\eta}^{I}(\lambda_{V},\lambda_{W}),\phi^{\circ})_{\nu}}{\Delta_{T_{W}}^{\prime,-1}\Gamma(\chi,\eta)} = \sum_{w\in W_{G}} \mathbf{b}(w_{V}\chi,w_{W}\eta)\mathbf{d}_{V}(w_{V}\chi)\mathbf{d}_{W}(w_{W}\mu\eta)\left((w_{0}w_{V}\chi\boxtimes w_{W}\mu\eta)\,\delta_{B^{+}}^{-\frac{1}{2}}\right)(\lambda).$$

Proof. Write $\lambda = (\lambda_V, \lambda_W) \in \Lambda_V^+ \times \Lambda_W^-$. Note that ${}^{w_0}\lambda_V^{-1}B_V^0{}^{w_0}\lambda_V \subset B_V^0$ and $\lambda_W^{-1}B_W^0\lambda_W^0 \subset B_W^0$. By (5.3.2.2) we have

$$(R(\mu_+^*w_0\mathbf{1}_{I_V}w_0\lambda_V I_V)\otimes R(\mathbf{1}_{I_W}\lambda_W I_W)\Phi_{\chi,\eta}^I(1),\phi^\circ)_\nu = \operatorname{vol}(I_V^{w_0}\lambda_V I_V)\operatorname{vol}(I_W\lambda_W I_W)(\Phi_{\chi,\eta}^I(\lambda_V,\lambda_W),\phi^\circ)_\nu$$
(5.5.1.2)

But by (5.1.1.2), (5.1.2.2) and (5.5.1.1) the LHS is

$$\frac{\operatorname{vol}(I_V^{w_0}\lambda_V I_V)\operatorname{vol}(I_W\lambda_W I_W)}{\operatorname{vol}(I_V)\operatorname{vol}(I_W)} \sum_{(w_V,w_W)\in W_G} c_{w_0}^V(w_V\chi)c_{w_0}^W(w_V\mu\eta)(w_0w_V\chi)(\lambda_V)(w_W\mu\eta)(\lambda_W)\delta_{B^+}^{-\frac{1}{2}}(\lambda) \times \mathcal{L}_{\chi,\eta}\circ\overline{T}_{w^{-1},w_V\chi,w_W\eta}\left(R(\mu_+^*w_0)\Psi_{1,w_V\chi}\otimes\Psi_{1,w_W\eta}^J\right).$$

It remains to use Theorem 5.2.0.1 and Lemma 5.4.2.1 and to do elementary computations. \Box

5.5.2. We now state the formula for $\mathcal{W}_{\chi,\eta}^{I}(1) = (\Phi_{\chi,\eta}^{I}(1), \phi^{\circ})_{\nu}$. We postpone the proof until §7.2. **Proposition 5.5.2.1.** We have

$$\mathcal{W}_{\chi,\eta}^{I}(1) = \Delta_{T_{W}}^{\prime,-1} \Gamma(\chi,\eta) \Delta_{T_{W}} \Delta_{\mathrm{U}(W)}^{-1}$$

5.5.3. We end the proof of Theorem 1.0.2.3.

Lemma 5.5.3.1. The rational map

$$(\chi,\eta) \mapsto \Phi^{\circ}_{\chi,\eta} := (\Delta_{T_W}^{\prime,-1} \Gamma(\chi,\eta) \Delta_{T_W} \Delta_{\mathrm{U}(W)}^{-1})^{-1} \Phi^I_{\chi,\eta}$$
(5.5.3.1)

extends to a regular function. In particular, for every (χ, η) there exists a unique $\mathcal{W}^{\circ}_{\chi,\eta} \in WS_{\chi,\eta}$ such that $\mathcal{W}^{\circ}_{\chi,\eta}(1) = 1$.

Proof. Let (.,.) be the natural pairing between $\operatorname{Ind}_{H}^{G}\nu_{\mu,\psi}$ and $\operatorname{Ind}_{H}^{G}\overline{\nu}_{\mu,\psi}$. We have to show that for every $\Phi \in (\operatorname{ind}_{H}^{G}\overline{\nu}_{\mu,\psi})^{K}$ the map $(\chi,\eta) \mapsto (\Phi_{\chi,\eta}^{\circ}, \Phi)$ is regular. Let Φ_{0} be the generator of $(\operatorname{ind}_{H}^{G}\overline{\nu}_{\mu,\psi})^{K}$ built in Proposition 3.2.2.3. It is enough to show that $(\chi,\eta) \mapsto (\Phi_{\chi,\eta}^{\circ}, \Phi_{0}) = (\Phi_{\chi,\eta}^{\circ}(1), \phi^{\circ})_{\nu}$ is regular. But by Proposition 5.5.2.1 this map is constant equal to 1 so the result follows.

To conclude the proof of Theorem 1.0.2.3, note that by Theorem 1.0.2.1 and Lemma 5.5.3.1 we have $\Phi_{\chi,\eta} \neq 0$ if and only if $(\Phi_{\chi,\eta}(1), \phi^{\circ})_{\nu} \neq 0$, and moreover in that case $\Phi_{\chi,\eta}/(\Phi_{\chi,\eta}(1), \phi^{\circ})_{\nu} = \Phi_{\chi,\eta}^{\circ}$. Theorem 1.0.2.3 therefore follows from (5.5.0.2), Proposition 5.5.1.1, (5.5.3.1) and elementary computations.

5.5.4. Recall that in Proposition 1.0.2.4 we have stated in alternative version of Theorem 1.0.2.3 in the split case. We now prove a key result towards this formula. The proof of Proposition 1.0.2.4 will be ended in §7.2.5 once we define the representation $\mathcal{R}_{\overline{\mu}}$. Recall that ϕ^{\times} and ϕ^{1} were defined in §3.1.3.

Proposition 5.5.4.1. Assume that we are in the split case. For every $\lambda \in \Lambda^-$ and (χ, η) we have

$$(\Phi_{\chi,\eta}^{\circ}(\lambda),\phi^{\times})_{\nu} = \Delta_{\mathcal{U}(W)} \sum_{w \in W_{G}} \mathbf{b}^{\times}(w_{V}\chi,w_{W}\eta)\mathbf{d}_{V}(w_{V}\chi)\mathbf{d}_{W}(w_{W}\eta) \left((w_{0}w_{V}\chi \boxtimes w_{W}\mu\eta) \,\delta_{B^{+}}^{-\frac{1}{2}} \right)(\lambda)$$

where

$$\mathbf{b}^{\times}(\chi,\eta) = \prod_{i+j \le r'+1} L_F^{-1}(\frac{1}{2},\chi_i\eta_j) \prod_{i+j>r'+1} L_F^{-1}(\frac{1}{2},\chi_i^{-1}\eta_j^{-1}).$$
(5.5.4.1)

Proof. Using Lemma 3.1.3.1 we have, as with (5.5.1.2), that for generic (χ, η)

$$(R(w_0 1_{I_V w_0 \lambda_V I_V}) \otimes R(1_{I_W \lambda_W I_W}) \Phi^I_{\chi,\eta}(1), \phi^{\times})_{\nu} = \operatorname{vol}(I_V w_0 \lambda_V I_V) \operatorname{vol}(I_W \lambda_W I_W) (\Phi^I_{\chi,\eta}(\lambda_V, \lambda_W), \phi^{\times})_{\nu}.$$

Because $\phi^1 \in \nu_{\mu,\psi}^{K_W}$ we know that $S_{\eta}(\Psi_{\mu\eta} \otimes \phi^1) = \Psi_{\eta}^{1,J}$. As $\Delta'_{T_W} = \Delta_{T_W}$, $\operatorname{vol}(T_W^0) = 1$ and $\operatorname{vol}(I_J^1) = q^{-m} \operatorname{vol}(I_W)$, we see by (3.1.3.1) and Lemma 5.3.2.2 that

$$\mathcal{L}_{\chi,\eta}(R(w_0)\Psi_{1,\chi}\otimes S_{\eta}(\Psi_{\mu\eta}^{\circ}\otimes\phi^{\times})) = q^m \mathcal{L}_{\chi,\eta}(R(\mu_+^*w_0)\Psi_{1,\chi}\otimes\Psi_{1,\eta}^{1,J}) = \operatorname{vol}(I_V)\operatorname{vol}(I_W).$$

We now conclude that Proposition 5.5.4.1 holds for generic (χ, η) as in the proof of Proposition 5.5.1.1. For general (χ, η) , we use Lemma 5.5.3.1.

6. Unfolding of tempered periods

In this section, always assume that $r \geq 1$ and temporarily leave the unramified setting: E/F is any quadratic extension of local fields of characteristic zero, or $E = F \times F$. All the other objects are as described in §2.1, except that contrary to the situation of §2.1.2 we only assume that W^{\perp} is split. This means that exist two free families (v_1, \ldots, v_r) and (v_1^*, \ldots, v_r^*) in V such that, with $X = \operatorname{Vect}(v_1, \ldots, v_r)$ and $X^* = \operatorname{Vect}(v_1^*, \ldots, v_r^*)$, we have $W^{\perp} = X \oplus X^*$ and for every $1 \leq i, j \leq r$, $\langle v_i, v_j \rangle_V = \langle v_i^*, v_j^* \rangle_V = 0$ and $\langle v_i, v_j^* \rangle_V = \delta_{i,j}$.

6.1. Preliminaries

6.1.1. For f and g two positive functions on a set X, we write $f(x) \ll g(x)$ if there exists C > 0 such that for all $x \in X$ we have $f(x) \leq Cg(x)$.

For \mathbb{G} the *F*-points of a linear algebraic group over *F*, let ς be a (class of) logarithmic height function on \mathbb{G} , as in [BP20, Section 1.2]. Note that if $\mathbb{G}' \leq \mathbb{G}$ we may take $\varsigma_{|\mathbb{G}'}$ as the logarithmic height function on \mathbb{G}' , hence the absence of reference to the group in the notation. Denote by $\Xi^{\mathbb{G}}$ the Harish-Chandra special spherical function on \mathbb{G} and $\mathcal{C}^w(\mathbb{G})$ the weak Harish-Chandra Schwartz space (see [BP20, § 1.5]). It contains $C_c^{\infty}(\mathbb{G})$ as a dense subset. By definition, $C^w(\mathbb{G}) = \bigcup_{d>0} C_d^w(\mathbb{G})$ where for every $f \in C_d^w(\mathbb{G})$ we have

$$|f(g)| \ll \Xi^{\mathbb{G}}(g)\varsigma(g)^d, \ g \in \mathbb{G}.$$

Realize U(V) as a subgroup of $U(V) \times U(V)$ by the diagonal embedding. By [Wal03, Lemma II.1.5.] in the *p*-adic case, [Var77, Proposition 31] in the Archimedean case, for all $\varepsilon > 0$ we have

$$\int_{\mathcal{U}(V)} \Xi^{\mathcal{U}(V) \times \mathcal{U}(V)}(g) e^{-\varepsilon_{\varsigma}(g)} dg < \infty.$$
(6.1.1.1)

6.1.2. In the unramified situation we have already equipped our groups \mathbb{G} with a left-invariant Haar measure dg in §2.1.9. In general, we have to make choices.

We give the vector spaces F^k and E^k the k-fold product of $d_{\psi x}$ and $d_{\psi_E} x$ the ψ - and ψ_E autodual measure respectively, which we simply write dx. This applies to $X^* \cong E^r$ via our basis $(v_i^*)_{i=1}^r$. This also yields measures on the root subgroups N_{α} by the parametrizations n_{α} (see §2.1.4). The unipotent groups N_k and N(X) are then given the product measure on the root subgroups. Note that this coincides with our choices in the unramified setting.

On G_k , U(V) and U(W), if our data are unramified we keep dg, and otherwise we take any Haar measure. On G_k there is another natural choice, namely $d_{\psi}g := \frac{\prod_{1 \le i,j \le r} d_{\psi_E}g_{i,j}}{|\det g|^k}$. Let $v(G_k) > 0$ be the quotient $dg(d_{\psi}g)^{-1}$. In the unramified situation we have $v(G_k) = \Delta_{G_k}$. For k = r, denote by ψ_r the restriction of ψ_U to N_r . Fourier inversion yields for $f \in C_c^{\infty}(G_r)$ and $g \in G_r$

$$f(g) = \upsilon(G_{r-1})^{-1} \int_{N_{r-1} \setminus G_{r-1}} \int_{N_r} f(\gamma^{-1} n_r \gamma g) \overline{\psi}_r(n_r) dn_r d\gamma.$$
(6.1.2.1)

Let $P_r = G_{r-1} \ltimes E^{r-1}$ be the mirabolic subgroup of G_r fixing v_r on the left equipped with dp = dgdx. Let U_r be its unipotent radical. The isomorphism $P_r \setminus G_r \cong X^* \setminus \{0\}$ implies that for $f \in C_c^{\infty}(X^*)$

$$\int_{X^*} f(x^*) dx^* = \frac{v(G_{r-1})}{v(G_r)} \int_{P_r \setminus G_r} |\det \gamma| f(\gamma^* v_r^*) d\gamma, \qquad (6.1.2.2)$$

where we recall that γ^* was defined in §2.1.7 as the transpose or conjugate transpose of γ .

All the other groups we integrate on will be products of the above-mentioned ones and we give them the product measure.

6.2. Fourier–Jacobi periods

6.2.1. Denote by $\overline{\omega}_V$ the Weil representation of U(V) associated to $(\overline{\mu}, \overline{\psi})$ defined in §3.1. By the mixed model described in [GI16, Section 7.4], we may realize it on $\mathcal{S}(X^*) \otimes \overline{\nu}_{\mu,\psi}$, where $\mathcal{S}(X^*)$ is the Schwartz space on X^* and $\hat{\otimes}$ is the completed tensor product. We can identify any $\Phi \in \overline{\omega}_V$ with a $\overline{\nu}_{\mu,\psi}$ -valued function on X^* , or a \mathbb{C} -valued function on $X^* \oplus Y^*_+$. Define

$$\Phi_{Y^*_{\perp}} := \Phi(v^*_r) \in \overline{\nu}_{\mu,\psi}$$

We equip $\overline{\omega}_V$ with the inner product

$$(\Phi,\Phi')_{\overline{\omega}} = \int_{X^*} (\Phi(x^*),\Phi'(x^*))_{\overline{\nu}} dx^*,$$

where $(.,.)_{\overline{\nu}}$ is the inner product on $\overline{\nu}_{\mu,\psi}$ defined in (3.1.1.2). By (3.1.1.4), we have

$$\overline{\omega}_V(\gamma)\Phi = \overline{\mu} |.|^{\frac{1}{2}} (\gamma)\Phi(\gamma^*.), \ \gamma \in G_r.$$
(6.2.1.1)

By [BLX24, Lemma 20.1], there exists $\varepsilon > 0$ such that

$$\left|\left(\overline{\omega}_{V}(g_{V})\Phi_{1},\Phi_{2}\right)_{\overline{\omega}}\right| \ll_{\Phi_{1},\Phi_{2}} e^{-\varepsilon\varsigma(g_{V})}, \ g_{V} \in \mathrm{U}(V), \ \Phi_{1},\Phi_{2} \in \overline{\omega}_{V},$$

$$(6.2.1.2)$$

$$(\overline{\nu}_{\mu,\psi}(g_Wh)\phi_1,\phi_2)_{\overline{\nu}}|\ll_{\phi_1,\phi_2} e^{-\varepsilon\varsigma(g_W)}, \ g_W \in \mathrm{U}(W), \ h \in \mathbb{H}(\mathbb{W}), \ \phi_1,\phi_2 \in \overline{\nu}_{\mu,\psi}.$$
(6.2.1.3)

Moreover, by [BLX24, Lemma 20.1] the linear forms

$$f \in C_c^{\infty}(G) \mapsto \int_H f(h)(\overline{\nu}_{\mu,\psi}(h)\phi_1,\phi_2)_{\overline{\nu}}dh, \text{ and } f \in C_c^{\infty}(G_r) \mapsto \int_{N_r} f(n_r)\psi_r(n_r)dn_r$$

extend by continuity to $C^w(G)$ and $C^w(G_r)$ respectively, where we recall that $G = U(V) \times U(W)$. We denote them by $\mathcal{P}_H(f \otimes \phi_1 \otimes \phi_2)$ and $\mathcal{P}_{N_r}(f)$. We also introduce

$$\mathcal{P}_{\mathrm{U}(V)}(f \otimes \Phi_1 \otimes \Phi_2) := \int_{\mathrm{U}(V)} f(g)(\overline{\omega}_V(g)\Phi_1, \Phi_2)_{\overline{\omega}} dg, \ f \in \mathcal{C}^w(\mathrm{U}(V) \times \mathrm{U}(V)).$$

This integral converges absolutely by (6.1.1.1) and (6.2.1.2).

Recall that $L = U(V) \times (U(W) \times G_r)$ and $H^L = H \times N_r \subset L$ (§2.1.6). Let $\overline{\nu}^L$ be the representation of H^L defined by

$$\overline{\nu}^L := \overline{\nu}_{\mu,\psi} \boxtimes \psi_r$$

Recall that P(X) and N(X) were defined in §2.1.6. It follows from [GI16, Section 7.4] that for $n_r, n'_r \in N_r, n \in N(X), g_W \in U(W)$, and $g_V \in U(V)$

$$\psi_r(n'_r)\overline{\psi}_r(n_r)(\overline{\omega}_V(n'_r ng_W g_V)\Phi)_{Y^*_+} = \overline{\nu}^L(n_r ng_W, n'_r)(\overline{\omega}_V(g_V)\Phi)_{Y^*_+}.$$
(6.2.1.4)

For $\phi_1, \phi_2 \in \nu^L$, set

$$\mathcal{P}_{H^L}(f \otimes \Phi_1 \otimes \Phi_2) := \int_{H^L} f(h)(\overline{\nu}^L(h)\phi_1, \phi_2)_{\overline{\nu}} dh, \ f \in C_c^\infty(L),$$

which we extend by continuity to $C^w(L)$. Note that $\mathcal{P}_{H^L} = \mathcal{P}_H \otimes \mathcal{P}_{N_r}$.

6.2.2. For $(\mathbb{G}, \mathbb{H}) \in \{(\mathbb{U}(V) \times \mathbb{U}(V), \mathbb{U}(V)), (G, H), (L, H^L)\}$, if σ is a tempered representation of \mathbb{G} equipped with an invariant inner product (., .), then for every $\varphi_1, \varphi_2 \in \sigma$ the map $c_{\varphi_1, \varphi_2} : g \mapsto (\sigma(g)\varphi_1, \varphi_2)$ belongs to $\mathcal{C}^w(\mathbb{G})$. We set for $\Phi_i \in \overline{\omega}_V, \overline{\nu}_{\mu,\psi}$ or $\overline{\nu}^L$

$$\mathcal{P}_{\mathbb{H}}(\varphi_1 \otimes \Phi_1, \varphi_2 \otimes \Phi_2) := \mathcal{P}_{\mathbb{H}}(c_{\varphi_1, \varphi_2} \otimes \Phi_1 \otimes \Phi_2).$$
(6.2.2.1)

If $\varphi_1 = \varphi_2$ and $\Phi_1 = \Phi_2$, we will simply write $\mathcal{P}_{\mathbb{H}}(\varphi \otimes \Phi)$ for $\mathcal{P}_{\mathbb{H}}(\varphi_1 \otimes \Phi_1, \varphi_2 \otimes \Phi_2)$.

6.3. Relations between local periods

We embed P(X) in L by inclusion on the U(V) factor and projection on $U(W) \times G_r$.

Lemma 6.3.0.1. For all $f \in \mathcal{C}^w(L)$, $\Phi_1, \Phi_2 \in \nu^L$ we have

$$\int_{P(X)} f(p)(\overline{\omega}_V(p)\Phi_1, \Phi_2)_{\overline{\omega}} \delta_{P(X)}(p)^{\frac{1}{2}} dp$$

$$= v(G_r)^{-1} \int_{(N_r \setminus G_r)^2} \mathcal{P}_{H^L} \left(R(g_1)L(g_2)f \otimes (\overline{\omega}_V(g_1)\Phi_1)_{Y_+^*} \otimes (\overline{\omega}_V(g_2)\Phi_2)_{Y_+^*} \right) \delta_{P(X)}(g_1g_2)^{-\frac{1}{2}} dg_1 dg_2,$$

$$(6.3.0.1)$$

where dp is a left invariant measure on P(X).

Proof. Fix Φ_1 and Φ_2 . It can be proved that both sides are absolutely convergent and define continuous linear functionals on $\mathcal{C}^w(L)$ by repeating the proof of [BPC23, Proposition 8.6.2.1] using (6.2.1.1), (6.2.1.2) and (6.2.1.3). Therefore, we are reduced to proving (6.3.0.1) in the case $f = f_G \otimes f_r \in C_c^{\infty}(G) \otimes C_c^{\infty}(G_r)$. The integral on the left becomes

$$\int_{G_r} f_r(g_1) \delta_{P(X)}(g_1)^{\frac{1}{2}} \int_{\mathcal{U}(W)} \int_{N(X)} f_G(g_1 h n) (\overline{\omega}_V(g_1 h n) \Phi_1, \Phi_2)_{\overline{\omega}} dn dh dg_1.$$
(6.3.0.2)

For fixed $g_1 \in G_r$, write

$$\int_{U(W)\times N(X)} f_G(g_1hn)(\overline{\omega}_V(g_1hn)\Phi_1,\Phi_2)_{\overline{\omega}}dndh \\
= \int_{U(W)\times N(X)} f_G(g_1hn) \int_{X^*} ((\overline{\omega}_V(g_1hn)\Phi_1)(x^*),\Phi_2(x^*))_{\overline{\nu}}dx^*dndh \\
= \frac{\upsilon(G_{r-1})}{\upsilon(G_r)} \int_{U(W)\times N(X)} \int_{P_r\setminus G_r} f_G(g_1hn) \left((\overline{\omega}_V(g_2g_1hn)\Phi_1)_{Y^*_+}, (\overline{\omega}_V(g_2)\Phi_2)_{Y^*_+} \right)_{\overline{\nu}} dg_2dndh, \quad (6.3.0.3)$$

where we have used (6.1.2.2) and (6.2.1.1). By applying the Fourier inversion formula (6.1.2.1) to $g \in G_r \mapsto f_G(g_2^{-1}gg_2g_1hn)$ for g = 1 we see that the inner integral in (6.3.0.3) is $v(G_{r-1})^{-1}$ times

$$\int_{P_r \setminus G_r} \int_{N_{r-1} \setminus G_{r-1}} \int_{N_r} f_G(g_2^{-1} \gamma^{-1} n_r \gamma g_2 g_1 h n) \overline{\psi}_r(n_r) \left((\overline{\omega}_V(g_2 g_1 h n) \Phi_1)_{Y_+^*}, (\overline{\omega}_V(g_2) \Phi_2)_{Y_+^*} \right)_{\overline{\nu}} dn_r d\gamma dg_2.$$

$$(6.3.0.4)$$

Let us prove that

$$\int_{P_r \setminus G_r} \int_{N_{r-1} \setminus G_{r-1}} \left| \int_{N_r} f_G(g_2^{-1} \gamma^{-1} n_r \gamma g_2 g_1 h n) \overline{\psi}_r(n_r) dn_r \left((\overline{\omega}_V(g_2 g_1 h n) \Phi_1)_{Y_+^*}, (\overline{\omega}_V(g_2) \Phi_2)_{Y_+^*} \right)_{\overline{\nu}} \right| d\gamma dg_2$$

$$(6.3.0.5)$$

is finite. By the description of the restriction of $\overline{\omega}_V$ to G_r in (3.1.1.4) and by (6.1.2.2) we know that

$$\int_{P_r \setminus G_r} \left| \left((\overline{\omega}_V(g_2g_1hn)\Phi_1)_{Y_+^*}, (\overline{\omega}_V(g_2)\Phi_2)_{Y_+^*} \right)_{\overline{\nu}} \right| dg_2 < \infty.$$

It is therefore enough to show that

$$g_2 \in G_r \mapsto \int_{N_{r-1} \setminus G_{r-1}} \left| \int_{N_r} f_G(g_2^{-1} \gamma^{-1} n_r \gamma g_2 g_1 h n) \overline{\psi}_r(n_r) dn_r \right| d\gamma$$
(6.3.0.6)

is bounded. By the Iwazawa decomposition every $g_2 \in G_r$ can be written as $g_2 = g_{r-1}u_rtk$ where $g_{r-1} \in G_{r-1}$, $u_r \in U_r$, $t = e_r^{\vee}(s)$ for $s \in E^{\times}$ (i.e. the diagonal matrix with only 1's and s at the last coordinate) and $k \in K_V \cap \operatorname{GL}_r$. Note that GL_{r-1} normalizes U_r , that N_r is unimodular, and that t commutes with GL_{r-1} . For any $n_r \in N_r$ we have $\overline{\psi}_r(u_r n_r u_r^{-1}) = \overline{\psi}_r(n_r)$ and $t^{-1}n_r t = t'n_r t'^{-1}$ where $t' = \prod_{i=1}^{r-1} e_i^{\vee}(s)$. Then t' is in the center of G_{r-1} . Therefore, we see that

$$(6.3.0.6) = \int_{N_{r-1}\backslash G_{r-1}} \left| \int_{N_r} f_G(k^{-1}\gamma^{-1}n_r\gamma kg_1hn) \overline{\psi}_r(n_r) dn_r \right| d\gamma.$$

This proves that (6.3.0.6) is bounded and therefore that (6.3.0.5) is finite.

We may therefore collapse the two outer integrals in (6.3.0.4) to see that it is equal to

$$\int_{P_r \setminus G_r} \int_{N_r \setminus P_r} \int_{N_r} f_G(g_2^{-1} p^{-1} n_r p g_2 g_1 h n) \overline{\psi}_r(n_r) \left((\overline{\omega}_V(p g_2 g_1 h n) \Phi_1)_{Y_+^*}, (\overline{\omega}_V(p g_2) \Phi_2)_{Y_+^*} \right)_{\overline{\nu}} dn_r dp dg_2$$

=
$$\int_{N_r \setminus G_r} \int_{N_r} f_G(g_2^{-1} n_r g_2 g_1 h n) \overline{\psi}_r(n_r) \left((\overline{\omega}_V(g_2 g_1 h n) \Phi_1)_{Y_+^*}, (\overline{\omega}_V(g_2) \Phi_2)_{Y_+^*} \right)_{\overline{\nu}} dn_r dg_2, \qquad (6.3.0.7)$$

where we have again used (6.2.1.1) to make p appear in $\overline{\omega}_V$ and to integrate on $N_r \setminus P_r$ rather than $N_{r-1} \setminus G_{r-1}$.

Note that in (6.3.0.2), the integrals over g_1 , n and h are all over compacts. We can therefore plug (6.3.0.7) back into (6.3.0.2) and do the change of variables $g_1 = g_2^{-1}g_1$ to see that (6.3.0.2) is

$$\begin{split} \upsilon(G_{r})^{-1} \int_{N_{r}\setminus G_{r}} \int_{G_{r}} \int_{\mathrm{U}(W)\times N(X)} \int_{N_{r}} f_{r}(g_{2}^{-1}g_{1})\delta_{P(X)}(g_{1}g_{2}^{-1})^{\frac{1}{2}}f_{G}(g_{2}^{-1}n_{r}g_{1}hn)\overline{\psi}_{r}(n_{r}) \\ & \left((\overline{\omega}_{V}(g_{1}hn)\Phi_{1})_{Y_{+}^{*}},(\overline{\omega}_{V}(g_{2})\Phi_{2})_{Y_{+}^{*}}\right)_{\overline{\nu}} dn_{r}dndhdg_{1}dg_{2}. \\ = \upsilon(G_{r})^{-1} \int_{(N_{r}\setminus G_{r})^{2}} \int_{\mathrm{U}(W)\times N(X)} \int_{N_{r}^{2}} f_{r}(g_{2}^{-1}n'_{r}g_{1})\delta_{P(X)}(g_{1}g_{2}^{-1})^{\frac{1}{2}}f_{G}(g_{2}^{-1}n_{r}n'_{r}g_{1}hn)\overline{\psi}_{r}(n_{r}) \\ & \left((\overline{\omega}_{V}(n'_{r}g_{1}hn)\Phi_{1})_{Y_{+}^{*}},(\overline{\omega}_{V}(g_{2})\Phi_{2})_{Y_{+}^{*}}\right)_{\overline{\nu}} dn_{r}dn'_{r}dndhdg_{1}dg_{2}. \end{split}$$

The changes of variables $n_r = n_r (n'_r)^{-1}$ and $g_1 nh = nhg_1$ (which gives $\delta_{P(X)}(g_1)^{-1}$) and (6.2.1.4) give

$$v(G_r)^{-1} \int_{(N_r \setminus G_r)^2} \int_{\mathcal{U}(W) \times N(X)} \int_{N_r^2} f_1(g_2^{-1}n'_r g_1) \delta_{P(X)}(g_1 g_2)^{-\frac{1}{2}} f_2(g_2^{-1}n_r nhg_1) \left(\overline{\nu}^L(n_r nh, n'_r)((\overline{\omega}_V(g_1)\Phi_1)_{Y_+^*}), (\overline{\omega}_V(g_2)\Phi_2)_{Y_+^*} \right)_{\overline{\nu}} dn_r dn'_r dn dh dg_1 dg_2.$$

This is exactly what we were after by definition of \mathcal{P}_{H^L} .

Proposition 6.3.0.2. Let τ , σ_W and σ_V be irreducible tempered representations of G_r , U(W) and U(V) respectively, equipped with invariant inner products. Let $\Sigma := I_{P(X)}^{U(V)} \tau \boxtimes \sigma_W$ be the parabolic induction equipped with its canonical inner product. Then for every $\varphi_V^1, \varphi_V^2 \in \sigma_V, \varphi_{\Sigma}^1, \varphi_{\Sigma}^2 \in \Sigma$, and $\Phi_1, \Phi_2 \in \overline{\omega}_V$ we have

$$\mathcal{P}_{\mathrm{U}(V)}(\varphi_{V}^{1} \otimes \varphi_{\Sigma}^{1} \otimes \Phi_{1}, \varphi_{V}^{2} \otimes \varphi_{\Sigma}^{2} \otimes \Phi_{2}) = v(G_{r})^{-1} \int_{(H \setminus \mathrm{U}(V))^{2}} \mathcal{P}_{H^{L}}\left((\sigma_{V}(h_{i})\varphi_{V}^{i} \otimes \varphi_{\Sigma}^{i}(h_{i}) \otimes (\overline{\omega}_{V}(h_{i})\Phi_{i})_{Y_{+}^{*}})_{i=1,2} \right) dh_{1}dh_{2}.$$

Proof. By definition of the invariant inner product on Σ we have

$$\mathcal{P}_{\mathrm{U}(V)}(\varphi_{V}^{1} \otimes \varphi_{\Sigma}^{1} \otimes \Phi_{1}, \varphi_{V}^{2} \otimes \varphi_{\Sigma}^{2} \otimes \Phi_{2}) = \int_{\mathrm{U}(V)} \int_{P(X) \setminus \mathrm{U}(V)} (\varphi_{\Sigma}^{1}(h_{2}h_{1}), \varphi_{\Sigma}^{2}(h_{2})) dh_{2}(\sigma_{V}(h_{1})\varphi_{V}^{1}, \varphi_{V}^{2}) (\overline{\omega}_{V}(h_{1})\Phi_{1}, \Phi_{2})_{\overline{\omega}} dh_{1}.$$
(6.3.0.8)

By [CHH88, Theorem 2] and [Wal03, Lemma II.1.6] we know that

$$\int_{P(X)\setminus \mathrm{U}(V)} \left| (\varphi_{\Sigma}^{1}(h_{2}h_{1}), \varphi_{\Sigma}^{2}(h_{2})) \right| dh_{2} \ll \Xi^{\mathrm{U}(V)}(h_{1}).$$

Moreover, we have $|(\sigma_V(h_1)\varphi_V^1,\varphi_V^2)| \ll \Xi^{U(V)}(h_1)$. This implies that (6.3.0.8) is absolutely convergent by (6.2.1.2) and (6.1.1.1). Then

$$\mathcal{P}_{\mathrm{U}(V)}(\varphi_{V}^{1} \otimes \varphi_{\Sigma}^{1} \otimes \Phi_{1}, \varphi_{V}^{2} \otimes \varphi_{\Sigma}^{2} \otimes \Phi_{2})$$

$$= \int_{(P(X)\setminus\mathrm{U}(V))^{2}} \int_{P(X)} (\sigma_{V}(ph_{1})\varphi_{V}^{1}, \sigma_{V}(h_{2})\varphi_{V}^{2})(\varphi_{\Sigma}^{1}(ph_{1}), \varphi_{\Sigma}^{2}(h_{2}))(\overline{\omega}_{V}(ph_{1})\Phi_{1}, \overline{\omega}_{V}(h_{2})\Phi_{2})_{\overline{\omega}}d_{L}pdh_{i}$$

$$= v(G_{r})^{-1} \int_{(P(X)\setminus\mathrm{U}(V))^{2}} \int_{(N_{r}\setminus G_{r})^{2}} \delta_{P(X)}^{-1}(g_{i})\mathcal{P}_{H^{L}}\left(\sigma_{V}(g_{i}h_{i})\varphi_{V} \otimes \varphi_{\Sigma}^{i}(g_{i}h_{i}) \otimes (\overline{\omega}_{V}(g_{i}h_{i})\Phi_{i})_{Y_{+}^{*}}\right) dg_{i}dh_{i}$$

$$= v(G_{r})^{-1} \int_{(H\setminus\mathrm{U}(V))^{2}} \mathcal{P}_{H^{L}}\left(\sigma_{V}(h_{i})\varphi_{V}^{i} \otimes \varphi_{\Sigma}^{i}(h_{i}) \otimes (\overline{\omega}_{V}(h_{i})\Phi_{i})_{Y_{+}^{*}}\right) dh_{i},$$

where in the second inequality we have applied Lemma 6.3.0.1 to the map

$$(g_V, g_W, g_r) \mapsto \left(\sigma_V(g_V h_1)\varphi_V^1, \sigma_V(h_2)\varphi_V^2\right) \left((\tau(g_r) \otimes \sigma_W(g_W))\varphi_{\Sigma}^1(h_1), \varphi_{\Sigma}^2(h_2)\right),$$

which is in $\mathcal{C}^w(L)$.

7. Unramified *L*-function for Fourier–Jacobi models

We now go back to the unramified setting described in §2 and prove Proposition 5.5.2.1 (value of $\mathcal{W}^{I}_{\chi,\eta}(1)$) and Theorem 1.0.3.1 (formula for $\mathcal{P}_{H}(\varphi^{\circ}, \phi^{\circ})$ in the tempered case).

7.1. Satake-parameters for unitary groups

7.1.1. Following [BPC23, Section 8.7], we introduce some notations on dual complex groups. Define W_F to be the Weil group of F, and let $Fr \in W_F$ be an absolute Frobenius. In the inert case,

let W_E be the Weil group of E. Note that Fr is sent to c in the quotient $W_F/W_E \simeq \text{Gal}(E/F)$. For convenience, we write W_E for W_F in the split case, so that W_F/W_E is trivial.

For V_k a k-dimensional nondegenerate split skew-Hermitian space over E/F, we identify the dual group $\widehat{U}(V_k)$ of $U(V_k)$ with $\operatorname{GL}_k(\mathbb{C})$ equipped with its standard pinning. The Langlands dual group of $U(V_k)$ is

$$^{L}\mathrm{U}(V_{k}) = \mathrm{GL}_{k}(\mathbb{C}) \rtimes W_{F},$$

where the Galois action factorises through W_F/W_E , with $c(g) = g^*$ in the inert case where

$$g^{\star} = J_k^{\ t} g^{-1} J_k^{-1}, \ J_k = \begin{pmatrix} & 1 \\ & \ddots & \\ (-1)^{k-1} & & \end{pmatrix}.$$

The dual group \widehat{G}_k of G_k is identified with $\operatorname{GL}_k(\mathbb{C}) \times \operatorname{GL}_k(\mathbb{C})$, and its Langlands dual group is

$${}^{L}G_{k} = (\mathrm{GL}_{k}(\mathbb{C}) \times \mathrm{GL}_{k}(\mathbb{C})) \rtimes W_{F},$$

where the Galois action factorises through W_F/W_E with $c(g_1, g_2) = (g_2, g_1)$ in the inert case. Note that in the split case there is also an action of $\operatorname{Gal}(E/F) = \{1, c\}$ on LG_k : c acts by $c(g_1, g_2) = (g_2, g_1)$ on \widehat{G}_k and by identity on W_F . For $S \in {}^LG_k$, we denote it by S^c .

We write $S \mapsto S^*$ for the automorphism of LG_k which is identity on W_F and sends $(g_1, g_2) \in \widehat{G}_k$ to (g_2^*, g_1^*) . There is a base-change embedding BC : ${}^L\mathrm{U}(V_k) \to {}^LG_k$ which is given by

$$g \in \widehat{\mathcal{U}}(V_k) \mapsto (g, g^\star) \in \widehat{G}_k \tag{7.1.1.1}$$

and which is identity on W_F . When the context is clear, we will identify an element $S \in {}^{L}\mathrm{U}(V_k)$ with its image in ${}^{L}G_k$.

We write $(\hat{T}_{V_K}, \hat{B}_{V_k})$ for the standard Borel pair in $\hat{U}(V_k)$, and (\hat{T}_k, \hat{B}_k) for the one in \hat{G}_k . These are subgroups of $\hat{U}(V_k)$ and \hat{G}_k respectively which are stable by the action of W_F . Set

$${}^{L}T_{V_{k}} = \widehat{T}_{V_{k}} \rtimes W_{F}, \ {}^{L}B_{V_{k}} = \widehat{B}_{V_{k}} \rtimes W_{F}, \ {}^{L}T_{k} = \widehat{T}_{k} \rtimes W_{F}, \ {}^{L}B_{k} = \widehat{B}_{k} \rtimes W_{F}.$$

Recall that $G = U(V) \times U(W)$ and set $\tilde{G} := U(W) \times U(W)$. Their dual groups (resp. Langlands dual groups) are

$$\begin{split} \widehat{G} &= \widehat{\mathrm{U}}(V) \times \widehat{\mathrm{U}}(W), \ \widehat{\widetilde{G}} &= \widehat{\mathrm{U}}(W) \times \widehat{\mathrm{U}}(W), \\ (\text{resp. } {}^{L}G &= {}^{L}\mathrm{U}(V) \times_{W_{F}} {}^{L}\mathrm{U}(W), \ {}^{L}\widetilde{G} &= {}^{L}\mathrm{U}(W) \times_{W_{F}} {}^{L}\mathrm{U}(W)). \end{split}$$

Set $B = B_V \times B_W$, $T = T_V \times T_W$, $\tilde{B} = B_W \times B_W$ and $\tilde{T} = T_W \times T_W$. Write $\hat{B} = \hat{B}_V \times \hat{B}_W$, $\hat{T} = \hat{T}_V \times \hat{T}_W$, $\hat{\tilde{B}} = \hat{B}_W \times \hat{B}_W$ and $\hat{\tilde{T}} = \hat{T}_W \times \hat{T}_W$ and set ${}^L T = \hat{T} \rtimes W_F$, ${}^L \tilde{T} = \hat{\tilde{T}} \rtimes W_F$.

Recall that P(X) is the parabolic subgroup of U(V) stabilizing X, with Levi decomposition P(X) = M(X)N(X) where M(X) is the Levi subgroup stabilizing X^* . Set $M = M(X) \times U(W) \subset G$. The corresponding L-groups are ${}^LM(X)$ and LM which are identified with the corresponding Levi subgroups in ${}^LU(V)$ and LG respectively. Note that ${}^LM(X)$ is isomorphic to ${}^LG_r \times_{W_F} {}^LU(W)$ via the map which is identity on W_F and

$$\begin{pmatrix} g_r^{(1)} & & \\ & g_W & \\ & & g_r^{(2)} \end{pmatrix} \mapsto ((g_r^{(1)}, g_r^{(2)\star}), g_W).$$

on $\widehat{M}(X)$. For $S \in {}^{L}M(X)$, we denote by $(S^{(r)}, S^{(m)}) \in {}^{L}G_r \times_{W_F} {}^{L}U(W)$ its image by this morphism. If $S = (S_V, S_W) \in {}^{L}T$, we have $S_V \in {}^{L}M(X)$ and we set $\widetilde{S} := (S_V^{(m)}, S_W) \in {}^{L}\widetilde{G}$.

For $\mathbb{G} \in \{G, \tilde{G}, U(V_k), G_k, M, M(X)\}$, let $W_{\mathbb{G}}$ be the Weyl group $\operatorname{Norm}_{\widehat{\mathbb{G}}}({}^{L}\mathbb{T})/\widehat{\mathbb{T}}$, where $\widehat{\mathbb{T}} \subset \widehat{\mathbb{G}}$ is the standard maximal torus. Note that we have isomorphisms $W_{U(V)} = W_V$, $W_{U(W)} = W_W$ and $W_G = W_V \times W_W$. These groups act by conjugation on ${}^{L}T_V$, ${}^{L}T_W$, and ${}^{L}T$ respectively, and we denote these actions by wS.

For a complex Lie groups ${}^{L}\mathbb{G}$ and a subgroup ${}^{L}\mathbb{Q}$ with respective identity components $\widehat{\mathbb{G}}$ and $\widehat{\mathbb{Q}}$, set

$$D_{\widehat{\mathbb{G}}/\widehat{\mathbb{Q}}}(S) = \det\left(1 - \operatorname{Ad}(S) \mid \operatorname{Lie}(\widehat{\mathbb{G}}) / \operatorname{Lie}(\widehat{\mathbb{Q}})\right), \ S \in {}^{L}\mathbb{Q}.$$

For any k, the choice of the Borel pair (T_k, B_k) in G_k allows us to identify Λ_k the group of cocharacters of T_k with the group of characters of LT_k trivial on W_F . We will denote by $\lambda_k \mapsto \chi_{\lambda_k}$ this identification. For $\lambda_k \in \Lambda_k^+$ (the cone of positive cocharacters with respect to B_k), we write ch_{λ_k} for the character of the irreducible representation of LG_k with highest weight χ_{λ_k} . If $S_k \in ^LT_k$, set $\overline{\mu}S_k = ((I_k, -I_k), \mathrm{id}).S_k$ in the inert case, and $\overline{\mu}S_k = ((\overline{\mu}(\varpi)I_k, \mu(\varpi)I_k), \mathrm{id}).S_k$ in the split case. Therefore, $\chi_{\lambda_k}(\overline{\mu}S_k) = \overline{\mu}(\lambda_k)\chi_{\lambda_k}(S_k)$.

7.1.2. For $k, l \in \mathbb{N}$, we define the representation

$${}^{L}(G_{k} \times G_{l}) = {}^{L}G_{k} \times_{W_{F}} {}^{L}G_{l} \to \operatorname{GL}(\mathbb{C}^{k} \otimes \mathbb{C}^{l} \oplus \mathbb{C}^{k} \otimes \mathbb{C}^{l})$$
$$(S_{k}, S_{l}) \mapsto S_{k} \overset{\mathrm{I}}{\otimes} S_{l}$$

which sends $((g_k^{(1)}, g_k^{(2)}), (g_l^{(1)}, g_l^{(2)})) \in \widehat{G}_k \times \widehat{G}_l$ to $g_k^{(1)} \otimes g_l^{(1)} \oplus g_k^{(2)} \otimes g_l^{(2)}$, factorises through W_F/W_E and sends c to the operator that swaps the two copies of $\mathbb{C}^k \otimes \mathbb{C}^l$ in the inert case.

By [LM14, Appendix A], for any quasisplit reductive group \mathbb{G} over F, we have an isomorphism $H^1(W_F, Z(\widehat{\mathbb{G}})) \simeq \operatorname{Hom}(\mathbb{G}, \mathbb{C}^{\times})$. Let $\overline{\mu}_k$ be the character $(g_l, g_k) \mapsto \overline{\mu}(\det g_k)$ of $G_l \times G_k$. It corresponds to a class $[z_{\overline{\mu}}] \in H^1(W_F, Z(\widehat{G_l} \times \widehat{G_k}))$, and we choose a representative cocycle $z_{\overline{\mu}}$. Set

$$a_{\overline{\mu}} : (\hat{g}, w) \in {}^{L}(G_{l} \times G_{k}) \mapsto (z_{\overline{\mu}}(w)\hat{g}, w) \in {}^{L}(G_{l} \times G_{k})$$

and set $\overset{I}{\otimes}_{\overline{\mu}} := \overset{I}{\otimes} \circ a_{\overline{\mu}}$. Note that if $S_l \in \widehat{G}_l$. Fr and $S_k \in \widehat{G}_k$. Fr, we have

$$S_l \stackrel{\mathrm{I}}{\otimes}_{\overline{\mu}} S_k = S_l \stackrel{\mathrm{I}}{\otimes} \overline{\mu} S_k. \tag{7.1.2.1}$$

Composing $\overset{\circ}{\otimes}_{\overline{\mu}}$ with the embedding BC of (7.1.1.1), we obtain representations

$$^{L}(\mathrm{U}(V)\times G_{k})\to \mathrm{GL}(\mathbb{C}^{n}\otimes\mathbb{C}^{k}\oplus\mathbb{C}^{n}\otimes\mathbb{C}^{k}), \quad ^{L}(\mathrm{U}(W)\times G_{k})\to\mathrm{GL}(\mathbb{C}^{m}\otimes\mathbb{C}^{k}\oplus\mathbb{C}^{m}\otimes\mathbb{C}^{k})$$

which we denote again by $\overset{1}{\otimes}_{\overline{\mu}}$ for simplicity, and

 $\mathcal{R}_{\overline{\mu}}: {}^{L}G \to \mathrm{GL}(\mathbb{C}^{n} \otimes \mathbb{C}^{m} \oplus \mathbb{C}^{n} \otimes \mathbb{C}^{m}), \quad \widetilde{\mathcal{R}}_{\overline{\mu}}: {}^{L}\widetilde{G} \to \mathrm{GL}(\mathbb{C}^{m} \otimes \mathbb{C}^{m} \oplus \mathbb{C}^{m} \otimes \mathbb{C}^{m}).$

Consider the subspace

$$\mathcal{V}_{-} = \left\langle (e_i \otimes e_j, 0) \middle| \begin{array}{c} \frac{1 \leq i \leq n}{1 \leq j \leq m} \\ i + j > r' + 1 \end{array} \right\rangle \oplus \left\langle (0, e_i \otimes e_j) \middle| \begin{array}{c} \frac{1 \leq i \leq n}{1 \leq j \leq m} \\ i + j > r' + 1 \end{array} \right\rangle,$$
$$\left(\text{resp. } \widetilde{\mathcal{V}}_{-} = \left\langle (e_i \otimes e_j, 0) \middle| \begin{array}{c} \frac{1 \leq i \leq m}{1 \leq j \leq m} \\ i + j > m + 1 \end{array} \right\rangle \oplus \left\langle (0, e_i \otimes e_j) \middle| \begin{array}{c} \frac{1 \leq i \leq m}{1 \leq j \leq m} \\ i + j > m + 1 \end{array} \right\rangle,$$

where $(e_i)_{i=1}^k$ denotes the canonical basis of \mathbb{C}^k for any k. It is stable by LT (resp. by ${}^L\widetilde{T}$) and we set $\mathcal{R}_{-,\overline{\mu}}(S) := \mathcal{R}_{\overline{\mu}}(S)_{|\mathcal{V}_-}$ for $S \in {}^LT$ (resp. $\widetilde{\mathcal{R}}_{-,\overline{\mu}}(\widetilde{S}) := \mathcal{R}_{\overline{\mu}}(\widetilde{S})_{|\widetilde{\mathcal{V}}_-}$ for $\widetilde{S} \in {}^L\widetilde{T}$).

7.1.3. We can equip $\mathbb{C}^n \otimes \mathbb{C}^m \oplus \mathbb{C}^n \otimes \mathbb{C}^m$ with the symplectic pairing

$$\langle (u_n \otimes u_m, v_n \otimes v_m), (u'_n \otimes u'_m, v'_n \otimes v'_m) \rangle := ({}^t u_n J_n^{-1} v'_n) ({}^t u_m J_m^{-1} v'_m) - ({}^t u'_n J_n^{-1} v_n) ({}^t u'_m J_m^{-1} v_m).$$

The representation $\mathcal{R}_{\overline{\mu}}$ is symplectic. The subspace \mathcal{V}_{-} is not a Lagrangian, but the subspace

$$\mathcal{Y}_{-} := \mathcal{V}_{-} \oplus \langle (e_{j+r} \otimes e_{m-j+1}, 0) \mid 1 \le j \le m \rangle$$

is. In the split case it is stable by the action of ${}^{L}T$. The Lagrangian subspace \mathcal{Y} appearing in Proposition 1.0.2.4 is

$$\mathcal{Y} = \mathcal{R}_{\overline{\mu}}(w_{0,V})\mathcal{Y}_{-}.$$
(7.1.3.1)

It is also stable by ${}^{L}T$, and we denote by $\mathcal{Y}_{\overline{\mu}}$ this representation of ${}^{L}T$.

7.2. Proof of Proposition 5.5.2.1

7.2.1. Let χ and η be generic unramified characters of T_V and T_W respectively as in §5. They correspond to semisimple conjugacy classes in ${}^L U(V)$ and ${}^L U(W)$ which we identify with representatives $S_V \in \widehat{T}_V$. Fr and $S_W \in \widehat{T}_W$. Fr. Set $S = (S_V, S_W) \in {}^L T$. Recall that we have defined the regular functions \mathbf{d}_V , \mathbf{d}_W and \mathbf{b} in (5.5.0.1) and (1.0.2.5). Then we have

$$\mathbf{d}_{V}(\chi)\mathbf{d}_{W}(\eta) = D_{\widehat{G}/\widehat{B}}(w_{0,G}S)^{-1}, \text{ and } \mathbf{b}(\chi,\overline{\mu}\eta) = \det(1 - q_{F}^{-\frac{1}{2}}\mathcal{R}_{-,\overline{\mu}}(w_{0,G}S)),$$
(7.2.1.1)

where $w_{0,G}$ is the longest element in W_G .

7.2.2. Proof of Proposition 5.5.2.1 for r = 0. We now write the proof of Proposition 5.5.2.1 assuming first that r = 0. We have to show that

$$\mathcal{W}_{\chi,\overline{\mu}\eta}^{I}(1) = \Delta_{T_{W}}^{\prime,-1} \Gamma(\chi,\overline{\mu}\eta) \Delta_{T_{W}} \Delta_{\mathrm{U}(W)}^{-1}$$

By Proposition 5.5.1.1 and (7.2.1.1), this amounts to proving that for generic $S \in \hat{T}$. Fr we have

$$\sum_{w \in W_G} \frac{\det(1 - q_F^{-\frac{1}{2}} \mathcal{R}_{-,\overline{\mu}}(wS))}{D_{\widehat{G}/\widehat{B}}(wS)} = \frac{\Delta_{T_W}}{\Delta_{\mathrm{U}(W)}}.$$
(7.2.2.1)

We claim that the LHS of (7.2.2.1) is constant in S. Indeed, note that the composition $S \mapsto S$.Fr \mapsto det $(1 - q_F^{-\frac{1}{2}}\mathcal{R}_{-,\overline{\mu}}(S$.Fr)) is a linear combination of W_F -invariant characters of \widehat{T} . Let $\lambda \in X^*(\widehat{T})^{W_F}$ be such a character. By the natural isomorphism $X^*(\widehat{T}) \cong \mathbb{Z}^n \times \mathbb{Z}^n$, we can write the coordinates $\lambda = (\lambda_{V,1}, \ldots, \lambda_{V,n}, \lambda_{W,1}, \ldots, \lambda_{W,n})$. For each $1 \leq i \leq n$ we have $i - n \leq \lambda_{V,i}, \lambda_{W,i} \leq i - 1$. Let $\rho \in \frac{1}{2}X^*(\widehat{T})$ be the half-sum of the positive roots in \widehat{T} with respect to \widehat{B} . The coordinates of $\lambda + \rho$ are all integers or all half-integers between $\frac{1-n}{2}$ and $\frac{n-1}{2}$, so that we have the following alternative.

• Either λ is singular, i.e. there exists a coroot α^{\vee} such that $\langle \lambda + \rho, \alpha^{\vee} \rangle = 0$, in which case $\sum_{w \in W_G} \frac{\lambda(wS)}{D_{\widehat{G}/\widehat{B}}(wS)} = 0$ by the Weyl character formula [BPC23, Proposition A.0.2.1].

• Either $\lambda + \rho$ is conjugated to ρ under W_G and $S \mapsto \sum_{w \in W_G} \frac{\lambda(wS)}{D_{\widehat{G}/\widehat{B}}(wS)}$ is constant.

This shows that (7.2.2.1) is constant in S, and the claim follows from elementary computations taking S corresponding to the characters $(\chi, \eta) = w_{0,G}(\delta_{B_V}^{-\frac{1}{2}}, \mu \delta_{B_V}^{-1} \delta_{B_J}^{\frac{1}{2}})$ in the inert case and $w_{0,G}(\delta_{B_V}^{-\frac{1}{2}}, \mu \delta_{B_V} \delta_{B_J}^{-\frac{1}{2}})$ in the split case, noting that with this choice $\det(1 - q_F^{-\frac{1}{2}}\mathcal{R}_{-,\overline{\mu}}(wS))$ is zero unless w = 1 in the split or inert odd case, and w = 1 or w is the element corresponding to $(1, (1, n)) \in W_G \subset \mathfrak{S}_n \times \mathfrak{S}_n$ in the inert even case.

7.2.3. Proof of Proposition 5.5.2.1 for $r \ge 1$. We now prove Proposition 5.5.2.1 for $r \ge 1$. We first state two lemmas that will also be used in the proof of Proposition 7.3.1.1. Denote by $\Lambda_r^{++} \subset \Lambda_r^+$ the subcone of cocharacters that are dominant with respect to B_{r+1} through the embedding $g \in G_r \mapsto \begin{pmatrix} g \\ 1 \end{pmatrix} \in G_{r+1}$. Note that $\Lambda_r^{++} \subset \Lambda_V^+$.

Lemma 7.2.3.1. For $\lambda_r \in \Lambda_r^{++}$ we have

$$\mathcal{W}_{\chi,\overline{\mu}\eta}^{I}(\lambda_{r}) = \Delta_{T_{W}}^{\prime,-1} \Gamma(\chi,\overline{\mu}\eta) \frac{\Delta_{T_{W}}}{\Delta_{\mathrm{U}(W)}} \sum_{w \in W_{V}} \frac{\det(1 - q_{F}^{-\frac{1}{2}} (wS_{V})^{(r)\star} \overset{\mathrm{I}}{\otimes_{\overline{\mu}}} S_{W})}{D_{\widehat{\mathrm{U}}(V)/\widehat{B}_{V}} (wS_{V})} ch_{\lambda_{r}} \left((wS_{V})^{(r)} \right) \delta_{B_{V}}(\lambda_{r})^{\frac{1}{2}}.$$
(7.2.3.1)

Proof. For any S_W the map $S_V \in \widehat{T}_V$. Fr $\mapsto \det(1 - q_F^{-\frac{1}{2}}(S_V)^{(r)\star} \bigotimes_{\overline{\mu}}^{\mathbf{I}} S_W)ch_{\lambda_r}((S_V)^{(r)})$ is invariant under the action of $W_{M(X)}$. By the Weyl character formula [BPC23, Proposition A.0.2.1], we have

$$\sum_{w \in W_{M(X)}} D_{\widehat{U}(V)/\widehat{B}_{V}}(wS_{V})^{-1} = D_{\widehat{U}(V)/\widehat{P}(X)}(S_{V})^{-1}.$$

It follows that the RHS of (7.2.3.1) is

$$\Delta_{T_W}^{',-1}\Gamma(\chi,\overline{\mu}\eta)\frac{\Delta_{T_W}}{\Delta_{\mathrm{U}(W)}}\sum_{w\in W_{M(X)}\setminus W_V}\frac{\det(1-q_F^{-\frac{1}{2}}(wS_V)^{(r)\star}\overset{\mathrm{I}}{\otimes_{\overline{\mu}}}S_W)}{D_{\widehat{\mathrm{U}}(V)/\widehat{P}(X)}(wS_V)}ch_{\lambda_r}\left((wS_V)^{(r)}\right)\delta_{B_V}(\lambda_r)^{\frac{1}{2}}.$$

By Remark 3.3.0.3 and Proposition 5.5.1.1 specialized to $\lambda_V = \lambda_r \in \Lambda_V^+$ and $\lambda_W = 1$, we know that

$$\mathcal{W}_{\chi,\overline{\mu}\eta}^{I}(\lambda_{r}) = \Delta_{T_{W}}^{\prime,-1}\Gamma(\chi,\overline{\mu}\eta) \sum_{w \in W_{G}} \frac{\det(1-q_{F}^{-\frac{1}{2}}\mathcal{R}_{-,\overline{\mu}}(wS))}{D_{\widehat{G}/\widehat{B}}(wS)} \chi_{\lambda_{r}}\left((wS_{V})^{(r)}\right) \delta_{B_{V}}(\lambda_{r})^{\frac{1}{2}}.$$

The natural projection $W_G \to W_V$ induces a bijection $W_M \setminus W_G = W_{M(X)} \setminus W_V$, where we recall that $M = M(X) \times U(W)$. Therefore we have to show that

$$\sum_{w \in W_M} \frac{\det(1 - q_F^{-\frac{1}{2}} \mathcal{R}_{-,\overline{\mu}}(wS))}{D_{\widehat{G}/\widehat{B}}(wS)} \chi_{\lambda_r} \left((wS_V)^{(r)} \right) = \frac{\Delta_{T_W}}{\Delta_{\mathrm{U}(W)}} \frac{\det(1 - q_F^{-\frac{1}{2}} S_V^{(r)\star} \stackrel{\mathrm{I}}{\otimes_{\overline{\mu}}} S_W)}{D_{\widehat{\mathrm{U}}(V)/\widehat{P}(X)}(S_V)} ch_{\lambda_r}(S_V^{(r)}).$$
(7.2.3.2)

We have the equalities $W_M = W_{\widetilde{G}} \times W_{G_r}$ and

$$D_{\widehat{G}/\widehat{B}}(S) = D_{\widehat{U}(V)/\widehat{P}(X)}(S_V) D_{\widehat{G}/\widehat{B}}(\widetilde{S}) D_{\widehat{G}_r/\widehat{B}_r}(S_V^{(r)}),$$
$$\mathcal{R}_{-,\overline{\mu}}(S) = S_V^{(r)\star} \overset{\mathrm{I}}{\otimes_{\overline{\mu}}} S_W \oplus \widetilde{\mathcal{R}}_{-,\overline{\mu}}(\widetilde{S}).$$

It follows that the LHS of (7.2.3.2) is

$$\frac{\det(1-q_F^{-\frac{1}{2}}S_V^{(r)\star}\overset{\mathrm{I}}{\otimes_{\overline{\mu}}}S_W)}{D_{\widehat{U}(V)/\widehat{P}(X)}(S_V)}\sum_{\widetilde{w}\in W_{\widetilde{G}}}\frac{\det(1-q_F^{-\frac{1}{2}}\widetilde{\mathcal{R}}_{-,\overline{\mu}}(\widetilde{w}\widetilde{S}))}{D_{\widehat{\widetilde{G}}/\widehat{\widetilde{B}}}(\widetilde{w}\widetilde{S})}\sum_{w_r\in W_{G_r}}\frac{\chi_{\lambda_r}(w_rS_V^{(r)})}{D_{\widehat{G}_r/\widehat{B}_r}(w_rS_V^{(r)})}$$

By the r = 0 case and the Weyl character formula [BPC23, Proposition A.0.2.1], we have

$$\sum_{\widetilde{w}\in W_{\widetilde{G}}}\frac{\det(1-q_F^{-\frac{1}{2}}\widetilde{\mathcal{R}}_{-,\overline{\mu}}(\widetilde{w}\widetilde{S}))}{D_{\widehat{\widetilde{G}}/\widehat{\widetilde{B}}}(\widetilde{w}\widetilde{S})} = \frac{\Delta_{T_W}}{\Delta_{\mathrm{U}(W)}}, \quad \text{and} \quad \sum_{w_r\in W_{G_r}}\frac{\chi_{\lambda_r}(w_rS_V^{(r)})}{D_{\widehat{G}_r/\widehat{B}_r}(w_rS_V^{(r)})} = ch_{\lambda_r}(S_V^{(r)}).$$

This concludes.

For $k \in \mathbb{N}$, denote by $\operatorname{As}^{(-1)^k}$ the Asai representation

$$\operatorname{As}^{(-1)^k} : {}^LG_r \to \operatorname{GL}(\mathbb{C}^r \otimes \mathbb{C}^r)$$

given by $\operatorname{As}^{(-1)^k}(g) = g^{(1)} \otimes g^{(2)}$ for $g = (g^{(1)}, g^{(2)}) \in \widehat{G}_r$, and which factorises through W_F/W_E with $\operatorname{As}^{(-1)^k}(c) = (-1)^k s$ in the inert case, where $s \in \operatorname{GL}(\mathbb{C}^r \otimes \mathbb{C}^r)$ is characterized by $s(u \otimes v) = v \otimes u$.

Lemma 7.2.3.2. Let $S_V \in \hat{T}_V$.Fr, $S_W \in \hat{T}_W$.Fr and $S_r \in \hat{T}_r$.Fr be generic. We have

$$\det(1 - q_F^{-1}S_W \overset{I}{\otimes} S_r^c) \det(1 - q_F^{-1}As^{(-1)^m}(S_r)) = \sum_{w \in W_V}$$

$$\frac{\det(1 - q_F^{-\frac{1}{2}}(wS_V)^{(r)\star} \overset{I}{\otimes}_{\overline{\mu}} S_W)}{D_{\widehat{U}(V)/\widehat{B}_V}(wS_V)} \det(1 - q_F^{-\frac{1}{2}}(wS_V)^{(r)\star} \overset{I}{\otimes}_{\overline{\mu}} S_r) \det(1 - q_F^{-\frac{1}{2}}(wS_V)^{(m)} \overset{I}{\otimes}_{\overline{\mu}} S_r),$$

where S_r^c is the conjugate of S_r by c.

Proof. By (7.1.2.1), this is the same proof as [BPC23, Equation (8.7.2.8)], noting that det(1 – $q_F^{-1}\overline{\mu}S_W \overset{\mathrm{I}}{\otimes} (\overline{\mu}S_r)^c) = \det(1 - q_F^{-1}S_W \overset{\mathrm{I}}{\otimes} S_r^c)$ and $\det(1 - q_F^{-1}\mathrm{As}^{(-1)^{m+1}}(\overline{\mu}S_r)) = \det(1 - q_F^{-1}\mathrm{As}^{(-1)^m}(S_r)).$

We now end the proof of Proposition 5.5.2.1 in the case $r \ge 1$. We have to show that $\mathcal{W}^{I}_{\chi,\overline{\mu}\eta}(1) = \Delta_{T_{W}}^{\prime,-1}\Gamma(\chi,\overline{\mu}\eta)\Delta_{T_{W}}\Delta_{U(W)}^{-1}$, which by Lemma 7.2.3.1 amounts to

$$\sum_{w \in W_V} \frac{\det(1 - q_F^{-\frac{1}{2}} (wS_V)^{(r)\star} \stackrel{1}{\otimes_{\overline{\mu}}} S_W)}{D_{\widehat{U}(V)/\widehat{B}_V} (wS_V)} = 1.$$

This follows from Lemma 7.2.3.2 by taking the limit $S_r \to 0$.

_

7.2.4. As the proof of Theorem 1.0.3.1 is now complete, we rewrite its formula using Satake parameters. In the next section we will use it for $\lambda_W = 1$.

Theorem 7.2.4.1. Let $\varphi_V^{\circ} \in \sigma_V^{K_V}$ and $\varphi_W^{\circ} \in \sigma_W^{K_W}$ and $\mathcal{L}_H \in \operatorname{Hom}_H(\sigma_V \otimes \sigma_W \otimes \overline{\nu}_{\mu,\psi}, \mathbb{C})$. For every $\lambda = (\lambda_V, \lambda_W) \in \Lambda^-$ we have

$$\mathcal{L}_{H}(\sigma_{V}(\lambda_{V})\varphi_{V}^{\circ}\otimes\sigma_{W}(\lambda_{W})\varphi_{W}^{\circ}\otimes\phi^{\circ}) = \frac{\Delta_{\mathrm{U}(W)}}{\Delta_{T_{W}}}\mathcal{L}_{H}(\varphi_{V}^{\circ}\otimes\varphi_{W}^{\circ}\otimes\phi^{\circ}) \\ \times \sum_{w\in W_{G}}\frac{\det(1-q_{F}^{-\frac{1}{2}}\mathcal{R}_{-,\overline{\mu}}(wS))}{D_{\widehat{G}/\widehat{B}}(wS)}\left(((w_{V},w_{0}w_{W})\chi\boxtimes\eta)\,\delta_{B^{+}}^{-\frac{1}{2}}\right)(\lambda).$$

7.2.5. Proof of Proposition 1.0.2.4. We end this section by completing the proof of the alternative formula of Proposition 1.0.2.4 in the split case. It involved the Lagrangian subspace \mathcal{Y} defined in (7.1.3.1). Proposition 1.0.2.4 is now a consequence of Proposition 5.5.4.1, (7.2.1.1) and the equalities $\mathbf{b}^{\times}(\chi, w_0 \overline{\mu} \eta) = \det(1 - q_F^{-\frac{1}{2}} \mathcal{Y}_{\overline{\mu}}(S))$ and $\mathbf{d}_V(\chi) \mathbf{d}_W(w_0 \eta) = D_{\widehat{G}/\widehat{B}^+}(S)^{-1}$, where \mathbf{b}^{\times} was defined in (5.5.4.1).

7.3. Proof of Theorem 1.0.3.1

7.3.1. In this section, let σ_V , σ_W and τ be irreducible unramified representations of U(V), U(W) and G_r respectively. Their Satake parameters are semisimple conjugacy classes in L U(V), L U(W) and LG_r which we identify with representatives $S_V \in \hat{T}_V$.Fr, $S_W \in \hat{T}_W$.Fr and $S_r \in \hat{T}_r$.Fr respectively. Set $S := (S_V, S_W) \in {}^LT$. Let $\varphi_V^{\circ} \in \sigma_V^{K_V}$, $\varphi_W^{\circ} \in \sigma_W^{K_W}$ and $\varphi_\tau^{\circ} \in \tau^{K_r}$ be spherical vectors. Let $\mathcal{L}_H \in \operatorname{Hom}_H(\sigma_V \otimes \sigma_W \otimes \overline{\nu}_{\mu,\psi}, \mathbb{C})$ and $\mathcal{L}_W \in \operatorname{Hom}_{N_r}(\tau, \overline{\psi}_r)$. Set $\Phi^{\circ} = 1_{X^*(\mathcal{O}_E)} \otimes \phi^{\circ} \in \overline{\omega}_V$. Write $\tau^c = \tau \circ c$. Recall that P is the parabolic subgroup of U(V) stabilizing the flag $0 \subset Ev_1 \subset \ldots \subset X$.

Proposition 7.3.1.1. Let $s \in \mathbb{C}$. For $\Re(s)$ large enough we have the absolutely convergent sum

$$\sum_{\lambda_r \in \Lambda_r} \mathcal{L}_W(\tau(\lambda_r)\varphi_{\tau}^{\circ}) \mathcal{L}_H\left(\sigma_V(\lambda_r)\varphi_V^{\circ} \otimes \varphi_W^{\circ} \otimes (\overline{\omega}_V(\lambda_r)\Phi^{\circ})_{Y_+^*}\right) (|.|^s \delta_{P(X)}^{\frac{1}{2}} \delta_P^{-1})(\lambda_r)$$
$$= \frac{L(\frac{1}{2} + s, \sigma_V \times \tau \otimes \overline{\mu})}{L(1 + s, \sigma_W \times \tau^c)L(1 + 2s, \tau, \operatorname{As}^{(-1)^m})} \mathcal{L}_W(\varphi_{\tau}^{\circ}) \mathcal{L}_H(\varphi_V^{\circ} \otimes \varphi_W^{\circ} \otimes \phi^{\circ}).$$
(7.3.1.1)

Moreover, if σ_V , σ_W and τ are tempered, (7.3.1.1) holds if $\Re(s) > -\frac{1}{2}$.

Proof. Let $\lambda_r \in \Lambda_r^+$. By (6.2.1.1), $(\overline{\omega}_V(\lambda_r)\Phi^\circ)_{Y_+^*}$ is $\overline{\mu}(\lambda_r) |\lambda_r|^{\frac{1}{2}} \phi^\circ$ if $\lambda_r \in \Lambda_r^{++}$ and 0 otherwise. By Lemma 7.2.3.1 and the normalization of Proposition 5.5.2.1, we therefore have

$$\mathcal{L}_{H}\left(\sigma_{V}(\lambda_{r})\varphi_{V}^{\circ}\otimes\varphi_{W}^{\circ}\otimes(\overline{\omega}_{V}(\lambda_{r})\Phi^{\circ})_{Y_{+}^{*}}\right) = \mathcal{L}_{H}(\varphi_{V}^{\circ}\otimes\varphi_{W}^{\circ}\otimes\phi^{\circ})\times \\ \begin{cases} \sum_{w\in W_{V}}\frac{\det(1-q_{F}^{-\frac{1}{2}}(wS_{V})^{(r)\star}\overset{\mathrm{I}}{\otimes\overline{\mu}}S_{W})}{D_{\widehat{U}(V)/\widehat{B}_{V}}^{\circ}(wS_{V})}ch_{\lambda_{r}}\left((w\overline{\mu}S_{V})^{(r)}\right)\delta_{B_{V}}(\lambda_{r})^{\frac{1}{2}}|\lambda_{r}|^{\frac{1}{2}} & \text{if } \lambda_{r}\in\Lambda_{r}^{++},\\ 0 & \text{otherwise.} \end{cases}$$

By Shintani and Casselman–Shalika formulas ([Shi76] and [CS80]), we know that for $\lambda_r \in \Lambda_r$

$$\mathcal{L}_{W}(\tau(\lambda_{r})\varphi_{\tau}^{\circ}) = \begin{cases} \mathcal{L}_{W}(\varphi_{\tau}^{\circ})\delta_{B_{r}}(\lambda_{r})^{\frac{1}{2}}ch_{\lambda_{r}}(S_{\tau}) & \text{if } \lambda_{r} \in \Lambda_{r}^{+}, \\ 0 & \text{otherwise.} \end{cases}$$

By Cauchy identity [Bum04, Theorem 43.3], for $\Re(s)$ large enough and $w \in W_V$ we have

$$\sum_{\lambda_r \in \Lambda_r^{++}} |\det \lambda_r|^{\frac{1}{2}+s} ch_{\lambda_r}((w\overline{\mu}S_V)^{(r)}) ch_{\lambda_r}(S_r) = \det(1 - q_F^{-\frac{1}{2}}(wS_V)^{(r)} \overset{\mathrm{I}}{\otimes_{\overline{\mu}}} S_{r,s}),$$

where we have set $S_{r,s} := q_F^{-s} S_r$. Moreover, if σ_V and τ are tempered, this series is absolutely convergent for $\Re(s) > -\frac{1}{2}$ as their Satake parameters have complex norm 1. Therefore, it follows from Lemma 7.2.3.1 and the equality $(\delta_{B_V}^{\frac{1}{2}} \delta_{B_r}^{\frac{1}{2}} \delta_{P(X)}^{\frac{1}{2}} \delta_P^{-1})(\lambda_r) = 1$ that the LHS of (7.3.1.1) is

$$\mathcal{L}_{W}(\varphi_{\tau}^{\circ})\mathcal{L}_{H}(\varphi_{V}^{\circ}\otimes\varphi_{W}^{\circ}\otimes\phi^{\circ})\sum_{w\in W_{V}}\frac{\det(1-q_{F}^{-\frac{1}{2}}(wS_{V})^{(r)\star}\overset{\mathrm{I}}{\otimes_{\overline{\mu}}}S_{W})}{D_{\widehat{\mathrm{U}}(V)/\widehat{B}_{V}}(wS_{V})}\det(1-q_{F}^{-\frac{1}{2}}(wS_{V})^{(r)}\overset{\mathrm{I}}{\otimes_{\overline{\mu}}}S_{r,s})^{-1}.$$

Proposition 7.3.1.1 now follows from Lemma 7.2.3.2, and the equality of L-functions for any $w \in W_V$

$$\begin{split} L(\frac{1}{2} + s, \sigma_V \times \tau \otimes \overline{\mu})^{-1} &= \det(1 - q_F^{-\frac{1}{2}} S_V \overset{\mathrm{I}}{\otimes}_{\overline{\mu}} S_{r,s}) \\ &= \det(1 - q_F^{-\frac{1}{2}} (wS_V)^{(r)} \overset{\mathrm{I}}{\otimes}_{\overline{\mu}} S_{r,s}) \det(1 - q_F^{-\frac{1}{2}} (wS_V)^{(m)} \overset{\mathrm{I}}{\otimes}_{\overline{\mu}} S_{r,s}) \det(1 - q_F^{-\frac{1}{2}} (wS_V)^{(r)\star} \overset{\mathrm{I}}{\otimes}_{\overline{\mu}} S_{r,s}). \end{split}$$

7.3.2. We now prove Theorem 1.0.3.1. Assume that σ_V , σ_W and τ are tempered. Set $\Sigma := I_{P(X)}^{\mathrm{U}(V)} \tau \boxtimes \sigma_W$ a tempered representation of $\mathrm{U}(V)$. Without loss of generality we assume that $(\varphi_V^\circ, \varphi_V^\circ) = (\varphi_W^\circ, \varphi_W^\circ) = (\varphi_\tau^\circ, \varphi_\tau^\circ) = 1$ for the chosen inner products. Let $\varphi_{\Sigma}^\circ \in \Sigma$ be the spherical vector with $\varphi_{\Sigma}^\circ(1) = \varphi_{\tau}^\circ \otimes \varphi_W^\circ$. Then $(\varphi_{\Sigma}^\circ, \varphi_{\Sigma}^\circ) = 1$. By Proposition 6.3.0.2 and Proposition 7.3.1.1 and the Iwasawa decomposition $\mathrm{U}(V) = T_r H K_V$ we have

$$\mathcal{P}_{\mathrm{U}(V)}(\varphi_{V}^{\circ}\otimes\varphi_{\Sigma}^{\circ}\otimes\Phi^{\circ}) = \left|\frac{L(\frac{1}{2},\sigma_{V}\times\tau\otimes\overline{\mu})}{L(1,\sigma_{W}\times\tau^{c})L(1,\tau,\mathrm{As}^{(-1)^{m}})}\right|^{2}\frac{\mathcal{P}_{N_{r}}(\varphi_{\tau}^{\circ})\mathcal{P}_{H}(\varphi_{V}^{\circ}\otimes\varphi_{W}^{\circ}\otimes\phi^{\circ})}{\Delta_{G_{r}}}.$$

By [CS80] and [Xue16, Proposition 1.1.1] we know that

$$\mathcal{P}_{N_r}(\varphi_{\tau}^{\circ}) = \frac{\Delta_{G_r}}{L(1,\tau,\mathrm{Ad})} \text{ and } \mathcal{P}_{\mathrm{U}(V)}(\varphi_V^{\circ} \otimes \varphi_{\Sigma}^{\circ} \otimes \Phi^{\circ}) = \Delta_{\mathrm{U}(V)} \frac{L(\frac{1}{2}, \sigma_V \times \Sigma \otimes \overline{\mu})}{L(1,\sigma_V,\mathrm{Ad})L(1,\Sigma,\mathrm{Ad})}.$$

Theorem 1.0.3.1 now follows from the following *L*-function equality

$$\frac{L(\frac{1}{2},\sigma_V\times\Sigma\otimes\overline{\mu})}{L(1,\Sigma,\mathrm{Ad})} = \frac{L(\frac{1}{2},\sigma_V\times\sigma_W\otimes\overline{\mu})}{L(1,\sigma_W,\mathrm{Ad})L(1,\tau,\mathrm{Ad})} \left|\frac{L(\frac{1}{2},\sigma_V\times\tau\otimes\overline{\mu})}{L(1,\sigma_W\times\tau^c)L(1,\tau,\mathrm{As}^{(-1)^m})}\right|^2.$$

References

- [Ban98] W. Banks. "A Corollary to Bernstein's Theorem and Whittaker Functionals on the Metaplectic Group". In: *Mathematical Research Letters* 5.6 (1998), pp. 781–790.
- [BZSV24] D. Ben-Zvi, Y. Sakellaridis, and A. Venkatesh. *Relative Langlands Duality*. 2024. arXiv: 2409.04677.

- [BP20] R. Beuzart-Plessis. "A local trace formula for the Gan-Gross-Prasad conjecture for unitary groups: the Archimedean case". In: Astérisque 418 (2020), pp. viii + 299. DOI: 10.24033/ast.
- [BPC23] R. Beuzart-Plessis and P-H. Chaudouard. The global Gan–Gross–Prasad conjecture for unitary groups. II. From Eisenstein series to Bessel periods. 2023. arXiv: 2302.12331.
- [BLX24] P. Boisseau, W. Lu, and H. Xue. The global Gan–Gross–Prasad conjecture for Fourier– Jacobi periods on unitary groups. 2024. arXiv: 2404.07342.
- [Bum04] D Bump. *Lie Groups*. Vol. 225. Graduate Texts in Mathematics. Springer-Verlag, New York, 2004.
- [Cas80] W. Casselman. "The unramified principal series of *p*-adic groups. I. The spherical function". In: *Compositio Mathematica* 40.3 (1980), pp. 387–406.
- [CS80] W. Casselman and J. Shalika. "The unramified principal series of *p*-adic groups. II. The Whittaker function". In: *Compositio Mathematica* 41.2 (1980), pp. 207–231.
- [CHH88] M. Cowling, U. Haagerup, and R. Howe. "Almost L2 matrix coefficients." In: Journal für die reine und angewandte Mathematik 387 (1988), pp. 97–110.
- [GGP12] W. T. Gan, B. Gross, and D. Prasad. "Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups". In: Astérisque 346 (2012), pp. 1–109.
- [GI16] W. T. Gan and A. Ichino. "The Gross-Prasad conjecture and local theta correspondence". In: Invent. Math. 206.3 (2016), pp. 705–799. DOI: 10.1007/s00222-016-0662-8.
- [GKT23] W. T. Gan, S. Kudla, and S. Takeda. The Local Theta Correspondence. 2023.
- [Gro97] B. Gross. "On the motive of a reductive group". In: *Invent. Math* 130 (1997), pp. 287–313.
- [Gro98] B. Gross. On the Satake isomorphism. Ed. by A. J. Scholl and R. L.Editors Taylor. London Mathematical Society Lecture Note Series. Cambridge University Press, 1998, pp. 223–238. DOI: 10.1017/CB09780511662010.006.
- [Har12] R. Neal Harris. "The Refined Gross-Prasad Conjecture for Unitary Groups". In: International Mathematics Research Notices 2014.2 (2012), pp. 303–389.
- [II10] A. Ichino and T. Ikeda. "On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture". In: Geom. Funct. Anal. 19.5 (2010), pp. 1378– 1425. DOI: 10.1007/s00039-009-0040-4.
- [KMS03] S. I. Kato, A Murase, and T. Sugano. "Whittaker-Shintani functions for orthgonal groups". In: Tohoku Math. J. 55 (2003), pp. 1–64. DOI: 10.2748/tmj/1113247445.
- [KMS] S. I. Kato, A Murase, and T. Sugano. "Whittaker-Shintani functions for GL_n : An explicit formula".
- [Kho08] M. Khoury Jr. "Multiplicity-one results and explicit formulas for quasi-split *p*-adic unitary groups". PhD thesis. The Ohio State University, 2008.
- [LM14] E. Lapid and Z. Mao. "A conjecture on Whittaker–Fourier coefficients of cusp forms".
 In: J. Number Theory (2014). DOI: 10.1016/j.jnt.2013.10.003.

- [MVW87] C. Mœglin, M-F. Vignéras, and J-L. Waldspurger. Correspondances de Howe sur un corps p-adique. Lecture Notes in Mathematics. Springer Berlin, Heidelberg, 1987. DOI: 10.1007/BFb0082712.
- [Ren10] D. Renard. *Représentations des groupes réductifs p-adiques*. Cours spécialisés. SMF, 2010.
- [Sak13] Y. Sakellaridis. "Spherical functions on spherical varieties". In: American Journal of Mathematics 135.5 (2013), pp. 1291–1381.
- [She13a] X. Shen. "The Whittaker–Shintani Functions for Symplectic Groups". In: International Mathematics Research Notices 2014.21 (2013), pp. 5769–5831. DOI: 10.1093/imrn/rnt144.
- [She13b] X. Shen. "Unramified computation of tensor L-functions on symplectic groups". PhD thesis. University of Minnesota, 2013.
- [Shi76] T. Shintani. "On an explicit formula for class-1 "Whittaker functions" on GLn over p-adic fields." In: *Proc. Japan Acad.* 52.5 (1976), pp. 180–182.
- [Var77] V. S. Varadarajan. Harmonic analysis on real reductive groups. Springer Berlin, Heidelberg, 1977. DOI: 10.1007/BFb0097814.
- [Wal03] J.-L. Waldspurger. "La formule de Plancherel pour les groupes p-adiques d'après Harish-Chandra". In: Journal of the Institute of Mathematics of Jussieu 2 (2003), pp. 235– 333.
- [Xue14] H. Xue. "The Gan-Gross-Prasad conjecture for U(n)×U(n)". In: Adv. Math. 262 (2014), pp. 1130–1191. DOI: 10.1016/j.aim.2014.06.010.
- [Xue16] H. Xue. "Fourier–Jacobi periods and the central value of Rankin–Selberg L-functions". In: Israel J. Math. 212.2 (2016), pp. 547–633. DOI: 10.1007/s11856-016-1300-2.
- [Zha18] L. Zhang. "Whittaker-Shintani functions for general linear groups over *p*-adic fields". In: Sci. China Math. 61 (2018), pp. 95–110. DOI: 10.1007/s11425-016-9064-6.

Paul Boisseau Aix Marseille Univ, CNRS, I2M, Marseille, France email: paul.boisseau@univ-amu.fr