

Supplementary material to reproduce the binned parameters for $D \rightarrow K^- \pi^+ \pi^+ \pi^-$

BESIII Collaboration
(Dated: February 27, 2022)

I INTRODUCTION

The binned results of the strong-phase parameters are non-Gaussian and have correlations with each other. In order for analyses that utilise these results to benefit from the full likelihood distribution, which includes these correlations, some additional information is required to that already published in Ref. [1]. This additional information comprises the efficiency-corrected signal yields tagged with CP eigenstates and flavour modes in the $D \rightarrow K^- \pi^+ \pi^+ \pi^-$ phase space bins and the migration matrices for all the tag modes. In combination with the information presented in Tables 2-7 in Ref. [1], the additional information contained here can be used to construct the full likelihood that leads to results quoted in Table 11, and the projections in Fig 10.

II SIGNAL YIELDS

The efficiency corrected signal yields in $D \rightarrow K^- \pi^+ \pi^+ \pi^-$ phase-space regions shown in Table. I are obtained with the same method described in Sec. 4.2 of Ref. [1] and corrected with the MC-determined efficiencies.

TABLE I: Efficiency-corrected signal yields for the CP -tagged $K^- \pi^+ \pi^+ \pi^-$ events. The uncertainties are statistical only.

Mode	Bin	1	2	3	4
$K^- \pi^+$		94±23	21±17	37±16	102±25
$K^- \pi^+ \pi^0$		310±63	190±47	156±91	245±63
$K^- \pi^+ \pi^+ \pi^-$	1	16±11	80±30	11±15	76±29
	2		6±10	26±19	80±29
	3			7±11	35±22
	4				38±23
$K^+ K^-$		1922±94	1654±88	1780±94	1983±96
$\pi^+ \pi^-$		761±57	505±49	603±53	744±60
$\pi^+ \pi^- \pi^0$		6154±237	5286±213	6020±242	6951±261
$K_S^0 \pi^0 \pi^0$		2943±251	2735±236	3028±262	3250±263
$K_L^0 \pi^0$		4740±253	4274±241	4512±428	5129±280
$K_L^0 \omega$		5594±431	4917±390	5340±427	5740±447
$K_S^0 \pi^0$		3353±162	2672±144	2802±151	4145±184
$K_S^0 \eta(\gamma\gamma)$		653±80	487±68	433±68	608±84
$K_S^0 \eta(\pi\pi\pi^0)$		236±65	228±64	178±63	306±82
$K_S^0 \eta'(\gamma\pi\pi)$		638±103	703±97	754±111	1008±128
$K_S^0 \eta'(\pi\pi\eta)$		503±108	441±99	525±112	669±123
$K_S^0 \omega$		2808±242	2966±253	2877±262	3885±289
$K_S^0 \phi$		846±140	905±132	923±147	1037±151
$K_L^0 \pi^0 \pi^0$		6031±748	3942±706	5742±851	8076±1177

III SYSTEMATIC UNCERTAINTIES AND MIGRATION MATRICES

The fit to determine the strong-phase parameters includes all the sources of systematic uncertainties associated with the detector and measurement procedures. Furthermore the external parameters are included via a Gaussian

constraint. The sources and sizes of systematic uncertainties to be considered on the binned yields in Table I are collated here in Table II from the information presented in Ref. [1].

The systematic uncertainty due to K/π tracking and identification is assigned independently for $D \rightarrow K^-\pi^+$, $D \rightarrow K^-\pi^+\pi^0$ and $D \rightarrow K^-\pi^+\pi^+\pi^-$ decay modes, but correlated across the four bins of phase space. The tracking and identification uncertainties are correlated between the different tag modes, and across the four bins of phase space. The uncertainties due to π^0 reconstruction, impact of resonance modelling on efficiency, fit method for signal yields are also correlated across the tag modes. The phase space region populated by the $D \rightarrow K_S^0 K^-\pi^+$ decay is excluded in the measurement of the strong-phase parameters in the binned analysis and therefore doesn't contribute to the systematic uncertainty. The central values and uncertainties of the external parameters used at the time of the analysis can be found in Ref. [1].

TABLE II: Systematic uncertainties contributing to the determination of the binned hadronic parameters.

Systematics	Sizes
Size of CP -tagged $D \rightarrow K^-\pi^+$ samples	See Table [3] in Ref. [1]
K/π tracking and identification	0.7% for each track
π^0 reconstruction	1%, only contributing in $D \rightarrow K^-\pi^+\pi^0$
Impact of resonance modelling on efficiency	3% for $D \rightarrow K^-\pi^+\pi^+\pi^-$ and 3.5% for $D \rightarrow K^-\pi^+\pi^0$
Size of Monte Carlo samples	1% for each tag mode
$D \rightarrow K_S^0 K^-\pi^+$ background	Negligible as excluded by the binning scheme
Fit method for signal yields	1.5% for $D \rightarrow K^-\pi^+\pi^+\pi^-$ and 0.8% for $D \rightarrow K^-\pi^+\pi^0$
External parameters	See Table [6] in Ref. [1]

The migration effects among $D \rightarrow K^-\pi^+\pi^+\pi^-$ phase-space bins for various tag modes are determined with dedicated MC simulation samples. The migration matrices are listed in Tables III-VI, and show that $\gtrsim 90\%$ events are reconstructed correctly in the corresponding bins for flavour and CP tags; 70-90% events are reconstructed correctly in the corresponding bins for $K_S^0\pi^+\pi^-$ tag. The wrongly assigned events lie mostly in the neighbouring bins. These effects are taken into account when calculating the efficiency-corrected signal yields and induce correlations among the four $K^-\pi^+\pi^+\pi^-$ bins, which must be taken into consideration when determining the covariance matrices.

[1] M. Ablikim et al. (BESIII collaboration), JHEP **05**, 164 (2021), arXiv:2103.05988 [hep-ex].

TABLE III: Migration effects (%) in the flavour and CP tag modes. The columns and rows are the truth and reconstructed bins, respectively.

		$K^-\pi^+$				$K^-\pi^+\pi^0$				K^+K^-				$\pi^+\pi^-$			
Bin		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1		96.73	2.47	0.07	0.73	96.17	2.71	0.17	0.95	96.57	2.46	0.07	0.90	96.60	2.39	0.09	0.93
2		2.33	94.28	3.29	0.10	2.36	93.59	3.85	0.21	2.21	93.93	3.76	0.09	2.18	94.21	3.55	0.06
3		0.09	2.48	94.20	3.23	0.14	2.94	93.05	3.87	0.13	2.27	93.91	3.69	0.09	2.19	94.04	3.69
4		0.88	0.05	1.72	97.35	0.90	0.19	1.88	97.02	0.86	0.07	1.68	97.39	0.89	0.07	1.79	97.25
		$\pi^+\pi^-\pi^0$				$K_S^0\pi^0\pi^0$				$K_L^0\pi^0$				$K_L^0\omega$			
Bin		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1		96.19	2.70	0.15	0.96	96.66	2.40	0.06	0.88	96.60	2.49	0.14	0.76	96.23	2.96	0.07	0.74
2		1.97	93.59	4.23	0.21	2.17	93.85	3.71	0.26	1.82	94.54	3.61	0.03	2.95	93.98	3.03	0.04
3		0.25	2.43	92.58	4.74	0.24	2.64	93.56	3.56	0.09	2.03	94.53	3.34	0.36	1.97	93.92	3.75
4		1.02	0.15	2.07	96.76	1.05	0.08	1.64	97.23	0.98	0.01	1.44	97.57	1.00	0.27	2.06	96.67
		$K_S^0\pi^0$				$K_S^0\eta(\gamma\gamma)$				$K_S^0\eta(\pi\pi\pi^0)$				$K_S^0\eta'(\gamma\pi\pi)$			
Bin		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1		95.98	2.96	0.06	1.00	96.61	2.39	0.19	0.80	96.36	2.49	0.13	1.02	95.63	2.82	0.24	1.31
2		2.49	94.03	3.40	0.08	2.20	94.35	3.40	0.05	2.36	93.43	4.14	0.07	1.99	94.40	3.52	0.08
3		0.13	2.53	93.89	3.45	0.21	2.76	94.43	3.00	0.35	2.65	93.30	3.70	0.25	2.77	92.55	4.42
4		1.02	0.09	1.95	96.94	1.05	0.07	1.96	96.92	1.37	0.07	1.53	97.02	1.04	0.15	2.06	96.75
		$K_S^0\eta'(\pi\pi\eta)$				$K_S^0\omega$				$K_S^0\phi$				$K_L^0\pi^0\pi^0$			
Bin		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1		95.53	3.37	0.31	0.79	95.53	3.01	0.33	1.13	95.82	3.04	0.23	0.91	96.67	2.90	0.04	0.40
2		2.14	94.19	3.54	0.13	3.35	92.35	3.92	0.37	2.17	94.09	3.64	0.10	2.10	94.14	3.74	0.01
3		0.33	2.38	94.08	3.21	0.17	2.56	93.32	3.96	0.24	3.04	92.44	4.28	0.05	2.07	93.52	4.35
4		0.69	0.06	2.03	97.21	0.96	0.16	2.10	96.78	0.80	0.12	2.24	96.83	0.90	0.03	1.64	97.43

TABLE IV: Migration effects (%) for the $K^-\pi^+\pi^+\pi^-$ events tagged with $K^-\pi^+\pi^+\pi^-$. The columns and rows are the truth and reconstructed bins, respectively. Values below 0.005 are recorded as -.

Bin	11	22	33	44	12	13	14	23	24	34
11	90.24	0.48	0.08	0.58	6.13	0.51	1.44	0.26	0.27	0.01
22	0.19	90.88	0.14	0.27	3.62	-	0.03	4.39	0.48	-
33	-	0.08	84.73	0.15	0.04	1.00	0.27	6.78	-	6.94
44	0.09	-	0.03	93.23	0.12	-	2.37	0.03	0.32	3.81
12	3.14	1.93	0.07	0.12	90.12	3.01	0.31	0.17	0.94	0.18
13	0.25	0.44	0.15	0.05	2.44	88.75	4.11	2.30	0.19	1.32
14	1.19	-	0.12	0.89	0.30	2.69	92.09	0.11	2.51	0.10
23	0.17	2.23	2.84	0.31	0.26	2.06	0.18	88.29	2.95	0.69
24	0.19	0.35	0.18	0.40	1.16	0.19	2.62	2.38	88.18	4.34
34	0.06	-	1.69	3.88	0.08	0.89	0.25	0.19	2.15	90.81

