

# Direct photon spectra in Pb-Pb at $\sqrt{s_{NN}} = 5.5$ TeV: hydrodynamics+pQCD predictions

François Arleo<sup>1,2</sup>, David d'Enterria<sup>2</sup>, Dmitri Peressounko<sup>3</sup>

1. LAPTH, UMR CNRS/Univ. Savoie, B.P. 110, 74941 Annecy-le-Vieux Cedex

2. CERN/PH, CH-1211 Geneva 23

3. RRC "Kurchatov Institute", Kurchatov Sq. 1, Moscow 123182

**Abstract.** The  $p_T$ -differential spectra for direct photons produced in Pb-Pb collisions at the LHC, including thermal (hydrodynamics) and prompt (pQCD) emissions are presented.


We present predictions for the transverse momentum distributions of direct- $\gamma$  (i.e. photons not coming from hadron decays) produced at mid-rapidity in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.5$  TeV based on a combined hydrodynamics+pQCD approach. Thermal photon emission in Pb-Pb at the LHC is computed with a hydrodynamical model successfully used in nucleus-nucleus collisions at RHIC energies [1]. The initial entropy density of the produced system at LHC is obtained by extrapolating empirically the hadron multiplicities measured at RHIC [2]. Above  $p_T \approx 3$  GeV/ $c$ , additional prompt- $\gamma$  production from parton-parton scatterings is computed perturbatively at next-to-leading-order (NLO) accuracy [3]. We use recent parton distribution functions (PDF) [4] and parton-to-photon fragmentation functions (FF) [5], modified resp. to account for initial-state shadowing+isospin effects [6] and final-state parton energy loss [7].

We follow the evolution of the hot and dense system produced in central Pb-Pb at LHC by solving the equations of (ideal) relativistic 2D+1 hydrodynamics [1, 2] starting at a time  $\tau_0 = 1/Q_s \approx 0.1$  fm/ $c$ . The system is assumed to have an initial entropy density of  $s_0 = 1120$  fm<sup>-3</sup>, which corresponds to a maximum temperature at the center of  $T_0 \approx 650$  MeV ( $\langle T_0 \rangle \approx 470$  MeV). We use a quark gluon plasma (QGP) and hadron resonance gas (HRG) equation of state above and below  $T_{\text{crit}} \approx 170$  MeV resp., connected by a standard Maxwell construction assuming a first-order phase transition at  $T_{\text{crit}}$ . Thermal photon emission is computed using the most recent parametrizations of the QGP and HRG  $\gamma$  rates. For the QGP phase we use the AMY complete leading-log emission rates including LPM suppression [8]. For the HRG phase, we employ the improved parametrization from Turbide *et al.* [9].

Our NLO pQCD predictions are obtained with the code of ref. [3] with all scales set to  $\mu = p_T$ . Pb-Pb yields are obtained scaling the NLO cross-sections by the number of incoherent nucleon-nucleon collisions:  $N_{\text{coll}} = 1670, 12.9$  for 0-10% central ( $\langle b \rangle = 3.2$  fm) and 60-90% peripheral ( $\langle b \rangle = 13$  fm). Nuclear (isospin and shadowing) corrections of the CTEQ6.5M PDFs [4] are introduced using the NLO nDSg parametrization [6]. At relatively low  $p_T$ , prompt photon yields have a large contribution from jet fragmentation processes. As a result,

final-state parton energy loss in central Pb-Pb affects also the expected prompt  $\gamma$  yields. We account for medium-effects on the  $\gamma$ -fragmentation component by modifying the BFG parton-to-photon FFs [5] with BDMPS quenching weights. The effects of the energy loss are encoded in a single parameter,  $\omega_c = \langle \hat{q} \rangle L^2 \approx 50$  GeV, extrapolated from RHIC. The combination of initial-state (shadowing) and final-state (energy loss) effects results in a quenching factor for prompt photons of  $R_{PbPb} \approx 0.2$  (0.8) at  $p_T = 10$  (100) GeV/c [7].

Our predictions for the direct photon spectra at  $y=0$  in Pb-Pb at 5.5 TeV are shown in Fig. 1. The thermal contribution dominates over the (quenched) pQCD one up to  $p_T \approx 4$  (1.5) GeV/c in central (peripheral) Pb-Pb. Two differences are worth noting compared to RHIC results [1]: (i) the thermal-prompt crossing point moves up from  $p_T \approx 2.5$  GeV/c to  $p_T \approx 4.5$  GeV/c, and (ii) most of the thermal production in this transition region comes solely from the QGP phase. Both characteristics make of semi-hard direct photons at LHC, a valuable probe of the thermodynamical properties of the system.



**Figure 1.** Direct- $\gamma$  spectra in 0-10% central (left) and 60-90% peripheral (right) Pb-Pb at  $\sqrt{s_{NN}} = 5.5$  TeV, with the thermal (QGP and HRG) and prompt (pQCD) contributions differentiated.

Dd'E. and D.P. acknowledge resp. support from 6th EU FP contract MEIF-CT-2005-025073 and MPN Russian Federation grant NS-1885.2003.2.

## References

- [1] D. d'Enterria and D. Peressounko, *Eur. Phys. J. C* **46**, 451 (2006) [arXiv:nucl-th/0503054] and refs. therein.
- [2] F. Arleo, D. d'Enterria and D. Peressounko, arXiv:0707.2356 [nucl-th].
- [3] P. Aurenche *et al.*, *Eur. Phys. J. C* **9** (1999) 107 [arXiv:hep-ph/9811382].
- [4] J. Pumplin *et al.*, *JHEP* **0207** (2002) 012 [arXiv:hep-ph/0201195].
- [5] L. Bourhis, M. Fontannaz, J. P. Guillet, M. Werlen, *Eur. Phys. J. C* **19**, 89 (2001) [arXiv:hep-ph/0009101].
- [6] D. de Florian and R. Sassot, *Phys. Rev. D* **69**, 074028 (2004) [arXiv:hep-ph/0311227].
- [7] F. Arleo, arXiv:0707.2320 [hep-ph].
- [8] P. Arnold, G. D. Moore and L. G. Yaffe, *JHEP* **0112**, 009 (2001).
- [9] S. Turbide, R. Rapp and C. Gale, *Phys. Rev. C* **69**, 014903 (2004).