

# Cascading DGP

Claudia de Rham<sup>a,b</sup>, Gia Dvali<sup>c,d</sup>, Stefan Hofmann<sup>a,e</sup>, Justin Khoury<sup>a</sup>,

Oriol Pujolas<sup>d</sup>, Michele Redi<sup>f</sup> and Andrew J. Tolley<sup>a</sup>

<sup>a</sup>Perimeter Institute for Theoretical Physics, 31 Caroline St. N., Waterloo, ON, N2L 2Y5, Canada

<sup>b</sup>Dept. of Physics & Astronomy, McMaster University, Hamilton ON, L8S 4M1, Canada

<sup>c</sup>CERN, Theory Division, CH-1211 Geneva 23, Switzerland

<sup>d</sup>Center for Cosmology and Particle Physics, New York University, New York, NY 10003 USA

<sup>e</sup>NORDITA, Roslagstullsbacken 23, 106 91 Stockholm, Sweden and

<sup>f</sup>Institut de Theorie des Phenomenes Physiques, EPFL, CH-1015, Lausanne, Switzerland

We present a higher codimension generalization of the DGP scenario which, unlike previous attempts, is free of ghost instabilities. The 4D propagator is made regular by embedding our visible 3-brane within a 4-brane, each with their own induced gravity terms, in a flat 6D bulk. The model is ghost-free if the tension on the 3-brane is larger than a certain critical value, while the induced metric remains flat. The gravitational force law "cascades" from a 6D behavior at the largest distances followed by a 5D and finally a 4D regime at the shortest scales.

The DGP model [1] provides a simple mechanism to modify gravity at large distances by adding a localized graviton kinetic term on a codimension 1 brane in a flat 5D bulk. The natural generalization to higher codimension, however, is not so straightforward. On one hand these models require some regularization due to the divergent behavior of the Green's functions in higher codimension. More seriously, most constructions seem to be plagued by ghost instabilities [2, 3] (see [4] for related work). The purpose of this letter is to show that both pathologies can be resolved by embedding a succession of higher-codimension DGP branes into each other.

## I. SCALAR

We shall focus on the codimension 2 case. As a warm-up exercise, we consider a real scalar field with action,

$$S = \frac{1}{2} \int d^6x \left[ M_6^4 \partial_\mu \phi \partial^\mu \phi + M_5^3 \phi \delta^4(x) + M_4^2 \phi \delta^3(x) \right] \quad (1)$$

describing a codimension 2 kinetic term embedded into a codimension 1 one in 6D. We will impose throughout the paper  $Z_2 \times Z_2$  orbifold projection identifying  $y \rightarrow -y$  and  $z \rightarrow -z$ . The model possesses the two mass scales,

$$m_5 = \frac{M_5^3}{M_4^2} \quad \text{and} \quad m_6 = \frac{M_6^4}{M_5^3} \quad (2)$$

In absence of the 4D kinetic term the propagator on the codimension 1 brane (4-brane) is the DGP propagator [1],

$$G^0(y, y^0) = \frac{1}{M_5^3} \int \frac{dq}{2} \frac{e^{iq(y-y^0)}}{p^2 + q^2 + 2m_6 \frac{p}{p^2 + q^2}} \quad ; \quad (3)$$

where  $p$  is 4D momentum and  $y$  the coordinate orthogonal to the codimension 2 brane (3-brane). To find the exact 5D propagator we can treat the 4D kinetic term (located at  $y = 0$ ) as a perturbation and then sum the series. One finds,

$$G^{\text{exact}} = G_0(y, y^0) + M_4^2 G^0(y) p^2 G^0(y^0) + \dots$$

$$= G^0(y, y^0) \frac{M_4^2 p^2}{1 + M_4^2 p^2 G^0(0)} G^0(y) G^0(y^0) \quad ; \quad (4)$$

In particular the 4D brane-to-brane propagator is determined in terms of the integral of the higher dimensional Green's function,

$$G_4^{\text{exact}} = \frac{G^0(0)}{M_4^2 G^0(0) p^2 + 1} \quad ; \quad (5)$$

For the case at hand,

$$G^0(0) = \frac{1}{M_5^3} \int \frac{dq}{2} \frac{1}{p^2 + q^2 + 2m_6 \frac{p}{p^2 + q^2}} = \frac{2}{M_5^3} \frac{1}{4m_6^2 p^2} \tanh^{-1} \frac{2m_6 p}{2m_6 + p} \quad ; \quad (6)$$

For  $p > 2m_6$ , the analytic continuation of this expression is understood.

Remarkably, the 5D kinetic term makes the 4D propagator finite, thereby regularizing the logarithmic divergence characteristic of pure codimension 2 branes. In particular, when  $M_5$  goes to zero one has  $G_4^{\text{exact}} \sim M_6^4 \log(p=m_6)$ , reproducing the codimension 2 Green's function with physical cutoff given by  $m_6$ . The corresponding 4D Newtonian potential scales as  $1=r^3$  at the largest distances, showing that the theory becomes six dimensional, and reduces to the usual  $1=r$  on the shortest scales. Its behavior at intermediate distances, however, depends on  $m_{5,6}$ . If  $m_5 > m_6$  there is an intermediate 5D regime; otherwise the potential directly turns 6D at a distance of order  $(m_5 m_6)^{-1/2} \log(m_5=m_6)$ .

## II. GRAVITY

Let us now turn to gravity. In analogy with the scalar we consider the action,

$$S = \frac{M_6^4}{2} \int \mathbb{P} \frac{1}{g_6 R_6} + \frac{M_5^3}{2} \int \mathbb{P} \frac{1}{g_5 R_5} + \frac{M_4^2}{2} \int \mathbb{P} \frac{1}{g_4 R_4}$$

where each term represents the intrinsic curvature. This guarantees that the model is fully 6D general covariant.

To find the propagator it is convenient to follow the same procedure as for the scalar and sum the diagrams with insertion of the lower dimensional kinetic term, i.e. the Einstein's tensor  $E$ . For our purpose we only compute the propagator on the 3-brane. Given the higher dimensional propagator, the brane-to-brane propagator due to the insertion of a codimension 1 term is in compact form,

$$\begin{aligned} G^{\text{exact}} &= G^0 \int_{n=0}^{\infty} (M_4^2 E G^0)^n = G^0 [1 - M_4^2 E G^0]^{-1} \\ &= G^0 H^{-1}; \end{aligned} \quad (6)$$

where  $G^0$  is the 4D part of the higher dimensional Green's function evaluated at zero. The tensor  $H$  satisfies by definition,

$$[1 - M_4^2 E G^0] H = \frac{1}{2} \delta + \dots; \quad (7)$$

To find  $H$  one can write the most general Lorentz covariant structure compatible with the symmetries,

$$\begin{aligned} H &= a(\delta_{\mu\nu} + \dots) + b \\ &+ c(p_\mu p_\nu + p_\nu p_\mu + p_\mu p_\nu + p_\nu p_\mu) \\ &+ d p_\mu p_\nu + e p_\mu p_\nu + f p_\mu p_\nu p_\mu; \end{aligned}$$

Requiring that this satisfies Eq. (7) leads to a system of linear equations whose solution determines the coefficients  $a; b; c; d; e; f$ . Using this information one then reconstructs the exact propagator from Eq. (6).

It is straightforward to apply this technique to the cascading DGP. Starting from 6D, the propagator on the 4-brane is [5],

$$\begin{aligned} G_{MNPQ} &= \frac{1}{M_5^3 p_5^2 + 2m_6 p_5} \\ &+ \frac{1}{2} \tilde{\gamma}_{MP} \tilde{\gamma}_{NQ} + \frac{1}{2} \tilde{\gamma}_{MQ} \tilde{\gamma}_{NP} - \frac{1}{4} \tilde{\gamma}_{MN} \tilde{\gamma}_{PQ}; \\ \tilde{\gamma}_{MN} &= \gamma_{MN} + \frac{p_M p_N}{2m_6 p_5}; \end{aligned} \quad (8)$$

where  $M; N \dots$  are 5D indices and  $p_5^2 = p_M p^M$ .  $G^0$  is obtained by integrating the 5D propagator with respect to the extra-momentum. To compute the propagator on the 3-brane, then, we determine the coefficients  $a; b; c; d; e; f$  through the system of linear equations (7).

One finds,

$$\begin{aligned} a &= \frac{1}{2(I_1 p^2 + 1)} \\ b &= \frac{I_1 p^2}{(I_1 p^2 + 1)(I_1 p^2 - 2)} \\ c &= \frac{I_1}{2(I_1 p^2 + 1)} \\ d &= \frac{I_1}{(I_1 p^2 + 1)(I_1 p^2 - 2)} \\ e &= \frac{1}{3(I_1 p^2 + 1)} - \frac{4I_1 + 3I_2 p^2}{3(I_1 p^2 - 2)} \\ f &= \frac{I_2 + 2I_1^2 + I_1 I_2 p^2}{(I_1 p^2 + 1)(I_1 p^2 - 2)} \end{aligned} \quad (9)$$

where

$$\begin{aligned} I_1 &= \frac{1}{m_5} \int \frac{dq}{2} \frac{1}{p^2 + q^2 + 2m_6} \frac{1}{p^2 + q^2} \\ I_2 &= \frac{1}{2m_6 m_5} \int \frac{dq}{2} \frac{1}{p^2 + q^2} \frac{1}{(p^2 + q^2 + 2m_6)} \frac{1}{p^2 + q^2} \end{aligned}$$

All these coefficients are finite showing that the regularization is also effective for the spin 2 case.

Having determined the coefficients of the tensor  $H$  the full propagator is given by Eq. (6). To linear order the amplitude between two conserved sources on the brane is rather simple,

$$\frac{1}{M_4^2} \frac{I_1}{I_1 p^2 + 1} T_{\mu\nu} T^{\mu\nu} - \frac{I_1 p^2}{2I_1 p^2} \frac{1}{4} T T^0; \quad (10)$$

and only depends on the first integral  $I_1$ .

The coefficient in front of the amplitude is exactly as for the scalar however there is a non-trivial tensor structure. One worrisome feature of this amplitude is that the relative coefficient of  $T_{\mu\nu} T^{\mu\nu}$  and  $T T^0$  interpolates between  $-1=4$  in the IR and  $-1=2$  in the UV. The  $-1=4$  in the IR gives the correct tensor structure of gravity in 6D and is unavoidable because at large distances the physics is dominated by 6D Einstein term. From the 4D point of view this can be understood as the exchange of massive gravitons and an extra-scalar. The  $-1=2$  in the UV on the other hand signals the presence of a ghost. This agrees with previous results [2, 3] which used a different regularization. From the 4D point of view the theory decomposes into massive spin 2 fields and scalars. Since the massive spin 2 gives an amplitude with relative coefficient  $-1=3$  the extra repulsion must be provided by a scalar with wrong sign kinetic term. Separating from Eq. (10) the massive spin two contribution, we identify the scalar propagator as

$$G_{\text{ghost}} = \frac{1}{6M_4^2} \frac{I_1}{I_1 p^2 - 2}; \quad (11)$$

This propagator has a pole with negative residue therefore it contains a localized (tachyonic) ghost mode in addition to a continuum of healthy modes.

### III. GHOST FREE THEORY

To clarify the origin of the ghost it is illuminating to consider the decoupling limit studied in [6]. This will allow us to show how a healthy theory can be obtained by simply introducing tension on the lower dimensional brane while retaining the intrinsic geometry at.

In the 6D case the decoupling limit [6] corresponds to taking  $M_5/M_6 \rightarrow 1$  with  $\alpha' \rightarrow (m_6^2 M_5^3)^{2=7} = (M_6^1 M_5^9)^{1=7}$  finite. In this limit, the physics on the 4-brane admits a local 5D description, where only the non-linearities in the helicity 0 part of the metric are kept, and are suppressed by the scale  $\alpha'$ . The effective 5D Lagrangian is given by

$$L_5 = \frac{M_5^3}{4} h^{MN} (Eh)_{MN} - 3M_5^3 (\partial_\mu)^2 + \frac{9}{32m_6^2} \alpha' + (z) \frac{M_4^2}{4} \mathfrak{h} (E\mathfrak{h}) + \mathfrak{h} T ; \quad (12)$$

where  $(Eh)_{MN} = h_{MN} + \dots$  is the linearized Einstein tensor,  $M, N \dots$  are 5D indices and  $\mu, \nu \dots$  are four dimensional. We have rescaled  $\alpha'$  and  $h$  so that they are dimensionless, and the physical 5D metric is

$$\mathfrak{h}_{MN} = h_{MN} + \alpha' M_{MN} ; \quad (13)$$

The first line of (12) is the 5D version of the 'Lagrangian' introduced in [6] for the DGP model. In addition to this, we have the localized curvature term on the 3-brane, which depends on 4D physical metric  $\mathfrak{h}$ . This introduces a kinetic mixing between  $\alpha'$  and the 5D metric.

We now take a further step and compute the boundary effective action valid on the 3-brane. At the quadratic order by integrating out the fifth dimension the 5D kinetic term of  $\alpha'$  produces a 4D mass term  $\alpha' M_5^3 \frac{P}{4}$  while the Einstein tensor gives rise to Pauli-Fierz (PF) structure for  $h$  on the boundary [10],

$$L_4 = \frac{M_5^3}{2} h^{\mu\nu} (h_{\mu\nu} - h^2) - 6M_5^3 \frac{P}{4} + \frac{M_4^2}{4} \mathfrak{h} (E\mathfrak{h}) + \mathfrak{h} T ; \quad (14)$$

where  $h$  and  $\mathfrak{h}$  now denote the 5D fields evaluated at the 3-brane location.

In terms of the physical metric, (14) takes the form ,

$$L_4 = \frac{M_5^3}{2} \mathfrak{h}^{\mu\nu} \mathfrak{h}_{\mu\nu} - 3M_5^3 \frac{P}{4} \mathfrak{h} + \frac{M_4^2}{4} \mathfrak{h} (E\mathfrak{h}) + \mathfrak{h} T ; \quad (15)$$

Note that the kinetic term for  $\alpha'$  is completely absorbed by that of  $\mathfrak{h}$  and only a cross term between  $\alpha'$  and  $\mathfrak{h}$  remains.

From this it is straightforward to show the presence of a ghost. The scalar longitudinal component of  $h$  acquires

a positive kinetic term by mixing with the graviton [6, 7]. By taking

$$\mathfrak{h} = \mathfrak{h} + \frac{\alpha' \partial_\mu \partial_\nu}{m_5^2} ; \quad (16)$$

one finds that there are in fact two 4D scalar modes whose kinetic matrix in the UV is

$$\begin{pmatrix} \frac{3}{2} M_4^2 & 1 \\ -\frac{1}{4} & 0 \end{pmatrix} \quad (17)$$

which has obviously a negative eigenvalue corresponding to a ghost.

Having understood the origin of the ghost we are now ready to show how to cure it. To achieve this we clearly need to introduce a positive localized kinetic term for  $\alpha'$ . This can arise from extrinsic curvature contributions. The simplest and most natural choice is to put a tension on the 3-brane. This produces extrinsic curvature while leaving the metric on the brane flat since the tension only creates a deficit angle.

The solution to the 5D equations following from (12) for a 3-brane with tension  $\alpha'$  is [8]

$$h^{(0)} = \frac{\alpha'}{6M_5^3} \delta_{ij} ; \quad h^{(0)} = \frac{\alpha'}{6M_5^3} \delta_{ij} ; \quad (18)$$

This is an exact solution including the non-linear terms for  $\alpha'$  (they vanish identically for this profile). The background corresponds to a locally flat 6D bulk with deficit angle  $\alpha' M_6^4$  and with  $\alpha'$  at 4D sections.

The crucial point is that on this background the Lagrangian acquires contributions from the non-linear terms. These can be found considering the perturbations,

$$\begin{aligned} \alpha' &= \alpha'^{(0)}(z) + \alpha'(z; x) ; \\ h &= h^{(0)}(z) + h(z; x) ; \\ T &= T + T : \end{aligned} \quad (19)$$

Plugging (19) in (12) and dropping  $\alpha'$ , one obtains at quadratic order (up to a total derivative),

$$\begin{aligned} L_5 &= \frac{27M_5^3}{4m_6^2} \partial_M \partial_N \alpha'^M \alpha'^N - \alpha'^{(0)} (\partial_M)^2 \alpha'^{(0)} \\ &= \frac{9}{4m_6^2} (z) (\partial_\mu)^2 ; \end{aligned} \quad (20)$$

This is a localized kinetic term for  $\alpha'$  that contributes to the 4D effective action with a healthy sign when  $\alpha' > 0$ . Therefore for large enough  $\alpha'$  the kinetic matrix for the 2 scalars (17) becomes positive and the ghost is absent.

This can also be seen by computing the one particle exchange amplitude. With the addition of (20), the effective 4D equations are

$$\begin{aligned} M_4^2 (E\mathfrak{h}) &= 2M_5^3 \frac{P}{4} \mathfrak{h} + \mathfrak{h} \\ &= 2T + 6M_5^3 \frac{P}{4} ; \end{aligned} \quad (21)$$

$$\frac{3}{2m_6^2} \alpha' = M_5^3 \frac{P}{4} \mathfrak{h} ; \quad (22)$$

Using the Bianchi identities and the conservation of  $T$ , the double divergence of (21) leads to

$$M_5^3 \overset{P}{\square} (E\mathfrak{H}) + 6 \square_4 = 0; \quad (23)$$

where we have used that  $(E\mathfrak{H}) = 2(\square \square \mathfrak{H})_4$ . On the other hand, the trace of (21) in conjunction with (22) and (23), leads to

$$M_4^2 O_{( )} = 2T; \quad (24)$$

where  $O_{( )} = 9 = m_6^2 M_4^2 \square_4 - 24m_5 \overset{P}{\square} \square_4$ .

Combining (21), (23) and (24), one derives that the physical metric is, up to pure gauge terms,

$$\mathfrak{H} = \frac{2}{M_4^2} \frac{1}{O} T + \frac{T}{3} + \frac{1}{O_{( )}} T; \quad (25)$$

where  $O = \square_4 - 2m_5 \overset{P}{\square} \square_4$ . The tensor structure of the amplitude interpolates between  $d=4$  in the IR and

$$\frac{1}{3} + \frac{1}{6} \frac{3}{2M_4^2 m_6^2} \square_4$$

in the UV. The amplitude above corresponds to the exchange of massive spin 2 fields and a scalar obeying Eq. (24). The (DGP-like) kinetic term for the scalar is positive as long as,

$$> \frac{2}{3} M_4^2 m_6^2; \quad (26)$$

In this regime we see that the localized ghost disappears and the scalar sector is composed of a healthy resonance.

In the limit we are considering the tension required is consistent with having six non compact dimensions. Indeed, requiring that the deficit angle in the bulk is less than  $2\pi$  leads to  $< 2 M_4^4$ . It follows that,

$$3 M_5^6 > M_4^4 M_6^4; \quad (27)$$

which is always satisfied in the 5D decoupling limit and displays the necessity of the induced term on the codimension 1 brane. Moreover the condition above is

equivalent to having  $m_6 < m_5$ , which suggests that in order to avoid the ghost one should cascade from the highest dimension down to 4D 'step by step'.

From a phenomenological point of view, observations require that  $m_5 \sim H_0$ , the present Hubble scale. The most interesting possibility is when the 6D crossover scale is larger but of similar order. Assuming that the formulas above can be extrapolated in this regime for a Planckian  $M_4$  this implies that  $M_5$  is of order  $10M_{\text{eV}}$  and  $M_6 \sim m_{\text{eV}}$ . The latter also sets the scale of  $\Lambda$ .

#### IV. DISCUSSION

In this note we have presented a six dimensional DGP model with cascading localized kinetic terms. The model interpolates between a 6D behavior at large distances and a 4D one at short distances with an intermediate 5D regime. The kinetic terms regularize the divergent codimension 2 behavior. The model is ghost-free at least for a certain range of parameters if on the codimension 2 brane there is a large enough tension.

We have left several questions for future study. At linear-level the tensor structure of the graviton propagator is inconsistent with observations. In the context of DGP this was shown not to be a problem because the non-linearities restore the correct tensor structure [9]. A hint of a similar phenomenon in the present model is given by the longitudinal terms of the graviton propagator (6). These are singular when the mass parameters  $m_{5,6}$  vanish and give large contributions to nonlinear diagrams. In fact, we expect a 'double' Vainshtein effect. For dense enough sources, the non-linearities should first decouple the extra 5D scalar mode restoring 5D behavior followed by another step to 4D. Another important direction to study is cosmology. The model has the intriguing codimension 2 feature that tension does not curve the space. This obviously of interest for the cosmological constant problem.

Acknowledgements We thank Gregory Gabadadze for useful discussions. This work is supported in part by NSF grant PHY-0245068 and by the David and Lucile Packard Foundation Fellowship for Science and Engineering (GD and MR), by DURSI under grant 2005 BP-A 10131 (OP), and by NSERC and MRI (JK, AJT, CdR and SH).

[1] G. R. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B 484, 112 (2000) [arXiv:hep-th/0002190].  
 [2] S. L. Dubovsky and V. A. Rubakov, Phys. Rev. D 67, 104014 (2003) [arXiv:hep-th/0212222].  
 [3] G. Gabadadze and M. Shifm an, Phys. Rev. D 69, 124032 (2004) [arXiv:hep-th/0312289].  
 [4] N. Kaloper and D. Kiley, JHEP 0705, 045 (2007) [arXiv:hep-th/0703190].  
 [5] G. R. Dvali and G. Gabadadze, Phys. Rev. D 63, 065007 (2001) [arXiv:hep-th/0008054].  
 [6] M. A. Luty, M. Porrati and R. Rattazzi, JHEP 0309 (2003) 029 [arXiv:hep-th/0303116]; A. Nicolis and

R. Rattazzi, JHEP 0406 (2004) 059.  
 [7] N. Arkani-Hamed, H. Georgi and M. D. Schwartz, Annals Phys. 305, 96 (2003) [arXiv:hep-th/0210184].  
 [8] G. Dvali, G. Gabadadze, O. Pujlas and R. Rahman, Phys. Rev. D 75, 124013 (2007) [arXiv:hep-th/0612016].  
 [9] C. Deayet, G. Dvali, G. Gabadadze and A. Vainshtein, Phys. Rev. D 65, 044026 (2002) [arXiv:hep-th/0106001].  
 [10] The massless 5D graviton decomposes in a continuum of massive 4D gravitons. Therefore in the unitary gauge the boundary effective action will have PF structure.