

MULTIPLE TIME-SCALE METHODS IN PARTICLE SIMULATIONS OF PLASMAS†

BRUCE I. COHEN

*Lawrence Livermore National Laboratory
University of California, Livermore, CA 94550*

(Received March 7, 1985)

This paper surveys recent advances in the application of multiple-time-scale methods to particle simulation of collective phenomena in plasmas. These methods dramatically improve the efficiency of simulating low-frequency kinetic behavior by allowing the use of a large time step while retaining accuracy. The numerical schemes surveyed provide selective damping of unwanted high-frequency waves and preserve numerical stability in a variety of physics models: electrostatic, magnetoinductive, Darwin, and fully electromagnetic. The paper reviews hybrid simulation models, the implicit moment-equation method, the direct implicit method, orbit averaging, and subcycling.

I. INTRODUCTION

Particle codes are the most versatile and reliable tools for the study of complex kinetic plasma behavior. These codes follow the trajectories of thousands of sample particles in electromagnetic fields calculated self-consistently from Maxwell's equations. The numerical stability of these codes previously required resolution of the fast time scales associated with high-frequency waves. This was a significant limitation when the phenomena of interest occurred on much longer time scales. The recent introduction of implicit time-integration schemes¹⁻³ has partially removed these restrictions. The enormous range of time scales for single-particle and collective phenomena in most plasmas (Fig. 1) provides ample motivation for implicit methods and multiple-time-scale techniques in general.

It has long been recognized that implicit time integration was needed to relax time-step constraints in particle codes.^{4,5} The major inhibition had been the very large number of nonlinear equations to be solved simultaneously. In 1981 three groups independently formulated practical implicit particle codes.¹⁻³ They exhibited the dependence of the plasma response (the charge and current densities ρ and \mathbf{J}) on the electric field, linearized, and obtained a sparse matrix equation of rank equal to the number of electric- and magnetic-field quantities defined on the spatial grid. Solution was then achieved by standard methods.

Currently there are two principal approaches to implicit particle simulation. Mason¹ and Denavit² introduced fluid-moment equations describing charge and momentum conservation as intermediaries between the particle and field equa-

† This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract No. W-7405-ENG-48.




FIGURE 1 Space and time scales for plasma phenomena. (a) Schematic of plasma phenomena for a plasma generated by a neodymium-glass laser. (b) Schematic for a deuterium plasma in a magnetic mirror.

tions. Brackbill and Forslund⁶ extended this method to two-dimensional electromagnetic simulation. The implicit moment-equation approach (Section 2) can be viewed as a kinetic extension of a class of fluid-particle hybrid-simulation techniques in which a fluid species is advanced implicitly.^{7,8}

A more direct implicit approach (Section 3) has been introduced and refined by Friedman and Langdon³ and Cohen *et al.*,⁹ in which *no* auxiliary equations are introduced. In an electrostatic model, the charge density at the advanced time level is linearized about an explicit approximate density; and an increment is computed that is linear in the advanced field. The resulting field equation is elliptic, with coefficients depending on particle data accumulated on the spatial grid in the form of a susceptibility. The particles are advanced serially in a conventional manner. The direct implicit particle method has been implemented in two dimensions and for electromagnetic simulation.⁹

Another class of methods that improves particle-code efficiency takes advantage of multiple time scales in selecting time steps (Section 4). In an orbit-averaged magnetoinductive algorithm,¹⁰ particles are advanced with a small time step to resolve their orbits. An explicit solution for the fields, omitting electrostatic fields, is obtained using currents accumulated from the particles and temporally averaged. This reduces the number of particles and permits a large time step for the field advance. Orbit averaging in an electrostatic model requires an implicit field solution if a long time step is desired.¹ In an explicit algorithm using electron sub-cycling,¹² ions are advanced with a large time step much less often than the electrons are advanced and Poisson's equation is solved, making the cost of the ions negligible. Application of the gyrokinetic formalism employing analytical gyrophase averaging has also extended explicit codes to longer time steps.¹³

An overview of design criteria, limitations, and current research activity is presented in Section 5. This paper updates an earlier survey.¹⁴

II. HYBRID SIMULATION MODELS AND IMPLICIT MOMENT EQUATIONS

Hybrid Models

Implicit time-integration schemes have been applied to models in which fluid equations are used to represent one species and a particle description is taken for another. Two representative models are described here that were used to study magnetically confined plasmas that exhibit strong diamagnetic effects.^{7,8}

Hewett formulated a two-dimensional ($r - z$) hybrid simulation code to study magnetic pinches, whereas Cohen and Brengle⁸ performed one-dimensional (r) hybrid simulations of (magnetic) field-reversed plasmas. In both schemes, ions are simulated as particles and advanced explicitly; the electrons are an inertialess

fluid with equation of motion

$$0 = -\mathbf{E}_e - \frac{\nabla n_e T_e}{en} - \frac{\mathbf{u}_e \times \mathbf{B}}{c_e} + \frac{m_e \nu_0}{n_s e} \sum n_s z_s^2 (\mathbf{u}_s - \mathbf{u}_e), \quad (1)$$

where T_e and \mathbf{u}_e are the electron temperature and drift velocity, \mathbf{E} and \mathbf{B} are the electric and magnetic fields, \mathbf{J} is the total current, n_e is the electron number density, ν_0 is the electron-ion collision frequency, \mathbf{u}_s is the average ion drift velocity, and z_s is the ion charge state. The plasma is quasineutral, $n_e \approx \sum_s z_s n_s$. The electron current is $\mathbf{J}_e = -en_e \mathbf{u}_e$. In Hewett's model,⁷ the last term in Eq. (1) is replaced by $\boldsymbol{\eta} \cdot \mathbf{J}$, where $\boldsymbol{\eta}$ is the resistivity.

By evaluating \mathbf{E} in Eq. (1) at the advanced time and combining with the equation of magnetostatics:

$$\nabla \times \mathbf{B} = 4\pi \mathbf{J}/c \quad (2)$$

and Faraday's law,

$$\nabla \times \mathbf{E} = -\partial \mathbf{B}/c \partial t. \quad (3)$$

Hewett obtained implicit field equations. The ion charge and current densities are gathered explicitly from the particles. At very low densities, Hewett set η equal to a large value to force $\nabla \times \mathbf{B} = 0$. He solved the field equations globally by using a noniterative alternating-direction implicit algorithm.

In contrast to Hewett,⁷ who solved Eq. (1) for \mathbf{E} , Cohen and Brengle⁸ solved for \mathbf{u}_e and, hence, \mathbf{J}_e . Then

$$\hat{\theta} \cdot \nabla^2 \mathbf{A} = -\frac{4\pi}{c} J_\theta(A_\theta, \phi) \quad (4)$$

globally, and

$$\int \frac{dl}{rB} \frac{\partial}{\partial r} \left[4\pi r J_r(A_\theta, \phi) - r \frac{\partial^2}{\partial t \partial r} \phi \right] = 0 \quad (5)$$

on closed magnetic-field lines, where $\mathbf{E} = -\hat{\theta} \partial A_\theta/c \partial t - \hat{r} \partial \phi/\partial r$ and $\mathbf{B} = \nabla \times A_\theta \hat{\theta}$. Line tying on open field lines justifies $\partial \phi/\partial r = 0$. Canonical angular momentum relates the ion current in Eq. (4) to the vector potential A_θ , $\mathcal{L}_{\theta i} \equiv m_i r v_\theta + z_i e r A_\theta$. Equations (4) and (5) are solved iteratively (≤ 5 iterations) after linearizing J_θ with respect to A_θ . Temporally averaging J_{ri} over past data reduces statistical noise.

Both Hewett's⁷ and Cohen and Brengle's⁸ algorithms avoided potential numerical difficulties arising from the infinite phase velocity of Alfvén waves as the mass density approaches zero, $\omega/k = v_A \equiv (B^2/4\pi n m_i)^{1/2} \rightarrow \infty$. In these implicit hybrid codes, the stability condition $\Delta x/\Delta t > v_A$ is relaxed. Time-step constraints are set instead by accuracy considerations and stability of the ion orbits.^{7,8}

Implicit Moment Equations

Mason¹ and Denavit² independently synthesized simulation schemes whose implicitness is derived from the introduction of fluid-moment equations. This

improves the hybrid models in that all species are now kinetic and implicitly coupled to the fields. The implicit moment method has good stability properties and is broadly applicable.

In a one-dimensional electrostatic model, the fluid equations in difference form are

$$n_s^{n+1} = n_s^n - \Delta t D_x J_s^{n+1/2} / q_s; \quad (6)$$

$$J_s^{n+1/2} = J_s^{n-1/2} + q_s \Delta t (-D_x P_s^{+n} + q_s n_s^n E^*) / m_s, \quad (7)$$

where $P_s^+ = \Delta x^{-1} \sum_i m_s v_i^2$ is the kinetic stress summed over the particles, D_x is the difference form of $\partial/\partial x$, n_s and J_s are implicit predictions of the fluid number and current densities, n_s and J_s are the explicit densities accumulated from the particles, and

$$E^* = \theta E^{n+1} + \frac{(1-\theta)}{4} (E^{n+1} + 2E^n + E^{n-1}); \quad 0 \leq \theta \leq 1. \quad (8)$$

The electric field is calculated from Gauss' law:

$$D_x E^{n+1} = 4\pi \sum q_s n_s^{n+1}, \quad (9)$$

rendered implicit by use of Eqs. (6) and (7). Particles are then advanced individually with the implicitly predicted electric field, and the necessary particle data are gathered.

An iteration can be performed to better time-center P_s^+ , whose convergence requires²

$$kv\Delta t < 1, \quad (10)$$

where v is a characteristic particle velocity and k is the largest wave number retained. This is also the condition for accurate particle trajectories and plasma dielectric response.^{5,9} The implicit field solution relaxes the stability constraint set by plasma waves, allowing $\Delta t > \omega_p^{-1}$, the inverse plasma frequency. With $\omega_p \Delta t > 1$, Eq. (10) restricts wavelengths to be long compared to the Debye length,

$$k\lambda_D \equiv k(T/4\pi n e^2)^{1/2} = kv_t \Delta t / \omega_p \Delta t \ll 1, \quad (11)$$

where $v_t \equiv (T/m)^{1/2}$. An implicit prediction of the kinetic stress relaxes the $kv\Delta t$ stability constraint^{1,6} but does not remove it as an accuracy constraint.^{5,9}

For $\omega_p \Delta t \ll 1$ and $\Delta x > \lambda_D$, there can be a grid-aliasing instability,¹⁵ which results in heating until $\lambda_D \sim \Delta x$. However, implicitness, dissipation,⁵ and use of $\omega_p \Delta t > 1$, or a change of the force law,¹⁵ control the grid instability.^{2,6} Thus, for $\Delta x \gg \lambda_D$, the time step in the implicit moment algorithm is bounded from above by Eq. (10) and from below by the finite-grid instability. These limits are not a serious hindrance.^{1,2,6}

Mason¹ and Denavit² have successfully applied this scheme to a number of problems. Of particular importance are simulations of electron transport in inertial-confinement fusion.¹⁶ Brackbill and Forslund⁶ have extended the implicit

moment method to impressive two-dimensional electromagnetic simulations of the Weibel and lower-hybrid-drift instabilities, shocks, and collisionless electron transport in laser fusion.⁶ The implicit moment method has been applied by Barnes and Kamimura¹⁷ to two-dimensional, electrostatic simulations of low-frequency phenomena in magnetized plasma using both guiding-center and Newton-Lorentz particle equations. J. A. Byers has applied the moment method to linearized electrostatic simulations of unstable Bernstein waves (unpublished).

III. DIRECT IMPLICIT PARTICLE METHODS

In the direct implicit particle method,^{3,9} an implicit solution of the field equations is achieved by relating linear increments to the charge and current density directly to the change in the particle motion induced by the fields or their increments at the advanced time. This differs markedly from the implicit moment method, but the two methods possess similar stability and accuracy properties.^{5,9}

In a simple one-dimensional electrostatic model,³ the direct implicit method has the following form. The particle position x^{n+1} at time t^{n+1} is

$$x^{n+1} = \beta \Delta t^2 a^{n+1} + \bar{x}^{n+1}, \quad (12)$$

where $0 < \beta < 1$ and \bar{x}^{n+1} is the position with a^{n+1} suppressed in the equation of motion. Thus, $x^{n+1} = \bar{x}^{n+1} + \delta x$, where $\delta x = \beta \Delta t^2 a^{n+1}$, and the charge density is $\rho^{n+1} = \bar{\rho}^{n+1}(x^{n+1}) + \delta \rho$, where

$$\delta \rho = -\nabla \cdot [\bar{\rho}^{n+1}(x) \delta \mathbf{x}(x)] \quad (13)$$

is the linearized increment to the charge density. Poisson's equation gives

$$-\nabla \cdot (1 + \chi) \nabla \phi^{n+1} = 4\pi \bar{\rho}^{n+1}, \quad (14)$$

where the effective susceptibility is

$$\chi(x) = 4\pi \beta (q \bar{\rho}^{n+1}/m) \Delta t^2 = \beta \omega_p^2(x) \Delta t^2. \quad (15)$$

The charge density is related to the particle positions by

$$\rho_j^{n+1} = (q/\Delta x) \sum_k S(x_k^{n+1} - x_j), \quad (16)$$

where j is the grid index, k is the particle index, q is the charge, and S is the "shape function" for particle-mesh interpolation. Expanding S gives

$$S(x_k^{n+1} - x_j) \approx S(\bar{x}_k^{n+1} - x_j) + (x_k^{n+1} - \bar{x}_k^{n+1}) \frac{\partial S(\bar{x}_k^{n+1} - x_j)}{\partial \bar{x}_k^{n+1}}, \quad (17)$$

and from Eq. (12),

$$x_k^{n+1} - \bar{x}_k^{n+1} \approx \beta (q/m) \Delta t^2 \sum_i S(\bar{x}_k^{n+1} - x_i) E_i^{n+1}, \quad (18)$$

with relative error of order $\omega_{tr}^2 \Delta t^2 = qE \Delta t^2 / m L_E < 1$, where ω_{tr} is the particle-trapping frequency and L_E is the length scale over which E varies.

With the use of Eqs. (14), (17), and (18), Poisson's equation becomes

$$-(\phi_{j-1}^{n+1} - 2\phi_j^{n+1} + \phi_{j+1}^{n+1})/\Delta x^2 = 4\pi\bar{\rho}_j^{n+1} + \sum_i W_{ij}E_i^{n+1}, \quad (19)$$

where $\bar{\rho}_j^{n+1}$ is the conventional charge density given by Eq. (17) with $x_k^{n+1} \approx \bar{x}_k^{n+1}$ and

$$W_{ij} = \frac{\beta 4\pi q \Delta_i^2}{m\Delta x} \sum_k S(\bar{x}_k^{n+1} - x_i) \frac{\partial S(\bar{x}_k^{n+1} - x_j)}{\partial x_k^{n+1}}. \quad (20)$$

For linear splines, $W_{ij} = 0$ whenever $|i - j| > 1$ and $\partial S/\partial x_k^{n+1} = \pm 1$ or 0. No additional particle data beyond that needed for ρ_j^{n+1} are needed for W_{ij} , and the field equation is a linear pentadiagonal system solved by direct Gaussian inversion.³

The direct method has been tested in simulations of wave propagation, two-stream instability, and free expansion of a plasma slab; and the implicit difference scheme has been varied.^{5,18} If the plasma is nearly uniform so that $\chi(x)$ approximately equals its spatial average, Fourier-transform methods can be used.¹⁹

Langdon *et al.*⁹ have formulated the direct implicit method with a magnetic field and in two dimensions. The spatial differencing of Eq. (14) can be simplified to reduce the bandedness of the matrix equation for the field. Langdon *et al.*⁹ address the subtle problem of self-consistent spatial filtering and also prove that the implicit field-particle matrix equation is positive and in some cases symmetric, useful properties for iterative solutions. Reference 9 also outlines a simple iteration scheme to obtain a more exact solution of Poisson's equation. The convergence of the iterations, the influence of electric-field extrapolation on convergence, and the influences of spatial smoothing on linear dispersion and convergence have been analysed in Ref. 20.

Langdon and Barnes describe the implementation of the direct implicit method in an electromagnetic code in Ref. 21. The implicit increment to the current density in the direct implicit electromagnetic code is analogous to the increment to the charge density given in Eq. (13):

$$\delta \mathbf{J} = \bar{\rho} \delta \mathbf{v} - (1/2) \nabla \times (\bar{\mathbf{J}} \times \delta \mathbf{x}), \quad (21)$$

where $\bar{\mathbf{J}}_j = \sum q \mathbf{v} S(\mathbf{x}_j - \bar{\mathbf{x}}_k)$ and $\delta \mathbf{x}$ and $\delta \mathbf{v}$ are the linearized increments to \mathbf{x} and \mathbf{v} due to the electric field \mathbf{E}^{n+1} at the advanced time step. In a leapfrog algorithm with positions and velocities known $\Delta t/2$ apart, a time average of successive position time levels achieves time centering in the current density. Reference 21 provides a detailed discussion of the differencing and the properties of the direct implicit electromagnetic code.

The direct method relaxes the usual stability constraints on $\omega_p \Delta t$ and $c \Delta t / \Delta x$. Where $kv_i \Delta t < 1$ is a stability constraint in the implicit-moment method, $\omega_r \Delta t < 1$ is required in the direct method. However, $kv_i \Delta t < 1$ remains as an accuracy constraint; and the grid-aliasing instability must be controlled. Finally, the direct method avoids the possible difficulty in the implicit-moment method that the fluid number and current densities are inconsistent with those accumulated from the particles.

IV. ORBIT AVERAGING AND SUBCYCLING

Orbit averaging^{10,11} and subcycling¹² take a multiple-time-scale approach to selecting time steps in particle simulation. Independent time scales are selected for advancing particles and fields according to natural separations that exist. The subcycling method succeeds in making the ions a negligible factor in the cost of simulating ions and electrons.¹² However, the algorithm described in Ref. 12 is explicit and therefore does *not* allow the use of a large $\omega_{pe}\Delta t$.

Orbit averaging^{10,11} differs from subcycling and is more complicated. In its original magnetoinductive form,¹⁰ a particle species gyrating in a magnetic field is explicitly advanced with a small time step, $\omega_c\Delta t < 1$, where $\omega_c \equiv qB/mc$. Currents are accumulated on a spatial grid after each small time step and are temporally averaged over ΔT , $\omega_c\Delta T \gg 1$, for use in Ampere's law to determine the vector potential and magnetic field. An inductive electric field is calculated from Faraday's law, and electrostatics are ignored. The fields are advanced with the large time step (Fig. 2). A corrector iteration through the particle and field equations is performed to improve the time centering. Biasing of the field equations to the field amplitudes at the advanced time level introduces dissipation.

Analysis and simulations show that the orbit-averaged magneto-inductive scheme, although explicit, is numerically stable with use of large $\Delta T : \Delta t \gg \Delta t$, $\omega_c\Delta T \gg 1$, and $kv_A\Delta T \gg 1$.¹⁰ However, this scheme is unstable for $\omega_c\Delta T \leq \mathcal{O}(1)$. The real triumph of orbit averaging is how it significantly reduces the number of particles and total operations; the averaged contributions from a single particle can substitute for those from many particles in a conventional code. This increased efficiency has allowed realistic two-dimensional simulations of mirror experiments at Lawrence Livermore National Laboratory.¹⁰

In contrast to the magnetoinductive algorithm, stability of an orbit-averaged electrostatic algorithm for $\omega_p\Delta T > 1$ requires implicit solution of Poisson's equation.¹¹ The combination of orbit averaging with the direct implicit or implicit moment methods can produce algorithms that are stable for $\omega_p\Delta T > 1$.¹¹ Orbit




FIGURE 2 Schematic of time levels in generic orbit averaging. The quantities $\langle n \rangle$ and $\langle J \rangle$ are averaged in time over the small time step quantities.

averaging in this application achieves a reduction of particles and allows the particles to be advanced serially over many time steps before incurring input/output penalties, an advantage for large simulations using disk storage. Even greater gains in efficiency can be achieved with a better separation of particle and field time steps, by using an implicit prediction of the kinetic stress⁶ or when wave propagation is perpendicular to B .¹¹

V. GENERAL CONSIDERATIONS

In developing new implicit difference schemes for particle simulation, design criteria have evolved:

- Accurate reproduction of low-frequency phenomena, $\omega < \Delta t^{-1}$, with minimal damping.
- Substantial damping of modes with $\omega > \Delta t^{-1}$.
- Minimal collection and storage of particle data.
- Minimal numerical cooling and heating.⁵
- Galilean invariance so as not to destabilize fast or slow space-charge waves.⁵
- Robust stability and accuracy properties with respect to approximate solution for the fields.

Analyses of wave-dispersion properties in Refs. 2, 5, 6, and 20 have led to a number of important conclusions. With proper choice of coefficients in the differencing schemes,⁵ damping of high-frequency oscillations $\omega\Delta t \geq 1$ can be enhanced, while the damping of low-frequency waves can be removed to high order in $\omega\Delta t < 1$. The wave-dissipation properties and unphysical secular particle acceleration are directly related.⁵

Both implicit moment and direct implicit algorithms have proven successful in performing simulations with large time step. Magnetic fields have been incorporated,^{5,6,9,21} and iterative refinement of the solution of the implicit field-particle equations has been studied.^{1,2,9,20} The two methods share residual constraints on the time step that generally coincide with those required to resolve physical processes, e.g., $k\lambda_D < 1$, $kv_i\Delta t < 1$.^{2,5,6,9} Some merging of the two methods has occurred,^{6,9,21,22} and collisional effects have been included.²²

Research on advanced particle-code techniques continues. Areas of interest are numerous. It would be desirable to relax residual constraints. The problem of self-consistent spatial smoothing⁹ has been resolved recently.²⁰ The potential for (and limitations of) simplified spatial differencing is being examined. More simulation experience is needed, especially with direct implicit and orbit-averaged algorithms. A two-dimensional electromagnetic direct implicit simulation code has been built, and relativity should be added. More needs to be done on how to best realize additional computational savings by combining orbit averaging or subcycling with implicit methods and by taking advantage of multitasking capabilities on the new supercomputers. Research continues, motivated by the impressive successes of these new techniques in dramatically extending the applicability of particle simulations and greatly improving their realism.^{6,10,16,20,22}

VI. ACKNOWLEDGMENTS

It is a pleasure to thank A. B. Langdon, R. P. Freis, and A. Friedman for their collaboration, and J. A. Byers and C. Gerich for their comments.

REFERENCES

1. R. J. Mason, *J. Comput. Phys.*, **41**, 233 (1981).
2. J. Denavit, *J. Comput. Phys.*, **42**, 337 (1981).
3. A. Friedman, A. B. Langdon, and B. I. Cohen, *Comments Plasma Phys. Contr. Fusion*, **6**, 225 (1981).
4. A. B. Langdon, *J. Comput. Phys.*, **30**, 202 (1979).
5. B. I. Cohen, A. B. Langdon, and A. Friedman, *J. Comput. Phys.*, **46**, 15 (1982).
6. J. U. Brackbill and D. W. Forslund, *J. Comput. Phys.*, **46**, 271 (1982); D. W. Forslund and J. U. Brackbill, *Phys. Rev. Lett.*, **48**, 1614 (1982); J. U. Brackbill and D. W. Forslund, in *Multiple Time Scales*, edited by J. U. Brackbill and B. I. Cohen (Academic Press, New York, 1985).
7. D. W. Hewett, *J. Comput. Phys.*, **38**, 378 (1980).
8. T. A. Brengle, B. I. Cohen, and M. E. Stewart, *Computer (IEEE)*, **16**, 44 (1983).
9. A. B. Langdon, B. I. Cohen, and A. Friedman, *J. Comput. Phys.*, **51**, 107 (1983).
10. B. I. Cohen, T. A. Brengle, D. B. Conley, and R. P. Freis, *J. Comput. Phys.*, **38**, 45 (1980); B. I. Cohen and R. P. Freis, *J. Comput. Phys.*, **45**, 367 (1982).
11. B. I. Cohen, R. P. Freis, and V. Thomas, *J. Comput. Phys.*, **45**, 345 (1982).
12. J. C. Adam, A. Gourdin Serveniére, and A. B. Langdon, *J. Comput. Phys.*, **47**, 229 (1982).
13. W. W. Lee, *Phys. Fluids*, **26**, 556 (1983).
14. B. I. Cohen, in *Energy Modeling and Simulation Proceedings*, edited by A. S. Kydes, 10th World Congress on Systems Simulation and Scientific Computation, Montreal, 1982 (North Holland, Amsterdam, 1983) p. 383.
15. A. B. Langdon, *J. Comput. Phys.*, **6**, 247 (1970).
16. R. J. Mason, *Phys. Rev. Lett.*, **47**, 652 (1981).
17. D. C. Barnes and T. Kamimura, *Bull. Am. Phys. Soc.*, **26**, 986 (1981).
18. A. Friedman, A. B. Langdon, and B. I. Cohen, *Bull. Am. Phys. Soc.*, **26**, 985 (1981).
19. T. Tajima and J. N. Leboeuf, *Bull. Am. Phys. Soc.*, **26**, 986 (1981).
20. B. I. Cohen, A. B. Langdon, and A. Friedman, *J. Comput. Phys.*, **56**, 51 (1984).
21. A. B. Langdon and D. C. Barnes, in *Multiple Time Scales*, edited by J. U. Brackbill and B. I. Cohen (Academic Press, New York, 1985).
22. R. J. Mason, in *Multiple Time Scales*, edited by J. U. Brackbill and B. I. Cohen (Academic Press, New York, 1985).