

O rientifolds and D -branes in N = 2 gauged linear sigma models

Ilka Brunner^{a,b}, Manfred Herbst^c

^a Arnold Sommerfeld Center
Ludwig Maximilians Universität
Theresienstr. 37
80333 München, Germany

^b Excellence Cluster Universe
Technische Universität München
Boltzmannstr. 2
85748 Garching, Germany

^c Theory Division
Department of Physics, CERN
CH -1211 Geneva 23
Switzerland

Ilka.Brunner@physik.uni-muenchen.de, Manfred.Herstb@cern.ch

Abstract

We study parity symmetries and boundary conditions in the framework of gauged linear sigma models. This allows us to investigate the Kähler moduli dependence of the physics of D-branes as well as orientifolds in a Calabi-Yau compactification. We first determine the parity action on D-branes and define the set of orientifold-invariant D-branes in the linear sigma model. Using probe branes on top of orientifold planes, we derive a general formula for the type (SO vs Sp) of orientifold planes. As applications, we show how compactifications with and without vector structure arise naturally at different real slices of the Kähler moduli space of a Calabi-Yau compactification. We observe that orientifold planes located at certain components of the fixed point locus can change type when navigating through the stringy regime.

1 Introduction and results

Orientifolds and D-branes play an important role for the consistency of type II string compactifications [1,7] as both classes of objects are needed to ensure a balance of Ramond (Ramond charges and to preserve spacetime supersymmetry at the same time.

In this paper we are interested in B-type orientifolds of Calabi-Yau manifolds, and in particular their dependence on the Kähler moduli. A suitable framework to investigate these issues are gauged linear sigma models [8], which provide the possibility to interpolate between the large and small radius regime of a Calabi-Yau compactification. Here, the compactification is described in terms of a $1+1$ dimensional abelian gauge theory; the stringy Landau Ginzburg point and the geometric limit are located at different limits of the Fayet-Iliopoulos-parameters r of the gauge theory. Together with the theta angles the combination $t = r + i\theta$ parametrizes the Kähler moduli space. Here, the theta angle contains in particular the information on the B-field at large volume.

The possibility of turning on a discrete B-field plays an important role in the discussion of type I string theory or, more generally, of orientifolds in type IIB string theory. In particular, it implies the possibility of compactifications without vector structure [9,12]. In the context of the linear sigma model, the different discrete values of the B-field descend from different real slices in the Kähler moduli space parametrized by [13,14]. In particular, the linear sigma model allows to understand large volume compactifications

distinguished by B-elds as extremal limit points of different branches of a stringy moduli space. In some cases the branches can get connected in the stringy regime, such that it becomes possible to navigate from one large volume point to another taking a path in the interior of the moduli space. However, the interior of the moduli space contains a singular locus, and the real slices singled out by the orientifold projection might pass through it, depending on the particular value of the theta angles; this was observed in [14] and will be reviewed and worked out in detail below.

An important problem is to understand the D-brane categories compatible with the orientifold projection [15,16]. At the Landau-Ginzburg point D-branes are described in terms of matrix factorizations of the superpotential, and the brane category relevant for the description of unoriented strings has been constructed in [17], cf. also [18]. On the other hand, a geometric description of branes on Calabi-Yau manifolds is provided by the derived category of coherent sheaves, and parities have been studied in this context in [19]. In this paper, we lift the constructions of these two approaches to the linear sigma model, thereby connecting different corners in the Kähler moduli space. For D-branes without orientifolds this analysis was already carried out in [20], and before in the mathematics literature (up to monodromies) in [21{26]. Earlier results on the level of Ramond/Ramond charges were obtained for D-branes in [27{31] and including orientifolds in [14].

Once the parity action on D-branes is understood, we can proceed and determine under certain assumptions the type of an orientifold plane ($S0$ vs Sp gauge group). Generically, the fixed point set of the parity action consists of several irreducible components, and the type of the individual orientifold planes can be tested by determining the gauge group on probe branes positioned on top of the fixed point set. We work out explicit formulas that determine the orientifold type (up to an overall sign to be fixed once and for all for each parity) from the linear sigma model data of the brane and the parity. With this at hand, we show that the orientifold type can change when navigating through the non-geometric regime. Similar effects have already been observed in [14,32,33] using tadpole cancellation conditions. In the cases where large volume regimes with different values of the B-eld are connected in the interior of the moduli space, we observe that the type of the orientifold plane changes along the path. This of course is in agreement with the fact that, at least for toroidal orientifolds, compactifications distinguished by a B-eld at large volume correspond to compactifications with or without vector structure. Interestingly, we also find non-trivial monodromies: starting out at large volume, continuing to the stringy regime and going back to the same large volume point with the same B-eld, a change of type can be observed in examples.

To give a further application of our techniques, we consider configurations of $O7$ -planes and singular $D7$ -branes with $SO(N)$ gauge group, which have been studied recently in the context of F -theory model building [34,36]. In fact, the $D7$ -brane carries a curve of ordinary double points that lies on the intersection with the orientifold plane and that pinches off at a collection of points. F -theory and probe branes in type IIB were used in [35,36] to argue that the $D7$ -brane geometry in the presence of the orientifold is constrained to be singular, admitting fewer deformation parameters than a $D7$ -brane on a generic hypersurface. We will give an explanation of the singularity that relies just on the requirement to have an orientifold-invariant D -brane with the right gauge group.

The issue of tadpole cancellation and the construction of consistent supersymmetric string vacua is one out of several interesting model building applications, which we omit at present, but hope to address in future work. This question has however been investigated in some detail, for instance at points of enhanced symmetry using explicit constructions in rational conformal field theory [14,37,42]. In our context the Gepner point corresponds to the Landau-Ginzburg point and all the RCFT branes considered in the papers cited correspond to very simple matrix factorizations of the superpotential. However, the techniques presented in this work provide many more possibilities of constructing consistent string vacua¹ and additionally give control over the Kähler moduli dependence.

The role of orientifolds and D -branes for tadpole cancellation in the topological string was revealed in [43], following earlier work on open string mirror symmetry [44,45]. We expect that the present paper paves the way to consider more general tadpole cancelling states in this context.

In the following, we give a brief outline of the paper and its main results in more detail.

D -branes

In order to set the stage we start this work with a brief review section on gauged linear sigma models with abelian gauge group $T = U(1)^k$ [8,20]. This section can be skipped by readers that are familiar with the results of [20].

¹For example, not all Landau-Ginzburg models correspond to rational conformal field theories. Even if the bulk theory is rational, most branes will break the enhanced symmetry making a conformal field theory construction hard, while a Landau-Ginzburg description is still possible.

In particular, we introduce the complexified Kähler moduli space $M_K = (\mathbb{C}^k)^n / S^2$, where S is the singular locus of complex codimension one on which the world sheet description breaks down in view of massless D-branes [46].

We define D-branes in the linear sigma model as matrix factorizations or complexes of Wilson line branes and explain the notion of D-isomorphism classes, or equivalently quasi-isomorphism classes, which define the set of low-energy D-branes in each phase of the linear sigma model. The transport of D-branes across phase boundaries is implemented in view of the grade restriction rule, which is a "gauge" fixing condition on the D-isomorphism classes and depends on the path between phases. We also briefly discuss the Brezinski-Knorrer map that relates the matrix factorizations of the linear sigma model to geometric D-branes on the hypersurface or complete intersection in the low-energy theory.

Orientifolds

After these preparations we proceed in Sec. 3 with defining and studying B-type parity actions and orientifolds in gauged linear sigma models, first on a world sheet without boundary. The world sheet parity action is the composition of three operators, $P = (-1)^{F_L}$ for $m \in \mathbb{Z}$. It flips the orientation of the world sheet, is a holomorphic involution acting on the chiral fields of the linear sigma model, and for m odd the operator $(-1)^{F_L}$ flips the sign for left-moving states in the Ramond sector.

We observe the well-known effect that only slices in M_K of real dimension k survive the orientifold projection [14]. In fact, there are 2^k such slices parametrized by \mathbb{Z}_2 -valued theta angles $\theta = (\theta_1; \dots; \theta_k)$ for $\theta_i \in \mathbb{Z}_2$; g. Each slice may or may not intersect the singular locus S , which is now real codimension one and cannot be avoided by any path. This leads to the observation that some phases of the linear sigma model are not connected to others, at least not in a world sheet description.³ Somewhat surprising, there are even non-perturbative regions "deep inside" the moduli space that are not connected to any of the phases where, at least in principle, perturbative string methods can be applied.

The fixed point set of the holomorphic involution takes a particularly simple form. For linear sigma models without superpotential (which have toric varieties as low-energy configurations) it splits into a finite number of irreducible components, the orientifold planes O , that are parametrized by a discrete choice of k phases $\theta = (\theta_1; \dots; \theta_k)$. For linear sigma models with

²Here, we mean the Kähler moduli space before orientifold projection.

³For an M-theory analysis that allows avoiding the singularity see [47].

superpotential the components O may become reducible at low-energies so that they split up into a finite number of irreducible components O_i . The explicit parametrization of the irreducible components of the fixed point locus turns out valuable for determining a simple formula for the types of the individual orientifold planes.

Orientifolds and D-branes

In Sec. 4 we investigate the world sheet parity action in the presence of boundaries and define the set of invariant D-branes in the gauged linear sigma model. The latter depends on the following data: (i) the slice on the Kähler moduli space, (ii) the integer m that controls the appearance of $(-1)^{F_L}$, (iii) the involution and (iv) a sign associated with the orientifold. In fact, changing the latter sign flips the gauge groups, $SO(n)$ to $Sp(n=2)$ or vice versa, of all invariant D-branes as well as the type of all orientifold planes simultaneously.

On a slice of M_K where two adjacent phases of the linear sigma model are not separated by the singular locus we can still move D-branes between the two phases by applying the grade restriction rule of [20]. We show that the latter is compatible with the world sheet parity action and can indeed be applied to invariant D-branes.

A particularly important piece of information on an invariant D-brane is the type of its gauge group [6,17,19]. Applying our formalism we are able to derive an explicit formula (81) for the sign that determines the gauge group (SO or Sp) of an important class of invariant D-branes, i.e. D-branes given by Koszul complexes (or Koszul-like matrix factorizations) that localize at the intersection of a finite number of holomorphic polynomials.

In Sec. 5 resp. 6 we proceed discussing non-compact models (without superpotential) and compact models (with superpotential) separately, as some of the results will depend on whether we deal with complexes or matrix factorizations.

In Sec. 6.1 we consider the effect of the (bre-wise) Konner-Map on the world sheet parity action and on the set of invariant D-branes.

In Sec. 5.1 and 6.2 we have a closer look at the Kähler moduli space M_K and its slicing by the discrete theta angles. In general, the slices are not connected. However, at special loci of the moduli space, such as orbifold points or Landau-Ginzburg orbifold points, they can be connected, cf. [14]. In the linear sigma model this can be seen by considering the set of invariant D-branes at these special loci. For higher-dimensional moduli spaces this leads to the phenomenon that large volume points corresponding to different

values of the discrete B-field can be connected through a path in moduli space.

We continue in Sec. 5.2 and 6.3 with computing explicit formulas (94) and (106) for the type $\sigma = \pm 1$ of an orientifold plane O_σ by testing the gauge group of a probe brane on top of the orientifold plane. We find that the relative types of the various fixed point components O_σ depend on the slice in M_k . In particular, the type σ is proportional to the character $\chi_\sigma = (\chi_1)^{\pm 1} \cdots (\chi_k)^{\pm k}$.

In Sec. 6.4 we discuss the simple example of O_7 -planes at four points on the torus. Depending on the choice of the B-field, this configuration is T-dual to an orientifold with or without vector structure. We reproduce the result of [9], where it was found that for vanishing B-field all four points carry the same type, whereas for non-vanishing B-field one point carries a type opposite to the other three points. In Sec. 5.3 and 6.5 we exemplify the phenomenon of type change along continuous paths in moduli space in two-parameter models. We close this work in Sec. 6.6 by commenting on the weak-coupling limit of a certain F-theory compactification that was discussed in [35,36].

2 A brief review of D-branes in gauged linear sigma models

In this section we introduce gauged linear sigma models and review the main results and concepts of [20] for describing D-branes.

The motivation to consider $N = (2;2)$ supersymmetric gauged linear sigma models relies on the observation that they provide an ultra-violet description for $N = (2;2)$ superconformal field theories such as a non-linear sigma model on Calabi-Yau hypersurfaces [8]. In that way the complicated non-linear sigma model is lifted to a model with linear target space C^N described by chiral multiplets X_i for $i = 1, \dots, N$, while all non-linear interactions are governed by the coupling of the chiral multiplets to gauge multiplets V_a for $a = 1, \dots, k$.

In this work we consider only abelian gauge groups $T = U(1)^k$. The action of the gauge group on the chiral multiplets is controlled by the integral charges Q_i^a , i.e. $g X_i = g^{Q_i^a} X_i$, where $g^{Q_i^a} = g_1^{Q_i^1} \cdots g_k^{Q_i^k}$ for an element $g \in T$.

The classical action involves a gauge-invariant F-term superpotential, $W(X) = g W(X)$, whose coefficients parametrize the complex structure moduli space M_C in the infra-red theory. In this work we are not interested

in deforming the complex structure and fix the coefficients in the superpotential once and for all.

The action furthermore includes a twisted superpotential $\int^P t^a v_a$ where $v_a = D + D - V_a$ is the gauge field strength. The parameters $t^a = r^a - i^a$ turn out to become coordinates on the (complexified) Kähler moduli space M_K of the low-energy theory. The Fayet-Iliopoulos parameters r^a take values in \mathbb{R}_{FI}^k , and the theta angles i^a enter in the action via a topological term that measures the instanton number of the gauge bundle and therefore take values in $(S^1)^k$. It is convenient to work with the parametrization $e^t = (e^{t^1}; \dots; e^{t^k}) \in (\mathbb{C}^*)^k$.

Phases in the classical Kähler moduli space

The main advantages of the gauged linear sigma model over the non-linear sigma model is its explicit dependence on the Kähler moduli space M_K , even more so as moving around in M_K involves generalized \mathbb{C}^* transitions between low-energy geometries, which are hard to control in the non-linear sigma model but can be studied easily in the gauged linear sigma model.

Classically the infra-red dynamics is governed by the zeros of the potential

$$(1) \quad U_{\text{pot}} = \sum_{i=1}^N \sum_{a=1}^k Q_i^a x_i^2 + \sum_{a=1}^k \frac{e_a^2}{2} \sum_{i=1}^N Q_i^a x_i^2 - r^a + \sum_{i=1}^N \int @_i W(x) dx^2;$$

where x_i are the lowest components of the chiral multiplets, and v_a are the complex scalars in the vector multiplets. Setting $U_{\text{pot}} = 0$ requires that each term in (1) has to vanish individually. The second one yields the D-term equations

$$(2) \quad \sum_{i=1}^N Q_i^a x_i^2 = r^a; \quad \text{for } a = 1; \dots; k;$$

and the last one the holomorphic F-term equations

$$(3) \quad @_i W(x) = 0; \quad \text{for } i = 1; \dots; N;$$

Let us first consider the situation without superpotential, $W(x) = 0$. The solutions to the D-term equations modulo gauge transformations restrict the chiral fields x_i to the symplectic quotient $\mathbb{C}^1(r) = T$, which is in fact a toric variety. It will suffice and in fact be more convenient in the following to drop the explicit dependence on the parameters r^a and work with the algebraic instead of the symplectic quotient. The latter is given by

$$(4) \quad X_r = \frac{\mathbb{C}^N}{(\mathbb{C}^*)^k};$$

where $(\mathbb{C}^*)^k$ is the complexification of the gauge group T . In fact, X_r is the space of $(\mathbb{C}^*)^k$ -orbits in \mathbb{C}^N that intersect the solution set of the D-term equation (2). The deleted set \tilde{X}_r contains precisely the subset of points in \mathbb{C}^N , whose $(\mathbb{C}^*)^k$ -orbits do not intersect (2).

For generic values of the parameters r^a the first term in the potential (1) provides a non-degenerate mass matrix $M^{ab}(x; x)$ for the scalars ϕ_a and therefore sets them to zero.

As we move around in R_{FI}^k the symplectic quotient changes and can undergo generalized flop transitions. The flops occur at (real) codimension one walls, which subdivide the FI-space into phases (or Kahler cones), and are usually referred to as phase boundaries. In terms of the algebraic quotient X_r the walls are the locations where the deleted set \tilde{X}_r changes.

In view of the potential (1) the positions of the phase boundaries in R_{FI}^k are the loci where the D-term equation (2) admits a solution such that the mass matrix $M^{ab}(x; x)$ degenerates. Consequently, a subgroup $U(1)_?$ remains unbroken and the corresponding scalar $\phi_?$ can take non-vanishing expectation values, thus leading to non-normalizable wave functions and therefore to a singularity in the low-energy theory.


If we turn on a superpotential $W(x)$ the F-term equations limit the low-energy dynamics to a holomorphic subvariety in X_r . Generically, the directions transverse to (3) are not massive, and the fields x_i can still fluctuate around (3) so that we end up with a Landau-Ginzburg model with potential $W(x)$ over the base toric variety X_r . In the other extreme, if all transverse directions are massive, the theory is confined to the subvariety given by $\partial_i W(x) = 0$ for $i = 1; \dots; N$. In the situation of both massive and massless directions the low-energy dynamics is described by a hybrid model.

The (quantum) Kahler moduli space M_K

In the classical analysis the singular locus is real codimension one in $(\mathbb{C}^*)^k$. However, when quantizing the system some of the flat directions for the scalars ϕ_a get lifted by an effective potential $W_{\text{eff}}(\phi; t)$ and only a singular locus $S \subset (\mathbb{C}^*)^k$ of complex codimension one remains. The complexified Kahler moduli space of the low-energy theory is then

$$M_K = (\mathbb{C}^*)^k \times S$$

Classical moduli space:



Kähler moduli space:




Figure 1. The classical and quantum moduli space of one-parameter models.

For $k = 1$ the moduli space is depicted in Fig. 1. The singular locus is a point at

$$(5) \quad S = \left(\frac{Y^N}{e^t = \prod_{i=1}^N Q_i^{Q_i}} \right) :$$

For the higher dimensional moduli spaces it suffices to note that for large values of r the singular locus between two adjacent phases is determined by the unbroken subgroup $U(1)_?$. A symptotically, it is $S = (C)_{\text{wall}}^{k-1} S_?$ $(C)^k$, where $S_?$ is given by (5) with respect to the Kähler parameter and the charges of the unbroken gauge group $U(1)_?$. At the boundary between two adjacent phases the singularity therefore reduces effectively to the one-dimensional situation.

R-symmetries

For the sake of completeness let us briefly note that a necessary condition to obtain a superconformal theory in the infra-red is the invariance of the gauged linear sigma model under an axial and a vector $U(1)$ R-symmetry, cf. for instance [48]. The former is ensured by requiring the conformal condition (or Calabi-Yau condition)

$$(6) \quad \sum_{i=1}^N Q_i^a = 0; \quad \text{for } a = 1, \dots, k :$$

We will henceforth impose this condition.

If no superpotential is present, we assign vector R-charge zero to all supermultiplets, which then turns into the standard R-charge assignment for the non-linear sigma model in the infra-red. If a superpotential is present, some of the chiral multiplets have to carry non-vanishing vector R-charge and the global symmetry is ensured by

$$(7) \quad W(\mathbf{x}) = {}^2W(\mathbf{x});$$

where $\mathbf{x} = (\mathbb{R}^1 x_1; \dots; \mathbb{R}^N x_N)$ for some phase \mathbb{R} . We shall henceforth assume an integrality condition on the R-charges of the fields in the linear sigma model, i.e. the R-charge is equal modulo 2 to the fermion number, $(-1)^F = (-1)^{R_i}$.

Some interesting examples

Example 1

Let us consider the gauged linear sigma models with the following chiral multiplets:

$$(8) \quad \begin{array}{|c|c|c|c|c|} \hline & x_1 & \dots & x_N & p \\ \hline U(1) & 1 & \dots & 1 & N \\ \hline \end{array}$$

The deleted sets at $r = 0$ resp. $r = 0$ are

$$(9) \quad \text{fp} = 0g \quad \text{and} \quad \text{+} = \text{fx}_1 = \dots = \text{fx}_N = 0g;$$

and the corresponding toric varieties in the infra-red are the orbifold $X_- = \mathbb{C}^N / \mathbb{Z}_N$ and its crepant resolution X_+ , which is the total space of the line bundle $\mathcal{O}(-N) \rightarrow \mathbb{C}P^{N-1}$.

Let us turn on a superpotential $W(p; \mathbf{x}) = pG(\mathbf{x})$ with a homogeneous degree N polynomial $G(\mathbf{x})$. A frequent choice is the Fermat type polynomial,

$$G(\mathbf{x}) = x_1^N + \dots + x_N^N;$$

We assign R-charge $+2$ to p and 0 to all other fields. In the small volume limit the theory becomes a Landau-Ginzburg model with potential $G(\mathbf{x})$ on the orbifold X_- . At large volume we obtain a Landau-Ginzburg model over X_+ , whose potential however induces F-term masses. The low-energy theory therefore localizes at $\text{fp} = G(\mathbf{x}) = 0g$ and becomes a non-linear sigma model on a degree N hypersurface in projective space $\mathbb{C}P^{N-1}$.




Figure 2. The classical Kahler moduli space of Example 2 without theta angle directions of the two-parameter model.

Example 2

A frequently considered two-parameter model is given by the following fields and charges:

(10)

	x_1	x_2	x_3	x_4	x_5	x_6	p
$U(1)_1$	0	0	1	1	1	1	4
$U(1)_2$	1	1	0	0	0	2	0

Its classical phase diagram together with the deleted sets is shown in Fig. 2. Phase III contains the orbifold C^5/Z_8 , and phase I its smooth total resolution. Phases II and IV are partial resolutions, the former being a line bundle over weighted projective space, $O(-8) \times W P_{11222}^8$.

Let us turn on the superpotential $W(p; x) = pG(x)$ with a homogeneous polynomial $G(x)$ of bidegree $(4; 0)$, for example,

$$G(x) = x_6^4(x_1^8 + x_2^8) + x_3^4 + x_4^4 + x_5^4.$$

We assign R-charge + 2 to p and 0 to all other fields. In phase III this results in a Landau-Ginzburg model over the orbifold C^5/Z_8 and in phase IV in a LG model over the toric variety X_{IV} . Phases I and II are geometric in view of massive F-terms. In particular, phase II corresponds to a degree 8 hypersurface in $W P_{11222}^8$ and phase I to a smooth Calabi-Yau hypersurface.

2.1 D-branes from the ultra-violet to the infra-red

Let us consider boundary conditions that preserve B-type supersymmetry $N = 2_B$. The latter is characterized by the unbroken vector R-symmetry.

As usual in supersymmetric theories the variation of the bulk action gives rise to total derivatives and thus to boundary terms. The strategy in [20] was to introduce appropriate boundary counter terms prior to imposing boundary conditions. In fact, the supersymmetry variations of the bulk kinetic terms can be compensated by standard boundary terms that are equal for all D-branes. We are not interested in these and instead concentrate on the part that specifies the D-brane data.

Let us first consider the situation without superpotential. The modification to include W will turn out to be only minor from the ultra-violet perspective of the gauged linear sigma model.

D-branes in models without superpotential

A D-brane in the gauged linear sigma model is described by an $N = 2_B$ invariant Wilson line at the boundary of the world sheet,

$$(11) \quad P \exp i \oint_{\partial} ds A :$$

It carries a representation $\rho(g)$ of the gauge group T as well as a representation $R(\cdot)$ of the vector R-symmetry and a representation of the world sheet fermion number. In view of the integrality condition on the R-charges we may set $\rho = R(e^i)$.

The simplest choice for the Wilson line corresponds to an irreducible representation of the gauge group, $\rho(g) = g_1^{q_1^1} \dots g_k^{q_k^k}$, i.e.

$$(12) \quad A = \sum_s v_s \operatorname{Re}(\cdot) = \sum_{a=1}^{X^k} q^a (v_a)_s \operatorname{Re}(a) :$$

We call it a Wilson line brane and denote it by $W(q) = W(q^1; \dots; q^k)$. The representation of the R-symmetry is $R(\cdot) = \cdot^j$ for some integer j , and $\rho = (\cdot^{-1})^j$. We refer to a Wilson line brane with even and odd j as brane resp. antibrane.

The general D-brane B can be constructed by piling up a stack of Wilson line branes, $W = \prod_{i=1}^n W(q_i)$,⁴ and turning on a supersymmetric interaction,

⁴By abuse of notation we sometimes refer to W as the Chan-Paton space of the D-brane.

ie. a tachyon profile Q , among the individual components. The corresponding superconnection reads

$$(13) \quad A = v_s \text{Re}(\) + \frac{1}{2} f Q ; Q^y g \quad \frac{1}{2} \sum_i^X \partial_i Q + \frac{1}{2} \sum_i^X \partial_i Q^y ;$$

where ∂_i is the $N = 2_B$ superpartner of the chiral field x_i .

The Wilson line (11) is supersymmetric if and only if the tachyon profile $Q(x)$ depends holomorphically on the chiral fields x_i and squares to zero. Also, $Q(x)$ has to respect the representation of the gauge group,

$$(14) \quad (g)^{-1} Q(g \cdot x) (g) = Q(x):$$

In view of the R-symmetry representation the stack W splits up into components of definite R-degree, $W = \sum_j W^j$, and from A we find that $Q(x)$ has to carry R-charge one,

$$(15) \quad R(\) Q(\) R(\)^{-1} = Q(x):$$

This implies in particular that $Q(x)$ is odd,

$$(16) \quad Q(x) = -Q(x);$$

and therefore the interaction $Q(x)$ in the superconnection couples branes to antibranes only. Moreover, having R-charge one implies that the tachyon profile can be brought into the block-form

$$Q(x) = \begin{matrix} & 0 & & & & & & 1 \\ & 0 & d^{j_{ax}-1} & 0 & \dots & 0 & 0 & \\ & 0 & 0 & d^{j_{ax}-2} & \dots & 0 & 0 & \\ & \vdots & & \ddots & \ddots & \vdots & & \\ & 0 & 0 & 0 & \dots & 0 & d^{j_{in}} & \\ & 0 & 0 & 0 & \dots & 0 & 0 & \end{matrix} \begin{matrix} \\ \\ \\ \\ \\ \\ \\ \end{matrix} :$$

Each non-trivial map $d^j : W^j \rightarrow W^{j+1}$ increases the R-degree by one. The data for the D-brane, $B = (W ; (g); R(\) ; Q(x))$, can therefore conveniently be encoded in a complex of Wilson line branes,

$$(17) \quad \dots \xrightarrow{d^{j-2}} W^{j-1} \xrightarrow{d^{j-1}} W^j \xrightarrow{d^j} W^{j+1} \xrightarrow{d^{j+1}} \dots ;$$

where $W^j = \sum_{i=1}^{n_j} W^j(q_i^j)$. In explicit examples we will often drop the R-degree index and use the convention to underline the component of R-degree $j = 0$. We denote the set of D-branes in a gauged linear sigma model without superpotential by $D(C^N ; T)$.

RG-flow and D-isomorphisms

Let us study the RG-flow of the Wilson line (13) to the infra-red while staying deep inside of one of the phases in the Kähler moduli space. The discussion here will be independent of F-terms and is applicable to both complexes and matrix factorizations. In particular, we do not yet integrate out fields with F-term masses that constrain the low-energy dynamics to a holomorphic subvariety in X_r , i.e. in a model with superpotential $W(x)$ we consider the low-energy theory as a Landau-Ginzburg model over X_r in any phase.

As the gauge coupling constants are massive parameters in two dimensions they will blow up as the theory flows to the infra-red and as a consequence the equations of motion for the gauge multiplets become algebraic. In particular, integrating out the gauge fields v_a and the scalars ϕ_a shows that the superconnection (12) becomes the supersymmetric pullback of a connection A to the world sheet,

$$A = x^i A_{ij} dx^j - \frac{i}{2} F_{ij} \sigma_i \sigma_j ;$$

A is the connection of the holomorphic line bundle $\mathcal{O}(q) = \mathcal{O}(q_1; \dots; q_k)$ on the toric variety X_r , and F is its field strength. The charges q_a now determine the divisor class, or more physically, the world volume flux on the D-brane. The complex (17) then turns into a complex of holomorphic vector bundles over X_r , and the matrix factorization (20) couples together line bundles over the base space X_r of the LG-model.

In the following we are particularly interested in the interplay of the boundary RG-flow and the bulk D-term equations (2). Instead of considering the RG-flow explicitly we identify deformations of the Wilson line (13) that do not alter the infra-red fixed point. These deformations lead to equivalence relations between D-branes, called D-isomorphisms in [20]. The low-energy D-branes can then be defined as equivalence classes in the gauged linear sigma model. D-isomorphisms are composed of the following two kinds of manipulations.

(i) The first manipulation can be seen by noticing that the superconnection (13) contains a matrix valued boundary potential $f_Q; Q^Y g$.

Suppose a D-brane is reducible, $B = B_1 \cup B_2$, with tachyon profile

$$(21) \quad Q(x) = \begin{pmatrix} Q_1(x) & 0 \\ 0 & Q_2(x) \end{pmatrix} ;$$

the reader to compute the corresponding boundary potentials and to verify that table (25) is not altered.

(ii) For the second manipulation the essential idea is that renormalization group flow can change the boundary action by boundary D-term $s, Q Q^Y (:::)$, but not by boundary F-term $s, Q (:::)$. Here, Q and Q^Y are the $N = 2_B$ supercharges. The theory flows to an infra-red fixed point with a particular D-term, irrespective of the chosen D-term in the gauged linear sigma model, i.e. deforming boundary D-terms does not alter the infra-red D-brane.

In order to describe these D-term deformations it is convenient to consider the supersymmetry generator on the world sheet boundary from the Noether procedure. In the zero mode approximation it becomes

$$(26) \quad iQ = \int (\partial_t + iA_t) + Q :$$

and reduces to Quillen's superconnection [55,57]. It can be used to express the superconnection A of the low-energy theory as

$$(27) \quad A = x A + \frac{1}{2} f Q ; Q^Y g :$$

Quillen's superconnection in (26) is written in the unitary frame for the associated graded holomorphic vector bundle E with hermitian metric. In what follows it is more convenient to work in the holomorphic frame, for which $iQ^{hol} = \int (\partial_t + Q(x))$.

A D-term deformation $M = M(x; x; \cdot)$ in the holomorphic frame is then a transformation $(Q^0)^{hol} = M Q^{hol} M^{-1}$, or equivalently

$$(28) \quad Q^0(x) = M \int (\partial_t + M^{-1} + M Q(x) M^{-1}) :$$

We assume that M commutes with the representations of the gauge group and the global symmetries. In particular,

$$(29) \quad \begin{aligned} Q^0(g) &= g M^{-1} (g) M^{-1} ; \\ R^0(\cdot) &= M R(\cdot) M^{-1} ; \end{aligned}$$

In the special situation when M depends only on the chiral fields x_i it is simply a similarity transformation

$$(30) \quad Q^0(x) = M Q(x) M^{-1} ;$$

i.e. a change of the holomorphic frame of E .

An important example of the general transformation (28) is as follows. Consider

$$Q_s(x) = \begin{pmatrix} Q_1(x) & s(x) \\ 0 & Q_2(x) \end{pmatrix}$$

with $Q_1 + Q_2 = 0$, and assume that for $s = 0$ the D-brane becomes the reducible D-brane (21) with B_2 empty.

As long as we keep s non-zero we can deform $Q_s(x)$ by a similarity transformation (30),

$$Q_s(x) = M Q_{s^0}(x) M^{-1}; \quad \text{for } M = \begin{pmatrix} s & \text{id} & 0 \\ 0 & s^0 & \text{id} \end{pmatrix} :$$

However, setting $s = 0$ by a similarity transformation is not possible. Let us consider the general D-term deformation (28) in infinitesimal form $M = \text{id} + s(x; \cdot)$,

$$(31) \quad \frac{\partial}{\partial s} Q_s(x) \Big|_{s=0} = \text{tr} \left(Q_0^{\text{hol}}; (x; \cdot) \right) :$$

Inserting the D-brane under consideration on the left-hand side we obtain

$$\frac{\partial}{\partial s} Q_s(x) \Big|_{s=0} = \begin{pmatrix} 0 & (x) \\ 0 & 0 \end{pmatrix} :$$

The existence of a D-term deformation to set $s = 0$ therefore reduces to the requirement that $\text{tr} \left(Q_0^{\text{hol}} \right)$ is Q_0^{hol} -exact. To see that this is indeed true we merely remark that since $\text{tr} \left(Q_0^{\text{hol}} \right)$ is Q_0^{hol} -closed it corresponds to a state between the D-branes B_1 and B_2 . However, since the open string spectrum between any D-brane and an empty one is empty, it follows that $\text{tr} \left(Q_0^{\text{hol}} \right)$ must be Q_0^{hol} -exact. See [20] for a more detailed discussion of this point.

Example 1 with $N = 3$ and no superpotential

Consider $B = B_1 \cup B_2$, defined in (23) and (24). In both the orbifold and the large volume phase the D-brane B is D-isomorphic to

$$(32) \quad B^0 : W(1) \xrightarrow{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}} W(0)^3 \xrightarrow{\begin{pmatrix} 0 & x_3 & x_2 \\ x_3 & 0 & x_1 \\ x_2 & x_1 & 0 \end{pmatrix}} \underline{W(1)^3} \xrightarrow{(px_1; px_2; px_3)} W(1) :$$

To show this we start with B and first use relation (31) to turn on a constant map from the Wilson line components $W(2)$ in B_2 to $W(2)$ in B_1 . Then we use a change of basis (30) to transform it to $B^0 = (\underline{W(2)} \oplus W(2)) = B^0$. This shows the equivalence of B and B^0 in the infra-red.

Again we can add backward arrows in B^0 to make it into a matrix factorization of $W(p; x) = pG(x)$. Then B and B^0 are still D-isomorphic.

For later applications it turns out to be more convenient to reformulate the two manipulations from above in terms of quasi-isomorphisms on the set of linear sigma model D-branes (or the underlying category) [58]. Indeed, D-isomorphisms are nothing else but quasi-isomorphisms [20].

Recall that a quasi-isomorphism U between two D-branes B_1 and B_2 is a Q -closed map, i.e. $Q_2 U = U Q_1$, such that its cone,


$$(33) \quad Q_{C(U)} = \begin{pmatrix} Q_1 & 0 \\ U & Q_2 \end{pmatrix};$$

is empty. The following manipulations show that quasi-isomorphic D-branes, B_1 and B_2 , are indeed related by a chain of brane-antibrane annihilations and D-term deformations [20]:

$$Q_1 = \begin{pmatrix} 0 & 1 \\ Q_1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ Q_2 & \text{id } A \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ U & Q_2 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ Q_2 & \text{id } A \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ U & Q_2 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ Q_2 & 0 \end{pmatrix} = Q_2$$

In the first and last step we used brane-antibrane annihilation (22), in the second a similarity transformation (30) to turn on U , and in the third an infinitesimal D-term deformation (31) to turn on id .

Having introduced D-isomorphisms we can define now the set of low-energy D-branes on a toric variety X_r as D-isomorphism classes of linear sigma model branes. Let us denote the set of low-energy D-branes by $D(X_r)$ and $M F_W(X_r)$. We obtain the following two pyramids of maps, where the vertical maps correspond to modding out by D-isomorphisms in the respective phase:



Return to orbifolds or LG-orbifolds

In order to close the discussion of D-isomorphisms, let us briefly consider their role in the special case when the phase in M_K corresponds to an orbifold $X_r = C_S^{N-k}$. It occurs if the deleted set consists of k irreducible factors, $r = \prod_{l=1}^k f_l$, where $f_l \in \mathbb{Z}[x_1, \dots, x_N]$ contains k elements. The fields x_1 for $l=1, \dots, k$ get vacuum expectation values, say $x_1 = 1$, which break T to G . For D-branes the representation $\rho(g)$ of T then descends to a representation $\rho(g)$ of G .

How does this affect the D-isomorphisms? In view of the deleted set \mathcal{R} the empty D-branes are given by

$$W(q, Q_1) \xrightarrow{x_1} W(q) \quad \text{for } 1 \leq i :$$

After assigning expectation values this descends to the trivial complex,

$$O(q) \xrightarrow{1} O(q) :$$

where $q = q \bmod Q_1$ is now a representation of \mathcal{R} . Since the only empty D-branes in the orbifold model are given by such trivial complexes, we find that any quasi-isomorphism is a similarity transformation as in (30) and thus invertible, i.e. there are no non-trivial quasi-isomorphisms anymore. A similar argument holds for matrix factorizations in LG-orbifolds as well.

RG-flow and F-term masses

Before we turn to the question of how to relate the sets of low-energy D-branes across phase boundaries, let us consider another issue that is specific to models with a superpotential and thus to matrix factorizations. As elucidated above the superpotential can give rise to masses for some of the chiral multiplets, which then must be integrated out in the strict infra-red limit. As an example consider the superpotential $W(p; x) = pG(x)$ which gives masses to p and to the transverse mode of the hypersurface $fG(x) = 0$ at large volume. Let $N = (N^1; \dots; N^k)$ be the gauge charge of p .

The effect of the massive modes on matrix factorizations was studied in [20], cf. also [59{63]. Indeed, a bre-wise version of Korrer periodicity [64] implements the equivalence of the set of matrix factorizations in $M_{F_W}(X_{\mathcal{R}})$ and the set of complexes (of coherent sheaves) $D(M_{\mathcal{R}})$ on the hypersurface $M_{\mathcal{R}} = \{p = G(x) = 0\} \subset X_{\mathcal{R}}$.

Take a matrix factorization given by the data $(W; (g); R(\cdot); Q(p; x))$. Let j_m be the minimal R-degree in the representations $R(\cdot)$. The matrix factorization is mapped to the D-brane on $M_{\mathcal{R}}$ as follows. First impose $G(x) = 0$, which implies $Q(p; x)^2 = 0$. Second consider a Wilson line component $W(q)$ as a graded module $A(q)$, where $A = C[p; x]/(G)$. Here $C[p; x]$ is the graded coordinate ring of the linear sigma model and taking the quotient by the ideal (G) corresponds to imposing $G(x) = 0$. Now consider $A(q)$ as an infinite module over the ring $B = C[x]/(G)$, that is $A(q) = \sum_{m=0}^{\infty} p^m B(q+mN)$, i.e. every Wilson line component $W(q)$ becomes an infinite stack of line bundles on $M_{\mathcal{R}}$,

$$\sum_{m=0}^{\infty} O_{M_{\mathcal{R}}}(q+mN)[2m] :$$

At large volume the Konnerth map acts on them in the following way. B_1 becomes

$$\begin{array}{ccccccc}
 O(-1) & \xrightarrow{\frac{\mathcal{X}}{\mathcal{X}}} & O(0) & \xrightarrow[3]{\frac{\mathcal{X}}{\mathcal{X} \frac{\mathcal{X}^2}{\mathcal{X}}}} & O(1) & \xrightarrow[3]{\frac{\mathcal{X}}{\mathcal{X} \frac{\mathcal{X}^2}{\mathcal{X}}}} & O(2) \\
 & & & & & & \xrightarrow[3]{\frac{\mathcal{X}}{\mathcal{X} \frac{\mathcal{X}^2}{\mathcal{X}}}} & O(3) & \xrightarrow[3]{\frac{\mathcal{X}}{\mathcal{X} \frac{\mathcal{X}^2}{\mathcal{X}}}} & O(4) & \xrightarrow[3]{\frac{\mathcal{X}}{\mathcal{X} \frac{\mathcal{X}^2}{\mathcal{X}}}} & \dots \\
 & & & & & & & & & & & & O(5) & \xrightarrow{\frac{\mathcal{X}}{\mathcal{X}}} & \dots
 \end{array}$$

where the line bundles $O(q)$ are understood to be pulled back from CP^2 to the elliptic curve E . In fact, this complex is an empty D-brane, in accordance with table (25).

On the other hand, B_2 is mapped to

$$O(-1) \xrightarrow{0} O(2) \xrightarrow{1} O(2) \xrightarrow{0} O(5) \xrightarrow{1} O(5) \xrightarrow{0} \dots$$

Trivial brane-antibrane pairs can be dropped in the infra-red, and the single line bundle $O(-1)[1]$ remains.

2.2 Moving around in moduli space

So far we considered the renormalization group flow to the infra-red only deep inside of the phases in the Kähler moduli space. We defined the set of low-energy D-branes in the infra-red theory as the set of D-isomorphism classes of D-branes in the linear sigma model. Let us now turn to the question of how to transport low-energy D-branes across phase boundaries between adjacent phases.

Grade restriction rule

The analysis of the gauged linear sigma model on the cylinder, corresponding to a propagating closed string in the infra-red, shows that along the singular locus $S = M_K$ the world sheet description breaks down. Additional non-normalizable modes show up, which are due to noncompact directions in field space, i.e. the effective potential $W_{\text{eff}}(\phi; t)$ for large values of the scalar fields ϕ^a has flat directions along S .

In the presence of D-branes, the analysis of the effective potential was redone in [20] on a strip of width L and led to the following result. For the gauge group $T = U(1)$ a Wilson line component $W(q)$ in a D-brane B causes




Figure 3. Windows for the grade restriction rule. Here S is odd.

no singularity near the phase boundary if and only if the grade restriction rule,

$$(35) \quad \frac{S}{2} < \frac{P}{2} + q < \frac{S}{2};$$

is satisfied. Here, $S = 1 + 2 \sum_{i \in \mathcal{D}} Q_i$.

The grade restriction rule is illustrated in Fig. 3. Take a path through a window w between singular points in the FI-theta-plane. Then the condition (35) admits only a set N^w of S consecutive charges for the Wilson line components $W(q)$ of the D-brane B . Denote the set of grade restricted complexes and matrix factorizations by T^w resp. $M_{F_W}(T^w)$.

For higher rank gauge groups, $T = U(1)^k$, only the unbroken gauge group $U(1)_?$ at the respective phase boundary enters in the condition (35).

Combining D-isomorphisms and the grade restriction rule

As it stands the grade restriction rule is a condition on the D-branes in the gauged linear sigma model, i.e. $T^w = D(C^N; T)$ and $M_{F_W}(T^w) = M_{F_W}(C^N; T)$. For the low-energy D-branes the important observation is the fact that the grade restriction rule is a unique 'gauge choice' in the D-isomorphism class (for $k > 1$ unique up to D-isomorphisms that are common to both phases, see [20]). A low-energy D-brane can therefore be transported, say from phase I to phase II, by first imposing the grade restriction rule on the D-isomorphism class in phase I and then mapping the grade restricted representative to its D-isomorphism class in phase II.

In fact, the compositions of maps, $\pi_{II} \circ \rho_{I,II}$ and $\rho_{I,II} \circ \pi_{II}$ in the following diagrams are inverse to each other:

$$\begin{array}{ccc} & D(C^N; T) & \\ & \downarrow & \\ D(X_I) & \xrightarrow[\rho_{I,II}]{} & T^w \xrightarrow[\rho_{II,I}]{} D(X_{II}); \end{array}$$

$$\begin{array}{ccc} & MF_W(C^N; T) & \\ & \downarrow & \\ MF_W(X_I) & \xrightarrow[\rho_{I,II}]{} & MF_W(T^w) \xrightarrow[\rho_{II,I}]{} MF_W(X_{II}); \end{array}$$

Here, ρ denotes modding out by D -isomorphisms in the respective phase, and ρ is picking the representative of the D -isomorphism class in the grade restricted set. Let us illustrate this in our example.

Example 1 with $N = 3$ and no superpotential

Here, $S = 3$. We pick a window $w = f < < g$, which admits $N^w = f < 0 < g$ according to the grade restriction rule. At $r \ll 0$ we found the D -isomorphic D -branes, $B_1 = B = B^0$, and at $r \gg 0$ we found $B_2 = B = B^0$. From its definition (32) it follows that in both phases the D -brane B^0 is the grade restricted representative in the D -isomorphism class and can thus be transported across the phase boundary:

$r \ll 0$	GRR	$r \gg 0$
$B_1 = B^0$	B^0	$B_2 = B^0$

3 Orientifolds in linear sigma models

In this section we put aside D -branes and review and study world sheet parity actions in gauged linear sigma models without boundary. Let us pick the cylinder $\Sigma = \mathbb{R} \times S^1$ with coordinates $(t; x) \sim (t; x + 2\pi)$ as world sheet. The parity action is an orientation reversal $(t; x) \mapsto (t; -x)$ dressed by an involution of the target space coordinates and possibly dressed by a sign on the left-moving Ramond sector states, $(-1)^{F_L}$.

In theories preserving $N = 2_B$ supersymmetry the action of the orientation reversal needs to be extended to the $N = (2; 2)$ superspace coordinates by $(\theta; \bar{\theta}) \mapsto (\theta; \bar{\theta})$. This in particular implies that (anti)chiral multiplets are mapped to (anti)chiral multiplets, twisted chiral multiplets are mapped to twisted antichiral multiplets, and vector multiplets to vector

multiplets [13]. Also, the involution is holomorphic and acts non-trivially on the chiral superfields of the gauged linear sigma model.

In the following we will have a closer look at the fixed point locus of the holomorphic involution and at the orientifold Kahler moduli space.

3.1 The holomorphic involution and orientifold planes

Recall the bulk Lagrangian of $N = (2;2)$ gauged linear sigma models,

$$(36) \quad \mathcal{L} = \int d^4x \sum_{a=1}^k \frac{1}{2e_a^2} \text{Tr} \left(\sum_{i=1}^N X_i e^{Q_i} \sum_{i=1}^N X_i \right) + \text{Re} \int d^2e \sum_{a=1}^k t^a + \text{Re} \int d^2 W(X) :$$

The kinetic terms in the first line are invariant under the orientation reversal by itself, but we can dress the latter by the holomorphic involution,

$$(37) \quad \begin{aligned} X_i &\mapsto !_i X_{(i)} && \text{for } i = 1; \dots; N; \\ V_a &\mapsto V_a && \text{for } a = 1; \dots; k; \end{aligned}$$

The permutation is of order two, $!^2 = \text{id}$, and preserves the gauge charges, $Q_i = Q_{(i)}$. Invariance of the action requires the coefficients $!_i$ to be phase factors. We obtain the parity action on the component fields by combining the right-hand side of (37) with the transformation of the supercoordinates with respect to $!$. The result is

$$(38) \quad \begin{aligned} (\mathbb{X}; \mathbb{Y}; \mathbb{F}_i) &\mapsto (!_i X^{(i)}; !_i Y^{(i)}; !_i F^{(i)}); \\ (v; \mathbb{Y}; \mathbb{D}) &\mapsto (v; \mathbb{Y}; \mathbb{D}); \end{aligned}$$

The holomorphic involution needs to be involutive only up to gauge transformations,

$$(39) \quad !^2 X_i = g_i X_i;$$

that is $!_i !_{(i)} = g_i$, where $g_i = g_1^{Q_i^1} \dots g_k^{Q_i^k}$ is the character of the representation determined by the charges Q_i .

In view of (39) we see that only gauge equivalence classes of holomorphic involutions, g_i , matter. Note that we can always find a representative g_0 of the class so that $g_0^2 = 1$. It is however not unique. There are in fact 2^k gauge equivalent choices. If we take a reference involution g_0 , we can dress it by an element $g = (g_1; \dots; g_k) \in T$ for $a = 1$. The resulting involution,

$$(40) \quad g_0 = g_i ;$$

still satisfies the property $(\rho_0)^2 = 1$.

In the presence of a superpotential the phase factors in the B-type parity (37) are further constrained by

$$(41) \quad W(x) := W(\rho(x)) = W(x):$$

The minus sign on the right-hand side compensates the sign from $\rho: d^+ d^- \rightarrow d^+ d^-$ in the action (36). For a model with Fermat polynomial, $G(x) = \prod_{i=1}^N x_i^{k_i+2}$, and $W(p;x) = pG(x)$, this is ensured by

$$(42) \quad \rho_i^{k_i+2} = \rho_i^{-1} \quad \text{for } i = 1; \dots; N:$$

Here, ρ_i is an arbitrary phase factor. In particular, in ρ_0 the phase ρ_i must be a sign.

Orientifold planes

Orientifold planes are the irreducible components of the fixed point locus $\text{Fix}^T(\rho)$ of the holomorphic involution ρ . The fixed point locus can be determined by finding solutions to $\rho x_i = x_i$ for appropriate elements $\rho \in T$. Using the gauge symmetry it can be expressed in terms of special gauge choices ρ_0 , that is

$$(43) \quad \text{Fix}^T(\rho) = \bigcup_{\rho_0 \in T} \text{Fix}(\rho_0);$$

where $\text{Fix}(\rho_0) = \prod_{i=1}^N \{x_i = \rho_0 x_i\} = (C^*)^k$. In (43) the union is taken over a discrete subset of elements in the gauge group, $I \subset T$. In particular, any $\rho_0 = (\rho_1; \dots; \rho_k)$ with $\rho_a = 1$ is an element in I .

Note that in the low-energy conformation the deleted set ρ_r is removed in view of the D-term equations. Therefore, depending on the phase some of the components $\text{Fix}(\rho_0)$ of the fixed point locus may be removed in the infra-red. In models without superpotential the irreducible components of the fixed point locus, that is the orientifold planes, are then given by

$$(44) \quad O = \text{Fix}(\rho_0) \quad \rho_r = (C^*)^k:$$

In models with superpotential there are geometric phases where the low-energy conformation localizes on a holomorphic subvariety $M_r = \text{f} \in W = 0$. The orientifold planes are then given by

$$(45) \quad O = \text{Fix}(\rho_0) \setminus M_r \quad \rho_r = (C^*)^k:$$

As we will observe explicitly in examples later on, the intersection in (45) may be reducible (or even empty) and splits into a finite number of irreducible components O_i :

$$O = \sum_{i=1}^r O_i ;$$

Example 1 with $N = 3$

Orientifolds in local CP^2

Let us consider the allowed parity actions in Example 1 without superpotential. We can work in coordinates that diagonalize the holomorphic involution,

$$(\sigma_1 \sigma_2 \sigma_3 | p)(x_1; x_2; x_3; p) = (\sigma_1 x_1; \sigma_2 x_2; \sigma_3 x_3; p) ;$$

We find the following distinct choices of parity actions with corresponding orientifold planes O_i :

Involutions O_i	$Fix^T(\sigma)$	orbifold pt	large volume
$(\sigma_1 \sigma_2 \sigma_3)$	$O_{+1} = \text{space filling}$	$C^3 = Z_3$	O_9 on $O_{CP^2}(3)$
$(\sigma_1 \sigma_2; \sigma_3)$	$O_{+1} = fx_3 = 0g$ $O_{-1} = fx_1 = x_2 = p = 0g$	2 plane = Z_3	O_7 on $O_{CP^1}(3)$ O_3 on pt
$(\sigma_1; \sigma_2; \sigma_3)$	$O_{+1} = fx_2 = x_3 = 0g$ $O_{-1} = fx_1 = p = 0g$	line = Z_3	O_5 on C O_5 on CP^1
$(\sigma_1; \sigma_2; \sigma_3)$	$O_{+1} = fx_1 = x_2 = x_3 = 0g$ $O_{-1} = fp = 0g$	Z_3 fixed pt	O_7 on CP^2

Orientifolds on the elliptic curve

Let us next turn on a superpotential $W(p; x) = pG(x)$ with Fermat polynomial

$$G(x) = x_1^3 + x_2^3 + x_3^3 ;$$

In view of this superpotential we have to consider, besides the diagonal involution $(\sigma_1 \sigma_2 \sigma_3 | p)$, the additional involutions,

$$(\sigma_1 \sigma_2 | \sigma_3 | p)(x_1; x_2; x_3; p) = (\sigma_1 x_2; \sigma_2 x_1; \sigma_3 x_3; p) ;$$

where two coordinates are exchanged. However, we need to satisfy condition (42), which rules out some of the involutions that we considered in the non-compact situation. There are in fact only two independent involutions,




Figure 4. The slices of the moduli space in the presence of an orientifold. One slice connects the large volume and the ‘small’ volume point (sometimes Gepner point).

which are related by T-duality:

Involution σ	Fix ^T ()	Gepner point	large volume
$(-1; 1; 1; 1)$	$O_{+1} = fx_1 = x_2 = x_3 = 0g$ $O_{-1} = fp = 0g$	fixed pt of Z_3	E
$(-1; 1; 1; 1)$	$O_{+1} = fx_1 + x_2 = x_3 = 0g$ $O_{-1} = fx_1 - x_2 = p = 0g$	line = Z_3	1 pt E 3 pts E

In the table, $E \subset \mathbb{C}P^2$ denotes the elliptic curve at large volume. The locus O_{-1} for the second involution is reducible at large volume and consists of three points,

$$O_{-1} = \{fx_1 - x_2 = x_3 = p = 0g \subset E\};$$

where $3 = 2$.

3.2 Orientifolds and their constrained moduli space

So far we considered the parity action on the kinetic terms and the chiral superpotential in (36). As for the effect on the twisted chiral superpotential, t^a , we note that the orientation reversal maps the gauge field strength f_a to $-f_a$. Invariance of the path integral therefore requires $t^a = t^a \pmod{2\pi}$, where the $\pmod{2\pi}$ shift is due to the topological term for the theta angle. The theta angle is therefore restricted to

$$(46) \quad t^a = 0 \text{ or } \pi \pmod{2\pi};$$

According to these conditions the orientifold constrains the allowed complexified Kähler moduli to 2^k real half-dimensional slices in M_K . Note that the orientifold slices may or may not intersect the singular locus S .




Figure 5. The slices $= (; 0)$ resp. $= (0; 0)$ divide the moduli space in distinct sectors.

One-parameter models

For $k = 1$ the slices in the Kähler moduli space are depicted in Fig. 4. The singularity S sits at the theta angle $= S \pmod{2\pi}$, where $S = \sum_{Q_i > 0} Q_i$. The slice at $= S$ is therefore divided by S into two disconnected components. Otherwise, on the slice $\notin S$ the singularity is avoided and the slice connects large and small volume limits. We will find later, after a detailed investigation of D-branes in the orientifold background, that under special circumstances the two slices may be joined at the small volume point.

Higher dimensional moduli spaces

For $k > 1$ the slices in moduli space are complicated but more interesting. Let us illustrate this in the two-parameter model Example 2.

Example 2

The singular locus S is the union of the following two loci [65]:

$$S_0 = fe^{-t^2} = 1 - 4(1 - 2^8 e^{t^2})^2 g; \quad \text{and} \quad S_1 = fe^{-t^2} = 1 - 4g;$$

The orientifold action admits four slices. Two of them, $= (0;)$ and $= (;)$, do not intersect the singular locus S so that we can move freely between the four phases. On the other hand, the intersections of the slices $= (; 0)$ resp. $= (0; 0)$ with S are depicted in Fig. 5. In the former slice we can move from phases I to IV. In the latter all phases are separated, even more, there is a non-perturbative regime that is not accessible from any of the four perturbative regions.

3.3 Orientifolds at orbifold points

At an orbifold point of M_K the gauge group T of the linear sigma model is broken to a discrete subgroup in view of vacuum expectation values $x_{1 \in I} = 1$. Let us examine the consequences for the holomorphic involution.

The vacuum expectation values require special gauge choices for σ , namely the ones that act trivially on the fields x_1 for $1 \in I$, so that a \mathbb{Z}_2 -equivalence class of parity actions, $\sigma \rightarrow \sigma \tau$ for $\tau \in \mathbb{Z}_2$, remains and acts on the orbifold C^N/k . The orientifold group and the orbifold group are combined in an extension [66],

$$1 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_2 \rightarrow 1$$

Notice that after breaking the gauge group to the discrete group it is in general not possible to find a representative for the holomorphic involution that satisfies $\sigma^2 = \text{id}$.

4 D-branes in the presence of orientifolds

In this section we study the parity actions on world sheets with boundary. We will combine the considerations of the previous two sections to define parity-invariant, low-energy D-branes. The presentation will follow the discussion in [17].

4.1 The world sheet parity action on D-branes

Let us pick the strip $\Sigma = \mathbb{R} \times [0; \pi]$ with coordinates $(t; x)$ as world sheet. The orientation reversal acts as $\sigma : (t; x) \mapsto (t; \pi - x)$. Let us study the effect of the world sheet parity $P^m := (-1)^{F_L}$ on the general Wilson line (13).

Recall that the boundary action on the strip for a single Wilson line brane $W(q^a)$ is given by

$$(47) \quad \int_{a=1}^k \frac{1}{2} + q^a \int_0^\pi dt (v_a)_t \text{Re}(a)_{x_1=0} - (v_a)_t \text{Re}(a)_{x_1=\pi} :$$

Here we included the contribution from the theta angle. The orientation reversal exchanges the two boundaries of the strip giving rise to an overall sign in front of (47) and, therefore, inverting the sign of $a=2 + q^a$, i.e. its

effect on the charges is

$$(48) \quad P^m : q^a \mapsto q^a \quad (a = +, -) :$$

Notice that this map is well-defined because of the integral values of q^a that we found earlier in (46).

Let us now consider a general Wilson line with tachyon profile $Q(x)$. For convenience we set

$$A[Q] = \frac{1}{2} f Q ; Q^Y g + \frac{1}{2} \sum_{i=1}^X i @_i Q + \frac{1}{2} \sum_{i=1}^X i @_i Q^Y :$$

The path ordered Wilson line on the strip is

$$(49) \quad \text{Str}_{j_1(x)} P e^{i \int_1^0 dt A[Q]_{k=0}} \text{Str}_{j_2(x)} P e^{i \int_1^0 dt A[Q]_{k=0}} ;$$

where the time t^0 on the boundary $x = 0$ is oppositely oriented to the time t on the strip itself. Here, the fields $j_1(x)$ and $j_2(x)$ take values in $\text{End}(W)$ and correspond to incoming and outgoing string states at minus and plus infinite time.

The orientation reversal swaps the two boundaries, and the involution acts on the connection as $A[Q] = A[Q(x)]$, resulting in

$$\begin{aligned} \text{Str}_{j_1} P e^{i \int_1^0 dt A[Q]_{k=0}} \text{Str}_{j_2} P e^{i \int_1^0 dt A[Q]_{k=0}} &= \\ \text{Str}_{j_1^T} P e^{i \int_1^0 dt A[Q]_{k=0}^T} \text{Str}_{j_2^T} P e^{i \int_1^0 dt A[Q]_{k=0}^T} &: \end{aligned}$$

In the second line we applied the graded transpose and its properties as summarized in the appendix A. The appearance of the transposition in the parity action tells us that the Chan-Paton vector space W is mapped to its dual vector spaces,

$$P^m : W \rightarrow W^* :$$

In order to extract the parity transform of Q we rewrite the transposed superconnection as

$$\begin{aligned} A[Q]^T &= \frac{1}{2} f Q^T ; (Q^Y)^T g + \frac{1}{2} i @_i Q^T + \frac{1}{2} i @_i (Q^Y)^T = \\ &= \frac{1}{2} f Q^T ; (Q^T)^Y g + \frac{1}{2} i @_i Q^T + \frac{1}{2} i @_i (Q^T)^Y = \\ &= A[Q^T] ; \end{aligned}$$

Here the hermitian conjugate of an endomorphism on the dual space W^* is defined by $(M^T)^Y = (M^Y)^T$. Comparing with (49) we can easily read off the action of the world sheet parity P^m on the tachyon profile,

$$(50) \quad P^m : Q(x) \mapsto Q(x)^T ;$$

and on $\mathfrak{e}l\mathfrak{d}s$,

$$(51) \quad P^m : (x) \mathfrak{T} \quad (x)^T :$$

Let us consider the generalization of the parity transform of the charges in equation (48) to the stack of Wilson line branes W . The representation (g) of the gauge group T on W is determined by the charges of its Wilson line components, so that (48) turns into

$$(52) \quad P^m : (g) \mathfrak{T} \quad = (g) (g)^T :$$

The graded transpose appears by the same reasoning as above and is consistent with relation (14).

In order to determine the effect on the representation $R(\)$ of the vector R -symmetry we take the graded transpose of relation (15) and compare it with (50). We find that the representation has to transform as $R(\) \mathfrak{T} R(\)^T$. Note however that the world sheet parity action P^m contains the operator $(-1)^{F_L}$, which, for m odd, inverts the sign of left-moving Ramond sector states and in particular the sign of Ramond $\mathfrak{e}l\mathfrak{d}s$. Since D -branes are sources for these $\mathfrak{e}l\mathfrak{d}s$, the overall sign of the D -brane charges is also flipped, i.e. branes are mapped to antibranes if m is odd. In the context of D -branes the operator $(-1)^{F_L}$ is called the anti-brane operator (or anti-brane functor on the D -brane category [17,19]). Since the components W^j with R -degree j even and odd correspond to branes and antibranes respectively, we conclude that the operator $(-1)^{F_L}$ induces the shift $[m] : j \mathfrak{T} m + j$. The parity action P^m on the representation $R(\)$ is therefore dressed by a character,

$$(53) \quad P^m : R(\) \mathfrak{T} \quad \chi_m(\) R(\)^T :$$

The relation $\chi_m = R(-1)$ implies furthermore

$$(54) \quad P^m : \mathfrak{T} \quad (-1)^m \mathfrak{T} :$$

Let us summarize our findings of this section. The world sheet parity action acts on a D -brane $B = (W ; (g); R(\); Q(x))$ in the linear sigma model as

$$(55) \quad P^m : W ; ; R ; Q \mathfrak{T} \quad W ; \quad = (g)^T ; \chi_m(\) R^T ; \quad Q^T :$$

We sometimes use the abbreviation $P^m(B)$ for the parity image of B .

4.2 Dressing by quasi-isomorphisms

A well-defined parity operator on D -branes should square to the identity, so that we can gauge it in order to obtain an orientifold background. However,

P^m does not square to the identity, rather it acts as

$$(P^m)^2 : Q(x) \mapsto (-1)^{m+1} (-2) Q(x) (-2)^{-1} :$$

Recall that -2 is an element of the gauge group and therefore the dressing by the representation (-2) arises from applying the gauge invariance condition (14) on $Q(-2x)$. The sign $(-1)^{m+1}$ is due to the graded double transpose (118) in the appendix.

The non-involutive property of P^m is cured in conjunction with our wish to describe low-energy D-branes as D-isomorphism classes in the gauge linear sigma model. We supplement the D-brane data by an arbitrary quasi-isomorphism $B = (W_2; (g); R; Q(x); U(x))$, and define a dressed parity operator P^m as follows:

$$(56) \quad \begin{aligned} P^m(Q(x))U &= U(-Q(x))^T; \\ P^m((g))U &= U(g)U(g)^T; \\ P^m(R)U &= R U R^T; \\ P^m(U) &= (-1)^m U^T. \end{aligned}$$

By abuse of notation we abbreviate these transformations by $P^m(B)U = U P^m(B)$. Note that quasi-isomorphisms in the "inverse" direction are also possible, that is $V P^m(B) = P^m(B)V$. A homomorphism taking values in $\text{Hom}(W_2; W_1)$ transforms as

$$(57) \quad P^m(U)_1 = U_2^T :$$

In order to ensure that the parity operator P^m depends only on the gauge equivalence class of the holomorphic involution, g , the quasi-isomorphism U must transform as

$$(58) \quad U_g := U(g)^T :$$

Inserting (58) in the definition of the parity operator P^m one can easily check that the latter does not depend on the gauge choice.

Two orientifold actions

The definition of the dressed parity operator P^m does not yet ensure that it squares to the identity. We may however utilize the dressing by the quasi-isomorphism U to ensure this property by an appropriate transformation behaviour of the quasi-isomorphism itself, i.e. we determine $P^m(U)$ so that $(P^m)^2 = \text{id}$.

Let us pick a homomorphism taking values in $\text{Hom}(W_2; W_1)$ and apply the parity operator (57) twice. With the Ansatz $P^m(U) = A U^T$ for

some constant invertible matrix A we have

$$\begin{aligned} P^m \left(P^m(U_2) \right) &= A_1 U_1^T P^m(U_2)^T = \\ &= A_1 (\eta^2)^{-1} U_2^T = \\ &= A_1^{-1} (\eta^2)^{m+1} U_2^{m+1} (\eta^2)^{-1} U_2^T \end{aligned}$$

Requiring equality with the original field determines A up to a constant, so that U has to transform as

$$(59) \quad P^m(U) = (\eta^2)^{-1} U^T :$$

A quasi-isomorphism in the inverse direction, i.e. $V_2 P^m(U) = V_1^T$, transforms as

$$(60) \quad P^m(V) = V^T (\eta^2)^{m+1} :$$

Here the inverse constant appears for consistency with the case when the quasi-isomorphism is invertible, $U = V^{-1}$.

The constant c is associated with the parity operator. It has to be the same for all mutually compatible D -branes. In order to stress this we henceforth denote the parity operator on D -branes by P^m . By abuse of notation we continue to denote the set of D -branes, now supplemented with U , by $D(C^N; T)$ or $MF_W(C^N; T)$.

Like the quasi-isomorphism, the constant c depends on the gauge choice of η . Comparing the transformation (59) for U and U_g and requiring $P^m(U_g) = P^m(U) P^m(\eta)^T$ as well as (58) reveals that

$$(61) \quad \eta_g = \eta (\eta^2)^{-1} :$$

The combined shift, $(\eta^a; q^a) \mapsto (\eta^{a+2n^a}; q^{a-n^a})$ for $n^a \in \mathbb{Z}^k$, alters the constant c as follows:

$$(62) \quad \eta^{2n^a} (\eta^2)^{-1} :$$

What we have considered so far ensures that P^m squares to the identity on the D -brane data $(W; R; Q)$. However, since the quasi-isomorphism transforms now under the parity operator we have to impose $(P^m)^2 = \text{id}$ on U as well. In fact, using (56) and (59) we obtain

$$\begin{aligned} U &= P^m(P^m(U)) = \\ &= P^m((\eta^2)^{-1} P^m(U)^T) = \\ &= P^m((\eta^2)^{-1} P^m(U)^{m+1} (\eta^2)^{-1} U^T) = \\ &= (\eta^2)^{-2} (\eta^2)^{-1} U^T : \end{aligned}$$

The constant c is therefore determined up to a sign,

$$c = \pm 1 ; \text{ with } c = 1 :$$

The constant c depends on ϵ and η . We will refer to the constant c as orientifold sign, although it is strictly speaking not a sign. In the following we will often pick an involution σ that squares exactly to the identity, which implies that $\sigma^2 = 1$. Also the combined shift of the theta angles and the gauge charges resulting in (62) does not alter the sign ϵ .

In summary, P^m is a parity operator on the set of D-branes in the gauged linear sigma model. It squares to the identity operator and is determined by the discrete theta angle θ , by the dressing with the antibrane operator, m odd or even, and by the orientifold sign ϵ whose role will be elucidated in a moment.

Shift of R-degree

As known in D-brane categories an overall shift of the R-degree, $[1] : j \mapsto j+1$, is unphysical because all measurable quantities depend on the difference of R-degrees [67]. In view of the interpretation of the Z_2 -grading $(-1)^j$ as distinguishing branes from antibranes the shift $[1]$ is indeed the antibrane operator as it appeared already in the previous subsection.

Let us study the effect of the antibrane operator on the quasi-isomorphism U and on the sign ϵ . First, the parity action commutes with the shift only if the latter is accompanied by $m \mapsto m+2$, i.e. $P^{m+2} [1] = [1] P^m$.

The R-symmetry representation and the Z_2 -grading operator are mapped as follows,

$$(63) \quad \begin{aligned} [1] : R(\epsilon) &\mapsto \epsilon R(\epsilon); \\ [1] : \mathbb{T}(-1)^j &\mapsto \mathbb{T}(-1)^{j+1}. \end{aligned}$$

In view of the sign change of the Z_2 -grading operator the graded transpose is altered to

$$[1] : M^T \mapsto (\epsilon^T)^{-1} M^T (\epsilon^T)^{-1}.$$

The transformation property (56) of $Q(x)$ then tells us that the quasi-isomorphism U is mapped as

$$[1] : U \mapsto \epsilon^{-1} U.$$

Inserting (63) in the defining equation (59) of the orientifold sign ϵ we observe that it is mapped as

$$(64) \quad \mathbb{T}(-1)^j \mapsto \mathbb{T}(-1)^{j+1}.$$

Alltogether we have shown that the shift $[1]$ transforms the parity operator as follows,

$$(65) \quad P^{(m+2)} \epsilon^{-1} [1] = [1] P^m \epsilon^{-1}.$$

Notice that the combination $(-1)^{m-2}$ is an invariant under the shift of R -degree. Here, the square bracket denotes taking the next lower integer. We therefore expect that physical quantities depend on this invariant combination.

4.3 Parity invariant D-branes

As a next step we define parity-invariant low-energy D-branes to be D-isomorphism classes that are preserved by the parity operator P^m , i.e. there exists a quasi-isomorphism U or V from $B = (W; (g); R(\cdot); Q(x))$ to its world sheet parity image $P^m(B)$, so that $P^m(B) = B$. Explicitly,

$$\begin{aligned}
 Q(x)U &= U Q(x)^T; \\
 (g)U &= (g)gU (g)^T; \\
 (66) \quad R(\cdot)U &= R(\cdot)U R(\cdot)^T; \\
 U &= (-1)^m U^T; \\
 U &= (-1)^{m+1} U^T.
 \end{aligned}$$

Analogous relations hold for quasi-isomorphisms V . Note that the quasi-isomorphism is now determined by the invariance conditions (66), whereas in the previous subsection it was completely arbitrary. We denote the sets of invariant low-energy D-branes in gauged linear sigma models without and with superpotential by $D^m; (X_r)$ and $M F_W^m; (X_r)$, respectively.

Let us point out a subtlety here. In general, a D-brane B is invariant if it can be related to its world sheet parity image $P^m(B)$ by a chain of quasi-isomorphisms. However, if both types of quasi-isomorphisms, U and V , appear in the chain it is not clear how to determine the sign. We will later find an elegant resolution to this problem.

4.4 Gauge groups on D-branes and the type of orientifold planes

All D-brane in an orientifold background have to carry the same orientifold sign. In this section we want to consider the situation when the D-brane is a stack of identical D-branes B_i , that is the Chan-Paton space is the tensor product of an external Chan-Paton space $V_e = C^n$ and the internal Chan-Paton space W_i . The orientifold sign can then be distributed appropriately on the two contributions,

$$= e_i :$$

Stacks of irreducible invariant D-branes

Let us consider a stack of irreducible and invariant D-brane, i.e. the tachyon profile $Q_i(x)$ on the internal Chan-Paton space W_i is an irreducible endomorphism. The internal sign ϵ_i , which is now associated with the irreducible D-brane (not the orientifold), is defined through

$$(67) \quad U_i = \epsilon_i \epsilon_i^{-1} \epsilon_i^{m+1} U_i^T :$$

The stack of D-branes on the Chan-Paton space $W = V_e \oplus W_i$ is defined as follows:

$$\begin{aligned} Q(x) &= \text{id} \oplus Q_i(x) ; \\ (g) &= \text{id} \oplus \epsilon_i(g) ; \\ R(\cdot) &= \text{id} \oplus R_i(\cdot) ; \\ U &= U_e \oplus U_i : \end{aligned}$$

In particular, the quasi-isomorphism splits into an external isomorphism U_e on V_e and an internal quasi-isomorphism U_i .

Comparing the last relation of (66) for U and relation (67) for U_i we find that the external sign ϵ_e enters in the symmetry condition of the external isomorphism s ,

$$U_e = \epsilon_e (U_e)^t :$$

It therefore determines the gauge group on the stack of D-branes [6,17],

$$(68) \quad SO(n); n \in 2\mathbb{Z} \quad \text{for} \quad \epsilon_e = +1 ;$$

$$(69) \quad Sp(n=2); n \in 2\mathbb{Z} \quad \text{for} \quad \epsilon_e = -1 :$$

In the following we will mainly work with the internal, irreducible part of a D-brane. We drop the index i for convenience, with the exception of ϵ_i .

The type of orientifold planes

Given a parity action with multiple components of the fixed point locus in the infra-red theory, we may consider a probe D-brane that sits on top of one of the components. According to the gauge group of the probe D-brane, $\epsilon_e = +1$ or $\epsilon_e = -1$, we follow the general convention in the literature to define the type of the orientifold plane by $o := \epsilon_e$. Indicating the type we refer to the orientifold plane as O^o -plane. We will have to say more on the type of orientifold planes in later sections.

Stacks of brane image-brane pairs

A special class of invariant D-branes is given by brane image-brane pairs, i.e. D-branes that are of the form irreducible brane plus parity image brane.

In particular, the internal Chan-Paton space reads $W_i = V \oplus V$. If we consider a stack of such branes we tensor $V \oplus V$ with the external Chan-Paton space V_e , which is equivalent to considering

$$W = V_e \oplus V \oplus V \oplus V ;$$

and setting

$$\begin{aligned} Q(x) &= \text{id} \oplus Q_i(x) \oplus \text{id} \oplus (Q_i(x)^T) ; \\ (g) &= \text{id} \oplus_i(g) \oplus \text{id} \oplus (g)_i(g)^T ; \\ R(\cdot) &= \text{id} \oplus R_i(\cdot) \oplus \text{id} \oplus_m(\cdot) R_i(\cdot)^T ; \end{aligned}$$

The quasi-isomorphism for this D-brane can be written in the brane image-brane basis as

$$(70) \quad U = \begin{pmatrix} 0 & \propto U_e^t & m+1 & (-2)^{-1} \\ U_e & \text{id} & 0 & \end{pmatrix} :$$

where c is an a priori arbitrary constant and c was introduced for later convenience. Let us try to match the sign of the parity action by computing

$$U U^T m+1 (-2) = \begin{pmatrix} \propto \text{id} & \text{id} & 0 \\ 0 & c^{-1} c \text{id} & \text{id} \end{pmatrix} = c \text{id} :$$

So we find that we can always adjust the constant c to be equal to the sign. In particular, this means that the brane image-brane pairs appear for both orientifold signs, and since there is no symmetry restriction on the isomorphism U_e the gauge group is $U(n)$.

4.5 Moving between phases | Orientifolds and the grade restriction rule

In view of the observations of Sec. 3.2 the transport of low-energy D-branes between phases in the gauged linear sigma model is not always possible in the presence of orientifolds, at least not within the world sheet description. The reason is the singular locus S , which is real codimension one on the orientifold slices in M_K .

Let us concentrate on linear sigma models with gauge group $T = U(1)$ in the subsequent discussion.

Avoiding the singularity

If the slice in Kahler moduli space does not intersect with the singular locus, that is $\notin S \pmod{2}$, we can apply the grade restriction rule of [20] to

transport D-branes between the phase. It reads

$$\frac{S}{2} < \frac{1}{2} + q < \frac{S}{2} :$$

In fact, the world sheet parity action P^m , mapping $(=2 + q) \mathbb{T}^2$ ($=2 + q$), preserves the grade restriction rule. Once we have found a grade restricted representative for a D-brane, its image under the world sheet parity action P^m is again grade restricted. This is important for consistency of the gauged linear sigma model near the phase boundary.

From [20] we recall that there are no non-trivial D-isomorphisms between D-branes in the grade restricted set. The grade restricted D-branes are unique up to a basis change (30) of the Chan-Paton space W . Since P^m does not map out of the grade restricted set this implies that a grade restricted invariant D-brane B and its parity image $P^m(B)$ must be related by an (invertible) basis change U . It is therefore convenient to work with the grade restricted representative of a given D-isomorphism class. In particular, since U is an isomorphism it is easy to compute the sign ϵ_i and the problems with chains of quasi-isomorphisms that we mentioned in Sec. 4.3 do not show up.

Colliding with the singularity

If we consider a slice in M_K that collides with the singularity, we cannot transport D-branes from one phase to the other. We want to add a remark on the sign ϵ_i though.

Suppose we sit on the slice $\Sigma = S$. The windows that are adjacent to the singularity at S , cf. Fig. 3, admit the charges $N^W = f0;:::;S$ lg resp. $N^{W^+} = f1;:::;Sg$. If we pick a representative B for the D-brane that is grade restricted with respect to N^W , its world sheet parity image $P^m(B)$ will be grade restricted with respect to N^{W^+} . In order to map $P^m(B)$ back to B the associated quasi-isomorphism has to remove all the Wilson line components $W(S)$. This can be achieved by a chain of quasi-isomorphisms, all of which are of the same type, either U or V . In fact, these can then be composed to a single quasi-isomorphism, which allows to compute the sign ϵ_i .

Example 1 with $N = 2$ and no superpotential

Let us consider a simple example of an invariant D-brane at large volume. We take the world sheet parity action with $\sigma = \text{id}$ and $m = 1$ and pick the

slice $x_2 = 0$ that collides with the singularity S . The adjacent windows at the phase boundary determine $N^W = f_1; 0g$ and $N^{W^+} = f_0; 1g$.

Regard the D-brane

$$B : \underline{W(-1)} \xrightarrow{x_2} W(0); \quad Q = \begin{pmatrix} 0 & x_2 \\ 0 & 0 \end{pmatrix};$$

which is an element of T^W . It is mapped by the world sheet parity to

$$P_0^1(B) : \underline{W(0)} \xrightarrow{x_2} W(1); \quad Q^T = \begin{pmatrix} 0 & 0 \\ x_2 & 0 \end{pmatrix};$$

In view of the different gauge charge assignments the latter is clearly not isomorphic to the original complex. However, at large volume, $r \gg 0$, we can bind to it an empty D-brane via a D-term deformation (31),

$$\begin{array}{ccc} \underline{W(0)} & \xrightarrow{x_2} & W(1) \\ \text{\scriptsize @ } \begin{matrix} 0 \\ 1 \end{matrix} & & \text{\scriptsize @ } \begin{matrix} 1 \\ 0 \end{matrix} \\ & \text{\scriptsize R} & \text{\scriptsize R} \\ \underline{W(-1)} & \xrightarrow{x_2} & W(0) \oplus \xrightarrow{(x_1, x_2)} W(1) \end{array};$$

After eliminating trivial pairs $W(-q) \oplus W(q)$ for $q = 0, 1$, we get back the original D-brane. The associated quasi-isomorphism is

$$V = \begin{pmatrix} 0 & x_1 \\ x_1 & 0 \end{pmatrix}; \quad \text{with } \epsilon_i = 1;$$

4.6 Orientifolding complexes and matrix factorizations

Let us formulate invariant D-branes in models without superpotential in terms of complexes (17). This will facilitate some of the subsequent, explicit computations in examples. In the low-energy interpretation the following makes contact with the discussion of orientifold projections in the derived category of coherent sheaves in [19].

Similarly, invariant matrix factorizations are described by merely adding "backward arrows" in the complexes, as in (20). This is straightforward, and we will skip the general discussion of matrix factorizations here.

The defining conditions (66) for an invariant D-brane can be rewritten for a complex (17) in terms of a commutative diagram ,

$$(71) \quad \begin{array}{ccccccc} \cdots & \xrightarrow{(d^{m-j})^T} & W^j & \xrightarrow{(d^{m-j-1})^T} & W^{j+1} & \xrightarrow{(d^{m-j-2})^T} & \cdots \\ & & \left| \begin{array}{c} u^j \\ ? \end{array} \right. & & \left| \begin{array}{c} u^{j+1} \\ ? \end{array} \right. & & : \\ \cdots & \xrightarrow{d^{j-1}} & W^j & \xrightarrow{d^j} & W^{j+1} & \xrightarrow{d^{j+1}} & \cdots \end{array}$$

The second line is the complex for $Q(x)$, and the first line represents the world sheet parity in age $P^m(Q(x)) = Q(x)^T$. Note that we have $(d^j)^T = (-1)^j (d^j)^t$ according to the definition of the graded transpose, where t is the ordinary transposition of matrices. $W^j = {}_i W^j(q_i^j)$ is the component with R-degree j , and

$$W^{m-j} = W^j = {}_i W^j \quad (= \quad {}_i W^j)$$

is its dual, now carrying R-degree $m-j$.

The chain maps, $u^j : W^j \rightarrow W^{j+1}$, preserve the global symmetries as well as the gauge charges. They are the components of the quasi-isomorphism

$$U = \begin{array}{ccccccc} & & & & & & 1 \\ & & & & & & \vdots \\ & & & & & & C \\ & & & & & & \vdots \\ & & & & & & A \\ & & & & & & \vdots \\ & & & & & & 0 \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} \quad \begin{array}{c} \vdots \\ u^j \\ \vdots \end{array} :$$

The last condition in (66) becomes

$$(72) \quad u^j = (-1)^{j-1} (-1)^{(m+1)j} (u^{m-j})^t :$$

Koszul complexes

As examples for invariant D-branes we consider coherent sheaves $\mathcal{O}_C(q)$ that are localized at the common zero locus C of n polynomials (f_1, \dots, f_n) . They can be described via tachyon condensation [58,68{70] by Koszul complexes, which we define below. We determine the chain maps u^j that render the Koszul complexes invariant and provide a simple formula for the type of gauge group that is supported on them.

Let us denote the gauge charges of the polynomials by $(Q_{f_1}, \dots, Q_{f_n})$ and introduce $Q_f = \prod_{i=1}^n Q_{f_i}$. The Wilson line (11) associated with the Koszul complex can be realized in terms of boundary fermions ψ_i for $i = 1, \dots, n$, which upon quantization satisfy the Clifford algebra relations $\psi_i \psi_j = -\psi_j \psi_i$.

The associated Fock space, built on the Fock vacuum defined by $\psi_i = 0$, is then the Chan-Paton space W . The boundary interaction is given by

$$Q(x) = \prod_{i=1}^{X^n} f_i(x)_i$$

and acts naturally on the Fock space W . Since $Q(x)$ needs to be gauge invariant, the boundary fermions ψ_i must carry the gauge charges Q_{f_i} . The resulting complex reads

$$(73) \quad C : W_{\frac{m+n}{2}}(q, Q_f) \xrightarrow{f} \cdots \prod_{i=1}^{M^n} W_{\frac{m+n}{2}-1}(q, Q_{f_i}) \xrightarrow{f} W_{\frac{m+n}{2}}(q) :$$

We assigned R-degree $(m+n)-2$ to the Fock vacuum ψ_i , i.e. to the rightmost entry in the complex. This assignment is necessary for an invariant D-brane. Note that it also requires

$$(74) \quad m = n \pmod{2}$$

In order to determine the chain maps u^j and the corresponding sign ϵ_j it is instructive to describe the Koszul complex C in the language of alternating (or exterior) algebras. Let $R = \prod_{i=1}^n C[x]_i$,⁵ where $C[x]$ is the graded coordinate ring of chiral fields. We introduce the interior product

$$(75) \quad \iota_v : \wedge^p R \rightarrow \wedge^{p-1} R ; \quad \forall v = (v_i)_{i=1}^p ;$$

where ι_v denotes a p -form in $\wedge^p R$ and $v = \sum_{i=1}^p v_i \psi_i$ a vector field in R . The tachyon profile $Q(x)$ in the complex C can then be realized as interior product ι_f ,

$$(76) \quad C : (\wedge^n R)_{\frac{m+n}{2}} \xrightarrow{f} \cdots \xrightarrow{f} (\wedge^1 R)_{\frac{m+n}{2}-1} \xrightarrow{f} (\wedge^0 R)_{\frac{m+n}{2}} :$$

For sake of brevity we did not indicate the gauge charges.

To calculate the parity image of this D-brane we need to determine the graded transpose of ι_f . The dual pairing of p -forms, $\wedge^p R$, with p -vectors, $\wedge^p R$, is

$$(77) \quad h(\psi_{i_1} \wedge \cdots \wedge \psi_{i_p}, \sum_{j_1, \dots, j_p} X_{j_1, \dots, j_p} \psi_{j_1} \wedge \cdots \wedge \psi_{j_p}) = \frac{1}{p!} \sum_{i_1, \dots, i_p} X_{i_1, \dots, i_p} \psi_{i_1} \wedge \cdots \wedge \psi_{i_p} ;$$

where ι is the natural generalization of the interior product to polyvectors. With the convention used in (77) it satisfies $\iota \circ \iota = \wedge$. Note that, according to the R-degree assignment in C , p -forms have R-degree $(m+n)-2-p$. Since the world sheet parity maps R-degrees as $j \mapsto n-j$, we assign the

⁵Henceforth, we drop the Fock vacuum ψ_i .

R -degree $(m+n)=2+p$ to p -vectors. The graded transpose f^T of the interior product is then defined by

$$h_f^T ; i_{p+1} := (-1)^{\frac{m+n}{2}+p+m} h ; f ; i_p ;$$

which is in accord with the definition (113) in the appendix. Inserting the right-hand side in the dual pairing (77) we readily find that

$$f^T : \wedge^p R \rightarrow \wedge^{p+1} R ; \quad \forall (-1)^{\frac{m+n}{2}+p} f \wedge ;$$

and the world sheet parity in the case of the boundary interaction $Q(x)^T$ is realized as $(-1)^{(m+n)=2+p} (f \wedge)$ on p -vectors.

Let us next construct the quasi-isomorphism U that makes C invariant. According to (71) we have to construct maps u^p such that the following diagram commutes,

$$(78) \quad \begin{array}{ccc} (\wedge^{n-p} R)_{\frac{m+n}{2}+p} & \xrightarrow{f} & (\wedge^{n-p-1} R)_{\frac{m+n}{2}+p+1} \\ u^p \uparrow & & \uparrow u^{p+1} \\ (\wedge^p R)_{\frac{m+n}{2}+p} & \xrightarrow{(-1)^{\frac{m+n}{2}+p} f \wedge} & (\wedge^{p+1} R)_{\frac{m+n}{2}+p+1} \end{array} ;$$

The idea is to choose a volume form $\omega \in \wedge^n R$ and try the Ansatz

$$(79) \quad u^p(\omega) = \mu_p f_0(x) \omega_{f^{-1}} ;$$

μ_p are constants to be determined below. ω_f is defined as the representation matrix of the holomorphic involution on the polynomials f_i , that is $\omega_f = (\omega_f)_{ij} f_j$ for $j = 1, \dots, n$. Its inverse is inserted in every contraction between ω and ω_f . The polynomial $f_0(x)$ must be such that U is a quasi-isomorphism, i.e. according to the definition of a quasi-isomorphism, around (33), the polynomials $(f_0; f_1; \dots; f_n)$ must give rise to an empty Koszul complex, or put in yet another way, the common zero locus of all polynomials must be contained in the deleted set Σ . Note also that in view of $P^m : q \rightarrow q$ the polynomial f_0 has to carry gauge charge

$$(80) \quad Q_{f_0} = -m + 2q - Q_f ;$$

The constants μ_p for $p = 0, \dots, n$ are fixed by inserting the Ansatz in the diagram (78) and requiring that it commutes. We obtain $\mu_{p+1} = (-1)^{(m+n)=2} \mu_p$. Using the freedom to normalize U , we set $\mu_0 = 1$ and obtain

$$\mu_p = (-1)^{\frac{m+n}{2}+1)p} ;$$

The internal sign ϵ_i of the D -brane is determined by constructing the graded transpose of u^p . Since the latter is even its graded transpose equals

its ordinary transpose,

$$h ; (u^p)^t ()_{ij} := h ; u^p ()_{in-p} ;$$

where $\mathbb{Z}^{\wedge p} \mathbb{R}$ and $\mathbb{Z}^{\wedge n-p} \mathbb{R}$. Inserting u^p on the right-hand side we end after some algebra,

$$(u^p)^t = \mathbb{1}_n (-1)^{p(m+1)} \det_{f-f}^2 u^p ;$$

where $(f^{-2})_{ij} = \varrho_{f_i} (-2)_{ij}$, which acts on a $(n-p)$ -form as $f^{-2} = \varrho (-2)_{p(-2)}$. Applying the holomorphic involution on both sides we obtain

$$u^{n-p} = \mathbb{1}_n (-1)^{p(m+1)} \frac{\varrho (-2)}{s_0 \det_{f-f}^0} p (-2) \mathbb{1} (u^p)^t ;$$

The factor s_0 is from $f_0 = s_0 f_0$ in u^{n-p} . Comparing with (72) we obtain the internal sign ϵ_i for the Koszul complex,

$$\epsilon_i = (-1)^{\frac{m-n}{2}} \frac{\varrho (-2)}{\det_{f-f}^0} ;$$

where we introduced $\det_{f-f}^0 = s_0 \det_{f-f}$. We have therefore succeeded in determining the gauge group for Koszul complexes, i.e.

$$(81) \quad \epsilon = \epsilon_i = (-1)^{(m-n)/2} \varrho (-2) \det_{f-f}^0 ;$$

Note that this result confirms the expectation that the gauge group does not depend on shifts of \mathbb{R} -degree [1], which exchanges branes and antibranes. Indeed, the external sign depends on the invariant combination $(-1)^{[m=2]}$, cf. relation (65). To see this note that in view of (74) we have $(m-n)/2 = [m=2] - [n=2]$.

Koszul-like matrix factorizations

Let us briefly comment on models with non-vanishing superpotential. A natural analog of Koszul complexes is provided by introducing additional polynomials $(g_1 ; \dots ; g_n)$ and defining a matrix factorization

$$Q(x) = \sum_{i=1}^{X^n} (f_i(x) \mathbb{1}_i + g_i(x) \mathbb{1}_i) ;$$

The condition $Q^2 = W$ is ensured by $W = \sum_i f_i g_i$.

On the level of complexes, we complete the factorization by including arrows "backwards"

$$(82) \quad C : (\wedge^n \mathbb{R})_{\frac{m-n}{2}} \xrightarrow{g} \dots \xrightarrow{g} (\wedge^1 \mathbb{R})_{\frac{m+n}{2}} \xrightarrow{1} \xrightarrow{g} (\wedge^0 \mathbb{R})_{\frac{m+n}{2}} ;$$

where $g = \sum_{i=1}^P g_i \omega_i$ is a 1-form in R . Obviously, $f_f; g \wedge g = W$ realizes the matrix factorization. The graded transpose can be found to be

$$(g^\wedge)^T = \left(1 \right)^{\frac{m+n}{2}+P} g :$$

The chain maps u^P and thus the sign ω_i turn out to be the same as for Koszul complexes.

4.7 Tensor products of invariant D-branes

Tensor products of complexes or matrix factorizations have been studied and used to construct special types of D-branes on many occasions [71{83]. In particular, all the known boundary states of Gepner models are realized as tensor products of simple matrix factorizations in the corresponding Landau-Ginzburg model.

Here we want to address the question of how the invariance of a D-brane under the orientifold action behaves under taking graded tensor products. The results of this subsection will be most important when we later study the bre-wise Korrer map that relates matrix factorizations of the linear sigma model to complexes of coherent sheaves at low energies.

Some properties of the graded tensor product are listed in appendix A. Let us briefly present its definition. For two endomorphisms of definite R -charge, $A \in \text{End}(W_1)$ and $B \in \text{End}(W_2)$, we define the graded tensor product as

$$A \otimes B = A \otimes_2^{\mathbb{A}^j} B ;$$

where we used the ordinary tensor product on the right-hand side. \mathbb{A}^j is the R -charge of A . In order to not forget subtle signs related to the insertion of $\otimes_2^{\mathbb{A}^j}$, which takes care of the grading, we will work explicitly with the ordinary tensor product.

Let us consider two invariant D-branes, $B_a = (W_a; \omega_a; R_a; Q_a; U_a)$ for $a = 1, 2$, satisfying (66) with $(\omega_a; \omega_a; m_a)$. We can form the tensor product brane $B = (W; \omega; R; Q; U)$ with $W = W_1 \otimes W_2$ and

$$\begin{aligned} (83) \quad Q &= Q_1 \otimes_2^{\omega_1} Q_2 ; \\ (g) &= \omega_1(g) \otimes_2^{\omega_2}(g) ; \\ R(\cdot) &= R_1(\cdot) \otimes_2^{\omega_2} R_2(\cdot) : \end{aligned}$$

For matrix factorizations the tensor product brane is associated with the sum of superpotentials $W = W_1 + W_2$.

These definitions together with (52) and (53) imply that

$$(84) \quad \begin{aligned} &= i_1 + i_2 ; \\ m &= m_1 + m_2 : \end{aligned}$$

A rather non-trivial question is to build the quasi-isomorphism U out of U_1 and U_2 , and in turn relate the sign i to i_1 and i_2 . U cannot just be the naive guess, that is $U_1 \otimes U_2$. To see this notice that upon using the graded transpose of tensor products, formula (119) in the appendix, the world sheet parity image of Q is

$$Q^T = Q_1^T \otimes_{\mathbb{Z}} \mathbb{Z} \otimes_{\mathbb{Z}} Q_2^T :$$

We find that $U_1 \otimes U_2$ cannot map Q^T back to Q , since it cannot turn the a 's into id_a 's and vice versa.

We need a more sophisticated quasi-isomorphism for the tensor product D-brane. To construct it we introduce the projection operators $p_a^r = 1/2(id_a + (-1)^f a)$ and note that they can be used to switch between a and id_a , i.e. $p_a^r a = a p_a^r = (-1)^f p_a^r id_a$. This suggests an Ansatz for U , which is a linear combination of four terms, $p_1^{r_1} U_1 \otimes p_2^{r_2} U_2$ with $r_a = 0;1$. Inserting the Ansatz in the invariance condition for Q in (66) it turns out that the quasi-isomorphism has to take the form

$$(85) \quad U = \sum_{r_1, r_2 = 0;1}^X (-1)^{(m_1 + r_1)r_2} p_1^{r_1} U_1 \otimes p_2^{r_2} U_2 :$$

U is furthermore compatible with all other equations in (66). In particular, the last one gives the simple sign relation

$$(86) \quad i = i_1 i_2 :$$

An application: The Koszul complex revisited

Let us reconsider the Koszul complexes of the previous subsection. Using the tensor product techniques we recompute the sign ϵ that determines the gauge group. Let us first assume that C is invariant with invertible quasi-isomorphism U_{inv} , which requires

$$(87) \quad \epsilon = q = q \otimes \mathbb{Z} :$$

Moreover, we work in a basis for the polynomials f_a that diagonalizes the action of the holomorphic involution, i.e. $f_a = s_a f_a$ for $a = 1; \dots; n$. Recall that an invariant D-brane requires $m = n \pmod{2}$.

We start by using the fact that a Koszul complex C of n polynomials $(f_1; \dots; f_n)$ is the tensor product of n Koszul complexes of a single polynomial,

$$C_a : W_{\frac{m_a-1}{2}}(Q_a - Q_{f_a}) \xrightarrow{f_a} W_{\frac{m_a+1}{2}}(Q_a) :$$

Each m_a has to be odd and we set $q_a := Q_{f_a} - 2Q_a$. By (84) the integers $(m_1; \dots; m_n)$ and the auxiliary theta angles $(\theta_1; \dots; \theta_n)$ must be chosen so that they sum up to

$$\begin{aligned} m &= m_1 + \dots + m_n ; \\ &= \theta_1 + \dots + \theta_n : \end{aligned}$$

Now let us compute the signs $i_{\mathcal{P}a}$ for the complexes C_a . The isomorphism U_a is given by the chain map

$$\begin{array}{ccc} W_{\frac{m_a-1}{2}}(Q_a - Q_{f_a}) & \xrightarrow{\binom{m_a-1}{2} s_a f_a} & W_{\frac{m_a+1}{2}}(Q_a) \\ \left| \begin{array}{c} 1 \\ ? \end{array} \right. & & \left| \begin{array}{c} \binom{m_a-1}{2} s_a^{-1} \\ ? \end{array} \right. \\ W_{\frac{m_a-1}{2}}(Q_a - Q_{f_a}) & \xrightarrow{f_a} & W_{\frac{m_a+1}{2}}(Q_a) \end{array}$$

Applying equation (72) we find that $i_{\mathcal{P}a} = \binom{m_a-1}{2} q_a \binom{2}{2} s_a^{-1}$. For the tensor product complex C we therefore have

$$i = \prod_{a=1}^n i_{\mathcal{P}a} = \binom{m-n}{2} q \binom{2}{2} \prod_{a=1}^n s_a^{-1} :$$

If the condition (87) for an invertible quasi-isomorphism U_{inv} is not satisfied, the Koszul complex may still be invariant provided that there exists a polynomial $f_0(x)$ so that $U = f_0 U_{inv}$ is a quasi-isomorphism, cf. equation (79). Setting $f_0 = s_0 f_0$ we find

$$i = \binom{m-n}{2} q \binom{2}{2} \prod_{a=0}^n s_a^{-1} :$$

In general, the polynomials f_a will not diagonalize \mathcal{F} . For an invariant D-brane we have $f_a = \sum_{b=0}^n f_{ab} f_b$ for $a = 0; 1; \dots; n$. The sign that determines the gauge group can then be written in terms of the determinant of \mathcal{F} ,

$$e = \text{sgn } i = \binom{m-n}{2} q \binom{2}{2} \det \mathcal{F}^0 :$$

5 Non-compact models

So far we discussed aspects of D-branes in gauged linear sigma models that are largely independent of the presence or absence of an F-term superpotential $W(x)$, the discussions included both, $Q(x)$ describing complexes and matrix factorizations. Let us now specialize to the case without superpotential and consider some examples of orientifolds and invariant D-branes described through complexes of Wilson line branes. As an application of the formula (81) we determine the type of orientifold planes by testing the gauge group of probe branes.

We are mainly interested in the dependence of the set of invariant D-branes and the orientifold planes on the slices of the Kähler moduli space. As a particular consequence of our linear sigma model approach we will find that the different slices may be connected along special loci in M_K . We investigate the phenomenon of type change of orientifold planes that was discussed in [14].

5.1 Orbifold phases and the orientifold moduli space

Let us consider linear sigma models with an orbifold phase and study the relation between the linear sigma model and the orbifold description. We illustrate the main points in Example 1 that becomes the quotient $C^N =$ with discrete group $= Z_N$ at the orbifold point. Recall the charges (8) and the moduli space in Fig. 4.

From the linear sigma model to the orbifold

As briefly reviewed in Sec. 3.3 the discrete group in the orbifold phase is due to a vacuum expectation value for the field p of gauge charge N . This expectation value also restricts the gauge equivalence class of holomorphic involutions to a p -equivalence class, for 2 , i.e. it requires $(p) =$
 $N p = p$.

By conveniently setting $p = 1$ an invariant D-brane in the orbifold theory is determined by the linear sigma model data through

$$\begin{aligned} Q(x) &:= Q(p=1;x); \\ (\) &:= (\) \quad \text{for } 2 \quad U(1); \\ R(\) &:= R(\); \\ U &:= U(p=1); \end{aligned}$$

A Wilson line component $W(q)$ becomes a \mathbb{Z}_N -equivariant line bundle $\mathcal{O}(q)$ with charge $q = q \bmod N$. We denote the \mathbb{Z}_N -equivariant Chan-Paton bundle descending from W by E .

It follows immediately that the orbifold data $(E; \mathbb{Z}_N; Q; U)$ of an invariant D-brane satisfies the invariance conditions (66), now with a representation χ of \mathbb{Z}_N , cf. [66].⁶ In particular, χ is a character of the orbifold group \mathbb{Z}_N , which implies that the theta angle is defined only modulo N at the orbifold point,

$$(88) \quad \theta = 2\pi \frac{q}{N} \bmod 2\pi :$$

This can be seen explicitly in the world sheet parity action on the charges,

$$q \mathbb{T} = q \bmod N :$$

From the orbifold to the linear sigma model

For the inverse map, lifting D-branes from the orbifold to the linear sigma model, we first have to decide to which slice of the Kähler moduli space we want to lift. For a given q we have a $\bmod N$ choice of theta angles in the linear sigma model. Let us pick one such choice.

The representation of the gauge group $U(1)$ is obtained by lifting the charges q in \mathbb{Z}_N to integers $q = q \bmod N$ in a fixed interval, say $0 \leq q < N$. Then the tachyon profile $Q(p; x)$ is constructed from $Q(x)$ by multiplying the entries in the latter by appropriate powers of p as to match the gauge charges determined by $U(1)$. The representation of the R-symmetry is simply $R(\mathbb{Z}_N) = R(\mathbb{Z}_N)$. Finally, the isomorphism $U(1)$ just lifts to $U(1)$ by lifting in appropriate powers of p .

The freedom of choosing the theta angle $\bmod N$ actually means that we can lift a D-brane from the orbifold point to different slices of the moduli space, i.e. a priori distinct slices of the Kähler moduli space are connected at the orbifold point. This can easily be picturized in the N -fold cover of the moduli space parametrized by the algebraic mirror coordinate z , defined by $e^{\pm i\theta} = (z/N)^{\pm 1}$. Since a shift $\mathbb{T} \rightarrow \mathbb{T} + N$ corresponds to a phase shift $\mathbb{T} \rightarrow e^i$, changing the slice means going straight through the orbifold point at $z = 0$. As depicted in Fig. 6 this leads to a qualitative difference for N odd and N even. In view of the combined shift $(\mathbb{T}; q) \mathbb{T} \rightarrow (\mathbb{T} + 2\pi; q - 1)$ we see that the slices $\theta = 0$ and $\theta = 2\pi$ are connected at the orbifold point for N odd, but remain disconnected for N even. So for N odd we have

⁶In [66] the representation χ for \mathbb{Z}_N and the isomorphism $U(1)$ are denoted by (g) for $g \in \mathbb{Z}_N$ and (χ) , respectively. In particular, the first two lines in their conditions (3.10) correspond to the second and to the last line in (66).




Figure 6. The N -fold cover of the Kähler moduli space, $e^t = (\mathbb{N})^N$. Relation (88) says that we can move straight through the orbifold point. For N odd (here 5) the two slices, at $\theta = 0$ and $\theta = \pi$, are connected at the orbifold point. For N even (here 6) the two slices, at $\theta = 0$ and $\theta = \pi$, remain disconnected.

two disconnected components of the orientifold moduli space, whereas for N even we have three.

When changing the slice at the orbifold point we have to be careful with relating the corresponding two sets of invariant D-branes properly. Let us pick an arbitrary invariant D-brane with quasi-isomorphism U in the orbifold phase. Recall that the sign ϵ_i is determined via

$$U = \epsilon_i^{m+1} (\epsilon^2)^{-1} U^T :$$

As we shift the theta angle to $\theta = \pi$ the parity action (48) on the gauge charges is changed to

$$q \mapsto -q :$$

Accordingly, in order to keep the D-brane invariant, we need to modify the quasi-isomorphism to $U^0 = pU$ with a new sign ϵ_i^0 . Using $p = \epsilon_p p$, we find

$$\epsilon_i^0 = \epsilon_p^{-1} \epsilon_i :$$

Since the gauge group on a stack of D-branes cannot be altered as we change the slice, the orientifold sign is modified in the same way,

$$(89) \quad \epsilon_i^0 = \epsilon_p^{-1} \epsilon_i :$$

Consequently, when we move straight through the orbifold point in Fig. 6 we have to take into account the orientifold sign change (89).

Higher-dimensional moduli spaces

In general, for gauge group $T = U(1)^k$ a necessary condition for a connection between different slices of the orientifold moduli space is that the deleted set X_r that determines X_r in the particular phase has at least one irreducible component of the form $\{x_1 = 0\}$ for some $1 \leq r \leq N$. The vacuum expectation value for x_1 then breaks the gauge group so that $q^a = \tilde{q}^a + Q_1^a$ and in particular,

$$(90) \quad q^a = \tilde{q}^a + Q_1^a \quad \text{and} \quad U^0 = U(1) :$$

Taking into account the change of quasi-isomorphism, $U^0 = x_1 U$, this shows an equivalence of the sets of invariant low-energy D-branes,

$$(91) \quad D^m(X_r) = D^0(X_r) :$$

For higher-dimensional moduli spaces this leads to the interesting phenomenon that large volume limits distinguished by different values of the theta angles may be connected through a path in Kähler moduli space. This will be illustrated later in the two-parameter model of Example 2.

Example 1

Let us consider the inequivalent spacetime involutions of Example 1. We can always choose coordinates so that the involution acts diagonally, $(x_1, \dots, x_N, p) = (x_1, \dots, x_N, p)$. For N odd we have

$$\cong (1, \dots, 1; \underbrace{1, \dots, 1}_N) ; \quad \text{for} \quad = 0, \dots, N :$$

In the orbifold phase the fixed point locus is an $(N-1)$ -dimensional plane through the fixed point $p = \{x_1 = \dots = x_N = 0\}$. For N even we have

$$\begin{aligned} &\cong (1, \dots, 1; \underbrace{1, \dots, 1}_N) ; \\ 0 &\cong (1, \dots, 1; \underbrace{1, \dots, 1}_{N/2}, 1, 1) ; \quad \text{for} \quad = 0, \dots, N=2 : \end{aligned}$$

The fixed point locus of the involution is a union of an $(N-1)$ -dimensional and an $(N-2)$ -dimensional plane in the orbifold phase, whereas the fixed point locus of U^0 is always the orbifold fixed point.

Fractional D-branes $O_p(q)$ on the orbifold $C^N = \mathbb{Z}_N$ are localized at p and carry \mathbb{Z}_N -charge q . They can be represented in the linear sigma model through the Koszul complexes (73) of N coordinate fields (x_1, \dots, x_N) . In the orientifold context fractional branes were studied before from different perspectives, see for instance [84, 87, 89]. Let us reexamine them from the linear sigma model point of view. In particular, we want to know which




Figure 7. Branes in an Z_3 orbifold and possible reflection planes, related by the Z_3 -symmetry

fractional branes are invariant for a given orientifold specified by $(\tau; m; \nu; \theta)$ or $(\tau; \theta)$.

From equation (74) we conclude that $m = N \bmod 2$, i.e. an invariant fractional brane must be dressed by $(-1)^F$ for N odd, whereas there is no dressing for N even.

In order to study the role of the theta angle, we note that the Z_N -representation q has to obey $2q = \nu \bmod N$ or formally

$$q = \frac{\nu + 2k}{2} \bmod N = 2 : \quad (75)$$

Recall that ν is defined $\bmod N$. For N even this has two solutions for q if $\nu/2$ is an integer and no solution if $\nu/2$ is half-integer. For N odd it always has only one solution. This has a nice pictorial representation in the quiver diagram corresponding to this orbifold. Here, the branes corresponding to irreducible representations of the orbifold group become red dots of the diagram, see Fig. 7 and 8. The N fundamental fractional branes are related by the quantum Z_N symmetry, i.e. 2π shifts of the theta angle at the orbifold point, which is depicted as a rotational symmetry in the corresponding diagram. Orientifolds are mirror-planes in these diagrams, respecting the symmetry.

It is now easy to see that for the case N odd there are N possible symmetry planes, each of them passing through exactly one point, as depicted in the figure for the case $N = 3$. The different orientifolds are related by rotational quantum symmetry. The corresponding invariant fractional brane can be lifted to both slices of the Kähler moduli space, see Fig. 6.

On the other hand, for N even there are two classes of orientifolds. The first class passes through precisely two points, leaving two of the fractional branes fixed, whereas the second class does not leave any point fixed, as shown in the figure for the case $N = 4$. The orientifold with two invariant fractional branes lies on the slice of M_K that collides with the singularity.




Figure 8. Branes in an Z_4 orbifold and possible inequivalent reflection planes: Either two or no branes are invariant under the parity.

The orientifold without invariant fractional branes is on the slice that is connected to the large volume point.

For the discussion of the gauge group let us distinguish between the two types of holomorphic involutions, σ and σ^0 . We use formula (81) to determine the gauge groups.

Since the involutions square to zero, the orientifold sign is indeed just a sign, $\epsilon = \pm 1$, and we can readily compute

$$\epsilon = (-1)^{\frac{m-N}{2}+}$$

For N even and on the slice $\theta = 0$ that collides with the singularity, the invariant fractional branes $O_p(0)$ and $O_p(N=2)$ therefore carry the same gauge group. For N odd the two slices in moduli space are connected at the orbifold point, which is reflected by the fact that ϵ does not depend on the theta angle.

For σ^0 (only N even) we use the representative $(1, \dots, 1; \dots, 1)$ for $N = 2$, which ensures $O_p = p$. However, $\epsilon = (-1)^{\frac{m-N}{2}}$ and the orientifold sign is actually a sign times a nontrivial constant, $\epsilon = (-1)^{\frac{m-N}{2}}$. Using this the external sign on the slice $\theta = 0$ becomes

$$(92) \quad \epsilon = \begin{cases} (-1)^{\frac{m-N}{2}+} & \text{for } O_p(0); \\ (-1)^{\frac{m-N}{2}} & \text{for } O_p(N=2): \end{cases}$$

The two invariant fractional branes carry opposite type of gauge group. These orientifolds appeared in the construction of six-dimensional RG fixed points from branes see for instance [87,88]. From a mirror perspective they have been discussed in [89].

5.2 The type of an orientifold plane

Let us use the results from Sec. 4.7 to compute the type \circ for the orientifold planes O as defined in (44). It will be convenient to work with the holomorphic involution $\sigma = \sigma_0$ that defines O through its fixed point locus. Furthermore, we use coordinates that diagonalize the involution. The probe brane that we will use to determine the type is given by the Koszul complex C of the coordinates for which $\sigma_0 x_i = \epsilon_i x_i$ with $\epsilon_i \in \{1, -1\}$. Setting these coordinates to zero gives the orientifold plane O . Let us denote them by $(f_1; \dots; f_d)$. d is the codimension of O .

We are already making several assumptions here. Indeed, a Koszul complex that corresponds to a D-brane that lies on top of O need not always exist. First the condition $m = d \pmod{2}$ has to be satisfied. Second a quasi-isomorphism U must exist for C in order to render it invariant. If it does not exist, it is sometimes possible to utilize a probe brane of higher codimension, that is $d + 2p$, which lies on O . Keeping in mind that the type of the gauge group alternates with p [6], we find that the type of the orientifold plane is given by

$$\circ = (-1)^{\sum \epsilon_i} \epsilon_e;$$

where ϵ_e is the external sign of the probe brane.

As we observed in the example (92), orientifold planes at orbifold singularities may lead to the effect that there exist two probe branes carrying opposite gauge group. The following result on the type of an orientifold plane can therefore be applied reliably only if we deal with a smooth orientifold geometry.

Under the above assumptions, the probe brane is a Koszul complex of the coordinates $f_1; \dots; f_d$, which are not invariant under σ_0 , and the σ_0 -invariant polynomials $f_{d+1}; \dots; f_{d+2p}$. The polynomial f_0 that enters the quasi-isomorphism (79) is σ_0 -invariant as well. We find

$$(93) \quad \circ = (-1)^{\sum \epsilon_i} \epsilon_e = \epsilon_0 (-1)^{(m-d)/2} \det(\epsilon) = \epsilon_{2q}(\epsilon)$$

where $\det(\epsilon) = \pm 1$ is the sign associated with the involution σ_0 ⁷, and q is the maximal charge in the Koszul complex (73). In the special situation that $\epsilon = (\epsilon_1; \dots; \epsilon_k)$ is given by signs $\epsilon_a = \pm 1$ we have $\det(\epsilon) = (-1)^{\sum \epsilon_a}$ and the formula for the type simplifies to

$$(94) \quad \circ = \epsilon_0 (-1)^{(m+d)/2} \epsilon_e(\epsilon) \quad \text{for } \epsilon_a = \pm 1$$

⁷It is a sign and independent of the gauge choice for ϵ only if the Calabi-Yau condition is satisfied.

We want to stress again that the result (93) requires the existence of a Koszul complex on \mathcal{O} . If such a Koszul complex fails to exist it is unclear which probe brane should be used to determine the type. Also, let us remark that the two probe branes (92) are reflected in formula (93) in a two-fold choice for the maximal charge q .

5.3 Type change in the orientifold moduli space

In this section we want to explore orientifolds and the dependence of their type on the slice in the Kahler moduli space. We will illustrate this point in a particular example. We will observe that the type of an orientifold plane is not an invariant concept and can change over the Kahler moduli space.

Example 2

Recall the list of chiral fields (10) and the moduli space for this example. Out of the list of possible target space involutions we consider:

	x_1	x_2	x_3	x_4	x_5	x_6	p
σ_0	1	+1	1	1	1	+1	1

The holomorphic involution acts diagonally with the indicated signs on the linear sigma model coordinates. For consistency with tadpole cancellation we pick $m = 1$, which is equal to the codimension of the fixed point locus mod 2.

Deep inside phases II and III, cf. the dotted line in Fig. 9, the field x_6 with charge $Q_6 = (1; 2)$ gets a vacuum expectation value, and relation (90) connects the slice $\tau = (0; \cdot)$ with $(\cdot; \cdot)$, and the slice $\tau = (0; 0)$ with $(\cdot; 2)$. Now recall from the discussion in Sec. 3.2 that the former two slices of the orientifold moduli space do not intersect the singular locus S . We can therefore move from the large volume point along Path A to the dashed line at infinity in phase II or III, change slice and move back to large volume. With our choice of σ_0 the orientifold sign σ_0 is not altered as we change slices.

On the other hand, deep inside phases III and IV, along the dotted line, the field p with charge $Q_p = (-4; 0)$ gets a vacuum expectation value. The associated shift of τ now does not correspond to a change of the slice. However, the orientifold sign is altered, $\sigma_0 = !_p^{-1} \sigma_0 = \sigma_0$. If we move along Path B in Fig. 9 we return to the original large volume point, but pick up a non-trivial monodromy on the D-branes. This discussion can be summarized in the following diagram, which shows how the various large volume




Figure 9. In the two-parameter model, Example 2, the two slices, $\tau = (0;)$ and $(;)$, are connected along the (dashed) line at infinity in phases II and III. This is due to the vacuum expectation value of the field x_6 . As a consequence, two large volume limits are connected via Path A in the moduli space. Path B induces a non-trivial monodromy but returns to the original large volume point.

points are connected via Paths A and B:

$$(95) \quad \begin{array}{ccc} D^{+1;1;(0;)}(X) & \xrightarrow{A} & D^{+1;1;(;)}(X) \\ \left| \begin{array}{c} B \\ ? \end{array} \right. & & \left| \begin{array}{c} B \\ ? \end{array} \right. \\ D^{-1;1;(0;)}(X) & \xrightarrow{A} & D^{-1;1;(;)}(X) \end{array}$$

Here, $D^{0^m;}(X)$ denotes the set of invariant D -branes on the toric variety X at large volume.

Let us analyse the orientifold planes $O_{(-1;2)}$. At large volume the fixed point locus of the involution is the union of two points,

$$p_a = O_{(+1; (-1)^a)} = \{x_a = x_3 = x_4 = x_5 = p = 0\} \quad \text{for } a = 1;2;$$

and two compact surfaces,

$$S_a = O_{(-1; (-1)^a)} = \{x_a = x_6 = p = 0\} \quad \text{for } a = 1;2;$$

At this point we could just use formula (94) in order to obtain the types. Let us be more explicit and put the probe branes on p_a resp. S_a .

The probe brane for the point p_a is the Koszul complex of $(x_a; x_3; x_4; x_5; p)$, which has $Q_f = (1; 1)$ and right-most gauge charge $q = (1; 0)$,

$$W_2(0; 1) \xrightarrow{f} \dots \xrightarrow{f} W_3(1; 0) :$$

The polynomial f_0 in the quasi-isomorphism (79) must therefore carry gauge charge according to the conditions

$$\begin{aligned} Q_{f_0} &= \dots = Q_f + 2q = (0; 0) \text{ on } \dots = (\dots); \\ Q_{f_0} &= \dots = Q_f + 2q = (1; 0) \text{ on } \dots = (0; \dots) : \end{aligned}$$

In the former situation $f_0 = 1$, thus the quasi-isomorphism is $U = U_{\text{inv}}$ and the type $\mathcal{O}_{p_a} = \dots_e$ of the orientifold plane on p_a is determined using (81),

$$\mathcal{O}_{p_a} = \dots_0 (1)^{(m-d)=2} \det \begin{pmatrix} 0 \\ f \end{pmatrix} = \dots (1^{\mathfrak{f}})_{\dots} :$$

In the latter case we can set $f_0 = x_a^2 x_6$, where $(x_1; x_2) = (x_2; x_1)$, with the quasi-isomorphism $V = x_a^2 x_6 (U_{\text{inv}})^{-1}$ and

$$\mathcal{O}_{p_a} = \dots_0 (1)^{(m-d)=2} \det \begin{pmatrix} 0 \\ f \end{pmatrix} = \dots (1^{\mathfrak{f}})_{\dots} :$$

For the surface S_a the naive Koszul complex does not provide an invariant D-brane. We therefore use a Koszul complex for $(x_a; x_3; x_4; x_6; p)$, which corresponds to a point on S_a . The type \mathcal{O}_{S_a} of the orientifold plane is then determined by $\mathcal{O}_{S_a} = \dots_e$. We have $Q_f = (1; 1)$ and right-most gauge charge $q = (1; 1)$,

$$W_2(0; 0) \xrightarrow{f} \dots \xrightarrow{f} W_3(1; 1) :$$

The condition on the gauge charge of f_0 reads

$$\begin{aligned} Q_{f_0} &= \dots = Q_f + 2q = (0; 0) \text{ on } \dots = (\dots); \\ Q_{f_0} &= \dots = Q_f + 2q = (1; 0) \text{ on } \dots = (0; \dots) : \end{aligned}$$

For $\dots = (\dots)$ the quasi-isomorphism is the invertible one, U_{inv} , and the type of the orientifold plane S_a is

$$\mathcal{O}_{S_a} = \dots (1^{\mathfrak{f}})_{\dots} :$$

For $\dots = (0; \dots)$ the quasi-isomorphism is $V = x_5 (U_{\text{inv}})^{-1}$ and

$$\mathcal{O}_{S_a} = \dots (1^{\mathfrak{f}})_{\dots} :$$

We observe that for both, the points and the surfaces, the two respective types are opposite, so that the total configuration of orientifold planes does not carry a net RR-charge in this example.

Let us summarize our results on the types of orientifold planes at large volume as follows:

large volume	$= (0;)$	$= (;)$
$o = +1$	$O_{p_1}^+ O_{p_2} O_{s_1} O_{s_2}^+$	$O_{p_1}^+ O_{p_2} O_{s_1}^+ O_{s_2}$
	 B ?	 B ?
$o = -1$	$O_{p_1} O_{p_2}^+ O_{s_1}^+ O_{s_2}$	$O_{p_1} O_{p_2}^+ O_{s_1} O_{s_2}^+$

From this diagram we find that all four different type assignments are connected through paths in moduli space. Path B leads back to the original large volume point, but still changes the overall type. Path A, which connects two different large volume points, swaps the types of the surfaces S_a .

6 Compact models

Let us next turn to linear sigma models with superpotential. They give rise to compact low-energy configurations. From now on we have to deal with matrix factorizations instead of complexes. In fact, many of the features that we observed for complexes in the previous section carry over to matrix factorizations, so that we elaborate on the peculiarities of the latter in the following.

First thing to keep in mind when turning on a superpotential is that in order to satisfy the homogeneity equation (7) some of the chiral fields have to carry non-vanishing R-charge. In particular, for a gauge-invariant potential of the form

$$(96) \quad W(p; x_i) = \sum p G(x_i);$$

we will assign R-charge +2 to the fields p and 0 to the fields x_i . This non-trivial charge assignment plays a special role in Landau-Ginzburg orbifold phases, which we discuss in Sec. 6.2.

Second, in a phase where the superpotential (96) gives rise to F-term masses for p and the transverse modes to $G(x_i) = 0$, the low-energy dynamics is restricted to the subvariety $M = \{p = G = 0\}$. In that case the matrix factorizations are mapped to geometric D-branes on M by the Konnermann map [20]. We start in Sec. 6.1 with investigating how the worldsheet parity action P^m on the matrix factorizations gets mapped to the parity action P_B^m on the geometric D-branes in the low-energy configuration.

6.1 The effect of the Konnerth map on the parity action

Let us review the standard Konnerth periodicity before we move on to orientifolds and the bre-wise version that is needed in the context of gauged linear sigma models. Consider phase space C^N with coordinates $x_1; \dots; x_N$ and C^{N+2} with coordinates $u; v; x_1; \dots; x_N$. Over the latter we consider the superpotential $W(u; v; x) = uv + \tilde{W}(x)$.

Konnerth periodicity then states that the set of (isomorphism classes of) matrix factorizations of $W(u; v; x)$ over C^{N+2} is equivalent to the set of matrix factorizations of $\tilde{W}(x)$ over C^N . Physically, in the Landau-Ginzburg model the superpotential $W(u; v; x)$ gives masses to the fields u and v so that they can be integrated out in the infra-red. A canonical matrix factorization of the term uv in $W(u; v; x)$ then establishes the equivalence of the two sets of matrix factorizations. The canonical matrix factorization reads

$$Q_c(u; v) = \begin{pmatrix} 0 & v \\ u & 0 \end{pmatrix} :$$

In fact, a matrix factorization $\mathcal{Q}(x)$ of $\tilde{W}(x)$ is mapped to a matrix factorization of $W(u; v; x)$ by taking the graded tensor product with Q_c ,

$$(97) \quad \mathcal{Q}(x) \otimes Q_c(u; v; x) = Q_c(u; v) \otimes \text{id}_c \mathcal{Q}(x) :$$

Conversely, Konnerth observed in [64] that by isomorphism any matrix factorization of $W(u; v; x)$ can be brought to the tensor product form as in (97), thus providing $\mathcal{Q}(x)$.

In the context of gauged linear sigma models the coordinate fields, here $u; v; x_1; \dots; x_N$, carry gauge charges as well as R-charges, and we have to determine how their representations on D-branes are mapped under Konnerth periodicity. The tensor product (97) dictates the following decomposition of representations,

$$(98) \quad \begin{aligned} \rho(g) &= \rho_c(g) \otimes \rho_e(g) ; \\ R(\rho) &= R_c(\rho) \oplus \mathbb{R}(\rho) : \end{aligned}$$

Let us set the R-charges for u and v to 0 and 2, the R-charge assignment of the remaining fields x_i does not play a role for the subsequent discussion. The representation of the R-symmetry $R_c(\rho)$ can be chosen up to an (unphysical) multiplication by a character $\chi(\rho)$. We set $R_c(\rho) = \text{diag}(1; \dots)$.

In order to determine the representation $\rho_c(g)$ in the uv -system we note that the canonical matrix factorization Q_c can be obtained by quantizing a boundary fermion ψ ; with canonical commutation relations $\{\psi, g\} = 1$.

The two states in the Hilbert space of the boundary fermion are $|j\rangle$ and $|\bar{j}\rangle$, where $\langle j|\bar{j}\rangle = 0$. The fermion is coupled to the bulk uv-system through the boundary supercharge $Q_c = u + v$.

If we assign to v and u the charges Q_v resp. $-Q_v$, then gauge invariance of the boundary supercharge requires $Q_c = Q_u = Q_v$. Accordingly, the Chan-Paton factors associated with the states $|j\rangle$ and $|\bar{j}\rangle$ have the canonical charge assignments $Q_v=2$ resp. $-Q_v=2$. Note however that $Q_v=2$ does not have to be an integer. We therefore shift it into the auxiliary theta angle $\theta_c = Q_v$ and obtain $\rho_c(g) = \text{diag}(1; g^{Q_v})$. The canonical matrix factorization is therefore

$$B_c : \underline{W}(0) \stackrel{u}{=} \overline{W}(-Q_v) :$$

To summarize, a matrix factorization of $\overline{W}(x)$ is mapped to a matrix factorization of $W(u; v; x)$ by taking the tensor product with B_c . The auxiliary theta angle gives rise to the non-trivial relation

$$(99) \quad \theta_c = \tilde{\theta} + Q_v ;$$

where θ_c and $\tilde{\theta}$ are the theta angles of the ultra-violet theory, including the uv-system, and the infra-red theory, respectively.

Orientifolds and the Korrer periodicity

Let us now check the compatibility of the Korrer map with the parity action. Note first that condition (41) on the superpotential requires that the involution acts in the uv-system as $(u; v) \mapsto (\iota_v^{-1}u; \iota_v v)$ for some phase ι_v .

We need to determine how the parity operator on matrix factorizations of $W(u; v; x)$ splits up in the tensor product (97) and (98). On the canonical matrix factorization the world sheet parity acts as

$$\begin{aligned} P_c(Q_c) &= Q_c^T ; \\ P_c(\rho_c(g)) &= \rho_{c^{-1}}(g) \rho_c(g)^T ; \\ P_c(R_c(\lambda)) &= m_c(\lambda) R_c(\lambda)^T ; \end{aligned}$$

where $m_c = 1$. The canonical quasi-isomorphism that makes B_c invariant is

$$U_c = \begin{pmatrix} 0 & 1 \\ \iota_v & 0 \end{pmatrix} :$$

It satisfies

$$(100) \quad U_c = \rho_c(\lambda)^{-1} U_c^T \text{ with } \rho_c = \iota_v^{-1} :$$

Using the results of Sec. 4.7 on tensor product branes we can construct the quasi-isomorphism U for the matrix factorization $Q(u;v;x)$ in terms of the quasi-isomorphism \tilde{U} for $\tilde{Q}(x)$. The relations between the constants associated with the orientifold action can be summarized as follows:

$$(101) \quad \begin{aligned} m &= m - 1 ; \\ \tilde{m} &= \tilde{m} - Q ; \\ \tilde{v} &= \tilde{v} : \end{aligned}$$

In summary, we found that the Konnermann map relates the sets of invariant D-branes as follows:

$$MF_W^m ; (C^{N+2}) \xrightarrow{=} MF_{\tilde{W}}^{\tilde{m}} ; (C^N)$$

In particular, the dressing of the parity action by the antibrane operator $(-1)^F_L$ changes under the Konnermann map, i.e. no dressing maps to dressing and vice versa.⁸ This result is in agreement with [18].

Fibre-wise Konnermann map

Let us return now to our original question. Given a parity operator in the linear sigma model we want to determine the parity operator on the compact hypersurface M_r in a geometric phase.

As pointed out in [20] the matrix factorizations in $MF_W(C^N; T)$ and the geometric D-branes in $D(M_r)$ are related by a fibre-wise version of Konnermann periodicity. For $W = pG(x)$ we can therefore adopt our previous discussion, replacing $(v;u)$ by $(p;G(x))$ and setting $\tilde{W} = 0$.

If we have a superpotential $W = \sum_{p=1}^P p G(x)$ that gives rise to a complete intersection M_r in the large volume phase we have to apply fibre-wise Konnermann periodicity P times,

$$MF_W^m ; (C^{N+2}) \xrightarrow{=} D^{\tilde{m} B} (M_r)$$

where the relation between the B-field and the theta angle is

$$B = \sum_{p=1}^P \frac{X'_p}{Q_p} ;$$

and the dressing by the antibrane operator is shifted according to

$$m = m' ;$$

⁸This shift was observed in the context of defects in [90]

The relation between orientifold signs is $\tilde{\cdot} = (-1)^{\sum_{i=1}^Q \epsilon_i}$. Here the phases ϵ_i are defined through the holomorphic involution on the field \mathbb{P}^1 , i.e. $\tilde{\cdot}(\epsilon) = \epsilon^*$. In a general coordinate basis the action of the involution on the ϵ_i 's may not be diagonal, so that the more invariant expression between the ultra-violet and the infra-red signs is in terms of the determinant of $\tilde{\cdot}$ acting on the ϵ_i 's or on the polynomials G_i , i.e.

$$\tilde{\cdot} = (-1)^{\sum_{i=1}^Q \epsilon_i} = \det(\tilde{\cdot})^{-1}$$

As we have reviewed in Sec. 2.1 the price to pay for applying the Konner normal map bre-wise is to deal with half-infinite complexes in $D(M_r)$. The worldsheet parity action on R -degrees, $\mathbb{Z}^m \rightarrow \mathbb{Z}^m$, implies that \mathbb{P}_B^m maps right-to-left-infinite complexes and vice versa. A D -brane is invariant if there exists a quasi-isomorphism between the left- and the right-infinite complex, i.e. the joined complex must be an infinite exact (i.e. empty) complex.

Since the description in terms of infinite complexes is cumbersome, in particular in the situation of complete intersections, we prefer to work directly with the matrix factorizations in the linear sigma model in the subsequent examples. We will make an exception if the low-energy D -brane in $D^{\text{irr}}(M_r)$ is expressible through a finite complex of vector bundles.

6.2 Landau-Ginzburg orbifolds and the orientifold moduli space

Landau-Ginzburg orbifolds are the 'compact' analog of the orbifold models that we discussed in Sec. 5.1. We therefore closely follow the discussion therein. We start with the one-parameter model, where the vacuum expectation value for the field ϕ of charge $Q_\phi = N$ breaks the gauge group from $U(1)$ to Z_N .

The main difference to the noncompact situation is the non-trivial R -charge assignment, $R_\phi = 2$, for the chiral field ϕ . When this field gets a vacuum expectation value, for instance $\phi = 1$, it is convenient to dress the R -symmetry by a global gauge transformation, i.e. in the Landau-Ginzburg model we use the shifted R -symmetry with charges $R_i = R_i + 2Q_i/N$ on the chiral fields. A matrix factorization is correspondingly mapped from the linear sigma model to the Landau-Ginzburg model through

$$\begin{aligned} Q(x) &:= Q(p=1;x); \\ \tilde{\cdot} &:= \cdot; & \text{for } 2 & \in U(1); \\ R(\cdot) &:= R(\cdot) \cdot (2/N)^{-1}; \\ U &:= U(p=1); \end{aligned}$$




Figure 10. The N -fold cover of the Kähler moduli space, $e^t = (\frac{N}{N})^N$. For both N even and odd only the large volume point with non-trivial B-field, $B = \pi$, is connected to the Gepner point, the large volume point with vanishing B-field is not. This is due to the relation $B = \frac{2\pi}{N} + N$.

The Landau-Ginzburg orbifold data of the D-brane clearly satisfies the invariance conditions (66) for the discrete group Z_N , provided that the theta angle is given by

$$\theta = 2\pi \frac{m}{N} \text{ mod } 2\pi ;$$

and the R-symmetry character $\chi_m(\theta)$ is determined by

$$(102) \quad \chi_m(\theta) = e^{i\theta m} + \frac{2\pi}{N} - 2\pi Q ;$$

In this way we obtain the set of invariant D-branes $M_{F_W}^m ; (C^N ; \theta)$ in the Landau-Ginzburg orbifold model, as it was studied before in [17]. Therein, the triangulated structure of the category of matrix factorizations was worked out in detail. In particular, the world sheet parity action was represented as a functor on the triangulated category.

In Sec. 5.1 we found that a shift of the theta angle by $2\pi/N$ leaves the theory invariant at the orbifold point. In view of (102) this shift has to be supplemented by a shift of m , so that χ_m is not altered, i.e.

$$(103) \quad \chi_m(\theta + \frac{2\pi}{N}) = \chi_{m+1}(\theta) = \chi_m(\theta) , \quad \chi_m(\theta) = \chi_{m+N}(\theta) ;$$

The lift of a matrix factorization from the Landau-Ginzburg orbifold to the linear sigma model can be found along the lines of Sec. 5.1 and is explained in detail in [20]. According to (103) we have a choice in lifting to different slices of the moduli space. Suppose we have a matrix factorization with quasi-isomorphism U for given $(\theta ; m)$. Then in view of the

relations (66) a combined shift (103) implies that we have to dress the quasi-isomorphism by the field p , i.e. $U^0 = pU$. Note that $m \mathbb{T} m + 2$ is conform with the R-charge $R_p = 2$. The sign of the orientifold action is altered according to (89),

$$^0 = !_p^1 :$$

Note that the implications for the moduli space are essentially the same as for the non-compact situation, see Fig. 10.

Higher-dimensional moduli spaces

Let us generalize this discussion to models with higher-rank gauge group, $T = U(1)^k$. We consider a phase where the deleted set x_r has one or several irreducible component of the form $fx_1 = 0g$ for some $l \in \{1, \dots, N\}$. The vacuum expectation value for x_1 then breaks the gauge group so that $q^a = q_1^a + Q_1^a$ and in particular,

$$(104) \quad q^a = q_1^a + Q_1^a ; \quad m^0 = m + R_1 ; \quad ^0 = !_1^1 :$$

Note that the field x_1 may or may not carry R-charge.

We obtain the equivalence

$$M_{F_W}^m ; (X_r) =! M_{F_W}^{m^0 ; ^0} (X_r) :$$

As distinguished from the non-compact situation (91) the integer m may get shifted by 2. In order to see that this can indeed have a non-trivial effect, recall that a common shift of R-degree $[1] : j \mathbb{T} j - 1$ is accompanied by $m \mathbb{T} m - 2$. This can be used to undo the shift of m in (104). However, the orientifold sign is then altered according to (64), that is $^0 \mathbb{T} ^0$.

6.3 The type of orientifold planes

Recall from Sec. 3.1 that in a geometric large volume phase the orientifold plane O is given by the intersection of the fixed point locus $\text{Fix}(g)$ with the holomorphic subvariety $M = \{p = G = 0\}_g$. This intersection may be reducible, $O = O ;$, which adds some subtleties as compared to the discussion of the type of orientifold planes for non-compact models in Sec. 5.2. The assumptions on the applicability of the type formulas are the same as in Sec. 5.2.

For the following let us denote the ambient space by $Y = \{p_1 = \dots ; p_r = 0\}_g \subset X$ and the complete intersection by $M = \{G_1(x) = \dots = G_r(x) = 0\}_g \subset Y$. For simplicity we work in a coordinate basis that diagonalizes g .

For a given component $O_{\mathbb{P}^1}$ in M , let us probe the type with a Koszul-like matrix factorization with tachyon profile $Q = \sum_i (f_i - i + g_i - i)$. The polynomials f_i are given by those coordinates x_i whose common zero locus is $O_{\mathbb{P}^1}$. Note that the fields p are not included, because in the geometric phase they obtain zero expectation values from the F -term equations no matter how the involution σ acts on them.

The polynomials that determine $O_{\mathbb{P}^1}$ can be separated in two sets. The first contains coordinate fields x_i that are not invariant under σ . Let us denote them by $f_1; \dots; f_s$. Their common zero locus gives $\text{Fix}(\sigma)$. In order to restrict to M and to pick an irreducible component we have to add a finite number of σ -invariant polynomials, $f_{s+1}; \dots; f_{s+r}$, where $r \geq 0; \dots; g$ is the number of polynomials needed to restrict to M and to pick an irreducible component.

As for the non-compact models in Sec. 5.2 the resulting Koszul-like matrix factorization may not be an invariant D-brane. In some cases a way out is to utilize a lower-dimensional probe brane to determine the type. For that we need to add the appropriate σ -invariant polynomials $f_{s+r+1}; \dots; f_{s+r+2p}$ to the Koszul complex. The type is then given by

$$(105) \quad \mathcal{O} = \mathcal{O}(\mathbb{P}^1)_e = \mathcal{O}(\mathbb{P}^1)^{(m-D+g)=2} =_{2q} \mathcal{O}(\mathbb{P}^1) \det(\mathcal{O}_{\mathbb{P}^1}) \det(\mathcal{O}):$$

Here, $D = s + r + g$ is the codimension of $O_{\mathbb{P}^1}$ in M . Notice that its codimension in the ambient space Y is $s + r$. For \mathcal{O} containing only signs we obtain the simpler expression

$$(106) \quad \mathcal{O} = \mathcal{O}(\mathbb{P}^1)^{(m+D+g)=2} =_{2q} \mathcal{O}(\mathbb{P}^1) \quad \text{for } a = 1:$$

6.4 Orientifolds with and without vector structure

Compactifications without vector structure have been introduced in [9,10], where they were investigated for toroidal compactifications, see [11,12] for recent works. The starting point was the observation that the gauge group for the heterotic string is $\text{Spin}(32) = \mathbb{Z}_2$ rather than $\text{SO}(32)$. This allows compactifications with gauge bundles which do not admit vectors of $\text{SO}(32)$. The obstruction to having vector structure is determined by a generalized Stiefel-Whitney class w_2 , defined modulo 2. On the dual type I side it was observed that the choice of w_2 corresponds to the choice of a discrete B-field that is still allowed by the orientifold projection, see [91].

Under T-duality these compactifications get mapped to IIB compactifications with $O7$ -planes. As opposed to the T-dual of a compactification with

vector structure, the different orientifold planes will have unequal type, leading effectively to a rank reduction of the gauge group. The orientifold action with fixed points on a two torus has four $O7$ -planes. In the case without vector structure three of them are O^- planes, and one is an O^+ , such that tadpole cancellation requires only 8 D7 branes, resulting in the gauge group $SO(8)$.

In the current paper we have developed a framework where the physics of orientifolds can be studied over the whole Kahler moduli space, in particular for all values of the discrete B-field. The earlier results on compactifications without vector structure should therefore be reproduced by our methods.

Example 1 with $N = 3$ and superpotential

To see this in the simplest example, we consider orientifolds of the cubic torus $E = \mathbb{C}P^2$. The superpotential is taken to be

$$(107) \quad W = p(x_1^3 + x_2^3 + x_3^3) :$$

We will focus on the holomorphic involution

$$(108) \quad \sigma(x_1; x_2; x_3; p) = (x_2; x_1; x_3; p) :$$

As was discussed in Sec. 3.2 its fixed point set at large volume consists of $3 + 1$ points on the torus,

$$\begin{aligned} O_{-1}; &= \{x_1 = x_2 = x_3 = 0\} \quad \text{for } p = 0; \\ O_{+1} &= \{x_1 + x_2 + x_3 = 0\} : \end{aligned}$$

The types can readily be computed using (106), where $r = 1$ for the three points and $r = 0$ for the single one. Taking into account the shift $B = +3$ we obtain

$$(109) \quad \begin{aligned} O_{-1}; &= O^-(1)^{r=2} (1)^{B=3} ; \\ O_{+1} &= O^-(1)^{r=2} : \end{aligned}$$

As expected the four points have equal type for vanishing B-field. Otherwise, for nonvanishing B-field the type of one point is different from the types of the other three points. Note that m being even in the gauged linear sigma model means m being odd in the geometric phase, so that the parity action is dressed in the infra-red by the anti-brane operator $(-1)^{F_L}$, as it should be for $O7$ -planes in the type IIB context.

Let us be more explicit and construct the probe branes that are used to test the type of each of these points. Since the fixed point set can in each case be described by two linear equations, $f_1 = f_2 = 0$, the matrix factorizations are of Koszul type with $W = f_1 g_1 + f_2 g_2$, see the general discussion in section 4.6.

The B-eld turned on

For $\epsilon = 0$ the matrix factorization and its parity in age take the form isomorphism between the orientifold and its image.

$$\begin{array}{ccc}
 W_{\frac{m-2}{2}}(1) \xrightarrow[\text{?}]{\begin{array}{c} f_1 \\ f_2 \\ \hline (g_1; g_2) \end{array}} W_{\frac{m}{2}}(0) \xrightarrow[\text{?}]{\begin{array}{c} (f_2; f_1) \\ \hline g_1 \\ g_2 \end{array}} W_{\frac{m+2}{2}}(1) \\
 \left| \begin{array}{c} u_1 \\ \text{?} \end{array} \right. & & \left| \begin{array}{c} u_0 \\ \text{?} \end{array} \right. \\
 W_{\frac{m-2}{2}}(1) \xrightarrow[\text{?}]{\begin{array}{c} f_2 \\ f_1 \\ \hline (g_1; g_2) \end{array}} W_{\frac{m}{2}}(0) \xrightarrow[\text{?}]{\begin{array}{c} (f_1; f_2) \\ \hline g_1 \\ g_2 \end{array}} W_{\frac{m+2}{2}}(1) \\
 \left| \begin{array}{c} u_1 \\ \text{?} \end{array} \right. & & \left| \begin{array}{c} u_1 \\ \text{?} \end{array} \right.
 \end{array}$$

The brane and its image t through the window $w = \dots < \dots$ with $N = f_1; 0; 1g$, see Sec. 2.2. In order to determine the isomorphism $(u_1; u_0; u_1)$ we need to consider the individual orientifold points separately.

To the single point O_{+1} we can associate the factorization $W = \mathbb{P}^1_a f_a^1 g_a^1$ with

$$\begin{aligned}
 f_1^1 &= x_1 + x_2; & f_2^1 &= x_3 \\
 g_1^1 &= p(x_1^2 - x_1 x_2 + x_2^2); & g_2^1 &= p x_3^2:
 \end{aligned}$$

The polynomials f_a^1 are odd under the holomorphic involution σ . The isomorphism is then given by

$$u_1^1 = 1; \quad u_0^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad u_1^1 = 1:$$

We find that $(U^1)^T = U^1$ and hence $o_{+1} = \dots_e = \dots_0(1)^{n=2}$, which confirms the result (109) for non-vanishing B-eld.

For the three orientifold points O_{-1} ; we have

$$\begin{aligned}
 f_1 &= x_1 - x_2; & f_2 &= x_3 - x_2 \\
 g_1 &= p(x_1^2 + x_1 x_2 + x_2^2); & g_2 &= p(x_3^2 + x_2 x_3 + x_2^2):
 \end{aligned}$$

The holomorphic involution acts on the polynomials as

$$f_1 = f_1; \quad f_2 = f_2 + f_1;$$

so that the isomorphism U is given by

$$u_1 = 1; \quad u_0 = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}; \quad u_1 = 1:$$

As a consequence, $(U)^T = U$ and therefore $o_{-1} = \dots_e = \dots_0(1)^{n=2}$. This confirms that for non-vanishing B-eld the type at the three orientifold points is opposite to the one at the single point calculated before.

To make contact with the discussion in Sec. 4.6 note that $\det f = -1$ for the three points, and $\det f = 1$ for the single point, such that our explicit calculation is in agreement with the general discussion.

Vanishing B-field

Let us next turn to the case $\tau = 0$. The D-brane and its image are related as follows,

$$\begin{array}{ccccc}
 W_{\frac{m-2}{2}}(2) & \xrightarrow[\text{(g}_1; \text{g}_2)]{\begin{array}{c} f_1 \\ f_2 \end{array}} & W_{\frac{m}{2}}(1)^2 & \xrightarrow[\text{g}_1]{\text{(f}_2; \text{f}_1)} & W_{\frac{m+2}{2}}(0) \\
 \left| \begin{array}{c} u_1 \\ ? \end{array} \right. & & \left| \begin{array}{c} u_0 \\ ? \end{array} \right. & & \left| \begin{array}{c} u_1 \\ ? \end{array} \right. \\
 W_{\frac{m-2}{2}}(1) & \xrightarrow[\text{(g}_1; \text{g}_2)]{\begin{array}{c} f_2 \\ f_1 \end{array}} & W_{\frac{m}{2}}(0)^2 & \xrightarrow[\text{g}_2]{\text{(f}_1; \text{f}_2)} & W_{\frac{m+2}{2}}(1)
 \end{array}$$

Obviously, U can in this case not be an isomorphism, it increases the degree by one, and therefore can only be a quasi-isomorphism linear in the coordinates x_i . That U is a quasi-isomorphism means that the bound state of the brane and its image brane obtained by binding them using the tachyon profile given by U is an empty brane. Which branes are empty depends on the phase under consideration. Since we are interested in relating our construction to compactifications without vector structure, we would like to make contact with the geometric regime at large volume. Here, the set $r = fx_1 = x_2 = x_3 = 0g$ is excluded and any brane located there flows to an empty brane. This means that the quasi-isomorphism should be of the form $f_0 U$, where U is the isomorphism considered previously, and f_0 is a polynomial in the fields x_i such that the common zero locus of $(f_0; f_1; f_2)$ is contained in r .

For the single fixed point O_{+1} one can choose

$$f_0^1 = x_1 - x_2 :$$

Since f_0 is symmetric under the holomorphic involution, we conclude that again $(f_0^1 U^1)^T = f_0^1 U^1$, such that the type does not change, $o_{+1} = o_0(1)^{\eta=2}$.

At the three fixed points O_{-1} ; this is different. Here, one can choose

$$f_0 = x_1 + x_2 :$$

Since the polynomial f_0 flips sign under parity transformation, we find that

$(f_0^1 U^1)^T = -f_0^1 U^1$ and the type of the orientifold will also flip, that is $o_{-1}; = o_0(1)^{\eta=2}$. All four points carry the same type for vanishing B-field.

To summarize, the transformation properties of the quasi-isomorphism between a D-brane and its parity image determine whether or not the orientifold type is changed when the theta angle is modified.

6.5 Type change in the orientifold moduli space

In models with higher-dimensional Kahler moduli space it may happen that different large volume points are connected via a path in moduli space. In this section we illustrate the change of orientifold type along paths in the compact version of Example 2.

Example 2

Recall the charges (10) and the moduli space from Fig. 9. The superpotential is $W = pG(x)$ with a quasi-homogeneous polynomial $G(x)$ of gauge charge $(4;0)$. For simplicity we pick the Fermat type polynomial $G(x) = x_6^4(x_1^8 + x_2^8) + x_3^4 + x_4^4 + x_5^4$. For the world sheet parity action we choose the holomorphic involution [14]

	x_1	x_2	x_3	x_4	x_5	x_6	p
σ	+1	+1	1	1	1	+1	1

which acts diagonally on the chiral fields. We set m to be even.

Let us follow the two paths in Fig. 9. The two slices of interest have theta angles $\theta = (0; \cdot)$ and $\theta = (\cdot; \cdot)$. Along Path A we meet the dashed line, which stretches between the Landau-Ginzburg point and the weighted projective moduli point. On this line x_6 obtains a vacuum expectation value, and according to the shifts (104) neither β_0 norm is altered when we change from slice $(0; \cdot)$ to $(\cdot; \cdot)$.

Following Path B is different. Along the dotted line between the Landau-Ginzburg point and the hybrid point the field p gets a vacuum expectation value, thus connecting $\theta = (0; \cdot)$ with $\theta = (\cdot; \cdot)$. The corresponding shifts are $\beta_0 = m + 2$ and $\beta_0 = \beta_0$. Indeed Path B connects the large volume theories $M_{F_W}^{m;(0; \cdot)}(X)$ and $M_{F_W}^{m+2;(0; \cdot)}(X)$. In the latter we can perform an overall shift of the R-degrees and use (64) to find that

$M F_W^{m+2;(0)}(X) \xrightarrow{[1]} M F_W^{m;(0)}(X)$. We obtain the following diagram:

$$\begin{array}{ccc} D^{m-1;(0)}(M) & & D^{m-1;(0)}(M) \\ \left\| \begin{array}{c} = \\ ? \end{array} \right. & & \left\| \begin{array}{c} = \\ ? \end{array} \right. \\ M F_W^{m;(0)}(X) & \xrightarrow{A} & M F_W^{m;(0)}(X) \\ \cdot K_B[1] & & \cdot K_B[1] \end{array}$$

The vertical map is the Konner map. Note that as compared to the diagram (95) for the non-compact model, now the sets of invariant D-branes $M F_W^{m+1;(0)}(X)$ and $M F_W^{m;(0)}(X)$ are not connected through a path in Kähler moduli space.

Let us investigate the fixed point locus of σ on the hypersurface $M = \text{fp} = G(x) = 0g$ at large volume. The non-trivial components O are

$$\begin{aligned} O_{(+1,1)} &= \{x_3 = x_4 = x_5 = 0g\} \subset M; \\ O_{(-1,1)} &= \{x_6 = 0g\} \subset M; \end{aligned}$$

The second is a divisor D , and the first is a union of eight points on M ,

$$p = O_{(+1,1)} = \{x_1 = x_2 = x_3 = x_4 = x_5 = 0g\} \subset M \quad \text{for } g = 1:$$

Let us compute the type of $O_{(+1,1)}$ first. The matrix factorization for the probe brane can be written in terms of boundary fermions as $Q = f_a(x) + g_a(p;x)$ with

$$f_1 = x_1; \quad f_2 = x_2; \quad f_3 = x_3; \quad f_4 = x_4; \quad f_5 = x_5;$$

The polynomials g_a are such that $W = \sum_a f_a g_a$. Using formula (106) with $r = 1$ we find for the type,

$$O_{(+1,1)} = O(-1)^{m=2}:$$

The type of $O_{(-1,1)}$ is computed using (106) with $r = 1$,

$$O_{(-1,1)} = O(-1)^{m=2}(-1)^{1=1}:$$

The model

The type IIB compactification at hand is a system of $O7$ -planes and $D7$ -branes on a degree eight hypersurface in weighted projective space $W\mathbb{P}_{11114}^8$. This is the large volume point of the following gauged linear sigma model.

	x_1	x_2	x_3	x_4	p	
$U(1)$	1	1	1	1	4	8
σ_0	+1	+1	+1	+1	1	1

The involution σ_0 acts diagonally on the coordinates with the signs given in the table. The superpotential is given by $W = pG(\mathbf{x})$, where

$$G(\mathbf{x}) = h(x) + \sum x_i^2;$$

and $h(x)$ is a degree eight polynomial. The low-energy configuration at large volume is the hypersurface $M = fG(\mathbf{x}) = 0$ in $W\mathbb{P}_{11114}^8$. The B-field vanishes. We set $m = 0$ in the gauged linear sigma model, which means $\kappa = 1$ in the non-linear sigma model on the hypersurface, i.e. the parity action is dressed by $(-1)^F$, as it should be for an $O7$ -plane.

The fixed point locus of σ_0 gives the orientifold plane at $(x = +1)$

$$O_{+1} = f = 0 \text{ on } M;$$

As the orientifold plane descends from F-theory, it is an $O7$ -plane, which means that $\kappa_{+1} = 1$. Let us apply the type formula (106) with the codimension of the orientifold plane $D = 1$, $\ell = 1$ for a Calabi-Yau hypersurface, and $r = 1$. We obtain $\tilde{\kappa}_0 = \kappa_0 = \kappa_1 = +1$. The set of invariant D -branes is therefore given by

$$M \cap F_W^{+1; \kappa=0}(X) = D^{+1; \kappa=0}(M):$$

All D -branes that we consider in the following must be contained in this set.

The $D7$ -brane descending from F-theory carries gauge group $SO(N)$ and is localized on the divisor

$$(110) \quad D = f(x)^2 - \sum x_i^2(x) = 0 \text{ on } M;$$

where $f(x)$ and $\sum x_i^2(x)$ are polynomials of degree n resp. $2n - 8$ for some integer $n > 4$.¹⁰

¹⁰In fact, in the configuration that descends from F-theory the integer takes the value $n = 16$ and the gauge group on the D -brane is $O(1)$, so that tadpole cancellation is automatic.

Invariant D-branes in the orientifold background

The D7-brane on D contains a curve of ordinary double points at $f = 0$ for $g \neq 0$, which lies on the intersection with the orientifold plane. This singular curve pinches o at the points $f = g = 0$, which are locally described by the Whitney umbrella $fu^2 = v^2wg \subset \mathbb{C}^3$. The number of these pinch points is given by multiplying the degrees of the polynomials $(h; g)$ that define their location, that is $8 \cdot n = (2n - 8)$. One of the goals of [34,36] was then to explain this singular behaviour and to find a mechanism that prohibits deforming the special divisor D to a generic degree $2n$ divisor,

$$D^0 = \text{fP}_{2n}(x) = 0 \text{g} \quad M :$$

In our approach we first check the gauge group for a D7-brane on the divisor D^0 , that is for a coherent sheaf $\mathcal{O}_{D^0}(n)$. In $D^{+1; 1\beta=0}(M)$ it can be described through a D9D9 system given by the complex

$$(111) \quad \mathcal{O}_M(n) \xrightarrow{\text{P}_{2n}(x)} \mathcal{O}_M(n) :$$

\mathcal{O}_M is the pull-back of the trivial holomorphic line bundle from the ambient space $W \subset \mathbb{P}_{11114}^8$ to the hypersurface M .¹¹ In order to determine the gauge group, let us compute the external sign for this D-brane. Since the D-brane (111) is a Koszul complex (73) of just one polynomial, we can apply formula (81) with $m = 1$, $\sim_0 = +1$, and $n = 1$. For illustration let us be more explicit here. In matrix form the tachyon profile Q and the isomorphism U that satisfies the invariance conditions (66) are given by

$$Q = \begin{pmatrix} 0 & \text{P}_{2n}(x) \\ 0 & 0 \end{pmatrix} ; \quad \text{and} \quad U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} :$$

The condition (67) then gives $\sim_{0i} = +1$. The external sign is therefore readily computed to be $\epsilon = \sim_0 = \sim_{0i} = -1$, and tells us that $\mathcal{O}_{D^0}(n)$ has to carry gauge group $Sp(N)$.¹² Note that we could even choose the polynomial $\text{P}_{2n}(x)$ to assume the special form $(x)^2 \cdot h(x) \cdot (x)$. The gauge group on this brane tells us however that the coherent sheaf $\mathcal{O}_{D^0}(n)$ is not the D-brane from the weak-coupling limit of F-theory, although it shares the same world

¹¹Note that in this simple situation the complex can be lifted to a matrix factorization by tensoring it with the canonical matrix factorization

$$\underline{W(0)} \xrightarrow{\quad G \quad} \underline{W(8)} :$$

Instead of doing so we will directly work in $D^{+1; 1\beta=0}(M)$.

¹²Instead of the even polynomial $\text{P}_{2n}(x)$ we could have considered a divisor determined through an odd polynomial $\text{P}_{2n-4}(x)$. This would lead to $\epsilon = +1$ and therefore to gauge group $SO(N)$. This divisor is however reducible into two components [34,36], one of them lying on the O7-plane. But this is again not the configuration.

volum e. In particular, there are no obstructions to deforming $\mathcal{O}_D(n)$ back to a generic divisor. We conclude that the D-brane that descends from F-theory on the divisor D cannot correspond to a single D9D9-system (111).

As suggested in [36] the next best guess for the actual D-brane that descends from F-theory is a rank two D9D9-system, that is a complex

$$(112) \quad \begin{array}{ccc} \mathcal{O}_M(a) & \underline{T(\cdot; \underline{x})} & \mathcal{O}_M(a) \\ \mathcal{O}_M(b) & & \underline{\mathcal{O}_M(b)} \end{array} ;$$

where $T(\cdot; \underline{x})$ is a rank two tachyon profile.

The D-brane (112) is invariant if we find an isomorphism U that satisfies condition (72). In fact, we have

$$u^0 = \sim_{0i} \sim_0 (u^1)^t = e \cdot \text{id} ;$$

where we used $e = \sim_0 = \sim_{0i} = \sim_{0i}$ and the freedom of choosing a basis for the Chan-Paton space to set $u^1 = \text{id}$. The invariance condition on $Q(\underline{x})$ in (66) becomes

$$T = e \cdot T^t$$

Recall that the D-brane should carry an orthogonal gauge group $SO(N)$, so $e = +1$ and the tachyon profile takes the form

$$T(\cdot; \underline{x}) = \dots + \dots ;$$

In the infrared the D-brane localizes on the determinant

$$\det T = \dots^2 (\dots^2) + \dots^2 ;$$

which is a polynomial of degree $2n = 2(a+b)$.

The determinant is already very similar to the polynomial in D . In fact [36], the D-brane on the divisor D corresponds to the tachyon profile T with the largest number of deformation parameters in the polynomials. It can be obtained by setting $a = 2$ and $b = n - 2$. Then the polynomials $(\cdot; \cdot; \cdot; \cdot)$ have degrees $(0; n-4; 2n-8; n)$. In that case the polynomial is redundant and can be set to zero by a similarity transformation of the Chan-Paton space. Finally setting $\dots = 1$, the tachyon profile becomes

$$T(\cdot; \underline{x}) = \dots ;$$

and its determinant is precisely the polynomial in (110), i.e.

$$D = f \det T = \dots^2 \dots^2 = 0g ;$$

We conclude that we have found strong indications that the D7-brane from the weak-coupling limit of F-theory corresponds to a rank two D9D9 system that carries gauge group $SO(N)$ and is localized on the divisor D . It does not however correspond to the coherent sheaf $\mathcal{O}_D(n)$. In fact, we found that the latter supports the gauge group $Sp(N)$.

Acknowledgment: We thank Robert Haslhofer for collaboration at an early stage of this project. We appreciated valuable discussions with Costas Bachas, Massimo Bianchi, Andres Collinucci, Mboyo Esole, Gabriele Honecker, Kentaro Hori, Hans Jockers, Daniel Kreier, Wolfgang Lerche, Fernando Marchesano, Greg Moore, Christian Romelsberger, Emanuel Scheidegger, Angel Uranga, and Johannes Walcher. M.H. thanks the ESI in Vienna, the LMU and the DFG cluster of excellence "Origin and Structure of the Universe" in Munich as well as the ETH Zurich for their hospitality. I.B. thanks CERN for hospitality. The work of I.B. is supported by a EURYI award of the European Science Foundation.

Appendix A \mathbb{Z}_2 -graded vector spaces and their dual

Let us consider a \mathbb{Z}_2 -graded complex vector space $V = V_+ \oplus V_-$ with involution $\sigma : V \rightarrow V$ that has eigenvalue ± 1 on V_\pm . An element $v \in V$ has degree $|v|$ so that $(\sigma)^{|v|} v = \pm v$.¹³ The dual vector space V^* is defined through the dual pairing $f; v$ for $v \in V$ and $f \in V^*$. It is non-vanishing for $|f| + |v| = m$. The pairing is called even/odd if m is even/odd.

The grading on V naturally induces a grading on the vector space of homomorphisms, $\text{Hom}(V_1; V_2) = \text{Hom}_+(V_1; V_2) \oplus \text{Hom}_-(V_1; V_2)$. For an element $M \in \text{Hom}(V_1; V_2)$ of definite degree we denote the degree by $|M|$ and we have

$$\sigma^{|M|} M = (\sigma)^{|M|} M :$$

The graded transpose

To an element $M \in \text{Hom}(V_1; V_2)$ we can associate a dual homomorphism in $\text{Hom}(V_2; V_1)$, the graded transpose M^T , via

$$(113) \quad M^T f; v_{V_1} = (\sigma)^{|f| - (|M| + |v|)} f; M v_{V_2} :$$

¹³In the main text the degree corresponds to the R-charge.

Here $v \in V_1$ and $f \in V_2$. In an even/odd basis, in which $\epsilon_1 = \epsilon_2 = \text{diag}(\text{id}; -\text{id})$, a homomorphism and its graded transpose are¹⁴

$$(114) \quad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}; \quad M^T = \begin{pmatrix} a^t & c^t \\ b^t & d^t \end{pmatrix};$$

where t is the ordinary transposition of matrices. In view of the shift m in grading between the vector space and its dual, the involution on the dual vector space V is $(-1)^{\epsilon_1} \tau$.

Let us summarize some properties for the graded transpose that are useful for the main part of this work. For compositions of homomorphisms we have

$$(115) \quad (AB)^T = (-1)^{\epsilon_A \epsilon_B} B^T A^T;$$

Its behaviour with respect to matrix inversion is

$$(116) \quad (M^T)^{-1} = {}^T(M^{-1})^T = (-1)^{\epsilon_M} (M^{-1})^T$$

For even homomorphisms we do not pick up a sign on the right-hand side and we can use the abbreviation $M^{-T} := (M^T)^{-1}$ unambiguously. The hermitian conjugation on the dual space is defined by requiring that hermitian conjugation commutes with the graded transpose,

$$(117) \quad (M^T)^y := (M^y)^T;$$

Double transpose

The double dual V^y of a vector space V is canonically isomorphic to V via the canonical isomorphism $e: V^y \rightarrow V$ defined by $e(v); f_v := f; v_v$.¹⁵ In the following and in the main part of this work we do not explicitly write out this isomorphism.

The double transpose of a homomorphism $M: V_1 \rightarrow V_2$ acts via the canonical isomorphism as $M^{TT}: V_1 \rightarrow V_2$. Let us determine its relation to M ,

$$\begin{aligned} M^{TT} v; f_v &= (-1)^{\epsilon_M} (j^y j^m) v; M^T f_v = \\ &= (-1)^{\epsilon_M} (j^y j^m) M^T f; v_v = \\ &= (-1)^{\epsilon_M} (j^y j^x j^j) f; M v_v = \\ &= (-1)^{\epsilon_M} (j^y j^x j^j) M v; f_v : \end{aligned}$$

¹⁴The slightly non-standard definition of the graded transpose in (113), including the sign $(-1)^{\epsilon_M \epsilon_n}$, ensures that M^T has the same form for both m even and odd.

¹⁵Note that this isomorphism is defined without sign as compared to [17], and therefore $\{e_{\text{here}} = e_{\text{there}}\}$.

We therefore find, using $\sum_{j=1}^m j = \frac{m(m+1)}{2}$, that

$$(118) \quad M^T = \begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^{\frac{m+1}{2}} M = \begin{pmatrix} m+1 & \dots & 1 \end{pmatrix}^{\frac{m+1}{2}} :$$

Alternatively, this can be seen directly with (114), keeping in mind that the grading operator on the dual vector space is $\begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^T$.

Graded tensor products

The graded tensor product, $V = V_1 \otimes V_2$ can be defined for endomorphisms A and B in terms of the ordinary (non-graded) tensor product,

$$(A \otimes B) := A \otimes_2 B :$$

The grading operator on the right-hand side ensures the multiplication rule

$$(A \otimes B)(C \otimes D) = \begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^{\otimes 2} (AC) \otimes (BD) :$$

However, the graded transpose is not the naive one, an explicit computation in the even/odd basis reveals

$$(119) \quad (A \otimes B)^T = A^T \begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^{\otimes 2} \otimes \begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^{\otimes 2} B^T = A^T \begin{pmatrix} 1 & \dots & 1 \end{pmatrix}^{\otimes 2} \otimes B^T :$$

References

- [1] A. Sagnotti, "Open strings and their symmetry groups," arXiv:hep-th/0208020.
- [2] P. Horava, "Strings on World Sheet Orbifolds," Nucl. Phys. B 327 (1989) 461.
- [3] J. Dai, R. G. Leigh and J. Polchinski, "New Connections Between String Theories," Mod. Phys. Lett. A 4 (1989) 2073.
- [4] M. Bianchi and A. Sagnotti, "On the systematics of open string theories," Phys. Lett. B 247 (1990) 517.
- [5] M. Bianchi and A. Sagnotti, "Twist symmetry and open string Wilson lines," Nucl. Phys. B 361 (1991) 519.
- [6] E. G. Gimon and J. Polchinski, "Consistency Conditions for Orientifolds and D-Manifolds," Phys. Rev. D 54 (1996) 1667 [arXiv:hep-th/9601038].
- [7] J. Polchinski, "Dirichlet-Branes and Ramond-Ramond Charges," Phys. Rev. Lett. 75 (1995) 4724 [arXiv:hep-th/9510017].
- [8] E. Witten, "Phases of $N = 2$ theories in two dimensions," Nucl. Phys. B 403, 159 (1993) [arXiv:hep-th/9301042].
- [9] E. Witten, "Toroidal compactification without vector structure," JHEP 9802 (1998) 006 [arXiv:hep-th/9712028].

- [10] M. Bianchi, \A note on toroidal compactifications of the type I superstring and other superstring vacuum configurations with 16 supercharges," Nucl. Phys. B 528 (1998) 73 [arXiv:hep-th/9711201].
- [11] I. Pesando, \A comment on discrete Kalb-Ramond field on orientifold and rank reduction," Phys. Lett. B 668 (2008) 324 [arXiv:0804.3931 [hep-th]].
- [12] C. Bachas, M. Bianchi, R. Blumenhagen, D. Lust and T. Weigand, \Comments on Orientifolds without Vector Structure," JHEP 0808 (2008) 016 [arXiv:0805.3696 [hep-th]].
- [13] I. Brunner and K. Hori, \Orientifolds and mirror symmetry," JHEP 0411 (2004) 005 [arXiv:hep-th/0303135].
- [14] I. Brunner, K. Hori, K. Hosomichi and J. Walcher, \Orientifolds of Gepner models," JHEP 0702 (2007) 001 [arXiv:hep-th/0401137].
- [15] K. Hori, D. Gao, to appear. See also <http://online.kitp.ucsb.edu/online/strings05/hori>
- [16] J. Distler, D. Freed and G. W. Moore, to appear. See also <http://www.phys.ethz.ch/~distler/mrg/Prestrings>
- [17] K. Hori and J. Walcher, \D-brane categories for orientifolds: The Landau-Ginzburg case," JHEP 0804 (2008) 030 [arXiv:hep-th/0606179].
- [18] J. Bertin and F. Rosay, \Periodicite de Knorrer etendue," Comptes Rendus Mathematique 345, 4 (2007) 187 [arXiv:math/0609400].
- [19] D. E. Diaconescu, A. Garcia-Raboso, R. L. Karp and K. Sinha, \D-brane superpotentials in Calabi-Yau orientifolds (projection)," arXiv:hep-th/0606180.
- [20] M. Herbst, K. Hori and D. Page, \Phases Of N=2 Theories In 1+1 Dimensions With Boundary," arXiv:0803.2045 [hep-th].
- [21] T. Bridgeland, \Flops and derived categories," Invent. Math. 147, 3 (2002) 613 [arXiv:math/0009053].
- [22] T. Bridgeland, A. King and M. Reid, \The McKay Correspondence As An Equivalence Of Derived Categories," J. Am. Math. Soc. 14, 535 (2001).
- [23] M. van den Bergh, \Non-commutative crepant resolutions," in \The legacy of Niels Henrik Abel," 749{770, Springer, Berlin, 2004, [arXiv:math/0211064].
- [24] Y. Kawamata, \Log Crepant Birational Maps and Derived Categories," [arXiv:math/0311139].
- [25] D. Orlov, \Derived categories of coherent sheaves and triangulated categories of singularities," [arXiv:math/0503632].

- [26] P. S. Aspinwall, \The Landau-Ginzburg to Calabi-Yau dictionary for D-branes," *J. Math. Phys.* 48 (2007) 082304 [arXiv:hep-th/0610209].
- [27] I. Brunner, M. R. Douglas, A. E. Lawrence and C. Romelsberger, \D-branes on the quintic," *JHEP* 0008 (2000) 015 [arXiv:hep-th/9906200].
- [28] S. G. ovindarajan and T. Jayaraman, \D-branes, exceptional sheaves and quivers on Calabi-Yau manifolds: From Mukai to McKay," *Nucl. Phys. B* 600 (2001) 457 [arXiv:hep-th/0010196];
- [29] P. Mayr, \Phases of supersymmetric D-branes on Kähler manifolds and the McKay correspondence," *JHEP* 0101 (2001) 018 [arXiv:hep-th/0010223];
- [30] T. Takayanagi, \Holomorphic tachyons and fractional D-branes," *Nucl. Phys. B* 603 (2001) 259 [arXiv:hep-th/0103021].
- [31] T. Takayanagi, \Tachyon condensation on orbifolds and McKay correspondence," *Phys. Lett. B* 519 (2001) 137 [arXiv:hep-th/0106142].
- [32] K. Becker, M. Becker, C. Vafa and J. Walcher, \Moduli stabilization in non-geometric backgrounds," *Nucl. Phys. B* 770 (2007) 1 [arXiv:hep-th/0611001].
- [33] K. Becker, M. Becker and J. Walcher, \Runaway in the Landscape," *Phys. Rev. D* 76 (2007) 106002 [arXiv:0706.0514 [hep-th]].
- [34] P. Alu and M. Esole, \Chemical identities from tadpole matching in type IIB and F-theory," arXiv:0710.2544 [hep-th].
- [35] A. P. Braun, A. Hebecker and H. Triendl, \D7-Brane Motion from M-Theory Cycles and Obstructions in the Weak Coupling Limit," *Nucl. Phys. B* 800 (2008) 298 [arXiv:0801.2163 [hep-th]].
- [36] A. Collinucci, F. Denef and M. Esole, \D-brane D reconstructions in IIB Orientifolds," arXiv:0805.1573 [hep-th].
- [37] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y. S. Stanev, \Comments on Gepner models and type I vacua in string theory," *Phys. Lett. B* 387 (1996) 743 [arXiv:hep-th/9607229].
- [38] R. Blumenhagen and A. Wiskirchen, \Spectra of 4D, $N = 1$ type I string vacua on non-toroidal CY threefolds," *Phys. Lett. B* 438 (1998) 52 [arXiv:hep-th/9806131].
- [39] G. Aldazabal, E. C. Andres, M. Leston and C. Nunez, \Type IIB orientifolds on Gepner points," *JHEP* 0309 (2003) 067 [arXiv:hep-th/0307183].
- [40] R. Blumenhagen and T. Weigand, \Chiral supersymmetric Gepner model orientifolds," *JHEP* 0402 (2004) 041 [arXiv:hep-th/0401148].

- [41] T. P. T. Dijkstra, L. R. Huiszoon and A. N. Schellekens, "Supersymmetric standard model spectra from RCFT orientifolds," Nucl. Phys. B 710 (2005) 3 [arXiv:hep-th/0411129].
- [42] K. Hosomichi, "Permutation orientifolds of Gepner models," JHEP 0701 (2007) 081 [arXiv:hep-th/0612109].
- [43] J. Walcher, "Evidence for Tadpole Cancellation in the Topological String," arXiv:0712.2775 [hep-th].
- [44] J. Walcher, "Opening mirror symmetry on the quintic," Commun. Math. Phys. 276 (2007) 671 [arXiv:hep-th/0605162].
- [45] D. R. Morrison and J. Walcher, "D-branes and Normal Functions," arXiv:0709.4028 [hep-th].
- [46] A. Strominger, "Massless black holes and conifolds in string theory," Nucl. Phys. B 451 (1995) 96 [arXiv:hep-th/9504090].
- [47] K. Hori, K. Hosomichi, D. C. Page, R. Rabadan and J. Walcher, "Non-perturbative orientifold transitions at the conifold," JHEP 0510 (2005) 026 [arXiv:hep-th/0506234].
- [48] K. Hori et al., "Mirror symmetry," SPIRES entry Providence, USA: AMS (2003) 929 p
- [49] N. P. Warner, "Supersymmetry in boundary integrable models," Nucl. Phys. B 450 (1995) 663 [arXiv:hep-th/9506064].
- [50] M. Kontsevich, unpublished.
- [51] D. Orlov, "Triangulated categories of singularities and D-branes in Landau-Ginzburg models," arXiv:math/0302304.
- [52] A. Kapustin and Y. Li, "D-branes in Landau-Ginzburg models and algebraic geometry," JHEP 0312 (2003) 005 [arXiv:hep-th/0210296].
- [53] I. Brunner, M. Herbst, W. Lerche and B. Scheuner, "Landau-Ginzburg realization of open string TFT," JHEP 0611 (2006) 043 [arXiv:hep-th/0305133].
- [54] A. Kapustin and Y. Li, "Topological Correlators in Landau-Ginzburg Models with Boundaries," Adv. Theor. Math. Phys. 7 (2004) 727 [arXiv:hep-th/0305136].
- [55] D. Quillen, "Superconnections and the Chern character," Topology 24 (1985) 89.
- [56] P. Kraus and F. Larsen, "Boundary string field theory of the DD-bar system," Phys. Rev. D 63 (2001) 106004 [arXiv:hep-th/0012198].
- [57] T. Takayanagi, S. Terashima and T. Uesugi, "Brane-antibrane action from boundary string field theory," JHEP 0103 (2001) 019 [arXiv:hep-th/0012210].

- [58] P. S. Aspinwall, "D-branes on Calabi-Yau manifolds," *arXiv:hep-th/0403166*.
- [59] J. Shamash, "The Poincare series of a local ring," *J. Algebra* 12 453 (1969).
- [60] D. Eisenbud, "Homological Algebra on a Complete Intersection, with an Application to Group Representations," *Transactions of the American Mathematical Society*, 260, 35 (1980), No. 1
- [61] L. L. Avramov, R. O. Buchweitz, "Homological algebra modulo a regular sequence with special attention to codimension two," *J. Algebra*, 230, 1 (2000) 24.
- [62] L. L. Avramov, D. R. Grayson, "Resolutions and cohomology over complete intersections," in *Computations in algebraic geometry with Macaulay 2*, 131-178, *Algorithms Comput. Math.*, 8, Springer, Berlin, 2002.
- [63] P. S. Aspinwall, "Topological D-Branes and Commutative Algebra," *arXiv:hep-th/0703279*.
- [64] H. K. Norrner, "Cohen-Macaulay modules on hypersurface singularities. I," *Invent. Math.* 88 (1987) 153.
- [65] D. R. Morrison and M. R. Plesser, "Summing the instantons: Quantum cohomology and mirror symmetry in toric varieties," *Nucl. Phys. B* 440 (1995) 279 [*arXiv:hep-th/9412236*].
- [66] M. R. Douglas and G. W. Moore, "D-branes, Quivers, and ALG Instantons," *arXiv:hep-th/9603167*.
- [67] M. R. Douglas, "D-branes, categories and $N = 1$ supersymmetry," *J. Math. Phys.* 42 (2001) 2818 [*arXiv:hep-th/0011017*].
- [68] A. Sen, "Tachyon condensation on the brane antibrane system," *JHEP* 9808, 012 (1998) [*arXiv:hep-th/9805170*].
- [69] E. Witten, "D-branes and K-theory," *JHEP* 9812 (1998) 019 [*arXiv:hep-th/9810188*].
- [70] K. Hori, "Linear models of supersymmetric D-branes," *arXiv:hep-th/0012179*.
- [71] S. K. Ashok, E. Dell'Aquila and D. E. Diaconescu, "Fractional branes in Landau-Ginzburg orbifolds," *Adv. Theor. Math. Phys.* 8 (2004) 461 [*arXiv:hep-th/0401135*];
- [72] S. K. Ashok, E. Dell'Aquila, D. E. Diaconescu and B. Florea, "Obstructed D-branes in Landau-Ginzburg orbifolds," *Adv. Theor. Math. Phys.* 8 (2004) 427 [*arXiv:hep-th/0404167*];
- [73] K. Hori and J. Walcher, "F-term equations near Gepner points," *JHEP* 0501 (2005) 008 [*arXiv:hep-th/0404196*];

- [74] K. Hori and J. Walcher, "D-branes from matrix factorizations," *Comptes Rendus Physique* 5 (2004) 1061 [arXiv:hep-th/0409204];
- [75] I. Brunner, M. Herbst, W. Lerche and J. Walcher, "Matrix factorizations and mirror symmetry: The cubic curve," *JHEP* 0611 (2006) 006 [arXiv:hep-th/0408243];
- [76] I. Brunner and M. R. Gaberdiel, "Matrix factorisations and permutation branes," *JHEP* 0507 (2005) 012 [arXiv:hep-th/0503207];
- [77] B. Ezhuthachan, S. Govindarajan and T. Jayaraman, "A quantum McKay correspondence for fractional 2p-branes on LG orbifolds," *JHEP* 0508 (2005) 050 [arXiv:hep-th/0504164];
- [78] H. Enger, A. Recknagel and D. Roggenkamp, "Permutation branes and linear matrix factorisations," *JHEP* 0601 (2006) 087 [arXiv:hep-th/0508053];
- [79] C. Caviezel, S. Fredenhagen and M. R. Gaberdiel, "The RR charges of A-type Gepner models," *JHEP* 0601 (2006) 111 [arXiv:hep-th/0511078];
- [80] S. Govindarajan, H. Jockers, W. Lerche and N. P. Warner, "Tachyon condensation on the elliptic curve," *Nucl. Phys. B* 765 (2007) 240 [arXiv:hep-th/0512208];
- [81] S. Fredenhagen and M. R. Gaberdiel, "Generalised $N = 2$ permutation branes," *JHEP* 0611 (2006) 041 [arXiv:hep-th/0607095];
- [82] C. Schmidt-Colinet, "Tensor product and permutation branes on the torus," *JHEP* 0707 (2007) 044 [arXiv:hep-th/0701128].
- [83] J. Knapp and E. Scheidegger, "Matrix Factorizations, Massey Products and F-Terms for Two-Parameter Calabi-Yau Hypersurfaces," arXiv:0812.2429 [hep-th].
- [84] K. A. Intriligator, "RG fixed points in six dimensions via branes at orbifold singularities," *Nucl. Phys. B* 496 (1997) 177 [arXiv:hep-th/9702038].
- [85] J. D. Blum and K. A. Intriligator, "Consistency conditions for branes at orbifold singularities," *Nucl. Phys. B* 506 (1997) 223 [arXiv:hep-th/9705030].
- [86] J. D. Blum and K. A. Intriligator, "New phases of string theory and 6d RG fixed points via branes at orbifold singularities," *Nucl. Phys. B* 506 (1997) 199 [arXiv:hep-th/9705044].
- [87] I. Brunner and A. Karch, "Branes at orbifolds versus Hanany Witten in six dimensions," *JHEP* 9803 (1998) 003 [arXiv:hep-th/9712143].
- [88] A. M. Urange, "A new orientifold of $C^2 = Z(N)$ and six-dimensional RG fixed points," *Nucl. Phys. B* 577 (2000) 73 [arXiv:hep-th/9910155].

- [89] S. Franco, A. Hanany, D. Kremer, J. Park, A. M. Uranga and D. Vegh, "Dimers and Orientifolds," JHEP 0709 (2007) 075 [arXiv:0707.0298 [hep-th]].
- [90] I. Brunner, H. Jockers and D. Roggenkamp, "D effects and D-Brane Monodromies," arXiv:0806.4734 [hep-th].
- [91] A. Sen and S. Sethi, "The mirror transform of type I vacua in six dimensions," Nucl. Phys. B 499 (1997) 45 [arXiv:hep-th/9703157].
- [92] A. Sen, "Orientifold limit of F-theory vacua," Phys. Rev. D 55 (1997) 7345 [arXiv:hep-th/9702165].