

# Nucleon Statistics in Holographic QCD : Aharonov-Bohm Effect in a Matrix Model

Koji HASHIMOTO<sup>1,\*</sup> and Norihiro IIZUKA<sup>2,†</sup>

<sup>1</sup> *Mathematical Physics Lab., RIKEN Nishina Center, Saitama 351-0198, Japan*

<sup>2</sup> *Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

We show that the Aharonov-Bohm effect in the nuclear matrix model [1] derives the statistical nature of nucleons in holographic QCD. For  $N_c = \text{odd}$  (even), the nucleon is shown to be a fermion (boson).

The statistics of baryons depends on the number of colors in QCD; in particular for large  $N_c$  QCD, as the baryons are bound states of  $N_c$  quarks, they are fermions for odd  $N_c$ , while bosons for even  $N_c$ . The nuclear matrix model [1] derived in holographic QCD offers a simple effective description of multi-baryon systems, where we can compute baryon spectra, short-distance nuclear forces, and even three-body nuclear forces [2]. However, since the nuclear matrix model has only bosonic variables, it is natural to ask how the fermionic nature of baryons comes out from the matrix model. In chiral soliton models, this question was answered from the properties of Wess-Zumino term [3].

To identify the statistics (fermionic/bosonic) of nucleons in the nuclear matrix model, we consider a  $2\pi$  rotation in the target space of the matrix model. The target space index is carried by  $X^M$  and  $w_{\dot{\alpha}i}$ . The effect on  $X^M$  is trivial, since  $X$  decouples from the system in the matrix model for a single baryon ( $k = 1$ ) once the ADHM constraint is solved. However, since we have a nontrivial gauge field  $A_0$ , there is a nontrivial effect on the  $w_{\dot{\alpha}i}$  sector. In fact, this gauge field  $A_0$  turns out to be responsible for the statistics of the baryons, as we will see.

In a pion effective lagrangian, it is known that the Wess-Zumino term is essential for showing the nucleon statistics, in the picture of solitonic nucleon of the system [3]. Now, in holographic QCD, this Wess-Zumino term is known to be from the 4-form Ramond-Ramond flux in the gravity background in the D4-D8 model of holographic QCD [4]. In the nuclear matrix model [1], the Ramond-Ramond flux generates a Chern-Simons term in 1 dimension, which is just a term consisting of a single gauge field  $A_0$ . The  $w_{\dot{\alpha}i}$  field is charged under the gauge symmetry, so it is natural to expect that the gauge dynamics in this 1 dimension with the Chern-Simons term gives the nucleon statistics.

In the nuclear matrix model, the terms including the fundamental field  $w_{\dot{\alpha}i}$ , except for the ADHM potential terms and the mass term, are

$$S = \frac{\lambda N_c M_{\text{KK}}}{54\pi} \int dt D_0 \bar{w}_i^{\dot{\alpha}} D_0 w_{\dot{\alpha}i} + N_c \int dt A_0. \quad (1)$$

$\dot{\alpha}$  is a spinor index which is for  $SU(2) \simeq SO(3)$  spatial rotation in the target space.  $i = 1, \dots, N_f$  is a flavor index. This is a one-dimensional gauge theory whose gauge field is  $A_0$ . The covariant derivative is defined as  $D_0 w_{\dot{\alpha}i} \equiv \partial_0 w_{\dot{\alpha}i} - i w_{\dot{\alpha}i} A_0$ . Note that  $A_0$  is a gauge field for  $U(k)$  gauge symmetry of the matrix model, so, for  $k = 1$  (single baryon),  $A_0$  does not carry any non-Abelian index.

Let us make a spatial rotation, for example along the  $x^3$  axis, by an angle  $2\pi$ . We look at how a wave function of a baryon transforms under this rotation, and if it acquires a phase  $n\pi$  with an odd integer  $n$ , *i.e.* it changes a sign, then the state is determined to be a fermion.

Since  $w_{\dot{\alpha}i}$  carries a spinor index of the target space, it is obvious that the spatial rotation acts for the case of the rotation around the  $x^3$  axis with an angle  $\theta$ , as

$$w_{\dot{\alpha}i} \rightarrow U_{\dot{\alpha}}^{\dot{\beta}} w_{\dot{\beta}i}, \quad U = \exp \left[ i \frac{\theta}{2} \tau^3 \right]. \quad (2)$$

Here  $\tau^3$  is the third component of Pauli matrices. Our spatial rotation by  $2\pi$  means that the angle  $\theta$  moves in the period  $0 \leq \theta \leq 2\pi$ .

As shown in [1], the vacuum of the matrix model for  $k = 1$  is quite simple,

$$w = \begin{pmatrix} \rho_0 & 0 \\ 0 & \rho_0 \end{pmatrix}. \quad (3)$$

After minimizing the hamiltonian, we obtain a certain nonzero value for this  $\rho_0$ . So the spatial rotation (2) corresponds to a certain path in the target space of  $w_{\dot{\alpha}i}$ . In the following, we would like to compute an Aharonov-Bohm phase with this path. For that, it is inconvenient that two nonzero entries in (3) moves simultaneously. So, we combine a gauge transformation  $\exp[-i\theta/2]$  together with the spatial rotation (2), so that we find a path

$$w_{\dot{\alpha}i} \rightarrow U_{\dot{\alpha}}^{\dot{\beta}} w_{\dot{\beta}i}, \quad U = \begin{pmatrix} 1 & 0 \\ 0 & e^{-i\theta} \end{pmatrix}. \quad (4)$$

With this, we find that only the lower-right corner of  $w_{\dot{\alpha}i}$  in (3) rotates. Indeed, the same change of the parameterization of the path was used in [3] for the soliton in the pion effective field theory.

We are interested in a phase change of a baryon wave function. The argument of the wave function is the moduli of this matrix model, and it is a part of  $w_{\dot{\alpha}i}$  configuration space. If we think of the path of  $w_{\dot{\alpha}i}$  defined by

\*koji(at)riken.jp

†norihiro.iizuka(at)cern.ch

(4), then the phase of the wave function of our concern is in fact an Aharonov-Bohm (AB) phase, for the path (4), as if we regard the lower-right entry of the matrix field  $w_{\hat{\alpha}i}$  as a position of a charged particle.

Let us write down the lagrangian for this charged particle, to compute the AB phase. Writing the lower-right component of  $w_{\hat{\alpha}i}$  as  $w_{\hat{\alpha}=2,i=2} = u + iv$  where  $u$  and  $v$  are real, then the relevant part of the matrix model is

$$S = \frac{\lambda N_c M_{\text{KK}}}{54\pi} \int dt |\partial_t(u + iv) - iA_0(u + iv)|^2. \quad (5)$$

It was shown in [1] that solving the equation of motion for  $A_0$ , gives  $A_0 = -27\pi/2\lambda M_{\text{KK}}\rho_0^2$ , which is a real constant. Then the action (5) can be rewritten with conjugate momenta in real coordinates as

$$S = \frac{1}{2M} \int dt [(P_u + A_0 v M)^2 + (P_v - A_0 u M)^2]. \quad (6)$$

Here we have defined the ‘‘mass’’  $M$  of the hypothetical particle moving in the  $u$ - $v$  space as  $M = \lambda N_c M_{\text{KK}}/27\pi$ . The expression shows that the particle is in a minimally-coupled gauge potential in the  $u$ - $v$  space, defined by

$$\tilde{A}_u \equiv -A_0 M v = \frac{N_c}{2\rho_0^2} v, \quad \tilde{A}_v \equiv A_0 M u = -\frac{N_c}{2\rho_0^2} u. \quad (7)$$

The magnetic flux made by this gauge potential is constant.

The path of this hypothetical charged particle is given by (4), which is

$$u + iv = \rho_0 e^{-i\theta} \quad (0 \leq \theta \leq 2\pi) \quad (8)$$

so the circle encloses the area  $\pi\rho_0^2$ , in a counter-clockwise way. The AB phase  $\Phi$  is given by an integration of the gauge potential (7) along this path,

$$\Phi = -\rho_0 \oint \tilde{A}_\theta d\theta = N_c \pi. \quad (9)$$

In the last equality, we have used (7) in a polar coordinate,  $\tilde{A}_\theta = -N_c/2\rho_0$ . The negative sign is from the orientation of the path.

This AB phase means that, when  $N_c$  is odd, the spatial rotation by the angle  $2\pi$  results in a sign  $(-1)$  multiplied to the baryon wave function. Therefore, when  $N_c$  is odd (even), the baryon is a fermion (boson).

It is intriguing that a simple mechanism, the AB phase, is encoded in the nuclear matrix model naturally to ensure the baryon statistics in holographic QCD.

— **Acknowledgment.** We would like to thank Y. Tachikawa for bringing us a question, which motivated the present work. K.H. is partly supported by the Japan Ministry of Education, Culture, Sports, Science and Technology.

- 
- [1] K. Hashimoto, N. Iizuka and P. Yi, ‘‘A Matrix Model for Baryons and Nuclear Forces,’’ arXiv:1003.4988 [hep-th].  
 [2] K. Hashimoto and N. Iizuka, ‘‘Three-Body Nuclear Forces from a Matrix Model,’’ arXiv:1005.4412 [hep-th].  
 [3] E. Witten, ‘‘Current Algebra, Baryons, And Quark Con-

- finement,’’ Nucl. Phys. B **223**, 433 (1983).  
 [4] T. Sakai and S. Sugimoto, ‘‘Low energy hadron physics in holographic QCD,’’ Prog. Theor. Phys. **113**, 843 (2005) [arXiv:hep-th/0412141].