



LHCb Upgrade Plans



Franz Muheim
University of Edinburgh
on behalf of the LHCb collaboration

Standard Model and New Physics Sensitivity

LHCb Experiment

Physics Programme the first 5 years

Running LHCb at 10 times design luminosity

Physics Reach with a 100 fb^{-1} data sample

CP violation in B_s decays

Probe New Physics in hadronic
and electroweak penguin decays

CKM angle γ

LHCb Upgrade Detector and Trigger Plans

LHCb Upgrade Detector

Vertex detector studies


Trigger and Read-out studies

Conclusions


Flavour in the LHC era
CERN Oct 9th 2006

Status of CKM Unitarity Triangles

- **ICHEP2006 Status**
 - including CDF Δm_s measurement



- **Tree diagrams**
 - Not sensitive to New Physics



- **Probe New Physics**
 - by comparing to SM predictions including loops
 - by measuring γ in loop diagrams
 - same for α , β and χ

- Standard Model is a very successful theory
- *We are very likely beyond the era of « alternatives » to the CKM picture. NP would appear as « corrections » to the CKM picture* Nir


Probing New Physics in B_s Mesons

- Flavour Changing Neutral Currents**


- NP appears as virtual particles in loop processes
- leading to observable deviations from SM expectations in flavour physics and CP violation (~~CP~~)
- New Physics parameterisation in B_s Oscillations

$$\Delta m_q = \left| 1 + h_q e^{2i\sigma_q} \right| \Delta m_q^{SM}$$

New Physics



$B_s \rightarrow \phi\phi$ penguin decay



$B_s - \bar{B}_s$ oscillations

- If New Physics is found at LHC**

- Probe NP flavour structure with FCNC

LHCb Sensitivities with 2 fb^{-1}



	Channel	Yield	B/S	Precision
γ	$B_s \rightarrow D_s^{-+} K^{+-}$	5.4k	< 1.0	$\sigma(\gamma) \sim 14^\circ$
	$B_d \rightarrow \pi^+ \pi^-$	36k	0.46	$\sigma(\gamma) \sim 4^\circ$
	$B_s \rightarrow K^+ K^-$	36k	< 0.06	
	$B_d \rightarrow D^0 (K\pi, KK) K^{*0}$	3.4 k, 0.5 k, 0.6 k	$< 0.3, < 1.7,$ < 1.4	$\sigma(\gamma) \sim 7^\circ - 10^\circ$
	$B^- \rightarrow D^0 (K^- \pi^+, K^+ \pi^-) K^-$	28k, 0.5k	0.6, 1.5	$\sigma(\gamma) \sim 5^\circ - 15^\circ$
	$B^- \rightarrow D^0 (K^+ K^-, \pi^+ \pi^-) K^-$	4.3 k	1.0	
	$B^- \rightarrow D^0 (K_S \pi^+ \pi^-) K^-$	1.5 - 5k	< 0.7	$\sigma(\gamma) \sim 8^\circ - 16^\circ$
α	$B_d \rightarrow \pi^+ \pi^- \pi^0$	14k	< 0.8	$\sigma(\alpha) \sim 10^\circ$
	$B \rightarrow \rho^+ \rho^0, \rho^+ \rho^-, \rho^0 \rho^0$	9k, 2k, 1k	1, $< 5, < 4$	
β	$B_d \rightarrow J/\psi(\mu\mu)K_S$	216k	0.8	$\sigma(\sin 2\beta) \sim 0.022$
Δm_s	$B_s \rightarrow D_s^- \pi^+$	120k	0.4	$\sigma(\Delta m_s) \sim 0.01 \text{ ps}^{-1}$
ϕ_s	$B_s \rightarrow J/\psi(\mu\mu)\phi$	131k	0.12	$\sigma(\phi_s) \sim 0.023$
Rare decays	$B_s \rightarrow \mu^+ \mu^-$	17	< 5.7	$\sigma(C_7^{\text{eff}}/C_9^{\text{eff}}) \sim 0.13$
	$B_d \rightarrow K^{*0} \mu^+ \mu^-$	4.4 k	< 2.6	
	$B_d \rightarrow K^{*0} \gamma$	35k	< 0.7	$\sigma(A_{\text{CP}}) \sim 0.01$
	$B_s \rightarrow \phi \gamma$	9.3 k	< 2.4	
charm	$D^{*+} \rightarrow D^0 (K^- \pi^+) \pi^+$	100 M		

Alessio Sarti

Frederic Teubert

Ulrik Egede

Patrick Spradlin


LHCb – The First Five Years

• LHCb Operations

- Luminosity tuneable by adjusting beam focus
- Design is to run at $\mathcal{L} \sim 2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ detectors up to $5 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
- little pile-up ($n = 0.5$)
- less radiation damage
- Luminosity will be achieved during 1st physics run

• LHCb Physics Goals

- Run five (nominal) years at $\mathcal{L} \sim 2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ and collect 6 to 10 fb⁻¹
- Exploit the B_s system
- Observation of CP violation in B_s mesons
- Precision measurements of B_s mass and lifetime difference
- Reduce error on CKM angle γ by a factor 5
- Probe New Physics in rare B meson decays with electroweak, radiative and hadronic penguin modes
- First observation of very rare decay $B_s \rightarrow \mu^+ \mu^-$



Physics Case for LHCb at High Luminosity



- **What's next?**
 - Many LHCb results will be statistically limited
 - New Physics effects are small -> require better precision measurements
 - LHCb is only B-physics experiment approved for running after 2010
 - Can LHCb exploit the full potential of B physics at hadron colliders?
- **LHCb Luminosity**
 - Running at $\mathcal{L} \sim 2 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$ is a LHCb design choice
 - LHC design luminosity is 50 times higher $\mathcal{L} \sim 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- **LHCb Upgrade Plans**
 - Upgrade LHCb detector such that it can operate at 10 times design luminosity of $\mathcal{L} \sim 2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$
 - Run ~5 yrs at $\mathcal{L} \sim 2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$
 - **Collect ~100 fb⁻¹ data sample**
 - Multiple interactions per beam crossing increases to $n \sim 4$
 - Is compatible with possible LHC luminosity upgrade (SLHC)
 - Does not require SLHC
 - Could be implemented ~2013

ϕ_s from $B_s \rightarrow J/\psi\phi$

• CP Violation in B_s mesons

- Interference in B_s mixing and decay
- ~~CP~~ B_s weak mixing phase ϕ_s is very small in SM

$$\phi_s = -\arg(V_{ts}^2) = -2\chi \approx -2\lambda^2\eta \approx -0.035$$


- \Rightarrow sensitive probe for **New Physics** e.g. stringent NMFV test
- NP parameterisation

$$\Delta m_q = \left| 1 + h_q e^{2i\sigma_q} \right| \Delta m_q^{SM}$$


- Angular analysis to separate $J/\psi\phi$ 2 CP-even and 1 CP-odd amplitudes

• ϕ_s Sensitivity

- at $\Delta m_s = 20 \text{ ps}^{-1}$
- Expect 131k $B_s \rightarrow J/\psi\phi$ signal events per 2 fb^{-1} (1 year)
- Expected precision $\sigma(\sin \phi_s) \sim 0.023$
- Small improvement in ϕ_s precision by adding pure CP modes



hep-ph/0604112
hep-ph/0509242



ϕ_s from $B_s \rightarrow J/\psi\phi$

- ϕ_s will be the ultimate SM test

- For ~~CP~~ in B mesons
- Similar to ε' in kaons for direct CP violation

- ϕ_s Sensitivity

- LHCb for 10 fb^{-1} (first 5 years)
- $\sim 3 \sigma$ SM evidence for $\phi_s \approx -0.035$
- ϕ_s precision statistically limited
- Theoretically clean

$$\sigma(\sin \phi_s) \sim 0.010$$

- Historical Aside

- 1988 NA31 measures $\sim 3 \sigma$ from zero
- Community approves NA48 & KTeV

$$\varepsilon'/\varepsilon = (3.3 \pm 1.1) 10^{-3}$$

- LHCb Upgrade Sensitivities

- Based on 100 fb^{-1} data sample
- Preliminary estimates by scaling with luminosity
- Potential trigger efficiency improvements not included

- $B_s \rightarrow J/\psi\phi$ - Key channel for LHCb Upgrade


- ϕ_s Sensitivity with 100 fb^{-1} data sample
- $\sim 10 \sigma$ SM measurement with 100 fb^{-1}

$$\sigma(\sin \phi_s) \sim 0.003$$

b → s Transitions in B_d Mesons




$\sin(2\beta^{\text{eff}}) \equiv \sin(2\phi_1^{\text{eff}})$ **HFAG**
ICHEP 2006
PRELIMINARY



- Compare $\sin 2\beta$ measurements
 - in $B_d \rightarrow \phi K_S$ with $B_d \rightarrow J/\psi K_S$
 - Individually, each decay mode in reasonable agreement with SM
 - But all measurements lower than $\sin 2\beta$ from
- Naïve $b \rightarrow s$ penguin average
 - $\sin 2\beta_{\text{eff}} = 0.52 \pm 0.05$
 - 2.6 σ discrepancy from SM
- Theory models
 - Predict to increase $\sin 2\beta_{\text{eff}}$ in SM


some of recent QCDF estimates



PLB 620 (2005) 143
hep-ph/0506268

$b \rightarrow s$ Transitions in $B_s \rightarrow \phi\phi$

- $B_s \rightarrow \phi\phi$ hadronic penguin decay
 - In SM weak mixing phase ϕ_s is identical in $B_s \rightarrow \phi\phi$ and $B_s \rightarrow J/\psi\phi$
 - Define $\Delta S(\phi\phi) = \sin\phi_s(\phi\phi) - \sin\phi_s(J/\psi\phi)$
 - Measurement of $\Delta S(\phi\phi) \approx \sin\phi_s(\phi\phi) \neq 0$ is clear signal for New Physics (NMFV)
- $\Delta S(\phi\phi)$ Sensitivity
 - Best $b \rightarrow s$ penguin mode for LHCb
 - Expect 1.2 k $B_s \rightarrow \phi\phi$ events per 2 fb^{-1}
 - Estimate sensitivity by scaling with $B_s \rightarrow J/\psi\phi$
 - $\sigma(\Delta S(\phi\phi)) \sim 0.14$ in 10 fb^{-1}



- Key channel for LHCb Upgrade
 - $\Delta S(\phi\phi)$ precision statistically limited
 - With 100 fb^{-1} estimate precision $\sigma(\Delta S(\phi\phi)) \sim 0.04$ exciting NP probe
 - Requires **1st level detached vertex trigger** for hadronic decay
 - Expect similar precision for $\Delta S(\phi K_S)$ in decay $B_d \rightarrow \phi K_S$

γ from $B^0 \rightarrow DK^{*0}$, $B^\pm \rightarrow DK^\pm$ & $B_s^0 \rightarrow D_s^\mp K^\pm$



- **LHCb goals for measuring CKM angle γ**

- $B^0 \rightarrow D^0 K^{*0}$, $B^\pm \rightarrow D^0 K^\pm$
Two **interfering tree processes** in neutral or charged B decay
- Use decays common to D^0 and anti- D^0
Cabbibo favoured self-conjugate D decays
e.g. $D^0 \rightarrow K_S \pi \pi$, $K_S K K$, $K K \pi \pi$ **Dalitz analysis**
Cabbibo favoured, single & doubly Cabbibo suppressed D decays
e.g. $D^0 \rightarrow K \pi$, $K K$, $K \pi \pi \pi$ **ADS (GLW) method**
- $B_s \rightarrow D_s^\mp K^\pm$ - two **tree decays** ($b \rightarrow c$ and $b \rightarrow u$) of $O(\lambda^3)$
Interference via B_s mixing

- **γ Sensitivity**

- Expected precision for ADS and Dalitz $\sigma(\gamma) \sim 5^\circ - 15^\circ$ in 2 fb^{-1}

- **Motivation for LHCb Upgrade**


- Theoretical error in SM is very small $< 1^\circ$
- Large statistics helps to reduce systematic error to similar level
- With 100 fb^{-1} estimate precision $\sigma(\gamma) \sim 1^\circ$
- Requires **1st level detached vertex trigger** for hadronic decays

Asymmetry A_{FB} in $B_d \rightarrow K^{*0} \mu^+ \mu^-$



- **Forward-backward asymmetry $A_{FB}(s)$**
 - Asymmetry angle - B flight direction wrt μ^+ direction in $\mu^+ \mu^-$ rest-frame

hep-ph/0003238
PRD61, 074024 (2000)




- **Sensitive probe of New Physics**
 - Deviations from SM by SUSY, graviton exchanges, extra dimensions
 - $A_{FB}(s_0) = 0$ - predicted at LO without hadronic uncertainties
 - Zero point s_0 and integral at high s sensitive to Wilson coefficients

- **Expected Signal Yield**
 - 4.4 k events per 2 fb⁻¹
 - Large statistics allows to measure additional transversity amplitudes
 - Sensitive to right-handed currents
- **A_{FB} zero point sensitivity**
 - $s_0 = 4.0 \pm 0.5 \text{ GeV}^2$ in 10 fb⁻¹

- **LHCb Upgrade Sensitivity**
 - $s_0 = 4.00 \pm 0.16 \text{ GeV}^2$ in 100 fb⁻¹

4% error on $C_7^{\text{eff}}/C_9^{\text{eff}}$




- **What are key measurements?**
 - Selection of four discussed above
 - Importance of different decays could change again with **additional data** from LHC, Tevatron and B-factories
- **LHCb measurements**
 - Many more are **statistics limited**
 - can be improved with **LHCb Upgrade**
 - many of these are very **sensitive to New Physics**
- **Additional LHCb Upgrade measurements**
 - Semileptonic charge asymmetry A_{SL}
 - **Very rare decays**
e.g. **observation of $B_d \rightarrow \mu^+ \mu^-$** and precision measurement of $B_s \rightarrow \mu^+ \mu^-$
 - **Electroweak and radiative penguin decays**
e.g. $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$
 - **Other hadronic penguin decays**
e.g. $B_d \rightarrow \phi K_S$ $B_d \rightarrow \eta' K_S$
 - **CP violation and mixing in charm meson decays**
 - **Lepton flavour violation** in B, charm and tau decays
e.g. $B^0 \rightarrow \mu^+ e^-$, $D^0 \rightarrow \mu^+ e^-$, $\tau^+ \rightarrow \mu^+ \gamma$, $\tau^+ \rightarrow \mu^+ \mu^- \mu^+$

Comparison with Super-B factory



Sensitivity Comparison ~2020
LHCb 100 fb⁻¹ vs Super-B factory 50 ab⁻¹

SuperB numbers from
 M Hazumi - Flavour in
 LHC era workshop



B_s only accessible to LHCb

Common

**No IP
Neutrals, ν**

Preliminary

LHCb Upgrade Detector and Trigger



- **LHCb Luminosity**

- Running at $\mathcal{L} \sim 2 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$ is default
- Make use of learning experience in running LHCb
- Will operate at luminosity up to $\mathcal{L} \sim 5 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$



- **LHCb Detectors**

- Detectors able to cope with $\mathcal{L} \sim 5 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
- Vertex detector sensors require replacing after 6 - 8 fb⁻¹ (~3 years)
- Default replacement - same geometry, similar slightly improved sensors

- **Level-0 Trigger - LO**

- High p_T - $\mu, \mu\mu, e, \gamma, \text{ hadron} + \text{ pileup}$
- Read-out at 40 MHz 4 μs latency
- Existing Front-End electronics limits LO Trigger output to 1.1 MHz

L0 efficiency



- **L0 muon trigger**
 - ~90% efficiency
 - scales with luminosity
- **L0 hadron trigger**
 - ~40% efficient
 - does not scale with luminosity
 - Required for $B_s \rightarrow \phi\phi$ and $B^\pm \rightarrow D^0 K^\pm$

- **The Big Question**

- How do we upgrade LHCb detector such that it can operate at 10 times design luminosity of $\mathcal{L} \sim 2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$?
- Physics, Detector and Trigger studies have started
- Several approaches under investigation

- **Vertex Detector**

- VELO sensors require replacing with radiation-hard sensors

- **LO Detached Vertex Trigger**

- Add Vertex Detector (VELO) and Trigger Tracker (TT) to LO Trigger
- Requires 40 MHz readout of VELO and TT
- Implementation in FPGAs
- Is Magnetic field in VELO region required?

- **Other LHCb Detectors**


- need upgrade due to occupancy and/or irradiation
- Replace inner most region of RICH photo detectors
- Replace inner most region of ECAL with crystal calorimeter
- Possibly add other sub-detectors to 40 MHz readout

- **Readout full detector at 40 MHz**
 - Requires new readout architecture
 - All trigger decisions in CPU farm
 - All Front-end electronics must be redesigned
 - Increased radiation hardness required
 - Electronics R&D can profit from common LHC development
- **Detectors for 40 MHz Readout**
 - VELO sensors require replacing with radiation-hard sensors
 - Silicon tracker sensors (TT and IT) need to be replaced
 - Outer tracker occupancy likely prohibitive
Increase (decrease) area of Inner/Outer Tracker
 - RICH photo detectors need to be replaced
- **Additional Considerations**
 - for running LHCb at $\mathcal{L} \sim 2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$
 - Costs expected to compare favourably with existing infrastructure and complementary approaches

Vertex Detector Upgrade


- Critical for LHCb upgrade physics programme

Radiation Hard Vertex Detector with Displaced Vertex Trigger




VESPA

VElo Superior Performance Apparatus





Radiation Hard Vertex Detector

- **Vertex Detector for LHCb Upgrade**
 - requires high radiation tolerance device $>10^{15}$ 1 MeV $\text{neutron}_{\text{eq}} / \text{cm}^2$
- **Geometry - Strixels / Pixels**
 - remove RF foil
 - 3% X_0 before 1st measurement
 - move closer to beam from 8 \rightarrow 5mm



VELO Module




Radiation Hard Technologies





- Active Technology R&D for LHC upgrades
- Applicable to strixels & pixels


Czochralski



n-on-p




3D



Extreme radiation hard
 For 4.5×10^{14} 24 GeV p/cm²
 Depletion voltage = 19V

F. Muheim



LHCb Upgrade Trigger Studies




• Method

- Combine L0 with detached vertex trigger
L0 - hadron $E_T > 3 \text{ GeV}$
track with largest $p_T > 2 \text{ GeV}$
impact parameter $|IP| > 50 \text{ um}$
- Run L1 trigger algorithm at L0

• Preliminary results

- $B_s \rightarrow D_s^\mp K^\pm$ at $L = 6 \times 10^{32}$
- For L0+vtx Min. bias efficiency does not depend strongly on # of interactions n_r
- L0 - hadron rate: $r = 0.8 \text{ MHz}$
 $B_s \rightarrow D_s^\mp K^\pm$ efficiency $\varepsilon = 66\%$
- Better efficiency than L0 trigger at $L = 2 \times 10^{32}$ (baseline)
 $r = 0.7 \text{ MHz}$ $\varepsilon = 39\%$
- Yield $B_s \rightarrow D_s^\mp K^\pm$ is 5 times baseline
- Yield scales linearly with luminosity





- **Standard Model is very successful**
 - Require precision measurements to probe/establish flavour structure of New Physics
- **Many LHCb results will be statistically limited**
 - LHCb plans to run initially for five years at $\mathcal{L} \sim 2 \dots 5 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
 - 6 - 10 fb^{-1} data set will not reach full potential of B physics at hadron colliders
- **LHCb Upgrade Plans**
 - Replace VELO with radiation hard vertex detector
 - Add first level detached vertex trigger to LHCb experiment to trigger efficiently on hadronic modes at high luminosities
 - Readout of all LHCb detectors at 40 MHz
 - Requires new front-end electronics, silicon sensors, RICH photo detectors
 - Run five years at $\mathcal{L} \sim 2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ and collect 100 fb^{-1} data sample
- **LHCb Physics reach with 100 fb^{-1}**
 - Perform $\sim 10\sigma$ measurement of SM weak B_s mixing phase $\phi_s = -0.035$ in $B_s \rightarrow J/\psi\phi$
 - Probe or establish New Physics by measuring ϕ_s in hadronic penguin decay $B_s \rightarrow \phi\phi$ with a precision of $\sigma(\Delta S(\phi\phi)) = 0.040$
 - Measure CKM angle γ to a precision of $\sigma(\gamma) \sim 1^\circ$
 - Probe New Physics in rare B meson decays
Measure Wilson coefficient C_7/C_9 to 4% in electroweak decay $B \rightarrow K^*0\mu^+\mu^-$
 - Measure $B_d \rightarrow \mu^+\mu^-$

Backup Slides



LHCb Physics Programme



B production,
 B_c , b-baryon physics
 Charm decays
 Tau Lepton flavour violation

- Rare decays - very sensitive to NP
- Radiative penguin e.g. $B_d \rightarrow K^* \gamma$, $B_s \rightarrow \Phi \gamma$
 - Electroweak penguin e.g. $B_d \rightarrow K^{*0} \mu^+ \mu^-$
 - Gluonic penguin e.g. $B_s \rightarrow \Phi \Phi$, $B_d \rightarrow \Phi K_s$
 - Rare box diagram e.g. $B_s \rightarrow \mu^+ \mu^-$