

Matrix Theory for Baryons: An Overview of Holographic QCD for Nuclear Physics

Sinya Aoki¹, Koji Hashimoto² and Norihiro Iizuka³

¹ *Graduate School of Pure and Applied Sciences, University of Tsukuba, Ibaraki 305-8571, Japan, and
Center for Computational Sciences, University of Tsukuba, Ibaraki 305-8577, Japan
E-mail: [saoki\(at\)het.ph.tsukuba.ac.jp](mailto:saoki@het.ph.tsukuba.ac.jp)*

² *Mathematical Physics Lab., RIKEN Nishina Center, Saitama 351-0198, Japan
E-mail: [koji\(at\)riken.jp](mailto:koji@riken.jp)*

³ *Theory Division, CERN, CH-1211 Geneva 23, Switzerland
E-mail: [norihiro.iizuka\(at\)cern.ch](mailto:norihiro.iizuka@cern.ch)*

ABSTRACT: We provide, for non-experts, a brief overview of holographic QCD and a review of a recent proposal of matrix-description [1] of multi-baryon systems in holographic QCD. Based on the matrix model, we derive the baryon interaction at short distances in multi-flavor holographic QCD. We show that there is a very universal repulsive core of inter-baryon forces for generic number of flavors. This is consistent with a recent lattice QCD analysis for $N_f = 2, 3$ where repulsive core looks universal. We also provide a comparison of our results with the lattice QCD and the operator product expansion (OPE) analysis.

Contents

1. M-theory for Nuclear Physics?	1
2. Universal Problems in Holographic QCD	4
2.1 How holographic QCD is different from QCD	4
2.2 Popular holographic models and their problems	7
3. Review : M(atrix) Description of Multi-Baryon System	11
3.1 Baryons are matrices	12
3.2 Derived baryon spectrum	15
3.3 Universal repulsive core of nucleons	16
3.4 Toward a description of atomic nuclei	17
4. Multi-Flavor Nuclear Forces Via Holography	19
4.1 Strangeness and holography	19
4.2 The effective model of multi-baryon system	20
4.3 Two-baryon configuration	21
4.4 Explicit inter-baryon potential	22
4.5 Universal repulsive core	24
5. A comparison with lattice QCD and OPE	24

1. M-theory for Nuclear Physics?

What is “M-theory” for nuclear physics? Although the “M-theory”¹ stands for a theory of everything which unifies all string theories [2], one can generalize the use of the word “M-theory” not only for string theories but also for other subjects in physics. What is M-theory for nuclear physics, if exists?

This kind of question brings us to a bigger picture of relations between various subjects within physics, so it is not of no use. The question, however, sounds ridiculous, because the answer for it is obvious: The M-theory for nuclear physics is QCD, or more precisely, the Standard Model of elementary particles. Nucleons, which are the building blocks of nuclei, are bound states of quarks and gluons in

¹M is for mystery, mother, matrix. [2]

QCD. Supposing that one could solve QCD completely, in principle one should be able to derive all the properties of nuclei, which is nothing but the nuclear physics. Therefore, in this sense, QCD is the M-theory for the nuclear physics. However, QCD is notorious as being difficult to solve, due to its strong coupling nature: the strong force makes quarks bound to each other. Therefore we need a new tool for solving QCD to “derive” nuclear physics. Once the new tool is available, we may then say that we “understand” the real-world nuclear physics phenomena from M-theory.

Since this new tool has been missing for long years in research, apparently we have a hierarchical structure between studying perturbative QCD, nuclear physics and hadron physics (see Fig. 1). Standard nuclear physics starts with a quantum mechanics of multi nucleons, with inter-nucleon potential (nuclear force) given by experiments, or by hand to match phenomena. The quantum mechanics Lagrangian becomes

$$S = \int dt \left[\sum_{s=1}^A \frac{M}{2} (\partial_t x_{(s)}^M(t))^2 - \sum_{s_1 \neq s_2} V[x_{(s_1)}^M - x_{(s_2)}^M] + \dots \right], \quad (1.1)$$

where we have A nucleons whose locations are given by $x_{(s)}^M(t)$ with $s = 1, \dots, A$. The first term is the kinetic term of the nucleons with mass M , while the second term is the nuclear force. The problem lying in the unification of our concern is the fact that in nuclear physics the nuclear force V is given by experiments, and not by fundamental theory, *i.e.*, QCD. In principle, the potential should have been derived from QCD, as we all know that nucleons and hadrons are made of quarks and gluons — but it is very difficult.

It is very recent that the nuclear force was calculated from QCD with use of numerical methods: lattice QCD [3, 4]. The lattice QCD has accomplished a great success in hadron physics. In particular for hadron spectroscopy and hadron interactions, the lattice QCD is now very close to the physical parameters of QCD, the real world. Furthermore, there is a progress in this direction toward nuclear physics itself [5]. Once the lattice QCD comes to deal with a system of multi-baryons, a part of nuclear physics becomes accessible directly from QCD. A huge number of quark contractions in large nuclei, which requires almost unrealistically high power of supercomputers, however, is a big obstacle in this direction. Furthermore, it is of course more ideal if we can understand physics without relying on the computers. Unfortunately we have not yet reached that stage. Therefore, we are facing at a situation where the hadron physics and the nuclear physics are disconnected each other in a sense, due to a difficulty in solving the strongly coupled QCD.

At this occasion, the new tool using string theory comes into play. The renowned AdS/CFT correspondence [6, 7] makes it possible to solve a certain limit of QCD-like gauge theories, and it offers a certain direct path from QCD to nuclear physics. If one can derive an action like (1.1) from QCD, it can be regarded as an effective theory for nuclear physics derived from M-theory.




Figure 1: A conceptual merit of the AdS/CFT correspondence, for a possible bridge between elementary particle physics, hadron physics and nuclear physics.

In this paper, we review the recent progress along this direction as an application of the AdS/CFT correspondence. In [1], two of the present authors (K.H. and N.I.), together with Piljin Yi, derived an action of a multi-baryon system, by using the AdS/CFT correspondence applied to large N_c QCD. The action indeed has the form of (1.1), and it serves as a candidate for the bridge between QCD and nuclear physics.

As it was derived from the large N_c QCD, the action is written only with two free parameters: the QCD scale and the QCD coupling. Therefore we can make a check of the derived theory by just calculating various observables in nuclear physics with this action and compare those with experiments, to test the validity of the action, up to the approximations of the large N_c and the strong coupling expansion. Explicitly demonstrated in the literature are:

- (i) Baryon spectroscopy [1]
- (ii) Universal repulsive core of nucleons [1]
- (iii) Three-body nuclear forces [8]
- (iv) Spin statistics of baryons [9]
- (v) Formation of atomic nuclei [10]

In all of these calculations, the results are qualitatively reasonable compared to the experiments². As we will describe in this paper, there are a lot more physical

²Within the same framework, using the flavor brane action, it has been reported that a wider class of results are compatible with experiments: baryon spectrum was originally derived in [11], and charge radii of baryons [12], suppression of multi-nucleon forces [13], baryon spectra with three flavors [14], *etc.* The repulsive core has been calculated in the same manner [15]. See also some alternative approach given in [16].

observables which can be calculated in the framework.

The first aim of this paper is to give a review of the effective action for the multi-baryon system [1], for non-experts of holographic methods. The second aim is to show a new result on the short-distance force between baryons with multi flavors where the number of flavors N_f is $N_f > 2$.

This paper is organized as follows. The first part of this paper is mainly a review. In section 2, we give a brief review of the status of holographic QCD, explaining the difference between the holographic QCD and real QCD, to emphasize what are remaining problems in holographic QCD. Then in section 3, we shall explain the nuclear physics action derived in AdS/CFT correspondence, with emphasis on its properties, new insights and connection to nuclear physics, for non-experts.

The second part of this paper consists of new results. In section 4, we calculate the short distance inter-baryon forces for the case of multi-flavors (the number of flavors larger than 2). We shall see that the repulsive core remains even for generic number of flavors, thus find a universal repulsive core. The result is consistent with recent lattice results with $N_f = 3$ where in most of the channels there appears an inter-baryon repulsive potential. Therefore our result would also serve as another nontrivial consistency check. This part of the paper includes technical details. Readers who know holographic QCD and the matrix model approach of [1] can start with section 4 as it is written independently of section 2 and 3. In the last section 5, we provide a review of the recent lattice results for multi-flavors and also the operator product expansion (OPE), as a comparison to our holographic results.

2. Universal Problems in Holographic QCD

For readers who are not familiar with the subject of the holographic QCD (the AdS/CFT correspondence applied to QCD),³ here in this section, we summarize important problems which are to be addressed in the holographic QCD. In particular, we make a stress on what are assumptions and what are ignored in holographic QCD. This would make clear an importance and a validity of the AdS/CFT matrix model approach to multi-baryon system and nuclear physics, which we shall review in the next section.

2.1 How holographic QCD is different from QCD

The holographic QCD is different from the real QCD. The holographic QCD, however, is very important setting as it provides us with a non-perturbative and analytic

³In this article, and in most of articles in this field, the word “AdS/CFT” is equivalent to the word “gauge/gravity” or “holography” in use. We say “bulk” for gravity or string side calculation and “boundary” for the gauge theory side calculations.

method to systematically study the strong coupling nature of gauge theories including QCD-like gauge theories. Therefore we first need to know why it is difficult to directly apply the holography method to the real QCD.⁴

2.1.1 Forced large N_c and λ limits

The AdS/CFT correspondence in string theory is a conjecture on the equivalence between a non-gravitational gauge theory and a *string theory in asymptotically AdS background*. Note that the duality is *not* between gauge theory and gravity, but rather string theory. Only when large N_c and large λ (which is a 't Hooft coupling of the gauge theory) are taken, the string theory side can be approximated by a gravity theory with background geometries of weakly curved spacetime. Low energy excitations of the string, such as gravitons, are light, while the long string itself becomes very heavy. This is almost an unique universal situation where one can describe low energy physics concretely in string theory as gravitational theory. Due to the technical difficulty of solving string theory in generically curved background, in many situation where we apply the holography to theories like QCD, the two limits, large N_c and large λ , are forced so that we can approximate string theory as a gravity theory.

It is this approximation which makes us face several difficulty in the comparison between the holographic QCD and realistic QCD or nuclear phenomena. According to old string models, hadrons with higher spins are stringy excitations. Holographic QCD follows and generalizes the old string models based on QCD strings. In the AdS/CFT correspondence, the tension of the strings in the gravity side is $\mathcal{O}(\lambda)$, so the stringy excitations become extremely heavy, and resultantly, the higher spin modes are parametrically heavy and decouple from the gravity excitations. This is the reason why in large λ , string theory is approximated as a gravitational theory. If we take λ to infinity, however, the highest spin excitation of the system be graviton, which has spin 2, and all the stringy modes whose spin are bigger than 2 be infinitely heavy. On the other hand, in nuclear physics, there are many hadronic excitations whose spin are bigger than 2 and all of these higher spin hadronic excitations have the same order mass scale compared with lower spin excitations. Therefore, in holographic QCD within the gravitational approximation, we should keep in our mind that there could be a contradiction for a comparison with data caused by the missing degrees of freedom whose spin are bigger than 2.

⁴Here keep in mind that we are talking about top-down approach of holography from string theory. Any bottom-up approach, to write down higher-dimensional gravity models as phenomenological models for QCD, does not have clear understanding on which gauge theory is dual to those bottom-up gravity models. Therefore, to be precise, we discuss top-down model only in the framework derived in string theory in this paper. Bottom-up models are criticized only through their comparison to real QCD data, while top-down models can be more concrete in criticisms as they are directly related to QCD through string theory or D-brane construction.

One may then wonder why we do not directly try to solve string theory in asymptotically AdS background, instead of using the gravity approximation. A problem is that a precise treatment of the fundamental strings in the curved geometry, *i.e.* the quantization of the string, is still missing in any formulations of string theory. We miss a fundamental tool to analyze the string side. This waits for a further development of methods to quantize strings in curved geometries.⁵

In addition, we have $N_c = 3$ in realistic QCD, while any quantities are computed at the leading order of the $1/N_c$ expansion around $N_c = \infty$ in holographic QCD. So, in comparison to experiments, we expect, at least, 33% or more errors generically due to the large N_c approximation. At present, computations of the sub-leading $1/N_c$ corrections, which correspond to that of string loop corrections in the gravity side, are technically difficult. These all imply that the holography methods are better applied to reveal some robust features of QCD, which are independent on the values of N_c , not to make a comparison to precision measurements.

We might wonder under what circumstance physical quantities could be more insensitive to N_c . In the confining phase of QCD, as the color degrees of freedom are confined, we cannot directly observe the number of colors. We therefore naively might expect that physics might be independent on the values of N_c . On the other hand, physics in the deconfining phase would suffer more defects from the large N_c limit. Interestingly, however, there are many successful examples of the calculations in the deconfining phase for the shear viscosity [18], quark energy loss [19] etc in the quark gluon plasma phase compared with the experiments at RHIC and LHC. At this moment, we do not have a clear picture under what situation the large N_c approximation are justified⁶.

2.1.2 Lack of the asymptotic freedom leading to multiple parameters

QCD is specified by a peculiar energy scale Λ_{QCD} as a result of the running coupling constant and the asymptotic freedom. In particular for the massless QCD, it has only this scale in the theory and there is no other parameter. In contrast to this, the scale in the holographic QCD is introduced into the system by hand as an input. In the holographic QCD, the operators which break the conformal invariance is introduced

⁵One major progress along this line would be a correspondence between vector models and higher-spin gauge theories [17], as an explicit toy model of the AdS/CFT correspondence in the $\lambda \rightarrow 0$ limit.

⁶There are several good examples which work beyond the large N_c and large λ limit in the holographic setting. One of the examples is the Wilson loop, which is nothing but a heavy string trajectory, where we can calculate, by using the localization technique, large N_c but any values of λ calculation and can see a precise matching between string side and gauge theory side. In addition, in the holographic QCD setting, there is an attempt even in the large λ limit to take into account the degrees of freedom corresponding to the massive open strings whose spins are bigger than 2, and quantize these massive stringy excitation in the weakly curved geometry [20]. This also gives a qualitatively very good comparison with the experimental data.

at some scale Λ_{cutoff} , so that the coupling constant in the theory becomes scale dependent (the running coupling).

This leads us to a strange situation where we have two scales in holographic system: One is the scale Λ_{cutoff} we introduced, and the other is the Λ_{QCD} . This Λ_{QCD} is an emergent scale in the low energy physics where hadron physics emerges. This Λ_{QCD} is determined as a function of two input parameters, Λ_{cutoff} and 't Hooft coupling constant $\lambda = g_s N_c$ (where g_s is a string theory coupling constant). In principle, if one can take the double-scaling limit at which the Λ_{QCD} is fixed while the holographic scale Λ_{cutoff} (at which typically particle fields which do not exist in QCD appear) is taken to infinity, by fine-tuning λ , then the above problem would be resolved.

This, however, is not an easy task: In order to make the gauge theory coupling constant at the scale Λ_{cutoff} to be weak, the corresponding geometry in the holographic side becomes highly curved that the supergravity description is no more reliable. As mentioned in the previous section, however, it is technically difficult to go beyond large N_c and λ . As a result, the difficulty of taking the double scaling limit remains in any holographic QCD models in the top-down approach.

Of course, one can say that the number of the parameters, two given by Λ_{cutoff} and λ , is significantly small, and it is good enough to have nontrivial check and predictions in QCD, compared to many other phenomenological models.

It is noted that the coupling constant in the gauge theory becomes strong again beyond Λ_{cutoff} due to supersymmetric particles which appear above Λ_{cutoff} ⁷. This property is completely different from the ordinary QCD, where the coupling constant becomes smaller and smaller at higher and higher energy (the asymptotic freedom).

2.2 Popular holographic models and their problems

Next we shall look at popular holographic models which are widely used for various purposes, in particular from the viewpoints of their strong points and limitations. We make emphasis on the point that, depending on physical quantities of interest, one can choose a holographic model among many. We here briefly review five models popularly used in the top-down approach of the holographic QCD.

- Supersymmetric D3-brane model (Asymptotic AdS_5)

The gauge-theory counterpart of this model is $\mathcal{N} = 4$ supersymmetric Yang-Mills theory. This theory is highly supersymmetric and so is far from the realistic QCD. However, to see robust results of deconfined gluons in high temperature, where we expect the effect of supersymmetry is not crucial, the theory would be sometimes good enough to extract typical phenomena of strong

⁷On the other hand, the Λ_{QCD} is seen in gravity side as an IR cut-off of the geometry and no geometry exists below that IR cut-off scale (radius).

coupling gauge theories. The most successful result which came out of this is the computation of shear viscosity of quark gluon plasma in high temperature phase of QCD [18]. Although the computation has employed only a geometry representing a finite temperature phase of the $\mathcal{N} = 4$ supersymmetric gauge theory, the result is close to the experimental observation.

It is difficult to argue why this model works so well. In terms of the large N_c expansion, reasons why the $1/N_c$ corrections do not contribute and why they do not modify qualitative nature are still missing. In addition, there are many fields in the supersymmetric theory which are absent in QCD. An issue of the universality of the value of the shear viscosity is still to be settled. Nevertheless, other physical quantities have been calculated so far and results give insightful suggestion for heavy ion experiments.

- D3D7 model

Introducing D7-branes as flavor D-branes [21] makes it possible to include supersymmetric quark fields (hyper multiplets in fundamental representation) in above D3-brane model. This make it possible to calculate the quark energy loss in quark gluon plasma and drag forces [19].

One can also discuss $U(1)$ part of chiral symmetry breaking in this D3D7 setting [22]. The position of flavor D7-brane represents the symmetry in the Yang-Mills theory on D3-branes. If the position of flavor D7-brane are symmetric, we have that symmetry in Yang-Mills theory, however if not, we have corresponding symmetry breaking. By embedding the $U(1)$ part of chiral symmetry as a geometrical rotational symmetry in D3-brane-setting, we can discuss how this rotational symmetry is spontaneously broken from the position of D7-brane at low temperature, and restored at high temperature. The position of D7-branes is determined in order to minimize the free energy of the system. See for example, Fig. 6.2 and Fig. 6.6 of [23].

- Witten's non-supersymmetric model [24]

The corresponding geometry is called Gibbons-Maeda geometry [25], and corresponds to a 1+3-dimensional pure bosonic Yang-Mills theory (*i.e.* the theory of gluons) at low energy without supersymmetry. The geometry is made of N_c D4-branes wrapping a circle. This circle compactification brings the 1+4-dimensional theory down to the 1+3-dimensional Yang-Mills theory. It breaks the supersymmetry by imposing anti-periodic boundary conditions for fermions, and at low energy all fermions are massive and only massless gluons survive. Adjoint scalars obtain masses through quantum corrections which are roughly of order of the scale defined by the radius of the circle.

This geometry captures important gauge-theory property: the confinement. In fact, using the bulk equations of motion from the gravity side, one can demonstrate as [24] that the fluctuation spectrum corresponding to the glueball spectra is discrete and mass-gapped, and that the calculated wilson loop shows the area law. Furthermore, above a critical temperature a phase transition occurs and is interpreted as a confinement-deconfinement transition since the spectrum becomes continuous at the high temperature phase.

There is one caveat here: the phase transition scale is nothing but the scale of the extra circle on which D4-branes wrapping, as it is the unique dimensionful physical parameter. So naively speaking, the higher-dimension cutoff scale is re-interpreted as the scale of the gluon theory. The excitation of the massive gluinos and adjoint scalars and their superpartners should come into the spectrum above the scale, therefore the theory be no more purely bosonic Yang-Mills theory. If one naively ignore those and regard the gravity fluctuation as the glueball spectrum made solely of the gluons above this scale, we might get some mismatches for the spectrum comparison.

Therefore the intrinsic problem of this geometry interpreted as a dual of the pure Yang-Mills theory is the double meaning of the dynamical scale and the compactification scale. In order to remove additional degrees of freedom, we have to take the double scaling limit; We keep Λ_{QCD} fixed and at the same time, take the scale, associated with the D4-brane wrapping circle, to infinity. Any proper scaling limit where the dynamical scale is fixed while the compactification scale is taken to infinity, has not been formulated yet.

- D4D6 model [26]

In the Witten's geometry, flavor D6-branes can be added to include quarks in the theory. The string connecting the N_c D4-branes and the N_f D6-branes give a low energy excitation which behaves like a quark. In the gravity description, the shape of the D6-branes is deformed and it can be interpreted as a spontaneous breaking of the (anomalous) $U(1)$ axial symmetry. The model can include various quark masses, so in particular quark mass dependences of various low energy quantities can be studied.

- D4D8 model (Sakai-Sugimoto model) [27]

This theory adds flavor D8-branes in the Witten's geometry. One of the superior point of this model compared with others is that by adding D8-branes, one can obtain only left-chirality fermions at the intersection points between D4 and D8-branes. On the other hand, by adding anti-D8-brane, one can obtain only right-chirality fermions at the intersection points between D4 and anti-D8 branes. This implies that by adding N_f number of both D8 and anti-D8-branes,




Figure 2: A schematic picture of two phases in the gravity dual of the bosonic pure Yang-Mills theory. On the geometry specified by the surface of the cylinder, flavor D8-branes (red lines) are put. The vertical direction in the figure is a holographic dimension, while the circular direction is for the compact circle which brings the 1+4-dimensional theory down to the 1+3-dimensional pure Yang-Mills theory. The left figure shows a geometry corresponding to a confining phase, while the right one is that for a deconfined geometry. (Left) The geometry consistently truncated at a certain place along the holographic direction corresponding to Λ_{QCD} , and the D8-brane and the anti-D8-brane are connected due to the geometry, which shows the spontaneous chiral symmetry breaking. (Right) The geometry ends with a horizon of a black hole (shaded region). The D8-brane and the anti-D8-brane are independent, which is a chiral symmetry restoration.

we can have both left and right chiral fermions (quarks) in the system with explicit dependence on the chiral symmetry $U(N_f)_L \times U(N_f)_R$, which are very close to the realistic QCD.

Similar to the D3D7 system, the chiral symmetry $U(N_f)_L \times U(N_f)_R$ is seen from the position of flavor D8 and anti-D8-branes. Due to the warped factor of Gibbons-Maeda geometry, one can demonstrate that the free energy at low temperature is lower if both N_f D8-branes and N_f anti-D8-brane are combined into N_f 8-branes. See the left figure of Fig. 2. In high temperature, these combined effects of D8 and anti-D8 are hidden behind the horizon (right figure of Fig. 2), and we have chiral symmetry restoration, which can be seen geometrically. In this way, this model shows the spontaneous chiral symmetry breaking at low temperature and its restoration at high temperature in a geometrical way.

Except for the point that the quark mass is difficult to be introduced⁸ due to the non-supersymmetric nature and the existence of chiral matter, this holographic

⁸See [28] for a possible way to introduce the quark masses to the model.

model is the most successful model in view of the study for the low energy hadron physics. In addition to the meson spectrum and interactions, baryon spectrum and its chiral dynamics can be systematically studied.

This model again suffers from the same problem as the Witten's geometry has: the unnecessary modes, such as squarks in addition to the gluinos, exist in the theory at high energy scale⁹. So we tentatively ignore modes which are expected to be absent in QCD, to compare the holographic results with experiments.

In summary, in all holographic models popularly known, there remains a problem of having fields which are absent in real QCD above some scale of the theory. And relatedly, the low energy scale of the theory which one would like to interpret as the QCD scale is shared with the scale where the unnecessary fields show up.

Naively, at very low energy, the effects of these unnecessary fields would be small, so the prediction from holography should be better at the low energy. This simple fact would motivate us strongly to visit nuclear physics. Nuclear physics treats nuclei: bound states of nucleons at the energy scale much lower than the QCD scale. However in order to make the comparison with data more precise, we have to take the double scaling limit in holographic QCD, where we take the scale, beyond which unnecessary fields be dynamical, to infinity while keeping the QCD scale fixed.

Due to the reasons explained in section 1, nuclear physics includes a lot to be explained by QCD. Standard nuclear physics has many assumptions, and the origins of those fundamental assumptions may be explained directly from QCD, once we apply the holographic methods to QCD.

3. Review : M(atrrix) Description of Multi-Baryon System

The upshot of the theory [1] for the multi-baryon system, derived in AdS/CFT, is that it is a theory of matrix degrees of freedom, with the following robust form of the action:

$$S = \frac{M}{2} \int dt \operatorname{tr} \left[(\partial_t X^M(t))^2 - g[X^M, X^N]^2 + \dots \right] \quad (3.1)$$

Let us clarify the relation between (3.1) and the nuclear physics action (1.1). The matrix X^M is a hermitian $A \times A$ matrix, where A is the number of baryons (which resultantly becomes the mass number of a nucleus if all the baryons are bounded together as a big nucleus). Once it is diagonalized, the eigenvalues are nothing but the locations of the baryons which are given by $x_{(s)}^M$ (for $s = 1, \dots, A$) in the nuclear physics action (1.1). There are off-diagonal entries in X^M , which we interpret the degrees of freedom associated with the nuclear force mediator (such as pion, massive

⁹Here holographic scale, which is determined by the scale on which D4-branes are wrapping, gives the scale beyond which these additional “junk” are excited.




Figure 3: Derivation of the matrix description of nuclear physics.

vector mesons etc). Classically integrating those degrees of freedom in the action gives rise to the interaction between the eigenvalues of X^M . For the detail of nuclear force derivations, see section 4 of [1]. This interaction is interpreted as the inter-nucleon potential (nuclear force). The terms which are not written in the action (3.1) (specified as “+...”) are fields representing spins and isospins (flavor degrees of freedom of the baryons). Again, the precise form of the action is given in [1] and presented in (4.1).

In this section, we provide a review of the matrix formulation of the multi-baryon system in simple terms. First, we shall explain below the reason why we have the matrix degrees of freedom for the baryons in AdS/CFT, and the origin of the action written above. Then we come to a review of the concrete analysis for a single baryon system to obtain the baryon spectrum, and also a review of two and three baryon systems for deriving the short distance nuclear force. These were done in the original paper [1]. Then in the final part of this section, we review the importance of the matrix model action (3.1) for providing a possible unified view of nuclear physics.

3.1 Baryons are matrices

As we outlined above, the most important and novel part of the new description of the multi-baryon system (3.1) is the fact that *baryons are described by $A \times A$ matrices*. In fact, this is a robust result once one applies the AdS/CFT correspondence to QCD for the multi-baryon system.

There are two key points to derive this fact, which are shown in Fig. 3.

- *A baryon is a D-brane.*

In AdS/CFT correspondence, we need a large N_c expansion to use the dual gravity description. For large N_c QCD, baryons are heavy object whose mass

is of order of $\mathcal{O}(N_c)$, since a single baryon consists of N_c quarks. In the gravity side of the AdS/CFT, what is the object whose mass is so large? The answer is D-branes. D-branes are solitonic objects in string theory, whose mass are order of $\mathcal{O}(N_c)$. Therefore the baryons are expected to correspond to the D-branes in the gravity side of the AdS/CFT correspondence. In fact, baryons are D-branes, and technically speaking, these baryon D-branes are wrapping on the closed surface like higher dimensional sphere on which string theoretic RR flux is penetrating. Through the D-brane action, the wrapped D-branes on some closed surface with penetrating flux induces N_c unit of $U(1)$ charges on that closed surface. On the closed surface, total charges must be zero to satisfy Gauss's law. This implies that we need to add compensating charged objects on that surface, which turns out to be N_c number of fundamental strings [29]. Therefore these D-branes behaves as baryons.

- *Multi-D-branes are matrices.*

D-branes are defined as surfaces on which open strings can end. When D-branes are on top of each other, fundamental and anti-fundamental strings connecting between those D-branes can be arbitrarily short, and can be massless. The low-energy excitation of those light modes are classified by an $A \times A$ matrix when A is the number of the D-branes, since each open string has two ends labeled as (a, b) where $a, b = 1, \dots, A$. Therefore the low energy degrees of freedom on the coincident A D-branes are $A \times A$ matrices.

Combining these two, we arrive at the inevitable conclusion that nuclei (or the multi-baryon system) in the AdS/CFT correspondence should be described by matrices.

Furthermore, the effective action of D-branes has the universal form of (3.1). The interpretation is definite: the eigenvalues of the field X are location of the A number of D-branes. Therefore, we come to a conjecture that the effective action (3.1) describes nuclear physics.

One of the most important properties of nuclei is its crucial dependence on isospins. Nuclear force strongly depends on whether the nucleon is a proton or a neutron. Consequently, we have a nuclear chart and stable/unstable nuclei. How the isospin dependence can come in in this formulation? The answer is quite simple: another matrix w which is an $A \times N_f$ complex matrix joins the effective action. The isospins are nothing but the quark flavor degrees of freedom, and N_f is the number of the quark flavors.

Fig. 4 clarifies why this new matrix shows up in the gravity side of the AdS/CFT correspondence. As we reviewed in the previous section, the flavor can be represented by an introduction of “flavor D-branes” into the gravity geometry. Then, in addition to the baryon D-branes, we have the flavor D-branes, so there appears an open string

Gravity side of AdS/CFT




Figure 4: The appearance of two kinds of matrices X and w . X connects baryon D-branes (depicted as two black blobs), while w connects a baryon D-brane and a flavor D-brane (depicted as three parallel sheets). w obtains nonzero VEV to satisfy the Gauss’s law on each baryon D-brane, which is equivalent to N_c open strings.

which connects the two kinds of D-branes. This string should be described by $A \times N_f$ matrices, as in the same manner as the $A \times A$ matrix X for the string among the baryon D-branes.

Although the species of the fields appearing in the low energy of the multi-baryon system in the AdS/CFT are just X and w , the precise interaction between these fields, and also the coefficients in the effective action, depends on what kind of D-brane configurations (holographic models) we use for the large N_c QCD. When we use the most popular D4D8 model (Sakai-Sugimoto model) described in section 2.2, the baryon D-branes are D4-branes wrapping S^4 , and the flavor D-branes are D8-branes wrapping S^4 . This means that the baryon D4-branes can be located inside the flavor D8-branes. The Dp - $D(p+4)$ system in superstring theory is well-understood, as a geometric realization of the instanton construction: the Dp -brane can be seen as a Yang-Mills instanton through the gauge fields on the $D(p+4)$ -brane, where the dynamics of the Dp -brane can be determined by a so-called ADHM matrices used for the instanton construction [30, 31]. Therefore, within the D4D8 holographic model, our low energy effective action for the multi-baryon system is nothing but a generalization of the ADHM matrix models.

The matrix effective action is concretely written in (4.1) for D4D8 model, but in

this review part we don't need the explicit form, as we explain only the conceptual part to show the robustness of the derivation. Furthermore, it is straightforward to construct explicit forms for the matrix effective action for another D-brane models described in subsection 2.2. However for concreteness, in this paper we concentrate on the model constructed for D4D8 model in [1].

Next, we give a review of a single baryon spectrum ($A = 1$), and also a derivation of the short distance nuclear force ($A = 2, 3$). The important fact for the application is that the matrix action has only two free parameters:

3.2 Derived baryon spectrum

The simplest case is $A = 1$ where we have only a single baryon. In this case, the quantum mechanics should give the baryon spectrum. Excited states of a baryon emerges from the quantum mechanics.

Let us recall the Skyrme model [32, 33]. In the Skyrme model, a baryon appears as a soliton of the Skyrme model which is nothing but a peculiar effective action of low energy pions. Any soliton has fluctuation modes, massive or massless (zero modes). The fluctuation modes, which are just a function of time, obey a hamiltonian, and they can be quantized. The resulting quantized fluctuation spectrum is interpreted in the Skyrme model as the baryon spectrum. Here in the AdS/CFT matrix model approach, the hamiltonian of the fluctuation modes are directly given as our matrix model hamiltonian (3.1). So, one easy interpretation of the matrix model is a moduli hamiltonian of generalized Skyrmons. However very small number of parameters (only two parameters) in our holographic setting gives the superiority of our construction compared with generic Skyrme model which have many parameters.

For $A = 1$, the matrix model becomes extremely simple. The hamiltonian for two flavors ($N_f = 2$) looks [1]

$$H = \frac{\lambda N_c M_{\text{KK}}}{54\pi} \left[\left(\frac{27\pi}{\lambda M_{\text{KK}}} \right)^2 \frac{1}{2\rho^2} + \frac{1}{3} M_{\text{KK}}^2 \rho^2 + \frac{2}{3} M_{\text{KK}}^2 (X^4)^2 \right], \quad (3.2)$$

where

$$w_\alpha^i = \rho(t) U_\alpha^i(t). \quad (3.3)$$

Here $\rho(t)$ and $X^4(t)$ are scalar degrees of freedom, and $U(t)$ is a 2×2 unitary matrix degree of freedom. $\rho(t)$ represents dissolved size of the D-brane (which is roughly the size of the baryons), and $X^4(t)$ is a displacement of the D-brane along the holographic direction. The matrix U is nothing but the moduli degrees of freedom appearing in the Skyrme model, and its quantization gives higher spins and isospins.

These three modes provides almost-independent harmonic oscillators, and the quantization results in the following spectra:

$$M = M_0 + \frac{M_{\text{KK}}}{\sqrt{6}} \left[\sqrt{(I/2 + 1)^2 + N_c^2} + 2n_\rho + 2n_{X^4} + 2 \right]. \quad (3.4)$$




Figure 5: A schematic picture of the baryon D-brane on the flavor D-brane. The harmonic oscillator excitation on the baryon D-brane is the size fluctuation ρ and the fluctuation of the D-brane location along the holographic direction X^4 . The horizontal direction is our space x^1, x^2, x^3 .

Here I is the isospin which is equal to the spin in the present case, and n_ρ (n_{X^4}) is a non-negative integer coming from the harmonic oscillator $\rho(t)$ ($X^4(t)$).

The baryon spectrum (3.4), as well as its calculation from the quantum hamiltonian, is quite close to what has been obtained in the soliton quantization approach in the Sakai-Sugimoto model [11].

We again would like to stress that the result is qualitatively robust in the AdS/CFT approach: because the baryon in the gravity dual should be represented by X and w strings, the spectrum should be given by its low energy quantization. So we are inevitably led to the quantum number n_{X^4} which is the oscillation of the D-brane along the holographic directions, and also the quantum number n_ρ which is the fluctuation of the magnitude of the string connecting the baryon D-brane and the flavor D-brane, and also the spin operator U which is the internal orientation of the same string. The coefficients appearing in the mass spectrum formula may differ among holographic models, but its structure should be shared in all the holographic models.

3.3 Universal repulsive core of nucleons

Once the baryon state can be identified within the matrix model degrees of freedom, it is straightforward to calculate the inter-baryon potential. Since the short-distance behavior of the nuclear force is one of the most important problems in nuclear physics, to derive it analytically is a very important issue. As we have the matrix model action

for $A = 2$ at hand, whether it reproduces the empirically-known repulsive core of nucleons would be a good touchstone for the validity of the matrix model approach.

Since the matrix model action (3.1) is explicitly given for $A = 2$, we just need to: first derive the off-diagonal term classically by solving ADHM constraints¹⁰ for a given set of two diagonal entries which defines the locations and the spin/isospins of the two baryons, and then substitute all back into the hamiltonian to derive the inter-baryon potential energy.

The calculation is straightforward and was given in [1]. The result for the inter-nucleon potential is as follows:¹¹

$$V_{\text{central}}(r) = \frac{\pi N_c}{\lambda M_{\text{KK}}} \left(\frac{27}{2} + 8\vec{I}_1 \cdot \vec{I}_2 \vec{J}_1 \cdot \vec{J}_2 \right) \frac{1}{r^2}, \quad (3.5)$$

$$V_{\text{tensor}}(r) = \frac{2\pi N_c}{\lambda M_{\text{KK}}} \vec{I}_1 \cdot \vec{I}_2 \frac{1}{r^2}. \quad (3.6)$$

This short-distance potential is positive for any choice of the spins and the isospins, therefore we conclude for $N_f = 2$ case, there are universal repulsive cores for the nuclear forces at short distance. This repulsive core behaves as $1/r^2$, showing very strong repulsive core at $r \rightarrow 0$ limit. We concluded that qualitatively the matrix model approach for multi-baryon system is consistent with experiments, in this sense. Whether the repulsive core behaves as $1/r^2$ or not should be tested in future.

The three-body nuclear force can be evaluated in the same manner. The short distance contribution to the intrinsic three-body force, which does not come from the effective integration of massive states (for example the famous Fujita-Miyazawa force [34]), is important as it cannot be evaluated using chiral perturbations. Using the matrix model approach, one can straightforwardly evaluate the three-body interaction. It was shown in [8] that the proton-proton-neutron aligned on a line gives a positive three-body potential, and that a spin-averaged three-neutron aligned on a line is positive too. These are consistent with experiments¹². In particular, the latter is relevant for neutron stars as it has a dense neutron system, and the effective repulsion would give a hard equation of state which is a good tendency for a recent observation of heavy neutron stars.

3.4 Toward a description of atomic nuclei

As we have the effective action (3.1) of the multi-baryon system, in principle, atomic nuclei and their properties, *i.e.* the nuclear physics, should emerge from the action. The action (3.1) has only two parameters, so once one can solve the effective action

¹⁰For detail, see section 4.1.2. of [1].

¹¹Using the soliton approach in the holographic D4D8 model, the short distance nuclear force was calculated [15]. The result is qualitatively similar to the result of the matrix model.

¹²See also the recent attempt in lattice QCD [35].



Figure 6: A plot of a nuclear density inside the nucleus, calculated in AdS/CFT [10].

completely in a quantum mechanical fashion, one can compare the results with experiments in principle. Whether this action provides us with an efficient and good description of atomic nuclei is a very important question, as the AdS/CFT connects directly the nuclear physics and QCD.

In particular, properties of heavy nuclei are yet to be uncovered, and they are far from QCD. The aim of the holographic approach is to uncover the relation between the nuclear physics and QCD directly, to make clear how observables in nuclear physics may depend on quantities defined in QCD. One of the important targets in nuclear physics in this sense is the nuclear radius. It has been known for many decades that stable nuclei are subject to a relation

$$r \sim 1.2 \times A^{1/3} \text{ [fm]} \quad (3.7)$$

where A is the mass number (the number of baryons) of the nucleus. This has been explained as a result of the nuclear density saturation: the nucleon density inside nuclei is almost constant and takes a universal value, so the nuclear radius is proportional to $A^{1/3}$.

The repulsive core of nucleons is thought to be a component to explain the A dependence of the nuclear radius in nuclear physics. If a nucleon can be regarded as a hard ball which is almost equivalent to the repulsive core, the total nucleus should have a volume proportional to A , therefore the A dependence follows. Since in the holographic QCD approach the repulsive core was reproduced as explained in the previous subsection, this nuclear radius would be a natural consequence.

In [10], one of the authors (K.H.) together with T. Morita demonstrated that indeed the above (3.7) is reproduced from the quantum mechanical matrix action (3.1), with a certain approximation employed. The result obtained in [10] is an analytic formula for the nuclear radius,

$$\sqrt{r_{\text{mean}}^2} = \frac{3^{5/2} \pi^{2/3}}{2^{5/6} 5^{1/6}} \frac{1}{M_{\text{KK}} \lambda^{2/3} N_c^{1/3}} A^{1/3}. \quad (3.8)$$

A nontrivial point is that the formula has the correct $A^{1/3}$ dependence.

The approximation used for deriving (3.8) is: a large A limit with quenching (ignoring the w degrees of freedom), and a large dimension limit (which is almost equivalent to a mean field limit), in addition to the standard limits taken in holographic QCD, such as the large N_c limit and the strong coupling limit $\lambda \gg 1$. Whether these approximations are appropriate or not should be studied in the future study. However, the message here is that we now have a good starting point (3.1) for calculating various quantities in nuclear physics from QCD.

It would be possible that the action (3.1) itself may be modified to include higher order terms, or that one needs to apply different approximations to (3.1) to correctly derive physical observables from (3.1). Especially higher order terms are crucial when we consider the effect of large A limit. The original action (3.1) is derived at the limit where baryon D-branes are at the bottom of the warped geometry, see Fig. 5. As the size be bigger, we need to consider the effect of curved geometry more precisely and this gives the higher derivative corrections to the action (3.1)¹³. All of those efforts can be a good bridge between QCD and nuclear physics.

4. Multi-Flavor Nuclear Forces Via Holography

4.1 Strangeness and holography

QCD at high density is a final frontier, which is still remains to be unveiled. It is expected that at the core of neutron stars, high density matter would be supplemented with strange quarks, in order to relax the Fermi energy of the ordinary two-flavor matter of neutrons and protons. To judge whether the strangeness really kicks in to the high density core of the neutron stars, we need to know inter-baryon interaction with multi-flavors. It has been known experimentally for many decades that nucleons are accompanied with repulsive cores, the short distance repulsion. To reveal whether there exists the repulsive core even for baryons including strange quark(s) is an indispensable cornerstone to reach the truth in the high density QCD.

The inter-baryon potential is a non-perturbative regime of QCD, even at the short distances of concern. Thus we need to rely on analytic method to solve the QCD approximately. In this section, we shall extend the analysis [1] to the case of multiple flavors $N_f > 2$, to find the short distance properties of the inter-baryon potential.

Another non-perturbative framework of QCD, lattice simulations, recently uncovered interesting features of the short-distance inter-baryon potential for the case of three flavors. It was demonstrated [37, 38] that indeed there remains a repulsive

¹³In addition, one may take a gravity dual of the matrix model under a certain assumption (such that A is large and also that the inter-nucleon distance is small) to investigate giant resonances in nuclei [36].

field	index	$U(k)$	$SU(N_f)$	$SU(2) \times SU(2)$
$X^M(t)$	$M = 1, 2, 3, 4$	adj.	$\mathbf{1}$	$(\mathbf{2}, \mathbf{2})$
$w_{\dot{\alpha}i}(t)$	$\dot{\alpha} = 1, 2; i = 1, \dots, N_f$	\mathbf{k}	\mathbf{N}_f	$(\mathbf{1}, \mathbf{2})$
$A_0(t)$		adj.	$\mathbf{1}$	$(\mathbf{1}, \mathbf{1})$
$D_s(t)$	$s = 1, 2, 3$	adj.	$\mathbf{1}$	$(\mathbf{1}, \mathbf{3})$

Figure 7: Fields in the nuclear matrix model.

core, so the repulsion is universal.¹⁴ In this section, we find that the holographic QCD shows the universal repulsive core for generic states in multi flavors.

4.2 The effective model of multi-baryon system

To extract the non-perturbative potential among baryons at short distances, the nuclear matrix model [1] derived in holographic QCD should provide a good sense of the generic nature. The action of the model is a quantum mechanics,

$$\begin{aligned}
S = & \frac{\lambda N_c M_{\text{KK}}}{54\pi} \int dt \text{tr}_k \left[(D_0 X^M)^2 - \frac{2}{3} M_{\text{KK}}^2 (X^4)^2 \right. \\
& + D_0 \bar{w}_i^{\dot{\alpha}} D_0 w_{\dot{\alpha}i} - \frac{1}{6} M_{\text{KK}}^2 \bar{w}_i^{\dot{\alpha}} w_{\dot{\alpha}i} + \frac{3^6 \pi^2}{4\lambda^2 M_{\text{KK}}^4} (\vec{D})^2 \\
& + \vec{D} \cdot \vec{\tau}^{\dot{\alpha}}_{\dot{\beta}} \bar{X}^{\dot{\beta}\alpha} X_{\alpha\dot{\alpha}} + \vec{D} \cdot \vec{\tau}^{\dot{\alpha}}_{\dot{\beta}} \bar{w}_i^{\dot{\beta}} w_{\dot{\alpha}i} \left. \right] \\
& + N_c \int dt \text{tr}_k A_0. \tag{4.1}
\end{aligned}$$

The system possesses a gauge symmetry $U(k)$ where k is the number of the baryons in the system. The table 1 shows the field content of the model.

Here, the dynamical fields are X^M and $w_{\dot{\alpha}i}$, while A_0 and D_s are auxiliary fields. In writing these fields, the indices for the gauge group $U(k)$ are implicit. The symmetry of this matrix quantum mechanics is $U(k)_{\text{local}} \times SU(N_f) \times SO(3)$ where the last factor $SO(3)$ is the spatial rotation, which, together with a holographic dimension, forms a broken $SO(4) \simeq SU(2) \times SU(2)$ shown in the table. The breaking is due to the mass terms for X^4 and $w_{\dot{\alpha}i}$. In the action, the trace is over these $U(k)$ indices, and the definition of the covariant derivatives is $D_0 X^M \equiv \partial_0 X^M - i[A_0, X^M]$, $D_0 w \equiv \partial_0 w - iwA_0$ and $D_0 \bar{w} \equiv \partial_0 \bar{w} + iA_0 \bar{w}$. The spinor indices of X are defined as $X_{\alpha\dot{\alpha}} \equiv X^M (\sigma_M)_{\alpha\dot{\alpha}}$ and $\bar{X}^{\dot{\alpha}\alpha} \equiv X^M (\bar{\sigma}_M)^{\dot{\alpha}\alpha}$ where $\sigma_M = (i\vec{\tau}, 1)$ and $\bar{\sigma}_M = (-i\vec{\tau}, 1)$, with Pauli matrices τ . The model has a unique scale M_{KK} , and $\lambda = N_c g_{\text{QCD}}^2$ is the 'tHooft coupling constant of QCD, with the number of colors N_c . The diagonal

¹⁴There is a channel at which the repulsive core disappears, for an appropriate choice of the baryon states. It is closely related to the conjectured two-baryon bound state called H-dibaryon [39], as demonstrated in lattice simulations [38, 40]. Whether such dibaryon exists or not should be confirmed by future experiments.

entries of X^i ($i = 1, 2, 3$) specify the location of the baryons. The location of the baryon D-brane in the holographic direction X^4 is stabilized at $X^4 = 0$ around which the harmonic excitations label excited baryon states. The w fields are responsible for spins and isospins (and flavor representations) of each baryon.

In [1], explicitly demonstrated is the nuclear force for the two-flavor case. There, a universal repulsive core was found. We here simply extend the two-flavor calculation to the case with a generic number of massless flavors, and will see the consequence.

The procedure we employ in the following is as follows. First, we look at the configuration which minimizes the potential of the matrix model. When taking a large λ , the D-term condition is required to be satisfied, which is nothing but the ADHM constraint. Then, we obtain a classical potential with a solution of the ADHM constraint, which depends on the inter-baryon distance and the moduli parameters of the two baryons. Taking an expectation value of this potential with respect to the product of the wave functions for each baryon, we obtain the inter-baryon potential.

4.3 Two-baryon configuration

The large λ limit lets only configurations satisfying the ADHM constraint remain. The ADHM constraint is equivalent to the D-term condition concerning D_s , and is given by

$$\vec{\tau}^{\dot{\alpha}\dot{\beta}} \left(\bar{X}^{\dot{\beta}\alpha} X_{\alpha\dot{\alpha}} + \bar{w}^{\dot{\beta}}{}_i w_{i\dot{\alpha}} \right)_{BA} = 0. \quad (4.2)$$

Here A, B are $U(k)$ indices. For a single baryon with generic number of flavors, the ADHM constraint is simply solved by $X = \text{constant}$ (baryons located anywhere), and

$$w = U \begin{pmatrix} \rho & 0 \\ 0 & \rho \\ 0 & 0 \\ \cdots & \cdots \\ 0 & 0 \end{pmatrix}, \quad (4.3)$$

which shows the $(i, \dot{\alpha})$ entry. Here U is a $U(N_f)$ unitary matrix specifying the baryon spin and isospin (flavor dependence). The flavor symmetry acts on U as $U \mapsto GU$. The baryon wave function is given as $\psi(U)$, as in the same manner as the famous Skyrme model.

We want to put two baryons located at $x_M = \pm r_M/2$, so that the distance between the two baryons is r_M . For the two baryons, now the coordinate field X_M is two by two matrices, so we parameterize them as

$$X_M = \frac{1}{2} r_M^a \tau_a \quad (4.4)$$

where τ_a is a Pauli matrix with index (A, B) . We specify the baryon location by the diagonal entries r_M^3 , while assuming the off-diagonal r^1 and r^2 are small as $1/(r^3)$, so that the distance defined by r_3 makes sense at large r^3 . The two baryons can have independent spins and flavor representations, so we allow

$$w^A = U^{(A)} \begin{pmatrix} \rho_A & 0 \\ 0 & \rho_A \\ 0 & 0 \\ \dots & \dots \\ 0 & 0 \end{pmatrix} (\mathbf{1}_{2 \times 2} + \epsilon^{(A)}),$$

for each baryon, $A = 1, 2$. (In this expression we don't make a summation over the index A .) And $\epsilon^{(A)}$ is taken to be a 2×2 traceless matrix at $\mathcal{O}(1/(r^3)^2)$. Note that at the large inter-baryon distance limit $r^3 \rightarrow \infty$, the ADHM data above reduces to just a set of two single-baryon ADHM data, (4.3) and a constant diagonal X .

It is quite straightforward to solve the ADHM constraint (4.2) with the above generic ansatz, and the solution is given as follows.

$$r_M^1 \sigma_M = \frac{-\rho_1 \rho_2}{|r^3|^2} r_M^3 \sigma_M (P_{12} - P_{12}^\dagger), \quad (4.5)$$

$$r_M^2 \sigma_M = \frac{-i \rho_1 \rho_2}{|r^3|^2} r_M^3 \sigma_M (P_{12} + P_{12}^\dagger), \quad (4.6)$$

$$\epsilon^{(1)} = \frac{-\rho_2^2}{4|r^3|^2} [P_{12}, P_{12}^\dagger], \quad \epsilon^{(2)} = \frac{\rho_1^2}{4|r^3|^2} [P_{12}, P_{12}^\dagger]. \quad (4.7)$$

Here we have defined

$$P_{12} \equiv \mathbf{P} [(U^{(1)})^\dagger U^{(2)}], \quad (4.8)$$

with \mathbf{P} being a projection of the $N_f \times N_f$ matrix to its upper-left 2×2 components, so that P_{12} is a 2×2 matrix. We can easily see that, when $N_f = 2$, the result here can reproduce the two-flavor result of [1].

4.4 Explicit inter-baryon potential

Let us substitute the above ADHM data, the two-baryon configuration with the inter-baryon distance r_M^3 and the spin/flavor dependence $U^{(A)}$, into the action (4.1) and derive the inter-baryon potential as a function of r^3 and $U^{(A)}$. As was done in [1], we need to integrate out the $U(2)$ auxiliary gauge field A_0 of the quantum mechanics,

$$A_0 = A_0^0 \mathbf{1}_{2 \times 2} + A_0^a \tau^a. \quad (4.9)$$

Since the model includes only the linear and quadratic terms in A_0 , it is straightforward to perform the integration. In the action (4.1), the terms relevant to A_0

are

$$\begin{aligned}
S_{A_0} = & \frac{\lambda N_c M_{\text{KK}}}{54\pi} \int dt \left[2(A_0^a)^2 (r_M^b)^2 - 2(A_0^a r_M^b)^2 \right. \\
& + ((A_0^0)^2 + A_0^a)^2 (|w^{A=1}|^2 + |w^{A=2}|^2) \\
& + 2A_0^0 A_0^1 (w^{A=1} \bar{w}^{A=2} + w^{A=2} \bar{w}^{A=1}) \\
& - 2i A_0^0 A_0^2 (w^{A=1} \bar{w}^{A=2} - w^{A=2} \bar{w}^{A=1}) \\
& \left. + 2A_0^0 A_0^3 (|w^{A=1}|^2 - |w^{A=2}|^2) + \frac{108\pi}{\lambda M_{\text{KK}}} A_0^0 \right]. \tag{4.10}
\end{aligned}$$

We integrate out all the components¹⁵ of the auxiliary field A_0 and write the potential as $S_{A_0} = -\int dt V_{A_0,2\text{-body}}$. We expand the result in terms of small ρ/r^3 (note that r^3 is the distance between the baryons in x^3 direction, not the cubic power of r !), to obtain the leading term

$$V_{A_0,2\text{-body}} = \frac{27\pi}{4} \frac{N_c}{\lambda M_{\text{KK}}} \frac{1}{(r^3)^2} |\text{tr} P_{12}|^2. \tag{4.11}$$

Here we have already put $\rho_1 = \rho_2 = \rho$ which is ensured at large N_c .

The remaining contributions to the inter-baryon potential, from the matrix model action, is the mass terms $\text{tr}(X_4^2)$ and $|w|^2$. Substituting the two-baryon configuration, we obtain

$$\begin{aligned}
V_{X^4,2\text{-body}} = & \frac{\lambda N_c M_{\text{KK}}^3}{162\pi} \frac{-\rho_1^2 \rho_2^2}{((r^3)^2)^2} \\
& \times \text{tr} \left[r_M^3 \sigma_M P_{12}^\dagger \right] \text{tr} \left[r_M^3 \sigma_M P_{12} \right]. \tag{4.12}
\end{aligned}$$

It turns out that the mass term for w does not give rise to an extra potential.

So, in total, the inter-baryon potential V in the small ρ/r^3 expansion is given as a sum of (4.11) and (4.12),

$$V_{2\text{-body}} = \frac{27\pi N_c}{4\lambda M_{\text{KK}}} \frac{1}{|\vec{r}|^2} \left(|\text{tr} P_{12}|^2 + |\text{tr} [\vec{\hat{r}} \cdot \vec{\tau} P_{12}]|^2 \right). \tag{4.13}$$

Here, we already substituted $r_{M=4}^3 = 0$ which is satisfied by the baryon wave functions at large N_c [1], and denoted $r_{M=1,2,3}^3$ as \vec{r} , the inter-baryon vector. $\vec{\hat{r}}$ is the unit vector along \vec{r} , and we also used the classical size ρ of a single baryon [1], $\rho_1^2 = \rho_2^2 = 3^{7/2}\pi/(\sqrt{2}\lambda M_{\text{KK}}^2)$.

We immediately notice that by taking $N_f = 2$ the potential (4.13) reduces that of the two-flavor inter-nucleon potential given in [1]. The potential has the $1/|\vec{r}|^2$ behavior which is peculiar to the holographic QCD [15, 12, 41], which is nothing but a harmonic potential in 4-dimensional space (our spatial 3 dimensions plus the holographic direction).

¹⁵Note that (4.5) and (4.6) satisfy $r_M^1 r_M^3 = r_M^2 r_M^3 = 0$ which may help reducing the A_0 action.

4.5 Universal repulsive core

It is already manifest that the inter-baryon potential for generic number of flavors, (4.13), is positive-semi-definite, since (4.13) is a sum of two positive semi-definite terms. Therefore, we conclude that holographic QCD predicts positive-semi-definite repulsive core for combination of any two baryon states.

Looking at the magnitude of the potential, we notice the following important fact: As the number of the flavors is larger than 3, the classical potential (4.13) can vanish, for appropriate choice of the baryon state. This is simply because we can choose a set of the unitary matrices $U^{(1)}$ and $U^{(2)}$ such that $\mathbf{P}((U^{(2)})^\dagger U^{(1)})$ vanishes. As the projection operator \mathbf{P} refers only the upper-left corner of the unitary matrices, once the size of the matrix N_f gets larger, the configuration of the baryon can evade the upper-left 2×2 corner, and thus does not contribute to the inter-baryon potential (4.13).

Substituting some particular values of constant U corresponds to a classical evaluation of the potential (as in the same manner as the Skyrme model), but in reality we need to take into account the baryon wave function $\psi_1(U^{(1)})\psi_2(U^{(2)})$. A generic wave function has a wide distribution over the space of the unitary matrices normalized. So the magnitude of the repulsive core depends on the two baryon states. The situation is the same as what has been known for the nucleon case ($N_f = 2$) [1].

In the next section, we review briefly the recent lattice calculations of the inter-baryon potential for three-flavor QCD, and discuss a comparison with our holographic result.

5. A comparison with lattice QCD and OPE

In the previous section, we have calculated a short-distance potential between two baryons in multi-flavor holographic QCD. We have found a universal repulsive potential for generic baryon states. In this section, we shall compare our results with ones obtained by a completely different technique: the lattice QCD.

First, we shall review the results in lattice QCD. Potentials between two octet baryons have been investigated in lattice QCD in the flavor $SU(3)$ symmetric limit [37, 42], where all quark mass in the 3 flavor QCD are artificially taken to be equal, $m_u = m_d = m_s$, with the lattice spacing $a \simeq 0.12$ fm and the spatial extension $L \simeq 2-4$ fm. Simulations employ six different values of quark mass, which correspond to the pseudo-scalar meson mass $m_{\text{PS}} \simeq 470, 670, 840, 1020, 1170$ MeV, where the relation that $m_{\text{PS}}^2 = Am_q$ holds for a small quark mass m_q with a common coefficient A . There are 6 independent potentials between two octet baryons, which correspond

to irreducible representations of the flavor SU(3) group as

$$\mathbf{8} \otimes \mathbf{8} = \underbrace{\mathbf{27} \oplus \mathbf{8}_s \oplus \mathbf{1}}_{\text{symmetric}} \oplus \overbrace{\mathbf{10}^* \oplus \mathbf{10} \oplus \mathbf{8}_a}_{\text{anti-symmetric}}, \quad (5.1)$$

where the first 3 representations are symmetric under the exchange of two octet baryons, while the last 3 are anti-symmetric. To satisfy a condition that a total wave function is odd under exchange of two octet baryons, the first 3 states have spin zero ($S = 0$, odd under the exchange) while the last 3 states must have spin one ($S = 1$, even under the exchange) if the orbital angular momentum between two baryons is zero ($L = 0$).

A typical example of corresponding potentials is shown in Fig. 8, taken from Ref. [37], where the central potentials (left three) and the effective central potential (right three), where an effect of the tensor potential is included, are plotted.

As can be seen from Fig. 8, inter-baryon potentials strongly depend on the representations. In top panels, $V^{(\mathbf{27})}$ and $V^{(\mathbf{10}^*)}$, which correspond to isospin-triplet and isospin-singlet nucleon-nucleon (NN) potentials in the $N_f = 2$ case¹⁶ [3, 4], respectively, have a repulsive core at short distance and an attractive pocket at medium distance. These features qualitatively agree with those of the NN potentials in quenched QCD, shown in Fig. 9. For $L = 0$, $V^{(\mathbf{27})}$ is isospin-triplet ($I = 1$) at $N_f = 2$ and spin-singlet ($S = 0$) while $V^{(\mathbf{10}^*)}$ is isospin-singlet ($I = 0$) at $N_f = 2$ and spin-triplet ($S = 1$). Therefore, the flavor singlet potential at $N_f = 2$ can not have spin-zero for $L = 0$.

On the other hand if the strange quark is introduced in the flavor representation, we have more varieties of potentials: $V^{(\mathbf{10})}$ has a stronger repulsive core and a weaker attractive pocket than $V^{(\mathbf{27})}$, $V^{(\mathbf{10}^*)}$, and $V^{(\mathbf{8}_s)}$ has only a repulsion with the strongest repulsive core among all, while $V^{(\mathbf{8}_a)}$ has a strongest attractive pocket with the weakest repulsive core. In contrast to these five cases, the singlet potential, $V^{(\mathbf{1})}$ shows attraction at all distances without repulsive core, which produces one bound state, the H-dibaryon, in this channel [38, 40]. Note that the flavor singlet potential has spin-zero for $L = 0$ in this case, contrary to the $N_f = 2$ case.

Increasing the number of flavor from 2 to 3, we observe that repulsive core becomes weaker in some channel ($\mathbf{8}_a$) and it even disappears in the singlet ($\mathbf{1}$), as seen in Tab. 5, where we summarize features of inter-baryonic potential in the flavor SU(3) limit.

Now let us discuss a comparison between our holographic QCD results and the lattice QCD results. In the previous section, we have found a universal repulsive core for multi-flavor inter-baryon potential. On the other hand, in the three-flavor lattice QCD, in most of the channels there appears repulsive cores. Therefore we conclude that our holographic results are consistent with the lattice QCD results, generically.

¹⁶This is because the Young tableau of both 27 and 10* do not have three rows.




Figure 8: The 6 independent potentials in the flavor SU(3) limit obtained in lattice QCD at $m_{\text{PS}} = 1014$ MeV (red) and 835 MeV (green) [37].

Only one exception is the existence of an attractive channel. In the lattice QCD result, the flavor-singlet combination of the baryons in the $\mathbf{8}$ representation for $N_f = 3$ is found to have a vanishing repulsive core. In the holographic side, as we work with the large N_c , it is not clear how the lattice QCD with $N_f = N_c = 3$ can be mapped to the holographic QCD. However, in the previous section, we have seen that a classical inter-baryon potential can vanish. So the disappearance of the repulsive core in the lattice QCD is not a contradiction with the holographic QCD. We leave a more detailed comparison to a future work.

In table 5, we summarize qualitative features of baryon-baryon potentials, to-




Figure 9: NN potentials in quenched QCD at $m_\pi \simeq 730$ MeV [4]. The spin singlet sector (1S_0) belongs to the $\mathbf{27}$ representation while the triplet to the $\mathbf{10}^*$ in the flavor $SU(3)$.

gether with the prediction from the operator product expansion (OPE) in perturbative QCD for their short distance behaviors [43, 44, 45]. Although the OPE analysis is consistent with the attractive core for the singlet potential in the lattice QCD, it disagrees with the strong repulsion of the $\mathbf{8}_s$ potential in the lattice QCD.¹⁷ Obviously it is desirable to investigate the short distance behaviors of the inter-baryon potential by various methods, including holographic QCD, in more details.

representation	$\mathbf{27}$	$\mathbf{8}_s$	$\mathbf{1}$	$\mathbf{10}^*$	$\mathbf{10}$	$\mathbf{8}_a$
repulsion	yes	strongest	no	yes	strong	weak
attraction	yes	no	strongest	yes	weak	strong
comment	$NN(I=1)$		H-dibaryon	$NN(I=0)$		
OPE	rpl.	att.	atr.	rpl.	rpl.	atr.

Table 1: Overall feature of inter-baryon potential in each representation. The last line shows the short distance behavior of the potential from OPE, where rpl.=repulsive and atr.=attractive.

Acknowledgments

We would like to thank T. Hatsuda, M. Hidaka, T. Morita, K. Yazaki, and P. Yi. for valuable discussions. S. A. is supported in part by Grant-in-Aid for Scientific Research on Innovative Areas (No. 2004: 20105001,20105003) and by SPIRE (Strategic Program for Innovative REsearch). This research was partially supported by KAKENHI Grant-in-Aid 23654096, 23105716, 22340069, 21105514. K. H. is partly

¹⁷This disagreement may be related to the fact that non-relativistic 6 quark operators are absent in this channel.

supported by the Japan Ministry of Education, Culture, Sports, Science and Technology. N.I. would like to thank mathematical physics laboratory at RIKEN for very kind hospitality where part of this work is done.

References

- [1] K. Hashimoto, N. Iizuka and P. Yi, “A Matrix Model for Baryons and Nuclear Forces,” arXiv:1003.4988 [hep-th].
- [2] E. Witten, “String theory dynamics in various dimensions,” Nucl. Phys. B **443**, 85 (1995) [hep-th/9503124];
T. Banks, W. Fischler, S. H. Shenker and L. Susskind, “M theory as a matrix model: A Conjecture,” Phys. Rev. D **55**, 5112 (1997) [hep-th/9610043].
- [3] N. Ishii, S. Aoki and T. Hatsuda, “The Nuclear Force from Lattice QCD,” Phys. Rev. Lett. **99**, 022001 (2007) [arXiv:nucl-th/0611096].
- [4] S. Aoki, T. Hatsuda and N. Ishii, “Theoretical Foundation of the Nuclear Force in QCD and its applications to Central and Tensor Forces in Quenched Lattice QCD Simulations,” Prog. Theor. Phys. **123**, 89 (2010) [arXiv:0909.5585 [hep-lat]].
- [5] T. Yamazaki, Y. Kuramashi, A. Ukawa, for the PACS-CS Collaboration, “Helium Nuclei in Quenched Lattice QCD,” Phys. Rev. D **81**, 111504 (2010) [arXiv:0912.1383 [hep-lat]].
- [6] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [7] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109];
E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [8] K. Hashimoto and N. Iizuka, “Three-Body Nuclear Forces from a Matrix Model,” arXiv:1005.4412 [hep-th].
- [9] K. Hashimoto and N. Iizuka, “Nucleon Statistics in Holographic QCD : Aharonov-Bohm Effect in a Matrix Model,” Phys. Rev. D **82**, 105023 (2010) [arXiv:1006.3612 [hep-th]].
- [10] K. Hashimoto and T. Morita, “Nucleus from String Theory,” Phys. Rev. D **84**, 046004 (2011) [arXiv:1103.5688 [hep-th]].
- [11] H. Hata, T. Sakai, S. Sugimoto and S. Yamato, “Baryons from instantons in holographic QCD,” Prog. Theor. Phys. **117**, 1157 (2007) [arXiv:hep-th/0701280].

- [12] K. Hashimoto, T. Sakai and S. Sugimoto, “Holographic Baryons: Static Properties and Form Factors from Gauge/String Duality,” *Prog. Theor. Phys.* **120**, 1093 (2008) [arXiv:0806.3122 [hep-th]].
- [13] K. Hashimoto, N. Iizuka and T. Nakatsukasa, “N-Body Nuclear Forces at Short Distances in Holographic QCD,” *Phys. Rev. D* **81**, 106003 (2010) arXiv:0911.1035 [hep-th].
- [14] K. Hashimoto, N. Iizuka, T. Ishii and D. Kadoh, “Three-flavor quark mass dependence of baryon spectra in holographic QCD,” *Phys. Lett. B* **691**, 65 (2010) [arXiv:0910.1179 [hep-th]].
- [15] K. Hashimoto, T. Sakai and S. Sugimoto, “Nuclear Force from String Theory,” *Prog. Theor. Phys.* **122**, 427 (2009) [arXiv:0901.4449 [hep-th]].
- [16] D. K. Hong, M. Rho, H. -U. Yee and P. Yi, “Chiral Dynamics of Baryons from String Theory,” *Phys. Rev. D* **76**, 061901 (2007) [hep-th/0701276 [HEP-TH]]; “Dynamics of baryons from string theory and vector dominance,” *JHEP* **0709**, 063 (2007) [arXiv:0705.2632 [hep-th]]; “Nucleon form-factors and hidden symmetry in holographic QCD,” *Phys. Rev. D* **77**, 014030 (2008) [arXiv:0710.4615 [hep-ph]].
- [17] E. S. Fradkin and M. A. Vasiliev, “On the Gravitational Interaction of Massless Higher Spin Fields,” *Phys. Lett. B* **189**, 89 (1987); “Cubic Interaction in Extended Theories of Massless Higher Spin Fields,” *Nucl. Phys. B* **291**, 141 (1987) ; X. Bekaert, S. Cnockaert, C. Iazeolla and M. A. Vasiliev, “Nonlinear higher spin theories in various dimensions,” hep-th/0503128 ; I. R. Klebanov and A. M. Polyakov, “AdS dual of the critical O(N) vector model,” *Phys. Lett. B* **550**, 213 (2002) [hep-th/0210114]; S. Giombi and X. Yin, “Higher Spin Gauge Theory and Holography: The Three-Point Functions,” *JHEP* **1009**, 115 (2010) [arXiv:0912.3462 [hep-th]]; S. Giombi and X. Yin, “Higher Spins in AdS and Twistorial Holography,” *JHEP* **1104**, 086 (2011) [arXiv:1004.3736 [hep-th]].
- [18] G. Policastro, D. T. Son and A. O. Starinets, “The Shear viscosity of strongly coupled N=4 supersymmetric Yang-Mills plasma,” *Phys. Rev. Lett.* **87**, 081601 (2001) [hep-th/0104066]; P. Kovtun, D. T. Son and A. O. Starinets, “Viscosity in strongly interacting quantum field theories from black hole physics,” *Phys. Rev. Lett.* **94**, 111601 (2005) [hep-th/0405231].
- [19] C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, “Energy loss of a heavy quark moving through N=4 supersymmetric Yang-Mills plasma,” *JHEP* **0607**, 013 (2006) [hep-th/0605158]; S. S. Gubser, “Drag force in AdS/CFT,” *Phys. Rev. D* **74**, 126005 (2006) [hep-th/0605182].
- [20] T. Imoto, T. Sakai and S. Sugimoto, “Mesons as Open Strings in a Holographic Dual of QCD,” *Prog. Theor. Phys.* **124**, 263 (2010) [arXiv:1005.0655 [hep-th]].

- [21] A. Karch and E. Katz, “Adding flavor to AdS / CFT,” JHEP **0206**, 043 (2002) [hep-th/0205236].
- [22] J. Babington, J. Erdmenger, N. J. Evans, Z. Guralnik and I. Kirsch, “Chiral symmetry breaking and pions in nonsupersymmetric gauge / gravity duals,” Phys. Rev. D **69**, 066007 (2004) [hep-th/0306018].
- [23] J. Erdmenger, N. Evans, I. Kirsch and E. Threlfall, “Mesons in Gauge/Gravity Duals - A Review,” Eur. Phys. J. A **35**, 81 (2008) [arXiv:0711.4467 [hep-th]].
- [24] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” Adv. Theor. Math. Phys. **2**, 505 (1998) [hep-th/9803131].
- [25] G. W. Gibbons and K. -i. Maeda, “Black Holes and Membranes in Higher Dimensional Theories with Dilaton Fields,” Nucl. Phys. B **298**, 741 (1988).
- [26] M. Kruczenski, D. Mateos, R. C. Myers and D. J. Winters, “Towards a holographic dual of large N(c) QCD,” JHEP **0405**, 041 (2004) [hep-th/0311270].
- [27] T. Sakai and S. Sugimoto, “Low energy hadron physics in holographic QCD,” Prog. Theor. Phys. **113**, 843 (2005) [arXiv:hep-th/0412141]; “More on a holographic dual of QCD,” Prog. Theor. Phys. **114**, 1083 (2005) [arXiv:hep-th/0507073].
- [28] O. Aharony and D. Kutasov, “Holographic Duals of Long Open Strings,” Phys. Rev. D **78**, 026005 (2008) [arXiv:0803.3547 [hep-th]]; K. Hashimoto, T. Hirayama, F. -L. Lin and H. -U. Yee, “Quark Mass Deformation of Holographic Massless QCD,” JHEP **0807**, 089 (2008) [arXiv:0803.4192 [hep-th]].
- [29] E. Witten, “Baryons and branes in anti-de Sitter space,” JHEP **9807**, 006 (1998) [hep-th/9805112].
- [30] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld and Yu. I. Manin, “Construction of instantons,” Phys. Lett. A **65**, 185 (1978).
- [31] N. Dorey, T. J. Hollowood, V. V. Khoze and M. P. Mattis, “The calculus of many instantons,” Phys. Rept. **371**, 231 (2002) [arXiv:hep-th/0206063].
- [32] T. H. R. Skyrme, “A Unified Field Theory Of Mesons And Baryons,” Nucl. Phys. **31**, 556 (1962); “A Nonlinear field theory,” Proc. Roy. Soc. Lond. A **260**, 127 (1961); “Particle states of a quantized meson field,” Proc. Roy. Soc. Lond. A **262**, 237 (1961).
- [33] G. S. Adkins, C. R. Nappi and E. Witten, “Static Properties Of Nucleons In The Skyrme Model,” Nucl. Phys. B **228**, 552 (1983).
- [34] J. Fujita and H. Miyazawa, “Pion Theory of Three-Body Forces,” Prog. Theor. Phys. **17**, 360 (1957).
- [35] T. Doi [HAL QCD Collaboration], “Three-Nucleon Forces explored by Lattice QCD Simulations,” arXiv:1112.4103 [hep-lat].

- [36] K. Hashimoto, “Holographic Nuclei,” *Prog. Theor. Phys.* **121**, 241 (2009) [arXiv:0809.3141 [hep-th]]; “Holographic Nuclei : Supersymmetric Examples,” *JHEP* **0912**, 065 (2009) [arXiv:0910.2303 [hep-th]].
- [37] T. Inoue *et al.* [HAL QCD Collaboration], “Baryon-Baryon Interactions in the Flavor SU(3) Limit from Full QCD Simulations on the Lattice,” *Prog. Theor. Phys.* **124**, 591 (2010) [arXiv:1007.3559 [hep-lat]].
- [38] T. Inoue *et al.* [HAL QCD Collaboration], “Bound H-dibaryon in Flavor SU(3) Limit of Lattice QCD,” *Phys. Rev. Lett.* **106**, 162002 (2011) [arXiv:1012.5928 [hep-lat]].
- [39] R. L. Jaffe, “Perhaps a Stable Dihyperon,” *Phys. Rev. Lett.* **38**, 195 (1977) [Erratum-*ibid.* **38**, 617 (1977)]. Reviewed in, T. Sakai, K. Shimizu and K. Yazaki, “H dibaryon,” *Prog. Theor. Phys. Suppl.* **137**, 121 (2000) [nucl-th/9912063].
- [40] S. R. Beane *et al.* [NPLQCD Collaboration], “Evidence for a Bound H-dibaryon from Lattice QCD,” *Phys. Rev. Lett.* **106**, 162001 (2011) [arXiv:1012.3812 [hep-lat]].
- [41] Y. Kim, S. Lee and P. Yi, “Holographic Deuteron and Nucleon-Nucleon Potential,” *JHEP* **0904**, 086 (2009) [arXiv:0902.4048 [hep-th]].
- [42] T. Inoue *et al.* [HAL QCD Collaboration], “Two-Baryon Potentials and H-Dibaryon from 3-flavor Lattice QCD Simulations,” arXiv:1112.5926 [hep-lat].
- [43] S. Aoki, J. Balog and P. Weisz, “Application of the operator product expansion to the short distance behavior of nuclear potentials,” *JHEP* **1005**, 008 (2010) [arXiv:1002.0977 [hep-lat]].
- [44] S. Aoki, J. Balog and P. Weisz, “The repulsive core of the NN potential and the operator product expansion,” *PoS LAT2009*, 132 (2009) [arXiv:0910.4255 [hep-lat]].
- [45] S. Aoki, J. Balog and P. Weisz, “Operator product expansion and the short distance behavior of 3-flavor baryon potentials,” *JHEP* **1009**, 083 (2010) [arXiv:1007.4117 [hep-lat]].