

Non-universal gaugino masses and fine tuning implications for SUSY searches in the MSSM and the GNMSSM

Anna Kaminska,^{a,b} Graham G. Ross^a and Kai Schmidt-Hoberg^c

^a*Rudolf Peierls Centre for Theoretical Physics, University of Oxford,
1 Keble Road, Oxford OX1 3NP, U.K.*

^b*DESY,
Notkestrasse 85, D-22607, Hamburg, Germany*

^c*Theory Division, CERN,
1211 Geneva 23, Switzerland*

E-mail: anna.kaminska@desy.de, g.ross1@physics.ox.ac.uk,
kai.schmidt-hoberg@cern.ch

ABSTRACT: For the case of the MSSM and the most general form of the NMSSM (GNMSSM) we determine the reduction in the fine tuning that follows from allowing gaugino masses to be non-degenerate at the unification scale, taking account of the LHC8 bounds on SUSY masses, the Higgs mass bound, gauge coupling unification and the requirement of an acceptable dark matter density. We show that low-fine tuned points fall in the region of gaugino mass ratios predicted by specific unified and string models. For the case of the MSSM the minimum fine tuning is still large, approximately 1:60 allowing for a 3 GeV uncertainty in the Higgs mass (1:500 for the central value), but for the GNMSSM it is below 1:20. We find that the spectrum of SUSY states corresponding to the low-fine tuned points in the GNMSSM is often compressed, weakening the LHC bounds on coloured states. The prospect for testing the remaining low-fine-tuned regions at LHC14 is discussed.

KEYWORDS: Supersymmetry Phenomenology

ARXIV EPRINT: [1308.4168](https://arxiv.org/abs/1308.4168)

Contents

1	Introduction	1
2	The fine tuning measure	3
3	SUSY, Higgs and DM cuts	4
4	The MSSM case	4
4.1	The gaugino focus point — analytic discussion	5
4.2	Numerical analysis of fine tuning in the MSSM	8
4.3	Dark matter in the MSSM	10
5	Beyond the MSSM - the GNMSSM	11
5.1	Operator analysis	11
5.2	The GNMSSM superpotential	12
5.3	Supersymmetry breaking	13
5.4	The gaugino focus point — analytic discussion	13
5.5	Numerical analysis of fine tuning for the GNMSSM	13
5.6	GNMSSM phenomenology	15
5.6.1	Compressed spectra	15
5.6.2	Dark matter	16
5.6.3	Outlook	16
6	Summary and conclusions	18
A	Fine tuning in the MSSM	19

1 Introduction

Low energy supersymmetry (SUSY) was introduced as a way to solve the hierarchy problem, allowing for a consistent separation of the electroweak scale from high scales such as the Grand Unified scale, the string scale and the Planck scale. As a bonus the simplest supersymmetric extension of the Standard Model (SM), the Minimal Supersymmetric Standard Model (MSSM) [1], predicts unification of the gauge couplings (to within a few percent) at a scale $M_X \sim 10^{16}$ GeV [2, 3] and, together with R-parity conservation, provides a good dark matter candidate in the form of a neutralino. In addition radiative breaking naturally explains why the electroweak group and not QCD is spontaneously broken and offers an explanation for the observed scale of electroweak symmetry breaking (EWSB), relating it to the SUSY breaking scale [4–8]. However, despite extensive searches performed at particle accelerators and dark matter detection experiments, no direct evidence for supersymmetry has yet been observed with strong lower bounds on the masses of the coloured SUSY partners of the SM states.

In the context of models that unify the gauge symmetries¹ these bounds already rule out much of the parameter space that has no hierarchy problem, thus losing much of the motivation for low-scale SUSY. The discovery of a Higgs-like state at approximately 125 GeV makes the problem worse in the MSSM because the Higgs mass can only be driven this high through large radiative corrections that in turn require a large average stop mass and/or a large off diagonal entry of the stop mass matrix $X_t = A_t - \mu \cot \beta$, both of which exacerbate the hierarchy problem.

It has been pointed out that the hierarchy problem is reduced in extensions of the MSSM [10], such as the NMSSM [11] with an additional singlet super field and particularly in its generalised version (GNMSSM) [12], see also [13, 14]. This has been explored in detail for the simplified case of universal boundary conditions for the SUSY breaking parameters (CGNMSSM) [15]. However, even allowing for the additional contribution to the Higgs mass coming from the singlet couplings, the parameter space of this model corresponding to low fine tuning has essentially been ruled out by a combination of the non-observation at the LHC of SUSY and dark matter (DM) abundance. In particular the DM abundance has to be reduced below the “over-closure” limit and this is dominantly through stau co-annihilation that is only effective for relatively low m_0 and $m_{1/2}$ and hence for sparticle masses in the reach of LHC8. For more general initial conditions, in particular non-universal gaugino masses, the situation changes because the LSP can now have significant Wino/Higgsino components to ensure its efficient annihilation.

Given the need for large fine tuning in the simplest SUSY schemes, it is perhaps timely to investigate non-universal boundary conditions for gaugino masses. It has been observed that this can reduce the hierarchy problem through the appearance of a new “focus point” that makes the Higgs mass less sensitive to the gaugino mass scale [16–26]. Here we explore this possibility in detail for both the MSSM and GNMSSM in the context of the latest LHC bounds on SUSY states and the measurement of the Higgs mass.

Of course any considerations based on the hierarchy problem must address the question as to how it should be quantified. In this context there has been recent progress [27–30] showing that the “conventional” fine tuning measure of the hierarchy problem follows from a normal likelihood analysis in which the electroweak breaking scale is treated as an observable. Thus minimising the fine tuning measure is elevated from an aesthetic principle to an essential part of a fit to data and a probabilistic interpretation can be given to the magnitude of the fine tuning.

The constraints of Higgs mass, dark matter abundance and its non-detection together with the gaugino focus point restriction (for reduced fine tuning) and the LHC SUSY bounds largely determines the SUSY spectrum and the resulting phenomenology of the GNMSSM with non-universal gaugino masses. In particular the focus point prefers gauginos with similar masses at low energies and, as for the CGNMSSM, rather heavy singlet states. For the case that the squarks are heavier one finds a compressed spectrum for the states accessible at LHC8 that can significantly weaken the bounds on them. As a result there remains a significant low-fine tuned region to be probed at LHC14.

¹Models based on low-energy SUSY that do not address their UV completion have lower fine tuning [9] due to the absence of large logarithmic corrections in the RG running from the UV scale. As we are concerned with models that have gauge coupling unification, we do not consider such low-scale models here.

The paper is organised as follows. In the next section we define the fine tuning measure used in this analysis. In section 3 we discuss the experimental bounds that we use for our numerical analysis. In section 4 we analyse the fine tuning for the case of the MSSM with non-universal gaugino masses both analytically and numerically. In the latter case we consider the limits and resulting phenomenology coming from LHC8 on SUSY states, the Higgs mass range and dark matter. In section 5, after motivating and discussing the structure of the GNMSSM, we perform a similar analysis of its fine tuning. We present a discussion of the phenomenology and discovery potential of the GNMSSM, paying particular regard to the possibility that its spectrum is compressed. A summary and our conclusions are presented in section 6.

2 The fine tuning measure

The fine tuning measure, Δ_p , with respect to a given independent parameter, p , was introduced in [31, 32], with the form

$$\Delta_p \equiv \frac{\partial \ln v^2}{\partial \ln p} = \frac{p}{v^2} \frac{\partial v^2}{\partial p}, \quad (2.1)$$

where v is the electroweak scale.²

In [27–29] it was shown that, if one includes the Z mass as one of the ‘nuisance’ parameters, this measure naturally appears in a likelihood fit in which the electroweak breaking scale, v , is treated as an observable. In this case the likelihood is suppressed by the overall fine tuning measure Δ_q given by

$$\Delta_q = \left(\sum_p \Delta_p^2 \right)^{1/2}. \quad (2.2)$$

Thus we see that reducing the fine tuning measure rather than being an aesthetic requirement emerges naturally from a likelihood fit of SUSY to data.

Usually one term dominates in eq. (2.2) and a more commonly used fine tuning measure is given by

$$\Delta \equiv \max \text{Abs}[\Delta_p]. \quad (2.3)$$

To allow for comparison with previous work we will use Δ when computing the overall fine tuning.

The quantity Δ^{-1} gives a measure of the accuracy to which independent parameters, p , must be tuned to get the correct electroweak breaking scale. In the analysis presented here the parameters correspond to the independent parameters in the superpotential and the soft scalar potential plus the top Yukawa coupling³ all defined at the unification scale and chosen to be of mass dimension 2 where appropriate, e.g. μ^2 . From the connection of the fine tuning measure to the likelihood one finds that an acceptable likelihood requires Δ should be much less than 100 [27–29].

² $v^2 = v_u^2 + v_d^2$ where $v_{u,d}$ are the up and down sector Higgs vacuum expectation values. Here we work in conventions in which $v \simeq 246$ GeV.

³We use the modified definition of fine tuning for the top-Yukawa coupling, appropriate for measured parameters [33].

3 SUSY, Higgs and DM cuts

In this section we briefly describe the cuts that we impose for the numerical analyses of the MSSM and the GNMSSM. As discussed below, the non-universal gaugino mass case in the GNMSSM often leads to a compressed SUSY spectrum with small mass differences between gauginos and the LSP that makes SUSY discovery more difficult. To account for this in a manner consistent with the non-observation of superpartners at the LHC we implemented a cut on the gluino mass as a function of the gluino-LSP mass difference as presented in [34, 35]. In figure 7 of [34] two bounds are shown, a weaker one for decoupled squarks and a stronger one for $m_{\text{squark}} \sim m_{\text{gluino}}$. Most parameter space points of interest to us are in the intermediate regime, but to be sure they are not excluded we will use the stronger bound. We further require the chargino and slepton masses to be above 100 GeV. We also require that the lightest supersymmetric particle (LSP) is a neutralino which is a good dark matter candidate and its relic density does not exceed the 5σ PLANCK [36] upper bound of $\Omega h^2 \leq 0.1334$. While an under-abundance could always be compensated by the relic density of a multitude of other particles, an overabundance would require a deviation from the standard thermal history of the Universe (or at least a sufficiently low reheating temperature, such that the dark matter candidate never reaches thermal equilibrium). In addition to constraints from the relic density there are constraints from dark matter direct detection searches which limit the cross section of the lightest neutralino to nucleons. We require that the points are consistent with dark matter direct detection constraints, in particular with the latest constraint from XENON100 [37]. As we only require dark matter not to be overabundant, the inferred cross section should be multiplied with $(\Omega h^2)^{\text{th}}/(\Omega h^2)^{\text{obs}} = (\Omega h^2)^{\text{th}}/0.1199$ with $(\Omega h^2)^{\text{th}}$ the predicted relic density for the given point in parameter space to account for cases with underabundant neutralinos. Finally, for the Higgs mass we take the average of the CMS and ATLAS best fit values of 125.7 GeV [38] and 125.5 GeV [39] respectively and allow for a 3 GeV uncertainty, $m_h = 125.6 \pm 3$ GeV.

For our numerical analyses we use **SPheno** [40, 41] created by **SARAH** [42–45]. This version performs a complete one-loop calculation of all SUSY and Higgs masses [46, 47] and includes the dominant two-loop corrections for the scalar Higgs masses [48–51]. We have extended **SPheno** by incorporating routines to calculate the fine tuning as presented in [15]. The dark matter relic density as well as the direct detection bounds are calculated with **MicrOmegas** [52–54].

4 The MSSM case

We start with the determination of the fine tuning in the MSSM with non-universal gaugino masses, which we write as $M_1 = a \cdot m_{1/2}$, $M_2 = b \cdot m_{1/2}$ and $M_3 = m_{1/2}$.⁴ All other parameters we take to be (C)MSSM like, i.e. a universal scalar mass and all other soft terms proportional to their corresponding superpotential couplings.

⁴This scenario has recently been studied in [22] based on the 2012 LHC limits. However we include it here as it is important to use exactly the same methods and cuts to calculate the fine tuning when comparing the MSSM and the GNMSSM; our estimate of the fine tuning in the MSSM case is higher than in [22].

4.1 The gaugino focus point — analytic discussion

As pointed out in [20], having solved the RGEs, the soft masses in the Higgs sector at a given energy scale Q can be written as a polynomial in initial parameters

$$\begin{aligned}
 m_{h_u}^2(Q) &= z_{h_u}^{m_0}(Q)m_0^2 + z_{h_u}^{m_{1/2}}(Q)m_{1/2}^2 + z_{h_u}^{A_0}(Q)A_0^2 + 2z_{h_u}^{m_{1/2}A_0}(Q)m_{1/2}A_0 \\
 m_{h_d}^2(Q) &= z_{h_d}^{m_0}(Q)m_0^2 + z_{h_d}^{m_{1/2}}(Q)m_{1/2}^2 + z_{h_d}^{A_0}(Q)A_0^2 + 2z_{h_d}^{m_{1/2}A_0}(Q)m_{1/2}A_0.
 \end{aligned}
 \tag{4.1}$$

Using this equation the relation between the electroweak VEV, v , and the independent parameters defined at the unification scale can be written as [20]

$$\lambda^{(0)}v^2 = -\frac{\tan^2\beta}{\tan^2\beta - 1}\bar{m}_{h_u}^2 + \frac{1}{\tan^2\beta - 1}\bar{m}_{h_d}^2 - |\mu|^2,
 \tag{4.2}$$

where $\lambda^{(0)}$ is the tree-level quartic Higgs coupling and

$$\bar{m}_{h_x}^2 = m_{h_x}^2 + \frac{\partial V^{(1)}}{\partial h_x^2}
 \tag{4.3}$$

with $V^{(1)}$ the one-loop Coleman-Weinberg effective potential.

For the analytic discussion we will for simplicity concentrate on the case with small A_0 such that we can drop the terms proportional to $m_{1/2}A_0$ in eq. (4.1). As we will also be interested in the low $\tan\beta$ regime both $m_{h_u}^2$ and $m_{h_d}^2$ contributions are non-negligible for the derivation of the EW scale. In this case, with

$$z_{h_{ud}}^{m_{1/2}}(Q) = \frac{\tan^2\beta}{\tan^2\beta - 1}z_{h_u}^{m_{1/2}}(Q) - \frac{1}{\tan^2\beta - 1}z_{h_d}^{m_{1/2}}(Q),
 \tag{4.4}$$

the gaugino focus point is defined as the energy scale at which $z_{h_{ud}}^{m_{1/2}}$ vanishes. Threshold corrections to the RGE running are neglected in our analytical estimates.

In order to compute $z_{h_{ud}}^{m_{1/2}}(Q)$ analytically we assume that only the top Yukawa is non-zero. In this case exact solutions to the RGEs in a closed form can be found following [55] and [56]. The results for the gaugino focus point are given in figure 1. For large $\tan\beta$ they are consistent with [20]. For the case that the focus point is of order the electroweak breaking scale the sensitivity of v to the gaugino mass scale $m_{1/2}$ is significantly reduced and so the fine tuning is reduced. Approximate expressions for the fine tuning in the MSSM case are given in the appendix.

It may be seen that the reduction of $z_{h_{ud}}^{m_{1/2}}(v)$ when the gaugino focus point approaches the electroweak scale not only reduces $\Delta_{m_{1/2}}$ but also reduces the fine tuning with respect to μ . This happens because μ^2 is related to $m_{h_u}^2$ and $m_{h_d}^2$ through the electroweak symmetry breaking condition eq. (4.2). Contours of minimal overall fine tuning for large $\tan\beta$, obtained only requiring the existence of viable electroweak breaking and $m_{1/2} > 0.5$ TeV, are shown in figure 2. The situation before and after the Higgs mass cut is presented. The general behavior of fine tuning in both cases is similar but satisfying the Higgs mass cut demands larger values of $m_{1/2}$ and optimal A_0 , which increases the fine tuning. Analytic study shows that in the central part of the considered (a, b) plane Δ_{μ_0} and $\Delta_{m_{1/2}}$ dominate




Figure 1. Analytic results for the gaugino focus point scale contours (in units of TeV) in the MSSM for $\tan\beta = 2, 3, 10$ from left to right. The points and the gray line correspond to specific models with non-universal gaugino masses; see text for details. The contours do not change much for larger $\tan\beta$.

the overall fine tuning. In this region the gaugino focus point is much above the electroweak scale, hence $m_{\tilde{h}_u}^2$ depends strongly on $m_{1/2}$ forcing large negative values of $m_{\tilde{h}_u}^2$ and large values of μ around the TeV scale. As the absolute values of a and b grow, $|z_{h_{ud}}^{m_{1/2}}(v)|$ decreases and both Δ_{μ_0} and $\Delta_{m_{1/2}}$ become smaller. The smallest fine tuning corresponds to values of a and b for which the gaugino focus point scale is close to the electroweak scale. For moderate a , the value $|b| \sim 2.5 - 3$ corresponds to such a low-scale gaugino focus point. The outer region of the (a, b) plane corresponds to the gaugino focus point below the electroweak scale. In this case $z_{h_{ud}}^{m_{1/2}}(v)$ is positive and grows as the absolute values of a and b increase. This means that gauginos give a positive contribution to $m_{\tilde{h}_u}^2$ and values of μ^2 are very small. The fine tuning is dominated by $\Delta_{m_{1/2}}$ and Δ_{A_0} . As the gaugino focus point scale moves further below the electroweak scale the value of m_0 needed to guarantee $m_{\tilde{h}_u}^2 < 0$ becomes large and the fine tuning increases. If we demand that $m_0 < 5$ TeV, values of a and b below the black dashed line in the right panel of figure 2 are excluded.

Figure 3 shows the estimated minimal fine tuning for small values of $\tan\beta$ without imposing the Higgs mass cut (but still requiring $m_{1/2} > 0.5$ TeV). As in the large $\tan\beta$ case, the central region of the (a, b) plane corresponds to large values of μ^2 and the fine tuning with respect to μ dominates, along with $\Delta_{m_{1/2}}$. The smallest fine tuning is again obtained for values of a and b corresponding to the gaugino focus point near the electroweak scale. The large $|b|$ region of the plot is related to small values of μ^2 and the fine tuning becomes dominated by $\Delta_{m_{1/2}}$ growing as the gaugino focus point scale moves away from the electroweak scale. The fine tuning contours do not exactly follow the gaugino focus point scale contours because the $m_{1/2}A_0$ term in $m_{\tilde{h}_u}^2$ and $m_{\tilde{h}_d}^2$ plays a non-negligible role.

Of course, if arbitrary values of the parameters a and b are chosen, the contribution to the overall fine tuning from $\Delta_{a,b}$ should also be included in the analysis above, which typically spoils the improvement in fine tuning. However if a and b are fixed by the underlying theory such contributions are absent. As discussed in [20] values of a and b in the low-focus-point region occur naturally in a variety of models. To illustrate this we show in the first plot of figure 1 the predicted points for the SU(5), SO(10) and E_6




Figure 2. Analytic results for minimal fine tuning in the MSSM in the limit $\tan \beta \rightarrow \infty$ before (left) and after (right) the Higgs mass cut. The region below the black dashed line corresponds to very large scalar masses, $m_0 > 5$ TeV.




Figure 3. Analytic results for minimal fine tuning in the MSSM for $\tan \beta = 2, 3$ without the Higgs mass cut.

GUT models (denoted by circles, squares and diamonds respectively) considered in [57]. GUT models with F terms in **75** or **200** of SU(5), in **210** or **770** of SO(10) and in the corresponding representations of “flipped SO(10)” embedded in E_6 predict gaugino mass ratios in the intermediate and low fine tuning region. Green triangles represent the OII orbifold model for various choices of the discrete Green Schwarz parameter, δ_{GS} [58]. The values $\delta_{GS} = -5, -6, -7$ are optimal from the point of view of fine tuning. For comparison we also show points relevant for mirage mediation, where soft terms receive comparable contributions from gravity (modulus) and anomaly mediated SUSY breaking. In this case gaugino masses at the GUT scale have the following form

$$M_a = m_{3/2} (\varrho + b_a g_a^2) \tag{4.5}$$

where g_a is the relevant gauge coupling, b_a is its β -function coefficient, while ϱ describes the relation between modulus and anomaly mediated contributions. This prescription for gaugino masses as a function of ϱ generates the gray line in figure 1 in the (a, b) parameter space. If ϱ is a continuous parameter there should be an additional contribution Δ_ϱ to the overall fine tuning. However specific string models fix the value of ϱ . Four examples




Figure 4. Dependence of the fine tuning on the Higgs mass in the MSSM. The light red points are before any cuts while the dark red points take into account cuts on the SUSY masses and the relic neutralino abundance. The left plot is uniform in the density of the input parameters, their density reflects the likelihood for finding a viable point. The right plot shows additional points where we zoomed into regions of small fine tuning. The minimal fine tuning after all cuts is about 60 for a Higgs mass of 122.6 GeV; requiring universal gaugino masses, i.e. $a = b = 1$, it is about 350.

are shown in figure 1 by the blue inverted triangles: (i) the minimal setup of KKLT-type moduli stabilization in type II B string theory [59–61], (ii) a model with vacuum uplifting via hidden sector matter superpotentials [59] (iii) and (iv) the Mini Landscape of orbifold compactifications in heterotic string theory [24] with SU(4) and SU(5) hidden sector gauge groups; the type II B string theory model with vacuum stabilisation by F-terms of hidden sector matter superpotentials predicts values of a and b in the low fine tuning region.

4.2 Numerical analysis of fine tuning in the MSSM

We turn now to the full numerical analysis of fine tuning in the (C)MSSM for the case of non-universal gaugino masses. After imposing the electroweak symmetry breaking conditions the independent parameters, defined at the unification scale, are m_0 , A_0 , $\tan \beta$, $\text{Sign } \mu$, $m_{1/2}$, a , b . We compute the overall fine tuning, Δ , corresponding to these parameters apart from the contribution from a and b assuming, as discussed above, that they are fixed in the underlying theory. The results of a broad scan of the MSSM parameter space are shown in figure 4. We show the results both before and after any cuts are made as described in section 3.

In order to achieve a Higgs mass in agreement with experimental results large mixing corresponding to large A-terms is needed. Note that the fine tuning measure is large; even allowing for ± 3 GeV in the Higgs mass about its central value of 125.6 GeV the smallest fine tuning we find after cuts on the SUSY spectrum but before the cut on the Higgs mass is about 30 but after the cut on the Higgs mass it is about 60 (if we were to take the central value without error the smallest fine tuning is about 500!). Although still problematic, this is significantly smaller than the case of degenerate gaugino masses at the unification scale where, using the same method of evaluation, the *smallest* fine tuning for a 122.6 GeV Higgs is about 350.




Figure 5. Dependence of the fine tuning on the input parameters a and b with the additional parameters chosen such that at each point the smallest fine tuning is realised. The upper left plot is after SUSY and DM but before the Higgs mass cut - the upper right plot includes the cut on the Higgs mass. It can be seen that this significantly increases the fine tuning, as expected. To compare with the analytic estimate the plots in the lower row show the same but instead of the overall fine tuning (which is typically dominated by the fine tuning with respect to μ) the fine tuning with respect to the gaugino mass parameter $m_{1/2}$ is shown (note the change of scale). The corresponding ellipse is very close to the analytic estimate.

Of course it is important to check the values of the gaugino mass ratios, a and b , needed to get low fine tuning. This is shown in figure 5. For every shown point we select the smallest fine tuning, i.e. the additional parameters are chosen such that the smallest fine tuning is realised. To connect with our analytical discussion above we not only show the overall fine tuning, but also the fine tuning with respect to the gaugino mass parameter $m_{1/2}$. We see that the smallest fine tuning is realised in an elliptic structure in the a - b plane. Comparing with figure 1 we see that this structure is the one expected on the basis of the gaugino fixed point. For the fine tuning with respect to $m_{1/2}$ we see that very small values of the fine tuning are possible, $\Delta_{m_{1/2}} \ll 10$. For the overall fine tuning, which is larger and typically dominated by the fine tuning with respect to μ , there is a somewhat broader minimum ring. For moderate a and before the Higgs mass cut, the optimal value for the magnitude of the Wino to gluino mass at the unification scale is of $|b| = \mathcal{O}(3 - 4)$. In the simplified analytic calculation of section 4.1 (which neglects A terms) this region corresponds to a gaugino focus point below 10 TeV. Once the Higgs mass cut is imposed the low-fine tuned region requires a and b to lie in the first quadrant, usually with large a . This is due to the fact that A -terms are also sensitive to a, b and optimal mixing is required for the Higgs mass to be in agreement with experimental results. The numerical results for fine tuning are in agreement with the approximate analytic predictions of figure 2.




Figure 6. Fine tuning in the gluino-squark and gluino-LSP plane for the MSSM after all cuts.

In figure 6 we show the fine tuning in the gluino-squark and gluino-LSP plane. Due to the additional flexibility in the gaugino sector there is the interesting possibility of compressed spectra where the mass difference between the gluino and the lightest neutralino is not large. While in the CMSSM the ratio between $M_1 : M_2 : M_3$ is about $1 : 2 : 6$, for non-universal gaugino masses the Wino and bino can be as heavy as the gluino where we expect $M_1 \sim M_2 \sim M_3$ for $a \sim 6$ and $b \sim 3$. The only other relevant parameter is the μ term, which sets the mass for the Higgsino like neutralinos. In the MSSM, however, it turns out that in order to achieve the required Higgs mass the gluino mass is much larger than μ , and compressed spectra are typically not realised. This can also be seen in figure 6 as there are no points close to the diagonal.

4.3 Dark matter in the MSSM

Due to the additional flexibility in the gaugino sector, a large variety of LSP compositions is possible. To infer something about the typical phenomenology we concentrate on regions corresponding to the smallest fine tuning. As there are no points with particularly small fine tuning for the MSSM case we look at viable points with a fine tuning below 300. Also, in order to ascribe meaning to the density of points, we only show points from the scan with a uniform density in the input parameters. In figure 7 we show the direct detection cross section vs. the mass of the lightest neutralino together with the latest bound from XENON100 [37] as well as the dark matter composition as a function of the relic density. As has been observed in previous studies [62, 63], it may be seen that the mass is in the TeV range, most often Higgsino like. It can be seen that all of the points are below the XENON100 direct detection limit. Regarding the composition we see that for points satisfying the relic abundance upper bound the LSP is mainly composed of Wino and Higgsino, with typically only a very small bino component. For the case of a Wino-like LSP we see that the relic abundance is always below the one required for dark matter. On the other hand the correct relic abundance is often achieved with a Higgsino like LSP.




Figure 7. Scaled (i.e. multiplied by the ratio of the density of SUSY dark matter to the observed dark matter density, $(\Omega h^2)^{\text{th}}/0.1199$) dark matter direct detection cross section as a function of the neutralino mass together with the latest bound from XENON100 [37]. Also shown is the dark matter composition, i.e. the bino, Wino and Higgsino fraction of the LSP, as a function of the relic density. Red points correspond to a mainly Wino-like LSP while green points correspond to a mainly Higgsino like LSP. For all red and green points, in addition to the SUSY and Higgs cuts, a fine tuning cut, $\Delta < 300$, was imposed. The grey points are before the fine tuning cut.

5 Beyond the MSSM - the GNMSSM

5.1 Operator analysis

Given the present lack of evidence for the MSSM at the LHC and its associated fine tuning problem it is important to ask if alternative SUSY extensions of the SM have lower fine tuning. A useful way to look for such extensions is to allow for a general modification of the MSSM by adding higher dimension operators that correspond to the effective field theory that results from integrating out additional heavy degrees of freedom and ask if such operators can reduce fine tuning. There is a unique leading dimension 5 operator with the form [10]

$$L = \frac{1}{M_*} \int d^2\theta f(X)(H_u H_d)^2, \tag{5.1}$$

where $X = \theta\theta m_0$ and m_0 is the SUSY breaking scale in the visible sector.

This gives contributions to the scalar potential of the form

$$V = (|h_u|^2 + |h_d|^2) (\chi_1 h_u h_d + h.c.) + \frac{1}{2} (\chi_2 (h_u h_d)^2 + h.c.) \tag{5.2}$$

where $\chi_1 = 2f(0)\mu_{\text{eff}}/M_*$, $\chi_2 = -2f'(0)m_0/M_*$ and μ_{eff} is the effective μ term including the singlet contribution.

Note that the χ_1 term is supersymmetric so there are associated corrections involving Higgsinos that will generate Higgsino mass terms of the same order of magnitude as the correction to the Higgs mass terms (once the Higgs acquire their vevs). However in practice these corrections are going to be of $\mathcal{O}(10 \text{ GeV})$, important to get a Higgs mass of 125 GeV but small compared to the Higgsino mass coming from the μ_{eff} term which will have to be of $\mathcal{O}(1 \text{ TeV})$. For this reason we concentrate on the effect in the scalar sector.

The fine tuning of this model has been analysed in [10] where it was shown that the fine tuning is significantly reduced by the first term of eq. (5.2) while the second term only gives a modest reduction. The reason for this is because the dominant effect comes from the contribution of eq. (5.2) to the Higgs mass after electroweak breaking and, due to the fact that the first term involves an extra power of h_u , it gives the larger contribution.

The obvious question is what new physics can give rise to the first operator corresponding to this term. The answer is through the integration out of a new heavy gauge singlet or SU(2) triplet superfield coupling to the Higgs sector. Interestingly the operator is *not* generated in the NMSSM, the simplest singlet extension of the MSSM, as it requires an explicit mass term for the singlet super field. We refer to the model with the singlet mass term as the generalised NMSSM (the GNMSSM).

5.2 The GNMSSM superpotential

The most general extension of the MSSM by a gauge singlet chiral superfield consistent with the SM gauge symmetry has a superpotential of the form

$$\mathcal{W} = \mathcal{W}_{\text{Yukawa}} + \frac{1}{3}\kappa S^3 + (\mu + \lambda S)H_u H_d + \xi S + \frac{1}{2}\mu_s S^2 \tag{5.3}$$

$$\equiv \mathcal{W}_{\text{NMSSM}} + \mu H_u H_d + \xi S + \frac{1}{2}\mu_s S^2 \tag{5.4}$$

where $\mathcal{W}_{\text{Yukawa}}$ is the MSSM superpotential generating the SM Yukawa couplings and $\mathcal{W}_{\text{NMSSM}}$ is the “normal” NMSSM with a \mathbb{Z}_3 symmetry. Throughout this article capital letters refer to superfields while small letters refer to the corresponding scalar component. One of the dimensionful parameters can be eliminated by a shift in the vev v_s . We use this freedom to set the linear term in S in the superpotential to zero, $\xi = 0$.

The form of eq. (5.4) seems to make the hierarchy problem much worse as the SM symmetries do not prevent arbitrarily high scales for the dimensionful mass terms. However these terms can be naturally of order the SUSY breaking scale if there is an underlying \mathbb{Z}_4^R or \mathbb{Z}_8^R symmetry [64, 65]. Before SUSY breaking the superpotential is of the NMSSM form. However after supersymmetry breaking in a hidden sector with gravity mediation soft superpotential terms are generated but with a scale of order the supersymmetry breaking scale in the visible sector characterised by the gravitino mass, $m_{3/2}$. With these the renormalisable terms of the superpotential take the form [65]

$$\mathcal{W}_{\mathbb{Z}_4^R} \sim \mathcal{W}_{\text{NMSSM}} + m_{3/2}^2 S + m_{3/2} S^2 + m_{3/2} H_u H_d, \tag{5.5}$$

$$\mathcal{W}_{\mathbb{Z}_8^R} \sim \mathcal{W}_{\text{NMSSM}} + m_{3/2}^2 S \tag{5.6}$$

where the \sim denotes that the dimensional terms are specified up to $\mathcal{O}(1)$ coefficients. Clearly the \mathbb{Z}_4^R case is equivalent to the GNMSSM. After eliminating the linear term in S the \mathbb{Z}_8^R case gives a constrained version of the GNMSSM with $\mu_s/\mu = 2\kappa/\lambda$.

Note that the SUSY breaking also breaks the discrete R symmetry but leaves the subgroup \mathbb{Z}_2^R , corresponding to the usual matter parity, unbroken. As a result the lightest supersymmetric particle, the LSP, is stable and a candidate for dark matter.

5.3 Supersymmetry breaking

The general soft SUSY breaking terms associated with the Higgs and singlet sectors are

$$V_{\text{soft}} = m_s^2 |s|^2 + m_{h_u}^2 |h_u|^2 + m_{h_d}^2 |h_d|^2 + \left(b\mu h_u h_d + \lambda A_\lambda s h_u h_d + \frac{1}{3} \kappa A_\kappa s^3 + \frac{1}{2} b_s s^2 + \xi_s s + h.c. \right). \quad (5.7)$$

Note that the shift in the vev v_s that is used to eliminate the linear term in the superpotential does not imply that the corresponding soft term ξ_s is zero as well.

These terms and the soft breaking terms associated with the squarks, sleptons and gauginos depend on the details of the supersymmetry breaking sector. Here we will assume underlying GUT relations with non-universal gaugino masses which, as above, we write as $M_1 = a \cdot m_{1/2}$, $M_2 = b \cdot m_{1/2}$ and $M_3 = m_{1/2}$. All other parameters we take to be CMSSM like, i.e. a universal scalar mass and all other soft terms proportional to their corresponding superpotential couplings.

The independent supersymmetry breaking parameters are therefore m_0 , M_1 , M_2 , M_3 , A_0 , B_0 and ξ_s where A_0 and B_0 are the constants of proportionality associated with the trilinear and bilinear terms respectively. These parameters are defined at the unification scale, M_X , and must be evaluated at low scales using the renormalisation group running. Taking into account the supersymmetric parameters as well, the supersymmetry breaking scheme is specified by the following set of parameters μ , μ_s , λ , κ , m_0 , $m_{1/2}$, a , b , A_0 , B_0 and ξ_s . Trading B_0 , ξ_s and μ for v , $\tan\beta$ and v_s via the EWSB conditions, there are ten parameters defining this model.

5.4 The gaugino focus point — analytic discussion

As before it is instructive to determine the gaugino focus point analytically. Following section 4.1 we assume that only the top quark Yukawa coupling is non-zero. Due to the effect of the singlet-Higgs Yukawa coupling, λ it is not possible to get an exact solution for the RGE running. Fortunately it is possible to find very good approximate solutions by means of truncated iterations, as suggested in [56]. In the analytic study we also neglect the coupling κ as it does not directly affect the Higgs doublets. The results for the gaugino focus point are given in figure 8. The points shown in the first plot correspond to specific models predicting non-universal gaugino masses, as explained in section 4.1. The general structure is similar to that of the MSSM showing that the focus point is relatively insensitive to λ .

5.5 Numerical analysis of fine tuning for the GNMSSM

In the following we will present the results of our full loop-level numerical analysis. Note that this analysis goes beyond the operator analysis of [20] as we do not require that the singlet mass is large and thus, in general, it cannot be integrated out. We are particularly interested in regions which allow for a rather large Higgs mass. The largest Higgs masses can be achieved when the additional tree-level contribution to the Higgs mass is large, corresponding to large λ , (which implies smallish κ [66]) and small $\tan\beta$. We randomly scan over all the free parameters within this region. In the following the fine tuning is calculated with respect to all independent high scale input parameters.




Figure 8. Analytic results for the gaugino focus point scale (in units of TeV) in the GNMSSM for $\lambda = 1.5$ (at the GUT scale) and $\tan \beta = 2, 3, 10$ from left to right. The contours do not change much for larger $\tan \beta$. The points and line shown are the same as shown in figure 1.




Figure 9. Dependence of the fine tuning on the lightest Higgs mass. The light blue points are before any cuts while the dark blue points take into account cuts on the SUSY masses and the relic neutralino abundance. The left plot is uniform in the density of the input parameters, their density reflects the likelihood for finding a viable point. The right plot shows additional points where we zoomed into regions of small fine tuning. The minimal fine tuning after the cuts were imposed is below 20; requiring universal gaugino masses, i.e. $a = b = 1$, it is about 100.

Figure 9 shows the fine tuning as a function of the Higgs mass m_h . The minimal fine tuning after the cuts were imposed is below 20;⁵ requiring universal gaugino masses, i.e. $a = b = 1$, it is about 100. Thus we see that there are significant areas of low fine tuning remaining to be explored. Comparing with figure 8 shows that the low fine tuning region corresponds to the gaugino focus point scale close to the electroweak scale for small $\tan \beta$. This pattern is similar to the analytically obtained fine tuning contours in figure 3 corresponding to the low $\tan \beta$ MSSM case without the Higgs mass cut.

⁵If we strengthen the cut to include only points whose relic abundance lies within the PLANCK range, $0.1064 \leq \Omega h^2 \leq 0.1334$ [36], the minimal fine tuning is still around 20.




Figure 10. Dependence of the fine tuning on the input parameters a and b with the additional parameters chosen such at each point that the smallest fine tuning is realised. The upper left plot is after SUSY and DM but before the Higgs mass cut - the upper right plot includes the cut on the Higgs mass. Note that due to the smaller fine tuning in the GNMSSM the scale is different from the corresponding MSSM figures. The lower plots show the smallest mass difference in GeV between the gluino and the neutralino LSP with the same cuts as in the upper figures. It can be seen that this mass difference can be very small in the regions of low fine tuning, corresponding to very compressed spectra.

5.6 GNMSSM phenomenology

To infer something about the typical phenomenology of the low fine tuned regions we show the fine tuning in the gluino-squark and gluino-LSP planes in figure 11. It can be seen that points with fine tuning below 100 can have gluino masses beyond 2 TeV and squark masses around 3 TeV.

5.6.1 Compressed spectra

As in the MSSM there is the interesting possibility of compressed spectra due to the flexibility in the mass structure of the gaugino sector. In the MSSM this typically does not happen because often there is a Higgsino like neutralino considerably lighter than the gluino. However in the GNMSSM the effective μ term is often close to M_3 so the LSP mass is close to that of the gluino. The resulting compression can be seen in the lower panels of figure 10 as well as in the right panel of figure 11. For the case of heavy squarks the LHC signal is dominantly gluino pair production and decay but this is significantly reduced for the case of compressed spectra.




Figure 11. Fine tuning in the gluino-squark and gluino-LSP plane for the MSSM after all cuts.

5.6.2 Dark matter

In the universal scalar mass case considered here the singlet states are always heavy and thus play no role in the low-energy phenomenology. Their dominant effect is the change to the Higgs mass that reduces the fine tuning, as was found in the CGNMSSM with universal gaugino masses. However the region of parameter space of the CGNMSSM that solves the little hierarchy problem has essentially been ruled out by a combination of LHC non-observation of SUSY and dark matter abundance. In particular the dark matter abundance has to be reduced below the “over-closure” limit and this is dominantly through stau co-annihilation that is only effective for relatively low m_0 and $m_{1/2}$ and hence requires sparticle masses in the reach of LHC8 [15]. For the case of non-universal gaugino masses the situation changes because the LSP can now have a significant non-bino component that allows for its efficient annihilation. As a result the constraint on m_0 and $m_{1/2}$ is relaxed and there remains a significant part of the low-fine-tuned parameter space to be tested at LHC14.

What about the prospect of direct detection of dark matter? In figure 12 we plot, for relatively low-fine-tuned points, the effective direct detection cross section versus the mass of the lightest neutralino. Also shown is the latest bound from XENON100 [37] as well as the dark matter composition as a function of the relic density. In order to ascribe meaning to the density of points, we only show points from the scan with a uniform density in the input parameters. It can be seen that all of the points are below the XENON100 direct detection limit. Regarding the composition, we see that for the correct relic density or an underabundance the LSP is mainly composed of Wino and Higgsino, with typically only a very small bino component. As in the MSSM the correct relic abundance seems to be more easily achieved with a Higgsino like LSP.

5.6.3 Outlook

Our analysis has concentrated on the more fine-tuned UV complete case with parameters defined at the unification scale and requiring gauge coupling unification. Even so it is clear from the figures that models with non-universal gaugino masses still have a sizable region of parameter space with reasonably small fine tuning that remains to be tested. While LHC14




Figure 12. (i) The dark matter direct detection cross section as a function of the neutralino mass. It has been scaled (i.e. multiplied with $(\Omega h^2)^{\text{th}}/0.1199$) to account for cases with underabundant neutralinos. Also shown is the latest bound from XENON100 [37]. (ii) The dark matter composition, that is the bino, Wino and Higgsino fraction of the LSP, as a function of the relic density. Mostly bino-like LSPs are shown in blue, mostly Wino-like LSPs are shown in red and mostly Higgsino like LSPs are shown in green, where *mostly* means a fraction > 0.5 . (iii) The distribution of bino-, Wino-, and Higgsino like LSPs in the a - b plane. For all points, in addition to the SUSY and Higgs cuts, a fine tuning $\Delta < 100$ was imposed. The grey points are before the fine tuning cut.

will cover most of this region, some points lie beyond even its reach, particularly for the case of a compressed spectrum. In the context of gauge coupling unification an interesting feature of the low fine-tuned regions around $b \sim 3 - 4$ is that within these regions the precision of gauge coupling unification is much better than in the case of universal gaugino masses, see e.g. [67].

Similar to the case of the CGNMSSM we find that viable points have a large supersymmetric singlet mass parameter, leading to heavy singlet states. In detail this constraint comes from the need to achieve acceptable electroweak breaking consistent with the universality of scalar masses at the high scale and the observed Higgs mass. Indeed this is also why universal scalar masses are not possible in the NMSSM. Allowing for non-universal Higgs masses solves this problem for the NMSSM and in the case of the GNMSSM it will allow for light singlet states that can have interesting phenomenological implications within the GNMSSM [68, 69]. The case of more general boundary conditions will be considered elsewhere.

6 Summary and conclusions

The non-observation, to date, of evidence for the existence of supersymmetric partners of the SM states has cast doubt on the viability of a supersymmetric solution to the hierarchy problem. To quantify this it is useful to use a fine-tuning measure that determines the precision of cancellation needed between uncorrelated parameters. Through the connection of the measure to a likelihood fit to the data one obtains an upper bound significantly less than 100 on the fine-tuning measure, Δ , that is consistent with an acceptable SUSY fit to the data. For the case that one does not require a UV completion of the SUSY model the fine tuning is minimised because of the absence of large renormalisation group logarithmic terms. In this case, even for top squarks with TeV masses, perfectly acceptable values, $\Delta \sim 10$, are obtained for the MSSM [9]. However if one requires a (GUT) UV completion and the associated gauge coupling unification the fine tuning in the CMSSM becomes unacceptable, $\Delta \geq 350$, largely because it is difficult in the CMSSM to get a Higgs as heavy as 126 GeV.

In this paper we have determined the fine tuning for a simple extension of the CMSSM, that allows for non-universal gaugino masses at the unification scale, which preserves gauge coupling unification and is consistent with the LHC bounds on SUSY masses, a 126 GeV Higgs and acceptable dark matter abundance. We found that, due to a gaugino “focus point”, the fine-tuning is considerably reduced with a minimum $\Delta \sim 60$, just acceptable. We also determined the fine tuning for a further extension of the model, the (C)GNMSSM, that includes a singlet supermultiplet. In this case, due mostly to the additional contribution to the Higgs mass coming from the singlet F-term, the fine tuning is further reduced and can be as low as $\Delta = 20$, perfectly acceptable. This means that, even after the LHC8 results, simple SUSY models can still provide a solution to the hierarchy problem that preserves the successful prediction following from gauge coupling unification.

For the case of the (C)MSSM the regions of parameter space with $\Delta < 100$ have SUSY masses in the reach of LHC14 and a spectrum that gives the usual large missing energy signatures of SUSY. As a result the model will be fully tested by LHC14. However for the (C)GNMSSM there is a significant region of parameter space for which the spectrum is compressed giving signals below the present range of sensitivity of the LHC experiments. Moreover there are regions with quarks and gluinos beyond the reach of LHC14. As a result, while LHC14 will probe most of the remaining low-fine-tuned region of the (C)GNMSSM parameter space with $\Delta < 100$, it will not be able to cover the entire region. In both the (C)MSSM and the (C)GNMSSM models Higgsino like dark matter is favoured and the direct dark matter cross section can be significantly below the XENON100 bound.

Acknowledgments

We would like to thank Vinzenz Maurer for correspondence and Florian Staub for discussions. The research presented here was partially supported by the EU ITN grant UNILHC 237920 (Unification in the LHC era). One of us (GGR) would like to thank the Leverhulme foundation for the award of an emeritus fellowship that also supported this research.

A Fine tuning in the MSSM

In the limit of large $\tan\beta$ the fine tuning in the MSSM with respect to initial parameters is given approximately by

$$\begin{aligned}
\Delta_{\mu_0} &\approx -\frac{4\mu^2}{m_Z^2} \frac{1}{1+\delta} \\
\Delta_{m_0} &\approx -\frac{4m_0^2}{m_Z^2} z_{h_u}^{m_0}(v) \frac{1}{1+\delta} \\
\Delta_{m_{1/2}} &\approx -\frac{4m_{1/2}^2}{m_Z^2} z_{h_u}^{m_{1/2}}(v) \frac{1}{1+\delta} - \frac{4m_{1/2}A_0}{m_Z^2} z_{h_u}^{m_{1/2}A_0}(v) \frac{1}{1+\delta} \\
\Delta_{A_0} &\approx -\frac{4A_0^2}{m_Z^2} z_{h_u}^{A_0}(v) \frac{1}{1+\delta} - \frac{4m_{1/2}A_0}{m_Z^2} z_{h_u}^{m_{1/2}A_0}(v) \frac{1}{1+\delta}
\end{aligned} \tag{A.1}$$

where

$$\delta \approx \frac{3h_t^4}{g^2\pi^2} \left[\log \frac{m_{\tilde{t}}}{m_t} + \frac{1}{2} \frac{(A_t - \mu/\tan\beta)^2}{m_{\tilde{t}}^2} \left(1 - \frac{(A_t - \mu/\tan\beta)^2}{12m_{\tilde{t}}^2} \right) \right]. \tag{A.2}$$

Taking into account that

$$\frac{1}{2}m_Z^2 \approx - (m_{H_u}^2 + \mu^2) \frac{1}{1+\delta} \tag{A.3}$$

the fine tuning with respect to μ_0 , which is often dominating, can be approximated by

$$\Delta_{\mu_0} \approx 2 + \frac{4m_{h_u}^2}{m_Z^2} \frac{1}{1+\delta} \tag{A.4}$$

which shows that reducing $m_{h_u}^2$ by the gaugino focus point helps not only with lowering $\Delta_{m_{1/2}}$, but also with Δ_{μ_0} . The approximate value of $m_{h_u}^2$ at the EW scale is given by

$$\begin{aligned}
m_{h_u}^2(v) &\approx -0.108A_0^2 - 0.030m_0^2 \\
&\quad + (-2.024 - 0.027a + 0.006a^2 - 0.182b - 0.006ab + 0.214b^2) m_{1/2}^2 \\
&\quad + (0.320 + 0.013a + 0.078b) m_{1/2}A_0
\end{aligned} \tag{A.5}$$

which shows that the fine tuning with respect to m_0 will usually play a negligible role, while the FT both with respect to $m_{1/2}$ and A_0 will vary in the (a, b) plane.

For low $\tan\beta$, neglecting the dependence of $\tan\beta$ on initial parameters and noticing that $z_{h_d}^A = z_{h_d}^{MA} = 0$, the fine tuning can be approximated by

$$\begin{aligned}
\Delta_{\mu_0} &\approx -\frac{4\mu^2}{m_Z^2} \frac{\tan^2\beta - 1}{\tan^2\beta - 1 + \delta \tan^4\beta / (1 + \tan^2\beta)} \\
\Delta_{m_0} &\approx -\frac{4m_0^2}{m_Z^2} \frac{\tan^2\beta z_{h_u}^{m_0}(v) - z_{h_d}^{m_0}(v)}{\tan^2\beta - 1 + \delta \tan^4\beta / (1 + \tan^2\beta)} \\
\Delta_{m_{1/2}} &\approx -\frac{4}{m_Z^2} \frac{m_{1/2}^2 \left(\tan^2\beta z_{h_u}^{m_{1/2}}(v) - z_{h_d}^{m_{1/2}}(v) \right) - m_{1/2}A_0 \tan^2\beta z_{h_u}^{m_{1/2}A_0}(v)}{\tan^2\beta - 1 + \delta \tan^4\beta / (1 + \tan^2\beta)} \\
\Delta_{A_0} &\approx -\frac{4}{m_Z^2} \frac{A_0^2 \tan^2\beta z_{h_u}^{A_0}(v) - m_{1/2}A_0 \tan^2\beta z_{h_u}^{m_{1/2}A_0}(v)}{\tan^2\beta - 1 + \delta \tan^4\beta / (1 + \tan^2\beta)}.
\end{aligned} \tag{A.6}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Dimopoulos and H. Georgi, *Softly Broken Supersymmetry and SU(5)*, *Nucl. Phys. B* **193** (1981) 150 [INSPIRE].
- [2] L.E. Ibáñez and G.G. Ross, *Low-Energy Predictions in Supersymmetric Grand Unified Theories*, *Phys. Lett. B* **105** (1981) 439 [INSPIRE].
- [3] S. Dimopoulos, S. Raby and F. Wilczek, *Supersymmetry and the Scale of Unification*, *Phys. Rev. D* **24** (1981) 1681 [INSPIRE].
- [4] L.E. Ibáñez and G.G. Ross, *SU(2)-L \times U(1) Symmetry Breaking as a Radiative Effect of Supersymmetry Breaking in Guts*, *Phys. Lett. B* **110** (1982) 215 [INSPIRE].
- [5] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, *Aspects of Grand Unified Models with Softly Broken Supersymmetry*, *Prog. Theor. Phys.* **68** (1982) 927 [Erratum *ibid.* **70** (1983) 330] [INSPIRE].
- [6] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, *Renormalization of Supersymmetry Breaking Parameters Revisited*, *Prog. Theor. Phys.* **71** (1984) 413 [INSPIRE].
- [7] L. Álvarez-Gaumé, J. Polchinski and M.B. Wise, *Minimal Low-Energy Supergravity*, *Nucl. Phys. B* **221** (1983) 495 [INSPIRE].
- [8] J.R. Ellis, J. Hagelin, D.V. Nanopoulos and K. Tamvakis, *Weak Symmetry Breaking by Radiative Corrections in Broken Supergravity*, *Phys. Lett. B* **125** (1983) 275 [INSPIRE].
- [9] H. Baer et al., *Radiative natural supersymmetry: Reconciling electroweak fine-tuning and the Higgs boson mass*, *Phys. Rev. D* **87** (2013) 115028 [arXiv:1212.2655] [INSPIRE].
- [10] S. Cassel, D.M. Ghilencea and G.G. Ross, *Fine tuning as an indication of physics beyond the MSSM*, *Nucl. Phys. B* **825** (2010) 203 [arXiv:0903.1115] [INSPIRE].
- [11] R. Dermisek and J.F. Gunion, *The NMSSM Solution to the Fine-Tuning Problem, Precision Electroweak Constraints and the Largest LEP Higgs Event Excess*, *Phys. Rev. D* **76** (2007) 095006 [arXiv:0705.4387] [INSPIRE].
- [12] G.G. Ross and K. Schmidt-Hoberg, *The Fine-Tuning of the Generalised NMSSM*, *Nucl. Phys. B* **862** (2012) 710 [arXiv:1108.1284] [INSPIRE].
- [13] A. Delgado, C. Kolda, J.P. Olson and A. de la Puente, *Solving the Little Hierarchy Problem with a Singlet and Explicit μ Terms*, *Phys. Rev. Lett.* **105** (2010) 091802 [arXiv:1005.1282] [INSPIRE].
- [14] A. Delgado, C. Kolda, J.P. Olson and A. de la Puente, *Gauge-mediated embedding of the singlet extension of the minimal supersymmetric standard model*, *Phys. Rev. D* **82** (2010) 035006 [arXiv:1005.4901] [INSPIRE].
- [15] G.G. Ross, K. Schmidt-Hoberg and F. Staub, *The Generalised NMSSM at One Loop: Fine Tuning and Phenomenology*, *JHEP* **08** (2012) 074 [arXiv:1205.1509] [INSPIRE].
- [16] K. Choi, K.S. Jeong, T. Kobayashi and K.-i. Okumura, *Little SUSY hierarchy in mixed modulus-anomaly mediation*, *Phys. Lett. B* **633** (2006) 355 [hep-ph/0508029] [INSPIRE].

- [17] K. Choi, K.S. Jeong, T. Kobayashi and K.-i. Okumura, *TeV Scale Mirage Mediation and Natural Little SUSY Hierarchy*, *Phys. Rev. D* **75** (2007) 095012 [[hep-ph/0612258](#)] [[INSPIRE](#)].
- [18] H. Abe, T. Kobayashi and Y. Omura, *Relaxed fine-tuning in models with non-universal gaugino masses*, *Phys. Rev. D* **76** (2007) 015002 [[hep-ph/0703044](#)] [[INSPIRE](#)].
- [19] O. Lebedev, H.P. Nilles and M. Ratz, *A note on fine-tuning in mirage mediation*, [hep-ph/0511320](#) [[INSPIRE](#)].
- [20] D. Horton and G. Ross, *Naturalness and Focus Points with Non-Universal Gaugino Masses*, *Nucl. Phys. B* **830** (2010) 221 [[arXiv:0908.0857](#)] [[INSPIRE](#)].
- [21] M. Asano and T. Higaki, *Natural supersymmetric spectrum in mirage mediation*, *Phys. Rev. D* **86** (2012) 035020 [[arXiv:1204.0508](#)] [[INSPIRE](#)].
- [22] S. Antusch, L. Calibbi, V. Maurer, M. Monaco and M. Spinrath, *Naturalness of the Non-Universal MSSM in the Light of the Recent Higgs Results*, *JHEP* **01** (2013) 187 [[arXiv:1207.7236](#)] [[INSPIRE](#)].
- [23] H. Abe, J. Kawamura and H. Otsuka, *The Higgs boson mass in a natural MSSM with nonuniversal gaugino masses at the GUT scale*, *Prog. Theor. Exp. Phys.* **2013** (2013) 013B02 [[arXiv:1208.5328](#)] [[INSPIRE](#)].
- [24] M. Badziak, S. Krippendorff, H.P. Nilles and M.W. Winkler, *The heterotic MiniLandscape and the 126 GeV Higgs boson*, *JHEP* **03** (2013) 094 [[arXiv:1212.0854](#)] [[INSPIRE](#)].
- [25] I. Gogoladze, F. Nasir and Q. Shafi, *Non-Universal Gaugino Masses and Natural Supersymmetry*, *Int. J. Mod. Phys. A* **28** (2013) 1350046 [[arXiv:1212.2593](#)] [[INSPIRE](#)].
- [26] T.T. Yanagida and N. Yokozaki, *Focus Point in Gaugino Mediation — Reconsideration of the Fine-tuning Problem*, *Phys. Lett. B* **722** (2013) 355 [[arXiv:1301.1137](#)] [[INSPIRE](#)].
- [27] D. Ghilencea and G. Ross, *The fine-tuning cost of the likelihood in SUSY models*, *Nucl. Phys. B* **868** (2013) 65 [[arXiv:1208.0837](#)] [[INSPIRE](#)].
- [28] D. Ghilencea, *Fixing the EW scale in supersymmetric models after the Higgs discovery*, *Nucl. Phys. B* **876** (2013) 16 [[arXiv:1302.5262](#)] [[INSPIRE](#)].
- [29] D. Ghilencea, *A new approach to Naturalness in SUSY models*, *PoS Corfu2012* (2013) 034 [[arXiv:1304.1193](#)] [[INSPIRE](#)].
- [30] M. Cabrera, J. Casas and R. Ruiz de Austri, *Bayesian approach and Naturalness in MSSM analyses for the LHC*, *JHEP* **03** (2009) 075 [[arXiv:0812.0536](#)] [[INSPIRE](#)].
- [31] J.R. Ellis, K. Enqvist, D.V. Nanopoulos and F. Zwirner, *Observables in Low-Energy Superstring Models*, *Mod. Phys. Lett. A* **1** (1986) 57 [[INSPIRE](#)].
- [32] R. Barbieri and G. Giudice, *Upper Bounds on Supersymmetric Particle Masses*, *Nucl. Phys. B* **306** (1988) 63 [[INSPIRE](#)].
- [33] P. Ciafaloni and A. Strumia, *Naturalness upper bounds on gauge mediated soft terms*, *Nucl. Phys. B* **494** (1997) 41 [[hep-ph/9611204](#)] [[INSPIRE](#)].
- [34] ATLAS collaboration, *Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum and 20.3fb^{-1} of $\sqrt{s} = 8\text{TeV}$ proton-proton collision data*, *ATLAS-CONF-2013-047* (2013).
- [35] CMS collaboration, *Search for supersymmetry in pp collisions at $\sqrt{s} = 8\text{TeV}$ in events with three leptons and at least one b -tagged jet*, *CMS-PAS-SUS-13-008*.

- [36] PLANCK collaboration, P. Ade et al., *Planck 2013 results. XVI. Cosmological parameters*, [arXiv:1303.5076](#) [INSPIRE].
- [37] XENON100 collaboration, E. Aprile et al., *Dark Matter Results from 225 Live Days of XENON100 Data*, *Phys. Rev. Lett.* **109** (2012) 181301 [[arXiv:1207.5988](#)] [INSPIRE].
- [38] CMS collaboration, *Combination of standard model Higgs boson searches and measurements of the properties of the new boson with a mass near 125 GeV*, CMS-PAS-HIG-13-005.
- [39] ATLAS collaboration, *Combined measurements of the mass and signal strength of the Higgs-like boson with the ATLAS detector using up to 25fb^{-1} of proton-proton collision data*, ATLAS-CONF-2013-014 (2013).
- [40] W. Porod, *SPheno, a program for calculating supersymmetric spectra, SUSY particle decays and SUSY particle production at e^+e^- colliders*, *Comput. Phys. Commun.* **153** (2003) 275 [[hep-ph/0301101](#)] [INSPIRE].
- [41] W. Porod and F. Staub, *SPheno 3.1: Extensions including flavour, CP-phases and models beyond the MSSM*, *Comput. Phys. Commun.* **183** (2012) 2458 [[arXiv:1104.1573](#)] [INSPIRE].
- [42] F. Staub, *SARAH*, [arXiv:0806.0538](#) [INSPIRE].
- [43] F. Staub, *From Superpotential to Model Files for FeynArts and CalcHep/CompHEP*, *Comput. Phys. Commun.* **181** (2010) 1077 [[arXiv:0909.2863](#)] [INSPIRE].
- [44] F. Staub, *Automatic Calculation of supersymmetric Renormalization Group Equations and Self Energies*, *Comput. Phys. Commun.* **182** (2011) 808 [[arXiv:1002.0840](#)] [INSPIRE].
- [45] F. Staub, *SARAH 3.2: Dirac Gauginos, UFO output and more*, *Computer Physics Communications* **184** (2013) pp. 1792–1809 [[arXiv:1207.0906](#)] [INSPIRE].
- [46] D.M. Pierce, J.A. Bagger, K.T. Matchev and R.-j. Zhang, *Precision corrections in the minimal supersymmetric standard model*, *Nucl. Phys. B* **491** (1997) 3 [[hep-ph/9606211](#)] [INSPIRE].
- [47] F. Staub, W. Porod and B. Herrmann, *The electroweak sector of the NMSSM at the one-loop level*, *JHEP* **10** (2010) 040 [[arXiv:1007.4049](#)] [INSPIRE].
- [48] A. Dedes, G. Degrossi and P. Slavich, *On the two loop Yukawa corrections to the MSSM Higgs boson masses at large $\tan\beta$* , *Nucl. Phys. B* **672** (2003) 144 [[hep-ph/0305127](#)] [INSPIRE].
- [49] A. Dedes and P. Slavich, *Two loop corrections to radiative electroweak symmetry breaking in the MSSM*, *Nucl. Phys. B* **657** (2003) 333 [[hep-ph/0212132](#)] [INSPIRE].
- [50] A. Brignole, G. Degrossi, P. Slavich and F. Zwirner, *On the two loop sbottom corrections to the neutral Higgs boson masses in the MSSM*, *Nucl. Phys. B* **643** (2002) 79 [[hep-ph/0206101](#)] [INSPIRE].
- [51] A. Brignole, G. Degrossi, P. Slavich and F. Zwirner, *On the $O(\alpha_t^2)$ two loop corrections to the neutral Higgs boson masses in the MSSM*, *Nucl. Phys. B* **631** (2002) 195 [[hep-ph/0112177](#)] [INSPIRE].
- [52] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, *MicrOMEGAs 2.0: A program to calculate the relic density of dark matter in a generic model*, *Comput. Phys. Commun.* **176** (2007) 367 [[hep-ph/0607059](#)] [INSPIRE].
- [53] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, *MicrOMEGAs 2.0.7: A program to calculate the relic density of dark matter in a generic model*, *Comput. Phys. Commun.* **177** (2007) 894 [INSPIRE].

- [54] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, *MicrOMEGAs: A tool for dark matter studies*, *Nuovo Cim. C* **033N2** (2010) 111 [[arXiv:1005.4133](#)] [[INSPIRE](#)].
- [55] L.E. Ibáñez and C. Lopez, *$N=1$ Supergravity, the Weak Scale and the Low-Energy Particle Spectrum*, *Nucl. Phys. B* **233** (1984) 511 [[INSPIRE](#)].
- [56] Y. Mambrini, G. Moultaka and M. Rausch de Traubenberg, *General one loop renormalization group evolutions and electroweak symmetry breaking in the $(M+1)$ SSM*, *Nucl. Phys. B* **609** (2001) 83 [[hep-ph/0101237](#)] [[INSPIRE](#)].
- [57] S.P. Martin, *Non-universal gaugino masses from non-singlet F -terms in non-minimal unified models*, *Phys. Rev. D* **79** (2009) 095019 [[arXiv:0903.3568](#)] [[INSPIRE](#)].
- [58] A. Brignole, L.E. Ibáñez and C. Muñoz, *Towards a theory of soft terms for the supersymmetric Standard Model*, *Nucl. Phys. B* **422** (1994) 125 [Erratum *ibid.* **B 436** (1995) 747-748] [[hep-ph/9308271](#)] [[INSPIRE](#)].
- [59] K. Choi and H.P. Nilles, *The gaugino code*, *JHEP* **04** (2007) 006 [[hep-ph/0702146](#)] [[INSPIRE](#)].
- [60] W. Cho, Y. Kim, K. Lee, C. Park and Y. Shimizu, *LHC Signature of Mirage Mediation*, *JHEP* **04** (2007) 054 [[hep-ph/0703163](#)] [[INSPIRE](#)].
- [61] K. Choi, K.S. Jeong and K.-i. Okumura, *Phenomenology of mixed modulus-anomaly mediation in fluxed string compactifications and brane models*, *JHEP* **09** (2005) 039 [[hep-ph/0504037](#)] [[INSPIRE](#)].
- [62] L. Roszkowski, R. Ruiz de Austri, R. Trotta, Y.-L.S. Tsai and T.A. Varley, *Global fits of the Non-Universal Higgs Model*, *Phys. Rev. D* **83** (2011) 015014 [[arXiv:0903.1279](#)] [[INSPIRE](#)].
- [63] S. Akula, P. Nath and G. Peim, *Implications of the Higgs Boson Discovery for $mSUGRA$* , *Phys. Lett. B* **717** (2012) 188 [[arXiv:1207.1839](#)] [[INSPIRE](#)].
- [64] H.M. Lee et al., *A unique Z_4^R symmetry for the MSSM*, *Phys. Lett. B* **694** (2011) 491 [[arXiv:1009.0905](#)] [[INSPIRE](#)].
- [65] H.M. Lee et al., *Discrete R symmetries for the MSSM and its singlet extensions*, *Nucl. Phys. B* **850** (2011) 1 [[arXiv:1102.3595](#)] [[INSPIRE](#)].
- [66] U. Ellwanger and C. Hugonie, *The upper bound on the lightest Higgs mass in the NMSSM revisited*, *Mod. Phys. Lett. A* **22** (2007) 1581 [[hep-ph/0612133](#)] [[INSPIRE](#)].
- [67] S. Raby, M. Ratz and K. Schmidt-Hoberg, *Precision gauge unification in the MSSM*, *Phys. Lett. B* **687** (2010) 342 [[arXiv:0911.4249](#)] [[INSPIRE](#)].
- [68] K. Schmidt-Hoberg and F. Staub, *Enhanced $h \rightarrow \gamma\gamma$ rate in MSSM singlet extensions*, *JHEP* **10** (2012) 195 [[arXiv:1208.1683](#)] [[INSPIRE](#)].
- [69] K. Schmidt-Hoberg, F. Staub and M.W. Winkler, *Enhanced diphoton rates at Fermi and the LHC*, *JHEP* **01** (2013) 124 [[arXiv:1211.2835](#)] [[INSPIRE](#)].