



# Measurement of $B_c^+$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV

The LHCb collaboration<sup>†</sup>

## Abstract

Production of  $B_c^+$  mesons in proton-proton collisions at a center-of-mass energy of 8 TeV is studied with data corresponding to an integrated luminosity of  $2.0 \text{ fb}^{-1}$  recorded by the LHCb experiment. The ratio of production cross-sections times branching fractions between the  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$  decays is measured as a function of transverse momentum and rapidity in the regions  $0 < p_T < 20 \text{ GeV}/c$  and  $2.0 < y < 4.5$ . The ratio in this kinematic range is measured to be  $(0.683 \pm 0.018 \pm 0.009)\%$ , where the first uncertainty is statistical and the second systematic.

Published in Phys. Rev. Lett.

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

---

<sup>†</sup>Authors are listed at the end of this Letter.



In the Standard Model, the  $B_c$  mesons are the only states formed by two heavy quarks of different flavor, the  $\bar{b}$  and the  $c$  quarks. The production of  $B_c$  mesons in hadron collisions implies the simultaneous production of  $b\bar{b}$  and  $c\bar{c}$  pairs, therefore it is rarer than that of other  $b$  mesons. The production of  $b\bar{b}$  and  $c\bar{c}$  quarkonium states in hadron collisions has been studied for two decades, however, significant puzzles remain [1]. The relative role of competing production mechanisms [2–5] is poorly understood and theory is unable to predict all experimentally observed features [6–11]. The study of  $B_c$  production offers a promising way of shedding light over these discrepancies and gaining an insight on the underlying physics. In proton-proton ( $pp$ ) collisions at the Large Hadron Collider (LHC),  $B_c$  mesons are expected to be mainly produced through the gluon-gluon fusion process  $gg \rightarrow B_c + b + \bar{c}$ . The production cross-sections of the  $B_c$  mesons have been calculated in the fragmentation approach [12, 13] and in the complete order- $\alpha_s^4$  approach [14–21], where  $\alpha_s$  is the strong-interaction coupling. In the latter approach, the total production cross-section of the  $B_c$  ground state,  $B_c^+$ , at a center-of-mass energy of 8 TeV, integrated over the whole phase space and including contributions from intermediate excited states, is predicted to be about 0.2% [22, 23] of the inclusive  $b\bar{b}$  cross-section [24].

Previously, only the average ratios of  $B_c^+$  to  $B^+$  or  $B_s^0$  cross-sections in specific kinematic regions had been measured [25–27], and double-differential cross-sections have not yet been measured. The production cross-sections of  $b$ -hadrons show different transverse momentum dependencies [28–31]. A precise measurement of  $B_c^+$  production as a function of transverse momentum and rapidity will provide useful information on the largely unknown production mechanism of the  $B_c^+$  meson and other bound states of heavy quarks, and is also important to guide  $B_c^+$  studies at the LHC.

In this Letter we report on the first measurement of the ratio of double differential inclusive production cross-sections multiplied by branching fractions,

$$R(p_T, y) \equiv \frac{d\sigma_{B_c^+}(p_T, y) \mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)}{d\sigma_{B^+}(p_T, y) \mathcal{B}(B^+ \rightarrow J/\psi K^+)}, \quad (1)$$

where transverse momentum  $p_T$  and rapidity  $y$  refer to the  $b$  meson. The cross-section includes contributions from excited states. We use a sample of  $pp$  collision data at 8 TeV, corresponding to an integrated luminosity of  $2.0 \text{ fb}^{-1}$  recorded by the LHCb experiment. The  $B_c^+$  and  $B^+$  mesons are reconstructed in the exclusive decays  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$  respectively, with  $J/\psi \rightarrow \mu^+ \mu^-$ . The inclusion of charge conjugate modes is implied throughout this Letter.

The LHCb detector [32] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream of the magnet. The combined tracking system provides a momentum measurement with a relative uncertainty that varies from 0.4% at low momentum,  $p$ , to 0.6% at 100 GeV/ $c$ . The minimum distance of a track to a primary vertex, the impact parameter (IP), is measured

with a resolution of  $(15 + 29/p_T)$   $\mu\text{m}$ , where  $p_T$  is in  $\text{GeV}/c$ . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors. Photon, electron and hadron candidates are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers.

The trigger consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, in which all charged particles with  $p_T > 300 \text{ MeV}/c$  are reconstructed [33]. Events are first required to pass the hardware trigger, which requires one or two muons with high  $p_T$ . In the subsequent software trigger, the event is required to have one muon with high  $p_T$  and large IP with respect to all primary  $pp$  interaction vertices (PVs), or a pair of oppositely charged muons with an invariant mass consistent with the known  $J/\psi$  meson mass [34]. Finally, the tracks of two or more of the final state particles are required to form a vertex that is significantly displaced from the PVs. A multivariate algorithm [35] is also used to identify secondary vertices consistent with the decay of a  $b$  meson.

The  $b$ -meson candidate selection is performed in two steps, a preselection and a final selection on the output of a multivariate classifier based on a boosted decision tree algorithm (BDT) [36, 37]. Simulated  $B_c^+$  and  $B^+$  decays are used to optimize the  $b$ -meson candidate selection. Production of  $B^+$  mesons is simulated using PYTHIA 6.4 [38] with an LHCb specific configuration [39]. The generator BCVEGPY [40] is used to simulate  $B_c^+$  meson production. Decays of  $B_c^+$ ,  $B^+$  and  $J/\psi$  mesons are described by EVTGEN [41] and photon radiation is simulated using the PHOTOS package [42]. The decay products are traced through the detector by the GEANT4 package [43, 44]. Following Ref. [45], the  $B_c^+$  meson lifetime is set to  $\tau_{B_c^+} = 0.509 \text{ ps}$ . The selection requirements are the same for  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$  candidates.

In the preselection,  $J/\psi$  candidates are formed from pairs of oppositely charged particles with  $p_T$  larger than  $0.55 \text{ GeV}/c$ , with a good quality of the track fit and identified as muons. The two muons are required to originate from a common vertex. The  $J/\psi$  candidates with invariant mass between  $3.04 \text{ GeV}/c^2$  and  $3.14 \text{ GeV}/c^2$  are combined with a charged particle that has  $p_T > 1.0 \text{ GeV}/c$ , a good quality of the track fit and is separated from any PV. The pion mass hypothesis is assigned to the track for the selection of the  $B_c^+$  candidate and the kaon hypothesis for that of the  $B^+$  candidate. The  $J/\psi$  candidate and the hadron ( $\pi$  or  $K$ ) are required to originate from a common vertex. To improve the  $b$ -meson mass resolution, the mass of the muon pair is constrained to the known  $J/\psi$  meson mass [34] in this vertex fit. The  $b$ -meson candidates are required to have a decay time larger than  $0.2 \text{ ps}$ , and to point toward the primary vertex.

In the final selection, the BDT is trained using a simulated  $B_c^+$  signal sample and background events populating the data mass sideband  $6376 < M_{J/\psi \pi^+} < 6600 \text{ MeV}/c^2$ . The following variables are used as input to the BDT:  $\chi_{\text{IP}}^2$  of all particles;  $p_T$  of muons,  $J/\psi$  and  $\pi^+$ ; and the  $b$ -meson decay length, decay time, and the vertex fit  $\chi^2$  of a fit to the decay tree [46]. The quantity  $\chi_{\text{IP}}^2$  is defined as the difference in  $\chi^2$  of a given primary vertex reconstructed with and without the considered particle. The selection value on the

BDT output is chosen to maximize the signal significance  $N_S/\sqrt{N_S + N_B}$ , where  $N_S$  and  $N_B$  are the expected numbers of signal and background events, respectively. The same BDT requirements are used for the  $B^+$  meson.

The  $B_c^+$  and  $B^+$  candidates are subdivided into 10 bins of  $p_T$  and 3 bins of  $y$ . Bin sizes are chosen to contain approximately the same number of signal candidates, except for the highest  $p_T$  bin. The differential production ratio  $R$  is measured as

$$R(p_T, y) = \frac{N_{B_c^+}(p_T, y) \epsilon_{B^+}(p_T, y)}{N_{B^+}(p_T, y) \epsilon_{B_c^+}(p_T, y)}, \quad (2)$$

where  $N_B(p_T, y)$  is the number of reconstructed signal decays and  $\epsilon_B(p_T, y)$  is the total efficiency in a given  $(p_T, y)$  bin, including geometrical acceptance, reconstruction, selection and trigger effects.

In each  $p_T$  and  $y$  bin, the number of signal decays is determined by performing an extended maximum likelihood fit to the unbinned invariant mass distribution of  $B_c^+$  candidates reconstructed in  $6150 < M_{J/\psi \pi^+} < 6550 \text{ MeV}/c^2$  and  $B^+$  candidates in  $5150 < M_{J/\psi K^+} < 5550 \text{ MeV}/c^2$ . For both  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$  decays, the fit includes components for signal, combinatorial background, and Cabibbo-suppressed backgrounds  $B_c^+ \rightarrow J/\psi K^+$  and  $B^+ \rightarrow J/\psi \pi^+$ . Other sources of backgrounds, such as  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$ , are negligible. The  $B_c^+ \rightarrow J/\psi \pi^+$  signal is described by a double-sided Crystal Ball (DSCB) function, which is an empirical function with a Gaussian core and power-law tails on both sides. The  $B^+ \rightarrow J/\psi K^+$  signal is described by the sum of two DSCB functions, to account for different mass resolutions in different kinematic regions. The tail parameters are determined from simulation. The combinatorial background is described by an exponential function. The shapes of the Cabibbo-suppressed backgrounds are determined from simulation. The ratios of the yield of the Cabibbo-suppressed background to that of the signal are fixed to the central value of  $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+) = (6.9 \pm 2.0)\%$  for  $B_c^+$  candidates [47], and  $\mathcal{B}(B^+ \rightarrow J/\psi \pi^+)/\mathcal{B}(B^+ \rightarrow J/\psi K^+) = (3.83 \pm 0.13)\%$  for  $B^+$  candidates [48], respectively.

As an example, Fig. 1 shows the  $B_c^+$  and  $B^+$  mass distributions together with the fit results for the bin  $2.0 < p_T < 3.0 \text{ GeV}/c$  and  $2.0 < y < 2.9$ . The mass resolution is approximately  $11 \text{ MeV}/c^2$  for  $B_c^+$  signals and  $8.7 \text{ MeV}/c^2$  for  $B^+$  signals. Summing over all bins, a total signal yield of  $3.1 \times 10^3$   $B_c^+$  candidates and  $7.1 \times 10^5$   $B^+$  candidates is obtained. In each  $(p_T, y)$  bin the total efficiency is determined from simulation and ranges from 2.4% to 23.2% for  $B_c^+$  candidates and from 3.6% to 33.5% for  $B^+$  candidates.

The systematic uncertainties associated with the signal shape in each bin (0.1% – 2.6%) are estimated by comparing the ratios between input signal yields and fit results in simulation. The uncertainties from the combinatorial background shape (0.1% – 4.4%) are determined by varying the fit function. The input value for the ratio of branching fractions  $\mathcal{B}(B_c^+ \rightarrow J/\psi K^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$  is varied within its uncertainty and the resulting difference (0.1% – 0.9%) is taken as systematic uncertainty. The effect of  $\mathcal{B}(B^+ \rightarrow J/\psi \pi^+)/\mathcal{B}(B^+ \rightarrow J/\psi K^+)$  is found to be negligible. The systematic uncertainty associated with the relative trigger efficiency is estimated to be 1%. Other effects, such as




Figure 1: Invariant mass distribution of (*left*)  $B_c^+ \rightarrow J/\psi \pi^+$  and (*right*)  $B^+ \rightarrow J/\psi K^+$  candidates with  $2.0 < p_T < 3.0 \text{ GeV}/c$  and  $2.0 < y < 2.9$ . The results of the fit described in the text are superimposed.




Figure 2: Distributions of (*left*)  $p_T$  and (*right*)  $y$  of the  $B_c^+$  signal after event selection. The points with error bars are background-subtracted data, and the solid histogram is the simulation based on the complete order- $\alpha_s^4$  calculation, implemented in the  $B_c^+$  generator BCVEGPY [40]. The uncertainties are statistical.

the  $(p_T, y)$  binning scheme, the shapes of the Cabibbo-suppressed backgrounds, the  $B_c^+$  lifetime uncertainty and the uncertainty of tracking efficiency, are negligible.

Figure 2 shows that simulation provides a good description of  $p_T$  and  $y$  distributions of  $B_c^+$  mesons in data. The values of  $R(p_T, y)$  in the range  $0 < p_T < 20 \text{ GeV}/c$  and  $2.0 < y < 4.5$  are shown in Fig. 3 and Ref. [49]. Figure 4 shows the ratio  $R(p_T)$  integrated over  $y$  in the region  $2.0 < y < 4.5$  and  $R(y)$  integrated over  $p_T$  in the region  $0 < p_T < 20 \text{ GeV}/c$ . The ratios are found to vary as a function of  $p_T$  and  $y$ . The results are compared with the theoretical predictions in Ref. [49].

The resulting integrated value of  $R$  in the region  $0 < p_T < 20 \text{ GeV}/c$  and  $2.0 < y < 4.5$  is measured to be

$$R = (0.683 \pm 0.018 \pm 0.009)\%,$$




Figure 3: Ratio  $R(p_T, y)$  as a function of  $p_T$  in the regions (*top*)  $2.0 < y < 2.9$ , (*middle*)  $2.9 < y < 3.3$ , and (*bottom*)  $3.3 < y < 4.5$ . The error bars on the data show the statistical and systematic uncertainties added in quadrature.




Figure 4: Ratio (*left*)  $R(p_T)$  as a function of  $p_T$  integrated over  $y$  in the region  $2.0 < y < 4.5$  and (*right*)  $R(y)$  as a function of  $y$  integrated over  $p_T$  in the region  $0 < p_T < 20$  GeV/c. The error bars on the data show the statistical and systematic uncertainties added in quadrature.

where the first uncertainty is statistical and the second systematic. To enable comparison with the previous LHCb measurement [26],  $R$  and its total uncertainty are also reported in the range  $4 < p_T < 20$  GeV/c and  $2.5 < \eta < 4.5$  as  $(0.698 \pm 0.023)\%$ . The previous LHCb measurement of  $R$  at 7 TeV of Ref. [26] is updated using the recent measurement of the  $B_c^+$  lifetime [45] to be  $(0.61 \pm 0.12)\%$ .

In summary, we present the first measurement of the  $B_c^+$  double differential production cross-section ratio with respect to that of the  $B^+$  meson. The measurement is performed in three bins of rapidity and ten bins of  $p_T$  in  $pp$  collisions at  $\sqrt{s} = 8$  TeV on a data

sample collected with the LHCb detector. The relative production rates of  $B_c^+$  and  $B^+$  mesons are found to depend on their transverse momentum and rapidity. The measured transverse momentum and rapidity distributions of the  $B_c^+$  meson are well described by the complete order- $\alpha_s^4$  calculation. However, the theoretical predictions on the  $B_c^+$  and  $B^+$  production cross-sections suffer from big uncertainties [22, 50], and the prediction of the branching fraction of the  $B_c^+ \rightarrow J/\psi \pi^+$  decay has a big spread (see for example Ref. [51]), more work on the theoretical side is required to have concluding remarks on the  $B_c^+$  absolute production rate. These results will provide useful information on the  $B_c^+$  production mechanism, and help understand the quarkonium production, therefore deepen our understanding of QCD.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are indebted to the communities behind the multiple open source software packages on which we depend. We are also thankful for the computing resources and the access to software R&D tools provided by Yandex LLC (Russia). Individual groups or members have received support from EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR (Russia), XuntaGal and GENCAT (Spain), Royal Society and Royal Commission for the Exhibition of 1851 (United Kingdom). We would like to thank Matteo Cacciari, Chao-Hsi Chang, Xing-Gang Wu, and Rui-Lin Zhu for useful discussions.



## Supplementary material

Table 1 shows the ratio  $R(p_T, y)$  in each  $(p_T, y)$  bin.

Table 1:  $R(p_T, y)$  in units of  $10^{-2}$  as a function of  $p_T$  and  $y$ . The first uncertainty is statistical and the second systematic.

$p_T$ (GeV/c)	$2.0 < y < 2.9$	$2.9 < y < 3.3$	$3.3 < y < 4.5$	$2.0 < y < 4.5$
$0 < p_T < 2$	$0.67 \pm 0.10 \pm 0.01$	$0.73 \pm 0.10 \pm 0.01$	$0.35 \pm 0.06 \pm 0.01$	$0.54 \pm 0.05 \pm 0.01$
$2 < p_T < 3$	$0.70 \pm 0.09 \pm 0.02$	$0.72 \pm 0.09 \pm 0.02$	$0.50 \pm 0.06 \pm 0.01$	$0.62 \pm 0.05 \pm 0.01$
$3 < p_T < 4$	$0.62 \pm 0.08 \pm 0.01$	$0.58 \pm 0.08 \pm 0.01$	$0.57 \pm 0.07 \pm 0.02$	$0.59 \pm 0.05 \pm 0.01$
$4 < p_T < 5$	$0.83 \pm 0.08 \pm 0.02$	$0.60 \pm 0.07 \pm 0.01$	$0.81 \pm 0.08 \pm 0.02$	$0.79 \pm 0.05 \pm 0.01$
$5 < p_T < 6$	$0.90 \pm 0.09 \pm 0.02$	$0.78 \pm 0.09 \pm 0.01$	$0.76 \pm 0.09 \pm 0.02$	$0.83 \pm 0.06 \pm 0.01$
$6 < p_T < 7$	$0.84 \pm 0.09 \pm 0.01$	$0.99 \pm 0.11 \pm 0.02$	$0.64 \pm 0.08 \pm 0.01$	$0.79 \pm 0.06 \pm 0.01$
$7 < p_T < 8$	$0.95 \pm 0.10 \pm 0.01$	$0.74 \pm 0.11 \pm 0.01$	$0.65 \pm 0.09 \pm 0.01$	$0.82 \pm 0.06 \pm 0.01$
$8 < p_T < 10$	$0.80 \pm 0.08 \pm 0.01$	$0.57 \pm 0.08 \pm 0.01$	$0.80 \pm 0.09 \pm 0.02$	$0.77 \pm 0.05 \pm 0.01$
$10 < p_T < 14$	$0.70 \pm 0.06 \pm 0.01$	$0.75 \pm 0.09 \pm 0.01$	$0.60 \pm 0.08 \pm 0.01$	$0.68 \pm 0.05 \pm 0.01$
$14 < p_T < 20$	$0.74 \pm 0.09 \pm 0.01$	$0.68 \pm 0.15 \pm 0.03$	$0.55 \pm 0.13 \pm 0.02$	$0.68 \pm 0.07 \pm 0.01$
$0 < p_T < 20$	$0.76 \pm 0.03 \pm 0.01$	$0.70 \pm 0.03 \pm 0.01$	$0.58 \pm 0.03 \pm 0.01$	$0.68 \pm 0.02 \pm 0.01$

The results are compared with the theoretical predictions in Fig. 5 and Fig. 6. For  $B_c^+$  meson the predictions following the  $\alpha_s^4$  approach [40] are shown. We use the CTEQ6LL [52] parton distribution functions, and the leading order running  $\alpha_s$ , the characteristic energy scale  $Q^2 = p_T^2 + m_{B_c^+}^2$ , and the masses of the  $b$  and  $c$  quarks are set to  $m_b = 4.95 \text{ GeV}/c^2$  and  $m_c = 1.326 \text{ GeV}/c^2$ . The normalization of the theoretical predictions uses  $0.47 \mu\text{b}$  as the  $B_c^+$  production cross-section in the whole phase space and  $0.33\%$  for  $\mathcal{B}(B_c^+ \rightarrow J/\psi \pi^+)$  [51], corrected for the latest measurement of the  $B_c^+$  lifetime. The theoretical prediction on the  $B^+$  cross-section is based on the fixed order + next-to-leading log (FONLL) framework [50]. The uncertainties on the theory curves are the uncertainties of the FONLL calculation, including the uncertainties of the  $b$  quark mass, the renormalisation and factorisation scales, and CTEQ6.6 [53] functions. The FONLL predictions are scaled according to the measured branching fraction value  $\mathcal{B}(B^+ \rightarrow J/\psi K^+) = 0.106\%$  [34] and the  $B^+$  production cross-section  $38.9 \mu\text{b}$  measured at  $\sqrt{s} = 7 \text{ TeV}$  [54] increased by  $20\%$  due to higher collision energy [55].




Figure 5: Ratio  $R(p_T, y)$  as a function of  $p_T$  in the regions  $2.0 < y < 2.9$  (top left),  $2.9 < y < 3.3$  (top right), and  $3.3 < y < 4.5$  (bottom left), with theoretical predictions following the  $\alpha_s^4$  approach [40] overlaid.




Figure 6: Ratio  $R(p_T)$  as a function of  $p_T$  integrated over  $y$  in the region  $2.0 < y < 4.5$  (left) and  $R(y)$  as a function of  $y$  integrated over  $p_T$  in the region  $0 < p_T < 20$  GeV/c (right) are compared to the theoretical predictions following the  $\alpha_s^4$  approach [40].

## References

- [1] N. Brambilla *et al.*, *Heavy quarkonium: progress, puzzles, and opportunities*, Eur. Phys. J. **C71** (2011) 1534, [arXiv:1010.5827](#).
- [2] C.-H. Chang, *Hadronic production of  $J/\psi$  associated with a gluon*, Nucl. Phys. **B172** (1980) 425.
- [3] R. Baier and R. Ruckl, *Hadronic collisions: A quarkonium factory*, Z. Phys. **C19** (1983) 251.
- [4] G. T. Bodwin, E. Braaten, and G. P. Lepage, *Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium*, Phys. Rev. **D51** (1995) 1125, [arXiv:hep-ph/9407339](#), erratum *ibid.* **D55** (1997) 5853.
- [5] P. L. Cho and A. K. Leibovich, *Color octet quarkonia production*, Phys. Rev. **D53** (1996) 150, [arXiv:hep-ph/9505329](#); P. L. Cho and A. K. Leibovich, *Color octet quarkonia production. II*, Phys. Rev. **D53** (1996) 6203, [arXiv:hep-ph/9511315](#).
- [6] CDF collaboration, F. Abe *et al.*,  *$J/\psi$  and  $\psi(2S)$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. Lett. **79** (1997) 572; CDF collaboration, F. Abe *et al.*, *Production of  $J/\psi$  mesons from  $\chi_c$  meson decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. Lett. **79** (1997) 578.
- [7] CDF collaboration, A. Abulencia *et al.*, *Polarization of  $J/\psi$  and  $\psi(2S)$  mesons produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, Phys. Rev. Lett. **99** (2007) 132001, [arXiv:0704.0638](#).
- [8] ALICE collaboration, B. Abelev *et al.*,  *$J/\psi$  polarization in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, Phys. Rev. Lett. **108** (2012) 082001, [arXiv:1111.1630](#).
- [9] CMS collaboration, S. Chatrchyan *et al.*, *Measurement of the  $\Upsilon(1S)$ ,  $\Upsilon(2S)$  and  $\Upsilon(3S)$  polarizations in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, Phys. Rev. Lett. **110** (2013) 081802, [arXiv:1209.2922](#).
- [10] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $J/\psi$  polarization in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. **C73** (2013) 2631, [arXiv:1307.6379](#).
- [11] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $\psi(2S)$  polarisation in  $pp$  collisions at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. **C74** (2014) 2872, [arXiv:1403.1339](#).
- [12] E. Braaten, K. Cheung, and T. C. Yuan, *Perturbative QCD fragmentation functions for  $B_c$  and  $B_c^*$  production*, Phys. Rev. **D48** (1993) R5049 (R), [arXiv:hep-ph/9305206](#).
- [13] K. Cheung,  *$B_c$  meson production at the Tevatron revisited*, Phys. Lett. **B472** (2000) 408, [arXiv:hep-ph/9908405](#).

- [14] C.-H. Chang and Y.-Q. Chen, *Hadronic production of the  $B_c$  meson at TeV energies*, Phys. Rev. **D48** (1993) 4086.
- [15] C.-H. Chang, Y.-Q. Chen, G.-P. Han, and H.-T. Jiang, *On hadronic production of the  $B_c$  meson*, Phys. Lett. **B364** (1995) 78, arXiv:hep-ph/9408242.
- [16] C.-H. Chang, Y.-Q. Chen, and R. J. Oakes, *Comparative study of the hadronic production of  $B_c$  mesons*, Phys. Rev. **D54** (1996) 4344, arXiv:hep-ph/9602411.
- [17] C.-H. Chang, C.-F. Qiao, J.-X. Wang, and X.-G. Wu, *Color-octet contributions to P-wave  $B_c$  meson hadroproduction*, Phys. Rev. **D71** (2005) 074012, arXiv:hep-ph/0502155.
- [18] K. Kolodziej, A. Leike, and R. Ruckl, *Production of  $B_c$  mesons in hadronic collisions*, Phys. Lett. **B355** (1995) 337, arXiv:hep-ph/9505298.
- [19] A. V. Berezhnoy, A. K. Likhoded, and M. V. Shevlyagin, *Hadronic production of  $B_c$  mesons*, Phys. Atom. Nucl. **58** (1995) 672, arXiv:hep-ph/9408284.
- [20] A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, *Photonic production of S- and P-wave  $B_c$  states and doubly heavy baryons*, Z. Phys. **A356** (1996) 89.
- [21] S. P. Baranov, *Pair production of  $B_c^{(*)}$  mesons in pp and  $\gamma\gamma$  collisions*, Phys. Rev. **D55** (1997) 2756.
- [22] C.-H. Chang and X.-G. Wu, *Uncertainties in estimating  $B_c$  hadronic production and comparisons of the production at TEVATRON and LHC*, Eur. Phys. J. **C38** (2004) 267, arXiv:hep-ph/0309121.
- [23] Y.-N. Gao *et al.*, *Experimental prospects of the  $B_c$  studies of the LHCb experiment*, Chin. Phys. Lett. **27** (2010) 061302.
- [24] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $J/\psi$  production in pp collisions at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. **C71** (2011) 1645, arXiv:1103.0423.
- [25] CDF collaboration, F. Abe *et al.*, *Observation of the  $B_c$  meson in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. Lett. **81** (1998) 2432, arXiv:hep-ex/9805034; CDF collaboration, F. Abe *et al.*, *Observation of  $B_c$  mesons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, Phys. Rev. **D58** (1998) 112004, arXiv:hep-ex/9804014.
- [26] LHCb collaboration, R. Aaij *et al.*, *Measurements of  $B_c^+$  production and mass with the  $B_c^+ \rightarrow J/\psi\pi^+$  decay*, Phys. Rev. Lett. **109** (2012) 232001, arXiv:1209.5634.
- [27] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay  $B_c^+ \rightarrow B_s^0\pi^+$* , Phys. Rev. Lett. **111** (2013) 181801, arXiv:1308.4544.

- [28] CDF collaboration, T. Aaltonen *et al.*, *First measurement of the ratio of branching fractions  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- \bar{\nu}_\mu)/B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)$* , Phys. Rev. **D79** (2009) 032001, [arXiv:0810.3213](#).
- [29] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $b$  hadron production fractions in 7 TeV  $pp$  collisions*, Phys. Rev. **D85** (2012) 032008, [arXiv:1111.2357](#).
- [30] LHCb collaboration, R. Aaij *et al.*, *Study of the kinematic dependences of  $\Lambda_b^0$  production in  $pp$  collisions and a measurement of the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  branching fraction*, JHEP **08** (2014) 143, [arXiv:1405.6842](#).
- [31] LHCb collaboration, R. Aaij *et al.*, *Study of beauty hadron decays into pairs of charm hadrons*, Phys. Rev. Lett. **112** (2014) 202001, [arXiv:1403.3606](#).
- [32] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [33] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [34] Particle Data Group, K. A. Olive *et al.*, *The review of particle physics*, Chin. Phys. C **38** (2014) 090001.
- [35] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [36] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [37] R. E. Schapire and Y. Freund, *A decision-theoretic generalization of on-line learning and an application to boosting*, Jour. Comp. and Syst. Sc. **55** (1997) 119.
- [38] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [39] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE** (2010) 1155.
- [40] C.-H. Chang, J.-X. Wang, and X.-G. Wu, *BCVEGPY2.0: An upgraded version of the generator BCVEGPY with the addition of hadroproduction of the  $P$ -wave  $B_c$  states*, Comput. Phys. Commun. **174** (2006) 241, [arXiv:hep-ph/0504017](#).
- [41] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [42] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in  $Z$  and  $W$  decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).

- [43] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [44] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [45] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $B_c^+$  meson lifetime using  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$  decays*, Eur. Phys. J. **C74** (2014) 2839, arXiv:1401.6932.
- [46] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, arXiv:physics/0503191.
- [47] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay  $B_c^+ \rightarrow J/\psi K^+$* , JHEP **09** (2013) 075, arXiv:1306.6723.
- [48] LHCb collaboration, R. Aaij *et al.*, *Measurements of the branching fractions and CP asymmetries of  $B^\pm \rightarrow J/\psi \pi^\pm$  and  $B^\pm \rightarrow \psi(2S) \pi^\pm$  decays*, Phys. Rev. **D85** (2012) 091105(R), arXiv:1203.3592.
- [49] See supplementary material for details, which includes Refs. [52-55].
- [50] M. Cacciari *et al.*, *Theoretical predictions for charm and bottom production at the LHC*, JHEP **10** (2012) 137, arXiv:1205.6344.
- [51] C.-F. Qiao, P. Sun, D. Yang, and R.-L. Zhu,  *$B_c$  exclusive decays to charmonium and a light meson at next-to-leading order accuracy*, Phys. Rev. **D89** (2014) 034008, arXiv:1209.5859.
- [52] J. Pumplin *et al.*, *New generation of parton distributions with uncertainties from global QCD analysis*, JHEP **07** (2002) 012, arXiv:hep-ph/0201195.
- [53] P. M. Nadolsky *et al.*, *Implications of CTEQ global analysis for collider observables*, Phys. Rev. **D78** (2008) 013004, arXiv:0802.0007.
- [54] LHCb collaboration, R. Aaij *et al.*, *Measurement of  $B$  meson production cross-sections in proton-proton collisions at  $\sqrt{s} = 7$  TeV*, JHEP **08** (2013) 117, arXiv:1306.3663.
- [55] LHCb collaboration, R. Aaij *et al.*, *Production of  $J/\psi$  and  $\Upsilon$  mesons in  $pp$  collisions at  $\sqrt{s} = 8$  TeV*, JHEP **06** (2013) 064, arXiv:1304.6977.

## LHCb collaboration

R. Aaij<sup>41</sup>, B. Adeva<sup>37</sup>, M. Adinolfi<sup>46</sup>, A. Affolder<sup>52</sup>, Z. Ajaltouni<sup>5</sup>, S. Akar<sup>6</sup>, J. Albrecht<sup>9</sup>, F. Alessio<sup>38</sup>, M. Alexander<sup>51</sup>, S. Ali<sup>41</sup>, G. Alkhazov<sup>30</sup>, P. Alvarez Cartelle<sup>37</sup>, A.A. Alves Jr<sup>25,38</sup>, S. Amato<sup>2</sup>, S. Amerio<sup>22</sup>, Y. Amhis<sup>7</sup>, L. An<sup>3</sup>, L. Anderlini<sup>17,g</sup>, J. Anderson<sup>40</sup>, R. Andreassen<sup>57</sup>, M. Andreotti<sup>16,f</sup>, J.E. Andrews<sup>58</sup>, R.B. Appleby<sup>54</sup>, O. Aquines Gutierrez<sup>10</sup>, F. Archilli<sup>38</sup>, A. Artamonov<sup>35</sup>, M. Artuso<sup>59</sup>, E. Aslanides<sup>6</sup>, G. Auriemma<sup>25,n</sup>, M. Baalouch<sup>5</sup>, S. Bachmann<sup>11</sup>, J.J. Back<sup>48</sup>, A. Badalov<sup>36</sup>, C. Baesso<sup>60</sup>, W. Baldini<sup>16</sup>, R.J. Barlow<sup>54</sup>, C. Barschel<sup>38</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>47</sup>, V. Batozskaya<sup>28</sup>, V. Battista<sup>39</sup>, A. Bay<sup>39</sup>, L. Beaucourt<sup>4</sup>, J. Beddow<sup>51</sup>, F. Bedeschi<sup>23</sup>, I. Bediaga<sup>1</sup>, S. Belogurov<sup>31</sup>, K. Belous<sup>35</sup>, I. Belyaev<sup>31</sup>, E. Ben-Haim<sup>8</sup>, G. Bencivenni<sup>18</sup>, S. Benson<sup>38</sup>, J. Benton<sup>46</sup>, A. Berezhnoy<sup>32</sup>, R. Bernet<sup>40</sup>, M.-O. Bettler<sup>47</sup>, M. van Beuzekom<sup>41</sup>, A. Bien<sup>11</sup>, S. Bifani<sup>45</sup>, T. Bird<sup>54</sup>, A. Bizzeti<sup>17,i</sup>, P.M. Bjørnstad<sup>54</sup>, T. Blake<sup>48</sup>, F. Blanc<sup>39</sup>, J. Blouw<sup>10</sup>, S. Blusk<sup>59</sup>, V. Bocci<sup>25</sup>, A. Bondar<sup>34</sup>, N. Bondar<sup>30,38</sup>, W. Bonivento<sup>15,38</sup>, S. Borghi<sup>54</sup>, A. Borgia<sup>59</sup>, M. Borsato<sup>7</sup>, T.J.V. Bowcock<sup>52</sup>, E. Bowen<sup>40</sup>, C. Bozzi<sup>16</sup>, T. Brambach<sup>9</sup>, D. Brett<sup>54</sup>, M. Britsch<sup>10</sup>, T. Britton<sup>59</sup>, J. Brodzicka<sup>54</sup>, N.H. Brook<sup>46</sup>, H. Brown<sup>52</sup>, A. Bursche<sup>40</sup>, J. Buytaert<sup>38</sup>, S. Cadeddu<sup>15</sup>, R. Calabrese<sup>16,f</sup>, M. Calvi<sup>20,k</sup>, M. Calvo Gomez<sup>36,p</sup>, P. Campana<sup>18</sup>, D. Campora Perez<sup>38</sup>, L. Capriotti<sup>54</sup>, A. Carbone<sup>14,d</sup>, G. Carboni<sup>24,l</sup>, R. Cardinale<sup>19,38,j</sup>, A. Cardini<sup>15</sup>, L. Carson<sup>50</sup>, K. Carvalho Akiba<sup>2</sup>, G. Casse<sup>52</sup>, L. Cassina<sup>20,k</sup>, L. Castillo Garcia<sup>38</sup>, M. Cattaneo<sup>38</sup>, Ch. Cauet<sup>9</sup>, R. Cenci<sup>23,t</sup>, M. Charles<sup>8</sup>, Ph. Charpentier<sup>38</sup>, M. Chefdeville<sup>4</sup>, S. Chen<sup>54</sup>, S.-F. Cheung<sup>55</sup>, N. Chiapolini<sup>40</sup>, M. Chrzaszcz<sup>40,26</sup>, X. Cid Vidal<sup>38</sup>, G. Ciezarek<sup>53</sup>, P.E.L. Clarke<sup>50</sup>, M. Clemencic<sup>38</sup>, H.V. Cliff<sup>47</sup>, J. Closier<sup>38</sup>, V. Coco<sup>38</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, V. Cogoni<sup>15</sup>, L. Cojocariu<sup>29</sup>, G. Collazuol<sup>22</sup>, P. Collins<sup>38</sup>, A. Comerma-Montells<sup>11</sup>, A. Contu<sup>15,38</sup>, A. Cook<sup>46</sup>, M. Coombes<sup>46</sup>, S. Coquereau<sup>8</sup>, G. Corti<sup>38</sup>, M. Corvo<sup>16,f</sup>, I. Counts<sup>56</sup>, B. Couturier<sup>38</sup>, G.A. Cowan<sup>50</sup>, D.C. Craik<sup>48</sup>, A.C. Crocombe<sup>48</sup>, M. Cruz Torres<sup>60</sup>, S. Cunliffe<sup>53</sup>, R. Currie<sup>53</sup>, C. D'Ambrosio<sup>38</sup>, J. Dalseno<sup>46</sup>, P. David<sup>8</sup>, P.N.Y. David<sup>41</sup>, A. Davis<sup>57</sup>, K. De Bruyn<sup>41</sup>, S. De Capua<sup>54</sup>, M. De Cian<sup>11</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, W. De Silva<sup>57</sup>, P. De Simone<sup>18</sup>, C.-T. Dean<sup>51</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>9</sup>, L. Del Buono<sup>8</sup>, N. Déléage<sup>4</sup>, D. Derkach<sup>55</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>38</sup>, A. Di Canto<sup>38</sup>, H. Dijkstra<sup>38</sup>, S. Donleavy<sup>52</sup>, F. Dordei<sup>11</sup>, M. Dorigo<sup>39</sup>, A. Dosil Suárez<sup>37</sup>, D. Dossett<sup>48</sup>, A. Dovbnya<sup>43</sup>, K. Dreimanis<sup>52</sup>, G. Dujany<sup>54</sup>, F. Dupertuis<sup>39</sup>, P. Durante<sup>38</sup>, R. Dzhelyadin<sup>35</sup>, A. Dziurda<sup>26</sup>, A. Dzyuba<sup>30</sup>, S. Easo<sup>49,38</sup>, U. Egede<sup>53</sup>, V. Egorychev<sup>31</sup>, S. Eidelman<sup>34</sup>, S. Eisenhardt<sup>50</sup>, U. Eitschberger<sup>9</sup>, R. Ekelhof<sup>9</sup>, L. Eklund<sup>51</sup>, I. El Rifai<sup>5</sup>, Ch. Elsasser<sup>40</sup>, S. Ely<sup>59</sup>, S. Esen<sup>11</sup>, H.-M. Evans<sup>47</sup>, T. Evans<sup>55</sup>, A. Falabella<sup>14</sup>, C. Färber<sup>11</sup>, C. Farinelli<sup>41</sup>, N. Farley<sup>45</sup>, S. Farry<sup>52</sup>, R. Fay<sup>52</sup>, D. Ferguson<sup>50</sup>, V. Fernandez Albor<sup>37</sup>, F. Ferreira Rodrigues<sup>1</sup>, M. Ferro-Luzzi<sup>38</sup>, S. Filippov<sup>33</sup>, M. Fiore<sup>16,f</sup>, M. Fiorini<sup>16,f</sup>, M. Firlej<sup>27</sup>, C. Fitzpatrick<sup>39</sup>, T. Fiutowski<sup>27</sup>, P. Fol<sup>53</sup>, M. Fontana<sup>10</sup>, F. Fontanelli<sup>19,j</sup>, R. Forty<sup>38</sup>, O. Francisco<sup>2</sup>, M. Frank<sup>38</sup>, C. Frei<sup>38</sup>, M. Frosini<sup>17,g</sup>, J. Fu<sup>21,38</sup>, E. Furfaro<sup>24,l</sup>, A. Gallas Torreira<sup>37</sup>, D. Galli<sup>14,d</sup>, S. Gallorini<sup>22,38</sup>, S. Gambetta<sup>19,j</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>59</sup>, Y. Gao<sup>3</sup>, J. García Pardiñas<sup>37</sup>, J. Garofoli<sup>59</sup>, J. Garra Tico<sup>47</sup>, L. Garrido<sup>36</sup>, D. Gascon<sup>36</sup>, C. Gaspar<sup>38</sup>, R. Gauld<sup>55</sup>, L. Gavardi<sup>9</sup>, G. Gazzoni<sup>5</sup>, A. Geraci<sup>21,v</sup>, E. Gersabeck<sup>11</sup>, M. Gersabeck<sup>54</sup>, T. Gershon<sup>48</sup>, Ph. Ghez<sup>4</sup>, A. Gianelle<sup>22</sup>, S. Gianì<sup>39</sup>, V. Gibson<sup>47</sup>, L. Giubega<sup>29</sup>, V.V. Gligorov<sup>38</sup>, C. Göbel<sup>60</sup>, D. Golubkov<sup>31</sup>, A. Golutvin<sup>53,31,38</sup>, A. Gomes<sup>1,a</sup>, C. Gotti<sup>20,k</sup>, M. Grabalosa Gándara<sup>5</sup>, R. Graciani Diaz<sup>36</sup>, L.A. Granado Cardoso<sup>38</sup>, E. Graugés<sup>36</sup>, E. Graverini<sup>40</sup>, G. Graziani<sup>17</sup>, A. Grecu<sup>29</sup>, E. Greening<sup>55</sup>, S. Gregson<sup>47</sup>, P. Griffith<sup>45</sup>, L. Grillo<sup>11</sup>, O. Grünberg<sup>63</sup>, B. Gui<sup>59</sup>, E. Gushchin<sup>33</sup>, Yu. Guz<sup>35,38</sup>, T. Gys<sup>38</sup>, C. Hadjivasiliou<sup>59</sup>, G. Haefeli<sup>39</sup>, C. Haen<sup>38</sup>, S.C. Haines<sup>47</sup>, S. Hall<sup>53</sup>, B. Hamilton<sup>58</sup>, T. Hampson<sup>46</sup>, X. Han<sup>11</sup>,

S. Hansmann-Menzemer<sup>11</sup>, N. Harnew<sup>55</sup>, S.T. Harnew<sup>46</sup>, J. Harrison<sup>54</sup>, J. He<sup>38</sup>, T. Head<sup>38</sup>,  
 V. Heijne<sup>41</sup>, K. Hennessy<sup>52</sup>, P. Henrard<sup>5</sup>, L. Henry<sup>8</sup>, J.A. Hernando Morata<sup>37</sup>,  
 E. van Herwijnen<sup>38</sup>, M. Heß<sup>63</sup>, A. Hicheur<sup>2</sup>, D. Hill<sup>55</sup>, M. Hoballah<sup>5</sup>, C. Hombach<sup>54</sup>,  
 W. Hulsbergen<sup>41</sup>, P. Hunt<sup>55</sup>, N. Hussain<sup>55</sup>, D. Hutchcroft<sup>52</sup>, D. Hynds<sup>51</sup>, M. Idzik<sup>27</sup>, P. Ilten<sup>56</sup>,  
 R. Jacobsson<sup>38</sup>, A. Jaeger<sup>11</sup>, J. Jalocha<sup>55</sup>, E. Jans<sup>41</sup>, P. Jaton<sup>39</sup>, A. Jawahery<sup>58</sup>, F. Jing<sup>3</sup>,  
 M. John<sup>55</sup>, D. Johnson<sup>38</sup>, C.R. Jones<sup>47</sup>, C. Joram<sup>38</sup>, B. Jost<sup>38</sup>, N. Jurik<sup>59</sup>, S. Kandybei<sup>43</sup>,  
 W. Kanso<sup>6</sup>, M. Karacson<sup>38</sup>, T.M. Karbach<sup>38</sup>, S. Karodia<sup>51</sup>, M. Kelsey<sup>59</sup>, I.R. Kenyon<sup>45</sup>,  
 T. Ketel<sup>42</sup>, B. Khanji<sup>20,38,k</sup>, C. Khurewathanakul<sup>39</sup>, S. Klaver<sup>54</sup>, K. Klimaszewski<sup>28</sup>,  
 O. Kochebina<sup>7</sup>, M. Kolpin<sup>11</sup>, I. Komarov<sup>39</sup>, R.F. Koopman<sup>42</sup>, P. Koppenburg<sup>41,38</sup>, M. Korolev<sup>32</sup>,  
 A. Kozlinskiy<sup>41</sup>, L. Kravchuk<sup>33</sup>, K. Kreplin<sup>11</sup>, M. Kreps<sup>48</sup>, G. Krocker<sup>11</sup>, P. Krokovny<sup>34</sup>,  
 F. Kruse<sup>9</sup>, W. Kucewicz<sup>26,o</sup>, M. Kucharczyk<sup>20,26,k</sup>, V. Kudryavtsev<sup>34</sup>, K. Kurek<sup>28</sup>,  
 T. Kvaratskheliya<sup>31</sup>, V.N. La Thi<sup>39</sup>, D. Lacarrere<sup>38</sup>, G. Lafferty<sup>54</sup>, A. Lai<sup>15</sup>, D. Lambert<sup>50</sup>,  
 R.W. Lambert<sup>42</sup>, G. Lanfranchi<sup>18</sup>, C. Langenbruch<sup>48</sup>, B. Langhans<sup>38</sup>, T. Latham<sup>48</sup>,  
 C. Lazzeroni<sup>45</sup>, R. Le Gac<sup>6</sup>, J. van Leerdam<sup>41</sup>, J.-P. Lees<sup>4</sup>, R. Lefèvre<sup>5</sup>, A. Leflat<sup>32</sup>,  
 J. Lefrançois<sup>7</sup>, S. Leo<sup>23</sup>, O. Leroy<sup>6</sup>, T. Lesiak<sup>26</sup>, B. Leverington<sup>11</sup>, Y. Li<sup>7</sup>, T. Likhomanenko<sup>64</sup>,  
 M. Liles<sup>52</sup>, R. Lindner<sup>38</sup>, C. Linn<sup>38</sup>, F. Lionetto<sup>40</sup>, B. Liu<sup>15</sup>, S. Lohn<sup>38</sup>, I. Longstaff<sup>51</sup>,  
 J.H. Lopes<sup>2</sup>, N. Lopez-March<sup>39</sup>, P. Lowdon<sup>40</sup>, D. Lucchesi<sup>22,r</sup>, H. Luo<sup>50</sup>, A. Lupato<sup>22</sup>,  
 E. Luppi<sup>16,f</sup>, O. Lupton<sup>55</sup>, F. Machefert<sup>7</sup>, I.V. Machikhiliyan<sup>31</sup>, F. Maciuc<sup>29</sup>, O. Maev<sup>30</sup>,  
 S. Malde<sup>55</sup>, A. Malinin<sup>64</sup>, G. Manca<sup>15,e</sup>, G. Mancinelli<sup>6</sup>, A. Mapelli<sup>38</sup>, J. Maratas<sup>5</sup>,  
 J.F. Marchand<sup>4</sup>, U. Marconi<sup>14</sup>, C. Marin Benito<sup>36</sup>, P. Marino<sup>23,t</sup>, R. Märki<sup>39</sup>, J. Marks<sup>11</sup>,  
 G. Martellotti<sup>25</sup>, A. Martín Sánchez<sup>7</sup>, M. Martinelli<sup>39</sup>, D. Martinez Santos<sup>42,38</sup>,  
 F. Martinez Vidal<sup>65</sup>, D. Martins Tostes<sup>2</sup>, A. Massafferri<sup>1</sup>, R. Matev<sup>38</sup>, Z. Mathe<sup>38</sup>,  
 C. Matteuzzi<sup>20</sup>, A. Mazurov<sup>45</sup>, M. McCann<sup>53</sup>, J. McCarthy<sup>45</sup>, A. McNab<sup>54</sup>, R. McNulty<sup>12</sup>,  
 B. McSkelly<sup>52</sup>, B. Meadows<sup>57</sup>, F. Meier<sup>9</sup>, M. Meissner<sup>11</sup>, M. Merk<sup>41</sup>, D.A. Milanese<sup>62</sup>,  
 M.-N. Minard<sup>4</sup>, N. Moggi<sup>14</sup>, J. Molina Rodriguez<sup>60</sup>, S. Monteil<sup>5</sup>, M. Morandin<sup>22</sup>, P. Morawski<sup>27</sup>,  
 A. Mordà<sup>6</sup>, M.J. Morello<sup>23,t</sup>, J. Moron<sup>27</sup>, A.-B. Morris<sup>50</sup>, R. Mountain<sup>59</sup>, F. Muheim<sup>50</sup>,  
 K. Müller<sup>40</sup>, M. Mussini<sup>14</sup>, B. Muster<sup>39</sup>, P. Naik<sup>46</sup>, T. Nakada<sup>39</sup>, R. Nandakumar<sup>49</sup>, I. Nasteva<sup>2</sup>,  
 M. Needham<sup>50</sup>, N. Neri<sup>21</sup>, S. Neubert<sup>38</sup>, N. Neufeld<sup>38</sup>, M. Neuner<sup>11</sup>, A.D. Nguyen<sup>39</sup>,  
 T.D. Nguyen<sup>39</sup>, C. Nguyen-Mau<sup>39,q</sup>, M. Nicol<sup>7</sup>, V. Niess<sup>5</sup>, R. Niet<sup>9</sup>, N. Nikitin<sup>32</sup>, T. Nikodem<sup>11</sup>,  
 A. Novoselov<sup>35</sup>, D.P. O'Hanlon<sup>48</sup>, A. Oblakowska-Mucha<sup>27,38</sup>, V. Obraztsov<sup>35</sup>, S. Oggero<sup>41</sup>,  
 S. Ogilvy<sup>51</sup>, O. Okhrimenko<sup>44</sup>, R. Oldeman<sup>15,e</sup>, C.J.G. Onderwater<sup>66</sup>, M. Orlandea<sup>29</sup>,  
 J.M. Otalora Goicochea<sup>2</sup>, A. Otto<sup>38</sup>, P. Owen<sup>53</sup>, A. Oyanguren<sup>65</sup>, B.K. Pal<sup>59</sup>, A. Palano<sup>13,c</sup>,  
 F. Palombo<sup>21,u</sup>, M. Palutan<sup>18</sup>, J. Panman<sup>38</sup>, A. Papanestis<sup>49,38</sup>, M. Pappagallo<sup>51</sup>,  
 L.L. Pappalardo<sup>16,f</sup>, C. Parkes<sup>54</sup>, C.J. Parkinson<sup>9,45</sup>, G. Passaleva<sup>17</sup>, G.D. Patel<sup>52</sup>, M. Patel<sup>53</sup>,  
 C. Patrignani<sup>19,j</sup>, A. Pearce<sup>54</sup>, A. Pellegrino<sup>41</sup>, G. Penso<sup>25,m</sup>, M. Pepe Altarelli<sup>38</sup>,  
 S. Perazzini<sup>14,d</sup>, P. Perret<sup>5</sup>, M. Perrin-Terrin<sup>6</sup>, L. Pescatore<sup>45</sup>, E. Pesen<sup>67</sup>, K. Petridis<sup>53</sup>,  
 A. Petrolini<sup>19,j</sup>, E. Picatoste Olloqui<sup>36</sup>, B. Pietrzyk<sup>4</sup>, T. Pilar<sup>48</sup>, D. Pinci<sup>25</sup>, A. Pistone<sup>19</sup>,  
 S. Playfer<sup>50</sup>, M. Plo Casasus<sup>37</sup>, F. Polci<sup>8</sup>, A. Poluektov<sup>48,34</sup>, I. Polyakov<sup>31</sup>, E. Polcarpo<sup>2</sup>,  
 A. Popov<sup>35</sup>, D. Popov<sup>10</sup>, B. Popovici<sup>29</sup>, C. Potterat<sup>2</sup>, E. Price<sup>46</sup>, J.D. Price<sup>52</sup>, J. Prisciandaro<sup>39</sup>,  
 A. Pritchard<sup>52</sup>, C. Prouve<sup>46</sup>, V. Pugatch<sup>44</sup>, A. Puig Navarro<sup>39</sup>, G. Punzi<sup>23,s</sup>, W. Qian<sup>4</sup>,  
 B. Rachwal<sup>26</sup>, J.H. Rademacker<sup>46</sup>, B. Rakotomiaramanana<sup>39</sup>, M. Rama<sup>18</sup>, M.S. Rangel<sup>2</sup>,  
 I. Raniuk<sup>43</sup>, N. Rauschmayr<sup>38</sup>, G. Raven<sup>42</sup>, F. Redi<sup>53</sup>, S. Reichert<sup>54</sup>, M.M. Reid<sup>48</sup>,  
 A.C. dos Reis<sup>1</sup>, S. Ricciardi<sup>49</sup>, S. Richards<sup>46</sup>, M. Rihl<sup>38</sup>, K. Rinnert<sup>52</sup>, V. Rives Molina<sup>36</sup>,  
 P. Robbe<sup>7</sup>, A.B. Rodrigues<sup>1</sup>, E. Rodrigues<sup>54</sup>, P. Rodriguez Perez<sup>54</sup>, S. Roiser<sup>38</sup>,  
 V. Romanovsky<sup>35</sup>, A. Romero Vidal<sup>37</sup>, M. Rotondo<sup>22</sup>, J. Rouvinet<sup>39</sup>, T. Ruf<sup>38</sup>, H. Ruiz<sup>36</sup>,  
 P. Ruiz Valls<sup>65</sup>, J.J. Saborido Silva<sup>37</sup>, N. Sagidova<sup>30</sup>, P. Sail<sup>51</sup>, B. Saitta<sup>15,e</sup>,



V. Salustino Guimaraes<sup>2</sup>, C. Sanchez Mayordomo<sup>65</sup>, B. Sanmartin Sedes<sup>37</sup>, R. Santacesaria<sup>25</sup>, C. Santamarina Rios<sup>37</sup>, E. Santovetti<sup>24,l</sup>, A. Sarti<sup>18,m</sup>, C. Satriano<sup>25,n</sup>, A. Satta<sup>24</sup>, D.M. Saunders<sup>46</sup>, D. Savrina<sup>31,32</sup>, M. Schiller<sup>42</sup>, H. Schindler<sup>38</sup>, M. Schlupp<sup>9</sup>, M. Schmelling<sup>10</sup>, B. Schmidt<sup>38</sup>, O. Schneider<sup>39</sup>, A. Schopper<sup>38</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>38</sup>, B. Sciascia<sup>18</sup>, A. Sciubba<sup>25,m</sup>, A. Semennikov<sup>31</sup>, I. Sepp<sup>53</sup>, N. Serra<sup>40</sup>, J. Serrano<sup>6</sup>, L. Sestini<sup>22</sup>, P. Seyfert<sup>11</sup>, M. Shapkin<sup>35</sup>, I. Shapoval<sup>16,43,f</sup>, Y. Shcheglov<sup>30</sup>, T. Shears<sup>52</sup>, L. Shekhtman<sup>34</sup>, V. Shevchenko<sup>64</sup>, A. Shires<sup>9</sup>, R. Silva Coutinho<sup>48</sup>, G. Simi<sup>22</sup>, M. Sirendi<sup>47</sup>, N. Skidmore<sup>46</sup>, I. Skillicorn<sup>51</sup>, T. Skwarnicki<sup>59</sup>, N.A. Smith<sup>52</sup>, E. Smith<sup>55,49</sup>, E. Smith<sup>53</sup>, J. Smith<sup>47</sup>, M. Smith<sup>54</sup>, H. Snoek<sup>41</sup>, M.D. Sokoloff<sup>57</sup>, F.J.P. Soler<sup>51</sup>, F. Soomro<sup>39</sup>, D. Souza<sup>46</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>9</sup>, P. Spradlin<sup>51</sup>, S. Sridharan<sup>38</sup>, F. Stagni<sup>38</sup>, M. Stahl<sup>11</sup>, S. Stahl<sup>11</sup>, O. Steinkamp<sup>40</sup>, O. Stenyakin<sup>35</sup>, S. Stevenson<sup>55</sup>, S. Stoica<sup>29</sup>, S. Stone<sup>59</sup>, B. Storaci<sup>40</sup>, S. Stracka<sup>23,t</sup>, M. Straticiu<sup>29</sup>, U. Straumann<sup>40</sup>, R. Stroili<sup>22</sup>, V.K. Subbiah<sup>38</sup>, L. Sun<sup>57</sup>, W. Sutcliffe<sup>53</sup>, K. Swientek<sup>27</sup>, S. Swientek<sup>9</sup>, V. Syropoulos<sup>42</sup>, M. Szczekowski<sup>28</sup>, P. Szczypka<sup>39,38</sup>, T. Szumlak<sup>27</sup>, S. T'Jampens<sup>4</sup>, M. Teklishyn<sup>7</sup>, G. Tellarini<sup>16,f</sup>, F. Teubert<sup>38</sup>, C. Thomas<sup>55</sup>, E. Thomas<sup>38</sup>, J. van Tilburg<sup>41</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>39</sup>, J. Todd<sup>57</sup>, S. Tol<sup>42</sup>, L. Tomassetti<sup>16,f</sup>, D. Tonelli<sup>38</sup>, S. Topp-Joergensen<sup>55</sup>, N. Torr<sup>55</sup>, E. Tournefier<sup>4</sup>, S. Tourneur<sup>39</sup>, M.T. Tran<sup>39</sup>, M. Tresch<sup>40</sup>, A. Trisovic<sup>38</sup>, A. Tsaregorodtsev<sup>6</sup>, P. Tsopelas<sup>41</sup>, N. Tuning<sup>41</sup>, M. Ubeda Garcia<sup>38</sup>, A. Ukleja<sup>28</sup>, A. Ustyuzhanin<sup>64</sup>, U. Uwer<sup>11</sup>, C. Vacca<sup>15</sup>, V. Vagnoni<sup>14</sup>, G. Valenti<sup>14</sup>, A. Vallier<sup>7</sup>, R. Vazquez Gomez<sup>18</sup>, P. Vazquez Regueiro<sup>37</sup>, C. Vázquez Sierra<sup>37</sup>, S. Vecchi<sup>16</sup>, J.J. Velthuis<sup>46</sup>, M. Veltri<sup>17,h</sup>, G. Veneziano<sup>39</sup>, M. Vesterinen<sup>11</sup>, B. Viaud<sup>7</sup>, D. Vieira<sup>2</sup>, M. Vieites Diaz<sup>37</sup>, X. Vilasis-Cardona<sup>36,p</sup>, A. Vollhardt<sup>40</sup>, D. Volyanskyy<sup>10</sup>, D. Voong<sup>46</sup>, A. Vorobyev<sup>30</sup>, V. Vorobyev<sup>34</sup>, C. Voß<sup>63</sup>, J.A. de Vries<sup>41</sup>, R. Waldi<sup>63</sup>, C. Wallace<sup>48</sup>, R. Wallace<sup>12</sup>, J. Walsh<sup>23</sup>, S. Wandernoth<sup>11</sup>, J. Wang<sup>59</sup>, D.R. Ward<sup>47</sup>, N.K. Watson<sup>45</sup>, D. Websdale<sup>53</sup>, M. Whitehead<sup>48</sup>, J. Wicht<sup>38</sup>, D. Wiedner<sup>11</sup>, G. Wilkinson<sup>55,38</sup>, M.P. Williams<sup>45</sup>, M. Williams<sup>56</sup>, H.W. Wilschut<sup>66</sup>, F.F. Wilson<sup>49</sup>, J. Wimberley<sup>58</sup>, J. Wishahi<sup>9</sup>, W. Wislicki<sup>28</sup>, M. Witek<sup>26</sup>, G. Wormser<sup>7</sup>, S.A. Wotton<sup>47</sup>, S. Wright<sup>47</sup>, K. Wyllie<sup>38</sup>, Y. Xie<sup>61</sup>, Z. Xing<sup>59</sup>, Z. Xu<sup>39</sup>, Z. Yang<sup>3</sup>, X. Yuan<sup>3</sup>, O. Yushchenko<sup>35</sup>, M. Zangoli<sup>14</sup>, M. Zavertyaev<sup>10,b</sup>, L. Zhang<sup>59</sup>, W.C. Zhang<sup>12</sup>, Y. Zhang<sup>3</sup>, A. Zhelezov<sup>11</sup>, A. Zhokhov<sup>31</sup>, L. Zhong<sup>3</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

<sup>9</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>10</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>11</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>12</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>13</sup>Sezione INFN di Bari, Bari, Italy

<sup>14</sup>Sezione INFN di Bologna, Bologna, Italy

<sup>15</sup>Sezione INFN di Cagliari, Cagliari, Italy

<sup>16</sup>Sezione INFN di Ferrara, Ferrara, Italy

<sup>17</sup>Sezione INFN di Firenze, Firenze, Italy

<sup>18</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

<sup>19</sup>Sezione INFN di Genova, Genova, Italy

- <sup>20</sup> *Sezione INFN di Milano Bicocca, Milano, Italy*
- <sup>21</sup> *Sezione INFN di Milano, Milano, Italy*
- <sup>22</sup> *Sezione INFN di Padova, Padova, Italy*
- <sup>23</sup> *Sezione INFN di Pisa, Pisa, Italy*
- <sup>24</sup> *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- <sup>25</sup> *Sezione INFN di Roma La Sapienza, Roma, Italy*
- <sup>26</sup> *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- <sup>27</sup> *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- <sup>28</sup> *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- <sup>29</sup> *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- <sup>30</sup> *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- <sup>31</sup> *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>32</sup> *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>33</sup> *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- <sup>34</sup> *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- <sup>35</sup> *Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>36</sup> *Universitat de Barcelona, Barcelona, Spain*
- <sup>37</sup> *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>38</sup> *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>39</sup> *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>40</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>41</sup> *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>42</sup> *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>43</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>44</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>45</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>46</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>47</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>48</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>49</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>50</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>51</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>52</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>53</sup> *Imperial College London, London, United Kingdom*
- <sup>54</sup> *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>55</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>56</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>57</sup> *University of Cincinnati, Cincinnati, OH, United States*
- <sup>58</sup> *University of Maryland, College Park, MD, United States*
- <sup>59</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>60</sup> *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>61</sup> *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to <sup>3</sup>*
- <sup>62</sup> *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>8</sup>*
- <sup>63</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>11</sup>*
- <sup>64</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to <sup>31</sup>*
- <sup>65</sup> *Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain, associated to <sup>36</sup>*
- <sup>66</sup> *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to <sup>41</sup>*
- <sup>67</sup> *Celal Bayar University, Manisa, Turkey, associated to <sup>38</sup>*

<sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil*

- <sup>b</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>c</sup> *Università di Bari, Bari, Italy*
- <sup>d</sup> *Università di Bologna, Bologna, Italy*
- <sup>e</sup> *Università di Cagliari, Cagliari, Italy*
- <sup>f</sup> *Università di Ferrara, Ferrara, Italy*
- <sup>g</sup> *Università di Firenze, Firenze, Italy*
- <sup>h</sup> *Università di Urbino, Urbino, Italy*
- <sup>i</sup> *Università di Modena e Reggio Emilia, Modena, Italy*
- <sup>j</sup> *Università di Genova, Genova, Italy*
- <sup>k</sup> *Università di Milano Bicocca, Milano, Italy*
- <sup>l</sup> *Università di Roma Tor Vergata, Roma, Italy*
- <sup>m</sup> *Università di Roma La Sapienza, Roma, Italy*
- <sup>n</sup> *Università della Basilicata, Potenza, Italy*
- <sup>o</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- <sup>p</sup> *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- <sup>q</sup> *Hanoi University of Science, Hanoi, Viet Nam*
- <sup>r</sup> *Università di Padova, Padova, Italy*
- <sup>s</sup> *Università di Pisa, Pisa, Italy*
- <sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*
- <sup>u</sup> *Università degli Studi di Milano, Milano, Italy*
- <sup>v</sup> *Politecnico di Milano, Milano, Italy*