

Higgs boson gluon-fusion production in N³LO QCD

Charalampos Anastasiou,¹ Claude Duhr,^{2,3,*} Falko Dulat,¹ Franz Herzog,⁴ and Bernhard Mistlberger¹

¹*Institute for Theoretical Physics, ETH Zürich, 8093 Zürich, Switzerland*

²*CERN Theory Division, 1211 Geneva 23, Switzerland*

³*Center for Cosmology, Particle Physics and Phenomenology (CP3),
Université Catholique de Louvain, 1348 Louvain-La-Neuve, Belgium*

⁴*Nikhef, Science Park 105, 1098 XG Amsterdam, The Netherlands*

(Dated: March 23, 2015)

We present the cross-section for the production of a Higgs boson at hadron-colliders at next-to-next-to-next-to-leading order (N³LO) in perturbative QCD. The calculation is based on a method to perform a series expansion of the partonic cross-section around the threshold limit to an arbitrary order. We perform this expansion to sufficiently high order to obtain the value of the hadronic cross at N³LO in the large top-mass limit. For renormalisation and factorisation scales equal to half the Higgs mass, the N³LO corrections are of the order of +2.2%. The total scale variation at N³LO is 3%, reducing the uncertainty due to missing higher order QCD corrections by a factor of three.

PACS numbers: 12.38.Bx

The success of the Large Hadron Collider (LHC) experiments in the exploration and interpretation of phenomena at the TeV scale is due, on the one hand, to amazing experimental and technological advances and, on the other hand, extraordinary progress in perturbative QCD. In particular, the discovery of the Higgs boson [1] by the LHC experiments has initiated an era of precision studies of the properties of the Higgs boson, where precise theory predictions for Higgs observables play an indispensable role.

The inclusive gluon-fusion cross-section is a prototypical example of a theoretical input for the interpretation of the experimental observations. It enters not only into the extraction of the Higgs-boson couplings from the measurements, but it could also play an important role in identifying deviations from the Standard Model predictions in Higgs physics. Unfortunately, the theory predictions for the inclusive cross-section are plagued by significant theoretical uncertainties. Scale variations at NNLO indicate that missing higher order effects are of the order of $\pm 9\%$ at LHC energies [2, 3], and the size of this uncertainty is comparable to the experimental uncertainty from the LHC Run 1 [4, 5]. Hence, with a few more years of data taking the theoretical uncertainty will be dominant, demanding for an update of the current theoretical predictions.

In this context, a vigorous effort has recently been made to compute the inclusive gluon-fusion cross-section at next-to-next-to-next-to-leading order (N³LO) in perturbative QCD. The cross-section at N³LO receives contributions from many different building blocks, all of which have been computed over the last years, at least partially. The three-loop corrections to Higgs production in gluon fusion have been obtained in ref. [6], and the corrections from the emission of an additional parton at one

or two loops were computed in ref. [7–11]. In order to obtain a finite result, appropriate ultra-violet and infrared counterterms need to be included [12–14]. While all of these contributions had been computed in full generality, contributions from the emission of two partons at one loop and three partons at tree-level had only been computed in an approximate manner. In particular, for these contributions the first two terms in the expansion around threshold could be obtained [15–18], confirming previous results for logarithmically enhanced terms in the cross section [21] and resulting in the complete computation of the inclusive gluon-fusion cross-section at N³LO in the soft-virtual [17–19] and next-to-soft approximations [20]. Owing to the universality of soft emissions, the previous results have sparked various new results for QCD processes at N³LO in the soft-virtual approximation [22].

Despite this progress, the soft-virtual and next-to-soft approximations are insufficient to make reliable predictions for the cross-section, owing to a slow convergence of the threshold expansion [20]. In this Letter we close this gap, and we present the gluon-fusion Higgs production cross-section at N³LO in perturbative QCD. We emphasise that this is the first ever complete computation of a cross-section at N³LO at a hadron collider.

We will describe the main result of our computation in this Letter, while a detailed account of the mathematical and computational methods will be presented elsewhere. Here it suffices to say that we work in the framework of reverse-unitarity [23, 24], and we perform a complete reduction of the cross-section to master integrals, without any approximations. For the double and triple-emission contributions at N³LO, we can derive differential equations satisfied by the master integrals [23–25], which we solve as generalised power series around the threshold limit. In this way, we obtain at least 37 terms in the

threshold expansion of each master integral. An important part of our computation has been the evaluation of the boundary conditions which are needed for solving the differential equations for the master integrals. Many of the boundary conditions required in this project had already been derived in the context of the soft-virtual and next-to-soft results [15–18, 20]. Using similar techniques, we have computed the remaining few unknown boundary conditions for master integrals which start to be relevant only at a high order in the threshold expansion.

Having at our disposal the complete set of master integrals as expansions around the threshold limit, we can easily obtain the cross-sections at N³LO for all partonic channels contributing to Higgs production via gluon fusion. The partonic cross-sections are related to the hadronic cross-section at the LHC through the integral

$$\sigma = \sum_{i,j} \int dx_1 dx_2 f_i(x_1, \mu_f) f_j(x_2, \mu_f) \hat{\sigma}_{ij}(z, \mu_r, \mu_f), \quad (1)$$

where the summation indices i, j run over the parton flavors in the proton, f_i are parton densities and $\hat{\sigma}_{ij}$ are partonic cross-sections. Furthermore, we define $z = \frac{m_H^2}{s}$, where m_H is the mass of the Higgs boson and \sqrt{s} is the partonic center-of-mass energy, related to the hadronic center-of-mass energy \sqrt{S} through $s = x_1 x_2 S$. The renormalisation and factorisation scales are denoted by μ_r and μ_f . We work in an effective theory approach where the top-quark is integrated out. The effective Lagrangian describing the interaction of the Higgs boson and the gluons is,

$$\mathcal{L}_{\text{eff}} = -\frac{C}{4} H G_{\mu\nu}^a G^{a\mu\nu}, \quad (2)$$

where H is the Higgs field, $G_{\mu\nu}^a$ is the gluon field strength tensor and C the Wilson coefficient, known up to N⁴LO [26]. We expand the partonic cross-sections into a perturbative series in the strong coupling constant evaluated at the scale μ_r ,

$$\hat{\sigma}_{ij} = \hat{\sigma}_0 \left[\delta_{ig} \delta_{jg} \delta(1-z) + \sum_{\ell=1}^{\infty} \left(\frac{\alpha_s(\mu_r)}{\pi} \right)^\ell \hat{\sigma}_{ij}^{(\ell)} \right]. \quad (3)$$

In this expression $\hat{\sigma}_0$ denotes the leading order cross-section, and the terms through NNLO in the above expansion have been computed in [23, 30, 31]. The main result of this Letter is the result for the N³LO coefficient, corresponding to $\ell = 3$ in eq. (3), for all possible parton flavours in the initial state. We cast the N³LO coefficients in the form

$$\hat{\sigma}_{ij}^{(3)} = \lim_{N \rightarrow \infty} \hat{\sigma}_{ij}^{(3,N)}, \quad (4)$$

where we introduce the truncated threshold expansions defined by

$$\hat{\sigma}_{ij}^{(3,N)} = \delta_{ig} \delta_{jg} \hat{\sigma}_{\text{SV}}^{(3)} + \sum_{n=0}^N c_{ij}^{(n)} (1-z)^n. \quad (5)$$




FIG. 1: The N³LO correction from the gg channel to the hadronic cross-section as a function of the truncation order N in the threshold expansion for the scale choice $\mu = m_H$.

Here, $\hat{\sigma}_{\text{SV}}^{(3)}$ denotes the soft-virtual cross-section at N³LO of ref. [17–19] and $N = 0$ is the next-to-soft approximation of ref. [20]. Using our method for the threshold expansion of the master integrals, we were able to determine the $c_{ij}^{(n)}$ analytically up to at least $n = 30$. Note that at any given order in the expansion these coefficients are polynomials in $\log(1-z)$. While this approach does not cast the partonic cross-sections in a closed analytic form, we argue that it yields the complete result for the value of the hadronic cross-section. In Fig. 1 we show the contribution of the partonic cross-section coefficients N³LO to the hadronic cross-section for a proton-proton collider with 13 TeV center-of-mass energy as a function of the truncation order N . We use NNLO MSTW2008 [28] parton densities and a value for the strong coupling at the mass of the Z -boson of $\alpha_s(m_Z) = 0.117$ as initial value for the evolution, and we set the factorisation scale to $\mu_f = m_H$. We observe that the threshold expansion stabilises starting from $N = 4$, leaving a negligible truncation uncertainty for the hadronic cross-section thereafter. We note, though, that we observe a very small, but systematic, increase of the expansion in the range $N \in [15, 37]$, as illustrated in Fig. 1. We have observed that a similar behaviour is observed for the threshold expansion at NNLO. The systematic increase originates from values of the partonic cross-section at very small z . Indeed, this increase appears only in the contributions to the hadronic cross-section integral for values $z < 0.1$. It is natural that the terms of the threshold expansion computed here do not furnish a good approximation of the hadronic integral in the small z region due to the divergent high energy behaviour of the partonic cross-sections [29]. However, it is observed that this region is suppressed in the total hadronic integral and for $z < 0.1$ contributes less than 0.4% of the total N³LO correction. The same region at NLO and NNLO, where analytic ex-




FIG. 2: Scale variation of the gluon fusion cross-section at all perturbative orders through N³LO.

pressions valid for all regions are known, is similarly suppressed. We therefore believe that the uncertainty of our computation for the hadronic cross-section due to the truncation of the threshold expansion is negligible (less than 0.2%).

In Fig. 2 we present the hadronic gluon-fusion Higgs production cross-section at N³LO as a function of a common renormalisation and factorisation scale $\mu = \mu_r = \mu_f$. We observe a significant reduction of the sensitivity of the cross-section to the scale μ . Inside a range

$\mu \in [\frac{m_H}{4}, m_H]$ the cross-section at N³LO varies in the interval $[-2.7\%, +0.3\%]$ with respect to the cross-section value at the central scale $\mu = \frac{m_H}{2}$. For comparison, we note that the corresponding scale variation at NNLO is about $\pm 9\%$ [2, 3]. This improvement in the precision of the Higgs cross-section is a major accomplishment due to our calculation and will have a strong impact on future measurements of Higgs-boson properties. Furthermore, even though for the scale choice $\mu = \frac{m_H}{2}$ the N³LO corrections change the cross-section by about +2.2%, this correction is captured by the scale variation estimate for the missing higher order effects of the NNLO result at that scale. We illustrate this point in Fig. 3, where we present the hadronic cross-section as a function of the hadronic center-of-mass energy \sqrt{S} at the scale $\mu = \frac{m_H}{2}$. We observe that the N³LO scale uncertainty band is included within the NNLO band, indicating that the perturbative expansion of the hadronic cross-section is convergent. However, we note that for a larger scale choice, e.g., $\mu = m_H$, the convergence of the perturbative series is slower than for $\mu = \frac{m_H}{2}$.

In table I we quote the gluon fusion cross section in effective theory at N³LO for different LHC energies. The perturbative uncertainty is determined by varying the common renormalisation and factorisation scale in the interval $[\frac{m_H}{4}, m_H]$ around $\frac{m_H}{2}$ and in the interval $[\frac{m_H}{2}, 2m_H]$ around m_H .

σ/pb	2 TeV	7 TeV	8 TeV	13 TeV	14 TeV
$\mu = \frac{m_H}{2}$	$0.99^{+0.43\%}_{-4.65\%}$	$15.31^{+0.31\%}_{-3.08\%}$	$19.47^{+0.32\%}_{-2.99\%}$	$44.31^{+0.31\%}_{-2.64\%}$	$49.87^{+0.32\%}_{-2.61\%}$
$\mu = m_H$	$0.94^{+4.87\%}_{-7.35\%}$	$14.84^{+3.18\%}_{-5.27\%}$	$18.90^{+3.08\%}_{-5.02\%}$	$43.14^{+2.71\%}_{-4.45\%}$	$48.57^{+2.68\%}_{-4.24\%}$

TABLE I: The gluon fusion cross-section in picobarn in the effective theory for different collider energies in the interval $[\frac{m_H}{4}, m_H]$ around $\mu = \frac{m_H}{2}$ and in the interval $[\frac{m_H}{2}, 2m_H]$ around $\mu = m_H$.

Given the substantial reduction of the scale uncertainty at N³LO, the question naturally arises whether other sources of theoretical uncertainty may contribute at a similar level. In the remainder of this Letter we briefly comment on this issue, leaving a more detailed quantitative study for future work.

First, we note that given the small size of the N³LO corrections compared to NNLO, we expect that an estimate for the higher-order corrections at N⁴LO and beyond can be obtained from the scale variation uncertainty. Alternatively, partial N⁴LO results can be obtained by means of factorisation theorems for threshold resummation. However, we expect that the insight from resummation on the N⁴LO soft contributions is only

qualitative given the importance of next-to-soft, next-to-next-to-soft and purely virtual contributions observed at N³LO, as seen in Fig. 1.

Electroweak corrections to Higgs production have been calculated through two loops in ref. [32], and estimated at three loops in ref. [33]. They furnish a correction of less than +5% to the inclusive cross-section. Thus, they are not negligible at the level of accuracy indicated by the scale variation at N³LO and need to be combined with our result in the future. Mixed QCD-electroweak or purely electroweak corrections of even higher order are expected to contribute at the sub-percent level and should be negligible.

Next, we have to comment on our assumption that the




FIG. 3: The gluon fusion cross-section at all perturbative orders through N³LO in the scale interval $[\frac{m_H}{4}, m_H]$ as a function of the center-of-mass energy \sqrt{S} .

top-quark is infinitely heavy and can be integrated out, see eq. (2). Moreover, we assumed that all other quarks have a zero Yukawa coupling. Finite quark mass effects are important, but it is sufficient that they are included through NLO or NNLO. Indeed, finite quark-mass effects have been computed fully through NLO in QCD [30], while subleading top-quark mass corrections have been computed at NNLO systematically as an expansion in the inverse top-quark mass [34]. In these references it was observed that through NLO finite quark mass effects amount to about 8% of the K-factor. At NNLO, the known $\frac{1}{m_{\text{top}}}$ corrections affect the cross-section at the $\sim 1\%$ level. A potentially significant contribution at NNLO which has not yet been computed in the literature originates from diagrams with both a top and bottom quark Yukawa coupling. Assuming a similar perturbative pattern as for top-quark only diagrams in the effective theory, eq. (2), higher-order effects could be of the order of 2%. We thus conclude that the computation of the top-bottom interference through NNLO is highly desired in the near future.

Finally, the computation of the hadronic cross-section relies crucially on the knowledge of the strong coupling constant and the parton densities. After our calculation, the uncertainty coming from these quantities has become dominant. Further progress in the determination of parton densities must be anticipated in the next few years due to the inclusion of LHC data in the global fits and the impressive advances in NNLO computations, improving the theoretical accuracy of many standard candle processes.

To conclude, we have presented in this Letter the computation of the gluon-fusion Higgs production cross-section through N³LO in perturbative QCD. While a thorough study of the impact of electroweak and quark mass effects is left for future work, we expect that the remaining theoretical uncertainty on the inclusive Higgs production cross-section is expected to be reduced to roughly half, which will bring important benefits in the study of the properties of the Higgs boson at the LHC Run 2. Besides its direct phenomenological impact, we believe that our result is also a major advance in our understanding of perturbative QCD, as it opens the door to push the theoretical predictions for large classes of inclusive processes to N³LO accuracy, like Drell-Yan production, associated Higgs production and Higgs production via bottom fusion. Moreover, on the more technical side, our result constitutes the first independent validation of the gluon splitting function at NNLO [14], because the latter is required to cancel all the infrared poles in the inclusive cross-section. In addition, we expect that the techniques developed throughout this work are not restricted to inclusive cross-sections, but it should be possible to extend them to certain classes of differential distributions, like rapidity distributions for Drell-Yan and Higgs production, thereby paving the way to a new era of precision QCD.

Acknowledgements: We are grateful to Elisabetta Furlan, Thomas Gehrmann and A. Lazopoulos for our collaboration on the many aspects of the Higgs cross-section N³LO project which are not covered in this Letter. We thank A. Lazopoulos in particular for an independent implementation of our results in `ihixs` and numerical comparisons. Research supported by the Swiss National Science Foundation (SNF) under contract 200021-143781 and the European Commission through the ERC grants “IterQCD”, “HEPGAME” and “MathAm”.

* On leave from the “Fonds National de la Recherche Scientifique” (FNRS), Belgium.

- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012); S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).
- [2] C. Anastasiou, S. Bühler, F. Herzog and A. Lazopoulos, JHEP **1204**, 004 (2012).
- [3] C. Anastasiou, S. Bühler, F. Herzog and A. Lazopoulos, JHEP **1112**, 058 (2011).
- [4] V. Khachatryan *et al.* [CMS Collaboration], arXiv:1412.8662 [hep-ex].
- [5] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. D **90**, no. 11, 112015 (2014) [arXiv:1408.7084 [hep-ex]].
- [6] P.A. Baikov, K.G. Chetyrkin, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, Phys. Rev. Lett. **102**, 212002 (2009); T. Gehrmann, E. W. N. Glover, T. Huber, N. Iqizlerli, C. Studerus, JHEP **1006**, 094

- (2010).
- [7] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *JHEP* **1312**, 088 (2013) [arXiv:1311.1425 [hep-ph]];
- [8] W. B. Kilgore, *Phys. Rev. D* **89** (2014) 7, 073008 [arXiv:1312.1296 [hep-ph]].
- [9] T. Gehrmann, M. Jaquier, E. W. N. Glover and A. Kououtsakis, *JHEP* **1202**, 056 (2012), [arXiv:1112.3554 [hep-ph]].
- [10] C. Duhr and T. Gehrmann, *Phys. Lett. B* **727**, 452 (2013) [arXiv:1309.4393 [hep-ph]]; Y. Li and H. X. Zhu, *JHEP* **1311**, 080 (2013) [arXiv:1309.4391 [hep-ph]].
- [11] C. Duhr, T. Gehrmann and M. Jaquier, *JHEP* **1502** (2015) 077 [arXiv:1411.3587 [hep-ph]]; F. Dulat and B. Mistlberger, [arXiv:1411.3586 [hep-ph]].
- [12] O. V. Tarasov, A. A. Vladimirov and A. Y. Zharkov, *Phys. Lett. B* **93**, 429 (1980); S. A. Larin and J. A. M. Vermaseren, *Phys. Lett. B* **303**, 334 (1993) [hep-ph/9302208]; T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, *Phys. Lett. B* **400**, 379 (1997) [hep-ph/9701390]; M. Czakon, *Nucl. Phys. B* **710**, 485 (2005) [hep-ph/0411261].
- [13] C. Anastasiou, S. Bühler, C. Duhr and F. Herzog, *JHEP* **1211**, 062 (2012); M. Höschele, J. Hoff, A. Pak, M. Steinhauser, T. Ueda, *Phys. Lett. B* **721**, 244 (2013); S. Bühler and A. Lazopoulos, *JHEP* **1310**, 096 (2013).
- [14] S. Moch, J. A. M. Vermaseren and A. Vogt, *Nucl. Phys. B* **688**, 101 (2004), [hep-ph/0403192]; *Nucl. Phys. B* **691**, 129 (2004) [hep-ph/0404111].
- [15] C. Anastasiou, C. Duhr, F. Dulat, B. Mistlberger, *JHEP* **1307**, 003 (2013) [arXiv:1302.4379 [hep-ph]].
- [16] H. X. Zhu, *JHEP* **1502** (2015) 155 [arXiv:1501.00236 [hep-ph]].
- [17] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, *Phys. Lett. B* **737** (2014) 325 [arXiv:1403.4616 [hep-ph]].
- [18] Y. Li, A. von Manteuffel, R. M. Schabinger and H. X. Zhu, *Phys. Rev. D* **90** (2014) 5, 053006 [arXiv:1404.5839 [hep-ph]].
- [19] Y. Li, A. von Manteuffel, R. M. Schabinger and H. X. Zhu, *Phys. Rev. D* **91** (2015) 3, 036008 [arXiv:1412.2771 [hep-ph]].
- [20] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog and B. Mistlberger, [arXiv:1411.3584 [hep-ph]].
- [21] S. Moch and A. Vogt, *Phys. Lett. B* **631**, 48 (2005) [hep-ph/0508265]; E. Laenen and L. Magnea, *Phys. Lett. B* **632** (2006) 270 [hep-ph/0508284]; N. A. Lo Presti, A. A. Almasy and A. Vogt, *Phys. Lett. B* **737** (2014) 120 [arXiv:1407.1553 [hep-ph]]; D. de Florian, J. Mazzitelli, S. Moch and A. Vogt, *JHEP* **1410** (2014) 176 [arXiv:1408.6277 [hep-ph]].
- [22] T. Ahmed, M. Mahakhud, N. Rana and V. Ravindran, *Phys. Rev. Lett.* **113** (2014) 11, 112002 [arXiv:1404.0366 [hep-ph]]; T. Ahmed, M. K. Mandal, N. Rana and V. Ravindran, *Phys. Rev. Lett.* **113** (2014) 212003 [arXiv:1404.6504 [hep-ph]]; T. Ahmed, N. Rana and V. Ravindran, *JHEP* **1410** (2014) 139 [arXiv:1408.0787 [hep-ph]]; T. Ahmed, M. K. Mandal, N. Rana and V. Ravindran, *JHEP* **1502** (2015) 131 [arXiv:1411.5301 [hep-ph]]; M. C. Kumar, M. K. Mandal and V. Ravindran, [arXiv:1412.3357 [hep-ph]]; S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Nucl. Phys. B* **888** (2014) 75 [arXiv:1405.4827 [hep-ph]].
- [23] C. Anastasiou and K. Melnikov, *Nucl. Phys. B* **646**, 220 (2002) [hep-ph/0207004].
- [24] C. Anastasiou and K. Melnikov, *Phys. Rev. D* **67** (2003) 037501 [hep-ph/0208115]; C. Anastasiou, L. J. Dixon and K. Melnikov, *Nucl. Phys. Proc. Suppl.* **116** (2003) 193 [hep-ph/0211141]; C. Anastasiou, L. J. Dixon, K. Melnikov and F. Petriello, *Phys. Rev. Lett.* **91** (2003) 182002 [hep-ph/0306192]; C. Anastasiou, L. J. Dixon, K. Melnikov and F. Petriello, *Phys. Rev. D* **69** (2004) 094008 [hep-ph/0312266].
- [25] A. V. Kotikov, *Phys. Lett. B* **259**, 314 (1991); *Phys. Lett. B* **267**, 123 (1991).
- [26] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, *Nucl. Phys. B* **510**, 61 (1998); Y. Schroder and M. Steinhauser, *JHEP* **0601**, 051 (2006); K. G. Chetyrkin, J. H. Kuhn and C. Sturm, *Nucl. Phys. B* **744**, 121 (2006).
- [27] S. Catani, D. de Florian and M. Grazzini, *JHEP* **0105**, 025 (2001); R. V. Harlander and W. B. Kilgore, *Phys. Rev. D* **64**, 013015 (2001)
- [28] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, *Eur. Phys. J. C* **63**, 189 (2009)
- [29] F. Hautmann, *Phys. Lett. B* **535** (2002) 159 [hep-ph/0203140].
- [30] D. Graudenz, M. Spira and P. M. Zerwas, *Phys. Rev. Lett.* **70** (1993) 1372; S. Dawson, *Nucl. Phys. B* **359** (1991) 283; A. Djouadi, M. Spira and P. M. Zerwas, *Phys. Lett. B* **264** (1991) 440; M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, *Nucl. Phys. B* **453** (1995) 17 [hep-ph/9504378]; R. Harlander and P. Kant, *JHEP* **0512**, 015 (2005) [hep-ph/0509189]. C. Anastasiou, S. Beerli, S. Bucherer, A. Daleo and Z. Kunszt, *JHEP* **0701**, 082 (2007) [hep-ph/0611236]. U. Aglietti, R. Bonciani, G. Degrassi and A. Vicini, *JHEP* **0701**, 021 (2007) [hep-ph/0611266]. R. Bonciani, G. Degrassi and A. Vicini, *JHEP* **0711**, 095 (2007) [arXiv:0709.4227 [hep-ph]]. C. Anastasiou, S. Bucherer and Z. Kunszt, *JHEP* **0910**, 068 (2009) [arXiv:0907.2362 [hep-ph]].
- [31] R. V. Harlander and W. B. Kilgore, *Phys. Rev. Lett.* **88**, 201801 (2002) [hep-ph/0201206]. V. Ravindran, J. Smith and W. L. van Neerven, *Nucl. Phys. B* **665**, 325 (2003) [hep-ph/0302135].
- [32] S. Actis, G. Passarino, C. Sturm and S. Uccirati, *Nucl. Phys. B* **811**, 182 (2009) [arXiv:0809.3667 [hep-ph]]. S. Actis, G. Passarino, C. Sturm and S. Uccirati, *Phys. Lett. B* **670**, 12 (2008) [arXiv:0809.1301 [hep-ph]]. U. Aglietti, R. Bonciani, G. Degrassi and A. Vicini, *Phys. Lett. B* **595**, 432 (2004) [hep-ph/0404071].
- [33] C. Anastasiou, R. Boughezal and F. Petriello, *JHEP* **0904**, 003 (2009) [arXiv:0811.3458 [hep-ph]].
- [34] R. V. Harlander and K. J. Ozeren, *JHEP* **0911**, 088 (2009) [arXiv:0909.3420 [hep-ph]]. A. Pak, M. Rogal and M. Steinhauser, *JHEP* **1002**, 025 (2010) [arXiv:0911.4662 [hep-ph]].