

SCALE INVARIANT VOLKOV–AKULOV SUPERGRAVITY

S. FERRARA ^{a,b,c}, M. PORRATI ^{a,d} ¹ AND A. SAGNOTTI ^{a,e} ²^a*Th-Ph Department, CERN
CH - 1211 Geneva 23 SWITZERLAND*^b*INFN - Laboratori Nazionali di Frascati
Via Enrico Fermi 40, 00044 Frascati ITALY*^c*Department of Physics and Astronomy, University of California
Los Angeles, CA 90095-1547 USA*^d*CCPP, Department of Physics, NYU
4 Washington Pl., New York NY 10003, USA*^e*Scuola Normale Superiore and INFN
Piazza dei Cavalieri 7
56126 Pisa ITALY**sergio.ferrara@cern.ch, mp9@nyu.edu, sagnotti@sns.it*

ABSTRACT

A scale invariant Goldstino theory coupled to supergravity is obtained as a standard supergravity dual of a rigidly scale-invariant higher-curvature supergravity with a nilpotent chiral scalar curvature. The bosonic part of this theory describes a massless scalaron and a massive axion in a de Sitter Universe.

¹On sabbatical at CERN-Ph-Th until September 1, 2015.

²On sabbatical at CERN-Ph-Th until September 30, 2015.

1 Introduction

Motivated by single-field inflationary scenarios [1], several sgoldstinoless [2–5] supergravity extensions of inflationary models were recently considered [6–11] (for a recent review see [12]). Interestingly enough, in [7, 11] many of these models were linked to pure higher-derivative supergravity with a nilpotency constraint on the scalar curvature chiral superfield \mathcal{R} . These include the Volkov–Akulov–Starobinsky model [7] and the pure Volkov–Akulov theory coupled to supergravity [7]. Recently, the full component form of the latter theory was presented in [13, 14]

Along these lines, various authors considered R^2 theories of gravity [15] and their supergravity embeddings [15, 16], which possess a rigid scale invariance and naturally accommodate a de Sitter Universe. It is the aim of this note to give the sgoldstinoless version of these theories, which naturally combines an enhanced rigid scale invariance and a de Sitter geometry. This theory also emerges as a limiting case of the inflationary scenario.

2 Scale-Invariant Nilpotent Supergravity

The superspace action density of the scale-invariant theory that we consider ³,

$$\mathcal{A} = \frac{\mathcal{R}\overline{\mathcal{R}}}{g^2}\Big|_D + \sigma \mathcal{R}^2 S_0 \Big|_F, \quad (2.1)$$

where g is a dimensionless parameter, is invariant under the rigid scale transformations

$$\mathcal{R} \rightarrow \mathcal{R}, \quad S_0 \rightarrow e^{-\lambda} S_0, \quad \sigma \rightarrow e^\lambda \sigma. \quad (2.2)$$

This theory is equivalent to the theory considered in [16], supplemented with the nilpotency constraint

$$\mathcal{R}^2 = 0, \quad (2.3)$$

which is enforced by the chiral Lagrange multiplier σ present in the second term of eq. (2.1).

Using manipulations similar to those originally introduced in [17], we can now turn this model into a scale-invariant version of the Volkov–Akulov model coupled to standard supergravity. To this end, we first use the superspace identity

$$\sigma \mathcal{R}^2 S_0 + h.c. \Big|_F = \left(\sigma \frac{\mathcal{R}}{S_0} + \overline{\sigma} \frac{\overline{\mathcal{R}}}{\overline{S_0}} \right) S_0 \overline{S_0} \Big|_D + \text{tot. deriv.}, \quad (2.4)$$

³We use throughout the conventions of [7].

and then introduce two Lagrange chiral superfield multipliers T and S according to

$$\mathcal{A} = \left(\sigma S + \bar{\sigma} \bar{S} + \frac{S \bar{S}}{g^2} \right) S_0 \bar{S}_0 \Big|_D - T \left(\frac{\mathcal{R}}{S_0} - S \right) S_0^3 + \text{h.c.} \Big|_F . \quad (2.5)$$

The final result is the standard supergravity action density

$$\mathcal{A} = - \left(T + \bar{T} - \sigma S - \bar{\sigma} \bar{S} - \frac{S \bar{S}}{g^2} \right) S_0 \bar{S}_0 \Big|_D + T S S_0^3 + \text{h.c.} \Big|_F + \text{tot. deriv.} \quad (2.6)$$

A final shift and a redefinition according to

$$T \rightarrow T + \sigma S , \quad X = \frac{S}{g} \quad (2.7)$$

yield the standard supergravity action density

$$\mathcal{A} = - (T + \bar{T} - X \bar{X}) S_0 \bar{S}_0 \Big|_D + W(T, X) S_0^3 + \text{h.c.} \Big|_F , \quad (2.8)$$

where

$$W(T, X, \sigma) = g T X + g^2 \sigma X^2 . \quad (2.9)$$

This is tantamount to the scale-invariant superpotential

$$W(T, X) = g T X , \quad (2.10)$$

where X is subject to the nilpotency constraint

$$X^2 = 0 , \quad (2.11)$$

so that X describes the sgoldstinoless Volkov–Akulov multiplet [2–5]. The corresponding bosonic Lagrangian,

$$\mathcal{L} = \frac{R}{2} - \frac{3}{(T + \bar{T})^2} |\partial T|^2 - g^2 \frac{|T|^2}{3(T + \bar{T})^2} , \quad (2.12)$$

is a special case of the result displayed in [7], so that it describes an $SU(1,1)/U(1)$ Kählerian model of curvature $-2/3$ with a scale-invariant positive potential. As a result, in terms of the canonical variable

$$T = e^{\phi \sqrt{\frac{2}{3}}} + i a \sqrt{\frac{2}{3}} , \quad (2.13)$$

one finds

$$\mathcal{L} = \frac{R}{2} - \frac{1}{2} (\partial \phi)^2 - \frac{1}{2} e^{-2\phi \sqrt{\frac{2}{3}}} (\partial a)^2 - \frac{g^2}{12} - \frac{g^2}{18} e^{-2\phi \sqrt{\frac{2}{3}}} a^2 . \quad (2.14)$$

Note that in the Einstein frame the metric is inert under the scale transformation corresponding to eq. (2.2), while

$$\phi \rightarrow \phi + \gamma , \quad a \rightarrow e^{\gamma \sqrt{\frac{2}{3}}} a . \quad (2.15)$$

3 de Sitter Vacuum Geometry

Since a is stabilized at zero, this model results in a de Sitter vacuum geometry, with a corresponding scale-invariant realization of supersymmetry breaking induced by the non-linear sgoldstinoless multiplet. The supersymmetry breaking scale M_s^2 is

$$M_s^2 = \frac{g}{2\sqrt{3}} M_{Planck}^2, \quad (3.1)$$

up to a conventional numerical factor. Eq. (2.8) describes the minimal supergravity model that embodies a scale-invariant goldstino interaction and leads unavoidably to a de Sitter geometry. This model involves a single dimensionless parameter g , which determines its *positive* vacuum energy according to

$$V = \frac{g^2}{12} M_{Planck}^4. \quad (3.2)$$

In contrast, the Volkov–Akulov model coupled to supergravity, depends on the two parameters f and W_0 , and consequently leads to a vacuum energy [18] [19] [20] [7] [13]

$$V = \frac{1}{3} |f|^2 - 3 |W_0|^2 \quad (3.3)$$

of arbitrary sign.

Acknowledgments

We would like to thank I. Antoniadis, E. Dudas, R. Kallosh, A. Kehagias, A. Linde and A. Van Proeyen for collaboration on related issues. S.F. is supported in part by INFN (I.S. GSS). M.P. is supported in part by NSF grant PHY–1316452, while A.S. is supported in part by Scuola Normale and by INFN (I.S. Stefi).

References

- [1] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” Phys. Lett. **B 91** (1980) 99; D. Kazanas, “Dynamics of the Universe and Spontaneous Symmetry Breaking,” Astrophys. J. **241** (1980) L59; K. Sato, “Cosmological Baryon Number Domain Structure and the First Order Phase Transition of a Vacuum,” Phys. Lett. B **99** (1981) 66; A. H. Guth, “The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems,” Phys. Rev. **D 23** (1981) 347; V. F. Mukhanov and G. V. Chibisov, “Quantum Fluctuation and Nonsingular Universe. (In Russian),” JETP Lett. **33** (1981) 532 [Pisma Zh.

- Eksp. Teor. Fiz. **33** (1981) 549]; A. D. Linde, “A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems,” Phys. Lett. **B 108** (1982) 389; A. Albrecht and P. J. Steinhardt, “Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking,” Phys. Rev. Lett. **48** (1982) 1220; A. D. Linde, “Chaotic Inflation,” Phys. Lett. **B129** (1983) 177; For recent reviews see: N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, “Non-Gaussianity from inflation: Theory and observations,” Phys. Rept. **402** (2004) 103 [astro-ph/0406398]; V. Mukhanov, “Physical foundations of cosmology,” Cambridge, UK: Univ. Pr. (2005); S. Weinberg, “Cosmology,” Oxford, UK: Oxford Univ. Pr. (2008); D. H. Lyth and A. R. Liddle, “The primordial density perturbation: Cosmology, inflation and the origin of structure,” Cambridge, UK: Cambridge Univ. Pr. (2009); D. S. Gorbunov and V. A. Rubakov, “Introduction to the theory of the early Universe: Cosmological perturbations and inflationary theory,”; J. Martin, C. Ringeval and V. Vennin, “Encyclopaedia Inflationaris,” arXiv:1303.3787 [astro-ph.CO].
- [2] D. V. Volkov and V. P. Akulov, “Is the Neutrino a Goldstone Particle?,” Phys. Lett. B **46** (1973) 109.
- [3] M. Rocek, “Linearizing the Volkov-Akulov Model,” Phys. Rev. Lett. **41** (1978) 451; U. Lindstrom and M. Rocek, “Constrained Local Superfields,” Phys. Rev. D **19** (1979) 2300.
- [4] R. Casalbuoni, S. De Curtis, D. Dominici, F. Feruglio and R. Gatto, “Nonlinear Realization of Supersymmetry Algebra From Supersymmetric Constraint,” Phys. Lett. B **220** (1989) 569.
- [5] Z. Komargodski and N. Seiberg, “From Linear SUSY to Constrained Superfields,” JHEP **0909** (2009) 066 [arXiv:0907.2441 [hep-th]], “Comments on Supercurrent Multiplets, Supersymmetric Field Theories and Supergravity,” JHEP **1007** (2010) 017 [arXiv:1002.2228 [hep-th]].
- [6] L. Alvarez-Gaume, C. Gomez and R. Jimenez, “Minimal Inflation,” Phys. Lett. B **690** (2010) 68 [arXiv:1001.0010 [hep-th]], “A Minimal Inflation Scenario,” JCAP **1103** (2011) 027 [arXiv:1101.4948 [hep-th]].
- [7] I. Antoniadis, E. Dudas, S. Ferrara and A. Sagnotti, “The Volkov-Akulov-Starobinsky Supergravity,” Phys. Lett. B **733** (2014) 32 [arXiv:1403.3269 [hep-th]];

- [8] S. Ferrara, R. Kallosh and A. Linde, “Cosmology with Nilpotent Superfields,” JHEP **1410** (2014) 143 [arXiv:1408.4096 [hep-th]];
- [9] R. Kallosh and A. Linde, “Inflation and Uplifting with Nilpotent Superfields,” JCAP **1501** (2015) 01, 025 [arXiv:1408.5950 [hep-th]];
- [10] G. Dall’Agata and F. Zwirner, “On sgoldstino-less Supergravity models of inflation,” JHEP **1412** (2014) 172 [arXiv:1411.2605 [hep-th]].
- [11] E. Dudas, S. Ferrara, A. Kehagias and A. Sagnotti, “Properties of Nilpotent Supergravity,” arXiv:1507.07842 [hep-th].
- [12] S. Ferrara and A. Sagnotti, “Some Pathways in non-Linear Supersymmetry: Special Geometry Born-Infeld’s, Cosmology and dualities,” arXiv:1506.05730 [hep-th], to appear in a special issue of “p-Adic Numbers, Ultrametric Analysis and Applications” devoted to Prof. V. S. Varadarajan.
- [13] E. A. Bergshoeff, D. Z. Freedman, R. Kallosh and A. Van Proeyen, “Pure de Sitter Supergravity,” arXiv:1507.08264 [hep-th].
- [14] F. Hasegawa and Y. Yamada, “Component action of nilpotent multiplet coupled to matter in 4 dimensional $\mathcal{N} = 1$ supergravity,” arXiv:1507.08619 [hep-th].
- [15] C. Kounnas, D. Lst and N. Toumbas, “ R^2 inflation from scale invariant supergravity and anomaly free superstrings with fluxes,” Fortsch. Phys. **63** (2015) 12 [arXiv:1409.7076 [hep-th]]; L. Alvarez-Gaume, A. Kehagias, C. Kounnas, D. Lust and A. Riotto, “Aspects of Quadratic Gravity,” arXiv:1505.07657 [hep-th].
- [16] S. Ferrara, A. Kehagias and M. Porrati, “ \mathcal{R}^2 Supergravity,” JHEP **1508** (2015) 001 [arXiv:1506.01566 [hep-th]].
- [17] S. Cecotti, “Higher Derivative Supergravity Is Equivalent To Standard Supergravity Coupled To Matter. 1.,” Phys. Lett. B **190** (1987) 86.
- [18] D. Z. Freedman, P. van Nieuwenhuizen and S. Ferrara, “Progress Toward a Theory of Supergravity,” Phys. Rev. D **13** (1976) 3214; S. Deser and B. Zumino, “Consistent Supergravity,” Phys. Lett. B **62** (1976) 335. For a review see: D. Z. Freedman and A. Van Proeyen, “Supergravity,” (Cambridge Univ. Press, 2012).

- [19] S. Deser and B. Zumino, “Broken Supersymmetry and Supergravity,” *Phys. Rev. Lett.* **38** (1977) 1433.
- [20] E. Cremmer, B. Julia, J. Scherk, P. van Nieuwenhuizen, S. Ferrara and L. Girardello, “Superhiggs effect in Supergravity with general scalar interactions,” *Phys. Lett. B* **79** (1978) 231, “Spontaneous Symmetry Breaking and Higgs Effect in Supergravity Without Cosmological Constant,” *Nucl. Phys. B* **147** (1979) 105; E. Cremmer, S. Ferrara, L. Girardello and A. Van Proeyen, “Coupling Supersymmetric Yang-Mills Theories to Supergravity,” *Phys. Lett. B* **116** (1982) 231, “Yang-Mills Theories with Local Supersymmetry: Lagrangian, Transformation Laws and SuperHiggs Effect,” *Nucl. Phys. B* **212** (1983) 413.