


AB

CERN - L3 033

SAC 3138

C2



CERN LIBRARIES, GENEVA



CM-P00065332

Measurement of the Strong Coupling Constant α_s for Bottom Quarks at the Z^0 Resonance

The L3 Collaboration

ABSTRACT

We have measured the ratio of the strong coupling constants α_s for bottom quarks and light quarks at the Z^0 resonance, in order to test the flavour independence of the strong interaction. The coupling strength α_s has been determined from the fraction of events with three jets, measured for a sample of all hadronic events, and for inclusive muon and electron events. The b purity is evaluated to be 22% for the first data set and 87% for the inclusive lepton sample. We find $\alpha_s(b)/\alpha_s(u\bar{d} s\bar{c}) = 1.00 \pm 0.05(\text{stat.}) \pm 0.06(\text{syst.})$.

Introduction

Quantum Chromodynamics, the theory of strong interactions, predicts the coupling constant α_s to be independent of the quark flavour.

For the light quark species, up, down and strange, this prediction is supported by the approximate isospin and SU(3) flavour symmetries. The relative coupling strengths for charm and bottom quarks have been measured in e^+e^- annihilation at center of mass energies around 30 GeV [1,2], confirming the flavour independence of α_s within large uncertainties. Also the comparison of α_s values measured in charmonium and bottomonium decays [3] with those obtained for other processes involving only u,d and s quarks supports the flavour independence of the strong interaction. However, since these comparisons involve different energy scales and different systematic errors, the accuracy of such tests is limited. The flavour composition in hadronic events produced in e^+e^- collisions at the Z^0 pole is different from that at lower center of mass energies. By comparing the corresponding 3-jet fractions one finds that the strong coupling constants for quarks with different electric charge (u,c and d,s,b) agree within 10% [4]. Here we present a precise measurement of the relative coupling strength for bottom quarks at the Z^0 resonance using the L3 detector at LEP.

The measured fraction of b events in hadronic decays of the Z^0 boson is in agreement with the Standard Model prediction of 22% [5,6]. Bottom quarks can be tagged via their semileptonic decays $b \rightarrow l + X$: because of the large mass of the b quark, the leptons have large momentum p and large transverse momentum p_\perp with respect to the nearest jet. By selecting hadronic events containing electrons and muons with large values of p and p_\perp we obtain a sample of events with a high b purity.

As the 3-jet rate ($\sigma_{3\text{-jet}}/\sigma_{\text{tot}}$) is a measure of α_s , the comparison of the 3-jet fractions for event samples with different flavour composition provides a test of flavour independence. In this letter we determine the ratio of the 3-jet fraction for inclusive lepton events and of the 3-jet fraction for a sample of all hadronic events. In this double ratio, which is sensitive to the ratio of the coupling constants $\alpha_s(b)/\alpha_s(u\bar{d}sc)$, most systematic uncertainties cancel.

The analysis is based on the events collected in 1990 ($\simeq 5.5 \text{ pb}^{-1}$ of integrated luminosity) at center of mass energies in the range $88.2 < \sqrt{s} < 94.2 \text{ GeV}$.

The L3 Detector

The L3 detector covers 99% of 4π . The detector includes a central vertex chamber, a precise electromagnetic calorimeter composed of BGO crystals, a ring of scintillation counters, a uranium and brass hadron calorimeter with proportional wire chamber read-out, and a high accuracy muon chamber system. These detectors are installed in a magnet with an inner diameter of 12 m, which provides a uniform field of 0.5 T along the beam direction.

The central tracking chamber is a time expansion chamber consisting of two cylindrical layers of 12 and 24 sectors, with 62 wires measuring the R- ϕ coordinate. The single wire

resolution is 58 μm averaged over the entire cell. The double track resolution is 640 μm . The fine segmentation of the electromagnetic detector and the hadron calorimeter allows us to measure the axis of jets with an angular resolution of about 2.5°, and to measure the total energy of hadronic events from Z^0 decay with a resolution of 10% [7]. The muon detector consists of three layers of precise drift chambers. A detailed description of each detector subsystem, and its performance, is given in reference [8].

For the present analysis, we use the data collected in the following ranges of polar angles:

- for the central chamber, $41^\circ < \theta < 139^\circ$,
- for the electromagnetic calorimeter, $42^\circ < \theta < 138^\circ$,
- for the hadron calorimeter, $5^\circ < \theta < 175^\circ$,
- for the muon chambers, $36^\circ < \theta < 144^\circ$.

Event Selection

The primary trigger for hadronic events requires a total energy above 15 GeV in the calorimeters, while the trigger for muons requires the signal of one of the 16 scintillation counter ϕ sectors to be in coincidence with a track in the muon chambers. The two triggers are in a logical OR with a charged track trigger and a scintillation counter multiplicity trigger. The combined trigger efficiency exceeds 99.9% for hadronic events, with or without leptons.

The selection of $e^+e^- \rightarrow \text{hadrons}$ events is based on the energy measured in the electromagnetic and hadronic calorimeters:

$$\begin{aligned} 0.6 < E_{\text{vis}}/\sqrt{s} &< 1.4, \\ |E_{||}|/E_{\text{vis}} &< 0.40, \quad E_{\perp}/E_{\text{vis}} < 0.50, \\ N_{\text{cluster}} &> 12. \end{aligned}$$

E_{vis} is the total energy observed in the detector, $E_{||}$ and E_{\perp} are the parallel and transverse energy imbalances with respect to the beam direction. N_{cluster} is the number of clusters, groups of neighbouring hits in the calorimeters with an energy exceeding 100 MeV. The algorithm normally reconstructs one cluster for each particle produced near the interaction point. The cut on the number of clusters rejects low multiplicity events (e^+e^- , $\mu^+\mu^-$, $\tau^+\tau^-$). Applying the same cuts to simulated events, we estimate the acceptance to be 97%.

Events of the type $Z^0 \rightarrow b\bar{b}$ are selected identifying the muons or electrons coming from the semileptonic decay of the b quark.

Muons are identified and measured in the muon chamber system. A muon track must consist of track segments in at least two of the three layers of the muon spectrometer. To

reject the punch-through background, we require that the reconstructed track points to the interaction region.

Electrons are identified using the BGO, the central tracking chamber and the hadron calorimeter. We require the lateral shower shape of the cluster in the BGO to be consistent with that of an electromagnetic shower. The energy in the hadron calorimeter behind this cluster must be small, and the centroid of the cluster must be matched to a track in the vertex chamber. The selection for both inclusive muon and electron events is described in detail in reference [5].

Leptons coming from b semi-leptonic decays have a large momentum and a large transverse momentum with respect to the b quark direction. Therefore we require the muons and electrons to have a momentum greater than 4 GeV and 3 GeV respectively, and a transverse momentum with respect to the axis of the nearest jet greater than 1.5 GeV for muons and 1 GeV for electrons. For this selection jets are reconstructed using the algorithm described in reference [7]: the measured momentum of the lepton is excluded in the calculation of the jet direction. From 110,000 hadronic events we select approximately 1800 muon and 1100 electron events.

Monte Carlo distributions were generated by the program JETSET 7.2 [9] with parton shower and string fragmentation. For the heavy quarks we use the Peterson fragmentation function [10] with parameters adjusted to match our measured inclusive muon data [5]. The generated events were passed through the L3 detector simulation [11] which includes the effects of energy loss, multiple scattering, interactions and decays in the detector materials and beam pipe. From Monte Carlo studies we estimate the b purity to be 86% for the muon sample and 88% for the electron sample. The error on these numbers is estimated to be $\pm 5\%$ [5].

Determination of the 3-Jet Rate

In order to determine the number of jets in the selected samples of hadronic and inclusive lepton events, we use the 'JADE' algorithm [12]. For each pair of 'particles' (calorimetric clusters and muons) i and j we calculate the scaled invariant mass squared $y_{ij} = 2E_i E_j / E_{\text{vis}}^2 \cdot (1 - \cos \theta_{ij})$, where E_i and E_j are the particle energies and θ_{ij} is the angle between particles i and j . The pair for which y_{ij} is smallest is replaced by a pseudoparticle k with four-momentum $p_k = p_i + p_j$. This procedure is repeated until all y_{ij} exceed the jet resolution parameter y_{cut} . The remaining pseudoparticles are called jets.

We determine the 3-jet rate

$$f_3 = N_{\text{3-jet}} / N_{\text{tot}}$$

(where N is the number of events) at nine equidistant values of y_{cut} ranging from 0.02 to 0.10, for the total hadronic sample

$$f_3(\text{had}) = N_{\text{3-jet(had)}} / N_{\text{tot(had)}}$$

and for the inclusive muon and electron samples separately

$$f_3(\mu) = N_{\text{3-jet(incl. muons)}} / N_{\text{tot(incl. muons)}},$$

$$f_3(e) = N_{3\text{-jet}}(\text{incl. electrons})/N_{\text{tot}}(\text{incl. electrons}) .$$

From these quantities we compute the ratios $f_3(\mu)/f_3(\text{had})$ and $f_3(e)/f_3(\text{had})$ for each value of y_{cut} .

In order to correct the two ratios we use the Monte Carlo JETSET 7.2. The corrected ratios are obtained by multiplying with a y_{cut} dependent factor

$$C = C_d \cdot C_h \cdot C_b .$$

The factors C_d take into account the detector effects and are evaluated from the parton shower Monte Carlo before and after including detector resolution and acceptance: they range from 0.90 to 1. The factors C_h are due to fragmentation effects and are given by the ratio of $f_3(e, \mu)/f_3(\text{had})$ before and after hadronization: this gives a correction of typically 3%. The numbers C_b have been evaluated using the first order QCD matrix element, including mass effects, assuming a b quark mass of 5 GeV [13]: the correction is around +2%.

The systematic uncertainty in the correction factor C_d has been evaluated by changing the energy response in different components of the calorimeter up to 10% in the Monte Carlo simulation. We find a systematic uncertainty of 3%.

To study the systematic uncertainty in the hadronization correction, we vary the ϵ_b parameter of the Peterson fragmentation function of the b quark, and the fragmentation parameters σ_q and b , in a range compatible with our measured inclusive lepton spectra and event shape distributions [5,14]. We find a systematic uncertainty of 2% due to hadronization.

y_{cut}	$f_3(\mu)/f_3(\text{had})$	$f_3(e)/f_3(\text{had})$
0.02	$0.94 \pm 0.03 \pm 0.05$	$0.94 \pm 0.04 \pm 0.05$
0.03	$0.98 \pm 0.04 \pm 0.05$	$0.99 \pm 0.05 \pm 0.06$
0.04	$0.99 \pm 0.04 \pm 0.05$	$1.00 \pm 0.05 \pm 0.06$
0.05	$0.99 \pm 0.04 \pm 0.05$	$1.02 \pm 0.06 \pm 0.07$
0.06	$1.00 \pm 0.05 \pm 0.06$	$1.02 \pm 0.07 \pm 0.07$
0.07	$0.97 \pm 0.05 \pm 0.06$	$1.03 \pm 0.07 \pm 0.08$
0.08	$0.95 \pm 0.06 \pm 0.06$	$1.02 \pm 0.08 \pm 0.08$
0.09	$0.92 \pm 0.06 \pm 0.07$	$1.02 \pm 0.09 \pm 0.09$
0.10	$1.00 \pm 0.07 \pm 0.07$	$1.04 \pm 0.09 \pm 0.09$

Table 1: Measured 3-jet rates for the inclusive muon and the inclusive electron samples for the nine different values of y_{cut} . The ratios have been corrected for detector acceptance and resolution, for hadronization and bottom mass effect. The first error is statistical, the second is systematic.

Table 1 shows the ratios $f_3(\mu)/f_3(\text{had})$ and $f_3(e)/f_3(\text{had})$ with the statistical and systematic errors, for the values of y_{cut} considered. The systematic error also includes the Monte Carlo statistical error, which ranges from 2% to 5%. There is good agreement

between μ and e data, and the results are independent of the jet resolution parameter y_{cut} .

Results and Conclusions

The two ratios for muons and electrons are finally combined in order to calculate the ratio between the 3-jet fraction for bottom events and for all other flavours. Assuming that there is no difference in the strong interaction for the quarks u, d, s and c, we can write

$$f_3(e + \mu)/f_3(\text{had}) = \frac{(1 - p^{e+\mu}) \cdot f_3^{\text{udsc}} + p^{e+\mu} \cdot f_3^b}{(1 - p^{\text{had}}) \cdot f_3^{\text{udsc}} + p^{\text{had}} \cdot f_3^b}$$

where $p^{e+\mu}$ is the combined b purity for the two samples of inclusive lepton events ($87 \pm 5\%$) and p^{had} is the percentage of b in hadronic events ($22 \pm 0.5\%$).

In first order QCD, the three jet rate is proportional to α_s :

$$f_3 = A(y_{\text{cut}}) \cdot \alpha_s.$$

The function $A(y_{\text{cut}})$ has been calculated in reference [15]. From the ratio $f_3(b)/f_3(\text{udsc})$ we can then determine the relative coupling strength for b and the other quarks: figure 1 shows the ratio $\alpha_s(b)/\alpha_s(\text{udsc})$ as a function of y_{cut} .

Taking into account second order corrections to the relation between the 3-jet rate and α_s [15], we find that the result changes by less than 1%. For these corrections we used a value for $\alpha_s(M_Z) = 0.115$ as determined from our previous analysis [16], and we fix the renormalization scale μ^2 to $y_{\text{cut}} \cdot s$.

For a value of $y_{\text{cut}} = 0.05$, for which the 4-jet rate is small and the 3-jet fraction is large (29%), we measure $f_3(e + \mu)/f_3(\text{had}) = 1.00 \pm 0.03(\text{stat.}) \pm 0.04(\text{syst.})$, from which we obtain

$$\alpha_s(b)/\alpha_s(\text{udsc}) = 1.00 \pm 0.05(\text{stat.}) \pm 0.06(\text{syst.})$$

where the systematic error includes also the small contributions from the errors on the $p^{e+\mu}$ and p^{had} fractions. This result is consistent with the flavour independence predicted by QCD. The precision is significantly better than that previously achieved [2].

Acknowledgments

We wish to express our gratitude to the CERN accelerator divisions for the excellent performance of the LEP machine. We acknowledge the effort of all engineers and technicians who have participated in the construction and maintenance of this experiment.

The L3 Collaboration:

B.Adeva,¹⁵ O.Adriani,¹³ M.Aguilar-Benitez,²³ H.Akbari,⁵ J.Alcaraz,²³ A.Aloisio,²⁵ G.Alverson,⁹
M.G.Alvaggi,²⁵ G.Ambrosi,²⁹ Q.An,¹⁶ H.Anderhub,⁴¹ A.L.Anderson,¹² V.P.Andreev,¹⁴ T.Angelov,¹²
L.Antonov,³⁶ D.Antreasyan,⁷ P.Arce,²³ A.Arefiev,²⁴ T.Azemoon,³ T.Aziz,⁸ P.V.K.S.Baba,¹⁶ P.Bagnaia,³²
J.A.Bakken,³¹ L.Baksay,³⁷ R.C.Ball,³ S.Banerjee,⁸ J.Bao,⁵ R.Barillère,¹⁵ L.Barone,³² R.Battiston,²⁹
A.Bay,¹⁷ U.Becker,¹² F.Behner,⁴¹ J.Behrens,⁴¹ S.Beingessner,⁴ Gy.L.Bencze,^{10,15} J.Berdugo,²³ P.Berges,¹²
B.Bertucci,²⁹ B.L.Betev,³⁶ A.Biland,⁴¹ G.M.Bilei,²⁹ R.Bizzarri,³² J.J.Blaising,⁴ P.Blömeke,¹ B.Blumenfeld,⁵
G.J.Bobbinck,² M.Boccilini,¹³ R.Bock,¹ A.Böhm,^{1,15} B.Borgia,³² D.Bourilkov,³⁶ M.Bourquin,¹⁷ D.Boutigny,⁴
B.Bouwens,² E.Brambilla,²⁵ J.G.Branson,³³ I.C.Brock,³⁰ F.Bruyant,¹⁵ C.Buisson,²² A.Bujak,³⁸ J.D.Burger,¹²
J.P.Burq,²² J.Busenitz,³⁷ X.D.Cai,¹⁶ M.Capell,²⁰ M.Caria,²⁹ F.Carminati,¹³ A.M.Cartacci,¹³ M.Cerrada,²³
F.Cesaroni,³² Y.H.Chang,¹² U.K.Chaturvedi,¹⁶ M.Chemarin,²² A.Chen,⁴³ C.Chen,⁶ G.M.Chen,⁶ H.F.Chen,¹⁸
H.S.Chen,⁶ M.Chen,¹² M.L.Chen,³ W.Y.Chen,¹⁶ G.Chiefari,²⁵ C.Y.Chien,⁵ M.Chmeissani,³ C.Civinini,¹³
I.Clare,¹² R.Clare,¹² H.O.Cohn,²⁷ G.Coignet,⁴ N.Colino,¹⁵ V.Commichau,¹ G.Conforto,¹³ A.Contin,^{7,15}
F.Crijns,²⁶ X.Y.Cui,¹⁶ T.S.Dai,¹² R.D'Alessandro,¹³ R.de Asmundis,²⁵ A.Degré,^{15,4} K.Deiters,¹²
E.Dénes,^{10,15} P.Denes,³¹ F.DeNotaristefani,³² M.Dhina,⁴¹ D.DiBitonto,³⁷ M.Diemoz,³² H.R.Dimitrov,³⁶
C.Dionisi,³² M.T.Dova,¹⁶ E.Drago,²⁵ T.Driever,²⁶ D.Duchesneau,¹⁷ P.Duinker,² I.Duran,²³ H.El Mamouni,²²
A.Engler,³⁰ F.J.Eppeling,¹² F.C.Erné,² P.Extermann,¹⁷ R.Fabbretti,³⁹ M.Fabre,⁴¹ S.Falciano,³² Q.Fan,¹⁶
S.J.Fan,³⁵ O.Fackler,²⁰ J.Fay,²² T.Ferguson,³⁰ G.Fernandez,²³ F.Ferroni,^{32,15} H.Fesefeldt,¹ E.Fiandrini,²⁹
J.Field,¹⁷ F.Filthaut,²⁶ G.Finocchiaro,³² P.H.Fisher,⁵ G.Forconi,¹⁷ T.Foreman,² K.Freudenreich,⁴¹
W.Friebel,⁴⁰ M.Fukushima,¹² M.Gaillard,¹⁹ Yu.Galaktionov,²⁴ E.Gallo,¹³ S.N.Ganguli,⁸ P.Garcia-Abia,²³
S.S.Gau,⁴³ D.Gele,²² S.Gentile,³² M.Glaubman,⁹ S.Goldfarb,³ Z.F.Gong,¹⁸ E.Gonzalez,²³ A.Gordeev,²⁴
P.Göttlicher,¹ D.Goujon,¹⁷ G.Gratta,²⁸ C.Grinnell,¹² M.Gruenewald,²⁸ M.Guanzioli,¹⁶ J.K.Guo,³⁵ A.Gurtu,⁸
H.R.Gustafson,³ L.J.Gutay,³⁸ H.Haan,¹ A.Hasan,¹⁶ D.Hauschildt,² C.F.He,³⁵ T.Hebbeker,¹ M.Hebert,³³
G.Herten,¹² U.Herten,¹ A.Hervé,¹⁵ K.Hilgers,¹ H.Hofer,⁴¹ H.Hoorani,¹⁶ L.S.Hsu,⁴³ G.Hu,¹⁶ G.Q.Hu,³⁵ B.Ille,²²
M.M.Ilyas,¹⁶ V.Innocente,^{25,15} H.Janssen,¹⁵ S.Jezequel,⁴ B.N.Jin,⁶ L.W.Jones,³ A.Kasser,¹⁹ R.A.Khan,¹⁶
Yu.Kamyshkov,^{24,27} Y.Karyotakis,^{4,15} M.Kaur,¹⁶ S.Khokhar,¹⁶ V.Khoze,¹⁴ M.N.Kienzle-Focacci,¹⁷
W.Kinnison,²¹ D.Kirkby,²⁸ W.Kittel,²⁶ A.Klimentov,²⁴ A.C.König,²⁶ O.Kornadt,¹ V.Koutsenko,^{24,12}
R.W.Kraemer,³⁰ T.Kramer,¹² V.R.Krastev,³⁶ W.Krenz,¹ J.Krizmanic,⁵ K.S.Kumar,¹¹ V.Kumar,¹⁶
A.Kunin,^{11,24} V.Lalieu,¹⁷ G.Landi,¹³ K.Lanius,¹⁵ D.Lanske,¹ S.Lanzano,²⁵ P.Lebrun,²² P.Lecomte,⁴¹
P.Lecoq,¹⁵ P.Le Coultre,⁴¹ D.Lee,²¹ I.Leedom,⁹ J.M.Le Goff,¹⁵ L.Leistam,¹⁵ R.Leiste,⁴⁰ M.Lenti,¹³
E.Leonardi,³² J.Lettry,⁴¹ P.M.Levchenko,¹⁴ X.Leytens,² C.Li,^{18,16} H.T.Li,⁶ J.F.Li,¹⁶ L.Li,³⁹ P.J.Li,³⁵ Q.Li,¹⁶
X.G.Li,⁶ J.Y.Liao,³⁵ Z.Y.Lin,¹⁸ F.L.Linde,^{15,2} B.Lindemann,¹ D.Linnhofer,⁴¹ R.Liu,¹⁶ Y.Liu,¹⁶ W.Lohmann,⁴⁰
E.Longo,³² Y.S.Lu,⁶ J.M.Lubbers,¹⁵ K.Lübelsmeyer,¹ C.Luci,¹⁵ D.Luckey,^{7,12} L.Ludovici,³² L.Luminari,³²
W.G.Ma,¹⁸ M.MacDermott,⁴¹ R.Magahiz,³⁴ P.K.Malhotra,⁸ R.Malik,¹⁶ A.Malinin,^{24,4} C.Maña,²³ D.N.Mao,³
Y.F.Mao,⁶ M.Maolinbay,⁴¹ P.Marchesini,⁴¹ A.Marchionni,¹³ J.P.Martin,²² L.Martinez-Laso,¹⁵ F.Marzano,³²
G.G.G.Massaro,² T.Matsuda,¹² K.Mazumdar,⁸ P.McBride,¹¹ T.McMahon,³⁸ D.McNally,⁴¹ Th.Meinholtz,¹
M.Merk,²⁶ L.Merola,²⁵ M.Meschini,¹³ W.J.Metzger,²⁶ Y.Mi,¹⁶ G.B.Mills,²¹ Y.Mir,¹⁶ G.Mirabelli,³²
J.Mnich,¹ M.Möller,¹ B.Monteleoni,¹³ G.Morand,¹⁷ R.Morand,⁴ S.Morganti,³² N.E.Moulai,¹⁶ R.Mount,²⁸
S.Müller,¹ E.Nagy,¹⁰ M.Napolitano,²⁵ H.Newman,²⁸ C.Neyer,⁴¹ M.A.Niaz,¹⁶ L.Niessen,¹ H.Nowak,⁴⁰
D.Pandoulas,¹ M.Pauluzzi,²⁹ F.Pauss,⁴¹ F.Plasil,²⁷ G.Passaleva,¹³ S.Patricelli,²⁵ Y.J.Pei,¹ D.Perret-Gallix,⁴
J.Perrier,¹⁷ A.Pevsner,⁵ M.Pieri,¹³ P.A.Piroué,³¹ V.Plyaskin,²⁴ M.Pohl,⁴¹ V.Pojidaev,²⁴ N.Produit,¹⁷
J.M.Qian,³ K.N.Qureshi,¹⁶ R.Raghavan,⁸ G.Rahal-Callot,⁴¹ G.Raven,² P.Razis,³⁷ K.Read,²⁷ D.Ren,⁴¹
Z.Ren,¹⁶ S.Reucroft,⁹ A.Ricker,¹ S.Riemann,⁴⁰ O.Rind,³ C.Rippich,³⁰ H.A.Rizvi,¹⁶ B.P.Roe,³ M.Röhner,¹
S.Röhner,¹ L.Romero,²³ J.Rose,¹ S.Rosier-Lees,⁴ R.Rosmalen,²⁶ Ph.Rosselet,¹⁹ A.Rubbia,¹² J.A.Rubio,^{15,23}
W.Ruckstuhl,¹⁷ H.Rykaczewski,⁴¹ M.Sachwitz,^{40,15} J.Salicio,^{15,23} J.M.Salicio,²³ G.Sanders,²¹ A.Santocchia,²⁹
M.S.Sarakinos,¹² G.Sartorelli,^{7,16} G.Sauvage,⁴ A.Savin,²⁴ V.Schegelsky,¹⁴ K.Schmiemann,¹ D.Schmitz,¹
P.Schmitz,¹ M.Schneegans,⁴ H.Schopper,⁴² D.J.Schotanus,²⁶ S.Shotkin,¹² H.J.Schreiber,⁴⁰ R.Schulte,¹
S.Schulte,¹ K.Schultze,¹ J.Schütte,¹¹ J.Schwenke,¹ G.Schwering,¹ C.Sciacca,²⁵ I.Scott,¹¹ R.Sehgal,¹⁶
P.G.Seiler,³⁹ J.C.Sens,² L.Servoli,²⁹ I.Sheer,³³ D.Z.Shen,³⁵ V.Shevchenko,²⁴ S.Shevchenko,²⁴ X.R.Shi,²⁸
K.Shmakov,²⁴ V.Shoutko,²⁴ E.Shumilov,²⁴ N.Smirnov,¹⁴ E.Soderstrom,³¹ A.Sopczak,³³ C.Spartiotis,⁵
T.Spickermann,¹ P.Spillantini,¹³ R.Starosta,¹ M.Steuer,^{7,12} D.P.Stickland,³¹ F.Sticcozzi,¹² W.Stoeffl,²⁰
H.Stone,¹⁷ K.Strauch,¹¹ B.C.Stringfellow,³⁸ K.Sudhakar,^{8,1} G.Sultanov,¹⁶ R.L.Sumner,³¹ L.Z.Sun,^{18,16}
H.Suter,⁴¹ R.B.Sutton,³⁰ J.D.Swain,¹⁶ A.A.Syed,¹⁶ X.W.Tang,⁶ E.Tarkovsky,²⁴ L.Taylor,⁹ C.Timmermans,²⁶
Samuel C.C.Ting,¹² S.M.Ting,¹² Y.P.Tong,⁴³ F.Tonisch,⁴⁰ M.Tonutti,¹ S.C.Tonwar,⁸ J.Tóth,^{10,15}
G.Trowitzsch,⁴⁰ C.Tully,²⁸ K.L.Tung,⁶ J.Ulbricht,⁴¹ L.Urbán,¹⁰ U.Uwer,¹ E.Valente,³² R.T.Van de Walle,²⁶
I.Vetlitsky,²⁴ G.Viertel,⁴¹ P.Vikas,¹⁶ U.Vikas,¹⁶ M.Vivargent,^{4,12} H.Vogel,³⁰ H.Vogt,⁴⁰ G.Von Dardel,¹⁵
I.Vorobiev,²⁴ A.A.Vorobyov,¹⁴ An.A.Vorobyov,¹⁴ L.Vuilleumier,¹⁹ M.Wadhwa,¹⁶ W.Wallraff,¹ C.R.Wang,¹⁸
G.H.Wang,³⁰ J.H.Wang,⁶ Q.F.Wang,¹¹ X.L.Wang,¹⁸ Y.F.Wang,¹³ Z.Wang,¹⁶ Z.M.Wang,^{16,18} A.Weber,¹

J.Weber⁴¹ R.Weill¹⁹ T.J.Wenaus²⁰ J.Wenninger¹⁷ M.White¹² C.Willmott²³ F.Wittgenstein¹⁵
D.Wright³¹ R.J.Wu⁶ S.L.Wu¹⁶ S.X.Wu¹⁶ Y.G.Wu⁶ B.Wyslouch¹² Y.Y.Xie³⁵ Y.D.Xu⁶ Z.Z.Xu¹⁸
Z.L.Xue³⁵ D.S.Yan³⁵ X.J.Yan¹² B.Z.Yang¹⁸ C.G.Yang⁶ G.Yang¹⁶ K.S.Yang⁶ Q.Y.Yang⁶ Z.Q.Yang³⁵
C.H.Ye¹⁶ J.B.Ye^{41,18} Q.Ye¹⁶ S.C.Yeh⁴³ Z.W.Yin³⁵ J.M.You¹⁶ M.Yzerman² C.Zaccardelli²⁸ P.Zemp⁴¹
M.Zeng¹⁶ Y.Zeng¹ D.H.Zhang² Z.P.Zhang^{18,16} J.F.Zhou¹ R.Y.Zhu²⁸ H.L.Zhuang⁶ A.Zichichi^{7,15,16}

- 1 I. Physikalisches Institut, RWTH, Aachen, Federal Republic of Germany[§]
III. Physikalisches Institut, RWTH, Aachen, Federal Republic of Germany[§]
2 National Institute for High Energy Physics, NIKHEF, Amsterdam, The Netherlands
3 University of Michigan, Ann Arbor, Michigan, United States of America
4 Laboratoire de Physique des Particules, LAPP, Annecy, France
5 Johns Hopkins University, Baltimore, Maryland, United States of America
6 Institute of High Energy Physics, IHEP, Beijing, China
7 INFN-Sezione di Bologna, Italy
8 Tata Institute of Fundamental Research, Bombay, India
9 Northeastern University, Boston, Massachusetts, United States of America
10 Central Research Institute for Physics of the Hungarian Academy of Sciences, Budapest, Hungary
11 Harvard University, Cambridge, Massachusetts, United States of America
12 Massachusetts Institute of Technology, Cambridge, Massachusetts, United States of America
13 INFN Sezione di Firenze and University of Firenze, Italy
14 Leningrad Nuclear Physics Institute, Gatchina, Soviet Union
15 European Laboratory for Particle Physics, CERN, Geneva, Switzerland
16 World Laboratory, FBLJA Project, Geneva, Switzerland
17 University of Geneva, Geneva, Switzerland
18 University of Science and Technology of China, Hefei, China
19 University of Lausanne, Lausanne, Switzerland
20 Lawrence Livermore National Laboratory, Livermore, California, United States of America
21 Los Alamos National Laboratory, Los Alamos, New Mexico, United States of America
22 Institut de Physique Nucléaire de Lyon, IN2P3-CNRS/Université Claude Bernard, Villeurbanne, France
23 Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas, CIEMAT, Madrid, Spain
24 Institute of Theoretical and Experimental Physics, ITEP, Moscow, Soviet Union
25 INFN-Sezione di Napoli and University of Naples, Italy
26 University of Nymegen and NIKHEF, Nymegen, The Netherlands
27 Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States of America
28 California Institute of Technology, Pasadena, California, United States of America
29 INFN-Sezione di Perugia and Università Degli Studi di Perugia, Perugia, Italy
30 Carnegie Mellon University, Pittsburgh, Pennsylvania, United States of America
31 Princeton University, Princeton, New Jersey, United States of America
32 INFN-Sezione di Roma and University of Roma, "La Sapienza", Italy
33 University of California, San Diego, California, United States of America
34 Union College, Schenectady, New York, United States of America
35 Shanghai Institute of Ceramics, SIC, Shanghai, China
36 Central Laboratory of Automation and Instrumentation, CLANP, Sofia, Bulgaria
37 University of Alabama, Tuscaloosa, Alabama, United States of America
38 Purdue University, West Lafayette, Indiana, United States of America
39 Paul Scherrer Institut, PSI, Würenlingen, Switzerland
40 Institut für Hochenergiephysik, Zeuthen, Federal Republic of Germany[§]
41 Eidgenössische Technische Hochschule, ETH Zürich Switzerland
42 University of Hamburg, Federal Republic of Germany
43 High Energy Physics Group, Taiwan, China

[§] Supported by the German Bundesministerium für Forschung und Technologie

References

1. TASSO Collaboration, W. Braunschweig et al., *Z. Phys. C* **44** (1989) 365.
2. TASSO Collaboration, W. Braunschweig et al., *Z. Phys. C* **42** (1989) 17.
3. W. Kwong et al., *Phys. Rev. D* **37** (1988) 3210.
4. T. Hebbeker, "QCD at LEP/SLC", Aachen Report PITHA 91/11.
5. L3 Collaboration, B. Adeva et al., *Phys. Lett. B* **261** (1991) 177;
L3 Collaboration, B. Adeva et al., *Phys. Lett. B* **252** (1990) 713.
6. L3 Collaboration, B. Adeva et al., L3 preprint #28, February 1991.
7. O. Adriani et al., *Nucl. Instr. and Methods A* **302** (1991) 53.
8. L3 Collaboration, B. Adeva et al., *Nucl. Instrum. Methods A* **289** (1990) 35.
9. T. Sjöstrand, *Comput. Phys. Commun.* **39** (1986) 347;
T. Sjöstrand and M. Bengtsson, *Comput. Phys. Commun.* **43** (1987) 367.
10. C. Peterson et al., *Phys. Rev. D* **27** (1983) 105.
11. GEANT version 3.13, September, 1989. See R. Brun et al., "GEANT 3", CERN DD/EE/84-1 (Revised), Sept. 1987. Hadronic interactions are simulated using the GHEISHA program. See H. Fesefeldt RWTH Aachen Report PITHA 85/02 (1985).
12. JADE Collaboration, W. Bartel et al. *Z. Phys. C* **33** (1986) 23;
JADE Collaboration, S. Bethke et al. *Phys. Lett. B* **213** (1988) 235.
13. B.L. Ioffe, *Phys. Lett. B* **78** (1978) 277.
14. L3 Collaboration, B. Adeva et al., *Phys. Lett. B* **257** (1991) 469.
15. Z. Kunszt and P. Nason in "Z Physics at LEP 1", CERN Report CERN-89-08, Vol. I, p. 373.
16. L3 Collaboration, B. Adeva et al., *Phys. Lett. B* **248** (1990) 464.




Figure 1

Measured $\alpha_s(b)/\alpha_s(\text{udsc})$ ratios, for the inclusive lepton sample (muons and electrons), as a function of the jet resolution parameter y_{cut} . The error includes the statistical and the systematic contributions.