


The Compact Muon Solenoid Experiment
Conference Report

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland



11 April 2017 (v3, 24 April 2017)

Charge asymmetry dependence of elliptic and triangular flow in pPb and PbPb collisions with CMS

Sang Eon Park for the CMS Collaboration

Abstract

In nucleus-nucleus collisions, the linear dependence found for the elliptic flow harmonic of both positive or negative charged particles as a function of event charge asymmetry ($A_{ch} = (N^+ - N^-)/(N^+ + N^-)$, where N^+ and N^- are the number of positive and negative charged particles, respectively) is predicted by the phenomenon known as the Chiral Magnetic Wave (CMW) due to its induced electric quadrupole moment. However, other scenarios are also possible and may provide alternative explanations for the experimental results. New measurements of elliptic (v_2) and triangular (v_3) flow for positive and negative charged particles as a function of A_{ch} in pPb and PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV are presented, using data collected by the CMS experiment during the LHC runs 1 and 2. The slopes and intercepts of the charged-dependent v_n harmonics vs. A_{ch} are directly compared for pPb and PbPb collisions with similar charged-particle multiplicities, where a strong CMW effect is not expected in very high multiplicity pPb events. Moreover, a comparison is made of the slope parameters between v_2 and v_3 harmonics normalized by the inclusive charge particle v_n in PbPb collisions as a function of centrality. These results provide a means to discriminate between the CMW and other scenarios such as local charge conservation as possible explanations for the observed charge dependent behavior.

Presented at *QM2017 Quark Matter 2017, XXVI international conference on ultrarelativistic heavy-ion collisions*



Charge asymmetry dependence of anisotropic flow in pPb and PbPb collisions with CMS experiment

Sang Eon Park on behalf of the CMS Collaboration

Physics & Astronomy MS-61, Rice University, Houston, US, 77005

Abstract

In nucleus-nucleus collisions, the linear dependence found for the elliptic flow harmonic of both positive or negative charged particles as a function of event charge asymmetry is predicted by the phenomenon known as the Chiral Magnetic Wave (CMW) due to its induced electric quadrupole moment. Here, the event charge asymmetry A_{ch} is defined as $\frac{N_+ - N_-}{N_+ + N_-}$, where N_+ and N_- are the number of positive and negative charged particles, respectively. However, other scenarios are also possible and may provide alternative explanations for the experimental results. New measurements of elliptic (v_2) and triangular (v_3) flow for positive and negative charged particles as a function of A_{ch} in pPb and PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV are presented, using data collected by the CMS experiment during the LHC runs 1 and 2. The slopes and intercepts of the charged-dependent v_n harmonics vs. A_{ch} are directly compared for pPb and PbPb collisions with similar charged-particle multiplicities, where a strong CMW effect is not expected in very high multiplicity pPb events. Moreover, a comparison is made of the slope parameters between v_2 and v_3 harmonics normalized by the inclusive charge particle v_n in PbPb collisions as a function of centrality. These results provide a means to discriminate between the CMW and other scenarios such as local charge conservation as possible explanations for the observed charge dependent behavior.

Keywords: CMW, Chiral Magnetic Wave, LCC, local charge conservation, small systems, chiral magnetic effect

1. Introduction

In relativistic nucleus-nucleus collisions, the nonzero axial chemical potential, induced by the chiral anomaly with an imbalance of left- and right-handed quarks, can produce an electric current along the magnetic field that is produced by the spectator protons [1, 2, 3]. The phenomenon of electric charge separation along the axis of magnetic field is known as the chiral magnetic effect (CME) [1]. The chiral separation effect (CSE) is a similar process, where the chiralities can be separated along the magnetic field direction due to the presence of the electric charges. The CME and CSE together form a collective excitation, known as the chiral magnetic wave (CMW). The propagation of the CMW produces an electric quadrupole moment in the system, where additional positive (negative) charges are accumulated away from (close to) the reaction plane [4].

Email address: sp35@rice.edu (Sang Eon Park on behalf of the CMS Collaboration)

It has been proposed that this electric quadruple moment in the quark-gluon plasma fireball can induce an effect on the second-order azimuthal anisotropic coefficient (known as “elliptic flow”, v_2) of the final-state particles, which then exhibits a negative (positive) correlation to event charge asymmetry, $A_{\text{ch}} \equiv \frac{N_+ - N_-}{N_+ + N_-}$, for positive- (negative-) charged particles [4]. Here, N_+ and N_- denote the number of positive- and negative-charged particles in each event. Therefore, the elliptic flow coefficients (v_2) of positive- and negative-charged particles show a linear dependence on A_{ch} as follows, $v_{2,\pm} = v_{2,\pm}^{\text{base}} \mp r A_{\text{ch}}$, where $v_{2,\pm}^{\text{base}}$ represents the $v_{2,\pm}$ value when $A_{\text{ch}} = 0$ and the r denotes the slope parameter.

Recent measurements of charge asymmetry dependence of $v_{2,\pm}$ in AA collisions made at the BNL’s RHIC and the CERN’s LHC provide evidence in line with the CMW mechanism [5, 6, 7]. However, the interpretation of the results still remains highly debated. For example, the local charge conservation effect has been shown to qualitatively explain the charge-dependent v_2 data as a function of event charge asymmetry [8], where the signal arises from the interplay of finite detector acceptance and the dependence of v_n on pseudorapidity (η) and/or transverse momentum (p_T) for charged particles from decays of resonances, resulting in a correlation between p_T (η)-averaged v_n and A_{ch} .

This contribution presents the measurements of charge asymmetry dependence of v_2 coefficients in pPb collisions, and both v_2 and v_3 coefficients in PbPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, using data collected with the CMS detector at the LHC [9]. The CMW contribution to any charge asymmetry dependent v_n signal is expected to be very small in a pPb collision compared to a PbPb collision with similar event multiplicity. This is mainly due to the fact that the induced magnetic field in pPb is much reduced in strength, and most importantly, oriented randomly with respect to the event plane [10]. Furthermore, a direct comparison of A_{ch} -dependent v_2 and v_3 coefficients in PbPb collisions can provide important evidence to differentiate between the CMW and local charge conservation (LCC) mechanisms.

2. Results




Fig. 1. The linear slope for v_2 (left) and $\langle p_T \rangle$ (right) parameters $r^{\text{norm}}(v_2)$ and $r^{\text{norm}}(\langle p_T \rangle)$ as a function of event multiplicity in pPb and PbPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV. Statistical and systematic uncertainties are indicated by the error bars and shaded regions, respectively [11]

Figure 1 (left) shows the extracted slope parameters for v_2 , averaged over $0.3 < p_T < 3\text{GeV}/c$, as a function of event multiplicity in pPb and PbPb collisions at $\sqrt{s_{\text{NN}}} = 5.02\text{TeV}$. The linear slope parameter, $r^{\text{norm}}(v_2)$, is extracted by linearly fitting the normalized v_2 difference, $(v_2^- - v_2^+)/ (v_2^- + v_2^+)$, as a function of the corrected A_{ch} .

In the overlapping multiplicity range between pPb and PbPb, slope parameter for pPb was slightly larger than that of PbPb. As discussed previously, no CMW effect is expected in high-multiplicity pPb events. Therefore, significant slope parameter observed in pPb compared to that of PbPb poses a considerable challenge to the mechanism of the CMW.

The event-averaged p_T values ($\langle p_T \rangle$) for positive and negative charged particles as a function of A_{ch} show the same pattern as in v_2 case, therefore we can extract the slope parameter $r^{\text{norm}}(\langle p_T \rangle)$ for the event averaged p_T values the same way, by linearly fitting the normalized $\langle p_T \rangle$ difference as a function of the corrected A_{ch} . Fig. 1 (right) shows the $r^{\text{norm}}(\langle p_T \rangle)$ for pPb and PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV as a function of $N_{\text{trk}}^{\text{offline}}$. Non-zero slope parameters for $\langle p_T \rangle$ are observed for all measured multiplicities, which account for about 50% of the observed slope for v_2 at $0.3 < p_T < 3\text{ GeV}/c$. This observation is in line with the predicted local charge conservation effect.




Fig. 2. The triangular flow v_3 for positive- and negative-charged particles (left) and the normalized difference in v_n , $(v_n^- - v_n^+)/ (v_n^- + v_n^+)$, for $n = 2$ and $n = 3$ (right) as a function of corrected event charge asymmetry for 30–40% centrality of PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. Statistical uncertainties are covered by the data points [11].

The charge asymmetry dependence of triangular flow v_3 for positive- and negative-charged particles is also studied in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, shown in Fig. 2 (left) for 30–40% centrality range. Similar to the case of v_2 , the v_3^+ (v_3^-) decreases (increases) as A_{ch} increases.

The normalized v_3 difference, $(v_3^- - v_3^+)/ (v_3^- + v_3^+)$, is then plotted as a function of the corrected A_{ch} values in Fig. 2 (right), and also compared to that for v_2 . As shown, the slopes $r^{\text{norm}}(v_2)$ and $r^{\text{norm}}(v_3)$ agree well within statistical uncertainties. The $r^{\text{norm}}(v_n)$ for both $n = 2$ and $n = 3$ as a function of centrality of PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV are shown in Fig. 3 for the centrality range of 30–80%. Over the centrality range studied, the slope parameters for elliptic and triangular flow are consistent with each other within uncertainties.

In the CME and CSE, the charges are expected to separate with respect to the reaction plane (Ψ_{RP}), or the second-order event plane (Ψ_2). Therefore CMW effect is highly suppressed with respect to the third-order event plane (Ψ_3), leading to a vanishing slope parameter $r^{\text{norm}}(v_3)$. The similar $r^{\text{norm}}(v_2)$ and $r^{\text{norm}}(v_3)$ values between elliptic and triangular flow observed in the data again indicate an underlying physics mechanism not related to the CMW. This behavior is qualitatively consistent with the prediction from the local charge conservation effects [8].

3. Summary

In summary, the charge-dependent elliptic (v_2) and triangular (v_3) flow coefficients in pPb and PbPb have been measured as a function of event charge asymmetry at $\sqrt{s_{NN}} = 5.02$ TeV. The normalized slope parameters of v_2 in pPb and PbPb collisions are found to be similar at the same event charged-particle multiplicities. Similar charged asymmetry dependence is also observed for the event-averaged p_T values of positive- and negative-charged particles. Moreover, in PbPb collisions, the normalized slope parameters of the v_2 and v_3 coefficients show similar magnitudes in various centralities. All these observations would not be expected from the mechanism of chiral magnetic wave, but are qualitatively consistent with predictions from the local charge conservation mechanism.




Fig. 3. The linear slope ($r^{\text{norm}}(v_n)$) parameter for elliptic ($n = 2$) and triangular ($n = 3$) flow as a function of centrality for PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. Statistical and systematic uncertainties are indicated by the error bars and shaded regions, respectively [11].

References

- [1] D. Kharzeev, Parity violation in hot QCD: Why it can happen, and how to look for it, Phys. Lett. B 633 (2006) 260. arXiv:hep-ph/0406125, doi:10.1016/j.physletb.2005.11.075.
- [2] D. E. Kharzeev, L. D. McLerran, H. J. Warringa, The Effects of topological charge change in heavy ion collisions: 'Event by event P and CP violation', Nucl. Phys. A 803 (2008) 227. arXiv:0711.0950, doi:10.1016/j.nuclphysa.2008.02.298.
- [3] D. E. Kharzeev, J. Liao, S. A. Voloshin, G. Wang, Chiral magnetic and vortical effects in high-energy nuclear collisions—A status report, Prog. Part. Nucl. Phys. 88 (2016) 1. arXiv:1511.04050, doi:10.1016/j.pnpnp.2016.01.001.
- [4] Y. Burnier, D. E. Kharzeev, J. Liao, H.-U. Yee, Chiral magnetic wave at finite baryon density and the electric quadrupole moment of quark-gluon plasma in heavy ion collisions, Phys. Rev. Lett. 107 (2011) 052303. arXiv:1103.1307, doi:10.1103/PhysRevLett.107.052303.
- [5] B. Abelev, et al., Charge separation relative to the reaction plane in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 110 (2013) 012301. arXiv:1207.0900, doi:10.1103/PhysRevLett.110.012301.
- [6] L. Adamczyk, et al., Observation of charge asymmetry dependence of pion elliptic flow and the possible chiral magnetic wave in heavy-ion collisions, Phys. Rev. Lett. 114 (2015) 252302. arXiv:1504.02175, doi:10.1103/PhysRevLett.114.252302.
- [7] J. Adam, et al., Charge-dependent flow and the search for the chiral magnetic wave in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C 93 (2016) 044903. arXiv:1512.05739, doi:10.1103/PhysRevC.93.044903.
- [8] A. Bzdak, P. Bozek, Contributions to the event-by-event charge asymmetry dependence for the elliptic flow of π^+ and π^- in heavy-ion collisions, Phys. Lett. B 726 (2013) 239. arXiv:1303.1138, doi:10.1016/j.physletb.2013.08.003.
- [9] S. Chatrchyan, et al., The CMS Experiment at the CERN LHC, JINST 3 (2008) S08004. doi:10.1088/1748-0221/3/08/S08004.
- [10] V. Khachatryan, et al., Observation of charge-dependent azimuthal correlations in pPb collisions and its implication for the search for the chiral magnetic effect, Phys. Rev. Lett. 118, 122301 (2017). arXiv:1610.00263, doi:10.1103/PhysRevLett.118.122301.
- [11] Charge asymmetry dependence of elliptic and triangular flow in pPb and PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Tech. Rep. CMS-PAS-HIN-16-017, CERN, Geneva (2017). URL <https://cds.cern.ch/record/2244682>