EUROPEAN ORGANISATION FOR NUCLEAR RESEARCH (CERN)

CERN-EP-2017-250 16th November 2018

Search for heavy resonances decaying into a *W* or *Z* boson and a Higgs boson in final states with leptons and *^b*-jets in 36 fb[−]¹ of [√] *s* = 13 TeV *pp* collisions with the ATLAS detector

The ATLAS Collaboration

A search is conducted for new resonances decaying into a *W* or *Z* boson and a 125 GeV Higgs boson in the $\nu \bar{\nu} b \bar{b}$, $\ell^{\pm} \nu b \bar{b}$, and $\ell^{+} \ell^{-} b \bar{b}$ final states, where $\ell^{\pm} = e^{\pm}$ or μ^{\pm} , in *pp* collisions at $\Delta \ell_{\bar{b}} = 13$ TeV. The data used correspond to a total integrated luminosity o boson in the *VVDD*, ℓ *VDD*, and ℓ ℓ *DD* intervalses, where $\ell = \ell$ or μ , in *pp* considers at $\sqrt{s} = 13$ TeV. The data used correspond to a total integrated luminosity of 36.1 fb⁻¹ collected with the ATLAS detector at the Large Hadron Collider during the 2015 and 2016 data-taking periods. The search is conducted by examining the reconstructed invariant or transverse mass distributions of *Wh* and *Zh* candidates for evidence of a localised excess in the mass range of 220 GeV up to 5 TeV. No significant excess is observed and the results are interpreted in terms of constraints on the production cross-section times branching fraction of heavy W' and Z' resonances in heavy-vector-triplet models and the CP-odd scalar boson *A* in two-Higgs-doublet models. Upper limits are placed at the 95% confidence level and range between 9.0×10^{-4} pb and 8.1×10^{-1} pb depending on the model and mass of the resonance.

 c 2018 CERN for the benefit of the ATLAS Collaboration. Reproduction of this article or parts of it is allowed as specified in the CC-BY-4.0 license.

1 Introduction

The ATLAS [\[1\]](#page-27-0) and the CMS [\[2\]](#page-27-1) collaborations discovered a Higgs boson (*h*) with a mass near 125 GeV and properties consistent with the Standard Model (SM) predictions [\[3–](#page-27-2)[5\]](#page-27-3). Two of the most important questions that remain are how the Higgs boson mass is protected against large radiative corrections (the naturalness problem [\[6](#page-27-4)[–8\]](#page-27-5)) and whether the Higgs boson is part of an extended scalar sector [\[9\]](#page-27-6), thus making this particle important for searches for new physics beyond the SM.

Various models with dynamical electroweak symmetry breaking scenarios attempt to solve the naturalness problem by assuming a new strong interaction at a higher energy scale. These models generically predict the existence of new vector resonances that naturally decay into a vector boson and a Higgs boson, for example in Minimal Walking Technicolour [\[10–](#page-27-7)[12\]](#page-27-8), Little Higgs [\[13\]](#page-27-9), and composite Higgs models [\[14,](#page-27-10) [15\]](#page-27-11). The decays into a vector-boson and Higgs boson final state are frequently enhanced in these models.

Another possible extension of the SM includes the addition of a second Higgs doublet [\[16\]](#page-27-12). A second Higgs doublet arises in many theories beyond the SM, collectively called two-Higgs-doublet models (2HDMs), such as the minimal supersymmetric SM [\[17–](#page-27-13)[21\]](#page-28-0), axion models [\[22\]](#page-28-1), and baryogenesis models [\[23\]](#page-28-2). In 2HDMs with a CP-conserving Higgs potential, the scalar sector of the theory consists of five Higgs bosons: two charged (H^{\pm}) , two neutral CP-even (h, H) and one neutral CP-odd (A) .

This paper describes a search for the production of new heavy vector bosons, denoted hereafter by W' and *Z* 0 , that decay into a *W* or a *Z* boson and an *h* boson and a search for a heavy CP-odd scalar boson *A* that decays into a *Z* and an *h* boson. The analyses described here target leptonic decays of the vector bosons $(W^{\pm} \to \ell^{\pm} \nu, Z \to \ell^{\pm} \ell^{-}/\nu \bar{\nu}; \ell^{\pm} = e^{\pm}, \mu^{\pm}$ and decays of the *h* boson into a *b*-quark pair. This results in three search channels: $W' \to W^{\pm} h \to \ell^{\pm} \nu b \bar{b}$, $Z' \ell A \to Z h \to \ell^{\pm} \nu b \bar{b}$, and $Z' \ell A \to Z h \$ three search channels: $W' \to W^{\pm} h \to \ell^{\pm} \nu b \bar{b}$, $Z'/A \to Z h \to \ell^{\pm}$ \bar{b} *b* \bar{b} , and $Z'/A \rightarrow Zh \rightarrow \nu \bar{\nu} b \bar{b}$.

Resonance searches are typically not sensitive to all free parameters of the underlying theory, thus simplified models [\[24\]](#page-28-3) can be used to parameterise a broad class of models, wherein only the relevant couplings and mass parameters are retained in the Lagrangian. For the interpretation of the results in the context of models with heavy vector triplets (HVT), a simplified model [\[25,](#page-28-4) [26\]](#page-28-5), based on a phenomenological Lagrangian is used as a benchmark. This model incorporates an $SU(2)_L$ triplet of heavy vector bosons, which allows the results to be interpreted in a large class of models. The new heavy vector bosons, W' and Z' , collectively denoted by *V'*, couple to the Higgs and gauge bosons via a combination of parameters g_Vc_H
and to the fermions via the combination (a^2/α) , c_R where a is the SU(2), gauge coupling. The paraand to the fermions via the combination (g^2/g_V) c_F , where g is the SU(2)_L gauge coupling. The para-
meter g_V represents the strength of the new vector-boson interaction, and cu and cu represent corrections meter q_V represents the strength of the new vector-boson interaction, and c_H and c_F represent corrections to the coupling strength specific to Higgs bosons and fermions, respectively. Two benchmark models are used in this analysis. In the first model, referred to as *Model A*, the branching fractions to fermions and gauge bosons are comparable, as in some models with an extended gauge symmetry [\[27\]](#page-28-6). For *Model B*, fermionic couplings are suppressed, as in strong dynamical models such as the minimal composite Higgs model $[28]$. At low resonance masses and large q_V couplings, the HVT models fail to reproduce the SM parameters, thus this search focuses on high masses, from 500 GeV up to 5 TeV.

The results from the $A \rightarrow Zh$ search are interpreted as exclusion limits on the ratio of the vacuum expectation values of the two Higgs doublets, $tan(\beta)$, and on $cos(\beta - \alpha)$, where α is the mixing angle between the two CP-even Higgs bosons. The exclusion limits are evaluated for the Type I, Type II, Lepton-specific, and Flipped 2HDMs. These differ with respect to which doublets couple to the up-type and down-type quarks as well as to the charged leptons $[16]$. Both the production via gluon–gluon fusion and the production with associated *b*-quarks (*bbA*) are considered in this search. The $A \rightarrow Zh$ decay mode is mostly relevant below the $t\bar{t}$ production threshold and the cross-section falls steeply with increasing *A* boson mass. Therefore, this search starts at the *Zh* threshold of approximately 220 GeV and goes up to 2 TeV.

Previous searches in the same final states have been performed by the ATLAS and the CMS collaborations using data at \sqrt{s} = 8 TeV and 13 TeV. The ATLAS searches for $W' \rightarrow Wh$ ($Z' \rightarrow Zh$) exclude, at \sqrt{s} 95% confidence level (CL), W' (Z') resonances with masses below 1.75 (1.49) TeV assuming the HVT benchmark *Model A* ($g_V = 1$) and below 2.22 (1.58) TeV assuming *Model B* ($g_V = 3$) [\[29,](#page-28-8) [30\]](#page-28-9). Searches by the CMS Collaboration exclude resonances with masses less than 2.0 TeV at 95% CL assuming the HVT benchmark *Model B* ($q_V = 3$) [\[31\]](#page-28-10). Searches using the fully hadronic final state ($W/Zh \rightarrow qq'b\bar{b}$) have also been performed by CMS and ATLAS and exclude *W' (Z')* resonances below 3.15 TeV (2.6 TeV) assuming the HVT benchmark *Model B* ($g_V = 3$) [\[32–](#page-28-11)[34\]](#page-29-0). Previous searches for a CP-odd scalar boson *A* in the *Zh* decay mode are reported in Refs. [\[35–](#page-29-1)[39\]](#page-29-2).

The search presented in this paper is performed by looking for a localised excess in the distribution of the reconstructed mass of the $v\bar{v}b\bar{b}$, $\ell^{\pm}vb\bar{b}$, and $\ell^{\pm}\ell^{\pm}b\bar{b}$ systems. The mass range covered by the search, f_{rom} 220 GeV to 5 TeV, probes a wide range of Higgs boson transverse momenta. Thus, from 220 GeV to 5 TeV, probes a wide range of Higgs boson transverse momenta. Thus, two methods are used to reconstruct Higgs boson candidates. At low transverse momenta, the decay products of the Higgs boson are reconstructed as individual jets. At high transverse momenta, the decay products start to merge and are reconstructed as a single jet. The signal yield and background normalisations are determined from a binned maximum-likelihood fit to the data distribution for each of the V' and A boson models (W', Z', \mathbb{Z}') gluon–gluon fusion *A*, *bbA*) and are used to set upper limits on the production cross-section times decay branching fraction. A combined fit using all three lepton channels sets bounds on the HVT model in the case where the V' bosons are degenerate in mass.

This paper is structured as follows. Sections [2](#page-2-0) and [3](#page-3-0) provide a brief description of the ATLAS experiment and the data and simulated event samples. The event reconstruction and selections are discussed in Sections [4](#page-5-0) and [5.](#page-8-0) The background estimation and systematic uncertainties are described in Sections [6](#page-11-0) and [7.](#page-13-0) Finally, Sections [8](#page-15-0) and [9](#page-26-0) detail the statistical analysis and provide a discussion of the results and concluding remarks.

2 ATLAS detector

The ATLAS detector $[40]$ at the LHC covers nearly the entire solid angle^{[1](#page-2-1)} around the collision point. It consists of an inner tracking detector (ID) surrounded by a thin superconducting solenoid, electromagnetic and hadronic calorimeters, and a muon spectrometer incorporating three large superconducting toroid magnets.

The ID is immersed in a 2T axial magnetic field and provides charged-particle tracking in the range $|\eta|$ < 2.5. It consists of silicon pixel, silicon microstrip, and transition radiation tracking detectors. Prior to the data-taking at the increased centre-of-mass energy of 13 TeV, the ID was enhanced by adding a new layer of pixel detectors (the IBL [\[41\]](#page-29-4)) inside the existing pixel detector layers in the barrel region (at a radius of approximately 33 mm). The upgraded detector typically provides four three-dimensional

 $¹$ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector</sup> and the *z*-axis along the beam pipe. The *x*-axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the *z*-axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta \eta)^2 + (\Delta \phi)^2}.$

measurements for tracks originating from the luminous region. The silicon microstrip tracker provides four two-dimensional measurement points per track. The transition radiation tracker enables track reconstruction at large radii up to $|\eta| = 2.0$ and provides electron identification information based on the number of hits above the threshold for transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, electromagnetic calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) electromagnetic calorimeters. An additional thin LAr presampler, covering $|\eta| < 1.8$, is used to correct for energy loss in material upstream of the calorimeters. Hadronic calorimetry is provided by a steel/scintillatortile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadronic endcap calorimeters. The solid-angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic measurements, respectively.

The muon spectrometer is composed of separate trigger and high-precision tracking chambers, measuring the deflection of muons in a magnetic field generated by superconducting air-core toroids. The precision chamber system covers the region $|\eta| < 2.7$ with three layers of monitored drift tubes, complemented by cathode strip chambers in the forward region, where the particle flux is highest. The muon trigger system covers the range $|\eta|$ < 2.4 with resistive plate chambers in the barrel, and thin-gap chambers in the endcap regions.

A two-level trigger system is used to select interesting events [\[42\]](#page-29-5). The level-1 trigger is implemented in hardware and uses a subset of detector information to reduce the event rate to a design value of at most 100 kHz. This is followed by the software-based trigger level, the high-level trigger, which reduces the event rate further to about 1 kHz.

3 Data and simulated event samples

The data used in this analysis were recorded with the ATLAS detector during the 2015 and 2016 *pp*collision runs at $\sqrt{s} = 13$ TeV and correspond to a total integrated luminosity of 36.1 fb⁻¹. The data
are required to satisfy a number of criteria that ensure that the ATLAS detector was in good operating are required to satisfy a number of criteria that ensure that the ATLAS detector was in good operating condition. A number of Monte Carlo (MC) simulation samples are used to model the background and signal processes for this search.

For the W' and *Z'* processes, simulated events were generated with MADGRAPH5_AMC@NLO (MG5_aMC) 2.2.2 [\[43\]](#page-29-6) at leading-order (LO) accuracy using the NNPDF 2.3 LO parton density function (PDF) set [\[44\]](#page-29-7). The parton shower and hadronisation were simulated with Pythia 8.186 [\[45\]](#page-29-8) using the A14 set [\[46\]](#page-29-9) of tuned parameters ("tune") together with the NNPDF 2.3 LO PDF set. Events were generated for a range of resonance masses from 500 to 5000 GeV, assuming a zero natural width. Higgs boson decays into $b\bar{b}$ and $c\bar{c}$ pairs were simulated, with a relative branching fraction $B(h \to c\bar{c})/B(h \to b\bar{b}) = 0.05$ fixed to the SM prediction [\[47\]](#page-29-10).

Events for the gluon–gluon fusion production of *A* bosons were generated at LO accuracy using the same set-up as for the *W'* and *Z'* samples. The *b*-quark associated production of *A* bosons was simulated with MG5 aMC 2.2.3 using next-to-leading-order (NLO) matrix elements with massive *b*-quarks and the CT10F4 NLO PDF set [\[48\]](#page-29-11). The parton shower and hadronisation were simulated with Pythia 8.210 [\[49\]](#page-30-0). Events were generated for a range of *A* boson masses from 220 to 2000 GeV assuming a zero natural width. For the *A* boson signals, only decays of the Higgs boson into a $b\bar{b}$ pair were generated.

For the interpretation of the $A \to Zh$ search in the context of 2HDMs, the masses of the H^{\pm} and *H* bosons are assumed to be equal to the mass of the *A* boson. The cross-sections were calculated using up to next-to-next-to-leading-order (NNLO) QCD corrections for gluon–gluon fusion and *b*-quark associated production in the five-flavour scheme as implemented in Sushi [\[50](#page-30-1)[–53\]](#page-30-2). For the *b*-quark associated production, a cross-section in the four-flavour scheme was also calculated as described in Refs. [\[54,](#page-30-3) [55\]](#page-30-4) and the results were combined with the five-flavour scheme calculation following Ref. [\[56\]](#page-30-5). The *A* boson width and the branching fractions for $A \rightarrow Zh$ and $h \rightarrow b\bar{b}$ have been calculated using 2HDMC [\[57,](#page-30-6) [58\]](#page-30-7). The procedure for the calculation of the cross-section and branching fractions as well as the choice of the 2HDM parameters follows Ref. [\[9\]](#page-27-6).

The production of *W* and *Z* bosons in association with jets was simulated with SHERPA 2.2.1 [\[59\]](#page-30-8) using the NNPDF 3.0 NNLO PDF set [\[60\]](#page-30-9) for both the matrix element calculation and the dedicated parton-shower tuning developed by the SHERPA authors. The event generation utilises COMIX $[61]$ and OPENLOOPS $[62]$, for the matrix element calculation, matched to the Sherpa parton shower using the ME+PS@NLO prescription [\[63\]](#page-30-12). The matrix elements were calculated for up to two additional partons at NLO and for three and four partons at LO in QCD. The cross-sections for *W*/*Z*+jets were calculated at NNLO accuracy [\[64\]](#page-31-0).

The $t\bar{t}$ process was simulated with Powheg-Box v2 [\[65–](#page-31-1)[67\]](#page-31-2) interfaced to PYTHIA 6.428 [\[68\]](#page-31-3). The CT10 PDF set [\[48\]](#page-29-11) was used in the calculation of the matrix elements, while the parton shower used the Perugia 2012 tune [\[69\]](#page-31-4) with the CTEQ6L1 PDF set [\[70\]](#page-31-5). The cross-section was calculated at NNLO accuracy including the resummation of next-to-next-to-leading logarithmic (NNLL) soft gluon terms with Top++2.0 [\[71–](#page-31-6)[77\]](#page-31-7). The predicted transverse momentum spectra of top quarks and the $t\bar{t}$ system were reweighted to the corresponding NNLO parton-level spectra [\[78\]](#page-31-8).

The production of single top quarks (*t*-channel, *s*-channel, and *Wt*) was simulated using the Powheg-Box event generator with the CT10 PDF set. The shower and hadronisation was simulated using the same event generator and set-up as for the $t\bar{t}$ process. The cross-section for the t - and *s*-channel single-topquark production was calculated at NLO accuracy using Hathor v2.1 [\[79,](#page-32-0) [80\]](#page-32-1), while for the *Wt* process an approximate NNLO calculation was used $[81]$. The top mass is set fixed to 172.5 GeV in the $t\bar{t}$ and single-top-quark samples.

Diboson events (*WW*, *WZ*, *ZZ*) were simulated using the SHERPA 2.1.1 event generator using the CT10 PDFs. Matrix elements were calculated for up to one (*ZZ*) or no (*WW*, *WZ*) additional partons at NLO and up to three additional partons at LO, and the cross-sections were calculated at NLO accuracy.

Finally, the SM processes $Vh(h \to b\bar{b})$, $\bar{t}h$, $\bar{t}W$, and $\bar{t}Z$ are included in the total background estimation. The $q\bar{q} \rightarrow Zh$ and $q\bar{q} \rightarrow Wh$ processes were simulated at LO with PYTHIA 8.186 using the NNPDF 2.3 LO PDF set and the A14 tune. The $qq \rightarrow Zh$ process was simulated at NLO using the Powheg-Box v2 event generator with the CT10 PDF set. The modelling of the shower, hadronisation and underlying event was provided by Pythia 8.186 using the AZNLO tune [\[82\]](#page-32-3) with the same PDF set as for the matrix element calculation. The cross-sections for the *Wh* and *Zh* processes were taken from Ref. [\[9\]](#page-27-6). The *tth* and \overline{t} V samples were generated at NLO accuracy with MG5_aMC 2.3.2 interfaced to PYTHIA 8.210. The NNPDF 3.0 NLO PDF set was used in the matrix element calculation while for the parton shower the A14 tune was used with the NNPDF 2.3 LO PDF set.

A summary of event generators used for the simulation of signal and background processes is shown in Table [1.](#page-5-1)

All simulated event samples include the effect of multiple *pp* interactions in the same and neighbouring bunch crossings (pile-up) by overlaying simulated minimum-bias events on each generated signal or

Table 1: Event generators used for the simulation of the signal and background processes. The acronyms ME and PS are used for matrix element and parton shower, respectively.

background event. The minimum-bias events were simulated with the single-, double- and non-diffractive *pp* processes of PyTHIA 8.186 using the A2 tune [\[83\]](#page-32-4) and the MSTW2008 LO PDF [\[84\]](#page-32-5). For all MAD-Graph and Powheg samples, the EvtGen v1.2.0 program [\[85\]](#page-32-6) was used for the bottom and charm hadron decays. The generated samples were processed using the Geant-based ATLAS detector simulation [\[86,](#page-32-7) [87\]](#page-32-8) and the same event reconstruction algorithms were used as for the data.

4 Event reconstruction

This search makes use of the reconstruction of multi-particle vertices, the identification and the kinematic properties of reconstructed electrons, muons, τ leptons, jets, and the determination of missing transverse momentum.

Collision vertices are reconstructed from at least two ID tracks with transverse momentum $p_T > 400$ MeV. The primary vertex is selected as the one with the highest $\sum p_T^2$, calculated considering all associated tracks.

Electrons are reconstructed from ID tracks that are matched to energy clusters in the electromagnetic calorimeter. The clusters are reconstructed using the standard ATLAS sliding-window algorithm, which clusters calorimeter cells within fixed-size η/ϕ rectangles [\[88\]](#page-32-9). Electron candidates are required to satisfy requirements for the electromagnetic shower shape, track quality, and track–cluster matching; these requirements are applied using a likelihood-based approach. The "Loose" and "Tight" working points defined in Ref. [\[89\]](#page-32-10) are used.

Muons are identified by matching tracks found in the ID to either full tracks or track segments reconstructed in the muon spectrometer ("combined muons"), or by stand-alone tracks in the muon spectrometer [\[90\]](#page-32-11). Muons are required to pass identification requirements based on quality requirements applied to the ID and muon spectrometer tracks. The "Loose" and "Medium" identification working points defined in Ref. [\[90\]](#page-32-11) are used in this analysis. The "Loose" working point includes muons reconstructed with the muon spectrometer alone to extend the acceptance to $|\eta| = 2.7$.

Electron and muon candidates are required to have a minimum p_T of 7 GeV and to lie within a region where there is good reconstruction and identification efficiency ($|\eta| < 2.7$ for muons and $|\eta| < 2.47$ for electrons). The "Loose" lepton identification criterion has to be fulfilled and all candidates have to originate from the primary vertex. The last condition is satisfied by requiring that the significance of the transverse impact parameter $|d_0|/\sigma(d_0)$ is less than 5.0 for electrons (< 3.0 for muons) and $|z_0 \sin(\theta)|$ is less than 0.5 mm, where z_0 is the longitudinal impact parameter and θ is the polar angle defined in Section [2.](#page-2-0) The lepton candidates are required to be isolated using requirements on the sum of the p_T of the tracks lying in a cone around the lepton direction whose size, ∆*R*, decreases as a function of the lepton *p*_T [\[91\]](#page-32-12). The efficiency of the isolation selection is tuned to be larger than 99% in a sample of $Z \rightarrow \ell^+ \ell^-$ decays [\[88,](#page-32-9) [90,](#page-32-11) [92\]](#page-32-13). The identification and isolation efficiencies of both the electrons and muons are calibrated using a tag-and-probe method in $Z \to \ell^+$ $-$ data events $[88, 90]$ $[88, 90]$ $[88, 90]$.

Two types of calorimeter-based jets, "small-*R*" and "large-*R*" jets, are used to reconstruct Higgs boson candidates over a wide momentum spectrum. Small-*R* jets are reconstructed from noise-suppressed topological clusters in the calorimeter $[93]$ using the anti- k_t [\[94\]](#page-32-15) algorithm implemented in the FastJET package [\[95\]](#page-32-16) with a radius parameter $R = 0.4$ and are required to have a $p_T > 20$ GeV for $|\eta| < 2.5$ (central jets) or $p_T > 30$ GeV for 2.5 $\langle |\eta| \rangle$ < 4.5 (forward jets). To reduce the number of small-*R* jets originating from pile-up interactions, these jets are required to pass the jet vertex tagger [\[96\]](#page-33-0) selection, with an efficiency of about 90%, if they are in the range $p_T < 60$ GeV and $|\eta| < 2.4$.

Large-*R* jets are used to reconstruct Higgs boson candidates with high momenta for which the *b*-quarks are emitted close to each other. They are constructed using the anti-*k^t* algorithm with a radius parameter of $R = 1.0$ and are trimmed [\[97\]](#page-33-1) to remove the energy of clusters that originate from initial-state radiation, pile-up interactions or the underlying event. This is done by reclustering the constituents of the initial jet, using the k_t algorithm [\[98,](#page-33-2) [99\]](#page-33-3), into smaller $R_{sub} = 0.2$ subjets and then removing any subjet that has a p_T less than 5% of the p_T of the parent jet [\[100\]](#page-33-4). The jet mass resolution is improved at high momentum using tracking in addition to calorimeter information [\[101\]](#page-33-5). Large-*R* jets are required to have $p_T > 250$ GeV and $|\eta| < 2.0$.

The momenta of both the large-*R* and small-*R* jets are corrected for energy losses in passive material and for the non-compensating response of the calorimeter. Small-*R* jets are also corrected for the average additional energy due to pile-up interactions [\[102,](#page-33-6) [103\]](#page-33-7).

A third type of jet, built from tracks (hereafter referred to as a track-jet), is used in this analysis for the identification of *b*-jets from decays of boosted Higgs bosons. The jets are built with the anti-*k^t* algorithm with $R = 0.2$ from at least two ID tracks with $p_T > 400$ MeV associated with the primary vertex, or with a longitudinal impact parameter $|z_0 \sin(\theta)| < 3$ mm [\[104\]](#page-33-8). Track-jets are required to have $p_T > 10$ GeV and $|\eta|$ < 2.5, and are matched to the large-*R* jets via ghost-association [\[105\]](#page-33-9).

Small-*R* jets and track-jets containing *b*-hadrons are identified with the multivariate MV2c10 *b*-tagging algorithm [\[106,](#page-33-10) [107\]](#page-33-11), which makes use of information about the jet kinematics, the properties of tracks within jets, and the presence of displaced secondary vertices. The algorithm is used at the 70% efficiency working point and provides a factor of 380 (120) in rejecting small-*R* jets (track-jets) from gluons and light quarks, and a factor of 12 (7) in rejecting small-*R* jets (track-jets) from c-quarks. Jets satisfying these requirements are referred to as "*b*-tagged jets".

To improve the mass resolution of the Higgs boson candidate, dedicated energy corrections are applied for *b*-tagged small-*R* and large-*R* jets to account for the semileptonic decays of the *b*-hadrons. The momentum of the closest muon in ∆*R* with *p*T larger than 5 GeV inside the jet cone is added to the jet momentum after removing the energy deposited by the muon in the calorimeter (the muon-in-jet correction) [\[104\]](#page-33-8). For this correction, muons are not required to pass the isolation requirements. For small-*R* jets only, an additional p_T -dependent correction, denoted "PtReco", is applied to the jet four-momentum to account for biases in the response of *b*-jets, improving the resolution of the dijet mass. This correction is determined from *Vh*($h \rightarrow b\bar{b}$) simulated events by calculating the ratio of the p_T of the true *b*-jets from the Higgs boson decay to the p_T of the reconstructed *b*-tagged jets after the muon-in-jet correction. The resolution of the dijet mass, m_{ij} (m_J), in this process is improved by 18% (22%) for the resolved (merged) Higgs boson reconstruction after these corrections, as shown in Figure [1.](#page-7-0)

Figure 1: Reconstructed mass of the Higgs boson candidates for the (left) resolved and (right) merged event topologies in a sample of simulated signal events with $m_A = 500$ GeV and $m_A = 1500$ GeV respectively. The different distributions correspond to the different *b*-jet energy corrections applied in each case as described in the text. The distributions are fit to the asymmetric function described in Ref. [\[108\]](#page-33-12) and the resolution parameter is shown in the plots.

Hadronically decaying τ -lepton candidates (τ_{had}) are identified using small-*R* jets with $p_T > 20$ GeV and $|\eta|$ < 2.5, outside the transition region between the barrel and endcap calorimeters (1.37 < $|\eta|$ < 1.52). These τ_{had} candidates must have either one or three associated tracks and must satisfy the "Medium" identification criterion [\[109\]](#page-33-13). They are used in the $v\bar{v}b\bar{b}$ channel to reject backgrounds with real hadronic τ-leptons.

The presence of neutrinos in the $v\bar{v}b\bar{b}$ and $\ell^{\pm}vb\bar{b}$ final states can be inferred from a momentum imbalance
in the transverse plane. The missing transverse momentum (\vec{F}^{miss}) is calculated as the negat in the transverse plane. The missing transverse momentum $(\vec{E}_{\text{T}}^{\text{miss}})$ is calculated as the negative vectorial sum of the transverse momenta of all the muons, electrons, small-*R* jets, and ID tracks associated with the primary vertex but not associated with any of those leptons and jets [\[110,](#page-33-14) [111\]](#page-33-15). To suppress non-collision and multijet backgrounds in the $v\bar{v}b\bar{b}$ channel, an additional track-based missing transverse momentum estimator, $\vec{p}_{\text{T}}^{\text{miss}}$, is built independently as the negative vectorial sum of the transverse momenta of all tracks from the primary vertex tracks from the primary vertex.

An overlap-removal algorithm is applied to prevent double counting of the leptons and jets used for the

resonance reconstruction. A τ-lepton is removed if the [∆]*^R* between the τ-lepton and an electron or a muon is below 0.2. In the case of a muon, the τ -lepton is not removed if the τ -lepton has p_T above 50 GeV and the reconstructed muon is not a combined muon. If a reconstructed muon and electron share the same ID track then the electron is removed. Small-*R* jets are removed if they are within a cone of size $\Delta R = 0.2$ around an electron or muon that has passed the isolation requirements. To account for semi-muonic *b*-jet decays, the jet is only removed if it has fewer than three associated tracks, or if more than 70% of the sum of the p_T of its associated tracks comes from the muon and p_T^j $\frac{f}{T}/p_{\rm T}^{\mu}$ < 2, where $p_{\rm T}^{j}$
AR = 0.4 around a $p_{\rm T}^{\prime}$ ($p_{\rm T}^{\mu}$) is the $p_{\rm T}$ of the jet (muon). Next, electrons and muons within a cone of size [∆]*^R* ⁼ ⁰.4 around a surviving small-*^R* jet are discarded if their distance from the jet direction is smaller than $\Delta R = (0.04 + 10 \,\text{GeV}/p_{\text{T}}^t)$. The shripking cone size ensures a bigh efficiency for boosted topologies. Small-R jets are also removed if they shrinking cone size ensures a high efficiency for boosted topologies. Small-*R* jets are also removed if they are within $\Delta R = 0.2$ of the axis of a τ_{had} candidate. Finally, large-*R* jets within $\Delta R = 1.2$ of any surviving electron are removed.

5 Analysis strategy and event selection

The search for the *Z'* and *A* bosons in the $Zh \rightarrow \nu \overline{\nu} b \overline{b}$ and $Zh \rightarrow \ell^+ \ell^- b \overline{b}$ decay modes uses event
samples wherein the number of reconstructed charged leptons is exactly zero or two (0-lepton and 2-lepton samples wherein the number of reconstructed charged leptons is exactly zero or two (0-lepton and 2-lepton channels). For the *W'* search in the $Wh \rightarrow \ell^{\pm}vb\overline{b}$ channel, events with exactly zero or one charged lepton
are used (0-lepton or 1-lepton channels). The lepton selection requirements described in the previous are used (0-lepton or 1-lepton channels). The lepton selection requirements described in the previous section are applied, using the "Loose" identification working point. The selections outlined below define regions sensitive to the different models.

For the 0-lepton channel, an $E_{\rm T}^{\rm miss}$ trigger with a threshold of 70 GeV was used to record the data in 2015 runs; the threshold varied between 90 and 110 GeV in 2016 runs due to the increasing instantaneous luminosity. Events are required to have $E_T^{\text{miss}} > 150$ GeV, where E_T^{miss} is reconstructed with fully calibrated
leptons and jets. The efficiency of the trigger selection exceeds 80% above 150 GeV. In the 2-lepton leptons and jets. The efficiency of the trigger selection exceeds 80% above 150 GeV. In the 2-lepton channel, events were recorded using a combination of single-lepton triggers with isolation requirements. In 2015, the lowest p_T threshold was 24 GeV; in 2016, it ranged from 24 to 26 GeV. Additional triggers without an isolation requirement are used to recover efficiency for leptons with $p_T > 60$ GeV. In the single-electron channel, the same single-electron triggers as in the 2-lepton channel are applied. In the single-muon channel, the same E_T^{miss} triggers as in the 0-lepton channel are used because they are more efficient than the single-muon triggers for this analysis. For events selected by the lepton triggers, the lepton that satisfied the trigger is required to match a reconstructed electron (muon) with $p_T > 27$ GeV and $|\eta|$ < 2.47 ($|\eta|$ < 2.5).

The wide range of resonance masses probed by this search implies that the resonance decay products can be produced with a wide range of transverse momenta. When the Higgs boson has relatively low p_T , the *b*-quarks from its decay can be reconstructed as two small-*R* jets. As the momentum of the Higgs boson increases, the two *b*-quarks become more collimated and a selection using a single large-*R* jet becomes more efficient. Two different methods are used for the reconstruction of the Higgs boson candidate: a "resolved" category in which two small-*R* jets are used to build the Higgs boson candidate, and a "merged" category where the highest- p_T ("leading") large- R jet is selected as the Higgs boson candidate.

For the resolved signal region, two small-*R* jets are required to have an invariant mass (m_{ij}) in the range 110 –140 GeV for the 0- and 1-lepton channels and in the range 100 –145 GeV for the 2-lepton channel. The latter selection is relaxed to take advantage of the smaller backgrounds in this channel. This dijet

candidate is defined by the two leading *b*-tagged small-*R* jets when two or more *b*-tagged jets are present in the event. In the case where only one *b*-tagged jet is present, the dijet pair is defined by the *b*-tagged jet and the leading small-*R* jet in the remaining set. The leading jet in the pair must have $p_T > 45$ GeV. For the merged signal region, a large-*R* jet is required with mass (m_J) in the range 75 to 145 GeV and at least one associated *b*-tagged track-jet.

Events which satisfy the selection requirements of both the resolved and merged categories, are assigned to the resolved one, since its better dijet mass resolution and lower background contamination increases the expected sensitivity. Events failing to satisfy both the resolved and merged signal region requirements are assigned to control regions defined in Section 6 , with priority given to the resolved category. This procedure provides a higher sensitivity for resonances of mass near 1 TeV compared to a procedure in which the merged category is prioritised.

Higgs boson candidates with one or two *b*-tagged jets define the "1 *b*-tag" or "2 *b*-tag" categories, respectively. For the merged selection, only one or two leading track-jets associated with the large-*R* jet are considered in this counting. For the 0- and 2-lepton channels, resolved events with more than two *b*-tagged jets or merged events with additional *b*-tagged track-jets not associated with the large-*R* jet are used to define signal regions sensitive to $b\overline{b}A$ production. These are labeled as "3+ *b*-tag" in the resolved category, and "1 *b*-tag additional *b*-tag" or "2 *b*-tag additional *b*-tag" in the merged category. In the 2-lepton channel, the latter two are merged and labeled as "1+2 *b*-tag additional *b*-tag".

The calculation of the reconstructed resonance mass depends on the decay channel. In the 0-lepton channel, where it is not possible to reconstruct the *Zh* system fully due to the presence of two neutrinos from the *Z* boson decay, the transverse mass defined as

$$
m_{\rm T, Vh} = \sqrt{(E_{\rm T}^h + E_{\rm T}^{\rm miss})^2 - (\vec{p}_{\rm T}^h + \vec{E}_{\rm T}^{\rm miss})^2},
$$

is used as the final discriminant. In order to reconstruct the invariant mass of the $Wh \to \ell^{\pm}vb\bar{b}$ system in the 1 lepton channel, the momentum of the neutrino in the z direction, n, is obtained by imposing a W the 1-lepton channel, the momentum of the neutrino in the *z*-direction, p_z , is obtained by imposing a *W* boson mass constraint on the lepton– E_T^{miss} system. In the resulting quadratic equation, the neutrino p_z is taken as the real component in the case of complex solutions, or as the smaller of the two solutions if both solutions are real. The mass resolution of the *Vh* system is improved in the resolved signal regions of all channels by rescaling the four-momentum of the dijet system by 125 GeV/ *m*jj. In the 2-lepton channel, the four-momentum of the dimuon system is scaled by 91.2 GeV / $m_{\mu\mu}$ as well in all signal regions. This helps to address the worse momentum resolution of high-momentum muons which are measured solely by the tracking detectors.

Additional selections are applied for each lepton channel, as outlined below, to reduce the main backgrounds and enhance the signal sensitivity. These selections are summarised in Table [2.](#page-11-1)

For the resolved and merged categories in the 0-lepton channel, the following selections are applied to reduce multijet and non-collision backgrounds to a negligible level:

- $p_T^{\text{miss}} > 30$ GeV (not applied in the resolved 2 and 3+ *b*-tag categories);
- the azimuthal angle between $\vec{E}_{\rm T}^{\rm miss}$ and $\vec{p}_{\rm T}^{\rm miss}$, $\Delta\phi(\vec{E}_{\rm T}^{\rm miss}, \vec{p}_{\rm T}^{\rm miss}) < \pi/2$;
- the azimuthal angle between $\vec{E}_{\rm T}^{\rm miss}$ and the Higgs boson candidate momentum direction, $\Delta\phi(\vec{E}_{\rm T}^{\rm miss}, h) >$
 $2\pi/3$. $2\pi/3$;

• the azimuthal angle between $\vec{E}_{\text{T}}^{\text{miss}}$ and the nearest small-*R* jet momentum direction, $\min[\Delta\phi(\vec{E}_{\text{T}}^{\text{miss}}, \text{small-}R \text{ jet})] > \pi/9$ (for the resolved category with four or more jets, > $\pi/6$ is used).

For the resolved category, $\min[\Delta\phi(\vec{E}_{\text{T}}^{\text{miss}}, \text{small-}R \text{ jet})]$ is calculated using the small-*R* jets that constitute the Higgs boson candidate and an additional small-*R* jet, which is the third leading h tagged ist tute the Higgs boson candidate and an additional small-*R* jet, which is the third leading *b*-tagged jet (if the event contains at least three *b*-tagged jets), the leading central jet which is not *b*-tagged (if the event contains only two *b*-tagged jets) and the leading forward jet (if the event contains only two central small-*R* jets). For the merged category, all central and forward small-*R* jets are used in the $\min[\Delta\phi(\vec{E}_{\text{T}}^{\text{miss}}, \text{small-}R]$ jet)] calculation. For the *Z'*/*A* search, the *tt* and *W*+jets backgrounds are fur-
ther reduced by rejecting events with at least one identified τ_{obs} condidate. This veto is not a ther reduced by rejecting events with at least one identified τ_{had} candidate. This veto is not applied when searching for the *W'* boson or in the HVT combined search, because it leads to a loss of signal events in the $Wh \to \tau^{\pm} \nu b \bar{b}$ final state.

For the 0-lepton resolved category, two additional selections are applied:

- the scalar sum of the p_T of the three leading central small-*R* jets, $\sum p_T^{\text{jet}}$ $T_T^{\text{ct}_i}$, is greater than 150 GeV. In the case where there are only two central small-*R* jets, the sum of the p_T of these two jets and of the leading forward small-*R* jet, if any, is required to be greater than 120 GeV;
- the azimuthal angle between the two jets used to reconstruct the Higgs boson candidate, $\Delta\phi(i, j)$, is required to be less than $7\pi/9$.

Finally, for the merged category, the missing transverse momentum must be larger than 200 GeV.

For the 1-lepton channel, a selection on the transverse momentum of the *W* boson candidate ($p_{T,W}$), which increases as a function of the reconstructed resonance mass, is applied to reduce the contribution of *W*+jets: $p_{\text{T},W}$ > max[150, 710 – 3.3 × 10⁵ GeV/*m*_{*Vh*}] GeV for the resolved category, while $p_{\text{T},W}$ > max[150, 394 · $ln(mv_h/(1 \text{ GeV}))$ – 2350] GeV for the merged category. This selection is optimised taking advantage of the larger transverse momentum of *W* bosons expected to be produced in the decays of high-mass resonances. The $t\bar{t}$ background is reduced in the resolved category by requiring fewer than four central jets in the event and in the merged category by rejecting events with additional *b*-tagged track-jets not associated with the large-*R* jet. For all categories, the transverse mass of the *W* candidate $(m_{\text{T},W})$, calculated from the transverse components of the lepton and $E_{\text{T}}^{\text{miss}}$ momentum vectors, is required to be less than 300 GeV.

In the 1-lepton channel, a significant contribution of multijet events arises mainly from non-prompt leptons from hadron decays and from jets misidentified as electrons. This background is significantly reduced by applying tighter selection requirements on the lepton isolation and identification, as well as on $E_{\rm T}^{\rm miss}$. Muons must satisfy the "Medium" identification and electrons must satisfy the "Tight" identification requirements. Stringent lepton isolation requirements are applied: the scalar sum of the p_T of tracks within a variable-size cone around the lepton (excluding its own track) must be less than 6% of the lepton p_T . In addition, in the case of electrons the sum of the transverse energy of the calorimeter energy clusters in a cone of $\Delta R = 0.2$ around the electron must be less than 6% of the electron p_T [\[90,](#page-32-11) [92\]](#page-32-13). Finally, the $E_{\rm T}^{\rm miss}$ value is required to be greater than 100 GeV for the merged category and greater than 80 (40) GeV for the resolved category in the electron (muon) channel.

In the 2-lepton channel, same-flavour leptons (*ee* or $\mu\mu$) are used. For both the resolved and merged categories, three kinematic selections are optimised as a function of the resonance mass to reduce the *tt* and *Z*+jets backgrounds. Selections on the mass of the dilepton system, max[40 GeV, 87 GeV − 0.030 · m_{Vh}] $< m_{\ell\ell} < 97 \text{ GeV} + 0.013 \cdot m_{Vh}$, and on $E_{\text{T}}^{\text{miss}}/$ $\sqrt{(1 \text{ GeV}) \cdot H_T}$ < 1.15 + 8 × 10⁻³ · *m_{Vh}* / (1 GeV) are relaxed

Table 2: Topological and kinematic selections for each channel and category as described in the text. ^(∗) Applies in the case of only two central jets. ^(**) Tau veto only applied for the *Z'* /*A* search. ^(†) Tighter threshold (80 GeV) is used for the single-electron channel. (††) Applied only for *mZh* > 320 GeV. (‡) Not applied in the resolved 2 and 3+ *b*-tag categories.

Variable	Resolved	Merged					
Common selection							
Number of jets	\geq 2 small- <i>R</i> jets (0, 2-lep.)	≥ 1 large-R jet					
	2 or 3 small- R jets (1-lep.)						
Leading jet p_T [GeV]	> 45	> 250					
$m_{\rm ii}$, $m_{\rm J}$ [GeV]	$110-140$ (0,1-lep.), $100-145$ (2-lep.)	$75 - 145$					
0-lepton selection							
$E_{\rm T}^{\rm miss}$ [GeV]	> 150	> 200					
$\sum p_{\rm T}^{\rm jet_i}$ [GeV]	$> 150(120^*)$						
$\Delta\phi(j, j)$	$< 7\pi/9$						
$p_{\rm T}^{\rm miss}$ [GeV]	$> 30^{\ddagger}$						
$\Delta\phi(\vec{E}_{\mathrm{T}}^{\mathrm{miss}}, \vec{p}_{\mathrm{T}}^{\mathrm{miss}})$	$\langle \pi/2$						
$\Delta\phi(\vec{E}_{\rm T}^{\rm miss},h)$	$> 2\pi/3$						
$min[\Delta\phi(\vec{E}_{\text{T}}^{\text{miss}}, \text{small-}R \text{ jet})]$	$> \pi/9$ (2 or 3 jets), $> \pi/6$ (≥ 4 jets)						
$N_{\tau_{\textrm{had}}}$	0^{**}						
	1-lepton selection						
Leading lepton p_T [GeV]	> 27	> 27					
$E_{\rm T}^{\rm miss}$ [GeV]	$> 40 (80^{\dagger})$	>100					
$p_{T,W}$ [GeV]	$>$ max[150, 710 – (3.3 \times 10 ⁵ GeV)/ m_{Vh}]	$> max[150, 394 \cdot ln(m_{Vh}/(1 \text{ GeV})) - 2350]$					
$m_{\text{T},W}$ [GeV]	< 300						
2-lepton selection							
Leading lepton p_T [GeV]	> 27	> 27					
Sub-leading lepton p_T [GeV]	>7	> 25					
$E_{\rm T}^{\rm miss}/\sqrt{H_{\rm T}}$ [$\sqrt{\rm GeV}$]	< 1.15 + 8 × 10 ⁻³ $\cdot m_{Vh}/(1 \text{ GeV})$						
$p_{\text{T,ell}}$ [GeV]	$> 20 + 9 \cdot \sqrt{m_{Vh}/(1 \text{ GeV}) - 320}$ ^{††}						
$m_{\ell\ell}$ [GeV]	$\left[\max[40 \text{ GeV}, 87 - 0.030 \cdot m_{Vh}/(1 \text{ GeV})], 97 + 0.013 \cdot m_{Vh}/(1 \text{ GeV})\right]$						

for higher-mass resonances to account for resolution effects and smaller backgrounds. The variable H_T is calculated as the scalar sum of the p_T of the leptons and small-*R* jets in the event. The momentum of the dilepton system $(p_{T,\ell\ell})$ is required to be greater than 20 GeV + 9 GeV · $\sqrt{m_{Vh}/(1 \text{ GeV}) - 320}$ for m_{Vh} and m_{Vh} and m_{Vh} and m_{Vh} and m_{Vh} and m_{Vh} for m_{Vh} and m_{Vh} and m_{Vh} and m m_{Vh} greater than 320 GeV. In the resolved dimuon category, an opposite-charge requirement is applied since the probability to mis-reconstruct the charge of individual muons is extremely low. Additionally, in this category the leading muon is required to have $|\eta|$ less than 2.5. Finally, for the merged category, the sub-leading lepton is required to have $p_T > 25$ GeV and for muons $|\eta|$ is restricted to be less than 2.5.

6 Background estimation

The background contamination in the signal regions is different for each of the three channels studied. In the 0-lepton channel, the dominant background sources are $Z + jets$ and $t\bar{t}$ events with a significant contribution from *W*+jets. In the 1-lepton channel, the largest backgrounds are $t\bar{t}$, single-top-quark and *W*+jets production. In the 2-lepton channel, *Z*+jets production is the predominant background followed by the $t\bar{t}$ background. The contribution from diboson, SM *Vh*, $t\bar{t}h$, and $t\bar{t}V$ production is small in all three channels. The multijet background, due to semileptonic hadron decays or misidentified jets, is found to be negligible in the 0- and 2-lepton channels after applying the event selections described in Section 5. In the 1-lepton channel, the multijet background remains significant only in the resolved 1 *b*-tag category. All background distribution shapes except those for multijet are estimated from the samples of simulated events with normalisations of the main backgrounds estimated from the data; the multijet shape and normalisation is determined using data.

The *^W*/*Z*+jets simulated event samples are split into different components. In the resolved category, the samples are split according to the true flavour of the two small-*R* jets forming the Higgs boson candidate. In the merged category, they are split according to the true flavour of the one or two leading track-jets associated with the large-*R* jet. The true jet flavour is determined by counting true heavy-flavour hadrons with $p_T > 5$ GeV within the cone of the reconstructed jet. If a true *b*-hadron is found, the jet is labelled as a *b*-jet, otherwise if a true *c*-hadron is found the jet is labelled as a *c*-jet. If neither a true *b*-hadron nor a true *c*-hadron is associated with the reconstructed jet, it is labelled as a light jet. For large-*R* jets with only one track-jet, the true hadrons are counted within this track-jet. Based on this association scheme, the *^W*/*Z*+jets simulated event samples are split into six components: *^W*/*Z*+*bb*, *^W*/*Z*+*bc*, *^W*/*Z*+*bl*, $W/Z + cc$, $W/Z + cl$ and $W/Z + ll$; in this notation *l* refers to a light jet. In the statistical analysis described in Section [8,](#page-15-0) the components *^W*/*Z*+*bb*, *^W*/*Z*+*bc*, and *^W*/*Z*+*cc* are treated as a single component denoted by *W*/*Z*+(*bb*, *bc*, *cc*). The combination of *W*/*Z*+*bl* and *W*/*Z*+*cl* is denoted by *W*/*Z*+(*bl*, *cl*). For the HVT, *Z*['], and *A* hoson interpretations, the pormalisations of the largest components *Z*+(*bb* and *A* boson interpretations, the normalisations of the largest components $Z+(bb, bc, cc)$ and $Z+(bl, cl)$ are determined from data. In the *^A* boson interpretation, the *^Z*+(*bb*, *bc*, *cc*) background normalisation in the 3⁺ *^b*-tag region is determined from this region independently. The normalisations of *^W*+(*bb*, *bc*, *cc*) and $W+(bl, cl)$ are determined from data for the W' and HVT interpretations.

The normalisation of the *tt* background is determined from the fits to data separately for the 0-, 1-, and 2 lepton channels. In the 0-lepton channel, only the signal regions are used in the fit. In the 1- and 2-lepton channels, dedicated control regions enhanced in *tt* events are used in addition to the signal regions. In the 1-lepton channel, resolved events in the sidebands of the *m*jj distribution between 50 GeV and 200 GeV (excluding the signal region with $110 < m_{ij} < 140$ GeV) are primarily composed of $t\bar{t}$ and $W +$ jets events. These control regions are included in the fit for the 1 and 2 *b*-tag categories. In the 2-lepton channel, a *tt* control region is defined using resolved events with different-flavour $(e\mu)$, oppositely charged leptons,
and without the $F^{miss} / \sqrt{H_E}$ requirement. The $t\bar{t}$ purity of this selection is greater than 90%. Thi and without the $E_T^{\text{miss}} / \sqrt{H_T}$ requirement. The $t\bar{t}$ purity of this selection is greater than 90%. This region combining the 1 and 2 *b*-tag events is used in the *A*, Z' , and HVT interpretations; for the *A* interpretation, a control region with $e\mu$ events and $3+b$ -tags is also included in the fit to provide an independent constraint on *tt* production with associated heavy-flavour jets.

The shape of the multijet background in the 1-lepton channel is estimated from a sample of data events orthogonal to the signal regions, the anti-isolated lepton region. In the muon channel, this region is defined by events where the sum of the transverse momentum of tracks in a cone of [∆]*^R* ⁼ ⁰.2 around the muon is between 6% and 15% of the muon p_T . In the electron channel, this region is defined by events where the sum of the calorimeter energy deposits in a cone of [∆]*^R* ⁼ ⁰.2 around the electron is larger than 6% of the electron p_T ; this region is defined after applying the track isolation requirement described in Section [4.](#page-5-0) A template shape for the multijet background is extracted from the anti-isolated lepton region after removing the contribution from the simulated electroweak and top-quark backgrounds. In this subtraction, the normalisation of the simulated electroweak and top-quark backgrounds is estimated

Table 3: Relative systematic uncertainties in the normalisation, cross-region extrapolation, and shape of the signal and background processes included in the fits described in the text. An "S" indicates a shape variation is included for the sources listed, "/" indicates a ratio of two regions, and "norm." is the sum of cross-section and acceptance variations. A range of values means the value depends on the lepton channel. Parentheses indicate when the uncertainty applies only to a given fit or a given region.

Process	Quantity/source	Value		Process	Quantity/source	Value
Signal	acceptance	$3 - 7\%$			norm.	19%
SM $Vh, t\bar{t}V, t\bar{t}h$	norm.			$Z+l$	resolved / merged	23%
Diboson	norm.	11%		0 -lep. $/ 2$ -lep.		8%
Multijet (1-lep.)	norm.	50%			generator, PDF, scale	S
	template method	S			norm. (A,Z')	26%
	norm.	19%		$W+(bb, bc, cc)$	resolved / merged	18-43%
Single top quark	resolved / merged 24%				$m_{\rm ii}$ SR / $m_{\rm ii}$ CR (1-lep.)	6%
	m_{ii} SR / m_{ii} CR (1-lep.)	7%			0 -lep. $/ 1$ -lep.	26%
$t\overline{t}$	resolved / merged	15-46%			generator, PDF, scale	S
	m_{ii} SR / m_{ii} CR (1-lep.)	7%		$W+(bl, cl)$	norm. (A,Z')	23%
	$SR / e\mu$ CR (2-lep.)	2%			resolved / merged	$15 - 35\%$
	PS, ISR/FSR, ME	S			$m_{\rm ii}$ SR / $m_{\rm ii}$ CR (1-lep.)	5%
	p_T reweight	S			0 -lep. $/ 1$ -lep.	22%
$Z+(bb, bc, cc)$	resolved / merged	19%			generator, PDF, scale	S
	0 -lep. $/ 2$ -lep.	15%	$W+l$		norm.	20-30%
	generator, PDF, scale	S			resolved / merged	16-20%
$Z+(bl, cl)$	resolved / merged	28%			$m_{\rm ii}$ SR / $m_{\rm ii}$ CR (1-lep.)	2%
	0-lep. $/$ 2-lep.	12%			0 -lep. $/ 1$ -lep.	19%
	generator, PDF, scale	S			generator, PDF, scale	S

by fitting them to data in the region $E_{\text{T}}^{\text{miss}} > 200$ GeV where the contribution from multijet events is negligible. In the signal and control regions used in the statistical analysis, the multijet pormalisation negligible. In the signal and control regions used in the statistical analysis, the multijet normalisation is determined by fitting the E_T^{miss} multijet template and the E_T^{miss} combined template of the electroweak and top-quark backgrounds to data in the 1 and 2 *b*-tag categories separately. Using this method, the multijet contribution is estimated to be less than 6% in all signal and control regions and is included in the statistical analysis.

7 Systematic uncertainties

Two types of systematic uncertainties, experimental and modelling, affect the reconstruction of the *mVh* and $m_{T,Vh}$ observables. Experimental uncertainties arise due to the trigger selection, the reconstruction, identification, energy/momentum, mass, and resolution for the leptons, jets and missing transverse momentum. Modelling uncertainties result in shape and normalisation uncertainties of the different MC samples used to model the signal and backgrounds. These stem from uncertainties in the matrix element calculation, the choice of parton shower and hadronisation models and their free parameters, the PDF set and the choice of renormalisation and factorisation scales.

The largest experimental systematic uncertainties are associated with the calibration and resolution of the small-*R* and large-*R* jet energy, the calibration and resolution of the large-*R* jet mass, and the determination of the jet *b*-tagging efficiency and misidentification rate. The uncertainties in the small-*R* jet energy scale have contributions from *in situ* calibration studies, from the dependency on the pile-up activity and on the flavour composition of jets $[103, 112]$ $[103, 112]$ $[103, 112]$. The small-*R* jet uncertainties are propagated to the E_T^{miss} measurement. The uncertainty in the scale and resolution of large-*R* jet energy and mass is estimated by comparing the ratio of calorimeter-based to track-based measurements in dijet data and simulation [\[100,](#page-33-4) [101\]](#page-33-5). The flavour tagging efficiency and its uncertainty for *b*-jets and *c*-jets is estimated in $t\bar{t}$ and $W + c$ jet events, respectively, while the light-jet misidentification rate and uncertainty is determined using dijet events [\[106,](#page-33-10) [107,](#page-33-11) [113,](#page-34-1) [114\]](#page-34-2). Other experimental systematic uncertainties with a smaller impact are those in the lepton energy and momentum scales, in lepton reconstruction and identification efficiency, and in the efficiency of the triggers. Finally, a global normalisation uncertainty of 3.2% is assigned due to the luminosity measurement from a preliminary calibration of the luminosity scale using *^x*–y beamseparation scans performed in August 2015 and May 2016, following a methodology similar to that detailed in Ref. [\[115\]](#page-34-3). Experimental uncertainties have an impact on the shape of the mass distributions and account for possible migration of events across the different regions.

Modelling uncertainties are assigned to each signal and background process and lead to variations in the normalisation and in the case of main backgrounds also in the shape of the templates in the different regions. In addition, for all MC samples, the statistical uncertainty arising from the number of simulated events is considered by introducing shape variations determined from the uncertainty in each bin of the m_{Vh} or $m_{T,Vh}$ distributions. The modelling uncertainties considered are shown in Table [3](#page-13-1) and described below.

For the signal processes, the uncertainties in the acceptance were derived by considering the following variations: the renormalisation and factorisation scales were varied by a factor of two, the nominal PDF set was replaced by the MSTW2008 LO PDF set and the tuned parameters were varied according to the variations derived from the eigentune method [\[46\]](#page-29-9). For both the *A* and *V'* signals, the total variations are less than 3% at resonance masses above 500 GeV. The variations increase to 7% for the *A* boson masses below 500 GeV.

The modelling uncertainties affecting $t\bar{t}$ and single-top-quark processes are derived as follows [\[116\]](#page-34-4). A variation of the parton shower, hadronisation, and the underlying-event model is obtained by replacing Pythia 6.428 by Herwig++ (version 2.7.1) [\[117\]](#page-34-5) with the UE-EE-5 tune and the CTEQ6L1 PDF set [\[70\]](#page-31-5). To assess potential differences in the matrix element calculation, a comparison is made to a sample where Powheg is replaced by MG5_aMC [\[43\]](#page-29-6). A comparison is also made to samples with smaller and larger amount of initial- and final-state radiation (ISR/FSR) by changing the renormalisation and factorisation scales by a factor of two and switching to the corresponding low- and high-radiation Perugia 2012 tunes. Finally, the difference between the nominal and corrected distributions due to the top-quark and \bar{t} p_T reweighting described in Section [3](#page-3-0) is included as a symmetrised shape uncertainty.

Similarly, for *^W*/*Z*+jets backgrounds, the following comparisons have been performed. The PDF set in the nominal samples was replaced by the alternative PDF sets: the hundred NNPDF 3.0 NNLO replicas, including the sets resulting from variations of α_s [\[60\]](#page-30-9), the MMHT2014 NNLO set, and the CT14 NNLO PDF set [\[118\]](#page-34-6). The scale uncertainties are estimated by comparing samples where the renormalisation and factorisation scales were modified by a factor of two. Finally, a comparison was made to a sample generated using MG5_aMC v2.2.2 interfaced to Pythia 8.186 and using the A14 tune together with the NNPDF 2.3 LO PDF set [\[49,](#page-30-0) [119,](#page-34-7) [120\]](#page-34-8).

For the $t\bar{t}$, single-top-quark, and $W/Z +$ jets backgrounds, the acceptance differences that affect the relative normalisation across regions with a common background normalisation are estimated by summing in

Table 4: A list of the signal and control regions (separated by commas below) included in the statistical analysis of the *A* and HVT model hypotheses. The notation 1+2 *b*-tag indicates the 1 and 2 *b*-tag regions are combined, and add. *b*-tag indicates the regions with additional *b*-tags not associated with the large-*R* jet.

Fit		Channel	Resolved	Merged	Resolved	
			signal regions	signal regions	control regions	
A		0-lepton	$1, 2, 3 + b$ -tag	1, 2 b -tag, and 1, 2 b -tag add. b -tag		
		2-lepton	$1, 2, 3 + b$ -tag	1, 2 b -tag, and $1+2$ b -tag add. b -tag	1+2 <i>b</i> -tag, $3 + b$ -tag <i>eµ</i>	
	Z', W'	0-lepton	$1, 2 b$ -tag	1, 2 b -tag		
HVT	W'	1-lepton	$1, 2 b$ -tag	$1, 2 b$ -tag	1, 2 <i>b</i> -tag m_{ii} sideband	
	Z^{\prime}	2-lepton	$1, 2 b$ -tag	$1, 2 b$ -tag	$1+2 b$ -tag e μ	

quadrature the relative yield variations between the different regions. These uncertainties are assigned to all regions used in the fit as shown in Table [4](#page-15-1) and across the different lepton channels as shown in Table [3.](#page-13-1)

For the multijet background included in the 1-lepton channel, a 50% uncertainty in the normalisation is estimated from the fit to the $E_{\text{T}}^{\text{miss}}$ distribution described in Section [6.](#page-11-0) Also, a shape variation is included to account for uncertainties in the determination of the template in the anti-isolated lepton region, arising from differences in the trigger scheme between isolated and anti-isolated regions and uncertainties in the normalisation of the top-quark and electroweak backgrounds in this region.

Finally, for the remaining small backgrounds only a normalisation uncertainty is assigned. For the diboson backgrounds a normalisation uncertainty of 11% is applied [\[121\]](#page-34-9). For the SM *Vh*, *ttV*, and *tth* production, a 50% uncertainty is assigned which covers the uncertainty in the cross-sections.

8 Results

In order to test for the presence of a massive resonance, the $m_{\text{T},Vh}$ and m_{Vh} templates obtained from the signal and background simulated event samples are fit to data using a binned maximum-likelihood approach based on the RooStats framework $[122-124]$ $[122-124]$. A total of five different fits are performed according to the signal interpretation: Z', W', HVT, A in gluon–gluon fusion, and A in b-quark associated production. The list of channels and regions used for the different fits is shown in Table [4.](#page-15-1)

The fits are performed on the $m_{T,Vh}$ distribution in the 0-lepton channel and the m_{Vh} distribution in the 1and 2-lepton channels using a binning of the distributions chosen to optimise the search sensitivity while minimising statistical fluctuations. As described in Section [6,](#page-11-0) the normalisations of the $t\bar{t}$, $Z+(bb, bc, cc)$, and $Z+(bl, cl)$ backgrounds are free parameters in all fits, as are the normalisations of $W+(bb, bc, cc)$ and $W+(bl, cl)$ in the *W*^{\prime} and HVT fits. The systematic uncertainties described in Section [7](#page-13-0) are incorporated in the fit as nuisance parameters with correlations across regions and processes taken into account. The in the fit as nuisance parameters with correlations across regions and processes taken into account. The signal normalisation is a free parameter in the fit. In order to account for migrations of signal events across different channels due to lepton reconstruction and selection inefficiencies, the $Zh \rightarrow \ell^{+} (Wh \rightarrow \ell^{\pm} v h \bar{h})$ signal samples are included in the 1(0)-lepton categories [−]*bb* $(Wh \to \ell^{\pm} \nu b\overline{b})$ signal samples are included in the 1(0)-lepton categories.

The total uncertainty in the signal yield is dominated by different sources of systematic uncertainty depending on the mass of the resonance used in the fit. The uncertainties in the *^W*/*Z*+(*bb*, *bc*, *cc*) shape and normalisation, *tt* normalisation, and in the flavour tagging efficiencies constitute the dominant sources of systematic uncertainty for low-mass resonances. For all interpretations, the statistical uncertainty dominates for resonances above 1 TeV. The uncertainties in the large-*R* jet mass resolution and in the track-jet *b*-tagging efficiency constitute the dominant systematic uncertainties at high masses.

The expected and observed event yields after the HVT fit are shown in Table [5.](#page-18-0) The m_T _{*Vh*} and m_V _{*N*} distributions after the HVT fit are shown in Figures [2](#page-19-0) and [3.](#page-20-0) Similar distributions are obtained from the W', *Z'* and *A* fits with background yields consistent within the uncertainties. The mass distributions for the resolved 3+ *b*-tag category and the merged categories with additional *b*-tagged jets, used in the *A* boson fits, are shown in Figure [4.](#page-21-0)

As no significant excess over the background prediction is observed, upper limits at the 95% CL are set on the production cross-section times the branching fraction for each model. The limits are evaluated using a modified frequentist method known as CL_s [\[125\]](#page-34-12) and the profile-likelihood-ratio test statistic [\[126\]](#page-34-13) using the asymptotic approximation.

The 95% CL upper limits on the production cross-section multiplied by the branching fraction for $W' \rightarrow$ *Wh* and $Z' \rightarrow Zh$ and the sum of branching fractions $B(h \rightarrow b\overline{b}) + B(h \rightarrow c\overline{c})$, which is fixed to 60.6% [\[47\]](#page-29-10), are shown in Figure [5\(](#page-22-0)a) and Figure 5(b) as a function of the resonance mass. The existence of *W'* and *Z'* bosons with masses $m_{W'} < 2.67$ TeV and $m_{Z'} < 2.65$ TeV, respectively, are excluded for the HVT benchmark *Model A* with counting constant $a_V = 1.525$ For *Model B* with counting constant the HVT benchmark *Model A* with coupling constant $g_V = 1$ [\[25\]](#page-28-4). For *Model B* with coupling constant $g_V = 3$ [\[25\]](#page-28-4), the corresponding excluded masses are $m_{W'} < 2.82$ TeV and $m_{Z'} < 2.83$ TeV.

To study the scenario in which the masses of charged and neutral resonances are degenerate, a likelihood fit over all the signal regions and control regions is performed. The 95% CL upper limit on the combined signal strength for the processes $W' \to Wh$ and $Z' \to Zh$, assuming $m_{W'} = m_{Z'}$, relative to the HVT model predictions, is shown in Figure [5\(](#page-22-0)c). For *Model A* (*Model B*), $m_{V'}$ < 2.80 TeV (2.93 TeV) is excluded.

The exclusion contours in the HVT parameter space $\{g_Vc_H, (g^2/g_V)c_F\}$ for resonances of mass 1.2 TeV, 2.0 TeV, and 3.0 TeV, are shown in Figure 5(d) where all three lepton channels are combined, taking into 2.0 TeV and 3.0 TeV are shown in Figure [5\(](#page-22-0)d) where all three lepton channels are combined, taking into account the branching fractions to *Wh* and *Zh* from the HVT model prediction. Here, the parameter c_F is assumed to be the same for quarks and leptons, including third-generation fermions, and other parameters involving more than one heavy vector boson, g_Vc_{VVV} , $g_V^2c_{VVVh}$ and c_{VVW} , are assumed to have negligible contributions to the overall cross sections for the processes of interest contributions to the overall cross-sections for the processes of interest.

Figures [6\(](#page-23-0)a) and [6\(](#page-23-0)b) show the 95% CL upper limits on the production cross-section of the *A* boson times its branching fraction to *Zh* and the branching fraction of $h \to b\overline{b}$ as a function of the resonance mass. Upper limits are placed separately for a signal arising from pure gluon–gluon fusion production (Figure [6\(](#page-23-0)a)) and from pure *b*-quark associated production (Figure [6\(](#page-23-0)b)). In the search for the *A* boson with *b*-quark associated production, a mild excess of events is observed around 440 GeV, mainly driven by the dimuon channel in the resolved category with 3+ *b*-tags. The local significance of this excess with respect to the background-only hypothesis is estimated to be 3.6 σ , and the global significance, accounting for the look-elsewhere effect [\[127\]](#page-34-14) is estimated to be 2.4 σ .

The data are also interpreted in terms of limits at 95% CL on the 2HDM parameters tan(β) and cos(β – α). The admixture of gluon–gluon fusion and *^b*-quark associated production, and the variation of the *^A* and *h* boson widths and branching fractions are taken into account according to the predictions of the different models. In this interpretation, the $m_{T,Vh}$ and m_{Vh} distributions of the simulated signal events are

smeared according to a Breit–Wigner function with a width predicted by the parameters of the model. This procedure has been verified to produce the same line-shape as the one including non-resonant and interference effects for widths $\Gamma_A/m_A < 10\%$.

Figure [7](#page-24-0) shows the excluded parameter space for a resonance mass of m_A = 300 GeV in four 2HDM types: I, II, Lepton-Specific, and Flipped. Greater sensitivity is observed at high tan(β) for the Type-II and Flipped models, due to an increased cross-section for *b*-quark associated production. The narrow regions with no exclusion power in Type-I and Type-II at low tan(β) that are far from cos($\beta - \alpha$)=0 are caused by the vanishing branching fraction of $h \to b\overline{b}$.

Figure [8](#page-25-0) shows the parameter exclusion for the four models in the tan(β)– m_A plane for cos($\beta - \alpha$) = 0.1. For the interpretation in Type-II and Flipped 2HDMs, the *b*-quark associated production is included in addition to the gluon–gluon fusion production. The shape of the expected exclusions is determined by the interplay of the expected cross-section limit, which decreases as a function of m_A , and the signal production cross-section times the $A \rightarrow Zh$ branching fraction at a given m_A and tan(β). This branching fraction decreases significantly at $m_A = 350$ GeV due to the opening of the $A \rightarrow t\bar{t}$ channel, but increases again at higher m_A , maintaining similar sensitivity into this m_A region. The variable tan(β) controls the admixture of the gluon–gluon fusion and *b*-quark associated production thereby affecting the rate at which the signal cross-section falls as a function of m_A , which leads to a varying sensitivity as a function of tan (β) . The excesses or deficits in the data visible in Figure [6](#page-23-0) are also reflected in Figure [8.](#page-25-0)

Table 5: The predicted and observed event yields in the signal regions defined in the text. The yields in the 1 and 2 *b*-tag regions correspond to the HVT fit for a signal of mass 1.5 TeV. In the 3+ *b*-tag and 1 and 2 *b*-tag with additional *b*-tags regions, the yields are from the fit using the *A* boson produced in association with *b*-quarks as signal with a mass of 1.5 TeV. The quoted uncertainties are the statistical and systematic uncertainties combined in quadrature after the fit. The uncertainties in the individual background predictions are larger than the total background uncertainty due to correlations in the normalisation parameters in the fit.

		Resolved		Merged				
0-lepton						$1 b$ -tag	$2 b$ -tag	
	1 b -tag	$2 b$ -tag	$3 + b$ -tag	$1 b$ -tag	$2 b$ -tag	add. b-tag	add. b-tag	
$t\bar{t}$	22900 ± 890	6640 ± 180	1000 ± 34	1650 ± 160	68 ± 12	2110 ± 70	105 ± 11	
Single top quark	2440 ± 330	552 ± 76	25.8 ± 5.6	217 ± 52	15.4 ± 4.1	136 ± 50	5.6 ± 2.4	
Diboson	317 ± 41	41.2 ± 5.8	4.5 ± 1.1	188 ± 30	34.8 ± 4.8	12.9 ± 2.3	1.6 ± 0.4	
$Z+l$	580 ± 210	1.3 ± 1.3		310 ± 130	0.38 ± 0.29	11.8 ± 8.2	0.1 ± 0.1	
$Z+(bl, cl)$	8240 ± 840	50 ± 17	5.4 ± 1.8	910 ± 160	10.1 ± 3.7	118 ± 27	0.6 ± 0.4	
$Z+(bb,bc,cc)$	1280 ± 170	1270 ± 150	41 ± 8	238 ± 45	101 ± 16	16.8 ± 4.2	8.6 ± 2.3	
$W+l$	960 ± 300	3 ± 2		227 ± 95	1.0 ± 0.6	5.4 ± 3.9	0.02 ± 0.02	
$W+(bl, cl)$	5960 ± 1100	56 ± 17	3.7 ± 2.3	770 ± 230	6.6 ± 3.2	65 ± 21	0.1 ± 0.1	
$W+(bb,bc,cc)$	530 ± 150	470 ± 130	16.5 ± 4.7	112 ± 44	40 ± 16	10.2 ± 5.1	3 ± 2	
SM Vh	55 ± 21	102 ± 39	1.04 ± 0.57	7.4 ± 2.9	4.7 ± 1.8	0.4 ± 0.2	0.06 ± 0.04	
\bar{t} h	10.4 ± 5.3	7.8 ± 3.9	6 ± 3	1.4 ± 0.7	0.2 ± 0.1	4 ± 2	0.6 ± 0.3	
$t\overline{t}V$	102 ± 54	41 ± 22	8.7 ± 4.5	17.7 ± 9.5	1.4 ± 0.8	24 ± 12	1.8 ± 1.0	
Total	43400 ± 200	9240 ± 95	$\frac{1}{1110} \pm 30$	4650 ± 79	282 ± 14	2510 ± 50	127 ± 11	
Data	43387	9236	1125	4657	283	2516	127	
1-lepton	$1 b$ -tag	$2 b$ -tag		$1 b$ -tag	$2 b$ -tag			
$t\bar{t}$	16300 ± 600	3900 ± 120		8100 ± 300	400 ± 50			
Single top quark	4100 ± 600	860 ± 130		1100 ± 300	120 ± 30			
Diboson	110 ± 20	12 ± 2		220 ± 30	34 ± 5			
$Z+l$	40 ± 10	0.09 ± 0.05		14 ± 6	0.2 ± 0.1			
$Z+(bl, cl)$	170 ± 10	0.7 ± 0.5		38 ± 6	0.4 ± 0.2			
$Z+(bb,bc,cc)$	27 ± 4	17 ± 2		11 ± 2	4.5 ± 0.6			
$W+l$	550 ± 180	3 ± 3		590 ± 230	0.2 ± 0.2			
$W+(bl, cl)$	5700 ± 440	24 ± 8		1800 ± 300	30 ± 10			
$W+(bb,bc,cc)$	820 ± 140	420 ± 70		350 ± 80	180 ± 40			
SM Vh	60 ± 20	90 ± 30		14 ± 6	11 ± 4			
Multijet	200 ± 100	1.7 ± 0.9						
Total	28100 ± 170	5320 ± 70		12200 ± 120	780 ± 30			
Data	28073	5348		12224	775			
2-lepton	$1 b$ -tag	$2 b$ -tag	$3 + b$ -tag	$1 b$ -tag	$2 b$ -tag		$1+2$ <i>b</i> -tag add. <i>b</i> -tag	
$\overline{t}\overline{t}$	2570 ± 80	1940 ± 110	58 ± 9	5.3 ± 2.6	0.4 ± 0.2		11 ± 5	
Single top quark	185 ± 25	58 ± 9	1.5 ± 0.4	0.7 ± 0.1	0.2 ± 0.2	0.5 ± 0.3		
Diboson	570 ± 80	159 ± 24	5.2 ± 1.3	35 ± 5	8.5 ± 1.3	4.6 ± 0.8		
$Z+l$	2210 ± 950	2 ± 3		85 ± 34	1.0 ± 0.5	6 ± 4		
$Z+(bl, cl)$	37200 ± 1100	130 ± 50	12 ± 5	240 ± 40	2.3 ± 0.8	55 ± 11		
$Z+(bb,bc,cc)$	7840 ± 690	6320 ± 170	150 ± 20	74 ± 12	34 ± 5	12 ± 3		
$W\!+\!l$	1.9 ± 0.7			0.03 ± 0.01		0.01 ± 0.01		
$W+(bl, cl)$	37 ± 9	0.9 ± 0.7		0.4 ± 0.1		0.01 ± 0.01		
$W+(bb,bc,cc)$	5.4 ± 1.4	1.9 ± 0.3	0.03 ± 0.01	0.17 ± 0.06	0.02 ± 0.01	0.06 ± 0.05		
SM Vh	105 ± 40	140 ± 60	1.3 ± 0.7	1.6 ± 0.6	0.8 ± 0.3	0.2 ± 0.1		
\bar{t} h	0.9 ± 0.5	1.6 ± 0.8	1.1 ± 0.5	0.05 ± 0.02	0.01 ± 0.01	0.15 ± 0.07		
$t\overline{t}V$	140 ± 80	60 ± 30	6 ± 3	10 ± 5	0.6 ± 0.3	12 ± 6		
Total	50900 ± 230	8810 ± 90	240 ± 20	450 ± 20	47 ± 5		101 ± 9	
Data	50876	8798	235	439	50		101	

Figure 2: Event distributions of $m_{T,Vh}$ for the 0-lepton channel, and m_{Vh} for the 1-lepton and 2-lepton channels in the resolved categories. The quantity on the vertical axis is the number of data events divided by the bin width in GeV. The background prediction is shown after a background-only maximum-likelihood fit to the data. The signal for the benchmark HVT *Model A* with $m_V = 1.5$ TeV is normalised to 10 times the theoretical cross-section. The background uncertainty band shown includes both the statistical and systematic uncertainties after the fit added in quadrature. The lower panels show the ratio of the observed data to the estimated SM background.

Figure 3: Event distributions of $m_{T,Vh}$ for the 0-lepton channel, and m_{Vh} for the 1-lepton and 2-lepton channels in the merged categories. The quantity on the vertical axis is the number of data events divided by the bin width in GeV. The background prediction is shown after a background-only maximum-likelihood fit to the data. The signal for the benchmark HVT *Model A* with $m_V = 1.5$ TeV is normalised to 10 times the theoretical cross-section. The background uncertainty band shown includes both the statistical and systematic uncertainties after the fit added in quadrature. The lower panels show the ratio of the observed data to the estimated SM background.

Figure 4: Event distributions of $m_{T,Vh}$ for the 0-lepton channel and m_{Vh} for the 2-lepton channel after a backgroundonly fit to the categories used in the search for the *A* boson produced in association with *b*-quarks. The quantity on the vertical axis is the number of data events divided by the bin width in GeV. The distribution for an *A* boson with mass of 500 GeV (1.5 TeV) is shown for illustration in the resolved (merged) regions normalised using a cross-section of 5 pb. The background uncertainty band shown includes both the statistical and systematic uncertainties after the fit added in quadrature. The lower panels show the ratio of the observed data to the estimated SM background.

Figure 5: Upper limits as a function of the resonance mass at the 95% CL for (a) the production cross-section of *Z'* times its branching fraction to *Zh* and the branching fraction $B(h \to b\bar{b}, c\bar{c})$ and (b) the production cross-
section of *W'* times its branching fraction to *Wh* and the branching fraction $B(h \to b\bar{b}, c\bar{c})$ (section of *W*^{\prime} times its branching fraction to *Wh* and the branching fraction *B*(*h* $\rightarrow b\bar{b}$, *cc*). (c) Upper limits at the 95% CI for the scaling factor of the production cross-section for *V*^{\prime} times its the 95% CL for the scaling factor of the production cross-section for *V'* times its branching fraction to *Wh*/*Zh* in
Model A. The production cross-sections predicted by *Model A* and *Model B* are shown for comparison *Model A*. The production cross-sections predicted by *Model A* and *Model B* are shown for comparison in (a)–(c). (d) Observed 95% CL exclusion contours in the HVT parameter space $\{g_Vc_H, (g^2/g_V)c_F\}$ for resonances of mass
1.2 TeV, 2.0 TeV, and 3.0 TeV. The areas outside the curves are excluded. Also shown are the benchmark model 1.2 TeV, 2.0 TeV and 3.0 TeV. The areas outside the curves are excluded. Also shown are the benchmark model parameters $A(g_V = 1)$, $A(g_V = 3)$ and $B(g_V = 3)$.

Figure 6: Upper limits at the 95% CL on the product of the production cross-section for $pp \rightarrow A$ and the branching fractions for $A \to Zh$ and $h \to b\bar{b}$ evaluated by combining the 0-lepton and 2-lepton channels. The possible signal components of the data are interpreted assuming (a) pure gluon–gluon fusion production, and (b) pure *b*-quark associated production.

(c) 2HDM Lepton-specific (d) 2HDM Flipped

Figure 7: The interpretation of the cross-section limits in the context of the various 2HDM types as a function of the parameters tan(β) and cos($\beta - \alpha$) for $m_A = 300$ GeV: (a) Type-I, (b) Type-II, (c) Lepton-specific, and (d) Flipped. Variations of the natural width up to $\Gamma_A/m_A=10\%$ have been taken into account. For the interpretation in Type-II and Flipped 2HDM, the *b*-quark associated production is included in addition to the gluon–gluon fusion production.

(c) 2HDM Lepton-specific (d) 2HDM Flipped

Figure 8: The interpretation of the cross-section limits in the context of 2HDMs of Type (a) I , (b) II, (c) Leptonspecific, and (d) Flipped, as a function of the parameters $tan(\beta)$ and m_A for $cos(\beta - \alpha) = 0.1$. Variations of the natural width up to $\Gamma_A/m_A = 10\%$ have been taken into account.

9 Conclusion

A search for *W*^{*'*} and *Z'* bosons and for a CP-odd Higgs boson *A* in the *ν* $\bar{v}b\bar{b}$, $\ell^{\pm}v b\bar{b}$ and $\ell^{\pm} \ell^- b\bar{b}$ final states is performed using 36.1 fb⁻¹ of 13 TeV, np collision data collected with ` is performed using 36.1 fb−¹ of 13 TeV *pp* collision data collected with the ATLAS detector at the LHC. No significant excess of events is observed above the SM predictions in all three channels.

Upper limits are placed at the 95% CL on the cross-section times branching fraction, $\sigma(pp \to V' \to V h) \times R(h \to hh \bar{G})$ ranging between 1.1×10^{-3} and 2.8×10^{-1} pb for the W' boson and between *Vh*) \times *B*(*h* \rightarrow *bb*, *cc*), ranging between 1.1 \times 10⁻³ and 2.8 \times 10⁻¹ pb for the *W*^{*'*} boson and between 0.0 \times 10⁻⁴ and 1.3 \times 10⁻¹ pb for the *Z'* boson in the mass range of 500 GeV to 5 Te 9.0×10^{-4} and 1.3×10^{-1} pb for the *Z'* boson in the mass range of 500 GeV to 5 TeV. The *W'* and *Z'* bosons with masses $m_{\text{max}} \le 2.67$ TeV. (2.82 TeV.) and $m_{\text{max}} \le 2.65$ TeV. (2.83 TeV.) are excluded for *Z*^{\prime} bosons with masses $m_{W'} < 2.67$ TeV (2.82 TeV) and $m_{Z'} < 2.65$ TeV (2.83 TeV) are excluded for the benchmark HVT *Model A (Model B)* while for the combined HVT search masses up to 2.80 TeV the benchmark HVT *Model A (Model B)*, while for the combined HVT search masses up to 2.80 TeV (2.93 TeV) are excluded.

For an *A* boson, upper limits are placed on $\sigma(pp \to A \to Zh) \times B(h \to b\bar{b})$ between 1.9×10^{-3} and 8.1×10^{-1} pb for gluon-gluon fusion production and between 3.4×10^{-3} and 7.3×10^{-1} pb for production 8.1×10^{-1} pb for gluon–gluon fusion production and between 3.4×10^{-3} and 7.3×10^{-1} pb for production
with associated h-quarks in the mass range 220 GeV to 2 TeV with associated *b*-quarks in the mass range 220 GeV to 2 TeV.

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTC, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; CONICYT, Chile; CAS, MOST and NSFC, China; COLCIENCIAS, Colombia; MSMT CR, MPO CR and VSC CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS, CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF, and MPG, Germany; GSRT, Greece; RGC, Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW and NCN, Poland; FCT, Portugal; MNE/IFA, Romania; MES of Russia and NRC KI, Russian Federation; JINR; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DST/NRF, South Africa; MINECO, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TAEK, Turkey; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, CRC and Compute Canada, Canada; COST, ERC, ERDF, Horizon 2020, and Marie Skłodowska-Curie Actions, European Union; Investissements d' Avenir Labex and Idex, ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and GIF, Israel; CERCA Programme Generalitat de Catalunya, Spain; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [\[128\]](#page-34-15).

References

- [1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2012.08.020) 716 (2012) 1, arXiv: [1207.7214 \[hep-ex\]](http://arxiv.org/abs/1207.7214).
- [2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2012.08.021) 716 (2012) 30, arXiv: [1207.7235 \[hep-ex\]](http://arxiv.org/abs/1207.7235).
- [3] ATLAS Collaboration, *Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-015-3685-1) 75 (2015) 476, arXiv: [1506.05669 \[hep-ex\]](http://arxiv.org/abs/1506.05669), Erratum: [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-016-3934-y) 76 (2016) 152.
- [4] CMS Collaboration, *Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at* 7 *and* 8 *TeV*, Phys. Rev. D 92 [\(2015\) 012004,](http://dx.doi.org/10.1103/PhysRevD.92.012004) arXiv: [1411.3441 \[hep-ex\]](http://arxiv.org/abs/1411.3441).
- [5] ATLAS and CMS Collaborations, *Combined measurement of the Higgs boson mass in pp collisions at* [√] *s* = 7 *and* 8 *TeV with the ATLAS and CMS experiments*, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.114.191803) 114 (2015) 191803, arXiv: [1503.07589 \[hep-ex\]](http://arxiv.org/abs/1503.07589).
- [6] S. Weinberg, *Gauge hierarchies*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(79)90248-X) 82 (1979) 387.
- [7] M. J. G. Veltman, *The infrared ultraviolet connection*, Acta Phys. Polon. B 12 (1981) 437.
- [8] C. H. Llewellyn Smith and G. G. Ross, *The real gauge hierarchy problem*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(81)90035-6) 105 (1981) 38.
- [9] LHC Higgs Cross Section Working Group, *Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector*, (2016), arXiv: [1610.07922 \[hep-ph\]](http://arxiv.org/abs/1610.07922).
- [10] F. Sannino and K. Tuominen, *Orientifold theory dynamics and symmetry breaking*, Phys. Rev. D 71 [\(2005\) 051901,](http://dx.doi.org/10.1103/PhysRevD.71.051901) arXiv: [0405209 \[hep-ph\]](http://arxiv.org/abs/0405209).
- [11] R. Foadi, M. T. Frandsen, T. A. Ryttov and F. Sannino, *Minimal walking technicolor: set up for collider physics*, Phys. Rev. D 76 [\(2007\) 055005,](http://dx.doi.org/10.1103/PhysRevD.76.055005) arXiv: [0706.1696 \[hep-ph\]](http://arxiv.org/abs/0706.1696).
- [12] A. Belyaev, R. Foadi, M. T. Frandsen, M. Jarvinen, F. Sannino et al., *Technicolor walks at the LHC*, Phys. Rev. D 79 [\(2009\) 035006,](http://dx.doi.org/10.1103/PhysRevD.79.035006) arXiv: [0809.0793 \[hep-ph\]](http://arxiv.org/abs/0809.0793).
- [13] M. Schmaltz and D. Tucker-Smith, *Little Higgs review*, [Ann. Rev. Nucl. Part. Sci.](http://dx.doi.org/10.1146/annurev.nucl.55.090704.151502) 55 (2005) 229, arXiv: [0502182 \[hep-ph\]](http://arxiv.org/abs/0502182).
- [14] M. J. Dugan, H. Georgi and D. B. Kaplan, *Anatomy of a composite Higgs model*, [Nucl. Phys. B](http://dx.doi.org/10.1016/0550-3213(85)90221-4) 254 (1985) 299.
- [15] K. Agashe, R. Contino and A. Pomarol, *The minimal composite Higgs model*, [Nucl. Phys. B](http://dx.doi.org/10.1016/j.nuclphysb.2005.04.035) 719 (2005) 165, arXiv: [0412089 \[hep-ph\]](http://arxiv.org/abs/0412089).
- [16] G. C. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, [Phys. Rept.](http://dx.doi.org/10.1016/j.physrep.2012.02.002) 516 (2012) 1, arXiv: [1106.0034 \[hep-ph\]](http://arxiv.org/abs/1106.0034).
- [17] P. Fayet, *Supersymmetry and weak, electromagnetic and strong interactions*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(76)90319-1) 64 (1976) 159.
- [18] P. Fayet,
	- *Spontaneously broken supersymmetric theories of weak, electromagnetic and strong interactions*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(77)90852-8) 69 (1977) 489.
- [19] G. R. Farrar and P. Fayet, *Phenomenology of the production, decay, and detection of new hadronic states associated with supersymmetry*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(78)90858-4) 76 (1978) 575.
- [20] P. Fayet, *Relations between the masses of the superpartners of leptons and quarks, the goldstino couplings and the neutral currents*, [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(79)91229-2) 84 (1979) 416.
- [21] S. Dimopoulos and H. Georgi, *Softly broken supersymmetry and SU(5)*, [Nucl. Phys. B](http://dx.doi.org/10.1016/0550-3213(81)90522-8) 193 (1981) 150.
- [22] J. E. Kim, *Light pseudoscalars, particle physics and cosmology*, [Phys. Rept.](http://dx.doi.org/10.1016/0370-1573(87)90017-2) 150 (1987) 1.
- [23] M. Joyce, T. Prokopec and N. Turok, *Nonlocal electroweak baryogenesis. Part 2: The classical regime*, [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.53.2958) 53 (1996) 2958, arXiv: [hep-ph/9410282](http://arxiv.org/abs/hep-ph/9410282).
- [24] D. Alves et al., *Simplified models for LHC new physics searches*, J. Phys. G 39 [\(2012\) 105005,](http://dx.doi.org/10.1088/0954-3899/39/10/105005) arXiv: [1105.2838 \[hep-ph\]](http://arxiv.org/abs/1105.2838).
- [25] D. Pappadopulo, A. Thamm, R. Torre and A. Wulzer, *Heavy vector triplets: bridging theory and data*, JHEP 09 [\(2014\) 060,](http://dx.doi.org/10.1007/JHEP09(2014)060) arXiv: [1402.4431 \[hep-ph\]](http://arxiv.org/abs/1402.4431).
- [26] J. de Blas, J. M. Lizana and M. Perez-Victoria, *Combining searches of Z' and W' bosons*, JHEP 01 [\(2013\) 166,](http://dx.doi.org/10.1007/JHEP01(2013)166) arXiv: [1211.2229 \[hep-ph\]](http://arxiv.org/abs/1211.2229).
- [27] V. D. Barger, W.-Y. Keung and E. Ma, *A gauge model with light W and Z bosons*, [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.22.727) 22 (1980) 727.
- [28] R. Contino, D. Marzocca, D. Pappadopulo and R. Rattazzi, *On the e*ff*ect of resonances in composite Higgs phenomenology*, JHEP 10 [\(2011\) 081,](http://dx.doi.org/10.1007/JHEP10(2011)081) arXiv: [1109.1570 \[hep-ph\]](http://arxiv.org/abs/1109.1570).
- [29] ATLAS Collaboration, *Search for a new resonance decaying to a W or Z boson and a Higgs boson in the* $\ell\ell/\ell\gamma/\gamma + b\bar{b}$ final states with the ATLAS detector, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-015-3474-x) 75 (2015) 263, arXiv: [1503.08089 \[hep-ex\]](http://arxiv.org/abs/1503.08089).
- [30] ATLAS Collaboration, *Search for new resonances decaying to a W or Z boson and a Higgs boson in the* ℓ^+
and $\nu\bar{\nu}b\bar{b}$ channels with np collisions at $\sqrt{s} = 13$ TeV with the ATI AS detector *W* or *Z* boson and a Higgs boson in the $\ell^+ \ell^- b\bar{b}$, $\ell v b\bar{b}$,
 \sqrt{s} = 13 TeV, with the ATI AS detector *and* $v\bar{v}b\bar{b}$ channels with pp collisions at $\sqrt{s} = 13$ *TeV with the ATLAS detector*,
Phys. Lett. B.765 (2017) 32, arXiv: 1607, 05621 [ben-ex] [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2016.11.045) 765 (2017) 32, arXiv: [1607.05621 \[hep-ex\]](http://arxiv.org/abs/1607.05621).
- [31] CMS Collaboration, *Search for heavy resonances decaying into a vector boson and a Higgs boson in final states with charged leptons, neutrinos, and b quarks*, [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2017.02.040) 768 (2017) 137, arXiv: [1610.08066 \[hep-ex\]](http://arxiv.org/abs/1610.08066).
- [32] CMS Collaboration, *Search for a massive resonance decaying into a Higgs boson and a W or Z boson in hadronic final states in proton–proton collisions at* [√] *s* = 8 *TeV*, JHEP 02 [\(2016\) 145,](http://dx.doi.org/10.1007/JHEP02(2016)145) arXiv: [1506.01443 \[hep-ex\]](http://arxiv.org/abs/1506.01443).
- [33] CMS Collaboration, *Search for heavy resonances that decay into a vector boson and a Higgs boson in hadronic final states at* [√] *s* = 13 *TeV*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-017-5192-z) 77 (2017) 636, arXiv: [1707.01303 \[hep-ex\]](http://arxiv.org/abs/1707.01303).
- [34] ATLAS Collaboration, *Search for heavy resonances decaying to a W or Z boson and a Higgs* √ *boson in the* $q\bar{q}^{(\prime)}$ *bb final state in pp collisions at* $\sqrt{s} = 13$ *TeV with the ATLAS detector*, (2017), arXiv: [1707.06958 \[hep-ex\]](http://arxiv.org/abs/1707.06958).
- [35] ALEPH, DELPHI, L3, OPAL Collaborations, *Search for neutral MSSM Higgs bosons at LEP*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s2006-02569-7) 47 (2006) 547, arXiv: [hep-ex/0602042](http://arxiv.org/abs/hep-ex/0602042).
- [36] ATLAS Collaboration, *Search for neutral Higgs bosons of the minimal supersymmetric standard model in pp collisions at* [√] *s* = 8 *TeV with the ATLAS detector*, JHEP 11 [\(2014\) 056,](http://dx.doi.org/10.1007/JHEP11(2014)056) arXiv: [1409.6064 \[hep-ex\]](http://arxiv.org/abs/1409.6064).
- [37] CMS Collaboration, *Search for neutral MSSM Higgs bosons decaying to a pair of tau leptons in pp collisions*, JHEP 10 [\(2014\) 160,](http://dx.doi.org/10.1007/JHEP10(2014)160) arXiv: [1408.3316 \[hep-ex\]](http://arxiv.org/abs/1408.3316).
- [38] ATLAS Collaboration, *Search for a CP-odd Higgs boson decaying to Zh in pp collisions at* √ \sqrt{s} = 8 *TeV with the ATLAS detector*, *[Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2015.03.054)* 744 (2015) 163, arXiv: [1502.04478 \[hep-ex\]](http://arxiv.org/abs/1502.04478).
- [39] CMS Collaboration, *Search for a pseudoscalar boson decaying into a Z boson and the* 125 *GeV Higgs boson in* $\ell^+ \ell^- b\bar{b}$ final states, [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2015.07.010) 748 (2015) 221, 2.15×1504 , 0.4710 [ban-ax] ` arXiv: [1504.04710 \[hep-ex\]](http://arxiv.org/abs/1504.04710).
- [40] ATLAS Collaboration, *The ATLAS experiment at the CERN Large Hadron Collider*, JINST 3 [\(2008\) S08003.](http://dx.doi.org/10.1088/1748-0221/3/08/S08003)
- [41] ATLAS Collaboration, *ATLAS insertable B-layer technical design report*, ATLAS-TDR-19, 2010, url: <https://cds.cern.ch/record/1291633>, *ATLAS insertable B-layer technical design report addendum*, ATLAS-TDR-19-ADD-1, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [42] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-017-4852-3) 77 (2017) 317, arXiv: [1611.09661 \[hep-ex\]](http://arxiv.org/abs/1611.09661).
- [43] J. Alwall et al., *The automated computation of tree-level and next-to-leading order di*ff*erential cross sections, and their matching to parton shower simulations*, JHEP 07 [\(2014\) 079,](http://dx.doi.org/10.1007/JHEP07(2014)079) arXiv: [1405.0301 \[hep-ph\]](http://arxiv.org/abs/1405.0301).
- [44] R. D. Ball et al., *Impact of heavy quark masses on parton distributions and LHC phenomenology*, [Nucl. Phys. B](http://dx.doi.org/10.1016/j.nuclphysb.2011.03.021) 849 (2011) 296, arXiv: [1101.1300 \[hep-ph\]](http://arxiv.org/abs/1101.1300).
- [45] T. Sjöstrand, S. Mrenna and P. Z. Skands, *A brief introduction to PYTHIA 8.1*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2008.01.036) 178 (2008) 852, arXiv: [0710.3820 \[hep-ph\]](http://arxiv.org/abs/0710.3820).
- [46] ATLAS Collaboration, *ATLAS Pythia 8 tunes to* 7 *TeV data*, ATL-PHYS-PUB-2014-021, 2014, url: <https://cds.cern.ch/record/1966419>.
- [47] LHC Higgs Cross Section Working Group, *Handbook of LHC Higgs cross sections: 3. Higgs properties*, (2013), arXiv: [1307.1347 \[hep-ph\]](http://arxiv.org/abs/1307.1347).
- [48] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky et al., *New parton distributions for collider physics*, Phys. Rev. D 82 [\(2010\) 074024,](http://dx.doi.org/10.1103/PhysRevD.82.074024) arXiv: [1007.2241 \[hep-ph\]](http://arxiv.org/abs/1007.2241).
- [49] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2015.01.024) 191 (2015) 159, arXiv: [1410.3012 \[hep-ph\]](http://arxiv.org/abs/1410.3012).
- [50] R. Harlander, S. Liebler and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2013.02.006) 184 (2013) 1605, arXiv: [1212.3249 \[hep-ph\]](http://arxiv.org/abs/1212.3249).
- [51] R. Harlander and P. Kant, *Higgs production and decay: analytic results at next-to-leading order QCD*, JHEP 12 [\(2005\) 015,](http://dx.doi.org/10.1088/1126-6708/2005/12/015) arXiv: [hep-ph/0509189](http://arxiv.org/abs/hep-ph/0509189).
- [52] R. Harlander and W. B. Kilgore, *Higgs boson production in bottom quark fusion at next-to-next-to leading order*, Phys. Rev. D 68 [\(2003\) 013001,](http://dx.doi.org/10.1103/PhysRevD.68.013001) arXiv: [hep-ph/0304035](http://arxiv.org/abs/hep-ph/0304035).
- [53] R. Harlander and W. B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.88.201801) 88 (2002) 201801, arXiv: [hep-ph/0201206](http://arxiv.org/abs/hep-ph/0201206).
- [54] S. Dawson, C. B. Jackson, L. Reina and D. Wackeroth, *Exclusive Higgs boson production with bottom quarks at hadron colliders*, [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.69.074027) 69 (2004) hep, arXiv: [hep-ph/0311067](http://arxiv.org/abs/hep-ph/0311067).
- [55] S. Dittmaier, M. Kramer and M. Spira, *Higgs radiation o*ff *bottom quarks at the Tevatron and the CERN LHC*, [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.70.074010) 70 (2004) hep, arXiv: [hep-ph/0309204](http://arxiv.org/abs/hep-ph/0309204).
- [56] R. Harlander, M. Kramer and M. Schumacher, *Bottom-quark associated Higgs-boson production: reconciling the four- and five-flavour scheme approach*, (2011), arXiv: [1112.3478 \[hep-ph\]](http://arxiv.org/abs/1112.3478).
- [57] D. Eriksson, J. Rathsman and O. Stal, *2HDMC: two-Higgs-doublet model calculator physics and manual*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2009.09.011) 181 (2010) 189, arXiv: [0902.0851 \[hep-ph\]](http://arxiv.org/abs/0902.0851).
- [58] H. E. Haber and O. Stal, *New LHC benchmarks for the* CP *-conserving two-Higgs-doublet model*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-015-3697-x, 10.1140/epjc/s10052-016-4151-4) 75 (2015) 491, Erratum: Eur. Phys. J. C 76 (2016) 312, arXiv: [1507.04281 \[hep-ph\]](http://arxiv.org/abs/1507.04281).
- [59] T. Gleisberg, S. Höche, F. Krauss, M. Schönherr, S. Schumann et al., *Event generation with SHERPA 1.1*, JHEP 02 [\(2009\) 007,](http://dx.doi.org/10.1088/1126-6708/2009/02/007) arXiv: [0811.4622 \[hep-ph\]](http://arxiv.org/abs/0811.4622).
- [60] NNPDF Collaboration: R. D. Ball et al., *Parton distributions for the LHC run 2*, JHEP 04 [\(2015\) 040,](http://dx.doi.org/10.1007/JHEP04(2015)040) arXiv: [1410.8849 \[hep-ph\]](http://arxiv.org/abs/1410.8849).
- [61] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, JHEP 12 [\(2008\) 039,](http://dx.doi.org/10.1088/1126-6708/2008/12/039) arXiv: [0808.3674 \[hep-ph\]](http://arxiv.org/abs/0808.3674).
- [62] F. Cascioli, P. Maierhofer and S. Pozzorini, *Scattering amplitudes with Open Loops*, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.108.111601) 108 (2012) 111601, arXiv: [1111.5206 \[hep-ph\]](http://arxiv.org/abs/1111.5206).
- [63] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements* + *parton showers: the NLO case*, JHEP 04 [\(2013\) 027,](http://dx.doi.org/10.1007/JHEP04(2013)027) arXiv: [1207.5030 \[hep-ph\]](http://arxiv.org/abs/1207.5030).
- [64] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through* $O(\alpha_s^2)$ *,*
Phys. Rev. D.74 (2006) 114017, arXiv: 0609070 [hen-ph] Phys. Rev. D 74 [\(2006\) 114017,](http://dx.doi.org/10.1103/PhysRevD.74.114017) arXiv: [0609070 \[hep-ph\]](http://arxiv.org/abs/0609070).
- [65] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, JHEP 11 [\(2004\) 040,](http://dx.doi.org/10.1088/1126-6708/2004/11/040) arXiv: [0409146 \[hep-ph\]](http://arxiv.org/abs/0409146).
- [66] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, JHEP 11 [\(2007\) 070,](http://dx.doi.org/10.1088/1126-6708/2007/11/070) arXiv: [0709.2092 \[hep-ph\]](http://arxiv.org/abs/0709.2092).
- [67] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, JHEP 06 [\(2010\) 043,](http://dx.doi.org/10.1007/JHEP06(2010)043) arXiv: [1002.2581 \[hep-ph\]](http://arxiv.org/abs/1002.2581).
- [68] T. Sjöstrand, S. Mrenna and P. Z. Skands, *PYTHIA 6.4 physics and manual*, JHEP 05 [\(2006\) 026,](http://dx.doi.org/10.1088/1126-6708/2006/05/026) arXiv: [hep-ph/0603175](http://arxiv.org/abs/hep-ph/0603175).
- [69] P. Z. Skands, *Tuning Monte Carlo generators: the Perugia tunes*, Phys. Rev. D 82 [\(2010\) 074018,](http://dx.doi.org/10.1103/PhysRevD.82.074018) arXiv: [1005.3457 \[hep-ph\]](http://arxiv.org/abs/1005.3457).
- [70] J. Pumplin et al., *New generation of parton distributions with uncertainties from global QCD analysis*, JHEP 07 [\(2002\) 012,](http://dx.doi.org/10.1088/1126-6708/2002/07/012) arXiv: [hep-ph/0201195](http://arxiv.org/abs/hep-ph/0201195).
- [71] M. Beneke, P. Falgari, S. Klein and C. Schwinn, *Hadronic top-quark pair production with NNLL threshold resummation*, [Nucl. Phys. B](http://dx.doi.org/10.1016/j.nuclphysb.2011.10.021) 855 (2012) 695, arXiv: [1109.1536 \[hep-ph\]](http://arxiv.org/abs/1109.1536).
- [72] M. Cacciari, M. Czakon, M. Mangano, A. Mitov and P. Nason, *Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation*, [Phys. Lett. B](http://dx.doi.org/10.1016/j.physletb.2012.03.013) 710 (2012) 612, arXiv: [1111.5869 \[hep-ph\]](http://arxiv.org/abs/1111.5869).
- [73] P. Bärnreuther, M. Czakon and A. Mitov, *Percent level precision physics at the Tevatron: First genuine NNLO QCD corrections to* $q\bar{q} \rightarrow t\bar{t} + X$ *, [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.109.132001)* 109 (2012) 132001, arXiv: [1204.5201 \[hep-ph\]](http://arxiv.org/abs/1204.5201).
- [74] M. Czakon and A. Mitov, *NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels*, JHEP 12 [\(2012\) 054,](http://dx.doi.org/10.1007/JHEP12(2012)054) arXiv: [1207.0236 \[hep-ph\]](http://arxiv.org/abs/1207.0236).
- [75] M. Czakon and A. Mitov, *NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction*, JHEP 01 [\(2013\) 080,](http://dx.doi.org/10.1007/JHEP01(2013)080) arXiv: [1210.6832 \[hep-ph\]](http://arxiv.org/abs/1210.6832).
- [76] M. Czakon, P. Fiedler and A. Mitov, *Total top-quark-pair-production cross section at hadron colliders through* $O(\alpha_S^4)$ *,*
 Shus Bay Latt 110 (2013) 252004 arXiv: 1303 6254 [bon-ph] [Phys. Rev. Lett.](http://dx.doi.org/10.1103/PhysRevLett.110.252004) 110 (2013) 252004, arXiv: [1303.6254 \[hep-ph\]](http://arxiv.org/abs/1303.6254).
- [77] M. Czakon and A. Mitov, *Top*++*: A Program for the Calculation of the Top-Pair Cross-Section at Hadron Colliders*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2014.06.021) 185 (2014) 2930, arXiv: [1112.5675 \[hep-ph\]](http://arxiv.org/abs/1112.5675).
- [78] M. Czakon, D. Heymes and A. Mitov, *Dynamical scales for multi-TeV top-pair production at the LHC*, JHEP 04 [\(2017\) 071,](http://dx.doi.org/10.1007/JHEP04(2017)071) arXiv: [1606.03350 \[hep-ph\]](http://arxiv.org/abs/1606.03350).
- [79] M. Aliev et al., *HATHOR: HAdronic Top and Heavy quarks crOss section calculatoR*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2010.12.040) 182 (2011) 1034, arXiv: [1007.1327 \[hep-ph\]](http://arxiv.org/abs/1007.1327).
- [80] P. Kant et al., *HatHor for single top-quark production: Updated predictions and uncertainty estimates for single top-quark production in hadronic collisions*, [Comput. Phys. Commun.](http://dx.doi.org/10.1016/j.cpc.2015.02.001) 191 (2015) 74, arXiv: [1406.4403 \[hep-ph\]](http://arxiv.org/abs/1406.4403).
- [81] N. Kidonakis, *Two-loop soft anomalous dimensions for single top quark associated production with a W[−] or H[−]*, Phys. Rev. D **82** [\(2010\) 054018,](http://dx.doi.org/10.1103/PhysRevD.82.054018) arXiv: [1005.4451 \[hep-ph\]](http://arxiv.org/abs/1005.4451).
- [82] ATLAS Collaboration, *Measurement of the Z*/γ[∗] *boson transverse momentum distribution in pp collisions at* \sqrt{s} = 7 *TeV with the ATLAS detector*, JHEP 09 [\(2014\) 145,](http://dx.doi.org/10.1007/JHEP09(2014)145) arXiv: [1406.3660 \[hep-ex\]](http://arxiv.org/abs/1406.3660).
- [83] ATLAS Collaboration, *Summary of ATLAS Pythia 8 tunes*, ATL-PHYS-PUB-2012-003, 2012, url: <https://cds.cern.ch/record/1474107>.
- [84] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, *Parton distributions for the LHC*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-009-1072-5) 63 (2009) 189, arXiv: [0901.0002 \[hep-ph\]](http://arxiv.org/abs/0901.0002).
- [85] D. J. Lange, *The EvtGen particle decay simulation package*, [Nucl. Instrum. Meth. A](http://dx.doi.org/10.1016/S0168-9002(01)00089-4) 462 (2001) 152.
- [86] S. Agostinelli et al., *GEANT4 - a simulation toolkit*, [Nucl. Instrum. Meth. A](http://dx.doi.org/10.1016/S0168-9002(03)01368-8) 506 (2003) 250.
- [87] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-010-1429-9) 70 (2010) 823, arXiv: [1005.4568 \[hep-ex\]](http://arxiv.org/abs/1005.4568).
- [88] ATLAS Collaboration, *Electron and photon energy calibration with the ATLAS detector using LHC Run 1 data*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-014-3071-4) 74 (2014) 3071, arXiv: [1407.5063 \[hep-ex\]](http://arxiv.org/abs/1407.5063).
- [89] ATLAS Collaboration, *Electron reconstruction and identification e*ffi*ciency measurements with the ATLAS detector using the 2011 LHC proton–proton collision data*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-014-2941-0) 74 (2014) 2941, arXiv: [1404.2240 \[hep-ex\]](http://arxiv.org/abs/1404.2240).
- [90] ATLAS Collaboration, *Muon reconstruction performance of the ATLAS detector in proton–proton collision data at* [√] *s* = 13 *TeV*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-016-4120-y) 76 (2016) 292, arXiv: [1603.05598 \[hep-ex\]](http://arxiv.org/abs/1603.05598).
- [91] K. Rehermann and B. Tweedie, *Efficient identification of boosted semileptonic top quarks at the LHC*, JHEP 03 [\(2011\) 059,](http://dx.doi.org/10.1007/JHEP03(2011)059) arXiv: [1007.2221 \[hep-ph\]](http://arxiv.org/abs/1007.2221).
- [92] ATLAS Collaboration, *Electron e*ffi*ciency measurements with the ATLAS detector using the 2015 LHC proton–proton collision data*, ATLAS-CONF-2016-024, 2016, url: <https://cds.cern.ch/record/2157687>.
- [93] W. Lampl et al., *Calorimeter Clustering Algorithms: Description and Performance*, ATL-LARG-PUB-2008-002, 2008, url: <https://cds.cern.ch/record/1099735>.
- [94] M. Cacciari, G. P. Salam and G. Soyez, *The anti-k^t jet clustering algorithm*, JHEP 04 [\(2008\) 063,](http://dx.doi.org/10.1088/1126-6708/2008/04/063) arXiv: [0802.1189 \[hep-ph\]](http://arxiv.org/abs/0802.1189).
- [95] M. Cacciari, G. P. Salam and G. Soyez, *FastJet User Manual*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-012-1896-2) 72 (2012) 1896, arXiv: [1111.6097 \[hep-ph\]](http://arxiv.org/abs/1111.6097).
- [96] ATLAS Collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at* √ \sqrt{s} = 8 *TeV* using the ATLAS detector, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-016-4395-z) 76 (2016) 581, arXiv: [1510.03823 \[hep-ex\]](http://arxiv.org/abs/1510.03823).
- [97] D. Krohn, J. Thaler and L.-T. Wang, *Jet trimming*, JHEP 02 [\(2010\) 084,](http://dx.doi.org/10.1007/JHEP02(2010)084) arXiv: [0912.1342 \[hep-ph\]](http://arxiv.org/abs/0912.1342).
- [98] S. Catani, Y. L. Dokshitzer, M. H. Seymour and B. R. Webber, *Longitudinally invariant k*[⊥] *clustering algorithms for hadron hadron collisions*, [Nucl. Phys. B](http://dx.doi.org/10.1016/0550-3213(93)90166-M) 406 (1993) 187.
- [99] S. D. Ellis and D. E. Soper, *Successive combination jet algorithm for hadron collisions*, [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.48.3160) 48 (1993) 3160, arXiv: [hep-ph/9305266](http://arxiv.org/abs/hep-ph/9305266).
- [100] ATLAS Collaboration, *Identification of Boosted, Hadronically-Decaying W and Z Bosons in* √ *s* = 13 *TeV Monte Carlo Simulations for ATLAS*, ATL-PHYS-PUB-2015-033, 2015, url: <https://cds.cern.ch/record/2041461>.
- [101] ATLAS Collaboration, *Jet mass reconstruction with the ATLAS Detector in early Run 2 data*, ATLAS-CONF-2016-035, 2016, url: <https://cds.cern.ch/record/2200211>.
- [102] ATLAS Collaboration, *Jet energy measurement with the ATLAS detector in proton–proton collisions at* [√] *s* = 7 *TeV*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-013-2304-2) 73 (2013) 2304, arXiv: [1112.6426 \[hep-ex\]](http://arxiv.org/abs/1112.6426).
- [103] ATLAS Collaboration, *Jet Calibration and Systematic Uncertainties for Jets Reconstructed in* $\frac{1}{2}$ ATLAS Conaboration, *set Catioration and Systematic Uncertainties for*
the ATLAS Detector at $\sqrt{s} = 13$ *TeV*, ATL-PHYS-PUB-2015-015, 2015, url: <https://cds.cern.ch/record/2037613>.
- [104] ATLAS Collaboration, *Boosted Higgs (*→ *bb) boson identification with the ATLAS detector at* ¯ √ $\bar{s} = 13 \text{ TeV},$ ATLAS-CONF-2016-039, 2016, url: <https://cds.cern.ch/record/2206038>.
- [105] M. Cacciari, G. P. Salam and G. Soyez, *The catchment area of jets*, JHEP 04 [\(2008\) 005,](http://dx.doi.org/10.1088/1126-6708/2008/04/005) arXiv: [0802.1188 \[hep-ph\]](http://arxiv.org/abs/0802.1188).
- [106] ATLAS Collaboration, *Performance of b-jet identification in the ATLAS experiment*, JINST 11 [\(2016\) P04008,](http://dx.doi.org/10.1088/1748-0221/11/04/P04008) arXiv: [1512.01094 \[hep-ex\]](http://arxiv.org/abs/1512.01094).
- [107] ATLAS Collaboration, *Optimisation of the ATLAS b-tagging performance for the 2016 LHC Run*, ATL-PHYS-PUB-2016-012, 2016, url: <https://cds.cern.ch/record/2160731>.
- [108] A. D. Bukin, *Fitting function for asymmetric peaks*, (2007), arXiv: [0711.4449 \[physics.data-an\]](http://arxiv.org/abs/0711.4449).
- [109] ATLAS Collaboration, *Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at* [√] *s* = 8 *TeV*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-015-3500-z) 75 (2015) 303, arXiv: [1412.7086 \[hep-ex\]](http://arxiv.org/abs/1412.7086).
- [110] ATLAS Collaboration, *Performance of algorithms that reconstruct missing transverse momentum in* [√] *s* = 8 *TeV proton–proton collisions in the ATLAS detector*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-017-4780-2) 77 (2017) 241, arXiv: [1609.09324 \[hep-ex\]](http://arxiv.org/abs/1609.09324).
- [111] ATLAS Collaboration, *Expected performance of missing transverse momentum reconstruction for the ATLAS detector at* [√] *s* = 13 *TeV*, ATL-PHYS-PUB-2015-023, 2015, url: <https://cds.cern.ch/record/2037700>.
- [112] ATLAS Collaboration, *Jet global sequential corrections with the ATLAS detector in proton–proton collisions at* [√] *s* = 8 *TeV*, ATLAS-CONF-2015-002, 2015, url: <https://cds.cern.ch/record/2001682>.
- [113] ATLAS Collaboration, *Calibration of b-tagging using dileptonic top pair events in a combinatorial likelihood approach with the ATLAS experiment*, ATLAS-CONF-2014-004, 2014, url: <https://cds.cern.ch/record/1664335>.
- [114] ATLAS Collaboration, *Calibration of the performance of b-tagging for c and light-flavour jets in the 2012 ATLAS data*, ATLAS-CONF-2014-046, 2014, url: <https://cds.cern.ch/record/1741020>.
- [115] ATLAS Collaboration, *Luminosity determination in pp collisions at* [√] *s* = 8 *TeV using the ATLAS detector at the LHC*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-016-4466-1) 76 (2016) 653, arXiv: [1608.03953 \[hep-ex\]](http://arxiv.org/abs/1608.03953).
- [116] ATLAS Collaboration, *Simulation of top-quark production for the ATLAS experiment at* [√] *s* = 13 *TeV*, ATL-PHYS-PUB-2016-004, 2016, url: <https://cds.cern.ch/record/2120417>.
- [117] M. Bahr, S. Gieseke, M. Gigg, D. Grellscheid, K. Hamilton et al., *Herwig*++ *physics and manual*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-008-0798-9) 58 (2008) 639, arXiv: [0803.0883 \[hep-ph\]](http://arxiv.org/abs/0803.0883).
- [118] J. Butterworth et al., *PDF4LHC recommendations for LHC run 2*, J. Phys. G 43 [\(2016\) 023001,](http://dx.doi.org/10.1088/0954-3899/43/2/023001) arXiv: [1510.03865 \[hep-ph\]](http://arxiv.org/abs/1510.03865).
- [119] J. Alwall et al., *The automated computation of tree-level and next-to-leading order di*ff*erential cross sections, and their matching to parton shower simulations*, JHEP 07 [\(2014\) 158,](http://dx.doi.org/10.1007/JHEP07(2014)079) arXiv: [1405.0301 \[hep-ph\]](http://arxiv.org/abs/1405.0301).
- [120] R. D. Ball, V. Bertone, S. Carrazza, C. S. Deans, L. Del Debbio et al., *Parton distributions with LHC data*, [Nucl. Phys. B](http://dx.doi.org/10.1016/j.nuclphysb.2012.10.003) 867 (2013) 244, arXiv: [1207.1303 \[hep-ph\]](http://arxiv.org/abs/1207.1303).
- [121] J. M. Campbell and R. Ellis, *MCFM for the Tevatron and the LHC*, [Nucl. Phys. Proc. Suppl. \(2010\) 205,](http://dx.doi.org/10.1016/j.nuclphysbps.2010.08.011) arXiv: [1007.3492 \[hep-ph\]](http://arxiv.org/abs/1007.3492).
- [122] L. Moneta et al., *The RooStats project*, 2010, arXiv: [1009.1003 \[hep-ph\]](http://arxiv.org/abs/1009.1003).
- [123] W. Verkerke and D. Kirkby, *The RooFit toolkit for data modeling*, 2003, arXiv: [physics/0306116 \[physics.data-an\]](http://arxiv.org/abs/physics/0306116).
- [124] M. Baak et al., *HistFitter software framework for statistical data analysis*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-015-3327-7) 75 (2015) 153, arXiv: [1410.1280 \[hep-ex\]](http://arxiv.org/abs/1410.1280).
- [125] A. L. Read, *Presentation of search results: The CL^s technique*, J. Phys. G 28 [\(2002\) 2693.](http://dx.doi.org/10.1088/0954-3899/28/10/313)
- [126] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-011-1554-0) 71 (2011) 1554, arXiv: [1007.1727 \[physics.data-an\]](http://arxiv.org/abs/1007.1727), Erratum: [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-013-2501-z) 73 (2013) 2501.
- [127] E. Gross and O. Vitells, *Trial factors or the look elsewhere e*ff*ect in high energy physics*, [Eur. Phys. J. C](http://dx.doi.org/10.1140/epjc/s10052-010-1470-8) 70 (2010) 525, arXiv: [1005.1891 \[physics.data-an\]](http://arxiv.org/abs/1005.1891).
- [128] ATLAS Collaboration, *ATLAS Computing Acknowledgements 2016–2017*, ATL-GEN-PUB-2016-002, url: <https://cds.cern.ch/record/2202407>.

The ATLAS Collaboration

M. Aaboud^{34d}, G. Aad⁹⁹, B. Abbott¹²⁵, O. Abdinov^{13,*}, B. Abeloos¹²⁹, S.H. Abidi¹⁶⁵, O.S. AbouZeid¹⁴³, N.L. Abraham¹⁵³, H. Abramowicz¹⁵⁹, H. Abreu¹⁵⁸, R. Abreu¹²⁸, Y. Abulaiti^{43a,43b}, B.S. Acharya^{64a,64b,p}, S. Adachi¹⁶¹, L. Adamczyk^{81a}, J. Adelman¹¹⁹, M. Adersberger¹¹², T. Adye¹⁴¹, A.A. Affolder¹⁴³, Y. Afik¹⁵⁸, T. Agatonovic-Jovin¹⁶, C. Agheorghiesei^{27c}, J.A. Aguilar-Saavedra^{137f,137a,aj}, F. Ahmadov^{77,ah}, G. Aielli^{71a,71b}, S. Akatsuka⁸³, H. Akerstedt^{43a,43b}, T.P.A. Åkesson⁹⁴, E. Akilli⁵², A.V. Akimov¹⁰⁸, G.L. Alberghi^{23b,23a}, J. Albert¹⁷⁴, P. Albicocco⁴⁹, M.J. Alconada Verzini⁸⁶, S. Alderweireldt¹¹⁷, M. Aleksa³⁵, I.N. Aleksandrov⁷⁷, C. Alexa^{27b}, G. Alexander¹⁵⁹, T. Alexopoulos¹⁰, M. Alhroob¹²⁵, B. Ali¹³⁹, G. Alimonti^{66a}, J. Alison³⁶, S.P. Alkire³⁸, B.M.M. Allbrooke¹⁵³, B.W. Allen¹²⁸, P.P. Allport²¹, A. Aloisio^{67a,67b}, A. Alonso³⁹, F. Alonso⁸⁶, C. Alpigiani¹⁴⁵, A.A. Alshehri⁵⁵, M.I. Alstaty⁹⁹, B. Alvarez Gonzalez³⁵, D. Álvarez Piqueras¹⁷², M.G. Alviggi^{67a,67b}, B.T. Amadio¹⁸, Y. Amaral Coutinho^{78b}, C. Amelung²⁶, D. Amidei¹⁰³, S.P. Amor Dos Santos^{137a,137c}, S. Amoroso³⁵, G. Amundsen²⁶, C. Anastopoulos¹⁴⁶, L.S. Ancu⁵², N. Andari²¹, T. Andeen¹¹, C.F. Anders^{59b}, J.K. Anders⁸⁸, K.J. Anderson³⁶, A. Andreazza^{66a,66b}, V. Andrei^{59a}, S. Angelidakis³⁷, I. Angelozzi¹¹⁸, A. Angerami³⁸, A.V. Anisenkov^{120b,120a}, N. Anjos¹⁴, A. Annovi^{69a}, C. Antel^{59a}, M. Antonelli⁴⁹, A. Antonov^{110,*}, D.J.A. Antrim¹⁶⁹, F. Anulli^{70a}, M. Aoki⁷⁹, L. Aperio Bella³⁵, G. Arabidze¹⁰⁴, Y. Arai⁷⁹, J.P. Araque^{137a}, V. Araujo Ferraz^{78b}, A.T.H. Arce⁴⁷, R.E. Ardell⁹¹, F.A. Arduh⁸⁶, J-F. Arguin¹⁰⁷, S. Argyropoulos⁷⁵, M. Arik^{12c}, A.J. Armbruster³⁵, L.J. Armitage⁹⁰, O. Arnaez¹⁶⁵, H. Arnold⁵⁰, M. Arratia³¹, O. Arslan²⁴, A. Artamonov^{109,*}, G. Artoni¹³², S. Artz⁹⁷, S. Asai¹⁶¹, N. Asbah⁴⁴, A. Ashkenazi¹⁵⁹, L. Asquith¹⁵³, K. Assamagan²⁹, R. Astalos^{28a}, M. Atkinson¹⁷¹, N.B. Atlay¹⁴⁸, K. Augsten¹³⁹, G. Avolio³⁵, B. Axen¹⁸, M.K. Ayoub^{15a}, G. Azuelos^{107,ay}, A.E. Baas^{59a}, M.J. Baca²¹, H. Bachacou¹⁴², K. Bachas^{65a,65b}, M. Backes¹³², P. Bagnaia^{70a,70b}, M. Bahmani⁸², H. Bahrasemani¹⁴⁹, J.T. Baines¹⁴¹, M. Bajic³⁹, O.K. Baker¹⁸¹, P.J. Bakker¹¹⁸, E.M. Baldin^{120b,120a}, P. Balek¹⁷⁸, F. Balli¹⁴², W.K. Balunas¹³⁴, E. Banas⁸², A. Bandyopadhyay²⁴, S. Banerjee^{179,1}, A.A.E. Bannoura¹⁸⁰, L. Barak¹⁵⁹, E.L. Barberio¹⁰², D. Barberis^{53b,53a}, M. Barbero⁹⁹, T. Barillari¹¹³, M-S. Barisits³⁵, J. Barkeloo¹²⁸, T. Barklow¹⁵⁰, N. Barlow³¹, S.L. Barnes^{58c}, B.M. Barnett¹⁴¹, R.M. Barnett¹⁸, Z. Barnovska-Blenessy^{58a}, A. Baroncelli^{72a}, G. Barone²⁶, A.J. Barr¹³², L. Barranco Navarro¹⁷², F. Barreiro⁹⁶, J. Barreiro Guimarães da Costa^{15a}, R. Bartoldus¹⁵⁰, A.E. Barton⁸⁷, P. Bartos^{28a}, A. Basalaev¹³⁵, A. Bassalat¹²⁹, R.L. Bates⁵⁵, S.J. Batista¹⁶⁵, J.R. Batley³¹, M. Battaglia¹⁴³, M. Bauce^{70a,70b}, F. Bauer¹⁴², H.S. Bawa^{150,n}, J.B. Beacham¹²³, M.D. Beattie⁸⁷, T. Beau¹³³, P.H. Beauchemin¹⁶⁸, P. Bechtle²⁴, H.C. Beck⁵¹, H.P. Beck^{20,t}, K. Becker¹³², M. Becker⁹⁷, C. Becot¹²², A. Beddall^{12d}, A.J. Beddall^{12a}, V.A. Bednyakov⁷⁷, M. Bedognetti¹¹⁸, C.P. Bee¹⁵², T.A. Beermann³⁵, M. Begalli^{78b}, M. Begel²⁹, J.K. Behr⁴⁴, A.S. Bell⁹², G. Bella¹⁵⁹, L. Bellagamba^{23b}, A. Bellerive³³, M. Bellomo¹⁵⁸, K. Belotskiy¹¹⁰, O. Beltramello³⁵, N.L. Belyaev¹¹⁰, O. Benary^{159,*}, D. Benchekroun^{34a}, M. Bender¹¹², N. Benekos¹⁰, Y. Benhammou¹⁵⁹, E. Benhar Noccioli¹⁸¹, J. Benitez⁷⁵, D.P. Benjamin⁴⁷, M. Benoit⁵², J.R. Bensinger²⁶, S. Bentvelsen¹¹⁸, L. Beresford¹³², M. Beretta⁴⁹, D. Berge¹¹⁸, E. Bergeaas Kuutmann¹⁷⁰, N. Berger⁵, L.J. Bergsten²⁶, J. Beringer¹⁸, S. Berlendis⁵⁶, N.R. Bernard¹⁰⁰, G. Bernardi¹³³, C. Bernius¹⁵⁰, F.U. Bernlochner²⁴, T. Berry⁹¹, P. Berta⁹⁷, C. Bertella^{15a}, G. Bertoli^{43a,43b}, I.A. Bertram⁸⁷, C. Bertsche⁴⁴, G.J. Besjes³⁹, O. Bessidskaia Bylund^{43a,43b}, M. Bessner⁴⁴, N. Besson¹⁴², A. Bethani⁹⁸, S. Bethke¹¹³, A. Betti²⁴, A.J. Bevan⁹⁰, J. Beyer¹¹³, R.M. Bianchi¹³⁶, O. Biebel¹¹², D. Biedermann¹⁹, R. Bielski⁹⁸, K. Bierwagen⁹⁷, N.V. Biesuz^{69a,69b}, M. Biglietti^{72a}, T.R.V. Billoud¹⁰⁷, H. Bilokon⁴⁹, M. Bindi⁵¹, A. Bingul^{12d}, C. Bini^{70a,70b}, S. Biondi^{23b,23a}, T. Bisanz⁵¹, C. Bittrich⁴⁶, D.M. Bjergaard⁴⁷, J.E. Black¹⁵⁰, K.M. Black²⁵, R.E. Blair⁶, T. Blazek^{28a}, I. Bloch⁴⁴, C. Blocker²⁶, A. Blue⁵⁵, U. Blumenschein⁹⁰, Dr. Blunier^{144a}, G.J. Bobbink¹¹⁸, V.S. Bobrovnikov^{120b,120a}, S.S. Bocchetta⁹⁴, A. Bocci⁴⁷, C. Bock¹¹²,

M. Boehler⁵⁰, D. Boerner¹⁸⁰, D. Bogavac¹¹², A.G. Bogdanchikov^{120b,120a}, C. Bohm^{43a}, V. Boisvert⁹¹, P. Bokan¹⁷⁰, T. Bold^{81a}, A.S. Boldyrev¹¹¹, A.E. Bolz^{59b}, M. Bomben¹³³, M. Bona⁹⁰, M. Boonekamp¹⁴², A. Borisov¹²¹, G. Borissov⁸⁷, J. Bortfeldt³⁵, D. Bortoletto¹³², V. Bortolotto^{61a,61b,61c}, D. Boscherini^{23b}, M. Bosman¹⁴, J.D. Bossio Sola³⁰, J. Boudreau¹³⁶, E.V. Bouhova-Thacker⁸⁷, D. Boumediene³⁷, C. Bourdarios¹²⁹, S.K. Boutle⁵⁵, A. Boveia¹²³, J. Boyd³⁵, I.R. Boyko⁷⁷, A.J. Bozson⁹¹, J. Bracinik²¹, A. Brandt⁸, G. Brandt⁵¹, O. Brandt^{59a}, F. Braren⁴⁴, U. Bratzler¹⁶², B. Brau¹⁰⁰, J.E. Brau¹²⁸, W.D. Breaden Madden⁵⁵, K. Brendlinger⁴⁴, A.J. Brennan¹⁰², L. Brenner¹¹⁸, R. Brenner¹⁷⁰, S. Bressler¹⁷⁸, D.L. Briglin²¹, T.M. Bristow⁴⁸, D. Britton⁵⁵, D. Britzger⁴⁴, I. Brock²⁴, R. Brock¹⁰⁴, G. Brooijmans³⁸, T. Brooks⁹¹, W.K. Brooks^{144b}, J. Brosamer¹⁸, E. Brost¹¹⁹, J.H Broughton²¹, P.A. Bruckman de Renstrom⁸², D. Bruncko^{28b}, A. Bruni^{23b}, G. Bruni^{23b}, L.S. Bruni¹¹⁸, S. Bruno^{71a,71b}, B.H. Brunt³¹, M. Bruschi^{23b}, N. Bruscino¹³⁶, P. Bryant³⁶, L. Bryngemark⁴⁴, T. Buanes¹⁷, Q. Buat¹⁴⁹, P. Buchholz 148 , A.G. Buckley 55 , I.A. Budagov 77 , M.K. Bugge 131 , F. Bührer 50 , O. Bulekov 110 , D. Bullock⁸, T.J. Burch¹¹⁹, S. Burdin⁸⁸, C.D. Burgard¹¹⁸, A.M. Burger⁵, B. Burghgrave¹¹⁹, K. Burka⁸², S. Burke¹⁴¹, I. Burmeister⁴⁵, J.T.P. Burr¹³², D. Büscher⁵⁰, V. Büscher⁹⁷, P. Bussey⁵⁵, J.M. Butler²⁵, C.M. Buttar⁵⁵, J.M. Butterworth⁹², P. Butti³⁵, W. Buttinger²⁹, A. Buzatu¹⁵⁵, A.R. Buzykaev^{120b,120a}, S. Cabrera Urbán¹⁷², D. Caforio¹³⁹, H. Cai¹⁷¹, V.M.M. Cairo^{40b,40a}, O. Cakir^{4a}, N. Calace⁵², P. Calafiura¹⁸, A. Calandri⁹⁹, G. Calderini¹³³, P. Calfayan⁶³, G. Callea^{40b,40a}, L.P. Caloba^{78b}, S. Calvente Lopez⁹⁶, D. Calvet³⁷, S. Calvet³⁷, T.P. Calvet⁹⁹, R. Camacho Toro³⁶, S. Camarda³⁵, P. Camarri^{71a,71b}, D. Cameron¹³¹, R. Caminal Armadans¹⁷¹, C. Camincher⁵⁶, S. Campana³⁵, M. Campanelli⁹², A. Camplani^{66a,66b}, A. Campoverde¹⁴⁸, V. Canale^{67a,67b}, M. Cano Bret^{58c}, J. Cantero¹²⁶, T. Cao¹⁵⁹, M.D.M. Capeans Garrido³⁵, I. Caprini^{27b}, M. Caprini^{27b}, M. Capua^{40b,40a}, R.M. Carbone³⁸, R. Cardarelli^{71a}, F.C. Cardillo⁵⁰, I. Carli¹⁴⁰, T. Carli³⁵, G. Carlino^{67a}, B.T. Carlson¹³⁶, L. Carminati^{66a,66b}, R.M.D. Carney^{43a,43b}, S. Caron¹¹⁷, E. Carquin^{144b}, S. Carrá^{66a,66b}, G.D. Carrillo-Montoya³⁵, D. Casadei²¹, M.P. Casado^{14,g}, A.F. Casha¹⁶⁵, M. Casolino¹⁴, D.W. Casper¹⁶⁹, R. Castelijn¹¹⁸, V. Castillo Gimenez¹⁷², N.F. Castro^{137a}, A. Catinaccio³⁵, J.R. Catmore¹³¹, A. Cattai³⁵, J. Caudron²⁴, V. Cavaliere¹⁷¹, E. Cavallaro¹⁴, D. Cavalli^{66a}, M. Cavalli-Sforza¹⁴, V. Cavasinni^{69a,69b}, E. Celebi^{12b}, F. Ceradini^{72a,72b}, L. Cerda Alberich¹⁷², A.S. Cerqueira^{78a}, A. Cerri¹⁵³, L. Cerrito^{71a,71b}, F. Cerutti¹⁸, A. Cervelli^{23b,23a}, S.A. Cetin^{12b}, A. Chafaq^{34a}, D. Chakraborty¹¹⁹, S.K. Chan⁵⁷, W.S. Chan¹¹⁸, Y.L. Chan^{61a}, P. Chang¹⁷¹, J.D. Chapman³¹, D.G. Charlton²¹, C.C. Chau³³, C.A. Chavez Barajas¹⁵³, S. Che¹²³, S. Cheatham^{64a,64c}, A. Chegwidden¹⁰⁴, S. Chekanov⁶, S.V. Chekulaev^{166a}, G.A. Chelkov^{77,ax}, M.A. Chelstowska³⁵, C. Chen^{58a}, C.H. Chen⁷⁶, H. Chen²⁹, J. Chen^{58a}, S. Chen¹⁶¹, S.J. Chen^{15c}, X. Chen^{15b,aw}, Y. Chen⁸⁰, H.C. Cheng¹⁰³, H.J. Cheng^{15d}, A. Cheplakov⁷⁷, E. Cheremushkina¹²¹, R. Cherkaoui El Moursli^{34e}, E. Cheu⁷, K. Cheung⁶², L. Chevalier¹⁴², V. Chiarella⁴⁹, G. Chiarelli^{69a}, G. Chiodini^{65a}, A.S. Chisholm³⁵, A. Chitan^{27b}, Y.H. Chiu¹⁷⁴, M.V. Chizhov⁷⁷, K. Choi⁶³, A.R. Chomont³⁷, S. Chouridou¹⁶⁰, Y.S. Chow^{61a}, V. Christodoulou⁹², M.C. Chu^{61a}, J. Chudoba¹³⁸, A.J. Chuinard¹⁰¹, J.J. Chwastowski⁸², L. Chytka¹²⁷, A.K. Ciftci^{4a}, D. Cinca⁴⁵, V. Cindro⁸⁹, I.A. Cioară²⁴, A. Ciocio¹⁸, F. Cirotto^{67a,67b}, Z.H. Citron¹⁷⁸, M. Citterio^{66a}, M. Ciubancan^{27b}, A. Clark⁵², B.L. Clark⁵⁷, M.R. Clark³⁸, P.J. Clark⁴⁸, R.N. Clarke¹⁸, C. Clement^{43a,43b}, Y. Coadou⁹⁹, M. Cobal^{64a,64c}, A. Coccaro⁵², J. Cochran⁷⁶, L. Colasurdo¹¹⁷, B. Cole³⁸, A.P. Colijn¹¹⁸, J. Collot⁵⁶, T. Colombo¹⁶⁹, P. Conde Muiño^{137a,i}, E. Coniavitis⁵⁰, S.H. Connell^{32b}, I.A. Connelly⁹⁸, S. Constantinescu^{27b}, G. Conti³⁵, F. Conventi^{67a,az}, M. Cooke¹⁸, A.M. Cooper-Sarkar¹³², F. Cormier¹⁷³, K.J.R. Cormier¹⁶⁵, M. Corradi^{70a,70b}, F. Corriveau^{101,af}, A. Cortes-Gonzalez³⁵, G. Costa^{66a}, M.J. Costa¹⁷², D. Costanzo¹⁴⁶, G. Cottin³¹, G. Cowan⁹¹, B.E. Cox⁹⁸, K. Cranmer¹²², S.J. Crawley⁵⁵, R.A. Creager¹³⁴, G. Cree³³, S. Crépé-Renaudin⁵⁶, F. Crescioli¹³³, W.A. Cribbs^{43a,43b}, M. Cristinziani²⁴, V. Croft¹²², G. Crosetti^{40b,40a}, A. Cueto⁹⁶, T. Cuhadar Donszelmann¹⁴⁶, A.R. Cukierman¹⁵⁰, J. Cummings¹⁸¹, M. Curatolo⁴⁹, J. Cúth⁹⁷, S. Czekierda⁸², P. Czodrowski³⁵, M.J. Da Cunha Sargedas De Sousa^{137a,137b}, C. Da Via⁹⁸,

W. Dabrowski^{81a}, T. Dado^{28a,z}, T. Dai¹⁰³, O. Dale¹⁷, F. Dallaire¹⁰⁷, C. Dallapiccola¹⁰⁰, M. Dam³⁹, G. D'amen^{23b,23a}, J.R. Dandoy¹³⁴, M.F. Daneri³⁰, N.P. Dang^{179,1}, A.C. Daniells²¹, N.D Dann⁹⁸, M. Danninger¹⁷³, M. Dano Hoffmann¹⁴², V. Dao¹⁵², G. Darbo^{53b}, S. Darmora⁸, J. Dassoulas³, A. Dattagupta¹²⁸, T. Daubney⁴⁴, S. D'Auria⁵⁵, W. Davey²⁴, C. David⁴⁴, T. Davidek¹⁴⁰, D.R. Davis⁴⁷, P. Davison⁹², E. Dawe¹⁰², I. Dawson¹⁴⁶, K. De⁸, R. De Asmundis^{67a}, A. De Benedetti¹²⁵, S. De Castro^{23b,23a}, S. De Cecco¹³³, N. De Groot¹¹⁷, P. de Jong¹¹⁸, H. De la Torre¹⁰⁴, F. De Lorenzi⁷⁶, A. De Maria^{51,v}, D. De Pedis^{70a}, A. De Salvo^{70a}, U. De Sanctis^{71a,71b}, A. De Santo¹⁵³, K. De Vasconcelos Corga⁹⁹, J.B. De Vivie De Regie¹²⁹, R. Debbe²⁹, C. Debenedetti¹⁴³, D.V. Dedovich⁷⁷, N. Dehghanian³, I. Deigaard¹¹⁸, M. Del Gaudio^{40b,40a}, J. Del Peso⁹⁶, D. Delgove¹²⁹, F. Deliot¹⁴², C.M. Delitzsch⁷, M. Della Pietra^{67a,67b}, D. Della Volpe⁵², A. Dell'Acqua³⁵, L. Dell'Asta²⁵, M. Dell'Orso^{69a,69b}, M. Delmastro⁵, C. Delporte¹²⁹, P.A. Delsart⁵⁶, D.A. DeMarco¹⁶⁵, S. Demers¹⁸¹, M. Demichev⁷⁷, A. Demilly¹³³, S.P. Denisov¹²¹, D. Denysiuk¹⁴², L. D'Eramo¹³³, D. Derendarz⁸², J.E. Derkaoui^{34d}, F. Derue¹³³, P. Dervan⁸⁸, K. Desch²⁴, C. Deterre⁴⁴, K. Dette¹⁶⁵, M.R. Devesa³⁰, P.O. Deviveiros³⁵, A. Dewhurst¹⁴¹, S. Dhaliwal²⁶, F.A. Di Bello⁵², A. Di Ciaccio^{71a,71b}, L. Di Ciaccio⁵, W.K. Di Clemente¹³⁴, C. Di Donato^{67a,67b}, A. Di Girolamo³⁵, B. Di Girolamo³⁵, B. Di Micco^{72a,72b}, R. Di Nardo³⁵, K.F. Di Petrillo⁵⁷, A. Di Simone⁵⁰, R. Di Sipio¹⁶⁵, D. Di Valentino³³, C. Diaconu⁹⁹, M. Diamond¹⁶⁵, F.A. Dias³⁹, M.A. Diaz^{144a}, J. Dickinson¹⁸, E.B. Diehl¹⁰³, J. Dietrich¹⁹, S. Díez Cornell⁴⁴, A. Dimitrievska¹⁶, J. Dingfelder²⁴, P. Dita^{27b}, S. Dita^{27b}, F. Dittus³⁵, F. Djama⁹⁹, T. Djobava^{157b}, J.I. Djuvsland^{59a}, M.A.B. Do Vale^{78c}, D. Dobos³⁵, M. Dobre^{27b}, D. Dodsworth²⁶, C. Doglioni⁹⁴, J. Dolejsi¹⁴⁰, Z. Dolezal¹⁴⁰, M. Donadelli^{78d}, S. Donati^{69a,69b}, P. Dondero^{68a,68b}, J. Donini³⁷, M. D'Onofrio⁸⁸, J. Dopke¹⁴¹, A. Doria^{67a}, M.T. Dova⁸⁶, A.T. Doyle⁵⁵, E. Drechsler⁵¹, M. Dris¹⁰, Y. Du^{58b}, J. Duarte-Campderros¹⁵⁹, F. Dubinin¹⁰⁸, A. Dubreuil⁵², E. Duchovni¹⁷⁸, G. Duckeck¹¹², A. Ducourthial¹³³, O.A. Ducu^{107, y}, D. Duda¹¹⁸, A. Dudarev³⁵, A.C. Dudder⁹⁷, E.M. Duffield¹⁸, L. Duflot¹²⁹, M. Dührssen³⁵, C. Dülsen¹⁸⁰, M. Dumancic¹⁷⁸, A.E. Dumitriu^{27b,e}, A.K. Duncan⁵⁵, M. Dunford^{59a}, A. Duperrin⁹⁹, H. Duran Yildiz^{4a}, M. Düren⁵⁴, A. Durglishvili^{157b}, D. Duschinger⁴⁶, B. Dutta⁴⁴, D. Duvnjak¹, M. Dyndal⁴⁴, B.S. Dziedzic⁸², C. Eckardt⁴⁴, K.M. Ecker¹¹³, R.C. Edgar¹⁰³, T. Eifert³⁵, G. Eigen¹⁷, K. Einsweiler¹⁸, T. Ekelof¹⁷⁰, M. El Kacimi^{34c}, R. El Kosseifi⁹⁹, V. Ellajosyula⁹⁹, M. Ellert¹⁷⁰, S. Elles⁵, F. Ellinghaus¹⁸⁰, A.A. Elliot¹⁷⁴, N. Ellis³⁵, J. Elmsheuser²⁹, M. Elsing³⁵, D. Emeliyanov¹⁴¹, Y. Enari¹⁶¹, J.S. Ennis¹⁷⁶, M.B. Epland⁴⁷, J. Erdmann⁴⁵, A. Ereditato²⁰, M. Ernst²⁹, S. Errede¹⁷¹, M. Escalier¹²⁹, C. Escobar¹⁷², B. Esposito⁴⁹, O. Estrada Pastor¹⁷², A.I. Etienvre¹⁴², E. Etzion¹⁵⁹, H. Evans⁶³, A. Ezhilov¹³⁵, M. Ezzi^{34e}, F. Fabbri^{23b,23a}, L. Fabbri^{23b,23a}, V. Fabiani¹¹⁷, G. Facini⁹², R.M. Fakhrutdinov¹²¹, S. Falciano^{70a}, R.J. Falla⁹², J. Faltova³⁵, Y. Fang^{15a}, M. Fanti^{66a,66b}, A. Farbin⁸, A. Farilla^{72a}, C. Farina¹³⁶, E.M. Farina^{68a,68b}, T. Farooque¹⁰⁴, S. Farrell¹⁸, S.M. Farrington¹⁷⁶, P. Farthouat³⁵, F. Fassi^{34e}, P. Fassnacht³⁵, D. Fassouliotis⁹, M. Faucci Giannelli⁴⁸, A. Favareto^{53b,53a}, W.J. Fawcett¹³², L. Fayard¹²⁹, O.L. Fedin^{135,r}, W. Fedorko¹⁷³, S. Feigl¹³¹, L. Feligioni⁹⁹, C. Feng^{58b}, E.J. Feng³⁵, M.J. Fenton⁵⁵, A.B. Fenyuk¹²¹, L. Feremenga⁸, P. Fernandez Martinez¹⁷², J. Ferrando⁴⁴, A. Ferrari¹⁷⁰, P. Ferrari¹¹⁸, R. Ferrari^{68a}, D.E. Ferreira de Lima^{59b}, A. Ferrer¹⁷², D. Ferrere⁵², C. Ferretti¹⁰³, F. Fiedler⁹⁷, M. Filipuzzi⁴⁴, A. Filipčič⁸⁹, F. Filthaut¹¹⁷, M. Fincke-Keeler¹⁷⁴, K.D. Finelli²⁵, M.C.N. Fiolhais^{137a,137c,b}, L. Fiorini¹⁷², A. Fischer², C. Fischer¹⁴, J. Fischer¹⁸⁰, W.C. Fisher¹⁰⁴, N. Flaschel⁴⁴, I. Fleck¹⁴⁸, P. Fleischmann¹⁰³, R.R.M. Fletcher 134 , T. Flick 180 , B.M. Flierl 112 , L.R. Flores Castillo 61a , M.J. Flowerdew 113 , G.T. Forcolin⁹⁸, A. Formica¹⁴², F.A. Förster¹⁴, A.C. Forti⁹⁸, A.G. Foster²¹, D. Fournier¹²⁹, H. Fox⁸⁷, S. Fracchia¹⁴⁶, P. Francavilla^{69a,69b}, M. Franchini^{23b,23a}, S. Franchino^{59a}, D. Francis³⁵, L. Franconi¹³¹, M. Franklin⁵⁷, M. Frate¹⁶⁹, M. Fraternali^{68a,68b}, D. Freeborn⁹², S.M. Fressard-Batraneanu³⁵, B. Freund¹⁰⁷, D. Froidevaux³⁵, J.A. Frost¹³², C. Fukunaga¹⁶², T. Fusayasu¹¹⁴, J. Fuster¹⁷², O. Gabizon¹⁵⁸, A. Gabrielli^{23b,23a}, A. Gabrielli¹⁸, G.P. Gach^{81a}, S. Gadatsch³⁵, S. Gadomski⁵², G. Gagliardi^{53b,53a}, L.G. Gagnon¹⁰⁷, C. Galea¹¹⁷, B. Galhardo^{137a,137c}, E.J. Gallas¹³², B.J. Gallop¹⁴¹,

P. Gallus¹³⁹, G. Galster³⁹, K.K. Gan¹²³, S. Ganguly³⁷, Y. Gao⁸⁸, Y.S. Gao^{150,n}, C. García¹⁷², J.E. García Navarro¹⁷², J.A. García Pascual^{15a}, M. Garcia-Sciveres¹⁸, R.W. Gardner³⁶, N. Garelli¹⁵⁰, V. Garonne¹³¹, A. Gascon Bravo⁴⁴, K. Gasnikova⁴⁴, C. Gatti⁴⁹, A. Gaudiello^{53b,53a}, G. Gaudio^{68a}, I.L. Gavrilenko¹⁰⁸, C. Gay¹⁷³, G. Gaycken²⁴, E.N. Gazis¹⁰, C.N.P. Gee¹⁴¹, J. Geisen⁵¹, M. Geisen⁹⁷, M.P. Geisler^{59a}, K. Gellerstedt^{43a,43b}, C. Gemme^{53b}, M.H. Genest⁵⁶, C. Geng¹⁰³, S. Gentile^{70a,70b}, C. Gentsos¹⁶⁰, S. George⁹¹, D. Gerbaudo¹⁴, G. Gessner⁴⁵, S. Ghasemi¹⁴⁸, M. Ghneimat²⁴, B. Giacobbe^{23b}, S. Giagu^{70a,70b}, N. Giangiacomi^{23b,23a}, P. Giannetti^{69a}, S.M. Gibson⁹¹, M. Gignac¹⁷³, M. Gilchriese¹⁸, D. Gillberg³³, G. Gilles¹⁸⁰, D.M. Gingrich^{3,ay}, M.P. Giordani^{64a,64c}, F.M. Giorgi^{23b}, P.F. Giraud¹⁴², P. Giromini⁵⁷, G. Giugliarelli^{64a,64c}, D. Giugni^{66a}, F. Giuli¹³², C. Giuliani¹¹³, M. Giulini^{59b}, B.K. Gjelsten¹³¹, S. Gkaitatzis¹⁶⁰, I. Gkialas^{9,k}, E.L. Gkougkousis¹⁴, P. Gkountoumis¹⁰, L.K. Gladilin¹¹¹, C. Glasman⁹⁶, J. Glatzer¹⁴, P.C.F. Glaysher⁴⁴, A. Glazov⁴⁴, M. Goblirsch-Kolb²⁶, J. Godlewski⁸², S. Goldfarb¹⁰², T. Golling⁵², D. Golubkov¹²¹, A. Gomes^{137a,137b}, R. Goncalves Gama^{78b}, J. Goncalves Pinto Firmino Da Costa¹⁴², R. Gonçalo^{137a}, G. Gonella⁵⁰, L. Gonella²¹, A. Gongadze⁷⁷, J.L. Gonski 57 , S. González de la Hoz 172 , S. Gonzalez-Sevilla 52 , L. Goossens 35 , P.A. Gorbounov 109 , H.A. Gordon²⁹, I. Gorelov¹¹⁶, B. Gorini³⁵, E. Gorini^{65a,65b}, A. Gorišek⁸⁹, A.T. Goshaw⁴⁷, C. Gössling⁴⁵, M.I. Gostkin⁷⁷, C.A. Gottardo²⁴, C.R. Goudet¹²⁹, D. Goujdami^{34c}, A.G. Goussiou¹⁴⁵, N. Govender^{32b,c}, E. Gozani¹⁵⁸, I. Grabowska-Bold^{81a}, P.O.J. Gradin¹⁷⁰, J. Gramling¹⁶⁹, E. Gramstad¹³¹, S. Grancagnolo¹⁹, V. Gratchev¹³⁵, P.M. Gravila^{27f}, C. Gray⁵⁵, H.M. Gray¹⁸, Z.D. Greenwood^{93,al}, C. Grefe²⁴, K. Gregersen⁹², I.M. Gregor⁴⁴, P. Grenier¹⁵⁰, K. Grevtsov⁵, J. Griffiths⁸, A.A. Grillo¹⁴³, K. Grimm⁸⁷, S. Grinstein^{14,aa}, Ph. Gris³⁷, J.-F. Grivaz¹²⁹, S. Groh⁹⁷, E. Gross¹⁷⁸, J. Grosse-Knetter⁵¹, G.C. Grossi⁹³, Z.J. Grout⁹², A. Grummer¹¹⁶, L. Guan¹⁰³, W. Guan¹⁷⁹, J. Guenther³⁵, F. Guescini^{166a}, D. Guest¹⁶⁹, O. Gueta¹⁵⁹, B. Gui¹²³, E. Guido^{53b,53a}, T. Guillemin⁵, S. Guindon³⁵, U. Gul⁵⁵, C. Gumpert³⁵, J. Guo^{58c}, W. Guo¹⁰³, Y. Guo^{58a,u}, R. Gupta⁴¹, S. Gurbuz^{12c}, G. Gustavino¹²⁵, B.J. Gutelman¹⁵⁸, P. Gutierrez¹²⁵, N.G. Gutierrez Ortiz⁹², C. Gutschow⁹², C. Guyot¹⁴², M.P. Guzik^{81a}, C. Gwenlan¹³², C.B. Gwilliam⁸⁸, A. Haas¹²², C. Haber¹⁸, H.K. Hadavand⁸, N. Haddad^{34e}, A. Hadef⁹⁹, S. Hageböck²⁴, M. Hagihara¹⁶⁷, H. Hakobyan^{182,*}, M. Haleem⁴⁴, J. Haley¹²⁶, G. Halladjian¹⁰⁴, G.D. Hallewell⁹⁹, K. Hamacher¹⁸⁰, P. Hamal¹²⁷, K. Hamano¹⁷⁴, A. Hamilton^{32a}, G.N. Hamity¹⁴⁶, P.G. Hamnett⁴⁴, L. Han^{58a}, S. Han^{15d}, K. Hanagaki^{79,x}, K. Hanawa¹⁶¹, M. Hance¹⁴³, D.M. Handl¹¹², B. Haney¹³⁴, P. Hanke^{59a}, J.B. Hansen³⁹, J.D. Hansen³⁹, M.C. Hansen²⁴, P.H. Hansen³⁹, K. Hara¹⁶⁷, A.S. Hard¹⁷⁹, T. Harenberg¹⁸⁰, F. Hariri¹²⁹, S. Harkusha¹⁰⁵, P.F. Harrison¹⁷⁶, N.M. Hartmann¹¹², Y. Hasegawa¹⁴⁷, A. Hasib⁴⁸, S. Hassani¹⁴², S. Haug²⁰, R. Hauser¹⁰⁴, L. Hauswald⁴⁶, L.B. Havener³⁸, M. Havranek 139 , C.M. Hawkes 21 , R.J. Hawkings 35 , D. Hayakawa 163 , D. Hayden 104 , C.P. Hays 132 , J.M. Hays 90 , H.S. Hayward 88 , S.J. Haywood 141 , S.J. Head 21 , T. Heck 97 , V. Hedberg 94 , L. Heelan 8 , S. Heer²⁴, K.K. Heidegger⁵⁰, S. Heim⁴⁴, T. Heim¹⁸, B. Heinemann^{44,at}, J.J. Heinrich¹¹², L. Heinrich¹²², C. Heinz⁵⁴, J. Hejbal¹³⁸, L. Helary³⁵, A. Held¹⁷³, S. Hellman^{43a,43b}, C. Helsens³⁵, R.C.W. Henderson⁸⁷, Y. Heng¹⁷⁹, S. Henkelmann¹⁷³, A.M. Henriques Correia³⁵, S. Henrot-Versille¹²⁹, G.H. Herbert¹⁹, H. Herde²⁶, V. Herget¹⁷⁵, Y. Hernández Jiménez^{32c}, H. Herr⁹⁷, G. Herten⁵⁰, R. Hertenberger¹¹², L. Hervas³⁵, T.C. Herwig¹³⁴, G.G. Hesketh⁹², N.P. Hessey^{166a}, J.W. Hetherly⁴¹, S. Higashino⁷⁹, E. Higón-Rodriguez¹⁷², K. Hildebrand³⁶, E. Hill¹⁷⁴, J.C. Hill³¹, K.H. Hiller⁴⁴, S.J. Hillier²¹, M. Hils⁴⁶, I. Hinchliffe¹⁸, M. Hirose⁵⁰, D. Hirschbuehl¹⁸⁰, B. Hiti⁸⁹, O. Hladik¹³⁸, D.R. Hlaluku^{32c}, X. Hoad⁴⁸, J. Hobbs¹⁵², N. Hod^{166a}, M.C. Hodgkinson¹⁴⁶, P. Hodgson¹⁴⁶, A. Hoecker³⁵, M.R. Hoeferkamp¹¹⁶, F. Hoenig¹¹², D. Hohn²⁴, T.R. Holmes³⁶, M. Holzbock¹¹², M. Homann⁴⁵, S. Honda¹⁶⁷, T. Honda⁷⁹, T.M. Hong¹³⁶, A. Hönle¹¹³, B.H. Hooberman¹⁷¹, W.H. Hopkins¹²⁸, Y. Horii¹¹⁵, A.J. Horton¹⁴⁹, J-Y. Hostachy⁵⁶, A. Hostiuc¹⁴⁵, S. Hou¹⁵⁵, A. Hoummada^{34a}, J. Howarth⁹⁸, J. Hoya⁸⁶, M. Hrabovsky¹²⁷, J. Hrdinka 35 , I. Hristova 19 , J. Hrivnac 129 , A. Hrynevich 106 , T. Hryn'ova 5 , P.J. Hsu 62 , S.-C. Hsu 145 , Q. Hu²⁹, S. Hu^{58c}, Y. Huang^{15a}, Z. Hubacek¹³⁹, F. Hubaut⁹⁹, F. Huegging²⁴, T.B. Huffman¹³², E.W. Hughes³⁸, M. Huhtinen³⁵, R.F.H. Hunter³³, P. Huo¹⁵², N. Huseynov^{77,ah}, J. Huston¹⁰⁴, J. Huth⁵⁷,

R. Hyneman¹⁰³, G. Iacobucci⁵², G. Iakovidis²⁹, I. Ibragimov¹⁴⁸, L. Iconomidou-Fayard¹²⁹, Z. Idrissi^{34e}, P. Iengo³⁵, O. Igonkina^{118,ad}, T. Iizawa¹⁷⁷, Y. Ikegami⁷⁹, M. Ikeno⁷⁹, Y. Ilchenko¹¹, D. Iliadis¹⁶⁰, N. Ilic¹⁵⁰, F. Iltzsche⁴⁶, G. Introzzi^{68a,68b}, P. Ioannou^{9,*}, M. Iodice^{72a}, K. Iordanidou³⁸, V. Ippolito⁵⁷, M.F. Isacson¹⁷⁰, N. Ishijima¹³⁰, M. Ishino¹⁶¹, M. Ishitsuka¹⁶³, C. Issever¹³², S. Istin^{12c,aq}, F. Ito¹⁶⁷, J.M. Iturbe Ponce^{61a}, R. Iuppa^{73a,73b}, H. Iwasaki⁷⁹, J.M. Izen⁴², V. Izzo^{67a}, S. Jabbar³, P. Jackson¹, R.M. Jacobs²⁴, V. Jain², G. Jäkel¹⁸⁰, K.B. Jakobi⁹⁷, K. Jakobs⁵⁰, S. Jakobsen⁷⁴, T. Jakoubek¹³⁸, D.O. Jamin¹²⁶, D.K. Jana⁹³, R. Jansky⁵², J. Janssen²⁴, M. Janus⁵¹, P.A. Janus^{81a}, G. Jarlskog⁹⁴, N. Javadov^{77,ah}, T. Javůrek⁵⁰, M. Javurkova⁵⁰, F. Jeanneau¹⁴², L. Jeanty¹⁸, J. Jejelava^{157a,ai}, A. Jelinskas¹⁷⁶, P. Jenni^{50,d}, C. Jeske¹⁷⁶, S. Jézéquel⁵, H. Ji¹⁷⁹, J. Jia¹⁵², H. Jiang⁷⁶, Y. Jiang^{58a}, Z. Jiang^{150,s}, S. Jiggins⁹², J. Jimenez Pena¹⁷², S. Jin^{15c}, A. Jinaru^{27b}, O. Jinnouchi¹⁶³, H. Jivan^{32c}, P. Johansson¹⁴⁶, K.A. Johns⁷, C.A. Johnson⁶³, W.J. Johnson¹⁴⁵, K. Jon-And^{43a,43b}, R.W.L. Jones⁸⁷, S.D. Jones¹⁵³, S. Jones⁷, T.J. Jones⁸⁸, J. Jongmanns^{59a}, P.M. Jorge^{137a,137b}, J. Jovicevic^{166a}, X. Ju¹⁷⁹, A. Juste Rozas^{14,aa}, A. Kaczmarska⁸², M. Kado¹²⁹, H. Kagan¹²³, M. Kagan¹⁵⁰, S.J. Kahn⁹⁹, T. Kaji¹⁷⁷, E. Kajomovitz¹⁵⁸, C.W. Kalderon⁹⁴, A. Kaluza⁹⁷, S. Kama⁴¹, A. Kamenshchikov¹²¹, N. Kanaya¹⁶¹, L. Kanjir⁸⁹, V.A. Kantserov¹¹⁰, J. Kanzaki⁷⁹, B. Kaplan¹²², L.S. Kaplan¹⁷⁹, D. Kar^{32c}, K. Karakostas¹⁰, N. Karastathis¹⁰, M.J. Kareem^{166b}, E. Karentzos¹⁰, S.N. Karpov⁷⁷, Z.M. Karpova⁷⁷, K. Karthik¹²², V. Kartvelishvili⁸⁷, A.N. Karyukhin¹²¹, K. Kasahara¹⁶⁷, L. Kashif¹⁷⁹, R.D. Kass¹²³, A. Kastanas¹⁵¹, Y. Kataoka¹⁶¹, C. Kato¹⁶¹, A. Katre⁵², J. Katzy⁴⁴, K. Kawade⁸⁰, K. Kawagoe⁸⁵, T. Kawamoto¹⁶¹, G. Kawamura⁵¹, E.F. Kay⁸⁸, V.F. Kazanin^{120b,120a}, R. Keeler¹⁷⁴, R. Kehoe⁴¹, J.S. Keller³³, E. Kellermann⁹⁴, J.J. Kempster⁹¹, J. Kendrick²¹, H. Keoshkerian¹⁶⁵, O. Kepka¹³⁸, S. Kersten¹⁸⁰, B.P. Kerševan 89 , R.A. Keyes¹⁰¹, M. Khader¹⁷¹, F. Khalil-Zada¹³, A. Khanov¹²⁶, A.G. Kharlamov^{120b,120a}, T. Kharlamova^{120b,120a}, A. Khodinov¹⁶⁴, T.J. Khoo⁵², V. Khovanskiy^{109,*}, E. Khramov⁷⁷, J. Khubua^{157b}, S. Kido⁸⁰, C.R. Kilby⁹¹, H.Y. Kim⁸, S.H. Kim¹⁶⁷, Y.K. Kim³⁶, N. Kimura¹⁶⁰, O.M. Kind¹⁹, B.T. King⁸⁸, D. Kirchmeier⁴⁶, J. Kirk¹⁴¹, A.E. Kiryunin¹¹³, T. Kishimoto¹⁶¹, D. Kisielewska^{81a}, V. Kitali⁴⁴, O. Kivernyk⁵, E. Kladiva^{28b,*}, T. Klapdor-Kleingrothaus⁵⁰, M.H. Klein¹⁰³, M. Klein⁸⁸, U. Klein⁸⁸, K. Kleinknecht⁹⁷, P. Klimek¹¹⁹, A. Klimentov²⁹, R. Klingenberg^{45,*}, T. Klingl²⁴, T. Klioutchnikova³⁵, F.F. Klitzner¹¹², P. Kluit¹¹⁸, S. Kluth¹¹³, E. Kneringer⁷⁴, E.B.F.G. Knoops⁹⁹, A. Knue¹¹³, A. Kobayashi¹⁶¹, D. Kobayashi⁸⁵, T. Kobayashi¹⁶¹, M. Kobel⁴⁶, M. Kocian¹⁵⁰, P. Kodys¹⁴⁰, T. Koffas³³, E. Koffeman¹¹⁸, M.K. Köhler¹⁷⁸, N.M. Köhler¹¹³, T. Koi¹⁵⁰, M. Kolb^{59b}, I. Koletsou⁵, A.A. Komar^{108,*}, T. Kondo⁷⁹, N. Kondrashova^{58c}, K. Köneke⁵⁰, A.C. König¹¹⁷, T. Kono^{79,as}, R. Konoplich^{122,an}, N. Konstantinidis⁹², B. Konya⁹⁴, R. Kopeliansky⁶³, S. Koperny^{81a}, A.K. Kopp⁵⁰, K. Korcyl⁸², K. Kordas¹⁶⁰, A. Korn⁹², A.A. Korol^{120b,120a,ar}, I. Korolkov¹⁴, E.V. Korolkova¹⁴⁶, O. Kortner¹¹³, S. Kortner¹¹³, T. Kosek¹⁴⁰, V.V. Kostyukhin²⁴, A. Kotwal⁴⁷, A. Koulouris¹⁰, A. Kourkoumeli-Charalampidi^{68a,68b}, C. Kourkoumelis⁹, E. Kourlitis¹⁴⁶, V. Kouskoura²⁹, A.B. Kowalewska⁸², R. Kowalewski¹⁷⁴, T.Z. Kowalski^{81a}, C. Kozakai¹⁶¹, W. Kozanecki¹⁴², A.S. Kozhin¹²¹, V.A. Kramarenko¹¹¹, G. Kramberger⁸⁹, D. Krasnopevtsev¹¹⁰, M.W. Krasny¹³³, A. Krasznahorkay³⁵, D. Krauss¹¹³, J.A. Kremer^{81a}, J. Kretzschmar⁸⁸, K. Kreutzfeldt⁵⁴, P. Krieger¹⁶⁵, K. Krizka¹⁸, K. Kroeninger⁴⁵, H. Kroha¹¹³, J. Kroll¹³⁸, J. Kroll¹³⁴, J. Kroseberg²⁴, J. Krstic¹⁶, U. Kruchonak⁷⁷, H. Krüger²⁴, N. Krumnack⁷⁶, M.C. Kruse⁴⁷, T. Kubota¹⁰², H. Kucuk⁹², S. Kuday^{4b}, J.T. Kuechler¹⁸⁰, S. Kuehn³⁵, A. Kugel^{59a}, F. Kuger¹⁷⁵, T. Kuhl⁴⁴, V. Kukhtin⁷⁷, R. Kukla⁹⁹, Y. Kulchitsky¹⁰⁵, S. Kuleshov^{144b}, Y.P. Kulinich¹⁷¹, M. Kuna^{70a,70b}, T. Kunigo⁸³, A. Kupco¹³⁸, T. Kupfer⁴⁵, O. Kuprash¹⁵⁹, H. Kurashige⁸⁰, L.L. Kurchaninov^{166a}, Y.A. Kurochkin¹⁰⁵, M.G. Kurth^{15d}, E.S. Kuwertz¹⁷⁴, M. Kuze¹⁶³, J. Kvita¹²⁷, T. Kwan¹⁷⁴, D. Kyriazopoulos¹⁴⁶, A. La Rosa¹¹³, J.L. La Rosa Navarro^{78d}, L. La Rotonda^{40b,40a}, F. La Ruffa^{40b,40a}, C. Lacasta¹⁷², F. Lacava^{70a,70b}, J. Lacey⁴⁴, D.P.J. Lack⁹⁸, H. Lacker¹⁹, D. Lacour¹³³, E. Ladygin⁷⁷, R. Lafaye⁵, B. Laforge¹³³, S. Lai⁵¹, S. Lammers⁶³, W. Lampl⁷, E. Lançon²⁹, U. Landgraf⁵⁰, M.P.J. Landon⁹⁰, M.C. Lanfermann⁵², V.S. Lang⁴⁴, J.C. Lange¹⁴, R.J. Langenberg³⁵,

A.J. Lankford¹⁶⁹, F. Lanni²⁹, K. Lantzsch²⁴, A. Lanza^{68a}, A. Lapertosa^{53b,53a}, S. Laplace¹³³, J.F. Laporte¹⁴², T. Lari^{66a}, F. Lasagni Manghi^{23b,23a}, M. Lassnig³⁵, T.S. Lau^{61a}, P. Laurelli⁴⁹, W. Lavrijsen¹⁸, A.T. Law¹⁴³, P. Laycock⁸⁸, T. Lazovich⁵⁷, M. Lazzaroni^{66a,66b}, B. Le¹⁰², O. Le Dortz¹³³, E. Le Guirriec⁹⁹, E.P. Le Quilleuc¹⁴², M. LeBlanc¹⁷⁴, T. LeCompte⁶, F. Ledroit-Guillon⁵⁶, C.A. Lee²⁹, G.R. Lee^{144a}, L. Lee⁵⁷, S.C. Lee¹⁵⁵, B. Lefebvre¹⁰¹, G. Lefebvre¹³³, M. Lefebvre¹⁷⁴, F. Legger¹¹², C. Leggett¹⁸, G. Lehmann Miotto³⁵, X. Lei⁷, W.A. Leight⁴⁴, M.A.L. Leite^{78d}, R. Leitner¹⁴⁰, D. Lellouch¹⁷⁸, B. Lemmer⁵¹, K.J.C. Leney⁹², T. Lenz²⁴, B. Lenzi³⁵, R. Leone⁷, S. Leone^{69a}, C. Leonidopoulos⁴⁸, G. Lerner¹⁵³, C. Leroy¹⁰⁷, R. Les¹⁶⁵, A.A.J. Lesage¹⁴², C.G. Lester³¹, M. Levchenko¹³⁵, J. Levêque⁵, D. Levin¹⁰³, L.J. Levinson¹⁷⁸, M. Levy²¹, D. Lewis⁹⁰, B. Li^{58a,u}, C-Q. Li^{58a,am}, H. Li¹⁵², L. Li^{58c}, Q. Li^{15d}, Q.Y. Li^{58a}, S. Li⁴⁷, X. Li^{58c}, Y. Li¹⁴⁸, Z. Liang^{15a}, B. Liberti^{71a}, A. Liblong¹⁶⁵, K. Lie^{61c}, J. Liebal²⁴, W. Liebig¹⁷, A. Limosani¹⁵⁴, C.Y. Lin³¹, K. Lin¹⁰⁴, S.C. Lin¹⁵⁶, T.H. Lin⁹⁷, R.A. Linck⁶³, B.E. Lindquist¹⁵², A.L. Lionti⁵², E. Lipeles¹³⁴, A. Lipniacka¹⁷, M. Lisovyi^{59b}, T.M. Liss^{171,av}, A. Lister¹⁷³, A.M. Litke¹⁴³, B. Liu⁷⁶, H.B. Liu²⁹, H. Liu¹⁰³, J.B. Liu^{58a}, J.K.K. Liu¹³², J. Liu^{58b}, K. Liu⁹⁹, L. Liu¹⁷¹, M. Liu^{58a}, Y.L. Liu^{58a}, Y.W. Liu^{58a}, M. Livan^{68a,68b}, A. Lleres⁵⁶, J. Llorente Merino^{15a}, S.L. Lloyd⁹⁰, C.Y. Lo^{61b}, F. Lo Sterzo⁴¹, E.M. Lobodzinska⁴⁴, P. Loch⁷, F.K. Loebinger⁹⁸, K.M. Loew²⁶, T. Lohse¹⁹, K. Lohwasser¹⁴⁶, M. Lokajicek¹³⁸, B.A. Long²⁵, J.D. Long¹⁷¹, R.E. Long⁸⁷, L. Longo^{65a,65b}, K.A. Looper¹²³, J.A. Lopez^{144b}, I. Lopez Paz¹⁴, A. Lopez Solis¹³³, J. Lorenz¹¹², N. Lorenzo Martinez⁵, M. Losada²², P.J. Lösel¹¹², A. Lösle⁵⁰, X. Lou^{15a}, A. Lounis¹²⁹, J. Love⁶, P.A. Love⁸⁷, H. Lu^{61a}, N. Lu¹⁰³, Y.J. Lu⁶², H.J. Lubatti¹⁴⁵, C. Luci^{70a,70b}, A. Lucotte⁵⁶, C. Luedtke⁵⁰, F. Luehring⁶³, W. Lukas⁷⁴, L. Luminari^{70a}, O. Lundberg^{43a,43b}, B. Lund-Jensen¹⁵¹, M.S. Lutz¹⁰⁰, P.M. Luzi¹³³, D. Lynn²⁹, R. Lysak¹³⁸, E. Lytken⁹⁴, F. Lyu^{15a}, V. Lyubushkin⁷⁷, H. Ma²⁹, L.L. Ma^{58b}, Y. Ma^{58b}, G. Maccarrone⁴⁹, A. Macchiolo¹¹³, C.M. Macdonald¹⁴⁶, J. Machado Miguens^{134,137b}, D. Madaffari¹⁷², R. Madar³⁷, W.F. Mader⁴⁶, A. Madsen⁴⁴, N. Madysa⁴⁶, J. Maeda⁸⁰, S. Maeland¹⁷, T. Maeno²⁹, A.S. Maevskiy¹¹¹, V. Magerl⁵⁰, C. Maiani¹²⁹, C. Maidantchik^{78b}, T. Maier¹¹², A. Maio^{137a,137b,137d}, O. Majersky^{28a}, S. Majewski¹²⁸, Y. Makida⁷⁹, N. Makovec¹²⁹, B. Malaescu¹³³, Pa. Malecki⁸², V.P. Maleev¹³⁵, F. Malek⁵⁶, U. Mallik⁷⁵, D. Malon⁶, C. Malone³¹, S. Maltezos¹⁰, S. Malyukov³⁵, J. Mamuzic¹⁷², G. Mancini⁴⁹, I. Mandić⁸⁹, J. Maneira^{137a,137b}, L. Manhaes de Andrade Filho^{78a}, J. Manjarres Ramos⁴⁶, K.H. Mankinen⁹⁴, A. Mann¹¹², A. Manousos³⁵, B. Mansoulie¹⁴², J.D. Mansour^{15a}, R. Mantifel¹⁰¹, M. Mantoani⁵¹, S. Manzoni^{66a,66b}, L. Mapelli³⁵, G. Marceca³⁰, L. March⁵², L. Marchese¹³², G. Marchiori¹³³, M. Marcisovsky¹³⁸, C.A. Marin Tobon³⁵, M. Marjanovic³⁷, D.E. Marley¹⁰³, F. Marroquim^{78b}, S.P. Marsden⁹⁸, Z. Marshall¹⁸, M.U.F Martensson¹⁷⁰, S. Marti-Garcia¹⁷², C.B. Martin¹²³, T.A. Martin¹⁷⁶, V.J. Martin⁴⁸, B. Martin dit Latour¹⁷, M. Martinez^{14,aa}, V.I. Martinez Outschoorn¹⁷¹, S. Martin-Haugh¹⁴¹, V.S. Martoiu^{27b}, A.C. Martyniuk⁹², A. Marzin³⁵, L. Masetti⁹⁷, T. Mashimo¹⁶¹, R. Mashinistov¹⁰⁸, J. Masik⁹⁸, A.L. Maslennikov^{120b,120a}, L.H. Mason¹⁰², L. Massa^{71a,71b}, P. Mastrandrea⁵, A. Mastroberardino^{40b,40a}, T. Masubuchi¹⁶¹, P. Mättig¹⁸⁰, J. Maurer^{27b}, B. Maček⁸⁹, S.J. Maxfield⁸⁸, D.A. Maximov^{120b,120a}, R. Mazini¹⁵⁵, I. Maznas¹⁶⁰, S.M. Mazza^{66a,66b}, N.C. Mc Fadden¹¹⁶, G. Mc Goldrick¹⁶⁵, S.P. Mc Kee¹⁰³, A. McCarn¹⁰³, R.L. McCarthy¹⁵², T.G. McCarthy¹¹³, L.I. McClymont⁹², E.F. McDonald¹⁰², J.A. Mcfayden³⁵, G. Mchedlidze⁵¹, S.J. McMahon¹⁴¹, P.C. McNamara¹⁰², C.J. McNicol¹⁷⁶, R.A. McPherson^{174,af}, S. Meehan¹⁴⁵, T.M. Megy⁵⁰, S. Mehlhase¹¹², A. Mehta⁸⁸, T. Meideck⁵⁶, B. Meirose⁴², D. Melini^{172,h}, B.R. Mellado Garcia^{32c}, J.D. Mellenthin⁵¹, M. Melo^{28a}, F. Meloni²⁰, A. Melzer²⁴, S.B. Menary⁹⁸, L. Meng⁸⁸, X.T. Meng¹⁰³, A. Mengarelli^{23b,23a}, S. Menke¹¹³, E. Meoni^{40b,40a}, S. Mergelmeyer¹⁹, C. Merlassino²⁰, P. Mermod⁵², L. Merola^{67a,67b}, C. Meroni^{66a}, F.S. Merritt³⁶, A. Messina^{70a,70b}, J. Metcalfe⁶, A.S. Mete¹⁶⁹, C. Meyer¹³⁴, J. Meyer¹¹⁸, J-P. Meyer¹⁴², H. Meyer Zu Theenhausen^{59a}, F. Miano¹⁵³, R.P. Middleton¹⁴¹, S. Miglioranzi^{53b,53a}, L. Mijović⁴⁸, G. Mikenberg¹⁷⁸, M. Mikestikova 138 , M. Mikuž 89 , M. Milesi 102 , A. Milic 165 , D.A. Millar 90 , D.W. Miller 36 , C. Mills 48 ,

A. Milov¹⁷⁸, D.A. Milstead^{43a,43b}, A.A. Minaenko¹²¹, Y. Minami¹⁶¹, I.A. Minashvili^{157b}, A.I. Mincer¹²², B. Mindur^{81a}, M. Mineev⁷⁷, Y. Minegishi¹⁶¹, Y. Ming¹⁷⁹, L.M. Mir¹⁴, A. Mirto^{65a,65b}, K.P. Mistry¹³⁴, T. Mitani¹⁷⁷, J. Mitrevski¹¹², V.A. Mitsou¹⁷², A. Miucci²⁰, P.S. Miyagawa¹⁴⁶, A. Mizukami⁷⁹, J.U. Mjörnmark 94 , T. Mkrtchyan 182 , M. Mlynarikova 140 , T. Moa 43a,43b , K. Mochizuki 107 , P. Mogg 50 , S. Mohapatra³⁸, S. Molander^{43a,43b}, R. Moles-Valls²⁴, M.C. Mondragon¹⁰⁴, K. Mönig⁴⁴, J. Monk³⁹, E. Monnier⁹⁹, A. Montalbano¹⁵², J. Montejo Berlingen³⁵, F. Monticelli⁸⁶, S. Monzani^{66a}, R.W. Moore³, N. Morange¹²⁹, D. Moreno²², M. Moreno Llácer³⁵, P. Morettini^{53b}, S. Morgenstern³⁵, D. Mori¹⁴⁹, T. Mori¹⁶¹, M. Morii⁵⁷, M. Morinaga¹⁷⁷, V. Morisbak¹³¹, A.K. Morley³⁵, G. Mornacchi³⁵, J.D. Morris⁹⁰, L. Morvaj¹⁵², P. Moschovakos¹⁰, M. Mosidze^{157b}, H.J. Moss¹⁴⁶, J. Moss^{150,o}, K. Motohashi¹⁶³, R. Mount¹⁵⁰, E. Mountricha²⁹, E.J.W. Moyse¹⁰⁰, S. Muanza⁹⁹, F. Mueller¹¹³, J. Mueller¹³⁶, R.S.P. Mueller¹¹², D. Muenstermann⁸⁷, P. Mullen⁵⁵, G.A. Mullier²⁰, F.J. Munoz Sanchez⁹⁸, W.J. Murray^{176,141}, H. Musheghyan³⁵, M. Muškinja⁸⁹, A.G. Myagkov^{121,ao}, M. Myska¹³⁹, B.P. Nachman¹⁸, O. Nackenhorst⁵², K. Nagai¹³², R. Nagai^{79,as}, K. Nagano⁷⁹, Y. Nagasaka⁶⁰, K. Nagata¹⁶⁷, M. Nagel⁵⁰, E. Nagy⁹⁹, A.M. Nairz³⁵, Y. Nakahama¹¹⁵, K. Nakamura⁷⁹, T. Nakamura¹⁶¹, I. Nakano¹²⁴, R.F. Naranjo Garcia⁴⁴, R. Narayan¹¹, D.I. Narrias Villar^{59a}, I. Naryshkin¹³⁵, T. Naumann⁴⁴, G. Navarro²², R. Nayyar⁷, H.A. Neal^{103,*}, P.Y. Nechaeva¹⁰⁸, T.J. Neep¹⁴², A. Negri^{68a,68b}, M. Negrini^{23b}, S. Nektarijevic¹¹⁷, C. Nellist⁵¹, A. Nelson¹⁶⁹, M.E. Nelson¹³², S. Nemecek¹³⁸, P. Nemethy¹²², M. Nessi^{35,f}, M.S. Neubauer¹⁷¹, M. Neumann¹⁸⁰, P.R. Newman²¹, T.Y. Ng^{61c}, Y.S. Ng¹⁹, T. Nguyen Manh¹⁰⁷, R.B. Nickerson¹³², R. Nicolaidou¹⁴², J. Nielsen¹⁴³, N. Nikiforou¹¹, V. Nikolaenko^{121,ao}, I. Nikolic-Audit¹³³, K. Nikolopoulos²¹, P. Nilsson²⁹, Y. Ninomiya⁷⁹, A. Nisati^{70a}, N. Nishu^{58c}, R. Nisius¹¹³, I. Nitsche⁴⁵, T. Nitta¹⁷⁷, T. Nobe¹⁶¹, Y. Noguchi⁸³, M. Nomachi¹³⁰, I. Nomidis³³, M.A. Nomura²⁹, T. Nooney⁹⁰, M. Nordberg³⁵, N. Norjoharuddeen¹³², O. Novgorodova⁴⁶, M. Nozaki⁷⁹, L. Nozka¹²⁷, K. Ntekas¹⁶⁹, E. Nurse⁹², F. Nuti¹⁰², F.G. Oakham^{33,ay}, H. Oberlack¹¹³, T. Obermann²⁴, J. Ocariz¹³³, A. Ochi⁸⁰, I. Ochoa³⁸, J.P. Ochoa-Ricoux^{144a}, K. O'Connor²⁶, S. Oda⁸⁵, S. Odaka⁷⁹, A. Oh⁹⁸, S.H. Oh⁴⁷, C.C. Ohm¹⁵¹, H. Ohman¹⁷⁰, H. Oide^{53b,53a}, H. Okawa¹⁶⁷, Y. Okumura¹⁶¹, T. Okuyama⁷⁹, A. Olariu^{27b}, L.F. Oleiro Seabra^{137a}, S.A. Olivares Pino^{144a}, D. Oliveira Damazio²⁹, M.J.R. Olsson³⁶, A. Olszewski⁸², J. Olszowska⁸², D.C. O'Neil¹⁴⁹, A. Onofre^{137a,137e}, K. Onogi¹¹⁵, P.U.E. Onyisi¹¹, H. Oppen¹³¹, M.J. Oreglia³⁶, Y. Oren¹⁵⁹, D. Orestano^{72a,72b}, N. Orlando^{61b}, A.A. O'Rourke⁴⁴, R.S. Orr¹⁶⁵, B. Osculati^{53b,53a,*}, V. O'Shea⁵⁵, R. Ospanov^{58a}, G. Otero y Garzon³⁰, H. Otono⁸⁵, M. Ouchrif^{34d}, F. Ould-Saada¹³¹, A. Ouraou¹⁴², K.P. Oussoren¹¹⁸, Q. Ouyang^{15a}, M. Owen⁵⁵, R.E. Owen²¹, V.E. Ozcan^{12c}, N. Ozturk⁸, K. Pachal¹⁴⁹, A. Pacheco Pages¹⁴, L. Pacheco Rodriguez¹⁴², C. Padilla Aranda¹⁴, S. Pagan Griso¹⁸, M. Paganini¹⁸¹, F. Paige^{29,*}, G. Palacino⁶³, S. Palazzo^{40b,40a}, S. Palestini³⁵, M. Palka^{81b}, D. Pallin³⁷, E.St. Panagiotopoulou¹⁰, I. Panagoulias¹⁰, C.E. Pandini⁵², J.G. Panduro Vazquez⁹¹, P. Pani³⁵, S. Panitkin²⁹, D. Pantea^{27b}, L. Paolozzi⁵², T.D. Papadopoulou¹⁰, K. Papageorgiou^{9,k}, A. Paramonov⁶, D. Paredes Hernandez¹⁸¹, A.J. Parker⁸⁷, K.A. Parker⁴⁴, M.A. Parker³¹, F. Parodi^{53b,53a}, J.A. Parsons³⁸, U. Parzefall⁵⁰, V.R. Pascuzzi¹⁶⁵, J.M.P. Pasner¹⁴³, E. Pasqualucci^{70a}, S. Passaggio^{53b}, F. Pastore⁹¹, S. Pataraia⁹⁷, J.R. Pater⁹⁸, T. Pauly³⁵, B. Pearson¹¹³, S. Pedraza Lopez¹⁷², R. Pedro^{137a,137b}, S.V. Peleganchuk^{120b,120a}, O. Penc¹³⁸, C. Peng^{15d}, H. Peng^{58a}, J. Penwell⁶³, B.S. Peralva^{78a}, M.M. Perego¹⁴², D.V. Perepelitsa²⁹, F. Peri¹⁹, L. Perini^{66a,66b}, H. Pernegger³⁵, S. Perrella^{67a,67b}, R. Peschke⁴⁴, V.D. Peshekhonov^{77,*}, K. Peters⁴⁴, R.F.Y. Peters⁹⁸, B.A. Petersen³⁵, T.C. Petersen³⁹, E. Petit⁵⁶, A. Petridis¹, C. Petridou¹⁶⁰, P. Petroff¹²⁹, E. Petrolo^{70a}, M. Petrov¹³², F. Petrucci^{72a,72b}, N.E. Pettersson¹⁰⁰, A. Peyaud¹⁴², R. Pezoa^{144b}, F.H. Phillips¹⁰⁴, P.W. Phillips¹⁴¹, G. Piacquadio¹⁵², E. Pianori¹⁷⁶, A. Picazio¹⁰⁰, M.A. Pickering¹³², R. Piegaia³⁰, J.E. Pilcher³⁶, A.D. Pilkington⁹⁸, M. Pinamonti^{71a,71b}, J.L. Pinfold³, H. Pirumov⁴⁴, M. Pitt¹⁷⁸, L. Plazak^{28a}, M.-A. Pleier²⁹, V. Pleskot⁹⁷, E. Plotnikova⁷⁷, D. Pluth⁷⁶, P. Podberezko^{120b,120a}, R. Poettgen⁹⁴, R. Poggi^{68a,68b}, L. Poggioli¹²⁹, I. Pogrebnyak¹⁰⁴, D. Pohl²⁴, I. Pokharel⁵¹, G. Polesello^{68a}, A. Poley⁴⁴, A. Policicchio^{40b,40a}, R. Polifka³⁵, A. Polini^{23b}, C.S. Pollard⁵⁵, V. Polychronakos²⁹, K. Pommès³⁵, D. Ponomarenko¹¹⁰, L. Pontecorvo^{70a}, G.A. Popeneciu^{27d}, D.M. Portillo Quintero¹³³, S. Pospisil¹³⁹, K. Potamianos⁴⁴, I.N. Potrap⁷⁷, C.J. Potter³¹, H. Potti¹¹, T. Poulsen⁹⁴, J. Poveda³⁵, M.E. Pozo Astigarraga³⁵, P. Pralavorio⁹⁹, A. Pranko¹⁸, S. Prell⁷⁶, D. Price⁹⁸, M. Primavera^{65a}, S. Prince¹⁰¹, N. Proklova¹¹⁰, K. Prokofiev^{61c}, F. Prokoshin^{144b}, S. Protopopescu²⁹, J. Proudfoot⁶, M. Przybycien^{81a}, A. Puri¹⁷¹, P. Puzo¹²⁹, J. Qian¹⁰³, G. Qin⁵⁵, Y. Qin⁹⁸, A. Quadt⁵¹, M. Queitsch-Maitland⁴⁴, D. Quilty⁵⁵, S. Raddum¹³¹, V. Radeka²⁹, V. Radescu¹³², S.K. Radhakrishnan¹⁵², P. Radloff¹²⁸, P. Rados¹⁰², F. Ragusa^{66a,66b}, G. Rahal⁹⁵, J.A. Raine⁹⁸, S. Rajagopalan²⁹, C. Rangel-Smith¹⁷⁰, T. Rashid¹²⁹, S. Raspopov⁵, M.G. Ratti^{66a,66b}, D.M. Rauch⁴⁴, F. Rauscher¹¹², S. Rave⁹⁷, I. Ravinovich¹⁷⁸, J.H. Rawling⁹⁸, M. Raymond³⁵, A.L. Read¹³¹, N.P. Readioff⁵⁶, M. Reale^{65a,65b}, D.M. Rebuzzi^{68a,68b}, A. Redelbach¹⁷⁵, G. Redlinger²⁹, R. Reece¹⁴³, R.G. Reed^{32c}, K. Reeves⁴², L. Rehnisch¹⁹, J. Reichert¹³⁴, A. Reiss⁹⁷, C. Rembser³⁵, H. Ren^{15d}, M. Rescigno^{70a}, S. Resconi^{66a}, E.D. Resseguie¹³⁴, S. Rettie¹⁷³, E. Reynolds²¹, O.L. Rezanova^{120b,120a}, P. Reznicek¹⁴⁰, R. Rezvani¹⁰⁷, R. Richter¹¹³, S. Richter⁹², E. Richter-Was^{81b}, O. Ricken²⁴, M. Ridel¹³³, P. Rieck 113 , C.J. Riegel 180 , J. Rieger 51 , O. Rifki 125 , M. Rijssenbeek 152 , A. Rimoldi 68a,68b , M. Rimoldi 20 , L. Rinaldi^{23b}, G. Ripellino¹⁵¹, B. Ristić³⁵, E. Ritsch³⁵, I. Riu¹⁴, F. Rizatdinova¹²⁶, E. Rizvi⁹⁰, C. Rizzi¹⁴, R.T. Roberts⁹⁸, S.H. Robertson^{101,af}, A. Robichaud-Veronneau¹⁰¹, D. Robinson³¹, J.E.M. Robinson⁴⁴, A. Robson⁵⁵, E. Rocco⁹⁷, C. Roda^{69a,69b}, Y. Rodina^{99,ab}, S. Rodriguez Bosca¹⁷², A. Rodriguez Perez¹⁴, D. Rodriguez Rodriguez¹⁷², S. Roe³⁵, C.S. Rogan⁵⁷, O. Røhne¹³¹, J. Roloff⁵⁷, A. Romaniouk¹¹⁰, M. Romano^{23b,23a}, S.M. Romano Saez³⁷, E. Romero Adam¹⁷², N. Rompotis⁸⁸, M. Ronzani⁵⁰, L. Roos¹³³, S. Rosati^{70a}, K. Rosbach⁵⁰, P. Rose¹⁴³, N-A. Rosien⁵¹, E. Rossi^{67a,67b}, L.P. Rossi^{53b}, J.H.N. Rosten³¹, R. Rosten¹⁴⁵, M. Rotaru^{27b}, J. Rothberg¹⁴⁵, D. Rousseau¹²⁹, D. Roy^{32c}, A. Rozanov⁹⁹, Y. Rozen¹⁵⁸, X. Ruan^{32c}, F. Rubbo¹⁵⁰, F. Rühr⁵⁰, A. Ruiz-Martinez³³, Z. Rurikova⁵⁰, N.A. Rusakovich⁷⁷, H.L. Russell¹⁰¹, J.P. Rutherfoord⁷, N. Ruthmann³⁵, E.M. Rüttinger^{44,m}, Y.F. Ryabov¹³⁵, M. Rybar¹⁷¹, G. Rybkin¹²⁹, S. Ryu⁶, A. Ryzhov¹²¹, G.F. Rzehorz⁵¹, A.F. Saavedra¹⁵⁴, G. Sabato¹¹⁸, S. Sacerdoti³⁰, H.F-W. Sadrozinski¹⁴³, R. Sadykov⁷⁷, F. Safai Tehrani^{70a}, P. Saha¹¹⁹, M. Sahinsoy^{59a}, M. Saimpert⁴⁴, M. Saito¹⁶¹, T. Saito¹⁶¹, H. Sakamoto¹⁶¹, Y. Sakurai¹⁷⁷, G. Salamanna^{72a,72b}, J.E. Salazar Loyola^{144b}, D. Salek¹¹⁸, P.H. Sales De Bruin¹⁷⁰, D. Salihagic^{113,*}, A. Salnikov¹⁵⁰, J. Salt¹⁷², D. Salvatore^{40b,40a}, F. Salvatore¹⁵³, A. Salvucci^{61a,61b,61c}, A. Salzburger³⁵, D. Sammel⁵⁰, D. Sampsonidis¹⁶⁰, D. Sampsonidou¹⁶⁰, J. Sánchez¹⁷², V. Sanchez Martinez¹⁷², A. Sanchez Pineda^{64a,64c}, H. Sandaker¹³¹, R.L. Sandbach⁹⁰, C.O. Sander⁴⁴, M. Sandhoff¹⁸⁰, C. Sandoval²², D.P.C. Sankey¹⁴¹, M. Sannino^{53b,53a}, Y. Sano¹¹⁵, A. Sansoni⁴⁹, C. Santoni³⁷, H. Santos^{137a}, I. Santoyo Castillo¹⁵³, A. Sapronov⁷⁷, J.G. Saraiva^{137a,137d}, B. Sarrazin²⁴, O. Sasaki⁷⁹, K. Sato¹⁶⁷, E. Sauvan⁵, G. Savage⁹¹, P. Savard^{165,ay}, N. Savic¹¹³, C. Sawyer¹⁴¹, L. Sawyer^{93,al}, J. Saxon³⁶, C. Sbarra^{23b}, A. Sbrizzi^{23a}, T. Scanlon⁹², D.A. Scannicchio¹⁶⁹, J. Schaarschmidt¹⁴⁵, P. Schacht¹¹³, B.M. Schachtner¹¹², D. Schaefer³⁶, L. Schaefer¹³⁴, R. Schaefer⁴⁴, J. Schaeffer⁹⁷, S. Schaepe³⁵, S. Schaetzel^{59b}, U. Schäfer⁹⁷, A.C. Schaffer¹²⁹, D. Schaile¹¹², R.D. Schamberger¹⁵², V.A. Schegelsky¹³⁵, D. Scheirich¹⁴⁰, F. Schenck¹⁹, M. Schernau¹⁶⁹, C. Schiavi^{53b,53a}, S. Schier¹⁴³, L.K. Schildgen²⁴, C. Schillo⁵⁰, M. Schioppa^{40b,40a}, S. Schlenker³⁵, K.R. Schmidt-Sommerfeld¹¹³, K. Schmieden³⁵, C. Schmitt⁹⁷, S. Schmitt⁴⁴, S. Schmitz⁹⁷, U. Schnoor⁵⁰, L. Schoeffel¹⁴², A. Schoening^{59b}, B.D. Schoenrock¹⁰⁴, E. Schopf²⁴, M. Schott⁹⁷, J.F.P. Schouwenberg¹¹⁷, J. Schovancova³⁵, S. Schramm⁵², N. Schuh⁹⁷, A. Schulte⁹⁷, M.J. Schultens²⁴, H-C. Schultz-Coulon^{59a}, H. Schulz¹⁹, M. Schumacher⁵⁰, B.A. Schumm¹⁴³, Ph. Schune¹⁴², A. Schwartzman¹⁵⁰, T.A. Schwarz¹⁰³, H. Schweiger⁹⁸, Ph. Schwemling¹⁴², R. Schwienhorst¹⁰⁴, A. Sciandra²⁴, G. Sciolla²⁶, M. Scornajenghi^{40b,40a}, F. Scuri^{69a}, F. Scutti¹⁰², J. Searcy¹⁰³, P. Seema²⁴, S.C. Seidel¹¹⁶, A. Seiden¹⁴³, J.M. Seixas^{78b}, G. Sekhniaidze^{67a}, K. Sekhon¹⁰³, S.J. Sekula⁴¹, N. Semprini-Cesari^{23b,23a}, S. Senkin³⁷, C. Serfon¹³¹, L. Serin¹²⁹, L. Serkin^{64a,64b}, M. Sessa^{72a,72b}, R. Seuster¹⁷⁴, H. Severini¹²⁵, F. Sforza¹⁶⁸, A. Sfyrla⁵², E. Shabalina⁵¹,

N.W. Shaikh^{43a,43b}, L.Y. Shan^{15a}, R. Shang¹⁷¹, J.T. Shank²⁵, M. Shapiro¹⁸, P.B. Shatalov¹⁰⁹, K. Shaw^{64a,64b}, S.M. Shaw⁹⁸, A. Shcherbakova^{43a,43b}, C.Y. Shehu¹⁵³, Y. Shen¹²⁵, N. Sherafati³³, A.D. Sherman²⁵, P. Sherwood⁹², L. Shi^{155,au}, S. Shimizu⁸⁰, C.O. Shimmin¹⁸¹, M. Shimojima¹¹⁴, I.P.J. Shipsey¹³², S. Shirabe⁸⁵, M. Shiyakova⁷⁷, J. Shlomi¹⁷⁸, A. Shmeleva¹⁰⁸, D. Shoaleh Saadi¹⁰⁷, M.J. Shochet³⁶, S. Shojaii¹⁰², D.R. Shope¹²⁵, S. Shrestha¹²³, E. Shulga¹¹⁰, M.A. Shupe⁷, P. Sicho¹³⁸, A.M. Sickles¹⁷¹, P.E. Sidebo¹⁵¹, E. Sideras Haddad^{32c}, O. Sidiropoulou¹⁷⁵, A. Sidoti^{23b,23a}, F. Siegert⁴⁶, Dj. Sijacki¹⁶, J. Silva^{137a,137d}, S.B. Silverstein^{43a}, V. Simak¹³⁹, L. Simic⁷⁷, S. Simion¹²⁹, E. Simioni⁹⁷, B. Simmons⁹², M. Simon⁹⁷, P. Sinervo¹⁶⁵, N.B. Sinev¹²⁸, M. Sioli^{23b,23a}, G. Siragusa¹⁷⁵, I. Siral¹⁰³, S.Yu. Sivoklokov¹¹¹, J. Sjölin^{43a,43b}, M.B. Skinner⁸⁷, P. Skubic¹²⁵, M. Slater²¹, T. Slavicek¹³⁹, M. Slawinska⁸², K. Sliwa¹⁶⁸, R. Slovak¹⁴⁰, V. Smakhtin¹⁷⁸, B.H. Smart⁵, J. Smiesko^{28a}, N. Smirnov¹¹⁰, S.Yu. Smirnov¹¹⁰, Y. Smirnov¹¹⁰, L.N. Smirnova¹¹¹, O. Smirnova⁹⁴, J.W. Smith⁵¹, M.N.K. Smith³⁸, R.W. Smith³⁸, M. Smizanska⁸⁷, K. Smolek¹³⁹, A.A. Snesarev¹⁰⁸, I.M. Snyder¹²⁸, S. Snyder²⁹, R. Sobie^{174,af}, F. Socher⁴⁶, A. Soffer¹⁵⁹, A. Søgaard⁴⁸, D.A. Soh¹⁵⁵, G. Sokhrannyi⁸⁹, C.A. Solans Sanchez³⁵, M. Solar¹³⁹, E.Yu. Soldatov¹¹⁰, U. Soldevila¹⁷², A.A. Solodkov¹²¹, A. Soloshenko⁷⁷, O.V. Solovyanov¹²¹, V. Solovyev¹³⁵, P. Sommer¹⁴⁶, H. Son¹⁶⁸, A. Sopczak¹³⁹, D. Sosa^{59b}, C.L. Sotiropoulou^{69a,69b}, S. Sottocornola^{68a,68b}, R. Soualah^{64a,64c,j}, A.M. Soukharev^{120b,120a}, D. South⁴⁴, B.C. Sowden⁹¹, S. Spagnolo^{65a,65b}, M. Spalla^{69a,69b}, M. Spangenberg¹⁷⁶, F. Spanò⁹¹, D. Sperlich¹⁹, F. Spettel¹¹³, T.M. Spieker^{59a}, R. Spighi^{23b}, G. Spigo³⁵, L.A. Spiller¹⁰², M. Spousta¹⁴⁰, R.D. St. Denis^{55,*}, A. Stabile^{66a,66b}, R. Stamen^{59a}, S. Stamm¹⁹, E. Stanecka⁸², R.W. Stanek⁶, C. Stanescu^{72a}, M.M. Stanitzki⁴⁴, B. Stapf¹¹⁸, S. Stapnes¹³¹, E.A. Starchenko¹²¹, G.H. Stark³⁶, J. Stark⁵⁶, S.H Stark³⁹, P. Staroba¹³⁸, P. Starovoitov^{59a}, S. Stärz³⁵, R. Staszewski⁸², M. Stegler⁴⁴, P. Steinberg²⁹, B. Stelzer¹⁴⁹, H.J. Stelzer³⁵, O. Stelzer-Chilton^{166a}, H. Stenzel⁵⁴, T.J. Stevenson⁹⁰, G.A. Stewart⁵⁵, M.C. Stockton¹²⁸, M. Stoebe¹⁰¹, G. Stoicea^{27b}, P. Stolte⁵¹, S. Stonjek¹¹³, A.R. Stradling⁸, A. Straessner⁴⁶, M.E. Stramaglia²⁰, J. Strandberg¹⁵¹, S. Strandberg^{43a,43b}, M. Strauss¹²⁵, P. Strizenec^{28b}, R. Ströhmer¹⁷⁵, D.M. Strom¹²⁸, R. Stroynowski⁴¹, A. Strubig⁴⁸, S.A. Stucci²⁹, B. Stugu¹⁷, N.A. Styles⁴⁴, D. Su¹⁵⁰, J. Su¹³⁶, S. Suchek^{59a}, Y. Sugaya¹³⁰, M. Suk¹³⁹, V.V. Sulin¹⁰⁸, D.M.S. Sultan^{73a,73b}, S. Sultansoy^{4c}, T. Sumida⁸³, S. Sun⁵⁷, X. Sun³, K. Suruliz¹⁵³, C.J.E. Suster¹⁵⁴, M.R. Sutton¹⁵³, S. Suzuki⁷⁹, M. Svatos¹³⁸, M. Swiatlowski³⁶, S.P. Swift², I. Sykora^{28a}, T. Sykora¹⁴⁰, D. Ta⁵⁰, K. Tackmann^{44,ac}, J. Taenzer¹⁵⁹, A. Taffard¹⁶⁹, R. Tafirout^{166a}, E. Tahirovic⁹⁰, N. Taiblum 159 , H. Takai 29 , R. Takashima 84 , E.H. Takasugi 113 , K. Takeda 80 , T. Takeshita 147 , Y. Takubo 79 , M. Talby 99 , A.A. Talyshev 120b,120a , J. Tanaka 161 , M. Tanaka 163 , R. Tanaka 129 , S. Tanaka 79 , R. Tanioka 80 , B.B. Tannenwald¹²³, S. Tapia Araya^{144b}, S. Tapprogge⁹⁷, S. Tarem¹⁵⁸, G.F. Tartarelli^{66a}, P. Tas¹⁴⁰, M. Tasevsky¹³⁸, T. Tashiro⁸³, E. Tassi^{40b,40a}, A. Tavares Delgado^{137a,137b}, Y. Tayalati^{34e}, A.C. Taylor¹¹⁶, A.J. Taylor⁴⁸, G.N. Taylor¹⁰², P.T.E. Taylor¹⁰², W. Taylor^{166b}, P. Teixeira-Dias⁹¹, D. Temple¹⁴⁹, H. Ten Kate³⁵, P.K. Teng¹⁵⁵, J.J. Teoh¹³⁰, F. Tepel¹⁸⁰, S. Terada⁷⁹, K. Terashi¹⁶¹, J. Terron⁹⁶, S. Terzo¹⁴, M. Testa⁴⁹, R.J. Teuscher^{165,af}, S.J. Thais¹⁸¹, T. Theveneaux-Pelzer⁹⁹, F. Thiele³⁹, J.P. Thomas²¹, J. Thomas-Wilsker⁹¹, A.S. Thompson⁵⁵, P.D. Thompson²¹, L.A. Thomsen¹⁸¹, E. Thomson¹³⁴, Y. Tian³⁸, M.J. Tibbetts¹⁸, R.E. Ticse Torres⁵¹, V.O. Tikhomirov^{108,ap}, Yu.A. Tikhonov^{120b,120a}, S. Timoshenko¹¹⁰, P. Tipton¹⁸¹, S. Tisserant⁹⁹, K. Todome¹⁶³, S. Todorova-Nova⁵, S. Todt⁴⁶, J. Tojo⁸⁵, S. Tokár^{28a}, K. Tokushuku⁷⁹, E. Tolley¹²³, L. Tomlinson⁹⁸, M. Tomoto¹¹⁵, L. Tompkins^{150,s}, K. Toms¹¹⁶, B. Tong⁵⁷, P. Tornambe⁵⁰, E. Torrence¹²⁸, H. Torres⁴⁶, E. Torró Pastor¹⁴⁵, J. Toth^{99,ae}, F. Touchard⁹⁹, D.R. Tovey¹⁴⁶, C.J. Treado¹²², T. Trefzger¹⁷⁵, F. Tresoldi¹⁵³, A. Tricoli²⁹, I.M. Trigger^{166a}, S. Trincaz-Duvoid¹³³, M.F. Tripiana¹⁴, W. Trischuk¹⁶⁵, B. Trocmé⁵⁶, A. Trofymov⁴⁴, C. Troncon^{66a}, M. Trottier-McDonald¹⁸, M. Trovatelli¹⁷⁴, L. Truong^{32b}, M. Trzebinski⁸², A. Trzupek⁸², K.W. Tsang^{61a}, J.C-L. Tseng¹³², P.V. Tsiareshka¹⁰⁵, G. Tsipolitis¹⁰, N. Tsirintanis⁹, S. Tsiskaridze¹⁴, V. Tsiskaridze⁵⁰, E.G. Tskhadadze^{157a}, I.I. Tsukerman¹⁰⁹, V. Tsulaia¹⁸, S. Tsuno⁷⁹, D. Tsybychev¹⁵², Y. Tu^{61b}, A. Tudorache^{27b}, V. Tudorache^{27b}, T.T. Tulbure^{27a}, A.N. Tuna⁵⁷, S. Turchikhin⁷⁷, D. Turgeman¹⁷⁸,

I. Turk Cakir^{4b,w}, R.T. Turra^{66a}, P.M. Tuts³⁸, G. Ucchielli^{23b,23a}, I. Ueda⁷⁹, M. Ughetto^{43a,43b}, F. Ukegawa¹⁶⁷, G. Unal³⁵, A. Undrus²⁹, G. Unel¹⁶⁹, F.C. Ungaro¹⁰², Y. Unno⁷⁹, K. Uno¹⁶¹, C. Unverdorben¹¹², J. Urban^{28b}, P. Urquijo¹⁰², P. Urrejola⁹⁷, G. Usai⁸, J. Usui⁷⁹, L. Vacavant⁹⁹, V. Vacek¹³⁹, B. Vachon¹⁰¹, K.O.H. Vadla¹³¹, A. Vaidya⁹², C. Valderanis¹¹², E. Valdes Santurio^{43a,43b}, M. Valente⁵², S. Valentinetti^{23b,23a}, A. Valero¹⁷², L. Valéry¹⁴, S. Valkar¹⁴⁰, A. Vallier⁵, J.A. Valls Ferrer¹⁷², W. Van Den Wollenberg¹¹⁸, H. Van der Graaf¹¹⁸, P. Van Gemmeren⁶, J. Van Nieuwkoop¹⁴⁹, I. Van Vulpen¹¹⁸, M.C. van Woerden¹¹⁸, M. Vanadia^{71a,71b}, W. Vandelli³⁵, A. Vaniachine¹⁶⁴, P. Vankov¹¹⁸, G. Vardanyan¹⁸², R. Vari^{70a}, E.W. Varnes⁷, C. Varni^{53b,53a}, T. Varol⁴¹, D. Varouchas¹²⁹, A. Vartapetian⁸, K.E. Varvell¹⁵⁴, G.A. Vasquez^{144b}, J.G. Vasquez¹⁸¹, F. Vazeille³⁷, D. Vazquez Furelos¹⁴, T. Vazquez Schroeder¹⁰¹, J. Veatch⁵¹, V. Veeraraghavan⁷, L.M. Veloce¹⁶⁵, F. Veloso^{137a,137c}, S. Veneziano^{70a}, A. Ventura^{65a,65b}, M. Venturi¹⁷⁴, N. Venturi³⁵, A. Venturini²⁶, V. Vercesi^{68a}, M. Verducci^{72a,72b}, W. Verkerke¹¹⁸, A.T. Vermeulen¹¹⁸, J.C. Vermeulen¹¹⁸, M.C. Vetterli^{149,ay}, N. Viaux Maira^{144b}, O. Viazlo⁹⁴, I. Vichou^{171,*}, T. Vickey¹⁴⁶, O.E. Vickey Boeriu¹⁴⁶, G.H.A. Viehhauser¹³², S. Viel¹⁸, L. Vigani¹³², M. Villa^{23b,23a}, M. Villaplana Perez^{66a,66b}, E. Vilucchi⁴⁹, M.G. Vincter³³, V.B. Vinogradov⁷⁷, A. Vishwakarma⁴⁴, C. Vittori^{23b,23a}, I. Vivarelli¹⁵³, S. Vlachos¹⁰, M. Vogel¹⁸⁰, P. Vokac¹³⁹, G. Volpi¹⁴, H. von der Schmitt¹¹³, E. Von Toerne²⁴, V. Vorobel¹⁴⁰, K. Vorobev¹¹⁰, M. Vos¹⁷², R. Voss³⁵, J.H. Vossebeld⁸⁸, N. Vranjes¹⁶, M. Vranjes Milosavljevic¹⁶, V. Vrba¹³⁹, M. Vreeswijk¹¹⁸, T. Šfiligoj⁸⁹, R. Vuillermet³⁵, I. Vukotic³⁶, T. Ženiš^{28a}, L. Živković¹⁶, P. Wagner²⁴, W. Wagner¹⁸⁰, J. Wagner-Kuhr¹¹², H. Wahlberg⁸⁶, S. Wahrmund⁴⁶, K. Wakamiya⁸⁰, J. Walder⁸⁷, R. Walker¹¹², W. Walkowiak¹⁴⁸, V. Wallangen^{43a,43b}, C. Wang^{15c}, C. Wang^{58b,e}, F. Wang¹⁷⁹, H. Wang¹⁸, H. Wang³, J. Wang¹⁵⁴, J. Wang⁴⁴, Q. Wang¹²⁵, R.-J. Wang¹³³, R. Wang⁶, S.M. Wang¹⁵⁵, T. Wang³⁸, W. Wang^{155,q}, W.X. Wang^{58a,ag}, Z. Wang^{58c}, C. Wanotayaroj⁴⁴, A. Warburton¹⁰¹, C.P. Ward³¹, D.R. Wardrope⁹², A. Washbrook⁴⁸, P.M. Watkins²¹, A.T. Watson²¹, M.F. Watson²¹, G. Watts¹⁴⁵, S. Watts⁹⁸, B.M. Waugh⁹², A.F. Webb¹¹, S. Webb⁹⁷, M.S. Weber²⁰, S.A. Weber³³, S.M. Weber^{59a}, S.W. Weber¹⁷⁵, J.S. Webster⁶, A.R. Weidberg¹³², B. Weinert⁶³, J. Weingarten⁵¹, M. Weirich⁹⁷, C. Weiser⁵⁰, H. Weits¹¹⁸, P.S. Wells³⁵, T. Wenaus²⁹, T. Wengler³⁵, S. Wenig³⁵, N. Wermes²⁴, M.D. Werner⁷⁶, P. Werner³⁵, M. Wessels^{59a}, T.D. Weston²⁰, K. Whalen¹²⁸, N.L. Whallon¹⁴⁵, A.M. Wharton⁸⁷, A.S. White¹⁰³, A. White⁸, M.J. White¹, R. White^{144b}, D. Whiteson¹⁶⁹, B.W. Whitmore⁸⁷, F.J. Wickens¹⁴¹, W. Wiedenmann¹⁷⁹, M. Wielers¹⁴¹, C. Wiglesworth³⁹, L.A.M. Wiik-Fuchs⁵⁰, A. Wildauer¹¹³, F. Wilk⁹⁸, H.G. Wilkens³⁵, H.H. Williams¹³⁴, S. Williams³¹, C. Willis¹⁰⁴, S. Willocq¹⁰⁰, J.A. Wilson²¹, I. Wingerter-Seez⁵, E. Winkels¹⁵³, F. Winklmeier¹²⁸, O.J. Winston¹⁵³, B.T. Winter²⁴, M. Wittgen¹⁵⁰, M. Wobisch⁹³, A. Wolf⁹⁷, T.M.H. Wolf¹¹⁸, R. Wolff⁹⁹, M.W. Wolter⁸², H. Wolters^{137a,137c}, V.W.S. Wong¹⁷³, N.L. Woods¹⁴³, S.D. Worm²¹, B.K. Wosiek⁸², J. Wotschack³⁵, K.W. Woźniak⁸², M. Wu³⁶, S.L. Wu¹⁷⁹, X. Wu⁵², Y. Wu¹⁰³, T.R. Wyatt⁹⁸, B.M. Wynne⁴⁸, S. Xella³⁹, Z. Xi¹⁰³, L. Xia^{15b}, D. Xu^{15a}, L. Xu²⁹, T. Xu¹⁴², W. Xu¹⁰³, B. Yabsley¹⁵⁴, S. Yacoob^{32a}, D. Yamaguchi¹⁶³, Y. Yamaguchi¹⁶³, A. Yamamoto⁷⁹, S. Yamamoto¹⁶¹, T. Yamanaka¹⁶¹, F. Yamane⁸⁰, M. Yamatani¹⁶¹, T. Yamazaki¹⁶¹, Y. Yamazaki⁸⁰, Z. Yan²⁵, H.J. Yang^{58c,58d}, H.T. Yang¹⁸, Y. Yang¹⁵⁵, Z. Yang¹⁷, W-M. Yao¹⁸, Y.C. Yap⁴⁴, Y. Yasu⁷⁹, E. Yatsenko⁵, K.H. Yau Wong²⁴, J. Ye⁴¹, S. Ye²⁹, I. Yeletskikh⁷⁷, E. Yigitbasi²⁵, E. Yildirim⁹⁷, K. Yorita¹⁷⁷, K. Yoshihara¹³⁴, C.J.S. Young³⁵, C. Young¹⁵⁰, J. Yu⁸, J. Yu⁷⁶, S.P.Y. Yuen²⁴, I. Yusuff^{31,a}, B. Zabinski 82 , G. Zacharis¹⁰, R. Zaidan¹⁴, A.M. Zaitsev^{121,ao}, N. Zakharchuk⁴⁴, J. Zalieckas¹⁷, A. Zaman¹⁵², S. Zambito⁵⁷, D. Zanzi¹⁰², C. Zeitnitz¹⁸⁰, G. Zemaityte¹³², A. Zemla^{81a}, J.C. Zeng¹⁷¹, Q. Zeng¹⁵⁰, O. Zenin¹²¹, D. Zerwas¹²⁹, D.F. Zhang^{58b}, D. Zhang¹⁰³, F. Zhang¹⁷⁹, G. Zhang^{58a,ag}, H. Zhang¹²⁹, J. Zhang⁶, L. Zhang⁵⁰, L. Zhang^{58a}, M. Zhang¹⁷¹, P. Zhang^{15c}, R. Zhang^{58a,e}, R. Zhang²⁴, X. Zhang^{58b}, Y. Zhang^{15d}, Z. Zhang¹²⁹, X. Zhao⁴¹, Y. Zhao^{58b,129,ak}, Z. Zhao^{58a}, A. Zhemchugov⁷⁷, B. Zhou 103 , C. Zhou 179 , L. Zhou 41 , M.S. Zhou 15d , M. Zhou 152 , N. Zhou 58c , Y. Zhou 7 , C.G. Zhu 58b , H. Zhu^{15a}, J. Zhu¹⁰³, Y. Zhu^{58a}, X. Zhuang^{15a}, K. Zhukov¹⁰⁸, A. Zibell¹⁷⁵, D. Zieminska⁶³,

N.I. Zimine⁷⁷, C. Zimmermann⁹⁷, S. Zimmermann⁵⁰, Z. Zinonos¹¹³, M. Zinser⁹⁷, M. Ziolkowski¹⁴⁸, G. Zobernig¹⁷⁹, A. Zoccoli^{23b,23a}, R. Zou³⁶, M. Zur Nedden¹⁹, L. Zwalinski³⁵.

¹Department of Physics, University of Adelaide, Adelaide; Australia.

²Physics Department, SUNY Albany, Albany NY; United States of America.

³Department of Physics, University of Alberta, Edmonton AB; Canada.

^{4(a)}Department of Physics, Ankara University, Ankara;^(b)Istanbul Aydin University, Istanbul;^(c)Division of Physics, TOBB University of Economics and Technology, Ankara; Turkey.

⁵LAPP, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.

⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.

¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.

^{12(a)}Bahcesehir University, Faculty of Engineering and Natural Sciences, Istanbul;^(b)Istanbul Bilgi

University, Faculty of Engineering and Natural Sciences, Istanbul;(*c*)Department of Physics, Bogazici

University, Istanbul;^(*d*)Department of Physics Engineering, Gaziantep University, Gaziantep; Turkey.

¹³Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

¹⁴Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

15(*a*) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing;(*b*)Physics Department, Tsinghua University, Beijing;(*c*)Department of Physics, Nanjing University, Nanjing;(*d*)University of Chinese Academy of Science (UCAS), Beijing; China.

¹⁶Institute of Physics, University of Belgrade, Belgrade; Serbia.

¹⁷Department for Physics and Technology, University of Bergen, Bergen; Norway.

¹⁸Physics Division, Lawrence Berkeley National Laboratory and University of California, Berkeley CA; United States of America.

¹⁹Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.

²⁰Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.

²¹School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.

²²Centro de Investigaciónes, Universidad Antonio Nariño, Bogota; Colombia.

23(*a*)Dipartimento di Fisica e Astronomia, Università di Bologna, Bologna;(*b*) INFN Sezione di Bologna; Italy.

²⁴Physikalisches Institut, Universität Bonn, Bonn; Germany.

²⁵Department of Physics, Boston University, Boston MA; United States of America.

²⁶Department of Physics, Brandeis University, Waltham MA; United States of America.

^{27(a)}Transilvania University of Brasov, Brasov;^(b)Horia Hulubei National Institute of Physics and

Nuclear Engineering, Bucharest;(*c*)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi;(*d*)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca;^(*e*)University Politehnica Bucharest, Bucharest;^(*f*)West University in

Timisoara, Timisoara; Romania.

^{28(a)}Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava;^(b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.

²⁹Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.

³⁰Departamento de Física, Universidad de Buenos Aires, Buenos Aires; Argentina.

³¹Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.

 $32(a)$ Department of Physics, University of Cape Town, Cape Town;^(b)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg;^(*c*)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.

³³Department of Physics, Carleton University, Ottawa ON; Canada.

34(*a*)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca;^(b)Centre National de l'Energie des Sciences Techniques Nucleaires (CNESTEN), Rabat;(*c*)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech;(*d*)Faculté des Sciences, Université Mohamed Premier and LPTPM, Oujda;(*e*)Faculté des sciences, Université Mohammed V, Rabat; Morocco.

³⁵CERN, Geneva; Switzerland.

³⁶Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.

³⁷LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.

³⁸Nevis Laboratory, Columbia University, Irvington NY; United States of America.

³⁹Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.

40(*a*)Dipartimento di Fisica, Università della Calabria, Rende;(*b*) INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy.

⁴¹Physics Department, Southern Methodist University, Dallas TX; United States of America.

⁴²Physics Department, University of Texas at Dallas, Richardson TX; United States of America.

^{43(*a*)}Department of Physics, Stockholm University;^(*b*)Oskar Klein Centre, Stockholm; Sweden.

⁴⁴Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.

⁴⁵Lehrstuhl für Experimentelle Physik IV, Technische Universität Dortmund, Dortmund; Germany.

⁴⁶Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.

⁴⁷Department of Physics, Duke University, Durham NC; United States of America.

⁴⁸SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom. ⁴⁹INFN e Laboratori Nazionali di Frascati, Frascati; Italy.

⁵⁰Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.

⁵¹II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.

⁵²Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

53(*a*)Dipartimento di Fisica, Università di Genova, Genova;(*b*) INFN Sezione di Genova; Italy.

⁵⁴II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.

⁵⁵SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.

⁵⁶LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.

⁵⁷Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.

58(*a*)Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei;^(b)Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao;(*c*)School of Physics and Astronomy, Shanghai Jiao Tong University, KLPPAC-MoE, SKLPPC, Shanghai;(*d*)Tsung-Dao Lee Institute, Shanghai; China.

59(*a*)Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg;(*b*)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.

⁶⁰Faculty of Applied Information Science, Hiroshima Institute of Technology, Hiroshima; Japan. 61(*a*)Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong;(*b*)Department

of Physics, University of Hong Kong, Hong Kong;(*c*)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong;

China.

⁶²Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.

⁶³Department of Physics, Indiana University, Bloomington IN; United States of America.

64(*a*) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine;(*b*) ICTP, Trieste;(*c*)Dipartimento di Chimica, Fisica e Ambiente, Università di Udine, Udine; Italy.

65(*a*) INFN Sezione di Lecce;(*b*)Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.

66(*a*) INFN Sezione di Milano;(*b*)Dipartimento di Fisica, Università di Milano, Milano; Italy.

67(*a*) INFN Sezione di Napoli;(*b*)Dipartimento di Fisica, Università di Napoli, Napoli; Italy.

68(*a*) INFN Sezione di Pavia;(*b*)Dipartimento di Fisica, Università di Pavia, Pavia; Italy.

69(*a*) INFN Sezione di Pisa;(*b*)Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.

70(*a*) INFN Sezione di Roma;(*b*)Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.

71(*a*) INFN Sezione di Roma Tor Vergata;(*b*)Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.

72(*a*) INFN Sezione di Roma Tre;(*b*)Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.

73(*a*) INFN-TIFPA;(*b*)Università degli Studi di Trento, Trento; Italy.

⁷⁴Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität, Innsbruck; Austria.

⁷⁵University of Iowa, Iowa City IA; United States of America.

⁷⁶Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America. ⁷⁷Joint Institute for Nuclear Research, Dubna; Russia.

78(*a*)Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora;(*b*)Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro;(*c*)Universidade Federal de São João del Rei (UFSJ), São João del Rei;(*d*) Instituto de Física, Universidade de São Paulo, São Paulo; Brazil.

⁷⁹KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.

⁸⁰Graduate School of Science, Kobe University, Kobe; Japan.

^{81(*a*)}AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow;(*b*)Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.

⁸²Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.

⁸³Faculty of Science, Kyoto University, Kyoto; Japan.

⁸⁴Kyoto University of Education, Kyoto; Japan.

⁸⁵Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.

⁸⁶Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina. ⁸⁷Physics Department, Lancaster University, Lancaster; United Kingdom.

⁸⁸Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.

⁸⁹Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.

⁹⁰School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.

⁹¹Department of Physics, Royal Holloway University of London, Egham; United Kingdom.

⁹²Department of Physics and Astronomy, University College London, London; United Kingdom.

⁹³Louisiana Tech University, Ruston LA; United States of America.

⁹⁴Fysiska institutionen, Lunds universitet, Lund; Sweden.

⁹⁵Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules (IN2P3), Villeurbanne; France.

⁹⁶Departamento de Física Teorica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain. ⁹⁷Institut für Physik, Universität Mainz, Mainz; Germany.

98School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.

⁹⁹CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.

¹⁰⁰Department of Physics, University of Massachusetts, Amherst MA; United States of America.

¹⁰¹Department of Physics, McGill University, Montreal QC; Canada.

¹⁰²School of Physics, University of Melbourne, Victoria; Australia.

¹⁰³Department of Physics, University of Michigan, Ann Arbor MI; United States of America.

¹⁰⁴Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.

¹⁰⁵B.I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk; Belarus.

¹⁰⁶Research Institute for Nuclear Problems of Byelorussian State University, Minsk; Belarus.

¹⁰⁷Group of Particle Physics, University of Montreal, Montreal OC: Canada.

¹⁰⁸ P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow; Russia.

¹⁰⁹Institute for Theoretical and Experimental Physics (ITEP), Moscow; Russia.

¹¹⁰National Research Nuclear University MEPhI, Moscow; Russia.

¹¹¹D.V. Skobeltsyn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University, Moscow; Russia.

¹¹²Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.

¹¹³Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.

¹¹⁴Nagasaki Institute of Applied Science, Nagasaki; Japan.

¹¹⁵Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.

¹¹⁶Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.

¹¹⁷Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen; Netherlands.

¹¹⁸Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.

¹¹⁹Department of Physics, Northern Illinois University, DeKalb IL: United States of America.

 $120(a)$ Budker Institute of Nuclear Physics and NSU, SB RAS, Novosibirsk;^(b)Novosibirsk State University Novosibirsk; Russia.

¹²¹Institute for High Energy Physics of the National Research Centre Kurchatov Institute, Protvino; Russia.

¹²²Department of Physics, New York University, New York NY; United States of America.

¹²³Ohio State University, Columbus OH; United States of America.

¹²⁴Faculty of Science, Okayama University, Okayama; Japan.

¹²⁵Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.

¹²⁶Department of Physics, Oklahoma State University, Stillwater OK; United States of America.

¹²⁷Palacký University, RCPTM, Joint Laboratory of Optics, Olomouc; Czech Republic.

¹²⁸Center for High Energy Physics, University of Oregon, Eugene OR; United States of America.

¹²⁹LAL, Université Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay; France.

¹³⁰Graduate School of Science, Osaka University, Osaka; Japan.

¹³¹Department of Physics, University of Oslo, Oslo; Norway.

¹³²Department of Physics, Oxford University, Oxford; United Kingdom.

¹³³LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris; France.

¹³⁴Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.

¹³⁵Konstantinov Nuclear Physics Institute of National Research Centre "Kurchatov Institute", PNPI, St. Petersburg; Russia.

¹³⁶Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.

137(*a*)Laboratório de Instrumentação e Física Experimental de Partículas - LIP;(*b*)Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa;(*c*)Departamento de Física, Universidade de Coimbra, Coimbra;(*d*)Centro de Física Nuclear da Universidade de Lisboa, Lisboa;(*e*)Departamento de Física, Universidade do Minho, Braga;(*f*)Departamento de Física Teorica y del Cosmos, Universidad de Granada, Granada (Spain);^(g)Dep Física and CEFITEC of Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Caparica; Portugal.

¹³⁸Institute of Physics, Academy of Sciences of the Czech Republic, Prague; Czech Republic.

¹³⁹Czech Technical University in Prague, Prague; Czech Republic.

¹⁴⁰Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.

¹⁴¹ Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.

¹⁴²IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.

¹⁴³Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.

144(*a*)Departamento de Física, Pontificia Universidad Católica de Chile, Santiago;(*b*)Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.

¹⁴⁵Department of Physics, University of Washington, Seattle WA; United States of America.

¹⁴⁶Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.

¹⁴⁷Department of Physics, Shinshu University, Nagano; Japan.

¹⁴⁸Department Physik, Universität Siegen, Siegen; Germany.

¹⁴⁹Department of Physics, Simon Fraser University, Burnaby BC; Canada.

¹⁵⁰SLAC National Accelerator Laboratory, Stanford CA; United States of America.

¹⁵¹Physics Department, Royal Institute of Technology, Stockholm; Sweden.

¹⁵²Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.

¹⁵³Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.

¹⁵⁴School of Physics, University of Sydney, Sydney; Australia.

¹⁵⁵Institute of Physics, Academia Sinica, Taipei; Taiwan.

¹⁵⁶Academia Sinica Grid Computing, Institute of Physics, Academia Sinica, Taipei; Taiwan.

157(*a*)E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi;(*b*)High Energy Physics Institute, Tbilisi State University, Tbilisi; Georgia.

¹⁵⁸Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.

¹⁵⁹Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.

¹⁶⁰Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.

¹⁶¹International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.

¹⁶²Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo; Japan.

¹⁶³Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.

¹⁶⁴Tomsk State University, Tomsk; Russia.

¹⁶⁵Department of Physics, University of Toronto, Toronto ON; Canada.

^{166(a)}TRIUMF, Vancouver BC;^(b)Department of Physics and Astronomy, York University, Toronto ON; Canada.

¹⁶⁷Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.

¹⁶⁸Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.

¹⁶⁹Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.

¹⁷⁰Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.

¹⁷¹Department of Physics, University of Illinois, Urbana IL; United States of America.

¹⁷²Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.

¹⁷³Department of Physics, University of British Columbia, Vancouver BC; Canada.

¹⁷⁴Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.

¹⁷⁵Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.

¹⁷⁶Department of Physics, University of Warwick, Coventry; United Kingdom.

¹⁷⁷Waseda University, Tokyo; Japan.

¹⁷⁸Department of Particle Physics, Weizmann Institute of Science, Rehovot; Israel.

¹⁷⁹Department of Physics, University of Wisconsin, Madison WI; United States of America.

¹⁸⁰Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.

¹⁸¹Department of Physics, Yale University, New Haven CT; United States of America.

¹⁸²Yerevan Physics Institute, Yerevan; Armenia.

^a Also at Department of Physics, University of Malaya, Kuala Lumpur; Malaysia.

^b Also at Borough of Manhattan Community College, City University of New York, NY; United States of America.

^c Also at Centre for High Performance Computing, CSIR Campus, Rosebank, Cape Town; South Africa.

^d Also at CERN, Geneva; Switzerland.

^e Also at CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.

^f Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

 θ Also at Departament de Fisica de la Universitat Autonoma de Barcelona, Barcelona; Spain.

^h Also at Departamento de Física Teorica y del Cosmos, Universidad de Granada, Granada (Spain); Spain.

ⁱ Also at Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal. *^j* Also at Department of Applied Physics and Astronomy, University of Sharjah, Sharjah; United Arab Emirates.

^k Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.

^{*l*} Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America.

^m Also at Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.

ⁿ Also at Department of Physics, California State University, Fresno CA; United States of America.

^o Also at Department of Physics, California State University, Sacramento CA; United States of America.

^p Also at Department of Physics, King's College London, London; United Kingdom.

^q Also at Department of Physics, Nanjing University, Nanjing; China.

^r Also at Department of Physics, St. Petersburg State Polytechnical University, St. Petersburg; Russia.

^s Also at Department of Physics, Stanford University; United States of America.

^t Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.

^u Also at Department of Physics, University of Michigan, Ann Arbor MI; United States of America.

^v Also at Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.

 w Also at Giresun University, Faculty of Engineering, Giresun; Turkey.

^x Also at Graduate School of Science, Osaka University, Osaka; Japan.

^y Also at Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; Romania.

^z Also at II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.

aa Also at Institucio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.

ab Also at Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

ac Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.

ad Also at Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen; Netherlands.

ae Also at Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Budapest; Hungary.

af Also at Institute of Particle Physics (IPP); Canada.

- *^a*^g Also at Institute of Physics, Academia Sinica, Taipei; Taiwan.
- *ah* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- *ai* Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.

^{a j} Also at Instituto de Física Teórica de la Universidad Autónoma de Madrid; Spain.

ak Also at LAL, Université Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay; France.

al Also at Louisiana Tech University, Ruston LA; United States of America.

am Also at LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris; France.

an Also at Manhattan College, New York NY; United States of America.

ao Also at Moscow Institute of Physics and Technology State University, Dolgoprudny; Russia.

ap Also at National Research Nuclear University MEPhI, Moscow; Russia.

aq Also at Near East University, Nicosia, North Cyprus, Mersin; Turkey.

ar Also at Novosibirsk State University, Novosibirsk; Russia.

as Also at Ochadai Academic Production, Ochanomizu University, Tokyo; Japan.

at Also at Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.

au Also at School of Physics, Sun Yat-sen University, Guangzhou; China.

^a^v Also at The City College of New York, New York NY; United States of America.

^a^w Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.

ax Also at Tomsk State University, Tomsk, and Moscow Institute of Physics and Technology State University, Dolgoprudny; Russia.

^a^y Also at TRIUMF, Vancouver BC; Canada.

az Also at Universita di Napoli Parthenope, Napoli; Italy.

[∗] Deceased