

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-PPE/93-151
August 9, 1993

Search for Lepton Flavour Violation in Z Decays

L3 Collaboration

Abstract

We have searched for lepton flavour violation in Z boson decays into lepton pairs using all data collected with L3 detector during the 1990, 1991 and 1992 runs on an event sample corresponding to 1,500,000 Z's produced. At the 95% confidence level the upper limits on the branching ratio for $Z \rightarrow e\mu$ is 0.6×10^{-5} , for $Z \rightarrow e\tau$ is 1.3×10^{-5} and for $Z \rightarrow \mu\tau$ is 1.9×10^{-5} .

(Submitted to *Physics Letters B*)

1 Introduction

In the Standard Model [1, 2] lepton flavour is conserved. However, there is no gauge principle requiring this conservation. Different models [3–9], beyond the Standard Model, allow processes which violate lepton flavour conservation. In theories where such violation arises through mixing with new particles [4, 6], the branching ratios for such processes, e.g. $Z \rightarrow \mu\tau$, have been calculated to be as large as 10^{-4} in certain models. The observation of such decays would be a clear indication of physics beyond the Standard Model. Here a search for lepton flavour violation in Z decays into $e\mu$, $e\tau$ and $\mu\tau$ is reported using the data collected with L3 detector during the 1990, 1991 and 1992 runs on an event sample corresponding to 1,500,000 Z 's produced. Direct searches for lepton flavour violation [10–17] have been performed previously by L3 and other experiments. Stringent constraints on violation of μ flavour exist from low- q^2 reactions, such as the absence of the decay $\mu \rightarrow eee$ [14], providing an upper limit [18] on $Br(Z \rightarrow e\mu)$ of 6.6×10^{-13} . Searches for neutrinoless τ decays $\tau \rightarrow eee$, $\tau \rightarrow \mu\mu\mu$ [12, 13] lead to much less stringent limits on $Br(Z \rightarrow e\tau)$ and $Br(Z \rightarrow \mu\tau)$.

2 The L3 Detector

The fiducial solid angle of the L3 detector [19] is 99% of 4π . The detector consists of a time expansion chamber (TEC) for tracking charged particles, a high resolution electromagnetic calorimeter of BGO crystals, a barrel of scintillation counters, a hadron calorimeter with uranium absorber and proportional wire chamber readout and a muon spectrometer. The luminosity is determined from small-angle Bhabha scattering using BGO electromagnetic calorimeters in the polar angle ranges θ and $\pi - \theta$ between 24.93 and 69.94 mrad. All subdetectors are installed inside a 12m diameter solenoidal magnet which provides a uniform 0.5 T field along the beam direction. A detailed description of each detector subsystem, and its performance, is given in [11, 19].

3 Signature and Background

The expected experimental signature of $Z \rightarrow e\mu$, $Z \rightarrow \mu\tau$ and $Z \rightarrow e\tau$ is an electron or muon with an energy close to the beam energy recoiling against a different type of lepton or hadrons from tau decay. Background arises from Standard Model leptonic final states and can be divided into two classes : (i) incorrectly reconstructed e^+e^- and $\mu^+\mu^-$ events; (ii) $\tau^+\tau^-$ events with one or both of the taus decaying into a muon or electron which carries almost all the energy of the tau. Good muon and electron energy resolution is essential to reduce the latter background while retaining a high detection efficiency. The energy resolution for electrons determined from Bhabha events is 1.3% at 45 GeV, while the muon momentum resolution determined from dimuon events is 2.5%.

About 300000 Monte Carlo events were generated to study Standard Model backgrounds from e^+e^- , $\mu^+\mu^-$, $\tau^+\tau^-$ and four-fermion channels using various Monte Carlo

generators [20–23]. A modified version of the KORALZ [20] Monte Carlo program was used to generate signal events.

4 Preselection and Lepton Identification

The preselection cuts, used to select a data sample containing high energy dilepton events of all types, are the following:

- The total energy is greater than 30 GeV.
- The number of jets is 2.
- The number of tracks in the TEC is between 1 and 6, to remove hadron events.
- The number of calorimeter clusters is less than 15, to remove hadron events.
- The acolinearity angle between the two jets is smaller than 20° , to remove radiative events.

Jets are reconstructed using a two step algorithm [24] which groups the energy deposited in calorimeters into clusters before collecting the clusters into jets. The clustering algorithm normally reconstructs one cluster for each muon, electron or photon shower, and a few clusters for a hadronic decay of a single τ . Under the above definition of a jet, particles with only one cluster, like electrons, are also considered as jets.

An electron is defined as a geometrical cluster in the electromagnetic calorimeter with an energy more than 2 GeV matched with a TEC track in the (R, ϕ) plane within 10 mrad. The cluster shower profile should be consistent with that of an electron, i.e. we require $0.97 < E_9/E_{25} < 1.025$, where $E_{9(25)}$ is the corrected sum of energies of 9(25) BGO crystals around the most energetic one, and $\chi^2 < 2$, where χ^2 is obtained by fitting the shape of the cluster to the mean electron shower profile. The electron candidate must be in the fiducial volume defined by $|\cos \theta| < 0.9$.

Muons are identified and their momenta measured in the muon chamber system surrounding the calorimeters. To be accepted, a muon track must have one track segment in each of the three chamber layers. The accepted muon track must extrapolate back to within 100 mm of the nominal vertex position in the transverse plane and 200 mm in the longitudinal plane. The muon candidate energy must be greater than 3 GeV and it must be in the fiducial volume defined by $|\cos \theta| < 0.75$.

5 $Z \rightarrow e\mu$ Channel

For the $Z \rightarrow e\mu$ channel we require one jet to be consistent with a beam energy electron and the other one to be consistent with a beam energy muon. This type of event is

essentially free of background and allows a less restrictive selection cut on lepton energy than $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ channels. Fig. 1 shows the normalised electron energy versus the normalised muon momentum for the data events selected. Signal events will populate the region where both normalised energies are close to unity. We require that for the electron candidate the normalised energy should be greater than 0.96 ($\approx 3\sigma$) and for the muon candidate greater than 0.92 ($\approx 3\sigma$). No events lie within this region giving the 95% confidence level limit on the signal of three events. We apply the above cuts to the sample of the signal and Standard Model background ($\tau^+\tau^-$, e^+e^- and $\mu^+\mu^-$) Monte Carlo events. We find no events in the Standard Model Monte Carlo sample. The efficiency of the analysis as determined by the signal Monte Carlo is $32.0 \pm 1.0\%$. This error includes the uncertainty due to the Monte Carlo statistics. The additional systematic error due to the corrected number of hadronic Z decays used for normalization and due to the uncertainties in the detector calibration constants is estimated to be 1.5% [25]. This yields a 95% confidence level limit on the branching ratio of

$$Br(Z \rightarrow e\mu) < 0.6 \times 10^{-5}.$$

6 $Z \rightarrow e\tau$ Channel

For $Z \rightarrow e\tau$ we require one jet to be consistent with a beam energy electron and the other one to be consistent with a τ decay. We consider the following τ decays :

- $\tau \rightarrow \mu\nu\bar{\nu}$

We require that the τ -jet consists of one muon with an energy greater than 3 GeV.

- $\tau \rightarrow e\nu\bar{\nu}$

We require that the τ -jet consists of one electron with an energy less than 30 GeV. The hadronic energy, contained in the τ jet, should be less than 0.5 GeV. To reject the $Z \rightarrow ee(\gamma)$ background we require that the acoplanarity angle between TEC tracks associated with the electron and the τ be more than 3.2 mrad. Fig. 2 shows the distribution of this variable for $Z \rightarrow ee(\gamma)$ as well as $Z \rightarrow e\tau$ events.

- $\tau \rightarrow 1$ prong

We require that the τ -jet, containing only one TEC track, has a total energy more than 3 GeV and electromagnetic energy less than 30 GeV. The hadronic energy contained in the τ jet should be greater than 0.5 GeV. To reject $Z \rightarrow ee(\gamma)$ events, where one electron passes close to the cracks in the electromagnetic calorimeter and therefore deposits some energy in the hadron calorimeter, we require that the energy in the hadron calorimeter beyond its first $25 X_0$ be greater than 10% of the total jet energy. The above cut on the acoplanarity angle between two TEC tracks is also applied.

- $\tau \rightarrow 3$ or 5 prongs

We require that the τ -jet contains at least two TEC tracks, has a total energy more than 3 GeV, has an electromagnetic energy less than 30 GeV and an energy deposition in the hadron calorimeter more than 1 GeV. To remove four-fermion events,

which have no missing energy, we require that energy in the hadronic calorimeter be less than $0.85 \times (E_{beam} - E_{ECAL})$.

Figure 3 shows the distribution of the electron energy after all cuts but the electron energy cut have been applied. To reject the remaining $Z \rightarrow \tau\tau$ background we require that the energy of the electron should be more than $0.985 \times E_{beam}$. After all cuts this yields an efficiency for the $Z \rightarrow e\tau$ channel of $23.8 \pm 0.9\%$.

Table 1 shows the estimated acceptance for the $Z \rightarrow e\tau$ channel for the different τ decay modes together with the number of surviving events after all cuts.

Channel	Acceptance (%)	Expected	Seen
$\tau \rightarrow e\nu\bar{\nu}$	4.0 ± 0.4	0	0
$\tau \rightarrow \mu\nu\bar{\nu}$	7.1 ± 0.5	0.7 ± 0.4	0
$\tau \rightarrow 1$ prong	8.5 ± 0.5	1.1 ± 0.4	2
$\tau \rightarrow 3$ or 5 prongs	4.2 ± 0.4	1.0 ± 0.3	0

Table 1: Acceptance, expected background events and number of events seen in the data for the $Z \rightarrow e\tau$ channel for different τ decay modes.

We find two events in the data, while we expect 2.8 ± 0.7 events from $Z \rightarrow \tau\tau$ and $Z \rightarrow ffff$ Monte Carlo. We set a 95% C.L. upper limit of 4.7 events for the $Z \rightarrow e\tau$ channel, yielding a 95% C.L. limit on the branching ratio of

$$Br(Z \rightarrow e\tau) < 1.3 \times 10^{-5}.$$

7 $Z \rightarrow \mu\tau$ Channel

For $Z \rightarrow \mu\tau$ we require one jet to be consistent with a beam energy muon and the other one to be consistent with a τ decay. We consider the following τ decays :

- $\tau \rightarrow \mu\nu\bar{\nu}$

Due to the background from dimuon events, this decay channel is not considered.

- $\tau \rightarrow e\nu\bar{\nu}$

We require that the τ -jet consists of one electron with energy greater than 5 GeV. The hadronic energy, contained in the τ jet, should be less than 0.5 GeV. To reject the $Z \rightarrow \mu\mu(\gamma)$ background we require : (i) the acoplanarity between TEC tracks associated with the muon and the τ to be greater than 3.2 mrad; (ii) there are no track segments in the muon chambers in the hemisphere opposite to the μ candidate.

- $\tau \rightarrow 1$ prong

We require that the τ -jet contains only one TEC track, has a total energy more than 5 GeV and an electromagnetic energy less than 35 GeV. The hadronic energy contained in the τ jet should be greater than 0.5 GeV. To reject the $Z \rightarrow \mu\mu(\gamma)$

background, the shower profile of the τ jet in the hadron calorimeter should be inconsistent with that of a muon [10]. Cuts (i) and (ii) from the previous decay channel are also applied.

- $\tau \rightarrow 3$ or 5 prongs

We require that the τ -jet contains at least two TEC tracks and has a total energy greater than 5 GeV. There must be no track segments in muon chambers in the hemisphere opposite to the μ candidate.

Figure 4 shows the distribution of the muon energy after all cuts but the electron energy cut have been applied. To reject the remaining $Z \rightarrow \tau\tau$ background we require that the energy of the muon should be more than $0.97 \times E_{beam}$. After all cuts this yields an efficiency for the $Z \rightarrow \mu\tau$ channel of $22.0 \pm 0.9\%$.

Table 2 shows the estimated acceptance for the $Z \rightarrow \mu\tau$ channel for the different τ decay modes together with the number of surviving events after all cuts.

Channel	Acceptance (%)	Expected	Seen
$\tau \rightarrow e\nu\bar{\nu}$	5.0 ± 0.4	1.8 ± 0.8	1
$\tau \rightarrow 1$ prong	11.8 ± 0.6	3.4 ± 0.8	3
$\tau \rightarrow 3$ or 5 prongs	5.2 ± 0.5	0.7 ± 0.3	1

Table 2: Acceptance, expected background events and number of events seen in the data for the $Z \rightarrow \mu\tau$ channel for different τ decay modes.

We find 5 events in data, while we expect 5.9 ± 1.2 events from $Z \rightarrow \tau\tau$, $Z \rightarrow \mu\mu$ and $Z \rightarrow ffff$ Monte Carlo. We set a 95% C.L. upper limit of 6.2 events for the $Z \rightarrow \mu\tau$ channel, yielding a 95% C.L. limit on the branching ratio of

$$Br(Z \rightarrow \mu\tau) < 1.9 \times 10^{-5}.$$

8 Conclusion

We have searched for lepton flavour violating decays of the Z boson, obtaining the limits :

$$Br(Z \rightarrow e\mu) < 0.6 \times 10^{-5}$$

$$Br(Z \rightarrow e\tau) < 1.3 \times 10^{-5}$$

$$Br(Z \rightarrow \mu\tau) < 1.9 \times 10^{-5}.$$

These limits are approximately 5 times better than previously published LEP results [15–17].

9 Acknowledgements

We wish to express our gratitude to the CERN accelerator divisions for the excellent performance of the LEP machine. We acknowledge the efforts of all engineers and technicians who have participated in the construction and maintenance of this experiment.

The L3 Collaboration:

O.Adriani,¹⁵ M.Aguilar-Benitez,²⁴ S.Ahlen,⁹ J.Alcaraz,¹⁶ A.Aloisio,²⁷ G.Alverson,¹⁰ M.G.Alvaggi,²⁷ G.Ambrosi,³² Q.An,¹⁷ H.Anderhub,⁴⁶ A.L.Anderson,¹⁴ V.P.Andreev,³⁶ T.Angelescu,¹¹ L.Antonov,⁴⁰ D.Antreasyan,⁷ P.Arce,²⁴ A.Arefiev,²⁶ A.Atamanchuk,³⁶ T.Azemoon,³ T.Aziz,⁸ P.V.K.S.Baba,¹⁷ P.Bagnaia,³⁵ J.A.Bakken,³⁴ R.C.Ball,³ S.Banerjee,⁸ J.Bao,⁵ R.Barillère,¹⁶ L.Barone,³⁵ A.Baschirotto,²⁵ R.Battiston,³² A.Bay,¹⁸ F.Becattini,¹⁵ J.Bechtluft,¹ R.Becker,¹ U.Becker,^{14,46} F.Behner,⁴⁶ J.Behrens,⁴⁶ Gy.L.Bencze,¹² J.Berdugo,²⁴ P.Berges,¹⁴ B.Bertucci,³² B.L.Betev,^{40,46} M.Biasini,³² A.Biland,⁴⁶ G.M.Bilei,³² R.Bizzarri,³⁵ J.J.Blaising,⁴ G.J.Bobbink,^{16,2} R.Bock,¹ A.Böhm,¹ B.Borgia,³⁵ M.Bosetti,²⁵ D.Bourilkov,²⁹ M.Bourquin,¹⁸ D.Boutigny,¹⁶ B.Bouwens,² E.Brambilla,²⁷ J.G.Branson,³⁷ I.C.Brock,³³ M.Brooks,²² A.Bujak,⁴³ J.D.Burger,¹⁴ W.J.Burger,¹⁸ J.Busenitz,⁴² A.Buytenhuys,²⁹ X.D.Cai,⁷ M.Capell,¹⁴ M.Caria,³² G.Carlino,²⁷ A.M.Cartacci,¹⁵ R.Castello,²⁵ M.Cerrada,²⁴ F.Cesaroni,³⁵ Y.H.Chang,¹⁴ U.K.Chaturvedi,¹⁷ M.Chemarin,²³ A.Chen,⁴⁸ C.Chen,⁶ G.Chen,⁶ G.M.Chen,⁶ H.F.Chen,¹⁹ H.S.Chen,⁶ M.Chen,¹⁴ W.Y.Chen,⁴⁸ G.Chiefari,²⁷ C.Y.Chien,⁵ M.T.Choi,⁴¹ S.Chung,¹⁴ C.Civinini,¹⁵ I.Clare,¹⁴ R.Clare,¹⁴ T.E.Coan,²² H.O.Cohn,³⁰ G.Coignet,⁴ N.Colino,¹⁶ A.Contin,⁷ F.Cotorobai,¹¹ X.T.Cui,¹⁷ X.Y.Cui,¹⁷ T.S.Dai,¹⁴ R.D'Alessandro,⁵ R.de Asmundis,²⁷ A.Degré,⁴ K.Deiters,⁴⁴ E.Dénes,¹² P.Denes,³⁴ F.DeNotaristefani,³⁵ M.Dhina,⁴⁶ D.DiBitonto,⁴² M.Diemoz,³⁵ H.R.Dimitrov,⁴⁰ C.Dionisi,³⁵ M.Dittmar,⁴⁶ L.Djambazov,⁴⁶ M.T.Dova,¹⁷ E.Drago,²⁷ D.Duchesneau,¹⁸ P.Duinker,² I.Duran,³⁸ S.Easo,³² H.El Mamouni,²³ A.Engler,³³ F.J.Eppling,¹⁴ F.C.Erné,² P.Extermann,¹⁸ R.Fabbretti,⁴⁴ M.Fabre,⁴⁴ S.Falciano,³⁵ S.J.Fan,³⁹ O.Fackler,²¹ J.Fay,²³ M.Felcini,¹⁶ T.Ferguson,³³ D.Fernandez,²⁴ G.Fernandez,²⁴ F.Ferroni,³⁵ H.Fesefeldt,¹ E.Fiandrini,³² J.H.Field,¹⁸ F.Filthaut,²⁹ P.H.Fisher,⁵ G.Forconi,¹⁸ L.Fredj,¹⁸ K.Freudenreich,⁴⁶ W.Friebel,⁴⁵ M.Fukushima,¹⁴ M.Gaillard,²⁰ Yu.Galaktionov,^{26,14} E.Gallo,¹⁵ S.N.Ganguli,^{16,8} P.Garcia-Abia,²⁴ D.Gele,²³ S.Gentile,³⁵ N.Gheordanescu,¹¹ S.Giagu,³⁵ S.Goldfarb,¹⁰ Z.F.Gong,¹⁹ E.Gonzalez,²⁴ A.Gougas,⁵ D.Goujon,¹⁸ G.Gratta,³¹ M.Gruenewald,¹⁶ C.Gu,¹⁷ M.Guanziroli,¹⁷ J.K.Guo,³⁹ V.K.Gupta,³⁴ A.Gurtu,⁸ H.R.Gustafson,³ L.J.Gutay,⁴³ K.Hangarter,¹ B.Hartmann,¹ A.Hasan,¹⁷ D.Hauschildt,² C.F.He,³⁹ J.T.He,⁶ T.Hebbeker,¹⁶ M.Hebert,³⁷ A.Hervé,¹⁶ K.Hilgers,¹ H.Hofe,⁴⁶ H.Hoorani,¹⁸ G.Hu,¹⁷ G.Q.Hu,³⁹ B.Ille,²³ M.M.Ilyas,¹⁷ V.Innocente,¹⁶ H.Janssen,¹⁶ S.Jezequel,⁴ B.N.Jin,⁶ L.W.Jones,³ I.Josa-Mutuberria,¹⁶ A.Kasser,²⁰ R.A.Khan,¹⁷ Yu.Kamyshkov,³⁰ P.Kapinos,^{36,45} J.S.Kapustinsky,²² Y.Karyotakis,¹⁶ M.Kaur,¹⁷ S.Khokhar,¹⁷ M.N.Kienzle-Focacci,¹⁸ J.K.Kim,⁴¹ S.C.Kim,⁴¹ Y.G.Kim,⁴¹ W.W.Kinnison,²² A.Kirkby,³¹ D.Kirkby,³¹ S.Kirsch,⁴⁵ W.Kittel,²⁹ A.Klimentov,^{14,26} R.Klöckner,¹ A.C.König,²⁹ E.Koffman,² O.Kornadt,¹ V.Koutsenko,^{14,26} A.Koulbardis,³⁶ R.W.Kraemer,³³ T.Kramer,¹⁴ V.R.Krastev,^{40,32} W.Krenz,¹ A.Krivshich,³⁶ H.Kuijten,²⁹ K.S.Kumar,¹³ A.Kunin,^{14,26} G.Landi,¹⁵ D.Lanske,¹ S.Lanzano,²⁷ A.Lebedev,¹⁴ P.Lebrun,²³ P.Lecomte,⁴⁶ P.Lecoq,¹⁶ P.Le Coultre,⁴⁶ D.M.Lee,²² J.S.Lee,⁴¹ K.Y.Lee,⁴¹ I.Leedom,¹⁰ C.Leggett,³ J.M.Le Goff,¹⁶ R.Leiste,⁴⁵ M.Lenti,¹⁵ E.Leonardi,³⁵ C.Li,^{9,17} H.T.Li,⁶ P.J.Li,³⁹ J.Y.Liao,³⁹ W.T.Lin,⁴⁸ Z.Y.Lin,¹⁹ F.L.Linde,² B.Lindemann,¹ L.Lista,²⁷ Y.Liu,¹⁷ W.Lohmann,⁴⁵ E.Longo,³⁵ Y.S.Lu,⁶ J.M.Lubbers,¹⁶ K.Lübelsmeyer,¹ C.Luci,³⁵ D.Luckey,^{7,14} L.Ludovici,³⁵ L.Luminari,³⁵ W.Lustermann,⁴⁵ J.M.Ma,⁶ W.G.Ma,¹⁹ M.MacDermott,⁴⁶ R.Malik,¹⁷ A.Malinin,²⁶ C.Maňa,²⁴ M.Maolinbay,⁴⁶ P.Marchesini,⁴⁶ F.Marion,⁴ A.Marin,⁹ J.P.Martin,²³ L.Martinez-Laso,²⁴ F.Marzano,³⁵ G.G.G.Massaro,² K.Mazumdar,¹⁸ P.McBride,¹³ T.McMahon,⁴³ D.McNally,⁴⁶ M.Merk,³³ L.Merola,²⁷ M.Meschini,¹⁵ W.J.Metzger,²⁹ Y.Mi,²⁰ A.Mihul,¹¹ G.B.Mills,²² Y.Mir,¹⁷ G.Mirabelli,³⁵ J.Mnich,¹ M.Möller,¹ B.Monteleoni,⁵ R.Morand,⁴ S.Morganti,³⁵ N.E.Moulai,¹⁷ R.Mount,³¹ S.Müller,¹ A.Nadtochy,³⁶ E.Nagy,¹² M.Napolitano,²⁷ F.Nessi-Tedaldi,⁴⁶ H.Newman,³¹ C.Neyer,⁴⁶ M.A.Niaz,¹⁷ A.Nippe,¹ H.Nowak,⁴⁵ G.Organtini,³⁵ D.Pandoulas,¹ S.Paoletti,³⁵ P.Paolucci,²⁷ G.Pascale,³⁵ G.Passaleva,^{15,32} S.Patricelli,²⁷ T.Paul,⁵ M.Pauluzzi,³² C.Paus,¹ F.Pauss,⁴⁶ Y.J.Pei,¹ S.Pensotti,²⁵ D.Perret-Gallix,⁴ J.Perrier,¹⁸ A.Pevsner,⁵ D.Piccolo,²⁷ M.Pieri,¹⁶ P.A.Piroué,³⁴ F.Plasil,³⁰ V.Plyaskin,²⁶ M.Pohl,⁴⁶ V.Pojidaev,^{26,15} H.Postema,¹⁴ Z.D.Qi,³⁹ J.M.Qian,³ K.N.Qureshi,¹⁷ R.Raghavan,⁸ G.Rahal-Callot,⁴⁶ P.G.Rancoita,²⁵ M.Rattaggi,²⁵ G.Raven,² P.Razis,²⁸ K.Read,³⁰ D.Ren,⁴⁶ Z.Ren,¹⁷ M.Rescigno,³⁵ S.Reucroft,¹⁰ A.Ricker,¹ S.Riemann,⁴⁵ B.C.Riemers,⁴³ K.Riles,³ O.Rind,³ H.A.Rizvi,⁷ S.Ro,⁴¹ F.J.Rodriguez,²⁴ B.P.Roe,³ M.Röhner,¹ L.Romero,²⁴ S.Rosier-Lees,⁴ R.Rosmalen,²⁹ Ph.Rosselet,²⁰ W.van Rossum,² S.Roth,¹ A.Rubbia,¹⁴ J.A.Rubio,¹⁶ H.Rykaczewski,⁴⁶ M.Sachwitz,⁴⁵ J.Salicic,¹⁶ J.M.Salicic,²⁴ G.S.Sanders,²² A.Santocchia,³² M.S.Sarakinos,¹⁴ G.Sartorelli,^{7,17} M.Sassowsky,¹ G.Sauvage,⁴ V.Schegelsky,³⁶ D.Schmitz,¹ P.Schmitz,¹ M.Schneegans,⁴ H.Schopper,⁴⁷ D.J.Shotanus,²⁹ S.Shotkin,¹⁴ H.J.Schreiber,⁴⁵ J.Shukla,³³ R.Schulte,¹ S.Schulte,¹ K.Schultze,¹ J.Schwenke,¹ G.Schowering,¹ C.Sciacca,²⁷ I.Scott,¹³ R.Sehgal,¹⁷ P.G.Seiler,⁴⁴ J.C.Sens,^{16,2} L.Servoli,³² I.Sheer,³⁷ D.Z.Shen,³⁹ S.Shevchenko,³¹ X.R.Shi,³¹ E.Shumilov,²⁶ V.Shoutko,²⁶ D.Son,⁴¹ A.Sopczak,¹⁶ V.Soulimov,²⁷ C.Spartiotis,⁵ T.Spickermann,¹ P.Spillantini,¹⁵ R.Starosta,¹ M.Steuer,^{7,14} D.P.Stickland,³⁴ F.Sticcozzi,¹⁴ H.Stone,³⁴ K.Strauch,¹³ B.C.Stringfellow,⁴³ K.Sudhakar,⁸ G.Sultanov,¹⁷ L.Z.Sun,^{19,17} G.F.Susinno,¹⁸ H.Suter,⁴⁶ J.D.Swain,¹⁷ A.A.Syed,²⁹ X.W.Tang,⁶ L.Taylor,¹⁰ G.Terzi,²⁵ Samuel C.C.Ting,¹⁴ S.M.Ting,¹⁴ M.Tonutti,¹ S.C.Tonwar,⁸ J.Tóth,¹² A.Tsaregorodtsev,³⁶ G.Tsipolitis,³³ C.Tully,³⁴ K.L.Tung,⁶ J.Ulbricht,⁴⁶ L.Urbán,¹² U.Uwer,¹ E.Valente,³⁵ R.T.Van de Walle,²⁹ I.Vetlitsky,²⁶ G.Viertel,⁴⁶ P.Vikas,¹⁷ U.Vikas,¹⁷ M.Vivargent,⁴ H.Vogel,³³ H.Vogt,⁴⁵ I.Vorobiev,²⁶ A.A.Vorobyov,³⁶ L.Vuilleumier,²⁰ M.Wadhwa,⁴ W.Wallraft,¹ C.Wang,¹⁴ C.R.Wang,¹⁹ X.L.Wang,¹⁹ Y.F.Wang,¹⁴ Z.M.Wang,^{17,19} C.Warner,¹ A.Weber,¹ J.Weber,⁴⁶ R.Weill,²⁰ T.J.Wenaus,²¹ J.Wenninger,¹⁸ M.White,¹⁴ C.Willmott,²⁴ F.Wittgenstein,¹⁶ D.Wright,³⁴ S.X.Wu,¹⁷ S.Wynhoff,¹ B.Wysłouch,¹⁴ Y.Y.Xie,³⁹ J.G.Xu,⁶ Z.Z.Xu,¹⁹ Z.L.Xue,³⁹ D.S.Yan,³⁹ B.Z.Yang,¹⁹ C.G.Yang,⁶ G.Yang,¹⁷ C.H.Ye,¹⁷ J.B.Ye,¹⁹ Q.Ye,¹⁷ S.C.Yeh,⁴⁸ Z.W.Yin,³⁹ J.M.You,¹⁷ N.Yunus,¹⁷ M.Yzerman,² C.Zaccardelli,³¹ N.Zaitsev,²⁷ P.Zemp,⁴⁶ M.Zeng,¹⁷ Y.Zeng,¹ D.H.Zhang,² Z.P.Zhang,^{19,17} B.Zhou,⁹ G.J.Zhou,⁶ J.F.Zhou,¹ R.Y.Zhu,³¹ A.Zichichi,^{7,16,17} B.C.C.van der Zwaan.²

-
- 1 I. Physikalisches Institut, RWTH, W-5100 Aachen, FRG[§]
 III. Physikalisches Institut, RWTH, W-5100 Aachen, FRG[§]
 2 National Institute for High Energy Physics, NIKHEF, NL-1009 DB Amsterdam, The Netherlands
 3 University of Michigan, Ann Arbor, MI 48109, USA
 4 Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941
 Annecy-le-Vieux CEDEX, France
 5 Johns Hopkins University, Baltimore, MD 21218, USA
 6 Institute of High Energy Physics, IHEP, 100039 Beijing, China
 7 INFN-Sezione di Bologna, I-40126 Bologna, Italy
 8 Tata Institute of Fundamental Research, Bombay 400 005, India
 9 Boston University, Boston, MA 02215, USA
 10 Northeastern University, Boston, MA 02115, USA
 11 Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania
 12 Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114,
 Hungary[‡]
 13 Harvard University, Cambridge, MA 02139, USA
 14 Massachusetts Institute of Technology, Cambridge, MA 02139, USA
 15 INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy
 16 European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland
 17 World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland
 18 University of Geneva, CH-1211 Geneva 4, Switzerland
 19 Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China
 20 University of Lausanne, CH-1015 Lausanne, Switzerland
 21 Lawrence Livermore National Laboratory, Livermore, CA 94550, USA
 22 Los Alamos National Laboratory, Los Alamos, NM 87544, USA
 23 Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne
 Cedex, France
 24 Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas, CIEMAT, E-28040 Madrid,
 Spain
 25 INFN-Sezione di Milano, I-20133 Milan, Italy
 26 Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
 27 INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
 28 Department of Natural Sciences, University of Cyprus, Nicosia, Cyprus
 29 University of Nymegen and NIKHEF, NL-6525 ED Nymegen, The Netherlands
 30 Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA
 31 California Institute of Technology, Pasadena, CA 91125, USA
 32 INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
 33 Carnegie Mellon University, Pittsburgh, PA 15213, USA
 34 Princeton University, Princeton, NJ 08544, USA
 35 INFN-Sezione di Roma and University of Rome, "La Sapienza", I-00185 Rome, Italy
 36 Nuclear Physics Institute, St. Petersburg, Russia
 37 University of California, San Diego, CA 92093, USA
 38 Dept. de Fisica de Particulas Elementales, Univ. de Santiago, E-15706 Santiago de Compostela, Spain
 39 Shanghai Institute of Ceramics, SIC, Shanghai, China
 40 Bulgarian Academy of Sciences, Institute of Mechatronics, BU-1113 Sofia, Bulgaria
 41 Center for High Energy Physics, Korea Advanced Inst. of Sciences and Technology, 305-701 Taejon,
 Republic of Korea
 42 University of Alabama, Tuscaloosa, AL 35486, USA
 43 Purdue University, West Lafayette, IN 47907, USA
 44 Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
 45 DESY-Institut für Hochenergiephysik, O-1615 Zeuthen, FRG
 46 Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
 47 University of Hamburg, W-2000 Hamburg, FRG
 48 High Energy Physics Group, Taiwan, China
 § Supported by the German Bundesministerium für Forschung und Technologie
 ‡ Supported by the Hungarian OTKA fund under contract number 2970.

References

- [1] S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264.
A. Salam, Nobel Symposium No. 8, ed. N.Svartholm, Almqvist and Wiksell, (1968) 367.
- [2] S. L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D2 (1970) 1285.
- [3] G. Mann and T. Riemann, Annalen Phys. (Leipzig) 40 (1983) 334. B187 (1987) 303.
- [4] T. K. Kuo and N. Nakagawa, Phys. Rev. D32 (1985) 306.
- [5] J. Bernabeu, A. Santamaria, J. Vidal, A. Mendez and J. W. F. Valle, Phys. Lett. B187 (1987) 303.
- [6] G. Eilam and T .G. Rizzo, Phys. Lett. B188 (1987) 91.
- [7] M. J. S. Levine, Phys. Rev. D36 (1987) 1329.
- [8] J. Bernabeu and A. Santamaria, Phys. Lett. B197 (1987) 418.
- [9] M. Dittmar and J. W. F. Valle, Preprint FTUV/90-19 B187 (1987) 303.
- [10] L3 Collaboration, B. Adeva et al., Physics Letters B **271** (1991) 453.
- [11] The L3 Collaboration, *Results from the L3 Experiment at LEP*, CERN-PPE/93-31, submitted to Physics Reports.
- [12] CLEO Collaboration, T. Bowcock et al., Phys. Rev. D41 (1990) 805.
- [13] ARGUS Collaboration, H. Albrecht et al., DESY 92-013.
- [14] Sindrum Collaboration, U. Bellgardt *et al.*, Nucl. Phys. B299 (1988) 1.
- [15] The OPAL Collaboration, M. Z. Akrawy et al., Phys. Lett. B 254 (1991) 293.
- [16] ALEPH Collab., D. Decamp et al., Phys. Rep. 216 (1992) 253.
- [17] DELPHI Collab., P. Abreu et al., Phys. Lett. B 298 (1992) 247.
- [18] S. Jadach et al., “Z Physics at LEP 1”, eds. G. Altarelli et al., CERN Report CERN-89-08, Vol. 2 (1989) 35.
- [19] L3 Collaboration, B. Adeva et al., Nuclear Instruments and Methods A **289** (1990) 35.
- [20] S. Jadach *et al.*, “Z Physics at LEP 1”, eds. G. Altarelli et al., CERN Report CERN-89-08, Vol. 3 (1989) 69.
- [21] M. Böhm, A. Denner and W. Hollik, Nucl. Phys. B304 (1988) 687;
F.A. Berends, R. Kleiss and W. Hollik, Nucl. Phys. B304 (1988) 712.
- [22] T. Sjöstrand and M. Bengtsson, Comput. Phys. Commun. 43 (1987) 367;
T. Sjöstrand, “Z Physics at LEP 1”, eds. G. Altarelli et al., CERN Report CERN-89-08, Vol. 3 (1989) 143.

- [23] J. Hilgart, R. Kleiss and F. Le Diberder, *Comput. Phys. Commun.* **75** (1993) 191.
- [24] O. Adriani et al., *Nucl. Instr. and Meth. A* **302** (1991) 53.
- [25] L3 Collaboration, O. Adriani et al., *Phys. Lett. B* **303** (1993) 391.

Figure Captions

Fig. 1 Muon energy versus electron energy for selected $Z \rightarrow e\mu$ candidate events. The shaded area represents the region corresponding to the signal events.

Fig. 2 The distribution of acoplanarity angle between two opposite TEC tracks in the dilepton events. The points are data, the histogram corresponds the $Z \rightarrow ee(\gamma)$ Monte Carlo events and the dashed line shows the distribution for the signal $Z \rightarrow e\tau$ Monte Carlo events. The normalisation for the signal distribution is arbitrary. The arrows indicate the cuts.

Fig. 3 The distribution of electron energy after all cuts but the electron energy cut have been applied in the $Z \rightarrow e\tau$ selection.

Fig. 4 The distribution of muon energy after all cuts but the muon energy cut have been applied in the $Z \rightarrow \mu\tau$ selection.




Figure 1:




Figure 2:




Figure 3:




Figure 4: