

BB

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN LIBRARIES, GENEVA

CERN-PPE/94-010 /  
26.01.1994



CM-P00070276

# NUCLEAR PROPERTIES OF THE EXOTIC HIGH- SPIN ISOMER $^{178m2}\text{Hf}$ FROM COLLINEAR LASER SPECTROSCOPY

N. Boos<sup>1</sup>, F. Le Blanc<sup>2</sup>, M. Krieg<sup>1</sup>, J. Pinard<sup>3</sup>, G. Huber<sup>1</sup>, M.D. Lunney<sup>2</sup>, D. Le Du<sup>4</sup>, R. Meunier<sup>4</sup>, M. Hussonnois<sup>2</sup>, O. Constantinescu<sup>2</sup>, J.B. Kim<sup>2</sup>, Ch. Briançon<sup>4</sup>, J.E. Crawford<sup>6</sup>, H.T. Duong<sup>3</sup>, Y.P. Gangrski<sup>7</sup>, T. Kühl<sup>5</sup>, B.N. Markov<sup>7</sup>, Yu.Ts. Oganessian<sup>7</sup>, P. Quentin<sup>4</sup>, B. Roussière<sup>2</sup> and J. Sauvage<sup>2</sup>

## ABSTRACT

The complete hyperfine spectrum in the optical transition  $5d^26s^2\ ^3P_2 \rightarrow 5d6s^26p\ ^1P_1$  was recorded for  $^{178m2}\text{Hf}$  by collinear laser spectroscopy using nanogram samples. The quadrupole moment and isomer shift were determined for the first time as well as a precise value and the sign of the magnetic dipole moment. The change in nuclear mean-square charge radius between the isomeric state  $^{178m2}\text{Hf}$  and the ground state  $^{178g}\text{Hf}$  was evaluated as  $\delta \langle r^2 \rangle^{178,178m2} = -0.059(9)\text{ fm}^2$ . From the hyperfine A- and B-factors the magnetic moment  $\mu_I^{178m2} = +8.16(4)\text{ n.m.}$  and the spectroscopic quadrupole moment  $Q_s^{178m2} = +6.00(7)\text{ b}$  were extracted.

(IS315)

Submitted to Phys. Rev. Lett.

- 
- 1) Institut für Physik, Universität Mainz, 55099 Mainz, Germany
  - 2) Institut de Physique Nucléaire, IN2P3-CNRS, 91406 Orsay, France
  - 3) Laboratoire Aimé Cotton, 91405 Orsay, France,
  - 4) CSNSM-Orsay, IN2P3-CNRS, 91405 Orsay, France
  - 5) GSI, 64291 Darmstadt, Germany
  - 6) Foster Radiation Laboratory, McGill University, Montréal, Canada
  - 7) JINR Dubna, POB 79, 101000 Moscow Region, Russia

# Nuclear properties of the exotic high-spin isomer $^{178m2}\text{Hf}$ from collinear laser spectroscopy

N. Boos<sup>1</sup>, F. Le Blanc<sup>2</sup>, M. Krieg<sup>1</sup>, J. Pinard<sup>3</sup>, G. Huber<sup>1</sup>, M.D. Lunney<sup>2</sup>, D. Le Du<sup>4</sup>, R. Meunier<sup>4</sup>, M. Hussonnois<sup>2</sup>, O. Constantinescu<sup>2</sup>, J.B. Kim<sup>2</sup>, Ch. Briançon<sup>4</sup>, J.E. Crawford<sup>6</sup>, H.T. Duong<sup>3</sup>, Y.P. Gangrski<sup>7</sup>, T. Kühl<sup>5</sup>, B.N. Markov<sup>7</sup>, Yu.Ts. Oganessian<sup>7</sup>, P. Quentin<sup>4</sup>, B. Roussière<sup>2</sup> and J. Sauvage<sup>2</sup>

<sup>(1)</sup> *Institut für Physik der Universität Mainz, 55099 Mainz, Germany*

<sup>(2)</sup> *Institut de Physique Nucléaire, IN2P3-CNRS, 91406 Orsay, France*

<sup>(3)</sup> *Laboratoire Aimé Cotton, 91405 Orsay, France*

<sup>(4)</sup> *CSNSM-Orsay, IN2P3-CNRS, 91405 Orsay, France*

<sup>(5)</sup> *GSI, 64291 Darmstadt, Germany*

<sup>(6)</sup> *Foster Radiation Laboratory, McGill University, Montréal, Canada*

<sup>(7)</sup> *JINR Dubna, POB 79, 101000 Moscow Region, Russia*

The complete hyperfine spectrum in the optical transition  $5d^26s^2\ ^3P_2 \rightarrow 5d6s^26p\ ^1P_1$  of  $^{178m2}\text{Hf}$  was recorded by collinear laser spectroscopy using nanogram samples. The quadrupole moment and isomer shift were determined for the first time as well as a precise value and the sign of the magnetic dipole moment. The change in nuclear mean-square charge radius between the isomeric state  $^{178m2}\text{Hf}$  and the ground state  $^{178g}\text{Hf}$  was evaluated as  $\delta\langle r^2 \rangle^{178,178m2} = -0.059(9)\text{ fm}^2$ . From the hyperfine A- and B-factors the magnetic moment  $\mu_I^{178m2} = +8.16(4)\text{ n.m.}$  and the spectroscopic quadrupole moment  $Q_s^{178m2} = +6.00(7)\text{ b}$  were extracted.

PACS numbers: 21.10.Ky, 27.70.+q, 32.30.Jc, 35.10.Fk, 42.62.Fi

A large number of high-spin isomers are observed in the region of the rare earth elements and beyond. Their high-spin values are interpreted as due to the alignment of several nucleon angular momenta. In this respect, the nucleus  $^{178}\text{Hf}$  is very interesting since it shows the four quasi-particle configuration  $^{178m2}\text{Hf}$  with spin  $I^\pi=16^+$  [1] and the extremely long half-life  $T_{1/2} = 31$  y [2]. Since this state has the lowest excitation energy  $E=2447.4$  keV for its angular momentum  $I$  and is located even below the  $I = 14$  levels, it cannot decay by low multipolarity transitions. The exceptionally long half-life allows to prepare ng samples and targets of this excited nuclear matter for the present optical measurements as well as for inelastic scattering and coulomb excitation experiments [3]. Thus, the nuclear properties can be studied in two states: the isomer with its four unpaired nucleons and the ground state with its fully paired structure ( $I=0$ ).

In the decay of the  $I^\pi = 16^+$  state a  $T_{1/2}=4$  s isomer at 1147.4 keV is populated and a second  $I^\pi = 8^-$  level is located at 1479.0 keV. They are interpreted as mixtures of the two proton configuration  $\{p[514]9/2 + p[404]7/2\}I^\pi = 8^-$  ( $K=8$ ,  $K$  being the projection of  $\vec{I}$  on the nuclear symmetry axis) and the two neutron configuration  $\{n[514]7/2 + n[624]9/2\}I^\pi = 8^-$  ( $K=8$ ) [2, 4]. The sum of excitation energies of these two states is close to the energy of the long-lived isomer. This suggests that the  $I^\pi = 16^+$  ( $K=16$ ) state is a superposition of the two  $8^-$  configurations and thus of a pure four quasi-particle nature.

In this letter, we present optical measurements on  $^{178m2}\text{Hf}$  together with corresponding systematic results obtained for the stable isotopes  $^{176-180}\text{Hf}$ . The measurements provide the A- and B-factors for the atomic levels of the transition which allow determination of the magnetic moment  $\mu_I$  and the spectroscopic quadrupole moment  $Q_s$  [5]. The isomer shift (IS) between the center-of-gravity of the hyperfine spectrum (HFS) of the isomeric state and the ground state resonance line allows determination of the change in the mean-square nuclear charge radius  $\delta\langle r^2 \rangle^{178,178m2}$  between the isomer and the ground state. These results include the first measurement of the quadrupole moment and the isomer shift as well as a more precise value and the sign of the magnetic dipole moment of  $^{178m2}\text{Hf}$ . The measurements are particularly challenging due to the extremely low production rate of the exotic Hf isomer that results in a microscopic sample.

The Hf isomer was produced in the reaction  $^{176}\text{Yb}(\alpha,2n)$  by bombarding a 96% enriched Yb target with a 36 MeV  $\alpha$ -beam provided by the U-200 cyclotron at Dubna

[6]. The two samples for our experiments were chemically separated and deposited on a quartz substrate. Each of the samples contained about  $2 \cdot 10^{13}$  atoms or 6 ng of  $^{178m2}\text{Hf}$  and a roughly 30 times larger amount of  $^{178g}\text{Hf}$  as deduced from the optical signals. By chlorination the sample was volatilized and fed into the ion source of the P.A.R.I.S. mass separator [7] at the CSNSM at Orsay. The 40 keV elemental ion beam was mass separated and transported to a collinear laser spectroscopy apparatus where the ions were neutralized in a charge exchange cell with sodium vapor and merged with a counterpropagating laser beam of typically 15 mW from a single-mode dye laser. This dye laser was stabilized to a molecular absorption line of  $\text{I}_2$ . Spectroscopy was performed by exciting the  $5d^26s^2 \ ^3P_2$  ( $8983.75 \text{ cm}^{-1}$ )  $\rightarrow$   $5d6s^26p \ ^1P_1$  ( $26463.93 \text{ cm}^{-1}$ ) transition ( $\lambda=572 \text{ nm}$ ) and then detecting the fluorescence of the subsequent decay through the strong resonance line ( $\lambda=378 \text{ nm}$ ) to the  $5d^26s^2 \ ^3F_2$  ground state. The hyperfine resonances were scanned via the beam velocity by applying a sweep voltage to the charge exchange cell. An axial magnetic field of  $2 \cdot 10^{-2} \text{ T}$  was applied to the charge exchange cell and a subsequent drift space in order to suppress optical pumping outside the detection zone. A large solid angle ellipsoidal mirror system has been used to collect the fluorescence light to a photon counting RCA8850 photomultiplier. An interference filter and a Schott BG3 colour glass were inserted between the collection system and the entrance window of the photomultiplier. Details are given in [8] but special care has been taken in order to reduce beam background signal due to scattering of the strong  $^{178g}\text{Hf}$  beam to the detection system. A signal/background ratio of 400 has been achieved for stable even isotopes.

The measured hyperfine spectrum of  $^{178m2+g}\text{Hf}$  is shown in fig. 1. The single peak corresponds to the  $I^\pi=0^+$  ground state. The positions were fitted to the HFS formula [9] yielding the hyperfine A- and B-factors for the lower  $^3P_2$  and upper  $^1P_1$  level as well as the isomer shift in the optical transition. In addition we have recorded hyperfine spectra of this transition for the stable Hf isotopes in order to determine their A- and B-factors and isotope shift needed for the extraction of the atomic parameters. The results are listed in tab. 1.

The IS can be decomposed into the normal mass shift (NMS), the specific mass shift (SMS) and the field shift (FS). The FS is expressed by  $FS^{A,A'} = F_i \cdot \lambda^{A,A'}$  where  $F_i$  is the electronic factor for the transition and the nuclear parameter  $\lambda^{A,A'} = K \cdot \delta \langle r^2 \rangle^{A,A'}$ , where

$\delta\langle r^2 \rangle^{A,A'}$  stands for the change in the mean-square charge radius of the two nuclei under consideration (for Hf,  $K=0.95$  [10]). The nuclear parameter  $\lambda^{178,180}$  has been obtained by a parametric analysis of 33 transitions in HfII [11] to be  $\lambda^{178,180} = +0.072(4)fm^2$  which, together with the model independent relative nuclear parameters  $\lambda_{rel}$  [12], produces the values in tab. 1 for the stable isotopes.  $SMS_{572nm}$  and  $F_{572nm}$  have been determined by a King plot [13], which relates the IS for one transition to that of another; in this case the 618nm transition of HfI [14].  $SMS_{618nm}$  and  $F_{618nm}$  were, in turn, evaluated by King plot procedures using 13 transitions between the ground state multiplet  $5d^26s^2\ ^3F$  and  $5d^26s6p$  configurations investigated by Cajko [15]. Since these configurations are pure [16], the SMS in these transitions can be estimated to be nearly zero [12]. Their electronic factors have been obtained by a fit of the measured  $IS^{A,A'}$  to the  $\lambda^{A,A'}$  from tab. 1. Finally, we determined  $F_{572nm} = +2396(347) MHz/fm^2$  and  $SMS_{572nm} = -3.00(1.09) \cdot NMS_{572nm}$ . (There are muonic data available [17] for the calibration of the electronic factor.  $F_{572nm}$  calibrated in this way differs by about  $-35\%$  from the above value.) For  $^{178}\text{Hf}$ , the change in the mean square charge radius between the fundamental and the isomeric state is thus determined to be

$$\delta\langle r^2 \rangle^{178,178m2} = -0.059(9)fm^2 \quad (1)$$

which almost corresponds to the subtraction of a neutron pair. The isotonic  $^{177,177m}\text{Lu}$  also exhibits a negative isomer shift but of a lower absolute value [18].

The absolute value of  $\delta\langle r^2 \rangle^{178,178m2}$  and its negative sign are very difficult to interpret. Due to the different pairing properties for the  $0^+$  and  $16^+$  states, there are a variety of contributions to this quantity. These contributions include monopole, quadrupole, hexadecapole and higher moments as well as coupling between the various modes. Moreover, current theoretical approaches generally do not deal with wavefunctions projected on good angular momentum values which immediately make them unsuitable for the description of small variations.

From the magnetic moments of  $^{177,179}\text{Hf}$  [19] and the  $A(^3P_2)$ -Factors in tab. 1 the hyperfine anomaly [20] is found to be  $^{177}\Delta^{179}(^3P_2) = -3.6(7.5) \cdot 10^{-3}$ , which is comparable to the corresponding value for the ground state  $^{177}\Delta^{179}(^3F_2) = -2.4(2.2) \cdot 10^{-3}$  [21]. Thus, neglecting hyperfine anomaly with respect to experimental uncertainties, the magnetic moment of  $^{178m2}\text{Hf}$  can be scaled according to the relation  $A = \mu_I \cdot \langle H(0) \rangle / (IJ)$ . With

the accurate value for  $\mu_I^{177}$  [19] and the A-factors from tab. 1 we obtain

$$\mu_I^{178m2} = +8.16(4)n.m. \quad (2)$$

An earlier rough measurement via nuclear orientation [22] yielded

$$|\mu_I^{178m2}| = 7.26(+20)(-00)n.m.$$

The magnetic moment of a nuclear state is mainly due to its single particle properties. The modified oscillator model [23] coupling the  $g_K$  values of the nuclear states involved has given a theoretical value of  $\mu_{theo}^{178m2} = +7.81$  n.m. Combining the experimental moments of neighbouring odd nuclei by the coupling rules given in [23] yields a semiempirical value  $\mu_{s.e.}^{178m2} = +8.17$  n.m. in excellent agreement with the experimental result (2), which further confirms the 4 quasi-particle assignment for the isomeric state [2].

For  $^{177}\text{Hf}$ , a spectroscopic quadrupole moment  $Q_s = 3.365(29)b$  has been obtained from muonic data [19], which is comparable to  $Q_s = 3.30(65)b$  (including Sternheimer correction) from ABMR measurements [20]. To determine  $Q_s$  for the isomer we use the first value for  $^{177}\text{Hf}$  and scale with the  $B(5d^26s^2^3P_2)$ -factors (tab. 1) to obtain

$$Q_s^{178m2} = +6.00(7)b \quad (3)$$

Assuming the strong coupling scheme for evaluating the intrinsic quadrupole moment from the measured  $Q_s$  (3), we obtain a value of  $Q_0=7.2(1)b$  for the  $I^\pi=16^+$  state. This is slightly higher than  $Q_0=6.961(43)b$  for the ground state derived from  $B(E2)$  values [24]. Theoretical calculations [25, 26] have predicted  $Q_0=7.3b$  for the ground state with very little variation for the  $16^+$  isomeric state.

Our collinear laser spectroscopy experiment has resulted in the determination of the magnetic dipole and electric quadrupole moment as well as the isomer shift of the high-spin isomer  $^{178m2}\text{Hf}$ . While the magnetic moment can be determined mainly by the single-particle nature of the nuclear states under consideration, the explanation for the negative isomer shift (and thus the change in mean-square charge radius) needs detailed microscopic calculations incorporating higher orders of deformation.

## References

- [1] T. L. Khoo, G. Løvholden; Phys. Lett. 67 B (1977), 271

- [2] R. G. Helmer, C. W. Reich; Nucl. Phys. A 114 (1968), 649  
R. G. Helmer, C. W. Reich; Nucl. Phys. A 211 (1973), 1
- [3] Ch. Briançon, M. Aiche, D. Le Du, R. Meunier, P. Quentin, Ch. Vieu, M. Hussonnois, J.B. Kim, D. Trubert, F. Le Blanc, M.D. Lunney, B. Roussière, J. Sauvage, G. Rotbard, G. Berrier-Ronsin, S. Fortier, S. Gales, J.M. Maison, L.H. Rosier, J. Vernotte, Yu.Ts. Oganessian, O. Constantinescu, Yu.P. Gangrski, B. Gorski, S.A. Karamian, B.N. Markov, Z. Szegłowski, V.M. Gorozhankin, V.B. Brudanin, V.G. Egorov, O.I. Kochetov, V.N. Pavlov, J. Pinard, H.T. Duong, N. Boos, M. Krieg, G. Huber, R. Kulesa, H.J. Wollersheim, H. Folger, J. Gerl, T. Happ, T. Kühl, G. Graw, D. Hofer, E. Müller-Zanotti, B.D. Valnion, J. De Boer, A. Gollwitzer, N. Gollwitzer, M. Wurfner, W.D. Hamilton, D. Mosbah, H.V. Muradian; *Proceedings of the Eighth International Symposium on "Capture Gamma-Ray Spectroscopy"*, Fribourg, Switzerland, Sept. 20-24, 1993 (to be published)
- [4] V. Soloviev, A. Shuskov; J. Phys. G 16 (1990), L57 and references quoted therein
- [5] E. W. Otten; in *Treatise on Heavy-Ion Physics*, ed. D. A. Bromley, Plenum Press, New York (1988)
- [6] Yu. Oganessian, S. Karamian, Y. Gangrski, B. Gorski, B. Markov, Z. Szegłowski, Ch. Briançon, D. Le Du, R. Meunier, M. Hussonnois, O. Constantinescu, M. Subbotin, J. Phys. G18 (1992), 393
- [7] J. Camplan, R. Meunier, C. Fatu; in *Proceedings of the 8th International EMIS conference*, ed. G. Andersson, G. Holmèn, Skörde, Sweden, June 12-15, 1973, 186
- [8] T. Kühl, R. Kirchner, O. Klepper, D. Marx, U. Dinger, J. Eberz, G. Huber, H. Lochmann, R. Menges, G. Ulm; Nucl. Instr. Meth. B 26 (1987), 419
- [9] H. Kopfermann; *Kernmomente*, Akademische Verlagsgesellschaft Frankfurt a. M. (1958)
- [10] G. Torbohm, B. Fricke, A. Rosén; Phys. Rev. A 31, 4 (1985), 2038
- [11] P. Aufmuth, I. Henneberg, A. Siminski, A. Steudel; Z. Phys. D 18 (1991), 107
- [12] P. Aufmuth, K. Heilig, A. Steudel; At. Data and Nucl. Tables 37 (1987), 455

- [13] W. H. King; *Isotope Shifts in Atomic Spectra*, Plenum Press, New York (1984)
- [14] J. Schecker, A. Berger, J. Das, S. Dutta, G. Gwinner, C. Holbrow, T. Kühl, T. Lauritsen, G. Sprouse, F. Xu; *Phys. Rev. A* 46 (1992), 3730
- [15] J. Cajko; *Z. Phys.* 234 (1970), 443
- [16] P. Aufmuth, R. Kirsch, A. Steudel, E. Wöbker; *Z. Phys. D* 7 (1987), 153
- [17] Y. Tanaka, R. Steffen, E. Shera, W. Reuter, M. Hoehn, J. Zumbro; *Phys. Rev. C* 30 (1984), 350
- [18] W. Borchers, U. Georg, A. Klein, P. Lievens, R. Neugart, M. Neuroth, Pushpa M. Rao, C. Schulz; in *Proc. Int. Conf. NFFS-6 and AMCO-9, IOP Conf. Ser. 132*, ed. R. Neugart, A. Wöhr, Bernkastel-Kues, Germany, July 19-24, 1992, 213
- [19] P. Raghavan; *At. Data and Nucl. Tables* 42 (1989), 189
- [20] S. Büttgenbach; *Hyperfine Structure in 4d- and 5d-Shell Atoms*, Springer Tracts in Modern Physics 96, Springer, Berlin, Heidelberg, New York (1982)
- [21] S. Büttgenbach, M. Herschel, G. Meisel, E. Schrödl, W. Witte; *Z. Phys.* 260 (1973), 157
- [22] H. Postma, B. Kasteleinen, N. Severijns, D. Vandeplassche, J. Vanhaverbeke, L. Vanneste, E. van Walle, J. Wouters, J. van Klinken; *Hyp. Int.* 52 (1989), 79
- [23] C. Ekström, H. Rubinsztein, P. Möller; *Physica Scripta* 14 (1976), 199
- [24] S. Raman, C. Malarkey, W. Milner, S. Kahane, P. Stelson; *At. Data and Nucl. Tables* 36 (1987), 1
- [25] P. Quentin, S. Krieger, J. Libert, M. Weiss, in *Nuclear Shapes and Nuclear Structure at low Excitation Energies*, ed. M. Vergnes, J. Sauvage, P. Heenen, H.T. Duong, NATO ASI Series B289 (1992) 163
- [26] J. P. Delaroche, private communication (1993)






Figure 1: Experimental hyperfine spectrum of  $^{178}\text{Hf}$ . The spectrum was recorded over four different ranges as shown. The single peak at 0 MHz having an amplitude of 14000 counts corresponds to the ground state  $^{178g}\text{Hf}$ . The nine hyperfine resonances of  $^{178m2}\text{Hf}$  are marked by arrows.

Table 1:

A (amu)	$I^\pi$	$A(^3P_2)$ [MHz]	$A(^1P_1)$ [MHz]	$B(^3P_2)$ [MHz]	$B(^1P_1)$ [MHz]	$IS^{178,A}$ [MHz]	$\lambda^{178,A}$ [fm <sup>2</sup> ]
176	$0^+$	-	-	-	-	-121.9(5.3)	-0.063(4)
177	$7/2^-$	+70.9(4)	-21.9(6)	-1206.5(3.7)	+687.7(2.0)	-87.1(3.0)	-0.045(3)
178	$0^+$	-	-	-	-	0.00	0.000
178m2	$16^+$	+159.5(2)	-51.3(4)	-2150.7(16.2)	+1234.8(9.6)	-134.6(3.1)	-0.056(8)
179	$9/2^+$	-44.7(2)	+13.9(2)	-1364.0(2.4)	+776.3(1.5)	-	+0.027(2)
180	$0^+$	-	-	-	-	+133.9(5.1)	+0.072(4)

Table 1: Hyperfine constants of the  $5d^26s^2\ ^3P_2$  and  $5d6s^26p\ ^1P_1$  atomic states, the isotope shift  $IS^{178,A}$  and the nuclear parameter  $\lambda^{178,A}$  for  $^{178m2}\text{Hf}$  and the stable isotopes  $^{176-180}\text{Hf}$