

LHC limits on axion-like particles from heavy-ion collisions

Simon Knapen^{1,2}, Tongyan Lin^{1,2,3}, Hou Keong Lou^{1,2} and Tom Melia^{1,2,4}

¹ Department of Physics, University of California, Berkeley, California 94720, USA

² Theoretical Physics Group, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

³ Department of Physics, University of California, San Diego, California 92093, USA

⁴ Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo Institutes for Advanced Study, University of Tokyo, Kashiwa 277-8583, Japan

Abstract

In these proceedings we use recent LHC heavy-ion data to set a limit on axion-like particles coupling to electromagnetism with mass in the range 10-100 GeV. We recast ATLAS data as per the strategy proposed in [1], and find results in-line with the projections given there.

Keywords

CERN report; axion-like particle; heavy-ion, ultra-peripheral collisions

1 Introduction

The LHC has completed its highest luminosity heavy-ion collision run (Pb-Pb), with ATLAS, CMS and ALICE all recording data at a centre-of-mass energy per nucleon of $\sqrt{s_{NN}} = 5.02$ TeV. In previous work [1] we showed that the large charge of the lead ions ($Z = 82$) results in a huge Z^4 coherent enhancement in the exclusive production of axion-like particles (ALPs) that couple to electromagnetism, which can lead to competitive limits for ALPs. This proceeding is an update to our previous work; we recast the analysis of the ATLAS $480 \mu\text{b}^{-1}$ data set [2] to provide limits on ALPs in the mass region $10 \text{ GeV} < m_a < 100 \text{ GeV}$. In line with the projections in [1], we find that the LHC heavy-ion data provides the strongest limits to date in this parameter range. While the physics potential of exclusive heavy ion collisions has been known for decades [3–5], to our knowledge this represents the first time LHC heavy-ion data sets the most stringent limit on a specific beyond the Standard Model physics scenario.

Ultra-peripheral collisions (UPCs) are quasi-elastic processes where the impact parameter is much greater than the ion radius (see *e.g.* Refs. [6–8]). The ions remain (largely) intact, and there is a large rapidity gap between any produced particle and the beam-line with very little detector activity. This clean environment, along with the Z^4 enhanced signal rate, provides a low background ALP search channel that can perform better than searches using the p-p run.

The production of an ALP in a UPC proceeds via photon fusion—see Fig. 1—where we consider a Lagrangian of the form

$$\mathcal{L}_a = \frac{1}{2}(\partial a)^2 - \frac{1}{2}m_a^2 a^2 - \frac{1}{4} \frac{a}{\Lambda} F \tilde{F}, \quad (1)$$

where $\tilde{F}^{\mu\nu} \equiv \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}/2$, and with a being the pseudoscalar ALP of mass m_a which couples to electromagnetism via the dimensionful coupling $1/\Lambda$. Such a coupling can be obtained through the $SU(2)_L$ invariant operator $-aB\tilde{B}/(4\cos^2\theta_W\Lambda)$ where B is the hypercharge field strength. Polarization effects of the incoming photons can lead to different scalar and pseudoscalar production rates, but the effects are relatively small when integrating over all impact parameters [9, 10]. Our limits therefore apply for scalar particles through the replacement $\tilde{F}(\tilde{B}) \rightarrow F(B)$ in Eq. (1).

Here we recap some of the details of our treatment in [1]. The ALP parameter space is already substantially constrained by cosmological and astrophysical observations, as well as by a broad range of collider and intensity frontier experiments (see *e.g.* [11, 12] for reviews and recent results). In the regime of interest for UPCs ($1 \text{ GeV} \lesssim m_a \lesssim 100 \text{ GeV}$), the existing constraints however come from




Fig. 1: Exclusive ALP production in ultra-peripheral Pb-Pb collisions.

LEP and LHC [13–15]. In Fig. 2, we show the expected sensitivity from performing a bump hunt in $m_{\gamma\gamma}$ for UPCs, assuming a luminosity for the current (1 nb^{-1}) and the high luminosity (10 nb^{-1}) Pb-Pb runs.¹ For each mass point we computed the expected Poisson limit [16]. The dominant backgrounds are estimated to be light-by-light scattering [3] and fake photons from electrons, and become negligible for $m_{\gamma\gamma} \gtrsim 20 \text{ GeV}$. In the region which there is background, we assume the entire signal falls into a bin of width 1 GeV. The signal selection criteria in this case are $E_T > 2 \text{ GeV}$ and $|\eta| < 2.5$ for the two photons and $|\phi_{\gamma\gamma} - \pi| < 0.04$. The analogous limit from the exclusive p-p analysis performed by CMS [17] is also shown, which is very weak due to low photon luminosities. For the $F\tilde{F}$ operator the heavy-ion limits are significantly stronger, whereas for the $B\tilde{B}$ operator, traditional p-p collider limits are enhanced due to additional production channels through the Z coupling.

Light-by-light scattering has been measured by the ATLAS collaboration [2], and the results were consistent with our estimates and those in earlier computations [18–20]. Using the observed $m_{\gamma\gamma}$ spectrum, we then derive an observed limit on ALPs for $F\tilde{F}$ and $B\tilde{B}$ couplings, which are shown in black in Fig. 2. In detail, we generated Monte Carlo samples for the ALP signal using a modified version of the STARlight code [21],² which assigns a small virtuality to the photons and as such leads to a typical $p_T^{\gamma\gamma} \lesssim 100 \text{ MeV}$ for the recoil of the $\gamma\gamma$ -system. We then follow the ATLAS analysis and apply the following selection cuts on the signal:

1. Require exactly two photons with $E_T > 3 \text{ GeV}$ and $|\eta| < 2.4$
2. Demand $|\phi_{\gamma\gamma} - \pi| < 0.03$, where $\phi_{\gamma\gamma}$ is the azimuthal angle between the two photons

The signal efficiency is $\sim 70\%$ near threshold and becomes fully efficient if the sum of the photon energies exceeds 9 GeV. The selection criteria are slightly different from our previous theoretical analysis, however we note that only the larger E_T cut leads to noticeable changes for the efficiencies. Given that we do not model photon identification at the detector level, we apply an extra total reconstruction efficiency of 90%, which roughly takes into account the per-photon ID efficiency of 95% measured by ATLAS.

The $m_{\gamma\gamma}$ spectrum measured by ATLAS is plotted in bin-widths of 3 GeV, starting at $m_{\gamma\gamma} = 6 \text{ GeV}$. For our exclusion, we generated samples with $m_{\gamma\gamma} = 7, 10, 13, 16, \dots \text{ GeV}$, and assume that all the events are contained in their respective bins after final selection. We further assume that ATLAS did not observe any events with $m_{\gamma\gamma} \gtrsim 30 \text{ GeV}$. The 95% exclusion limits on the coupling $1/\Lambda$ are obtained assuming only statistical uncertainties. A more detailed CL_s analysis that includes a proper treatment of systematics would yield slightly more conservative limits, and we encourage the experimental community to include such an analysis as it is beyond the scope of our simulation framework.

In summary, we have found that heavy-ion collisions at the LHC can provide the best limits on ALP-photon couplings for $7 \text{ GeV} < m_a < 100 \text{ GeV}$, confirming our previous estimates. The very

¹Limits from the p-Pb runs are not competitive despite their higher luminosity, because of the less advantageous Z^2 scaling of the production rate. Collisions with lighter elements, e.g. Ar-Ar, may set relevant limits if the luminosity could be enhanced by two to three orders of magnitude, as compared to current Pb-Pb run.

²Our patch for ALP production is now included in the latest STARlight release.




Fig. 2: *Left:* We show 95% exclusion limits on the operator $\frac{1}{4} \frac{1}{\Lambda} a F \tilde{F}$ using recent ATLAS results on heavy-ion UPCs [2] (solid black line). The expected sensitivity assuming a luminosity of 1 nb^{-1} (10 nb^{-1}) is shown in solid (dashed) green. For comparison, we also give the analogous limit from 36 pb^{-1} of exclusive p-p collisions [17] (red dot-dash). Remaining exclusion limits are recast from LEP II (OPAL 2γ , 3γ) [22] and from the LHC (ATLAS 2γ , 3γ) [23, 24] (see [1] for details). *Right:* The corresponding results for the operator $\frac{1}{4 \cos^2 \theta_W} \frac{1}{\Lambda} a B \tilde{B}$. The LEP I, 2γ (teal shaded) limit was obtained from [14].

large photon flux and extremely clean event environment in heavy-ion UPCs provides a rather unique opportunity to search for BSM physics.

Acknowledgements

We thank Bob Cahn, Lucian Harland-Lang, Yonit Hochberg, Joerg Jaeckel, Spencer Klein, Hitoshi Murayama, Michele Papucci, Dean Robinson, Sevil Salur, and Daniel Tapia Takaki for useful conversations. SK and TL also thank the participants of the 3rd NPKI workshop in Seoul for useful comments and discussions. This work was performed under DOE contract DE-AC02-05CH11231 and NSF grant PHY-1316783.

References

- [1] S. Knapen, T. Lin, H. K. Lou, and T. Melia, “Searching for Axionlike Particles with Ultraperipheral Heavy-Ion Collisions,” *Phys. Rev. Lett.* **118** (2017) no. 17, 171801, [arXiv:1607.06083 \[hep-ph\]](#).
- [2] ATLAS Collaboration, M. Aaboud *et al.*, “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC,” [arXiv:1702.01625 \[hep-ex\]](#).
- [3] G. Baur and C. A. Bertulani, “ γ - γ physics with peripheral relativistic heavy ion collisions,” *Zeitschrift für Physik A Atomic Nuclei* **330** (Mar, 1988) 77–81. <https://doi.org/10.1007/BF01287267>.
- [4] M. Drees, J. R. Ellis, and D. Zeppenfeld, “Can One Detect an Intermediate Mass Higgs Boson in Heavy Ion Collisions?,” *Phys. Lett.* **B223** (1989) 454–460.
- [5] E. Papageorgiu, “Coherent Higgs Production in Relativistic Heavy Ion Collisions,” *Phys. Rev.* **D40** (1989) 92.
- [6] G. Baur, K. Hencken, D. Trautmann, S. Sadovsky, and Y. Kharlov, “Coherent gamma gamma and gamma-A interactions in very peripheral collisions at relativistic ion colliders,” *Phys. Rept.* **364** (2002) 359–450, [arXiv:hep-ph/0112211 \[hep-ph\]](#).

- [7] C. A. Bertulani, S. R. Klein, and J. Nystrand, “Physics of ultra-peripheral nuclear collisions,” *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 271–310, [arXiv:nucl-ex/0502005 \[nucl-ex\]](#).
- [8] A. J. Baltz, “The Physics of Ultraperipheral Collisions at the LHC,” *Phys. Rept.* **458** (2008) 1–171, [arXiv:0706.3356 \[nucl-ex\]](#).
- [9] M. Greiner, M. Vidovic, J. Rau, and G. Soff, “Impact parameter dependence of the Higgs boson production in ultrarelativistic heavy ion collisions,” *J. Phys.* **G17** (1991) L45–L52.
- [10] G. Baur and L. G. Ferreira Filho, “Coherent particle production at relativistic heavy ion colliders including strong absorption effects,” *Nucl. Phys.* **A518** (1990) 786–800.
- [11] R. Essig *et al.*, “Working Group Report: New Light Weakly Coupled Particles,” in *Community Summer Study 2013: Snowmass on the Mississippi (CSS2013) Minneapolis, MN, USA, July 29-August 6, 2013*. 2013. [arXiv:1311.0029 \[hep-ph\]](#).
- [12] M. Bauer, M. Neubert, and A. Thamm, “Collider Probes of Axion-Like Particles,” [arXiv:1708.00443 \[hep-ph\]](#).
- [13] K. Mimasu and V. Sanz, “ALPs at Colliders,” *JHEP* **06** (2015) 173, [arXiv:1409.4792 \[hep-ph\]](#).
- [14] J. Jaeckel and M. Spannowsky, “Probing MeV to 90 GeV axion-like particles with LEP and LHC,” *Phys. Lett.* **B753** (2016) 482–487, [arXiv:1509.00476 \[hep-ph\]](#).
- [15] J. Jaeckel, M. Jankowiak, and M. Spannowsky, “LHC probes the hidden sector,” *Phys. Dark Univ.* **2** (2013) 111–117, [arXiv:1212.3620 \[hep-ph\]](#).
- [16] **Particle Data Group** Collaboration, K. A. Olive *et al.*, “Review of Particle Physics,” *Chin. Phys.* **C38** (2014) 090001.
- [17] **CMS** Collaboration, S. Chatrchyan *et al.*, “Search for exclusive or semi-exclusive photon pair production and observation of exclusive and semi-exclusive electron pair production in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **11** (2012) 080, [arXiv:1209.1666 \[hep-ex\]](#).
- [18] D. d’Enterria and G. G. da Silveira, “Observing light-by-light scattering at the Large Hadron Collider,” *Phys. Rev. Lett.* **111** (2013) 080405, [arXiv:1305.7142 \[hep-ph\]](#). [Erratum: *Phys. Rev. Lett.* 116, no. 12, 129901 (2016)].
- [19] M. Klusek-Gawenda, P. Lebedowicz, and A. Szczurek, “Light-by-light scattering in ultraperipheral Pb-Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys. Rev.* **C93** (2016) no. 4, 044907, [arXiv:1601.07001 \[nucl-th\]](#).
- [20] M. Klusek-Gawenda, W. Schaefer, and A. Szczurek, “Two-gluon exchange contribution to elastic $\gamma\gamma \rightarrow \gamma\gamma$ scattering and production of two-photons in ultraperipheral ultrarelativistic heavy ion and proton-proton collisions,” [arXiv:1606.01058 \[hep-ph\]](#).
- [21] S. R. Klein, J. Nystrand, J. Seger, Y. Gorbunov, and J. Butterworth, “STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions,” [arXiv:1607.03838 \[hep-ph\]](#).
- [22] **OPAL** Collaboration, G. Abbiendi *et al.*, “Multiphoton production in e^+e^- collisions at $\sqrt{s} = 181$ GeV to 209 GeV,” *Eur. Phys. J.* **C26** (2003) 331–344, [arXiv:hep-ex/0210016 \[hep-ex\]](#).
- [23] **ATLAS** Collaboration, G. Aad *et al.*, “Search for Scalar Diphoton Resonances in the Mass Range 65 – 600 GeV with the ATLAS Detector in pp Collision Data at $\sqrt{s} = 8$ TeV,” *Phys. Rev. Lett.* **113** (2014) no. 17, 171801, [arXiv:1407.6583 \[hep-ex\]](#).
- [24] **ATLAS** Collaboration, G. Aad *et al.*, “Search for new phenomena in events with at least three photons collected in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” *Eur. Phys. J.* **C76** (2016) no. 4, 210, [arXiv:1509.05051 \[hep-ex\]](#).