

Singlet Charge 2/3 Quark hiding the Top: Tevatron and LEP Implications

Wei-Shu Hou

*Department of Physics, National Taiwan University, Taipei, Taiwan 10764, R.O.C.**
Theory Division, CERN, CH-1211, Geneva 23, Switzerland

Hsuan-Cheng Huang

Department of Physics, National Taiwan University, Taipei, Taiwan 10764, R.O.C.

Abstract

If c and t quarks are strongly mixed with a weak singlet charge 2/3 quark, $BR(t \rightarrow \ell\nu + X)$ could be suppressed via the $t \rightarrow cH^0$ mode, thereby the top quark could still hide below M_W , whereas the heavy quark signal observed at the Tevatron is due to the dominantly singlet quark Q . This may occur without affecting the small m_c value. Demanding $m_Q \simeq 175$ GeV and $m_t \lesssim M_W$, we find that $BR(t \rightarrow \ell\nu + X)$ cannot be too suppressed. The heavy quark Q decays via W , H , and Z bosons. The latter can lead to b -tagged $Z + 4$ jet events, while the strong c - Q mixing is reflected in sizable $Q \rightarrow sW$ fraction. $Z \rightarrow t\bar{c}$ decay occurs at tree level and may be at the 10^{-3} order, leading to the signature of $Z \rightarrow \ell\nu b\bar{c}$, all isolated and with large p_T , at 10^{-5} order.

HEP-PH-9409227

Typeset using REVTeX

*Permanent address.

I. INTRODUCTION

Recently, the CDF collaboration has reported [1] some evidence for the production of heavy quarks with mass of order 174 GeV at the Tevatron. The most likely explanation is, of course, the standard model (SM) top quark. However, at present, in principle it is still possible [2] that the signal is due to some other heavy quark, whereas the actual top quark is hiding below M_W . This is because the top quark semileptonic branching ratio (BR) has not yet been measured. If, for some reason, $B_{s.l.} \equiv BR(t \rightarrow b\ell\nu) \ll \frac{1}{9}$, the SM expected value, the top quark may have evaded detection. This can arise basically only through scalar induced interactions [2].

One such scenario was proposed [3] earlier by Mukhopadhyaya and Nandi (MN). Following a suggestion [4] by Barbieri and Hall (BH), MN considered the existence of $SU(2)$ singlet charge $\frac{2}{3}$ quark Q alongside SM fermions. Since GIM is broken, the mixing of Q with up-type quarks induce tree level flavor changing neutral couplings of the SM Higgs boson. If $m_{H^0} < m_t$, $t \rightarrow cH^0$ transitions may trigger the aforementioned mechanism of suppressing $B_{s.l.}$. In a subsequent paper, facing criticisms of “naturalness” [5,6] ($1/m_Q$ suppressions of heavy Q effects), MN retracted, and considered $t \rightarrow cH^0$ dominance to be not very likely [7]. In this letter we study the precise conditions that $t \rightarrow cH^0$ dominance can be realised. We find that this requires Q to be strongly mixed with *both* charm and top, which can occur even with a small m_c eigenvalue. However, we find that although $t \rightarrow cH^0$ can be dominant, it is unlikely to be overwhelmingly dominant. Thus, $t \rightarrow bW^*$ should occur at reduced but still substantial fraction. This offers hope that, even if $m_t < M_W$, the top quark can be uncovered at the Tevatron by a renewed study with existing data.

The heavy quark Q can decay both via W and Z bosons [8], hence it could be the heavy quark observed by CDF. Although one could not explain the larger than expected cross section for 174 GeV quarks, one could plausibly account for the b -tagged $Z+4$ jets events [1]. Eventually, events with $ZZ+2$ jets should start to emerge with increased luminosity [9]. Another point of great phenomenological interest is $Z \rightarrow t\bar{c}$ decays, first stressed in

this context by BH [4]. These occur at tree level again because of GIM violation. Although widely known, the possibility apparently has not been studied with actual LEP data because of the standard expectation of a very heavy top. We estimate [4] that $Z \rightarrow t\bar{c}$ could occur at the 10^{-3} level, but *with* $BR(t \rightarrow b\ell\nu)$ of order a few percent. This results in a signal branching ratio of $Z \rightarrow \ell^+\nu b\bar{c}$ at the few $\times 10^{-5}$ level, and each LEP experiment could have a few tens of events at present. The background level can probably be managed, and LEP experiments are strongly urged to conduct such a search.

II. SINGLET QUARK INDUCED COUPLINGS

Besides the standard u -type quarks u_{iL}^0, u_{iR}^0 ($i = 1 - 3$), we add a left-right singlet charge $2/3$ quark u_{4L}^0, u_{4R}^0 . The left-handed singlet field u_{4L}^0 can pair up with the four right-handed fields to form gauge invariant singlet masses, which we denote as M'_i and M respectively. The right-handed singlet field u_{4R}^0 introduces three extra Yukawa couplings, resulting in off-diagonal masses which we denote as m'_i . Thus, the u -type quark mass matrix is

$$\mathbf{M} = \mathbf{Y} + \mathbf{S} = \begin{bmatrix} m & m' \\ M' & M \end{bmatrix}, \quad (1)$$

where

$$\mathbf{Y} = \begin{bmatrix} m & m' \\ 0 & 0 \end{bmatrix}, \quad \mathbf{S} = \begin{bmatrix} 0 & 0 \\ M' & M \end{bmatrix}, \quad (2)$$

are Yukawa and singlet masses. \mathbf{M} is diagonalised by a biunitary transform,

$$\bar{\mathbf{M}} = U^\dagger \mathbf{M} U' = \text{diag}(\bar{m}_u, \bar{m}_c, \bar{m}_t, \bar{M}_Q), \quad (3)$$

where, departing from the notation of MN [3],

$$U^\dagger = \begin{bmatrix} K & x^\dagger \\ y^\dagger & z^* \end{bmatrix}, \quad (4)$$

and K is a 3×3 matrix. The Yukawa matrix \mathbf{Y} is not simultaneously diagonalised,

$$\bar{\mathbf{Y}} = U^\dagger \mathbf{Y} U' = \bar{\mathbf{M}} - U^\dagger \mathbf{S} U', \quad (5)$$

and the off-diagonal term controls FCNC H^0 and Z^0 couplings. The apparent freedom due to the presence of U' rotation matrix on right-handed fields led MN originally to conclude that tcH^0 coupling could easily be rather large. However, from eq. (3), simple algebra gives

$$-U^\dagger \mathbf{S} U' = - \begin{bmatrix} x^\dagger x \bar{\mathbf{m}} & x^\dagger z M_Q \\ z^* x \bar{\mathbf{m}} & z^* z M_Q \end{bmatrix}, \quad (6)$$

where $\bar{\mathbf{m}}$ is the diagonal 3×3 mass matrix (see eq. (3)). We see that no reference to U' is left, and the off-diagonal couplings depend only on mass eigenvalues and Q -related mixing elements of U [6,7]. The relevant flavor changing Higgs couplings are [6] ($i \neq i'$)

$$\begin{aligned} & - (\bar{m}_i x_i^* x_i \bar{u}_{i'L} u_{iR} + \bar{m}_{i'} x_i^* x_{i'} \bar{u}_{iL} u_{i'R}) \frac{H}{v}, \\ & - (m_i z^* x_i \bar{Q}_L u_{iR} + m_Q x_i^* z \bar{u}_{iL} Q_R) \frac{H}{v}. \end{aligned} \quad (7)$$

The FCNC Z couplings are [3]

$$\begin{aligned} & \frac{g}{2 \cos \theta_W} x_i^* x_i \bar{u}_{i'L} \gamma_\mu u_{iL} Z^\mu + h.c., \\ & \frac{g}{2 \cos \theta_W} x_i^* z \bar{u}_{iL} \gamma_\mu Q_L Z^\mu + h.c., \end{aligned} \quad (8)$$

which is simply related to the Higgs couplings. The charged current becomes

$$\begin{aligned} & \frac{g}{\sqrt{2}} V_{ij} \bar{u}_{iL} \gamma_\mu d_{jL} W^\mu + h.c., \\ & \frac{g}{\sqrt{2}} y'_j \bar{Q}_L \gamma_\mu d_{jL} W^\mu + h.c., \end{aligned} \quad (9)$$

where

$$V \equiv K U^{(d)}, \quad y'_j \equiv y_i^* U_{ij}^{(d)}. \quad (10)$$

The 3×3 KM matrix V is no longer unitary. Both V and y' depend on the 3×3 left-handed down quark rotation matrix $U^{(d)}$.

III. DETAILS

We wish to explore the range of parameter space where tcH coupling could be sizable. To this end we make a special choice of basis to focus on the problem. First, we choose u_R fields such that $M' = 0$ in \mathbf{S} . Second, we choose u_{iL}^0 , $i = 1 - 3$, such that the matrix m is diagonal, hence the KM matrix largely comes from the down-type quark sector (we have checked that it is not possible to generate the observed KM matrix structure just by introducing u -type singlet quarks). Only the charged current is affected by the d -type quark sector, the FCNC Higgs and Z couplings depend only on x_i and z .

The u -type quark mass matrix is now in the form

$$\mathbf{M} = \begin{bmatrix} m_1 & 0 & 0 & \Delta_1 \\ 0 & m_2 & 0 & \Delta_2 \\ 0 & 0 & m_3 & \Delta_3 \\ 0 & 0 & 0 & M \end{bmatrix}. \quad (11)$$

The relevant freedom introduced by the singlet quark Q is parametrized as 3 new off-diagonal Yukawa terms, plus the diagonal, gauge invariant Dirac mass M . The parameters x_i , y_i and z can be found by diagonalizing $\mathbf{M}\mathbf{M}^\dagger$. Without loss of generality, we set $\Delta_1 = 0$ so u quark decouples from our discussion.

To illustrate the correlation between $\hat{m}_i \equiv m_i/M$ and $\hat{\Delta}_i \equiv \Delta_i/M$, we set $\Delta_2 = 0$ and plot, in Fig. 1, the mass eigenvalues m_t/M , M_Q/M vs. \hat{m}_3 for different $\hat{\Delta}_3$ values. Level repulsion is evident: $m_t < m_3$ and $M_Q > M$ for $m_3, \Delta_3 < M$. For larger m_3, Δ_3 values, we adopt the convention that, if $x_t > 0.5$, the heavier state is defined as the top quark. Thus, Fig. 1 depicts both the mass eigenvalues and the label for t and Q .

We are more interested in the effect of Δ_2 . With finite Δ_2 , but negligible m_1, Δ_1, m_2 , the heavy mass eigenvalues are

$$m_t^2, M_Q^2 = \frac{\Sigma^2 \mp \sqrt{\Sigma^4 - 4m_3^2(M^2 + \Delta_2^2)}}{2}, \quad (12)$$

where $\Sigma^2 = M^2 + m_3^2 + \Delta_3^2 + \Delta_2^2$. For sake of discussion, we consider the case where

$m_i, \Delta_i < M$ (top lighter). Note that in Fig. 1 when $\hat{\Delta}_3$ is not too large, the top mass eigenvalue is close to the diagonal term m_3 . This is a generic feature. When other Δ 's can be ignored and $\hat{\Delta}_i$ is not too big, the mass eigenvalue and mixing are roughly

$$\bar{m}_i^2 \sim \frac{m_i^2}{1 + \hat{\Delta}_i^2}, \quad x_i \sim \hat{\Delta}_i. \quad (13)$$

These relations become affected only when there are *two* $\hat{\Delta}_i$ values that are sizable, which follows largely as a consequence of unitarity of the 4×4 matrix U . In Fig. 2 we plot x_t, x_c as a function of $\hat{\Delta}_3$ for $\hat{m}_3 = 0.7$ and $\hat{\Delta}_2 = 0, 0.5, 1, 1.5$. Note the remarkable feature that x_c is almost independent of x_t , but x_t is suppressed by large x_c through unitarity. The physical reason for this can be traced back to the fact that $V_{cb} \sim 0.04$ is very small compared to 1, and that m_c is small.

Thus, the eigenvalue \bar{m}_c could be made small by choosing a small value for m_2 , but this does not forbid Δ_i from being sizable. This is precisely *counter* the ‘‘hierarchy principle’’ [10] advocated in ref. [6]. However, other than being a prejudice, there is really no reason why Δ_2 cannot be large, since it is an independent parameter. Of course, if $\Delta_2 \sim m_2$, then the conclusions of ref. [6] would hold.

IV. PHENOMENOLOGY

Inspection of eq. (7) suggests that $t_L \rightarrow c_R$ transitions are suppressed by m_c/v [6], but $t_R \rightarrow c_L$ transitions have the effective coupling $m_t x_c^* x_t/v$. Since m_t/v is not small, so long that $|x_c x_t|$ is not too suppressed, the $t \rightarrow cH^0$ mode has good probability to be dominant over $t \rightarrow bW^*$ [3]. The necessary condition is therefore that both x_c and x_t are sizable and neither are suppressed. Hence, Q, t and c all become rather arbitrarily mixed although the charm mass is fixed by m_2 . Such an unusual situation is bound to have unusual consequences beyond $t \rightarrow cH^0$ being sizable.

To illustrate the possibility of $t \rightarrow cH^0$ dominance, and at same time account for CDF's apparent observation of a heavy quark of mass 174 GeV, we demand that the heavier quark

(whether dominantly doublet or singlet) mass to be pinned to the CDF value. We then choose $\hat{\Delta}_2, \hat{\Delta}_3 = 0.7, 0.75, M = 110$ GeV, vary m_3 (to get m_t, M_Q etc.) and plot, in Fig. 3, $BR(t \rightarrow cH^0)$ vs. m_t (the physical mass) up to 90 GeV, for $m_H = 50-75$ GeV. We allow for m_H below the present LEP bound in case there are more than one Higgs doublet [12]. In producing Fig. 3, we compute the $t \rightarrow cH^0$ and bW^* decay width using the couplings of eqs. (7) and (9). We assume that $U^{(d)}$ amounts to a “small” rotation close to the “standard” KM matrix, ignoring all phases. We have also ignored $t \rightarrow cZ^*$ decay as this is a three body process subdominant compared to $t \rightarrow bW^*$. It is clear that, if the Higgs boson mass is sufficiently light, $t \rightarrow cH^0$ can be dominant. However, the combined demand of $m_Q \simeq 174$ GeV, and $m_t \lesssim M_W$, dictates that the $t \rightarrow cH^0$ mode cannot be overwhelmingly dominant. Thus, although suppressed, $BR(t \rightarrow bW^*)$ should not be vanishingly small. For larger m_H , $t \rightarrow cH^0$ dominance quickly fades, and the possibility is ruled out by CDF since $t \rightarrow bW^*$ is not drastically suppressed. In the following, we shall assume that one works in the domain where $B_{s,l}$ for the “top” (it could be the dominantly singlet quark, since we do not know the scale for M) is suppressed by 1/3 or more. That is, $B_{s,l} < 1/27$.

Fig. 3 corresponds to $x_c, x_t \simeq 0.59, 0.53$, with $m_t, M_Q = 75, 174$ GeV. The corresponding $Q \rightarrow bW, sW, tH, cH, tZ, cZ$ branching ratios are 0.51, 0.24, 0.04, 0.1, 0.02, 0.09, respectively. Note that, as a consequence of large x_c , $Q \rightarrow sW$ decay has a sizable rate! The modes $Q \rightarrow tH^0$ and tZ are suppressed by phase space, while $Q \rightarrow cH^0$ and cZ are suppressed by an extra power of $|x_c|^2$. Thus, W induced decays are still dominant, but the b content in final state is diluted slightly by the $Q \rightarrow sW$ mode. Although one cannot account for the large production cross section for the heavy quark (*one could always add another singlet u-type quark for this purpose*), other features reported by CDF can be accounted for [8,11], in particular, the appearance of b -tagged $Z + 4$ jet events. The Z boson comes from $Q \rightarrow cZ, tZ$, while a b -tag could come from $Q \rightarrow bW$, or from $t \rightarrow bW^*$ or $H \rightarrow b\bar{b}$, etc., in subsequent decays. This could be at 20% of the $Q\bar{Q}$ cross section, hence consistent to what is observed. On the other hand, single- W with b -tag is slightly depressed ($\sim 70\%$) compared to the standard top. We therefore conclude that the heavy quark observed by CDF may

well be a doublet-singlet mixed state Q . The scenario offers many signatures and can be checked experimentally. The light top with not too suppressed $B_{s,l}$ can perhaps be probed with existing Tevatron data [2].

The scenario has a consequence that may be studied at LEP. As first pointed out by Barbieri and Hall [4], $Z \rightarrow t\bar{c}$ can be quite sizable with existence of charge 2/3 singlet quarks. Using eq. (8) and x_c, x_t values for the example above, we estimate that $BR(Z \rightarrow t\bar{c} + \bar{t}c)$ is of order a few $\times 10^{-3}$, which is consistent with ref. [4]. Other phenomenological constraints are not particularly stringent, and can be found in ref. [4]. For example, $D^0-\bar{D}^0$ mixing constraint can be satisfied with small Δ_1 . Since $BR(t \rightarrow \ell\nu X)$ does not vanish, we estimate that the potentially observable signal of $Z \rightarrow t\bar{c} \rightarrow \ell\nu + 2$ jet (where the jets contain b and c) could have a branching ratio of order a few $\times 10^{-5}$. Since the lepton and neutrino should be well isolated with sizable (15 – 20 GeV) p_T or missing energy (they are *bona fide* virtual W decay events!), and that ℓ, ν and one jet should pair up to be the top mass, there should be sufficient handles for the suppression of background. The latter presumably comes from events with $Z \rightarrow b\bar{b}$ plus gluon bremsstrahlung.

ACKNOWLEDGMENTS

This work is supported in part by the National Science Council of the Republic of China under grant NSC-83-0208-M-002-023. WSH wishes to thank the CERN Theory Division for hospitality and a stimulating environment, and T. Ferguson for discussion.

REFERENCES

- [1] CDF Collaboration: F. Abe *et al.*, FERMILAB-PUB-94/097-E; and Phys. Rev. Lett. **73**, 225 (1994).
- [2] W.S. Hou, Phys. Rev. Lett. **72**, 3945 (1994).
- [3] B. Mukhopadhyaya and S. Nandi, Phys. Rev. Lett. **66**, 285 (1991).
- [4] R. Barbieri and L.J. Hall, Nucl. Phys. **B319**, 1 (1989).
- [5] T.P. Cheng and L.-F. Li, Phys. Rev. **D45**, 1708 (1992).
- [6] W.S. Hou, Phys. Rev. Lett. **69**, 3587 (1992).
- [7] B. Mukhopadhyaya and S. Nandi, Phys. Rev. **D46**, 5098 (1992).
- [8] See, for example, V. Barger and R.J.N. Phillips, preprint MAD/PHYS/830 (1994); and references therein.
- [9] P. Agrawal, S.D. Ellis and W.S. Hou, Phys. Lett. **B256**, 289 (1991).
- [10] F. del Aguila and M.J. Bowick, Nucl. Phys. **B224**, 107 (1983).
- [11] V. Barger *et al.*, preprint MAD/PH/844 and RAL-94-070.
- [12] K. Hikasa *et al.*, the Particle Data Group, Phys. Rev. **D45** (1992) S1.

FIGURES

FIG. 1. m_t/M , M_Q/M vs. \hat{m}_3 for various $\hat{\Delta}_3$ values. The solid(dash) lines stand for physical t(Q) quark.

FIG. 2. Mixing parameter x_t , x_c vs. $\hat{\Delta}_3$ for $\hat{m}_3 = 0.7$ and $\hat{\Delta}_2 = 0$ (dots), 0.5(solid), 1(dash), 1.5(dotdash). Note that x_c is basically independent of x_t .

FIG. 3. $BR(t \rightarrow cH^0)$ vs. m_t for $\hat{\Delta}_2$, $\hat{\Delta}_3 = 0.7, 0.75$, $M = 110$ GeV, and $m_H = 50-75$ GeV in 5 GeV intervals.