



Search for $bb\bar{b}\bar{b}$ tetraquark decays in 4 muons, B^+B^- , $B^0\bar{B}^0$ and $B_S^0\bar{B}_S^0$ channels at LHC



C. Becchi^a, A. Giachino^a, L. Maiani^{b,c,*}, E. Santopinto^a

^a Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Genova, via Dodecaneso 33, 16146 Genova, Italy

^b T. D. Lee Institute, Shanghai Jiao Tong University, Shanghai, 200240, China

^c Dipartimento di Fisica and INFN, Sapienza Università di Roma, Piazzale Aldo Moro 2, I-00185 Roma, Italy

ARTICLE INFO

Article history:

Received 26 February 2020

Received in revised form 28 April 2020

Accepted 15 May 2020

Available online 19 May 2020

Editor: B. Grinstein

ABSTRACT

We perform a quantitative analysis of the $bb\bar{b}\bar{b}$ tetraquark decays into hidden- and open-bottom mesons and calculate, for the first time, the $bb\bar{b}\bar{b}$ tetraquark total decay width. On the basis of our results, we propose the $bb\bar{b}\bar{b} \rightarrow B^+B^-(B^0\bar{B}^0)(B_S^0\bar{B}_S^0) \rightarrow l^+l^- + X$ decays as the most suitable channels to observe the $bb\bar{b}\bar{b}$ tetraquark states, since the calculated two-lepton cross section upper limit, $\simeq 39$ fb, is so large as to be potentially detectable with the 2018 LHCb sensitivity, paving the way to the observation of the $bb\bar{b}\bar{b}$ tetraquark in the future LHCb upgrade. The 4μ signal for the ground state, $J^{PC} = 0^{++}$, is likely to be too small even for the upgraded LHCb, but it may not be hopeless for the $J^{PC} = 2^{++}$ fully-bottom state.

© 2020 ISTITUTO NAZIONALE DI FISICA NUCLEARE. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The hypothetical existence of hadronic states with more than the minimal quark content ($q\bar{q}$ or qqq) was proposed by Gell-Mann in 1964 [1] and Zweig [2], followed by the construction of a quantitative model for two quarks plus two antiquarks by Jaffe [3], to describe the lightest scalar mesons. Recent years have seen considerable growth in the observations of four valence quark states that cannot be included in the well-known systematics of mesons made up of quark-antiquarks, $Z(4248)$, $Z(4430)$, etc. Similar particles have also been found in the bottom sector, $Z_b(10610)$ and $Z_b(10650)$, observed by the Belle collaboration [4] (see [5] or [6] for recent reviews).

The first predictions of a fully-bottom, $bb\bar{b}\bar{b}$ tetraquark below the 2Υ threshold were made in Refs. [7,8], and were supported by more recent contributions [9–13]. Theoretically, $J^{PC} = 0^{++}$ is expected for the $bb\bar{b}\bar{b}$ ground-state.

In 2018, LHCb performed a search for $bb\bar{b}\bar{b}$ decaying into four-muons in the mass range 17.5–20 GeV, but no significant excess was found, leading to the 95% CL upper limit [14]

$$\sigma(pp \rightarrow \mathcal{T}) \times B(\mathcal{T} \rightarrow \Upsilon(1S)\mu^+\mu^-) \times B(\Upsilon(1S) \rightarrow \mu^+\mu^-) < 20 \text{ fb.} \quad (1)$$

Ref. [15] estimates the $J^{PC} = 0^{++}$, fully-bottom tetraquark decay width into $\Upsilon + \mu^+\mu^-$ to be in the range: $10^{-3} - 10$ MeV [15]. Ref. [16] gives a total decay width of 1.2 MeV, a partial decay width into four-leptons in the range: $2.4 \cdot 10^{-3} - 2.4 \cdot 10^{-7}$ MeV with a branching ratio in the range: $2 \cdot 10^{-3} - 2 \cdot 10^{-7}$.

In this letter, assuming the mass of the $J^{PC} = 0^{++}$ fully-bottom tetraquark to lie below the $2\eta_b$ threshold, we present for the first time a calculation of decay widths and branching ratios of the main, hidden- and open-bottom channels. Our results are as follows.

1. Decay rates are all proportional to the parameter:

$$\xi = \frac{|\Psi_{\mathcal{T}}(0)|^2}{|\Psi_{\Upsilon}(0)|^2} \quad (2)$$

the ratio of overlap probabilities of the annihilating $b\bar{b}$ pairs in \mathcal{T} and Υ , respectively. Therefore, branching ratios are uniquely determined and reported in Table 1, which is the basis of our results.

2. In particular, we find

$$B(\mathcal{T} \rightarrow 4\mu) = 7.2 \cdot 10^{-7}. \quad (3)$$

The result (3) is not far from the lower limit of the range in [16], the reason being that the total tetraquark width is in the order of the η_b rather than of the $\Upsilon(1S)$ width, which gives a $\mathcal{O}(10^{-3})$ suppression. With (3), we obtain a realistic estimate of the cross section for $p + p \rightarrow \mathcal{T} \rightarrow \Upsilon + \mu^+\mu^- \rightarrow$

* Corresponding author.

E-mail address: luciano.maiani@roma1.infn.it (L. Maiani).

Table 1
Branching ratios of $J^{PC} = 0^{++}$ and 2^{++} fully-bottom tetraquarks, masses below $2\eta_b$ threshold, assuming S-wave decay.

$[bb][\bar{b}\bar{b}]$	$\eta_b + \text{any}$	$B_q \bar{B}_q$ ($q = u, d, s, c$)	$B_q^* \bar{B}_q^*$	$\Upsilon_b + \text{any}$	$\Upsilon_b + \mu^+ \mu^-$	4μ
$J^{PC} = 0^{++}$	0.65	0.022	0.066	$1.2 \cdot 10^{-3}$	$2.9 \cdot 10^{-5}$	$7.2 \cdot 10^{-7}$
$J^{PC} = 2^{++}$	0	0	0.25	$3.4 \cdot 10^{-3}$	$8.3 \cdot 10^{-5}$	$20 \cdot 10^{-7}$

4μ . Our result is about 350 times lower than the 95% CL upper limit quoted in (1). On the other hand, the calculated cross section for the tetraquark strong decays into two $B_q \bar{B}_q$ mesons, ($q = u, d, s, c$) is large enough, see Table 3, to be potentially detectable in the future LHCb upgrade [17].

3. The total width is expressed as: $\Gamma(\mathcal{T}(J = 0^{++})) = 7.7 \text{ MeV} \cdot \xi$. As a first estimate, we take $|\Psi_\Upsilon(0)|^2$ from the rate $\Upsilon \rightarrow \mu^+ \mu^-$, see also [18], and take the overlap probability in the tetraquark from the model in [13] to obtain:

$$|\Psi_\Upsilon(0)|^2 = 1.2 \text{ GeV}^3, \quad |\Psi_\mathcal{T}(0)|^2 = 0.31 \text{ GeV}^3$$

$$\xi_{th1} \sim 0.26, \quad \Gamma(\mathcal{T}) = 2.0 \text{ MeV} \quad (4)$$

To reduce systematic errors, one could use the overlap probability in Υ computed with a gaussian wave functions, the same method used in [13], to obtain

$$|\Psi_\Upsilon(0)|^2 = 0.16 \text{ GeV}^3$$

$$\xi_{th2} \sim 1.9, \quad \Gamma(\mathcal{T}) = 15 \text{ MeV} \quad (5)$$

The results (4), (5) indicate an uncertainty in the tetraquark width of almost one order of magnitude, which is for experiments to resolve.

We repeated the calculation for the $J^{PC} = 2^{++}$, fully-bottom tetraquark, assuming it lay below the $2\eta_b$ threshold. The $J = 2$ tetraquark is produced in $p + p$ collisions with a statistical factor of 5 with respect to the spin 0 state; the decay $\mathcal{T} \rightarrow \eta_b + \text{light hadrons}$ is suppressed. However, annihilation into two vector mesons $M_q^* \bar{M}_q^*$ takes place at a greater rate than for 0^{++} . Branching fractions of $J^{PC} = 2^{++}$ are listed in Table 1 and, using (4), (5),

$$\Gamma(\mathcal{T}(J = 2^{++})) = 3 - 21 \text{ MeV}. \quad (6)$$

2. Details of the calculation

The starting point is the Fierz transformation, which brings $b\bar{b}$ together [5]:

$$\begin{aligned} \mathcal{T}(J = 0) &= \left| (bb)_{\frac{3}{2}}^1 (\bar{b}\bar{b})_{\frac{3}{2}}^1 \right|_1^0 \\ &= -\frac{1}{2} \left(\sqrt{\frac{1}{3}} \left| (bb)_{\frac{1}{2}}^1 (\bar{b}\bar{b})_{\frac{1}{2}}^1 \right|_1^0 - \sqrt{\frac{2}{3}} \left| (bb)_{\frac{3}{2}}^1 (\bar{b}\bar{b})_{\frac{3}{2}}^1 \right|_1^0 \right) + \\ &+ \frac{\sqrt{3}}{2} \left(\sqrt{\frac{1}{3}} \left| (bb)_{\frac{1}{2}}^0 (\bar{b}\bar{b})_{\frac{1}{2}}^0 \right|_1^0 - \sqrt{\frac{2}{3}} \left| (bb)_{\frac{3}{2}}^0 (\bar{b}\bar{b})_{\frac{3}{2}}^0 \right|_1^0 \right), \quad (7) \end{aligned}$$

quark bilinears are normalised to unity, subscripts denote the dimension of colour representations, and superscripts the total spin. Similarly, for the $J = 2$ tetraquark, one finds:

$$\begin{aligned} \mathcal{T}(J = 2) &= \left| (bb)_{\frac{3}{2}}^1 (\bar{b}\bar{b})_{\frac{3}{2}}^1 \right|_1^2 \\ &= \left(\sqrt{\frac{1}{3}} \left| (bb)_{\frac{1}{2}}^1 (\bar{b}\bar{b})_{\frac{1}{2}}^1 \right|_1^2 - \sqrt{\frac{2}{3}} \left| (bb)_{\frac{3}{2}}^1 (\bar{b}\bar{b})_{\frac{3}{2}}^1 \right|_1^2 \right). \quad (8) \end{aligned}$$




Fig. 1. Feynman diagrams for the annihilation of a $b\bar{b}$ pair into gluons or into lighter quarks, corresponding to the decay mechanisms 1 to 4 considered in the text. J denotes the total angular momentum of the initial $b\bar{b}$ pair.

We describe \mathcal{T} decay as due to individual decays into lower mass states of one of the $b\bar{b}$ pairs in (7), described as follows, see Fig. 1.

1. The colour singlet, spin 0 pair decays into 2 gluons, which are converted into confined, light hadrons (i.e. not containing b flavour) with a rate of the order of α_S^2 ; taking the spectator $b\bar{b}$ pair into account, this decay leads to: $\mathcal{T} \rightarrow \eta_b + \text{light hadrons}$.
2. The colour singlet, spin 1 pair decays into 3 gluons, which are converted into confined light hadrons leading to: $\mathcal{T} \rightarrow \Upsilon + \text{light hadrons}$; final states $\Upsilon + \mu^+ \mu^-$ and 4μ are also produced. The rate is of the order of α_S^3 .¹
3. The colour octet, spin 1 pairs annihilate into one gluon, which materializes into a pair of light quark flavours, $q = u, d, s, c$, the latter recombine with the spectator pair to produce a pair of lower-lying, open-beauty mesons $B_q \bar{B}_q$ and $B_q^* \bar{B}_q^*$, with a rate of the order of α_S^2 .
4. The colour octet, spin 0 pairs annihilate into a pair of lighter quarks (necessary to neutralize the colour of the spectator $b\bar{b}$ pair) with amplitude of the order of α_S^2 and with a rate of the order of α_S^4 , which we neglect.

The total \mathcal{T} decay is the sum of these individual decay rates, which are obtained from the simple formula [21]

$$\Gamma((b\bar{b})_c^s) = |\Psi(0)_\mathcal{T}|^2 v \sigma((b\bar{b})_c^s \rightarrow f) \quad (9)$$

$|\Psi(0)_\mathcal{T}|^2$ is the overlap probability of the annihilating pair, v the relative velocity and σ the spin-averaged annihilation cross section in the final state f .² The spectator $b\bar{b}$ pair, given the lack of extra energy, appears as η_b or Υ on the mass shell, or combines with

¹ With respect to annihilation from the spin 0 state, there is an additional suppression factor: $\Gamma_{S=1}/\Gamma_{S=0} = [5(\pi^2 - 9)/(27\pi)]\alpha_S \sim \alpha_S/20$ due to gauge invariance and the 3-body phase space, see Refs. [20,21]. The extra suppression is evident in the branching fractions reported in Table 1.

² Our method of calculation is borrowed from the theory of K electron capture, where an atomic electron reacts with a proton in the nucleus to give a final nucleus and a neutrino. The rate is computed from formula (9), in which $|\psi(0)|^2$ is the overlap probability of the electron to the proton in the nucleus and σ the electron-proton weak cross section. The other electrons in the atom act as spectators, and rearrange later into a stable atom, after emission of radiation, with unit probability.

the outgoing $q\bar{q}$ pair into an open-beauty meson pair. Our results are valid in the situation where the tetraquark mass is just below the 2Υ threshold and each non-relativistic pair has mass very close to $2m_b$.

We normalise the overlap probabilities to that of $|\Psi_\Upsilon(0)|^2$, which can be derived from the Υ decay rate into lepton pairs. Eq. (9) applied to this case gives:

$$\Gamma(\Upsilon \rightarrow \mu^+ \mu^-) = Q_b^2 \frac{4\pi\alpha^2}{3} \frac{4}{m_\Upsilon^2} |\Psi_\Upsilon(0)|^2, \quad (Q_b = -1/3). \quad (10)$$

It is useful to connect with the Vector Meson Dominance parameter [22] defined by

$$J^\mu(x) = \bar{b}(x)\gamma^\mu b(x) = \frac{m_\Upsilon^2}{f} \Upsilon^\mu(x) \quad (11)$$

f being a pure number. One obtains [19]

$$|\Psi_\Upsilon(0)|^2 = \frac{m_\Upsilon^3}{4f^2}; \quad f = 13.2; \quad |\Psi_\Upsilon(0)|^2 \sim 1.2 \text{ GeV}^3. \quad (12)$$

Numerical results. From Eq. (9), the contribution to the \mathcal{T} decay rate of the colour singlet, spin 0 decay is

$$\begin{aligned} \Gamma_0 &= \Gamma(\mathcal{T} \rightarrow \eta_b + \text{light hadrons}) \\ &= 2 \cdot \frac{1}{4} \cdot |\Psi(0)_\mathcal{T}|^2 v\sigma((b\bar{b})_1^0 \rightarrow 2 \text{ gluons}) \\ &= \frac{1}{2} \Gamma(\eta_b) \cdot \xi = 5 \text{ MeV} \cdot \xi \end{aligned} \quad (13)$$

the factor 2 arises because of the two $(b\bar{b})_1^0$ pairs, 1/4 is the spectroscopic coefficient in (7) and we have approximated³

$$|\Psi(0)_\mathcal{T}|^2 v\sigma((b\bar{b})_1^0 \rightarrow 2 \text{ gluons}) \sim \Gamma(\eta_b) = 10 \text{ MeV}. \quad (14)$$

In a similar way, we obtain

$$\begin{aligned} \Gamma_1 &= \Gamma(\mathcal{T} \rightarrow \Upsilon + \text{light hadrons}) \\ &= 2 \cdot \frac{1}{12} \cdot |\Psi(0)_\mathcal{T}|^2 v\sigma((b\bar{b})_1^1 \rightarrow 3 \text{ gluons}) = \\ &= \frac{1}{6} \Gamma(\Upsilon) \cdot \xi = 9 \text{ keV} \cdot \xi \end{aligned}$$

$$\begin{aligned} \Gamma_2 &= \Gamma(\mathcal{T} \rightarrow \Upsilon + \mu^+ \mu^-) = B_{\mu\mu} \Gamma_1 = 0.22 \text{ keV} \cdot \xi \\ \Gamma_4 &= \Gamma(\mathcal{T} \rightarrow 4\mu) = B_{\mu\mu}^2 \Gamma_1 = 5.5 \cdot 10^{-3} \text{ keV} \cdot \xi \end{aligned} \quad (15)$$

where $B_{\mu\mu} = B(\Upsilon \rightarrow \mu^+ \mu^-)$. Finally, we consider the annihilation of the $(b\bar{b})_8^1$ in light quark pairs. This is illustrated in Fig. 2. Open circles represent the insertion of quark bilinears, and black dots the QCD vertices. Colour matrices and normalizations are explicitly indicated. The numerical factor associated to the traces of the colour matrices along fermion closed paths, C (the Chan-Paton factor [23]) gives the effective coupling constant of the process, $\alpha_{eff} = C\alpha_s$, which is what replaces $Q_b\alpha$ in Eq. (10). From Fig. 2 we read: $C = \sqrt{2}/3$ and we find:

$$\begin{aligned} \Gamma_5 &= \Gamma(\mathcal{T} \rightarrow M(b\bar{q}) + M(q\bar{b})) \\ &= 2 \cdot \frac{1}{6} \cdot \frac{2}{9} \cdot \left(\frac{4\pi\alpha_s^2}{3} \frac{4}{m_\Upsilon^2} \right) |\Psi(0)_\mathcal{T}|^2 \cdot \xi \end{aligned} \quad (16)$$

The factor 2 arises from the two choices of the annihilating bilinear,⁴ we have inserted the spectroscopic factor of the spin 1




Fig. 2. Colour flow in $b\bar{b}$ annihilation.

colour octet from (7), the Chan Paton factor and used $m_\mathcal{T} \sim 2m_\Upsilon$. In parentheses $v\sigma(b\bar{b} \rightarrow q\bar{q})$.

Using Eq. (12), $\alpha_s = 0.2$ and massless q , we obtain

$$\Gamma_5 = \frac{8\pi}{81} \left(\frac{\alpha_s}{f} \right)^2 m_\Upsilon \cdot \xi = 0.67 \text{ MeV} \cdot \xi \quad (17)$$

and

$$\Gamma(\mathcal{T}) = \Gamma_0 + \Gamma_1 + 4\Gamma_5 = 7.7 \text{ MeV} \cdot \xi \quad (18)$$

A non-vanishing mass of the final quark brings a negligible correction even for the charm. Eq. (17) refers to the total decay rate into pseudoscalar and vector meson pairs. We can separate the two rates according to the following argument. In non-relativistic notation, the spin-colour structure of the final state after annihilation corresponds to the operator (\mathcal{O} is normalized to unit norm, see [5]):

$$\mathcal{O}_{fin} = \frac{1}{4\sqrt{2}} \sum_A (b_c \lambda^A \sigma_2 \sigma b) \cdot (q_c \lambda^A \sigma_2 \sigma q) \quad (19)$$

Using the appropriate Fierz-rearranging relations, one sees that⁵ $B_q B_q^*$ and $\mathbf{B}_q \mathbf{B}_q^*$ pairs are produced in the spin combination, $\frac{1}{2} [(B_q B_q^*) + (\mathbf{B}_q \cdot \mathbf{B}_q^*)]$ and the rate in $\bar{b}q + q\bar{b}$ is shared between pseudoscalar and vector mesons in the ratio 1 : 3.

3. The value of $|\Psi_\mathcal{T}(0)|^2$

A value for $|\Psi_\mathcal{T}(0)|^2$ can be obtained from the calculation in [13,26]. Constituent coordinates are defined as

\mathbf{x}, \mathbf{y} : antiquarks; $\mathbf{z}, \mathbf{0}$: quarks

One defines the Jacobi coordinates

$$\xi_1 = \mathbf{x} - \mathbf{y}; \quad \xi_2 = \mathbf{z}; \quad \xi_3 = \mathbf{x} + \mathbf{y} - (\mathbf{z} + \mathbf{0}) \quad (20)$$

The \mathcal{T} wave function is a product of normalized gaussians with parameters $\beta_1 = \beta_2 = 0.77 \text{ GeV}$, $\beta_3 = 0.60 \text{ GeV}$, obtained by minimising the expectation of the Hamiltonian of Ref. [13,26]. The equality $\beta_1 = \beta_2$ is due to Charge conjugation invariance. By elementary integrations, one can obtain the wave function squared in the variable

$$\ell = \frac{1}{2} \xi_3, \quad (21)$$

i.e. the separation of the centers of gravity of quarks and antiquarks.

One finds

$$|\Psi_\mathcal{T}(\ell)|^2 = \left(\frac{2\beta_3}{\pi} \right)^{3/2} \cdot e^{-4\beta_3^2 \ell^2} \quad (22)$$

³ We understand that the same $|\Psi(0)_\mathcal{T}|^2$ applies to η_b and Υ .

⁴ given the symmetry of the tetraquark, we may call b_1 the annihilating b quark and pair it, in Eq. (7), to either \bar{b}_1 or \bar{b}_2 .

⁵ Fierzing colours produces $b\bar{q}$ and $q\bar{b}$ bilinears in colour singlets and colour octet; one may argue that gluons from the vacuum will screen colour octet charges [24,25].

Table 2
Estimates of overlap probability and radius, for Υ and \mathcal{T} .

Particle	Method	$ \Psi(0) ^2$ (GeV ³)	$\sqrt{\langle R^2 \rangle}$ (fm)
Υ	Eq. (12)	1.2	0.13, using Eq. (22)
Υ	Eq. (24)	0.16	0.25
Υ	Ref. [26]	0.24 (gaussian w.f.)	0.22
\mathcal{T}	$\langle \ell^2 \rangle$, Eq. (22)	0.31	0.20

For gaussian wave function there is a fixed relation $|\Psi(0)|^2 = (\frac{3}{2\pi R^2})^{3/2}$.

To compute $|\Psi_{\Upsilon}(0)|^2$, we follow Ref. [13]. We obtain the wave function

$$|\Psi_{\Upsilon}(\mathbf{x})|^2 = \left(\frac{\beta_{\Upsilon}}{\pi}\right)^{3/2} \cdot e^{-\beta_{\Upsilon}^2 \mathbf{x}^2}, \quad \beta_{\Upsilon} = 0.96 \text{ GeV} \quad (23)$$

$$|\Psi_{\Upsilon}(0)|^2 = 0.16 \text{ GeV}^3; \quad R_{\Upsilon} = \sqrt{\frac{3}{2\beta_{\Upsilon}}} = 0.25 \text{ fm}. \quad (24)$$

Table 2 reports the results obtained from different methods. As the table shows, the overlap probability from Υ leptonic decay is substantially larger than the one obtained in the gaussian model, which, for the radius, agrees reasonably with the independent evaluation of Ref. [26]. The discrepancy underlines the need to estimate $\xi = |\Psi_{\mathcal{T}}(0)|^2/|\Psi_{\Upsilon}(0)|^2$ by means of the same method for the numerator and denominator.

With the definition of the radius in Eqs. (22), for \mathcal{T} , and with Eq. (24) for Υ , we find $\sqrt{\langle \ell^2 \rangle} = 0.20 \text{ fm}$, $|\Psi(0)_{\mathcal{T}}|^2 = 0.31 \text{ GeV}^2$ leading to:

$$\xi_{\text{th1}} = 0.26 \quad (|\Psi_{\Upsilon}(0)|^2 = 1.2 \text{ GeV}^3)$$

$$\xi_{\text{th2}} = 1.9 \quad (|\Psi_{\Upsilon}(0)|^2 = 0.16 \text{ GeV}^3)$$

Branching ratios do not depend upon ξ and are not affected by this uncertainty.

4. Tetraquark cross section in the 4μ and $B_f^{(*)} B_f^{(*)}$ channels

By combining Eqs. (15) and (18) we obtain a very low branching fraction for $\mathcal{T} \rightarrow 4\mu$:

$$B_{4\mu} = B(\mathcal{T} \rightarrow 4\mu) = 7.2 \cdot 10^{-7} \quad (25)$$

From (25) and the assumption that the cross section for the production of 2Υ provides an upper bound to the tetraquark production cross section, one derives the upper limit:

$$\sigma_{\text{theo.}}(\mathcal{T} \rightarrow 4\mu) \leq \sigma(pp \rightarrow 2\Upsilon) B_{4\mu} = \begin{cases} 0.049 \text{ fb, with } \sigma(pp \rightarrow 2\Upsilon) \simeq 69 \text{ pb [27]} \\ 0.056 \text{ fb, with } \sigma(pp \rightarrow 2\Upsilon) \simeq 79 \text{ pb [28]} \end{cases} \quad (26)$$

$\sigma(pp \rightarrow 2\Upsilon) \simeq 69(79) \text{ pb}$ are the 2Υ production cross sections measured by CMS at 8 TeV [27] and 13 TeV [28]. According to (26), the four-muon tetraquark cross section is far below the current LHCb capabilities, the upper limit in Eq. (26) being more than 350 times lower than the CL of the 95% LHCb upper limit of $\simeq 20 \text{ fb}$ quoted in Eq. (1).

Summing over the four light flavours, we see from Table 1 that decays into meson pairs account for about 35% of \mathcal{T} decays. Decay into $B^+ B^-$ mesons may provide a promising channel to discover the $4b$ tetraquark. In Table 1 we report the tetraquark open-bottom branching fractions and the upper limits to the tetraquark two-lepton cross section, $\sigma_{\text{theo.}}(\mathcal{T} \rightarrow 2B_q \rightarrow 2l)$, calculated as

$$\begin{aligned} \sigma_{\text{theo.}}(\mathcal{T} \rightarrow 2B_f \rightarrow 2l) &= \sigma(pp \rightarrow \mathcal{T}) Br(\mathcal{T} \rightarrow 2B_q) [Br(B_q \rightarrow l + \bar{\nu} + X)]^2 \\ &\leq \sigma(pp \rightarrow 2\Upsilon) Br(\mathcal{T} \rightarrow 2B_q) [Br(B_q \rightarrow l + \bar{\nu} + X)]^2 \end{aligned} \quad (27)$$

Table 3

Upper limits of two- and four-lepton cross sections via \mathcal{T} production calculated using as inputs the two- Υ production cross sections measured by CMS at 8 TeV [27] and 13 TeV [28].

Decay channel	Predicted BF	Predicted two-lepton cross section upper limit (fb)	
		8 TeV	13 TeV
$\mathcal{T}(J=0) \rightarrow B^+ B^- (B^0 \bar{B}^0, B_s^0 \bar{B}_s^0) \rightarrow 2\ell + \dots$	0.022	34	39
$\mathcal{T}(J=0) \rightarrow 4\mu$	$7.2 \cdot 10^{-7}$	0.049	0.056
$\mathcal{T}(J=2) \rightarrow 4\mu$	$20 \cdot 10^{-7}$	0.14	0.16

Decays such as $B_f^* \rightarrow l + \bar{\nu} + X$ occur by means of intermediate electromagnetic decays, for example $B_f^* \rightarrow B_f + \gamma \rightarrow l + \bar{\nu} + X + \gamma$. For this reason, the excited open-bottom channels are not suited to the search for tetraquark states. Upper limits for \mathcal{T} production and decay into $B^+ B^-$ or $B^0 \bar{B}^0$ or $B_s^0 \bar{B}_s^0$ are reported in Table 3.

In conclusion, we propose the $B^+ B^-$, $B^0 \bar{B}^0$ and $B_s^0 \bar{B}_s^0$ channels in the search for full-bottom tetraquarks in future LHCb upgrades. The 4μ signal produced by the $J^{PC} = 2^{++}$ tetraquark may not be hopeless.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] M. Gell-Mann, Phys. Lett. 8 (1964) 214.
- [2] G. Zweig, An SU3 model for strong interaction symmetry and its breaking, CERNTH-401, 1964.
- [3] R.L. Jaffe, Phys. Rev. D 15 (1977) 267, <https://doi.org/10.1103/PhysRevD.15.267>.
- [4] A. Bondar, et al., Belle Collaboration, Phys. Rev. Lett. 108 (2012) 122001.
- [5] A. Ali, L. Maiani, A.D. Polosa, <https://doi.org/10.1017/9781316761465>.
- [6] A. Esposito, A. Pilloni, A.D. Polosa, Phys. Rep. 668 (2017) 1.
- [7] L. Heller, J.A. Tjon, Phys. Rev. D 32 (1985) 755.
- [8] A.V. Berezhnoy, A.V. Luchinsky, A.A. Novoselov, Phys. Rev. D 86 (2012) 034004.
- [9] J. Wu, Y.R. Liu, K. Chen, X. Liu, S.L. Zhu, Phys. Rev. D 97 (9) (2018) 094015.
- [10] W. Chen, H.X. Chen, X. Liu, T.G. Steele, S.L. Zhu, Phys. Lett. B 773 (2017) 247.
- [11] Y. Bai, S. Lu, J. Osborne, arXiv:1612.00012 [hep-ph].
- [12] Z.G. Wang, Eur. Phys. J. C 77 (7) (2017) 432, <https://doi.org/10.1140/epjc/s10052-017-4997-0>, arXiv:1701.04285 [hep-ph].
- [13] M.N. Anwar, J. Ferretti, F.K. Guo, E. Santopinto, B.S. Zou, Eur. Phys. J. C 78 (8) (2018) 647.
- [14] R. Aaij, et al., LHCb Collaboration, J. High Energy Phys. 1810 (2018) 086.
- [15] A. Esposito, A.D. Polosa, Eur. Phys. J. C 78 (9) (2018) 782.
- [16] M. Karliner, S. Nussinov, J.L. Rosner, Phys. Rev. D 95 (3) (2017) 034011.
- [17] R. Aaij, et al., LHCb Collaboration, arXiv:1808.08865.
- [18] E.J. Eichten, C. Quigg, Phys. Rev. D 52 (1995) 1726.
- [19] M. Tanabashi, et al., Particle Data Group, Phys. Rev. D 98 (2018) 030001 and 2019 update.
- [20] T. Appelquist, H.D. Politzer, Phys. Rev. Lett. 34 (1975) 43.
- [21] V.B. Berestetskii, E.M. Lifshitz, L.P. Pitaevskii, Quantum Electrodynamics, Pergamon Press, 1982, p. 371; a more recent application to particle physics is found in R. van Royen, V.F. Weisskopf, Nuovo Cimento A 50 (1967) 617.
- [22] See e.g. D. Schildknecht, Acta Phys. Pol. B 37 (2006) 595.
- [23] J.E. Paton, H.M. Chan, Nucl. Phys. B 10 (1969) 516.
- [24] G.S. Bali, Phys. Rep. 343 (2001) 1.
- [25] L. Maiani, A.D. Polosa, V. Riquer, Phys. Rev. D 100 (7) (2019) 074002.
- [26] B. Liu, P.N. Shen, H.C. Chiang, Phys. Rev. C 55 (1997) 3021.
- [27] V. Khachatryan, A.M. Sirunyan, et al., The CMS Collaboration, J. High Energy Phys. 2017 (2017) 13, [https://doi.org/10.1007/JHEP05\(2017\)013](https://doi.org/10.1007/JHEP05(2017)013).
- [28] A.M. Sirunyan, et al., CMS Collaboration, arXiv:2002.06393 [hep-ex].