



Search for dark matter produced in association with a single top quark in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector

The ATLAS Collaboration

This paper presents a search for dark matter in the context of a two-Higgs-doublet model together with an additional pseudoscalar mediator, a , which decays into the dark-matter particles. Processes where the pseudoscalar mediator is produced in association with a single top quark in the 2HDM+ a model are explored for the first time at the LHC. Several final states which include either one or two charged leptons (electrons or muons) and a significant amount of missing transverse momentum are considered. The analysis is based on proton–proton collision data collected with the ATLAS experiment at $\sqrt{s} = 13$ TeV during LHC Run 2 (2015–2018), corresponding to an integrated luminosity of 139 fb^{-1} . No significant excess above the Standard Model predictions is found. The results are expressed as 95% confidence-level limits on the parameters of the signal models considered.

1 Introduction

Strong evidence for the existence of a new, non-luminous matter component of the universe, dark matter (DM), arises from astrophysical observations such as precise measurements of the cosmic microwave background and from gravitational lensing measurements. Through its gravitational interactions, it is suggested that DM constitutes up to 26% of the matter–energy content of the universe [1, 2]. The nature and properties of DM remain largely unknown in the context of the Standard Model (SM) of particle physics. Under the hypothesis that DM behaves like a weakly interacting massive particle (WIMP) [3], searches are performed using multiple, complementary approaches. At hadron colliders, searches for WIMP-like DM production crucially rely on one or more visible particles being produced in association with the sought-after invisible DM candidate. The experimental signature for DM candidates is missing transverse momentum (\vec{p}_T^{miss} , its modulus denoted by E_T^{miss}) in collision events. Several models have been proposed in the past decades, with the details of the DM–SM production process depending on the model assumptions.

A class of simplified models for DM searches at the LHC is considered in this paper. It involves a two-Higgs-doublet extended sector together with an additional pseudoscalar mediator to DM, the 2HDM+ a model [4, 5]. This class of models represents one of the simplest ultraviolet-complete and renormalisable frameworks for investigating the broad phenomenology predicted by spin-0 mediator-based DM models [5–19]. For the present study, a type-II [20, 21] coupling structure of the Higgs sector to third-generation fermions is considered. The CP eigenstates are identified with the mass eigenstates, i.e. two scalars h and H , two pseudoscalars A and a , and charged scalars H^\pm . Three mixing angles are defined in the model: α denotes the mixing angle between the two CP-even weak spin-0 eigenstates, $\tan\beta$ is the ratio of the vacuum expectation values (VEVs) of the two Higgs doublets and θ represents the mixing angle of the two CP-odd weak spin-0 eigenstates. The alignment ($\cos(\beta - \alpha) = 0$) and decoupling limit is assumed, such that the lightest CP-even state of the Higgs sector, h , can be identified with the SM Higgs boson and the electroweak VEV is set to 246 GeV. The pseudoscalar mediator a couples the DM particles, χ , to the SM and mixes with the pseudoscalar partner of the SM Higgs boson, A . Following the prescriptions in Ref. [5], the masses of the heavy CP-even Higgs boson H and charged bosons H^\pm are set equal to the mass of the heavy CP-odd partner A .

This set of models offers a rich phenomenology, with a variety of final states that might arise depending on the production and decay modes of the various bosons composing the Higgs sector, as investigated in Ref. [22]. A recent study [23] has shown that final-state events characterised by the presence of E_T^{miss} and a single top quark provide promising sensitivity to 2HDM+ a models. As in SM single top production, three different types of processes contribute at leading order (LO) in QCD: t -channel production, s -channel production and associated production with a W boson (tW). In the following, these are collectively referred to as DM t processes. The t -channel process $pp \rightarrow tj\chi\bar{\chi}$ receives its dominant contributions from the two diagrams shown in Figures 1(a) and 1(b). These two diagrams interfere destructively, ensuring the perturbative unitarity of the $pp \rightarrow tj\chi\bar{\chi}$ process. The magnitude of the interference decreases with increasing H^\pm mass. In the case of the tW production channel, the two diagrams shown in Figures 1(c) and 1(d) provide the dominant contributions to the DM t cross section. As in t -channel production, these two diagrams interfere destructively. When the decays $H^\pm \rightarrow W^\pm a$ are kinematically possible, the charged Higgs bosons are produced on-shell and the cross section of $pp \rightarrow tW\chi\bar{\chi}$, assuming H^\pm masses of a few hundred GeV, increases to produce a sizeable event rate. Finally, s -channel production is relevant in regions of the parameter space characterised by low H^\pm masses (< 300 GeV) and it is not directly targeted by the analysis.

This paper presents a dedicated search for single top quarks produced in association with DM candidates, exploiting final-state signatures characterised by the presence of: large E_T^{miss} ; jets, possibly arising from the fragmentation of b -hadrons (b -jets); and one or two charged leptons, either electrons or muons ($\ell = e, \mu$). The analysis is conducted using proton–proton (pp) collisions at a centre-of-mass energy $\sqrt{s} = 13$ TeV produced at the LHC and collected by ATLAS between 2015 and 2018, for a dataset corresponding to 139 fb^{-1} . Three analysis channels, characterised by different lepton or jet multiplicities, are optimised to target different processes: tW_{1L} and tW_{2L} (single-lepton and dilepton final states, respectively) for the tW +DM events and tj_{1L} for t -channel DM production. The results are interpreted in the context of 2HDM+ a models, considering various assumptions about the most relevant parameters, m_a , m_{H^\pm} , and $\tan\beta$. Furthermore, the mutually exclusive tW_{1L} and tW_{2L} analysis channels are statistically combined to maximise the sensitivity to tW +DM processes.

Previous searches for 2HDM+ a models targeted associated production of DM candidates with Higgs or Z bosons, as well as DM and a $t\bar{t}$ pair (referred to as $DMt\bar{t}$) (see Ref. [24] for CMS and Ref. [22] and references therein for ATLAS). This search is targeting the unexplored models within ATLAS where DM produced in association with single top quarks (for CMS results, see Ref. [25]). The analysis is also sensitive to $DMt\bar{t}$ processes in regions of the parameter space where the DMt and $DMt\bar{t}$ production rates are similar.




Figure 1: Representative diagrams of the dark-matter particle χ pair production from the 2HDM+ a model considered in this analysis: (a)–(b) through the t -channel, and (c)–(d) through the tW channel.

2 ATLAS detector

The ATLAS detector [26] is a multipurpose particle detector with a forward–backward symmetric cylindrical geometry and nearly 4π coverage in solid angle.¹ The inner tracking detector consists of pixel and microstrip silicon detectors covering the pseudorapidity region $|\eta| < 2.5$, surrounded by a transition radiation tracker which enhances electron identification in the region $|\eta| < 2.0$. A new inner pixel layer, the insertable B-layer [27, 28], was added at a mean radius of 3.3 cm during the period between Run 1 and Run 2 of the LHC. The inner detector is surrounded by a thin superconducting solenoid providing an axial 2 T magnetic field and by a fine-granularity lead/liquid-argon (LAr) electromagnetic calorimeter covering

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point in the centre of the detector. The positive x -axis is defined by the direction from the interaction point to the centre of the LHC ring, with the positive y -axis pointing upwards, while the beam direction defines the z -axis. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity η is defined in terms of the polar angle θ by $\eta = -\ln \tan(\theta/2)$. Rapidity is defined as $y = 0.5 \ln[(E + p_z)/(E - p_z)]$ where E denotes the energy and p_z is the component of the momentum along the beam direction. The angular distance ΔR is defined as $\sqrt{(\Delta y)^2 + (\Delta\phi)^2}$.

$|\eta| < 3.2$. A steel/scintillator-tile calorimeter provides hadronic coverage in the central pseudorapidity range ($|\eta| < 1.7$). The endcap and forward regions ($1.5 < |\eta| < 4.9$) of the hadron calorimeter are made of LAr active layers with either copper or tungsten as the absorber material. A muon spectrometer with an air-core toroid magnet system surrounds the calorimeters. Three layers of high-precision tracking chambers provide coverage in the range $|\eta| < 2.7$, while dedicated fast chambers allow triggering in the region $|\eta| < 2.4$. The ATLAS trigger system consists of a hardware-based level-1 trigger followed by a software-based high-level trigger [29].

3 Data and Monte Carlo simulation

The data analysed in this paper correspond to an integrated luminosity of 139 fb^{-1} of pp collision data collected between 2015 and 2018 by the ATLAS detector with a centre-of-mass energy of 13 TeV and a 25 ns proton bunch crossing interval. The uncertainty in the combined 2015–2018 integrated luminosity is 1.7% [30], obtained using the LUCID-2 detector [31] for the primary luminosity measurements. All detector subsystems were required to be operational during data taking. The average number of interactions in the same and nearby bunch crossings (pile-up) increased from $\langle\mu\rangle = 13.4$ (2015 dataset) to $\langle\mu\rangle = 36.1$ (2018 dataset), with a highest $\langle\mu\rangle = 37.8$ (2017 dataset) and an average $\langle\mu\rangle = 33.7$.

Candidate events were recorded using a combined set of triggers [29] based on the presence of missing transverse momentum or charged leptons ($\ell = e, \mu$). The $E_{\text{T}}^{\text{miss}}$ trigger [32] is fully efficient for events with reconstructed $E_{\text{T}}^{\text{miss}} > 250 \text{ GeV}$ and it was used for the single-lepton analysis channels. Furthermore, an OR between $E_{\text{T}}^{\text{miss}}$ and single-lepton triggers was used for the tj_{IL} channel for events with reconstructed $E_{\text{T}}^{\text{miss}} < 250 \text{ GeV}$. Triggers based on a single muon (electron) require the presence of a muon (electron) with transverse momentum p_{T} (transverse energy E_{T}) above certain thresholds, and impose data quality and lepton isolation requirements. The lowest p_{T} (E_{T}) threshold without trigger prescaling is 24 (26) GeV for muons (electrons) and includes a lepton isolation requirement that is not applied for triggers with higher thresholds. In the two-lepton channel, lower thresholds for electrons and muons must be applied to retain sensitivity to the signal. A combined set of two-lepton triggers was used, with the muon (electron) p_{T} (E_{T}) trigger threshold depending on the data-taking period. The lepton trigger threshold ranged between 8 and 22 GeV for muons, and between 12 and 24 GeV for electrons. The analysis selections are chosen to guarantee maximum trigger efficiency, generally above 95%. Trigger matching requirements [29] are applied where the lepton(s) must lie in the vicinity of the corresponding trigger-level object.

Dedicated Monte Carlo (MC) simulated samples are used to model SM processes and estimate the expected signal yields. All samples were produced using the ATLAS simulation infrastructure [33] and GEANT4 [34], or a faster simulation based on a parameterisation of the calorimeter response and GEANT4 for the other detector systems [33]. The simulated events are reconstructed with the same algorithms as used for data. They contain a realistic modelling of pile-up interactions with pile-up profiles matching the ones of each dataset between 2015 and 2018, obtained by overlaying minimum-bias events simulated using the soft QCD processes of PYTHIA 8.186 [35] with the NNPDF2.3 LO set of parton distribution functions (PDFs) [36] and the A3 [37] set of tuned parameters (tune).

Signal MC samples for single top quark production in association with DM include tW , t -channel and s -channel processes. Samples were produced either varying the (m_a, m_{H^\pm}) parameters and assuming $\tan\beta$ equal to unity, or varying the $(\tan\beta, m_{H^\pm})$ parameters and setting $m_a = 250 \text{ GeV}$. Details of other parameter value assumptions are provided in Section 7. The samples were generated from leading-order (LO) matrix elements using the MADGRAPH5_aMC@NLO [38] v2.6.2 generator interfaced to PYTHIA 8.212 [39] with

Table 1: List of generators used for the different SM background processes. Diboson includes WW , WZ and ZZ production. Information is given about the underlying-event tunes, the PDF sets and the perturbative QCD highest-order accuracy (LO, NLO, next-to-next-to-leading order (NNLO), and next-to-next-to-leading-log (NNLL)) used for the normalisation of the different samples. Diboson cross sections are taken directly from SHERPA.

Process	Generator	PDF set	PS and frag./hadr.	UE tune	Cross-section accuracy
Top pair ($t\bar{t}$)	POWHEG-Box v2 [49–52]	NNPDF 3.0 NLO	PYTHIA 8	A14	NNLO+NNLL [53]
Single-top $\left\{ \begin{array}{l} t\text{-channel} \\ s\text{- and } Wt\text{-channel} \end{array} \right.$	POWHEG-Box v1 [54]	NNPDF 3.0 NLO	PYTHIA 8	A14	NNLO+NNLL [55]
	POWHEG-Box v2 [56]	NNPDF 3.0 NLO	PYTHIA 8	A14	NNLO+NNLL [57, 58]
V +jets ($V = W/Z$)	SHERPA 2.2.1 [43–47]	NNPDF 3.0 NNLO	SHERPA	Default	NNLO [59]
Diboson	SHERPA 2.2.1 or 2.2.2	NNPDF 3.0 NNLO	SHERPA	Default	NLO
$t\bar{t} + V$, $V = W, Z, h$	MADGRAPH5_aMC@NLO 2.3.3	NNPDF 3.0 NLO	PYTHIA 8	A14	NLO [38, 60]
tWZ	MADGRAPH5_aMC@NLO 2.6.7	NNPDF 3.0 NLO	PYTHIA 8	A14	NLO [38]

the A14 tune [40] for the modelling of parton showering (PS), hadronisation and the description of the underlying event. Parton luminosities are provided by the five-flavour scheme NNPDF3.0 NLO [41] PDF set. Signal cross sections are calculated to LO accuracy in QCD. Additional simulated samples are used for $DMt\bar{t}$ processes. They were generated using LO matrix elements, with up to one extra parton using the MADGRAPH5_aMC@NLO v2.6.7 generator interfaced to PYTHIA 8.244 with the same PDF set and tune as used for the tW , t - and s -channel processes. The top quark decay was simulated using MADSPIN [42]. In this case, signal cross sections are calculated to next-to-leading-order (NLO) accuracy using the same version of MADGRAPH5_aMC@NLO as suggested in Ref. [16].

Background samples were simulated using different MC event generators depending on the process. All background processes are normalised to the best available theoretical calculation of their respective cross sections. The event generators, the accuracy of theoretical cross sections, the underlying-event parameter tunes, and the PDF sets used in simulating the SM background processes most relevant for this analysis are summarised in Table 1. For all samples, except those generated using SHERPA [43–47], the EVTGEN v1.2.0 [48] program was used to simulate the properties of the b - and c -hadron decays.

4 Event reconstruction and object definitions

Common event-quality criteria and object reconstruction definitions are applied for all analysis channels, including standard data-quality requirements to select events taken during optimal detector operation. In addition, in each analysis channel, dedicated selection criteria, which are specific to the objects and kinematics of interest in those final states, are applied as described in Section 5.

Events are required to have at least one reconstructed interaction vertex with a minimum of two associated tracks each having $p_T > 500$ MeV. In events with multiple vertices, the one with the highest sum of squared transverse momenta of associated tracks is chosen as the primary vertex [61]. A set of baseline quality criteria are applied to reject events with non-collision backgrounds or detector noise [62].

Two levels of object identification requirements are defined for leptons and jets: baseline and signal. Baseline leptons and jets are selected with looser identification criteria, and are used in computing the missing transverse momentum as well as in resolving possible reconstruction ambiguities. Signal leptons and jets are a subset of the baseline objects, with tighter quality requirements which are used to define the

search regions. Isolation criteria, defined with a list of tracking-based and calorimeter-based variables, are used to select signal leptons by discriminating between semileptonic heavy-flavour decays and jets misidentified as leptons.

Electron candidates are reconstructed from energy deposits in the electromagnetic calorimeter that are matched to charged-particle tracks in the inner detector (ID) [63]. Baseline electrons are required to satisfy $p_T > 10$ GeV and $|\eta| < 2.47$, excluding the transition region between the barrel and endcap calorimeters ($1.37 < |\eta| < 1.52$). They are identified using the ‘loose’ likelihood identification operating point as described in Ref. [63]. The number of hits in the innermost pixel layer is used to discriminate between electrons and converted photons. The longitudinal impact parameter z_0 relative to the primary vertex is required to satisfy $|z_0 \sin \theta| < 0.5$ mm. Signal electrons are required to also satisfy $p_T > 20$ GeV and the ‘tight’ likelihood identification criteria as defined in Ref. [63]. The significance of the transverse impact parameter d_0 must satisfy $|d_0/\sigma(d_0)| < 5$ for signal electrons. Signal electrons with $p_T < 200$ GeV are further refined using the ‘FCLoose’ isolation working point, while those with larger p_T are required to pass the ‘FCHighPtCaloOnly’ isolation working point, as described in Ref. [63]. Corrections for energy contributions due to pile-up are applied.

Muon candidates are reconstructed from matching tracks in the ID and muon spectrometer, refined through a global fit which uses the hits from both subdetectors [64]. Baseline muons must have $p_T > 10$ GeV and $|\eta| < 2.5$, and satisfy the ‘medium’ identification criteria. Like the electrons, their longitudinal impact parameter z_0 relative to the primary vertex is required to satisfy $|z_0 \sin \theta| < 0.5$ mm. Signal muons are defined with tighter requirements on their transverse momentum and transverse impact parameter significance: $p_T > 20$ GeV and $|d_0/\sigma(d_0)| < 3$. The ‘FCLoose’ isolation working point is also required for signal muons [64].

Jets are reconstructed from topological clusters of energy depositions in the calorimeters using the anti- k_t algorithm [65], with a radius parameter $R = 0.4$ [66]. The average energy contribution from pile-up is subtracted according to the jet area and the jets are calibrated as described in Ref. [67]. To further reduce the effect of pile-up interactions, the jets with $|\eta| < 2.4$ and $p_T < 120$ GeV are required to satisfy the ‘medium’ working point of the jet vertex tagger (JVT), a tagging algorithm that identifies jets originating from the primary vertex using track information [68, 69]. Baseline jets are selected in the region $|\eta| < 4.5$ and have $p_T > 20$ GeV. The selection of signal jets requires them to be in the region $|\eta| < 2.5$ and to have $p_T > 30$ GeV.

Jets containing b -hadrons are identified as arising from b -quarks (‘ b -tagged’) using a multivariate algorithm (MV2c10), based on the track impact parameters, the presence of displaced secondary vertices and the reconstructed flight path of b - and c -hadrons inside the jet [70]. These b -tagged jets are reconstructed in the region $|\eta| < 2.5$ and have $p_T > 20$ GeV. The b -tagging working point provides an efficiency of 77% for jets containing b -hadrons in simulated $t\bar{t}$ events, with average rejection of 110 and 4.9 for light-flavour jets and jets containing c -hadrons, respectively [71].

To resolve the reconstruction ambiguities between electrons, muons and jets, an overlap removal procedure is applied to baseline objects in a prioritised sequence as follows. First, if an electron shares the same ID track with another electron, the one with lower p_T is discarded. Any electron sharing the same ID track with a muon is rejected. Next, jets that are not b -tagged are rejected if they lie within $\Delta R = 0.2$ of an electron. Similarly, jets that are not b -tagged are rejected if they lie within $\Delta R = 0.2$ of a muon if the jet has fewer than three associated tracks or the muon is matched to the jet through ghost association [72]. Finally, electrons and muons that are close to a remaining jet are discarded if their distance from the jet is $\Delta R < \min(0.4, 0.04 + 10 \text{ GeV}/p_T)$ as a function of the lepton p_T .

The missing transverse momentum \vec{p}_T^{miss} , with magnitude E_T^{miss} , is calculated as the negative vectorial sum of the transverse momentum of all baseline reconstructed objects (electrons, muons, jets and photons [73]) and the soft term. The soft term includes all tracks associated with the primary vertex but not matched to any reconstructed physics object. Tracks not associated with the primary vertex are not considered in the E_T^{miss} calculation, improving the E_T^{miss} resolution by suppressing the effect of pile-up [74, 75].

To compensate for differences between data and simulation in trigger, particle identification and reconstruction efficiency, correction factors that are usually functions of the relevant kinematic variables are derived from data and applied to the samples of simulated events.

5 Analysis strategy

The search is conducted in three independent analysis channels differing in lepton and jet multiplicities to maximise the sensitivity to distinct signal processes. The tW_{1L} analysis channel targets tW +DM events where one of the W bosons (directly produced or arising from the top quark decay) decays leptonically (Section 5.2). The tW_{2L} analysis channel targets the same signal processes, but considers events where both W bosons decay leptonically (Section 5.3). The two selections are designed to be mutually exclusive. The results of these two analysis channels are statistically combined to maximise the sensitivity to the tW +DM processes. Finally, the tj_{1L} analysis targets t -channel production of DM candidates and requires a single lepton in each event (Section 5.4). In all analysis channels, large missing transverse momentum and jets are required. Event selections and background estimation methods specific to each analysis channel are described in this section, as are the definitions of the signal, control, and validation regions (SR, CR, and VR, respectively).

Dedicated CRs are designed in each analysis channel for the major SM background processes in order to predict their expected contribution in the SRs. The CRs and SRs are mutually exclusive, with the CRs enriched in the major background processes relevant to each analysis channel while minimising the contamination from signal. The potential signal contamination in the CRs is found to be negligible, at the level of $< 3\%$ of the total SM expectation for all analysis channels.

The expected SM backgrounds are first determined independently for each channel, with a profile likelihood fit [76] in a background-only fit. In this fit, normalisation factors of the backgrounds, for which dedicated CRs are defined, are adjusted simultaneously to match the data in the associated CRs. The input to the background-only fit includes the number of events observed in the associated CRs and the number of events predicted by simulation in each CR for all background processes. They are both described by Poisson statistics. The systematic uncertainties, described in Section 6, are included in the fit as nuisance parameters. They are constrained by Gaussian distributions with widths corresponding to the sizes of the uncertainties and are treated as correlated, when appropriate, between the various regions. The product of the various probability density functions forms the likelihood, which the fit maximises by adjusting the background normalisation and the nuisance parameters.

Normalisation and nuisance parameters in the control regions are then used to constrain the corresponding backgrounds in the signal regions. The reliability of the MC extrapolation of the SM background estimates outside of the control regions is verified in dedicated validation regions. Statistically independent from the corresponding CRs and SRs, these VRs are designed to probe a kinematic region closer to that of the SRs. The potential signal contamination in the VRs is at the level of $< 1\%$ of the total SM expectation for most validation regions, and between 8% and 15% in a few validation regions in the tW_{1L} analysis channel.

In the absence of a significant event excess in the SRs, as determined by the background-only fit, upper limits on the interesting properties of the targeted signal models as well as a generic beyond the standard model (BSM) process are extracted from simultaneous fits of the CRs and SRs, referred to as model-dependent and model-independent signal fits as detailed in Section 7.

5.1 Kinematic requirements and event variables

The event selection criteria in each analysis channel are defined using the physics objects described in Section 4 and the event variables defined in this section.

The following variables are defined using simple combinations of the physics objects in the events.

- N_{jet} is the number of jets with $|\eta| < 2.5$ and $p_{\text{T}} > 30$ GeV.
- $N_{\text{jet}}^{\text{forward}}$ is the number of jets in the forward region, $2.5 < |\eta| < 4.5$ and $p_{\text{T}} > 30$ GeV.
- $N_{b\text{-jet}}$ is the number of b -jets with $|\eta| < 2.5$ and p_{T} above a given threshold defined in each analysis channel.
- The minimum azimuthal distance $\Delta\phi_{\text{min}}$ between the $\vec{p}_{\text{T}}^{\text{miss}}$ and the \vec{p}_{T} of each of the four leading jets in the event is useful for rejecting events with mismeasured jet energies leading to $E_{\text{T}}^{\text{miss}}$ in the event, and is defined as:

$$\Delta\phi_{\text{min}} = \min_{i \leq 4} \Delta\phi(\vec{p}_{\text{T}}^{\text{miss}}, \vec{p}_{\text{T},i}^{\text{jet}}),$$

where $\min_{i \leq 4}$ selects the jet that minimises $\Delta\phi$.

- $m_{\ell\ell}$ is the invariant mass of the dilepton system in the event.
- An iterative reclustering approach as defined in Ref. [77] is used to reconstruct the hadronically decaying W bosons. All the signal jets in the event are first reclustered using the anti- k_t algorithm with a large radius parameter of $R = 3.0$. The radius of each large-radius jet is then iteratively reduced to an optimal radius, $R(p_{\text{T}}) = 2 \times m_W / p_{\text{T}}$. The mass of the reclustered jet, $m_W^{\text{reclustered}}$, is used in the $tW_{1\text{L}}$ channel.
- $m_{\ell_1 b_1}$ is the invariant mass of the leading lepton and b -jet in the event.

A set of variables based on transverse mass are defined in order to distinguish between the signal and SM background processes in the following.

- The transverse mass formed by the $\vec{p}_{\text{T}}^{\text{miss}}$ and the leading lepton in the event, $m_{\text{T}}^{\text{lep}}$, is used to reduce the W +jets and semileptonic $t\bar{t}$ backgrounds. It is defined as:

$$m_{\text{T}}^{\text{lep}} = \sqrt{2p_{\text{T}}^{\ell} E_{\text{T}}^{\text{miss}} (1 - \cos \Delta\phi(\vec{p}_{\text{T}}^{\ell}, \vec{p}_{\text{T}}^{\text{miss}}))}.$$

- Similarly, the transverse mass $m_{\text{T}}^{\ell b}$ is formed by the $\vec{p}_{\text{T}}^{\text{miss}}$ and the system of the leading lepton and b -jet in the event to suppress the $W + b$ background, and is defined as:

$$m_{\text{T}}^{\ell b} = \sqrt{2p_{\text{T}}^{\ell+b_1} E_{\text{T}}^{\text{miss}} (1 - \cos \Delta\phi(\vec{p}_{\text{T}}^{\ell+b_1}, \vec{p}_{\text{T}}^{\text{miss}}))}.$$

- Closely related to m_T^{lep} , the transverse mass m_{T2} [78, 79] is used to bound the masses of pair-produced particles, such as in $t\bar{t}$ production, each of which decays so as to produce a visible particle that can be detected and an invisible particle that contributes to the missing transverse momentum. In the case of a dilepton final state, it is defined by:

$$m_{T2}(\vec{p}_T^{\ell_1}, \vec{p}_T^{\ell_2}, \vec{p}_T^{\text{miss}}) = \min_{\vec{q}_T} \left[\max \left(m_T^{\text{lep}}(\vec{p}_T^{\ell_1}, \vec{q}_T), m_T^{\text{lep}}(\vec{p}_T^{\ell_2}, \vec{p}_T^{\text{miss}} - \vec{q}_T) \right) \right],$$

where \vec{q}_T is the transverse momentum vector that minimises the larger of the two transverse masses m_T^{lep} , and $\vec{p}_T^{\ell_1}$ and $\vec{p}_T^{\ell_2}$ are the leading and subleading transverse momenta of the two leptons in the pair.

- The asymmetric transverse mass am_{T2} [80, 81], a variation of m_{T2} , is used in the tW_{1L} final state to reduce the number of dileptonic $t\bar{t}$ background events where one of the leptons is undetected. For these events, am_{T2} has a kinematic endpoint at the top quark mass.

To improve the selection of single-top events in the tW_{2L} channel, the following quantities based on invariant mass are defined.

- $m_{b\ell}^{\text{min}}$ is the minimum invariant mass found by combining the leading b -jet with each of the leptons, $m_{b\ell}^{\text{min}} = \min(m_{b_1\ell_1}, m_{b_1\ell_2})$. An upper endpoint at approximately 153 GeV or 160–170 GeV is expected for the events with one or two leptonic top quark decays, respectively.
- To further reduce the background with two leptonic top quark decays, such as $t\bar{t}$ and $t\bar{t}V$, $m_{b\ell}^t$, an extended variation of $m_{b\ell}^{\text{min}}$, is used in the tW_{2L} final state. It is defined as:

$$m_{b\ell}^t = \min[\max(m_{\ell_1 j_1}, m_{\ell_2 j_2}), \max(m_{\ell_1 j_2}, m_{\ell_2 j_1})],$$

where $m_{\ell_n j_m}$ is the invariant mass of lepton ℓ_n and jet j_m , where j_1 and j_2 are the two jets with highest b -tag discriminator value. For the $t\bar{t}$ and $t\bar{t}V$ backgrounds where both top quarks decay leptonically, $m_{b\ell}^t$ has a kinematic endpoint at approximately 160–170 GeV.

Additional variables based on angular separations of the objects are used in the tj_{1L} analysis to suppress SM background contributions, as defined below.

- $\Delta\eta(\ell_1, b_1)$, $\Delta\phi(\ell_1, b_1)$, and $\Delta R(\ell_1, b_1)$: the pseudorapidity difference, azimuthal angle difference, and angular distance between the leading lepton and b -jet in the event.
- $\Delta\phi(\ell_1, \vec{p}_T^{\text{miss}})$: the azimuthal angle difference between the \vec{p}_T^{miss} and the leading lepton in the event.

Table 2 summarises the trigger and preselection requirements for all analysis channels, in terms of lepton, jet and b -jet multiplicities, as well as transverse momenta and global kinematic variables. Events with extra baseline leptons are vetoed in addition.

Table 2: Summary of the trigger and preselection requirements for the three analysis channels. Opposite-sign leptons are indicated as (OS). Events with extra baseline leptons are vetoed in addition.

Variable	tW _{1L}	tW _{2L}	tj _{1L}
Trigger	E_T^{miss}	dilepton	E_T^{miss} OR one-lepton
N_ℓ^{signal}	= 1	= 2 (OS)	= 1
$p_T(\ell_1)$ [GeV]	> 30	> 25	> 30
$p_T(\ell_2)$ [GeV]	-	> 20	-
N_{jet}	≥ 3	≥ 1	$\in [1, 4]$
$p_T(\text{jet})$ [GeV]	> 30	> 30	> 30
$N_{b\text{-jet}}$	≥ 1	≥ 1	$\in [1, 2]$
$p_T(b_1)$ [GeV]	> 50	> 50	> 50
E_T^{miss} [GeV]	> 250	> 200	> 200
m_T^{lep} [GeV]	> 30	-	> 60
$m_{\ell\ell}$ [GeV]	-	$\geq 40, \notin [71, 111]$ ($ee/\mu\mu$)	-
$\Delta\phi_{\text{min}}$ [rad]	> 0.5	-	> 0.5

5.2 Single-lepton tW_{1L} analysis channel

Events with exactly one electron or muon are first selected for the SR if they also contain at least three jets, exactly one of which must be b -tagged, and satisfy the preselection requirements described in Table 2. The dominant SM background contributions in the channel are $t\bar{t}$, W +jets, and single top (Wt channel) production. Discriminating variables, E_T^{miss} , m_T^{lep} , $m_W^{\text{reclustered}}$ and the asymmetric transverse mass am_{T2} as described in Section 5.1, are used to further separate the signal from backgrounds. A ‘genetic algorithm’ [82] is used to optimise a baseline signal region defined as in Table 3. To increase the sensitivity to different signal model parameters, a binned E_T^{miss} distribution is used as the final input for the statistical analysis. The binning is chosen to be [250, 300] GeV, [300, 400] GeV, [400, 500] GeV, [500, 600] GeV and ≥ 600 GeV, referred to as Bins [0–4].

The acceptance times detector efficiency for the tW +DM signal processes after applying all selection criteria is between 0.3% and 5.1% in the parameter space of $\tan\beta = 1$, $m_a \in [100, 450]$ GeV and $m_{H^\pm} \in [400, 1500]$ GeV, and between 0.2% and 4.8% in the parameter space of $m_a = 250$ GeV, $\tan\beta \in [0.5, 30]$ and $m_{H^\pm} \in [400, 1500]$ GeV.

Dominant background contributions from the $t\bar{t}$ and W +jets processes are estimated using MC simulation and the dedicated CRs. The contribution from multijet production, where the lepton is a misidentified jet or originates from a heavy-flavour hadron decay or photon conversion, is found to be negligible. The remaining sources of background (single-top, Z +jets, diboson, $t\bar{t}V$, and tWZ production, as well as rarer processes such as triboson, $t\bar{t}t\bar{t}$, and $t\bar{t}WW$), are estimated from simulation.

Dedicated control regions $\text{CR}_{tW_{1L}}(t\bar{t})$ and $\text{CR}_{tW_{1L}}(W)$, defined in Table 3, are designed for the $t\bar{t}$ and W +jets background estimations. Compared to the SR, the acceptance for $t\bar{t}$ events is increased in $\text{CR}_{tW_{1L}}(t\bar{t})$ by requiring at least two b -jets, inverting the selection on am_{T2} and removing the requirement on $m_W^{\text{reclustered}}$. To increase the acceptance of the W +jets events and hence the sample size, $\text{CR}_{tW_{1L}}(W)$ is first selected by requiring $40 < m_T^{\text{lep}} < 100$ GeV and $m_W^{\text{reclustered}} < 60$ GeV. To exploit the lepton charge asymmetry of the

Table 3: Summary of signal, control and validation region definitions used in the tW_{1L} analysis channel. The ‘–’ entries represent an inclusive selection with no requirements. The W +jets control and validation regions are each split into two regions with opposite lepton charges as described in the text.

Variable	SR	CR($t\bar{t}$)	CR(W)	VR1($t\bar{t}$)	VR2($t\bar{t}$)	VR1(W)	VR2(W)
$N_{b\text{-jet}}$	= 1	≥ 2	= 1	= 1	= 1	= 1	= 1
$p_T(b_2)$ [GeV]	< 50	> 50	< 50	< 50	< 50	< 50	< 50
$m_W^{\text{reclustered}}$ [GeV]	> 60	–	< 60	–	< 60	> 60	< 60
m_T^{lep} [GeV]	> 200	> 200	$\in [40, 100]$	> 200	> 200	$\in [40, 100]$	> 100
am_{T2} [GeV]	> 220	< 220	> 220	< 220	> 220	> 220	> 220

W +jets events relative to the remaining backgrounds, it is subsequently split into two regions with opposite lepton charges, $CR_{tW_{1L}}(W^+)$ and $CR_{tW_{1L}}(W^-)$. Normalisation factors, $\mu_{t\bar{t}}$ and $\mu_{W\text{+jets}}$, defined as the ratio of the number of observed events to the SM prediction, are found to be 0.96 ± 0.08 and 1.01 ± 0.05 after the background-only fit for the $t\bar{t}$ and W +jets processes, respectively.

To validate the $t\bar{t}$ background predictions and the reliability of MC extrapolation in $m_W^{\text{reclustered}}$ and am_{T2} , two validation regions, VR1 $_{tW_{1L}}(t\bar{t})$ and VR2 $_{tW_{1L}}(t\bar{t})$, are defined by reversing the SR selection requirements on am_{T2} and $m_W^{\text{reclustered}}$ respectively, as shown in Table 3. To increase the sample size, the SR selection requirement on the $m_W^{\text{reclustered}}$ is removed in the VR1 $_{tW_{1L}}(t\bar{t})$ region. Similarly, for the W +jets background processes, two validation regions, VR1 $_{tW_{1L}}(W)$ and VR2 $_{tW_{1L}}(W)$, are defined by varying the SR selection requirements on m_T^{lep} and $m_W^{\text{reclustered}}$ shown in Table 3, respectively. Each of the W +jets validation regions is split into two regions with opposite lepton charge. Figure 2 shows the post-fit E_T^{miss} distributions in the representative validation regions. Good agreement is observed between data and SM expectation in all validation regions. The observed yield, post-fit background estimates and significance [83] in each CR and VR are shown in Figure 3 after the background-only fit. Since the W +jets CR is split into two regions with opposite lepton charges sharing the same normalisation factor, the significances in the CRs are shown explicitly. The data event yields are found to be consistent with background expectations.

5.3 Dilepton tW_{2L} analysis channel

Events with exactly two oppositely charged leptons (electron or muon) are first selected for the SR if they also contain at least one signal jet, at least one of which must be b -tagged with $p_T > 50$ GeV, and satisfy the preselection requirements described in Table 2. The dominant SM background contributions in the channel after these selections are from the $t\bar{t}$, $t\bar{t}Z$, and tWZ processes, followed by that of diboson events. The contribution from misidentified or non-prompt lepton backgrounds (referred to as ‘Fakes /non-prompt’ in Figures 4 and 5) is found to be negligible in the signal region.

Discriminating variables, $m_{b\ell}^{\text{min}}$, $m_{b\ell}^{\text{t}}$, m_{T2} and $\Delta\phi_{\text{min}}$ as defined in Section 5.1, are used to define the final signal region as shown in Table 4.

The acceptance times detector efficiency after applying all selection criteria for the tW +DM signal processes is between 0.07% and 0.7% in the parameter space of $\tan\beta = 1$, $m_a \in [100, 450]$ GeV and $m_{H^\pm} \in [400, 1500]$ GeV, and between 0.05% and 0.6% in the parameter space of $m_a = 250$ GeV, $\tan\beta \in [0.5, 30]$ and $m_{H^\pm} \in [400, 1500]$ GeV.




Figure 2: The E_T^{miss} distributions after the background-only fit (post-fit) are shown in three representative validation regions: (a) $\text{VR2}_{tW_{\text{IL}}}(\text{t}\bar{\text{t}})$, (b) $\text{VR1}_{tW_{\text{IL}}}(W^+)$ and (c) $\text{VR2}_{tW_{\text{IL}}}(W^-)$. The uncertainty bands plotted include all statistical and systematic uncertainties. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The overflow events, where present, are included in the last bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.




Figure 3: Comparison of the predicted backgrounds with the observed numbers of events in the CRs and VRs associated with the tW_{IL} channel. The normalisation of the backgrounds is obtained from the fit to the CRs. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The upper panel shows the observed number of events and the predicted background yield. All uncertainties are included in the uncertainty band. The lower panel shows the significances in each region.

The contributions from the $t\bar{t}$, $t\bar{t}V$ (with $V = W$ or Z boson) and diboson background processes are estimated from MC simulation and dedicated CRs. The remaining sources of background, including the

Table 4: Summary of signal, control and validation region definitions used in the tW_{2L} analysis channel. The ‘-’ entries represent an inclusive selection with no requirements. In the final states with three leptons, the corrected E_T^{miss} , $m_{b\ell}^{\text{min}}$ and m_{T2} variables are used instead as described in the main text. The selection requirement on the corrected $m_{b\ell}^{\text{min}}$ in the $\text{VR}(3\ell)$ region varies according to the jet and b -jet multiplicity as described in the main text. Events with additional baseline leptons are vetoed.

Variable	SR	CR($t\bar{t}$)	CR($t\bar{t}Z$)	CR(WZ)	VR($t\bar{t}$)	VR(3ℓ)
N_ℓ^{signal}	= 2 (OS)	= 2 (OS)	= 3 (≥ 1 SFOS)	= 3 (≥ 1 SFOS)	= 2 (OS)	= 3 (≥ 1 SFOS)
$p_T(\ell_3)$ [GeV]	-	-	> 20	> 20	-	> 20
$m_{ee/\mu\mu}$ [GeV]	$\notin [71, 111]$	$\notin [71, 111]$	$\in [71, 111]$	$\in [71, 111]$	$\notin [71, 111]$	$\in [71, 111]$
N_{jet}	≥ 1	≥ 1	≥ 3	$\in [1, 3]$	≥ 1	≥ 1
$N_{b\text{-jet}}$	≥ 1	≥ 1	≥ 1 (≥ 2 if $N_{\text{jet}} = 3$)	= 1	≥ 1	≥ 1
$m_{b\ell}^{\text{min}}$ [GeV]	< 170	< 170	< 170	> 170	< 170	varies
$m_{b\ell}^t$ [GeV]	> 150	< 150	-	-	> 150	-
m_{T2} [GeV]	> 130	$\in [40, 80]$	> 90	> 90	$\in [40, 80]$	> 90
$\Delta\phi_{\text{min}}$ [rad]	> 1.1	> 1.1	-	-	> 1.1	-

irreducible tWZ process, which is dominated by the $Z \rightarrow \nu\nu$ component, single top quark production, $t\bar{t}h$ production and other rarer processes such as $t\bar{t}t\bar{t}$ and $t\bar{t}WW$, are estimated from simulation.

The acceptance for $t\bar{t}$ events is increased in $\text{CR}_{tW_{2L}}(t\bar{t})$ by requiring a low value of m_{T2} and inverting the SR selection criteria on $m_{b\ell}^t$.

The $t\bar{t}V$ contribution is dominated by the $t\bar{t}Z$ component (about 80% of $t\bar{t}V$ in the SR). A dedicated control region, $\text{CR}_{tW_{2L}}(t\bar{t}Z)$, is defined by first selecting three leptons, where at least one same-flavour–opposite-charge (SFOS) pair is required to be consistent with coming from a Z boson decay with an invariant mass within a window of $[71, 111]$ GeV. If more than one such pair is present in the event, the pair which has an invariant mass closest to the Z boson mass is chosen. The purity of $t\bar{t}Z$ events is further increased by requiring at least three jets. To reduce the diboson background in this region, events with exactly one b -jet and three jets are rejected.

Due to the presence of three leptons in this region, the background contribution from misidentified or non-prompt leptons becomes non-negligible and is estimated using a data-driven matrix method (MM) as described in Refs. [84, 85]. Two types of lepton identification criteria, ‘tight’ and ‘loose’ are defined in the evaluation, corresponding to the baseline and signal lepton selections described in Section 4. The number of events containing misidentified or non-prompt leptons in the $t\bar{t}Z$ CR is estimated from the number of observed events with tight or loose leptons using as input the probability for loose prompt, misidentified or non-prompt leptons to satisfy the tight criteria. The probability for prompt loose leptons to pass the tight selection is determined from $t\bar{t}Z$ MC simulation. The equivalent probability for loose misidentified or non-prompt leptons to pass the tight selection is measured in a $t\bar{t}$ -enriched region with two same-sign leptons (electrons or muons) and a least one b -tagged jet, which is dominated by events with at least one misidentified or non-prompt lepton.

In the $\text{CR}_{tW_{2L}}(t\bar{t}Z)$ region, to mimic the event topology of the signal processes, a corrected \vec{p}_T^{miss} is obtained




Figure 4: The kinematic distributions in the $t\bar{t}$ and 3ℓ validation regions of the tW_{2L} analysis channel after the background-only fit: (a) $m_{b\ell}^t$ in $\text{VR}_{tW_{2L}}(t\bar{t})$, (b) E_T^{miss} in $\text{VR}_{tW_{2L}}(3L)$ and (c) m_{T2} in $\text{VR}_{tW_{2L}}(3L)$. The uncertainty bands plotted include all statistical and systematic uncertainties. The ‘Others’ category includes contributions from rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}W$, and Higgs boson production processes. The overflow events, where present, are included in the last bin. Since the $m_{b\ell}^t$ is defined for events with at least two jets, the events with exactly one jet are included in the overflow bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.

by vectorially adding the transverse momenta of the SFOS pair, and it is subsequently used to calculate the corrected m_{T2} . The two leptons from the SFOS pair are excluded in the calculation of $m_{b\ell}^{\min}$. To improve the estimation of the dominant background from the WZ process in the $\text{CR}_{tW_{2L}}(t\bar{t}Z)$, a dedicated WZ CR, $\text{CR}_{tW_{2L}}(WZ)$, is defined by inverting the $\text{CR}_{tW_{2L}}(t\bar{t}Z)$ selection requirements on the jet multiplicity and the corrected $m_{b\ell}^{\min}$. This CR is also used to aid in the estimation of all diboson processes in the SR. Normalisation factors $\mu_{t\bar{t}}$, $\mu_{t\bar{t}V}$ and μ_{Diboson} are found to be 1.00 ± 0.03 , 0.76 ± 0.26 and 0.80 ± 0.16 after the background-only fit for the $t\bar{t}$, $t\bar{t}V$ and diboson processes, respectively.

A validation region, $\text{VR}_{tW_{2L}}(t\bar{t})$, is defined in order to validate the $t\bar{t}$ background predictions by applying all the signal selection criteria, apart from requiring lower values of m_{T2} , as shown in Table 4. For the background predictions of the $t\bar{t}V$ and diboson processes, a 3ℓ validation region, $\text{VR}_{tW_{2L}}(3L)$, is defined with selection requirements similar to those of the $\text{CR}_{tW_{2L}}(t\bar{t}Z)$ and $\text{CR}_{tW_{2L}}(WZ)$. To ensure that the $\text{VR}_{tW_{2L}}(3L)$ is orthogonal to those two CRs, the selection on the corrected $m_{b\ell}^{\min}$ variable is varied according to the jet and b -jet multiplicities. For the events with exactly one b -jet, the corrected $m_{b\ell}^{\min}$ is required to be larger than 170 GeV if $N_{\text{jet}} > 3$, or smaller than 170 GeV if $N_{\text{jet}} \leq 3$. For the events with more than one b -jet and $N_{\text{jet}} > 2$, the corrected $m_{b\ell}^{\min}$ is required to be larger than 170 GeV. To increase the sample size in this region, the p_T threshold for the b -tagged jets is reduced to 40 GeV.

Figure 4 shows the post-fit kinematic distributions in the validation regions. Good agreement is observed between data and the SM expectation in all validation regions. The observed yield, post-fit background estimates and significance [83] in each CR and VR are shown in Figure 5 after the background-only fit. The data event yields are found to be consistent with background expectations.




Figure 5: Comparison of the predicted backgrounds with the observed numbers of events in the CRs and VRs associated with the tW_{2L} channel. The normalisation of the backgrounds is obtained from the fit to the CRs. The ‘Others’ category includes contributions from rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The upper panel shows the observed number of events and the predicted background yield. All uncertainties are included in the uncertainty band. The lower panel shows the significances estimated for each region.

5.4 Single-lepton tj_{1L} analysis channel

Events with exactly one electron or muon are first selected for the SR if they also contain 1–4 jets with $p_T > 30$ GeV, one or two of which must be b -tagged, and satisfy the preselection requirements described in Table 2. The fourth jet in the event, if present, is required to have $p_T < 50$ GeV. The second b -tagged jet is required to have $p_T > 30$ GeV. The dominant SM background contributions in this channel are from $t\bar{t}$, W +jets, and single top (Wt channel) production. Discriminating variables, E_T^{miss} , m_T^{lep} , $N_{\text{jet}}^{\text{forward}}$ and $\Delta\phi(\ell_1, b_1)$ as described in Section 5.1, are used to define the signal region as shown in Table 5.

To further improve the sensitivity, a boosted decision tree (BDT) is trained to distinguish between signal and background processes, using events passing the preselection defined in Table 2. The following nine kinematic variables defined in Section 5.1 are used as input:

- p_T and η of the highest- p_T jet: $p_T(j_1)$ and $\eta(j_1)$.
- The transverse masses: m_T^{lep} and $m_T^{\ell b}$.
- $\eta^{\ell b}$ of the leading lepton and b -jet system.
- The invariant mass of, and angular distances between, the highest- p_T lepton and b -jet: $m_{\ell_1 b_1}$, $\Delta\phi(\ell_1, b_1)$, and $\Delta R(\ell_1, b_1)$.
- The azimuthal separation between the highest- p_T lepton and missing transverse momentum, $\Delta\phi(\ell_1, \vec{p}_T^{\text{miss}})$.

To explore the kinematic features in different regions of the signal parameter space, samples with different signal model parameters are used as an ensemble in the training. A binned distribution of the BDT output score above 0.6 is then used to extract the final results in the signal regions. The binning of the distribution is optimised as [0.6, 0.75], [0.75, 0.85], [0.85, 0.9] and [0.9, 1.0], referred to as Bins [0–3].

The acceptance times detector efficiency after applying all selection criteria for t -channel production in the signal models is between 0.37% (0.36%) and 0.73% (0.67%) in the parameter space of $m_a = 250$ GeV, $\tan\beta = 0.3$ (0.5) and $m_{H^\pm} \in [500, 1750]$ GeV.

Table 5: Summary of signal, control and validation region definitions used in the $t_{j_{1L}}$ analysis channel. The ‘–’ entries represent an inclusive selection with no requirements.

Variable	SR	CR($t\bar{t}$)	CR(W)	VR($t\bar{t}$)	VR(W)
N_{jet}	$\in [1, 4]$	$\in [1, 4]$	$\in [1, 2]$	$\in [1, 4]$	$\in [1, 2]$
$N_{b\text{-jet}}$	$\in [1, 2]$	= 2	= 1	$\in [1, 2]$	= 1
$N_{\text{jet}}^{\text{forward}}$	> 0	–	–	–	–
$E_{\text{T}}^{\text{miss}}$ [GeV]	> 225	> 225	> 225	> 225	> 225
$m_{\text{T}}^{\text{lep}}$ [GeV]	> 100	> 100	$\in [60, 100]$	> 100	$\in [60, 100]$
$\Delta\phi(\ell_1, b_1)$ [rad]	< 1.2	> 1.5	> 1.5	< 1.5	< 1.5
BDT score	> 0.6	incl.	incl.	< 0.5	incl.

Similarly to the tW_{1L} analysis channel, dominant backgrounds from the $t\bar{t}$ and W +jets processes are estimated using MC simulation and dedicated CRs. The contribution from multijet production is found to be negligible. The remaining sources of background (single-top, Z +jets, diboson, $t\bar{t}V$, $t\bar{t}h$, tWZ production and rarer processes such as triboson, $t\bar{t}t\bar{t}$, and $t\bar{t}WW$) are estimated from simulation.

Dedicated control regions $\text{CR}_{t_{j_{1L}}}(t\bar{t})$ and $\text{CR}_{t_{j_{1L}}}(W)$ are designed to estimate the $t\bar{t}$ and W +jets background processes, respectively, as shown in Table 5. Compared to the SR, the acceptance for $t\bar{t}$ events is increased in $\text{CR}_{t_{j_{1L}}}(t\bar{t})$ by requiring exactly two b -jets and large $\Delta\phi(\ell_1, b_1)$ values. The contribution from W +jets events in the $\text{CR}_{t_{j_{1L}}}(W)$ is enhanced by selecting events with one or two jets, exactly one b -jet, and low $m_{\text{T}}^{\text{lep}}$ and large $\Delta\phi(\ell_1, b_1)$ values. No splitting based on the W boson charge is applied. The normalisation factors $\mu_{t\bar{t}}$ and $\mu_{W\text{+jets}}$ are found to be 1.00 ± 0.27 and 1.10 ± 0.13 for the $t\bar{t}$ and W +jets processes, respectively.

To validate the $t\bar{t}$ background predictions, a validation region $\text{VR}_{t_{j_{1L}}}(t\bar{t})$ is defined by requiring a BDT score that is lower than in the SR definition, as shown in Table 5. For the W +jets background, a validation region $\text{VR}_{t_{j_{1L}}}(W)$ is defined by requiring a lower $m_{\text{T}}^{\text{lep}}$ value than in the SR definition, as shown in Table 5. To ensure orthogonality to the corresponding CRs, events in these two VRs are required to have low $\Delta\phi(\ell_1, b_1)$. Figure 6 shows the post-fit distribution of representative kinematic variables and the BDT score for these two validation regions. Good agreement is observed between data and expectation in all validation regions. The observed yield, post-fit background estimates and significance [83] in each CR and VR are shown in Figure 7 after the background-only fit. The data event yields are found to be consistent with background expectations.




Figure 6: Distributions in the $t\bar{t}$ and W +jets validation regions in the tj_{IL} analysis channel: (a) $N_{jet}^{forward}$ in $VR_{tj_{IL}}(t\bar{t})$, (b) BDT score in $VR_{tj_{IL}}(t\bar{t})$ and (c) E_T^{miss} in $VR_{tj_{IL}}(W)$. The uncertainty bands plotted include all statistical and systematic uncertainties. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The overflow events, where present, are included in the last bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.




Figure 7: Comparison of the predicted backgrounds with the observed numbers of events in the CRs and VRs associated with the tj_{IL} channel. The normalisation of the backgrounds is obtained from the fit to the CRs. The upper panel shows the observed number of events and the predicted background yield. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. All uncertainties are included in the uncertainty band. The lower panel shows the significance for each region.

6 Systematic uncertainties

Several sources of experimental and theoretical systematic uncertainty in the signal and background estimates are considered. Their impact is reduced through the normalisation of the dominant backgrounds in the control regions defined with kinematic selections resembling those of the corresponding signal region. Uncertainties are included as nuisance parameters with Gaussian constraints in the likelihood fits, taking into account correlations between different regions. Uncertainties due to the numbers of events in the CRs are also included in the fit for each region. The magnitude of the contributions arising from detector, theoretical modelling and statistical uncertainties are summarised in Figure 8 as a relative uncertainty in the total background yield for each SR in the three analysis channels.

Dominant detector-related systematic uncertainties arise from the jet energy scale and resolution, and from the b -tagging efficiency and mis-tagging rates.

The uncertainties in the jet energy scale and resolution are based on their respective measurements in data [67] and are derived as a function of the p_T and η of the jet, as well as of the pile-up conditions and the jet flavour composition (light-quark, b -quark, or gluon-initiated jets) of the selected jet sample. Their contributions to the SRs are the dominant experimental uncertainty components and are almost equivalent in all analysis channels. The systematic uncertainty in the b -tagging efficiency is the second largest experimental uncertainty. It ranges from 4.5% for b -jets with $p_T \in [35, 40]$ GeV up to 7.5% for b -jets with high p_T (> 100 GeV). The b -tagging uncertainty is estimated by varying the η -, p_T - and flavour-dependent scale factors applied to each jet in the simulation within a range that reflects the systematic uncertainty in the measured tagging efficiency and mis-tag rates in data [86]. The uncertainties associated with trigger requirements, pile-up modelling, and lepton reconstruction and energy measurements have a small or negligible impact on the final results; however, the lepton, photon and jet-related uncertainties are propagated to the calculation of the E_T^{miss} , and additional uncertainties due to the energy scale and resolution of the soft term are included in the E_T^{miss} . Finally, uncertainties in estimates of the non-prompt




Figure 8: Relative uncertainties in the total background yield in each SR for the three analysis channels, including the contribution from the different sources of uncertainty. The ‘Detector’ category contains all detector-related systematic uncertainties and is dominated by the jet energy scale and resolution. The ‘Background normalisation’ represents the uncertainty in the fitted normalisation factors, including the available event counts in the CRs. Individual uncertainties can be correlated, and do not necessarily add up in quadrature to the total background uncertainty.

or misidentified leptons background are found to be below 1% in the tW_{2L} analysis channel and negligible for single-lepton selections.

Uncertainties in the modelling of the SM background processes in MC simulation and their theoretical cross-section uncertainties are also taken into account. Furthermore, for these processes the 1.7% uncertainty in the combined 2015–2018 integrated luminosity is included.

Modelling uncertainties in the $t\bar{t}$ and single-top backgrounds are dominant in all SRs for the tW_{1L} and tj_{1L} analysis channels, and the second leading source of uncertainty for the tW_{2L} SR. They are computed as the difference between the predictions from nominal samples and those from additional samples differing in hard-scattering generator and parameter settings, or by using internal weights assigned to the events depending on the choice of renormalisation and factorisation scales (μ_R and μ_F , respectively, varied independently by factors of 2 and 0.5), initial- and final-state radiation parameters, and PDF sets. The impact of the PS and hadronisation model is evaluated by comparing the nominal generator with a POWHEG-Box sample interfaced to HERWIG 7 [87, 88], using the H7UE set of tuned parameters [88]. To assess the uncertainty due to the choice of hard-scattering generator and matching scheme, an alternative generator set-up using MADGRAPH5_aMC@NLO interfaced to PYTHIA 8 is employed. For single-top Wt production, the impact of interference between single-resonant and double-resonant top quark production and on the implementation of the W lineshape in the generator is estimated in all analysis channels by comparing the nominal sample generated using the diagram removal method with alternative samples, including those generated using the diagram subtraction method [89]. For the tW_{2L} selection, this results in a 100% uncertainty in the subdominant Wt contribution.

For the $t\bar{t} + W/Z$ background, uncertainties due to parton shower and hadronisation modelling are evaluated by comparing the predictions from MADGRAPH5_aMC@NLO interfaced to PYTHIA 8 and HERWIG 7, while the uncertainties related to the choice of renormalisation and factorisation scales are assessed by varying the corresponding event generator parameters up and down by a factor of two around their nominal values. Their contribution is dominant in the tW_{2L} analysis channel and subdominant or small in all other SRs. A similar approach is used to assess the uncertainties in the tWZ process, with an additional 20% uncertainty assigned to account for uncertainties in the effects of interference between the $t\bar{t} + W/Z$ and tWZ processes. The 20% is assigned on the basis of preliminary comparisons of alternative approaches developed to evaluate interference effects in the $t\bar{t}-tW$ [90] and $t\bar{t}Z-tWZ$ processes [91].

Finally, modelling and normalisation uncertainties in minor backgrounds are also considered. For diboson and W/Z +jets events, they are estimated by varying the renormalisation, factorisation and resummation scales up and down by a factor of two around the values used to generate the nominal samples. For $t\bar{t}WW$, tZ , $t\bar{t}H$, Wh , Zh , $t\bar{t}t\bar{t}$, and triboson production processes, experimental and theoretical uncertainties in the event yields are also evaluated and found to be negligible.

For the DM signal processes, both the experimental and theoretical uncertainties in the expected signal yields are considered, including the aforementioned luminosity uncertainty. Experimental uncertainties are found to be 3–35% (2.5–11%) across the $m_a-m_{H^\pm}$ and $m_a-\tan\beta$ planes for the tW_{1L} (tW_{2L}) analysis channel, and in the range 3–14% as a function of m_{H^\pm} for the tj_{1L} selection, independently of $\tan\beta$. In all SRs, the dominant uncertainty in the signal yields is found to be from the jet energy scale and resolution, followed by uncertainties in b -tagging rates. Larger uncertainties for the tW_{1L} selections are found for the highest E_T^{miss} -binned region, where MC statistical fluctuations are also relevant. In the modelling of the signal samples, uncertainties due to the variations of the renormalisation and factorisation scales are dominant. They are evaluated using a variation scheme wherein μ_R and μ_F are scaled simultaneously by either a factor of 2 or 0.5. For the PS and hadronisation uncertainties, alternative samples with varied A14

tune parameter values are used. The effect of each systematic variation on the acceptance and efficiency is evaluated for each analysis channel SR by comparing the variation samples with the corresponding nominal sample. The impact on the total yields for $tW+DM$, $t\bar{t}+DM$ and t -channel production processes is also evaluated for each signal scenario and found to be between 5% and 15%. For the tW_{1L} and tW_{2L} analysis channels, the uncertainties vary between 5% and 30% across the $m_a-m_{H^\pm}$ and $m_a-\tan\beta$ planes, with the largest values obtained for samples characterised by low values of the H^\pm mass and independently of $\tan\beta$. For the tj_{1L} analysis channel, uncertainties are found to be between 15% and 5% as a function of increasing m_{H^\pm} for all $\tan\beta$ values considered.

7 Results

The event yields for all SRs in the three analysis channels are reported in Tables 6 and 7 and are summarised in Figure 9, where the significance for each of the SRs is also presented. The SM background expectations resulting from background-only fits are shown along with their statistical plus systematic uncertainties. No significant deviations from the expected yields are observed in any of the signal regions considered. The largest background contribution in the tW_{1L} and tj_{1L} analysis channel SRs arises from $t\bar{t}$ production, whilst the contribution from $t\bar{t}V$ is largest in the tW_{2L} SRs, with subdominant contributions from the $t\bar{t}$, single-top (including tWZ) and diboson processes. Other non-negligible background sources are W +jets and Z +jets production.

Table 6: Background-only fit results for the tW_{1L} and tW_{2L} signal regions. The backgrounds which contribute only a small amount (rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$ and Higgs boson production processes, and non-prompt or misidentified leptons background) are grouped and labelled as ‘Others’. The quoted uncertainties of the fitted SM background include both the statistical and systematic uncertainties.

	SR ^{Bin0} _{tW_{1L}}	SR ^{Bin1} _{tW_{1L}}	SR ^{Bin2} _{tW_{1L}}	SR ^{Bin3} _{tW_{1L}}	SR ^{Bin4} _{tW_{1L}}	SR _{tW_{2L}}
Observed events	182	191	60	24	12	12
Fitted SM bkg events	169 ± 14	171 ± 13	55 ± 6	20.1 ± 2.8	15.6 ± 2.8	5.9 ± 1.2
$t\bar{t}$	101 ± 12	84 ± 12	20 ± 5	5.1 ± 1.7	2.3 ± 1.5	1.2 ± 0.9
Single top	16.3 ± 5.2	17.3 ± 5.2	5.4 ± 3.2	2.0 ± 1.8	1.7 ^{+2.0} _{-1.7}	0.26 ^{+0.27} _{-0.26}
W +jets	28 ± 4	37.0 ± 4.3	14.2 ± 2.4	6 ± 1	5.9 ± 1.1	–
Z +jets	2.0 ± 0.9	1.1 ± 0.7	0.3 ± 0.1	0.15 ± 0.04	0.15 ± 0.02	–
Diboson	7.2 ± 1.7	9.6 ± 2.0	4.6 ± 1.0	2.2 ± 0.5	2.7 ± 0.6	0.5 ± 0.2
$t\bar{t}V$	12.3 ± 1.4	19.5 ± 3.5	8.7 ± 1.2	4.0 ± 0.7	2.5 ± 0.5	2.9 ± 0.7
tWZ	1.7 ± 0.2	2.4 ± 0.5	1.17 ± 0.15	0.42 ± 0.09	0.39 ± 0.09	0.8 ± 0.1
Others	0.6 ± 0.1	0.6 ± 0.1	0.17 ± 0.02	0.06 ± 0.02	0.03 ± 0.01	0.16 ± 0.08

Figures 10 to 12 show comparisons between the observed data and the post-fit SM predictions for some relevant kinematic distributions in the three analysis channels after applying all SR selection requirements except the one on the quantity shown (except for Figures 10(c) and 11(b), where all events in the SR are shown). For the tW_{1L} analysis channel, the m_T^{lep} and am_{T2} distributions are shown for all values of E_T^{miss} across the five bins. Similarly, for the tj_{1L} analysis channel, the number of forward jets and E_T^{miss} are shown for all BDT values above 0.6. The expected distributions for representative scenarios with different m_a , m_{H^\pm} , and $\tan\beta$ (depending on the analysis channel and SR considered) are shown for illustrative purposes.

Table 7: Background-only fit results for the $t_{j_{1L}}$ signal regions. The backgrounds which contribute only a small amount (Z +jets, rare processes such as tWZ , triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$ and Higgs boson production processes) are grouped and labelled as ‘Others’. The quoted uncertainties of the fitted SM background include both the statistical and systematic uncertainties.

	$SR_{t_{j_{1L}}}^{\text{Bin0}}$	$SR_{t_{j_{1L}}}^{\text{Bin1}}$	$SR_{t_{j_{1L}}}^{\text{Bin2}}$	$SR_{t_{j_{1L}}}^{\text{Bin3}}$
Observed events	360	178	69	29
Fitted SM bkg events	335 ± 74	187 ± 40	67 ± 18	37 ± 7
$t\bar{t}$	280 ± 75	151 ± 42	54 ± 16	30 ± 8
W +jets	14.4 ± 5.1	12.4 ± 8.7	$2.7^{+7.5}_{-2.7}$	<0.1
Single top	27 ± 14	13.2 ± 7.5	5.7 ± 3.7	3.8 ± 2.5
$t\bar{t}V$	5.0 ± 1.6	3.1 ± 1.0	1.4 ± 0.6	1.3 ± 0.5
Diboson	3.3 ± 1.0	2.5 ± 0.8	1.4 ± 0.5	0.8 ± 0.3
Others	4.9 ± 2.4	4.8 ± 2.0	2.0 ± 0.7	1.0 ± 0.4

Reasonable agreement is found between data and SM predictions in all distributions, although a mild excess of data events is found in the tW_{2L} distributions, accounting for a discrepancy lower than 2σ considering statistical and systematic uncertainties.




Figure 9: Results of the background-only fit extrapolated to all SRs. The normalisation of the backgrounds is obtained from the fit to the CRs. The upper panel shows the observed number of events and the predicted background yields. The ‘Others’ category includes contributions from rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. All uncertainties defined in Section 6 are included in the uncertainty band. The lower panel shows the significance in each SR. The significance calculation is performed as described in Ref. [83].




Figure 10: Representative distributions of (a) m_T^{lep} , (b) am_{T2} and (c) E_T^{miss} in the tW_{IL} channel. Observed data are compared with post-fit SM predictions. All SR selections except the one on the quantity shown are applied. The SR requirement is indicated by the arrow. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The expected distributions for representative scenarios with different m_a , m_{H^\pm} , and $\tan\beta$ are shown for illustrative purposes. The overflow events, where present, are included in the last bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.




Figure 11: Representative distributions of (a) $m_{b\ell}^t$, (b) E_T^{miss} and (c) m_{T2} in the tW_{2L} channel. Observed data are compared with post-fit SM predictions. All SR selections except the one on the quantity shown are applied. The SR requirement is indicated by the arrow. As the $m_{b\ell}^t$ is defined for events with at least two jets, the events with exactly one jet are included in the overflow bin. The ‘Others’ category includes contributions from rare processes such as triboson, $t\bar{t}\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The expected distributions for representative scenarios with different m_a , m_{H^\pm} , and $\tan\beta$ are shown for illustrative purposes. The overflow events, where present, are included in the last bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.

7.1 Statistical combination of the tW_{1L} and tW_{2L} analysis channels

A statistical combination of results from the tW_{1L} and tW_{2L} channels is performed to maximise the sensitivity to tW +DM models. The simultaneous fit is performed such that the individual background normalisation factors, $\mu_{t\bar{t}}^{1L}$, $\mu_{t\bar{t}}^{2L}$, $\mu_{W+\text{jets}}$ and $\mu_{t\bar{t}V}$, are constrained in the same regions as the respective, individual analyses to avoid extrapolations into a different phase space. Experimental uncertainties in the background and signal are evaluated using the same methods as described in Section 5 and correlated across channels. Modelling uncertainties from the same source for a given process are correlated, e.g. all modelling uncertainties for $t\bar{t}$ are correlated across the regions. Signal systematic uncertainties are also correlated for the exclusion fits described in the next section.

The predictions for SM backgrounds are, as expected, equivalent to those of the individual channels. In particular, the values for the $\mu_{t\bar{t}}$ background normalisation factor are found to be consistent for tW_{1L} and tW_{2L} estimates, $\mu_{t\bar{t}}^{1L} = 0.97 \pm 0.08$ and $\mu_{t\bar{t}}^{2L} = 1.00 \pm 0.03$, respectively.

7.2 Model-independent limits

The CL_s technique [92] is used to place 95% confidence level (CL) upper limits on event yields from physics BSM for each signal region (model-independent limits). The profile-likelihood-ratio test statistic is used to exclude the signal-plus-background hypotheses for specific signal models. When normalised to the integrated luminosity of the data sample, results can be interpreted as corresponding upper limits on the visible cross section, σ_{vis} , defined as the product of the BSM production cross section, the acceptance and the selection efficiency of a BSM signal. In the case of the tW_{1L} analysis channel, the E_T^{miss} bins are defined inclusively, i.e. all events above the lowest bin-threshold in E_T^{miss} are taken, to retain discovery potential. The SM predictions and their corresponding uncertainties are reported in Table 8. In the case of the tj_{1L} analysis channel, the last bin of the BDT score distribution, 0.9–1.0, is considered.

Table 8: Background-only fit results for tW_{1L} signal region bins for model-independent limits. Events with E_T^{miss} above the lowest bin-threshold are retained in each bin. This is indicated by value X in $SR_{tW_{1L}}^X$. The backgrounds which contribute only a small amount (rare processes such as tWZ , triboson, $t\bar{t} t\bar{t}$, $t\bar{t} WW$ and Higgs boson production processes) are grouped and labelled as ‘Others’. The quoted uncertainties of the fitted SM background include both the statistical and systematic uncertainties.

	$SR_{tW_{1L}}^{250}$	$SR_{tW_{1L}}^{300}$	$SR_{tW_{1L}}^{400}$	$SR_{tW_{1L}}^{500}$	$SR_{tW_{1L}}^{600}$
Observed events	469	287	96	36	12
Fitted SM bkg events	431 ± 27	262 ± 20	91 ± 10	36 ± 5	15.5 ± 2.8
$t\bar{t}$	213 ± 25	111 ± 18	28 ± 7	7.5 ± 2.9	2.3 ± 1.5
Single top	43 ± 15	27 ± 12	9 ± 7	3.9 ± 3.8	$1.7^{+2.0}_{-1.7}$
W +jets	91 ± 8	63 ± 5	26 ± 3	12.0 ± 1.7	5.9 ± 1.1
Z +jets	3.8 ± 1.0	1.7 ± 0.7	0.6 ± 0.1	0.30 ± 0.05	0.15 ± 0.02
Diboson	26 ± 5	19.0 ± 3.9	9 ± 2	4.8 ± 1.0	2.6 ± 0.6
$t\bar{t}V$	47 ± 3	34.7 ± 2.4	15.3 ± 1.2	6.6 ± 0.5	2.5 ± 0.3
Others	7.5 ± 0.5	5.3 ± 0.4	2.3 ± 0.1	0.9 ± 0.1	0.40 ± 0.05

Table 9: The 95% CL upper limits on the visible cross section ($\langle\epsilon\sigma\rangle_{\text{obs}}^{95}$) and on the number of signal events (S_{obs}^{95}) for all SRs and analysis channels as detailed in the text. The fourth column (S_{exp}^{95}) shows the 95% CL upper limit on the number of signal events, given the expected number (and $\pm 1\sigma$ exclusions of the expectation) of background events. The last two columns indicate the CL_B value, i.e. the confidence level observed for the background-only hypothesis, and the discovery p -value ($p(s=0)$). The associated significance is provided in parentheses.

Signal channel	$\langle\epsilon\sigma\rangle_{\text{obs}}^{95}$ [fb]	S_{obs}^{95}	S_{exp}^{95}	CL_B	$p(s=0)$ (Z)
SR _{tW_{1L}} ²⁵⁰	0.72	100.6	67 ⁺³³ ₋₁₆	0.85	0.12 (1.16)
SR _{tW_{1L}} ³⁰⁰	0.51	70.8	54 ± 16	0.85	0.15 (1.02)
SR _{tW_{1L}} ⁴⁰⁰	0.24	32.9	29 ⁺¹⁰ ₋₆	0.64	0.30 (0.52)
SR _{tW_{1L}} ⁵⁰⁰	0.14	18.9	19 ⁺⁸ ₋₅	0.52	0.45 (0.13)
SR _{tW_{1L}} ⁶⁰⁰	0.08	10.6	12 ⁺³ ₋₄	0.24	0.94 (-1.54)
SR _{tW_{2L}}	0.10	13.8	7.3 ^{+2.9} _{-1.1}	0.97	0.02 (2.07)
SR _{ij_{1L}} (BDT>0.9)	0.10	14.4	19 ⁺⁶ ₋₅	0.24	0.50 (0.00)

Table 9 summarises the observed (S_{obs}^{95}) and expected (S_{exp}^{95}) 95% CL upper limits on the number of BSM events and on σ_{vis} for all SRs. The p_0 -values, which represent the probability of the SM background to fluctuate to the observed number of events or higher, are also provided and are capped at $p_0 = 0.5$; the associated significance is provided in parentheses.

7.3 Model-dependent limits

Model-dependent exclusion limits are placed on the common signal parameters m_a , m_{H^\pm} , and $\tan\beta$ in the 2HDM+ a models considered in the analysis. Following the prescriptions in Ref. [5], the masses of the bosons H , H^\pm and A are set to be equal. The three quartic couplings between the scalar doublets and the a boson (λ_{P1} , λ_{P2} and λ_3) are all set equal to 3, in order to reduce the number of parameters and evade the constraints from electroweak precision measurements [18]. To further reduce the parameter space, unitary couplings between the a boson mediator and the DM particle χ ($g_\chi = 1$) are considered, with the DM particle mass set to $m_\chi = 10$ GeV. The mixing angle θ is fixed at $\sin\theta = 1/\sqrt{2}$, yielding full mixing between the a and A bosons and the largest cross sections for the processes of interest. Two sets of samples are considered,² varying either the (m_a, m_{H^\pm}) parameters and setting $\tan\beta$ to unity, or varying the $(m_{H^\pm}, \tan\beta)$ parameters and setting $m_a = 250$ GeV. The fit procedure takes into account correlations in the yield predictions between control and signal regions due to common background normalisation parameters and systematic uncertainties. The experimental systematic uncertainties in the signal are taken into account for the calculation and are assumed to be fully correlated with those in the SM background. The results of the combined fit for the tW_{1L} and tW_{2L} channels are interpreted using the sum of the respective signal yield estimates for each generated sample, with overlap between the samples removed according to the procedure illustrated in Ref. [22].

² As pointed out in Ref. [5], for the two considered sets of samples, only values $m_{H^\pm} < 600$ GeV provide a bounded-from-below scalar potential [20] for the 2HDM+ a model. This constraint can be relaxed up to a factor 2 if the quartic coupling λ_3 assumes a value closer to the perturbativity limit and they can be relaxed further in more general 2HDMs containing additional quartic couplings [93], as discussed in Ref. [94].

Figures 13(a) and 13(b) show the observed and expected exclusion contours as functions of (m_a, m_{H^\pm}) and $(m_{H^\pm}, \tan \beta)$, respectively, for the tW_{1L} and tW_{2L} channels, presented both individually and statistically combined. In this case, only the DMt contribution of the signal is taken into account to better illustrate the sensitivity to single-top signatures. Figures 14(a) and 14(b) show the observed and expected exclusion contours for the same models, but also include the expected contributions from the $DMt\bar{t}$ process. Figures 13 and 14 also report the 1σ and 2σ uncertainty bands around the observed limit contour, as well as the variations obtained by changing the theoretical cross-section predictions for signal to be 15% above or below the nominal value (as this is expected to be largest uncertainty in the signal yields across the plane). For low H^\pm masses, DMt production generally dominates $DMt\bar{t}$ production, due to the contribution from the resonant H^\pm diagrams, except when the mass difference $m_{H^\pm} - m_a$ is small enough to suppress the branching fraction of $H^\pm \rightarrow Wa$ decay relative to $H^\pm \rightarrow t\bar{b}$. On the other hand, $DMt\bar{t}$ contributions are dominant at high m_{H^\pm} . The width of H^\pm also increases at high m_{H^\pm} , and it is about 20% of its mass for $m_{H^\pm} = 1$ TeV. Moreover, as studied in Ref. [22], the $DMt\bar{t}$ cross section is proportional to $1/\tan^2 \beta$, whereas the H^\pm production cross section has a more complex dependence, with a minimum for $\tan \beta \sim 10$ and an enhancement for high values of $\tan \beta$. For $\tan \beta = 1$ and $m_{H^\pm} \sim m_a + m_W$, the $DMt\bar{t}$ cross section also dominates the DMt cross section. Assuming $m_\chi = 10$ GeV and $g_\chi = 1$, masses of a below 190 GeV are excluded at 95% CL for all values of m_{H^\pm} in the range 400–1400 GeV, and up to 330 GeV for m_{H^\pm} around 800 GeV. When only DMt contributions are taken into account, the constraints on m_a decrease by 20–50 GeV. In the case where $m_a = 250$ GeV, all values of m_{H^\pm} between 450 GeV and 1.5 TeV are excluded for $\tan \beta$ around and below unity, and scenarios with $\tan \beta$ below 1.5 are excluded for masses of H^\pm around 800 GeV.

The sensitivity of the tj_{1L} channel is small compared to the other analysis channels because of the smaller cross section of the DMt -channel production process. The observed and expected cross-section limits at 95% CL as a function of m_{H^\pm} for two representative values of $\tan \beta$ are shown in Figure 15 assuming a fixed value of $m_a = 250$ GeV. The limits are shown as a multiple of σ_{BSM} , the theoretical cross section of the t -channel DM production process. For $\tan \beta = 0.3$, H^\pm masses above 900 GeV are excluded under these hypotheses, whilst no exclusion is obtained for $\tan \beta = 0.5$.






Figure 12: Representative distributions of (a) BDT score, (b) $N_{\text{jet}}^{\text{forward}}$ and (c) $E_{\text{T}}^{\text{miss}}$ in the t_{J1L} channel. Observed data are compared with post-fit SM predictions. All SR selections except the one on the quantity shown are applied. The SR requirement is indicated by the arrow. The ‘Others’ category includes contributions from Z +jets and tWZ production, and rare processes such as triboson, $t\bar{t}t\bar{t}$, $t\bar{t}WW$, and Higgs boson production processes. The expected distributions for representative scenarios with different m_a , m_{H^\pm} , and $\tan\beta$ are shown for illustrative purposes. The overflow events, where present, are included in the last bin. The lower panels show the ratio of data to the background prediction. The hatched error bands indicate the combined experimental and MC statistical uncertainties on these background predictions.




(a) DMt, m_a - m_{H^\pm} plane




(b) DMt, m_{H^\pm} - $\tan\beta$ plane

Figure 13: The expected and observed exclusion contours as a function of (m_a, m_{H^\pm}) (top) and $(m_{H^\pm}, \tan\beta)$ (bottom), assuming only tW +DM contributions, for the individual tW_{1L} (purple line) and tW_{2L} (pink line) analysis channels, and for their statistical combination (green line). Experimental and theoretical systematic uncertainties, as described in Section 6, are applied to background and signal samples and illustrated by the ± 1 standard-deviation and ± 2 standard-deviation yellow bands and the green dotted contour lines, respectively, for the statistical combination.



(a) $DMt+DMt\bar{t}$, $m_a-m_{H^\pm}$ plane



(b) $DMt+DMt\bar{t}$, $m_{H^\pm}-\tan\beta$ plane

Figure 14: The expected and observed exclusion contours as a function of (m_a, m_{H^\pm}) (top) and $(m_{H^\pm}, \tan\beta)$ (bottom), assuming $DMt\bar{t}$ and DMt contributions, for the individual tW_{1L} (purple line) and tW_{2L} (pink line) analysis channels, and for their statistical combination (green line). Experimental and theoretical systematic uncertainties, as described in Section 6, are applied to background and signal samples and illustrated by the ± 1 standard-deviation and ± 2 standard-deviation yellow bands and the green dotted contour lines, respectively, for the statistical combination.

8 Conclusion

A search for dark matter has been performed in the context of a two-Higgs-doublet model together with an additional pseudoscalar mediator, a , which decays into the dark-matter particles. Processes where the pseudoscalar mediator is produced in association with a single top quark in the 2HDM+ a model are explored for the first time at the LHC. Several final states which include either one or two leptons (electrons or muons) and a significant amount of missing transverse momentum are considered. The analysis makes use of proton–proton collision data at $\sqrt{s} = 13$ TeV collected by the ATLAS experiment during LHC Run 2 (2015–2018), corresponding to an integrated luminosity of 139 fb^{-1} . No significant excess above the Standard Model predictions is found. The results are expressed as 95% confidence-level limits on the 2HDM+ a signal models considered. Assuming dark-matter particles with mass $m_\chi = 10$ GeV and coupling $g_\chi = 1$ to the mediator, and full mixing between the a and A bosons, masses of a below 200 GeV are excluded at 95% CL for all values of m_{H^\pm} in the range 400–1400 GeV, and up to 330 GeV for m_{H^\pm} around 900 GeV. For $m_a = 250$ GeV, all values of m_{H^\pm} below 1.5 TeV are excluded for $\tan\beta$ below unity, and scenarios with $\tan\beta$ below 1.5 are excluded for masses of H^\pm around 800 GeV.

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTC, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; COLCIENCIAS, Colombia; MSMT CR, MPO CR and VSC CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRT, Greece; RGC and Hong Kong SAR, China; ISF and Benozio Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW and NCN, Poland; FCT, Portugal; MNE/IFA, Romania; JINR; MES of Russia and NRC KI, Russian Federation; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DST/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TAEK, Turkey; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada, CRC and IVADO, Canada; Beijing Municipal Science & Technology Commission, China; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and GIF, Israel; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [95].

References

- [1] G. Hinshaw et al., *Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results*, *Astrophys. J. Suppl.* **208** (2013) 19, arXiv: 1212.5226 [astro-ph.CO].
- [2] Planck Collaboration, *Planck 2018 results. I. Overview and the cosmological legacy of Planck*, *Astron. Astrophys.* **641** (2020) A1, arXiv: 1807.06205 [astro-ph.CO].
- [3] G. Steigman and M. S. Turner, *Cosmological constraints on the properties of weakly interacting massive particles*, *Nucl. Phys. B* **253** (1985) 375.
- [4] M. Bauer, U. Haisch and F. Kahlhoefer, *Simplified dark matter models with two Higgs doublets: I. Pseudoscalar mediators*, *JHEP* **05** (2017) 138, arXiv: 1701.07427 [hep-ph].
- [5] T. Abe et al., *LHC Dark Matter Working Group: Next-generation spin-0 dark matter models*, *Phys. Dark Univ.* **27** (2020) 100351, arXiv: 1810.09420 [hep-ex].
- [6] P. J. Fox and E. Poppitz, *Leptophilic dark matter*, *Phys. Rev. D* **79** (2009) 083528, arXiv: 0811.0399 [hep-ph].
- [7] S. Cassel, D. M. Ghilencea and G. G. Ross, *Electroweak and dark matter constraints on a Z' in models with a hidden valley*, *Nucl. Phys. B* **827** (2010) 256, arXiv: 0903.1118 [hep-ph].
- [8] Y. Bai, P. J. Fox and R. Harnik, *The Tevatron at the frontier of dark matter direct detection*, *JHEP* **12** (2010) 048, arXiv: 1005.3797 [hep-ph].
- [9] J. Abdallah et al., *Simplified models for dark matter searches at the LHC*, *Phys. Dark Univ.* **9-10** (2015) 8, arXiv: 1506.03116 [hep-ph].
- [10] D. Abercrombie et al., *Dark Matter benchmark models for early LHC Run-2 Searches: Report of the ATLAS/CMS Dark Matter Forum*, *Phys. Dark Univ.* **27** (2020) 100371, ed. by A. Boveia, C. Doglioni, S. Lowette, S. Malik and S. Mrenna, arXiv: 1507.00966 [hep-ex].
- [11] K. Cheung, K. Mawatari, E. Senaha, P.-Y. Tseng and T.-C. Yuan, *The top window for dark matter*, *JHEP* **10** (2010) 081, arXiv: 1009.0618 [hep-ph].
- [12] U. Haisch, A. Hibbs and E. Re, *Determining the structure of dark-matter couplings at the LHC*, *Phys. Rev. D* **89** (2014) 034009, arXiv: 1311.7131 [hep-ph].
- [13] M. R. Buckley, D. Feld and D. Goncalves, *Scalar simplified models for dark matter*, *Phys. Rev. D* **91** (2015) 015017, arXiv: 1410.6497 [hep-ph].
- [14] M. R. Buckley and D. Goncalves, *Constraining the strength and CP structure of dark production at the LHC: The associated top-pair channel*, *Phys. Rev. D* **93** (2016) 034003, arXiv: 1511.06451 [hep-ph].
- [15] U. Haisch and E. Re, *Simplified dark matter top-quark interactions at the LHC*, *JHEP* **06** (2015) 078, arXiv: 1503.00691 [hep-ph].
- [16] M. Backovic et al., *Higher-order QCD predictions for dark matter production at the LHC in simplified models with s-channel mediators*, *Eur. Phys. J. C* **75** (2015) 482, arXiv: 1508.05327 [hep-ph].
- [17] C. Arina et al., *A comprehensive approach to dark matter studies: exploration of simplified top-philic models*, *JHEP* **11** (2016) 111, arXiv: 1605.09242 [hep-ph].
- [18] U. Haisch, P. Pani and G. Polesello, *Determining the CP nature of spin-0 mediators in associated production of dark matter and $t\bar{t}$ pairs*, *JHEP* **02** (2017) 131, arXiv: 1611.09841 [hep-ph].

- [19] S. Banerjee et al., *Cornering pseudoscalar-mediated dark matter with the LHC and cosmology*, *JHEP* **07** (2017) 080, arXiv: [1705.02327 \[hep-ph\]](#).
- [20] J. F. Gunion and H. E. Haber, *CP-conserving two Higgs doublet model: The approach to the decoupling limit*, *Phys. Rev. D* **67** (2003) 075019, arXiv: [hep-ph/0207010 \[hep-ph\]](#).
- [21] G. C. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1, arXiv: [1106.0034 \[hep-ph\]](#).
- [22] ATLAS Collaboration, *Constraints on mediator-based dark matter and scalar dark energy models using $\sqrt{s} = 13$ TeV pp collision data collected by the ATLAS detector*, *JHEP* **05** (2019) 142, arXiv: [1903.01400 \[hep-ex\]](#).
- [23] P. Pani and G. Polese, *Dark matter production in association with a single top-quark at the LHC in a two-Higgs-doublet model with a pseudoscalar mediator*, *Phys. Dark Univ.* **21** (2018) 8, arXiv: [1712.03874 \[hep-ph\]](#).
- [24] CMS Collaboration, *Search for dark matter produced in association with a Higgs boson decaying to a pair of bottom quarks in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 280, arXiv: [1811.06562 \[hep-ex\]](#).
- [25] CMS Collaboration, *Search for dark matter produced in association with a single top quark or a top quark pair in proton–proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2019) 141, arXiv: [1901.01553 \[hep-ex\]](#).
- [26] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [27] ATLAS Collaboration, *ATLAS Insertable B-Layer Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>.
- [28] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, *JINST* **13** (2018) T05008, arXiv: [1803.00844 \[physics.ins-det\]](#).
- [29] ATLAS Collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317, arXiv: [1611.09661 \[hep-ex\]](#).
- [30] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, ATLAS-CONF-2019-021, 2019, URL: <https://cds.cern.ch/record/2677054>.
- [31] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *JINST* **13** (2018) P07017.
- [32] ATLAS Collaboration, *2015 start-up trigger menu and initial performance assessment of the ATLAS trigger using Run-2 data*, ATL-DAQ-PUB-2016-001, 2016, URL: <https://cds.cern.ch/record/2136007>.
- [33] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, *Eur. Phys. J. C* **70** (2010) 823, arXiv: [1005.4568 \[physics.ins-det\]](#).
- [34] S. Agostinelli et al., *GEANT4 – a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250.
- [35] T. Sjöstrand, S. Mrenna and P. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852, arXiv: [0710.3820 \[hep-ph\]](#).
- [36] R. D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys. B* **867** (2013) 244, arXiv: [1207.1303 \[hep-ph\]](#).

- [37] ATLAS Collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie–Landshoff diffractive model*, ATL-PHYS-PUB-2016-017, 2016, URL: <https://cds.cern.ch/record/2206965>.
- [38] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079, arXiv: [1405.0301](https://arxiv.org/abs/1405.0301) [[hep-ph](#)].
- [39] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159, arXiv: [1410.3012](https://arxiv.org/abs/1410.3012) [[hep-ph](#)].
- [40] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [41] R. D. Ball et al., *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040, arXiv: [1410.8849](https://arxiv.org/abs/1410.8849) [[hep-ph](#)].
- [42] P. Artoisenet, R. Frederix, O. Mattelaer and R. Rietkerk, *Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations*, *JHEP* **03** (2013) 015, arXiv: [1212.3460](https://arxiv.org/abs/1212.3460) [[hep-ph](#)].
- [43] T. Gleisberg, S. Höche, F. Krauss, M. Schönherr, S. Schumann et al., *Event generation with SHERPA 1.1*, *JHEP* **02** (2009) 007, arXiv: [0811.4622](https://arxiv.org/abs/0811.4622) [[hep-ph](#)].
- [44] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, *JHEP* **12** (2008) 039, arXiv: [0808.3674](https://arxiv.org/abs/0808.3674) [[hep-ph](#)].
- [45] F. Cascioli, P. Maierhofer and S. Pozzorini, *Scattering Amplitudes with Open Loops*, *Phys. Rev. Lett.* **108** (2012) 111601, arXiv: [1111.5206](https://arxiv.org/abs/1111.5206) [[hep-ph](#)].
- [46] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani–Seymour dipole factorisation*, *JHEP* **03** (2008) 038, arXiv: [0709.1027](https://arxiv.org/abs/0709.1027) [[hep-ph](#)].
- [47] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements + parton showers: The NLO case*, *JHEP* **04** (2013) 027, arXiv: [1207.5030](https://arxiv.org/abs/1207.5030) [[hep-ph](#)].
- [48] D. J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152.
- [49] S. Frixione, P. Nason and G. Ridolfi, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126, arXiv: [0707.3088](https://arxiv.org/abs/0707.3088) [[hep-ph](#)].
- [50] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040, arXiv: [hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146).
- [51] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070, arXiv: [0709.2092](https://arxiv.org/abs/0709.2092) [[hep-ph](#)].
- [52] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043, arXiv: [1002.2581](https://arxiv.org/abs/1002.2581) [[hep-ph](#)].
- [53] M. Czakon and A. Mitov, *Top++: A program for the calculation of the top-pair cross-section at hadron colliders*, *Comput. Phys. Commun.* **185** (2014) 2930, arXiv: [1112.5675](https://arxiv.org/abs/1112.5675) [[hep-ph](#)].
- [54] R. Frederix, E. Re and P. Torrielli, *Single-top t -channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO*, *JHEP* **09** (2012) 130, arXiv: [1207.5391](https://arxiv.org/abs/1207.5391) [[hep-ph](#)].
- [55] N. Kidonakis, *Next-to-next-to-leading-order collinear and soft gluon corrections for t -channel single top quark production*, *Phys. Rev. D* **83** (2011) 091503, arXiv: [1103.2792](https://arxiv.org/abs/1103.2792) [[hep-ph](#)].

- [56] E. Re, *Single-top Wt -channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547, arXiv: [1009.2450 \[hep-ph\]](#).
- [57] N. Kidonakis, *Two-loop soft anomalous dimensions for single top quark associated production with a W - or H -*, *Phys. Rev. D* **82** (2010) 054018, arXiv: [1005.4451 \[hep-ph\]](#).
- [58] N. Kidonakis, *Next-to-next-to-leading logarithm resummation for s -channel single top quark production*, *Phys. Rev. D* **81** (2010) 054028, arXiv: [1001.5034 \[hep-ph\]](#).
- [59] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector Boson Production at Hadron Colliders: A Fully Exclusive QCD Calculation at Next-to-Next-to-Leading Order*, *Phys. Rev. Lett.* **103** (2009) 082001, arXiv: [0903.2120 \[hep-ph\]](#).
- [60] D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, (2016), arXiv: [1610.07922 \[hep-ph\]](#).
- [61] ATLAS Collaboration, *Vertex Reconstruction Performance of the ATLAS Detector at $\sqrt{s} = 13$ TeV*, ATL-PHYS-PUB-2015-026, 2015, URL: <https://cds.cern.ch/record/2037717>.
- [62] ATLAS Collaboration, *Selection of jets produced in 13 TeV proton–proton collisions with the ATLAS detector*, ATLAS-CONF-2015-029, 2015, URL: <https://cds.cern.ch/record/2037702>.
- [63] ATLAS Collaboration, *Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton–proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 639, arXiv: [1902.04655 \[hep-ex\]](#).
- [64] ATLAS Collaboration, *Muon reconstruction performance of the ATLAS detector in proton–proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **76** (2016) 292, arXiv: [1603.05598 \[hep-ex\]](#).
- [65] ATLAS Collaboration, *Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1*, *Eur. Phys. J. C* **77** (2017) 490, arXiv: [1603.02934 \[hep-ex\]](#).
- [66] M. Cacciari, G. P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063, arXiv: [0802.1189 \[hep-ph\]](#).
- [67] ATLAS Collaboration, *Jet energy scale measurements and their systematic uncertainties in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **96** (2017) 072002, arXiv: [1703.09665 \[hep-ex\]](#).
- [68] ATLAS Collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 581, arXiv: [1510.03823 \[hep-ex\]](#).
- [69] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896, arXiv: [1111.6097 \[hep-ph\]](#).
- [70] ATLAS Collaboration, *ATLAS b -jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970, arXiv: [1907.05120 \[hep-ex\]](#).
- [71] ATLAS Collaboration, *Optimisation and performance studies of the ATLAS b -tagging algorithms for the 2017-18 LHC run*, ATL-PHYS-PUB-2017-013, 2017, URL: <https://cds.cern.ch/record/2273281>.
- [72] M. Cacciari, G. P. Salam and G. Soyez, *The catchment area of jets*, *JHEP* **04** (2008) 005, arXiv: [0802.1188 \[hep-ph\]](#).
- [73] ATLAS Collaboration, *Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run 2 data collected in 2015 and 2016*, *Eur. Phys. J. C* **79** (2019) 205, arXiv: [1810.05087 \[hep-ex\]](#).

- [74] ATLAS Collaboration, *Performance of missing transverse momentum reconstruction with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **78** (2018) 903, arXiv: [1802.08168 \[hep-ex\]](#).
- [75] ATLAS Collaboration, *E_T^{miss} performance in the ATLAS detector using 2015–2016 LHC pp collisions*, ATLAS-CONF-2018-023, 2018, URL: <https://cds.cern.ch/record/2625233>.
- [76] M. Baak et al., *HistFitter software framework for statistical data analysis*, *Eur. Phys. J. C* **75** (2015) 153, arXiv: [1410.1280 \[hep-ex\]](#).
- [77] ATLAS Collaboration, *Search for top-squark pair production in final states with one lepton, jets, and missing transverse momentum using 36.1 fb⁻¹ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector*, *JHEP* **06** (2018) 108, arXiv: [1711.11520 \[hep-ex\]](#).
- [78] C. Lester and D. Summers, *Measuring masses of semi-invisibly decaying particles pair produced at hadron colliders*, *Phys. Lett. B* **463** (1999) 99, arXiv: [hep-ph/9906349](#).
- [79] A. Barr, C. Lester and P. Stephens, *A variable for measuring masses at hadron colliders when missing energy is expected; m_{T2} : the truth behind the glamour*, *J. Phys. G* **29** (2003) 2343, arXiv: [hep-ph/0304226 \[hep-ph\]](#).
- [80] P. Konar, K. Kong, K. T. Matchev and M. Park, *Dark matter particle spectroscopy at the LHC: generalizing M_{T2} to asymmetric event topologies*, *JHEP* **04** (2010) 086, arXiv: [0911.4126 \[hep-ph\]](#).
- [81] C. G. Lester and B. Nachman, *Bisection-based asymmetric M_{T2} computation: a higher precision calculator than existing symmetric methods*, *JHEP* **03** (2015) 100, arXiv: [1411.4312 \[hep-ph\]](#).
- [82] K. Sastry, D. Goldberg and G. Kendall, ‘Genetic Algorithms’, *Search Methodologies: Introductory Tutorials in Optimization and Decision Support Techniques*, ed. by E. K. Burke and G. Kendall, Springer US, 2005 97, ISBN: 978-0-387-28356-2, URL: https://doi.org/10.1007/0-387-28356-0_4.
- [83] G. Choudalakis and D. Casadei, *Plotting the differences between data and expectation*, *Eur. Phys. J. Plus* **127** (2012) 25, arXiv: [1111.2062 \[physics.data-an\]](#).
- [84] ATLAS Collaboration, *Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV*, *Eur. Phys. J. C* **71** (2011) 1577, arXiv: [1012.1792 \[hep-ex\]](#).
- [85] ATLAS Collaboration, *Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s} = 7$ TeV in dilepton final states with ATLAS*, *Phys. Lett. B* **707** (2012) 459, arXiv: [1108.3699 \[hep-ex\]](#).
- [86] ATLAS Collaboration, *ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970, arXiv: [1907.05120 \[hep-ex\]](#).
- [87] M. Bähr et al., *Herwig++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639, arXiv: [0803.0883 \[hep-ph\]](#).
- [88] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196, arXiv: [1512.01178 \[hep-ph\]](#).
- [89] S. Frixione, E. Laenen, P. Motylinski, C. White and B. R. Webber, *Single-top hadroproduction in association with a W boson*, *JHEP* **07** (2008) 029.
- [90] F. Demartin, B. Maier, F. Maltoni, K. Mawatari and M. Zaro, *tWH associated production at the LHC*, *Eur. Phys. J. C* **77** (2017) 34, arXiv: [1607.05862 \[hep-ph\]](#).

- [91] O. Bessidskaia Bylund, ‘Modelling Wt and tWZ production at NLO for ATLAS analyses’, *9th International Workshop on Top Quark Physics*, 2016, arXiv: [1612.00440 \[hep-ph\]](#).
- [92] A. L. Read, *Presentation of search results: the CL_S technique*, *J. Phys. G* **28** (2002) 2693.
- [93] M. Bauer, M. Klassen and V. Tenorth, *Universal properties of pseudoscalar mediators in dark matter extensions of 2HDMs*, *JHEP* **07** (2018) 107, arXiv: [1712.06597 \[hep-ph\]](#).
- [94] U. Haisch and G. Polesello, *Searching for heavy Higgs bosons in the $t\bar{t}Z$ and tbW final states*, *JHEP* **09** (2018) 151, arXiv: [1807.07734 \[hep-ph\]](#).
- [95] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2020-001, URL: <https://cds.cern.ch/record/2717821>.

The ATLAS Collaboration

G. Aad¹⁰², B. Abbott¹²⁸, D.C. Abbott¹⁰³, A. Abed Abud³⁶, K. Abeling⁵³, D.K. Abhayasinghe⁹⁴, S.H. Abidi¹⁶⁷, O.S. AbouZeid⁴⁰, N.L. Abraham¹⁵⁶, H. Abramowicz¹⁶¹, H. Abreu¹⁶⁰, Y. Abulaiti⁶, B.S. Acharya^{67a,67b,o}, B. Achkar⁵³, L. Adam¹⁰⁰, C. Adam Bourdarios⁵, L. Adamczyk^{84a}, L. Adamek¹⁶⁷, J. Adelman¹²¹, A. Adiguzel^{12c}, S. Adorni⁵⁴, T. Adye¹⁴³, A.A. Affolder¹⁴⁵, Y. Afik¹⁶⁰, C. Agapopoulou⁶⁵, M.N. Agaras³⁸, A. Aggarwal¹¹⁹, C. Agheorghiesei^{27c}, J.A. Aguilar-Saavedra^{139f,139a,ac}, A. Ahmad³⁶, F. Ahmadov⁸⁰, W.S. Ahmed¹⁰⁴, X. Ai¹⁸, G. Aielli^{74a,74b}, S. Akatsuka⁸⁶, M. Akbiyik¹⁰⁰, T.P.A. Åkesson⁹⁷, E. Akilli⁵⁴, A.V. Akimov¹¹¹, K. Al Houry⁶⁵, G.L. Alberghi^{23b,23a}, J. Albert¹⁷⁶, M.J. Alconada Verzini¹⁶¹, S. Alderweireldt³⁶, M. Aleksa³⁶, I.N. Aleksandrov⁸⁰, C. Alexa^{27b}, T. Alexopoulos¹⁰, A. Alfonsi¹²⁰, F. Alfonsi^{23b,23a}, M. Alhroob¹²⁸, B. Ali¹⁴¹, S. Ali¹⁵⁸, M. Aliev¹⁶⁶, G. Alimonti^{69a}, C. Allaire³⁶, B.M.M. Allbrooke¹⁵⁶, B.W. Allen¹³¹, P.P. Allport²¹, A. Aloisio^{70a,70b}, F. Alonso⁸⁹, C. Alpigiani¹⁴⁸, E. Alunno Camelia^{74a,74b}, M. Alvarez Estevez⁹⁹, M.G. Alvigi^{70a,70b}, Y. Amaral Coutinho^{81b}, A. Ambler¹⁰⁴, L. Ambroz¹³⁴, C. Amelung³⁶, D. Amidei¹⁰⁶, S.P. Amor Dos Santos^{139a}, S. Amoroso⁴⁶, C.S. Amrouche⁵⁴, F. An⁷⁹, C. Anastopoulos¹⁴⁹, N. Andari¹⁴⁴, T. Andeen¹¹, J.K. Anders²⁰, S.Y. Andreev^{45a,45b}, A. Andreazza^{69a,69b}, V. Andrei^{61a}, C.R. Anelli¹⁷⁶, S. Angelidakis⁹, A. Angerami³⁹, A.V. Anisenkov^{122b,122a}, A. Annovi^{72a}, C. Antel⁵⁴, M.T. Anthony¹⁴⁹, E. Antipov¹²⁹, M. Antonelli⁵¹, D.J.A. Antrim¹⁸, F. Anulli^{73a}, M. Aoki⁸², J.A. Aparisi Pozo¹⁷⁴, M.A. Aparo¹⁵⁶, L. Aperio Bella⁴⁶, N. Aranzabal³⁶, V. Araujo Ferraz^{81a}, R. Araujo Pereira^{81b}, C. Arcangeletti⁵¹, A.T.H. Arce⁴⁹, J-F. Arguin¹¹⁰, S. Argyropoulos⁵², J.-H. Arling⁴⁶, A.J. Armbruster³⁶, A. Armstrong¹⁷¹, O. Arnaez¹⁶⁷, H. Arnold¹²⁰, Z.P. Arrubarrena Tame¹¹⁴, G. Artoni¹³⁴, H. Asada¹¹⁷, K. Asai¹²⁶, S. Asai¹⁶³, T. Asawatavonvanich¹⁶⁵, N. Asbah⁵⁹, E.M. Asimakopoulou¹⁷², L. Asquith¹⁵⁶, J. Assahsah^{35d}, K. Assamagan²⁹, R. Astalos^{28a}, R.J. Atkin^{33a}, M. Atkinson¹⁷³, N.B. Atlay¹⁹, H. Atmani⁶⁵, P.A. Atmasiddha¹⁰⁶, K. Augsten¹⁴¹, V.A. Austrup¹⁸², G. Avolio³⁶, M.K. Ayoub^{15a}, G. Azuelos^{110,aj}, D. Babal^{28a}, H. Bachacou¹⁴⁴, K. Bachas¹⁶², F. Backman^{45a,45b}, P. Bagnaia^{73a,73b}, M. Bahmani⁸⁵, H. Bahrasemani¹⁵², A.J. Bailey¹⁷⁴, V.R. Bailey¹⁷³, J.T. Baines¹⁴³, C. Bakalis¹⁰, O.K. Baker¹⁸³, P.J. Bakker¹²⁰, E. Bakos¹⁶, D. Bakshi Gupta⁸, S. Balaji¹⁵⁷, R. Balasubramanian¹²⁰, E.M. Baldin^{122b,122a}, P. Balek¹⁸⁰, F. Balli¹⁴⁴, W.K. Balunas¹³⁴, J. Balz¹⁰⁰, E. Banas⁸⁵, M. Bandieramonte¹³⁸, A. Bandyopadhyay¹⁹, Sw. Banerjee^{181j}, L. Barak¹⁶¹, W.M. Barbe³⁸, E.L. Barberio¹⁰⁵, D. Barberis^{55b,55a}, M. Barbero¹⁰², G. Barbour⁹⁵, T. Barillari¹¹⁵, M-S. Barisits³⁶, J. Barkeloo¹³¹, T. Barklow¹⁵³, R. Barnea¹⁶⁰, B.M. Barnett¹⁴³, R.M. Barnett¹⁸, Z. Barnovska-Blenessy^{60a}, A. Baroncelli^{60a}, G. Barone²⁹, A.J. Barr¹³⁴, L. Barranco Navarro^{45a,45b}, F. Barreiro⁹⁹, J. Barreiro Guimarães da Costa^{15a}, U. Barron¹⁶¹, S. Barsov¹³⁷, F. Bartels^{61a}, R. Bartoldus¹⁵³, G. Bartolini¹⁰², A.E. Barton⁹⁰, P. Bartos^{28a}, A. Basalae⁴⁶, A. Basan¹⁰⁰, A. Bassalat^{65,ag}, M.J. Basso¹⁶⁷, R.L. Bates⁵⁷, S. Batlamous^{35e}, J.R. Batley³², B. Batool¹⁵¹, M. Battaglia¹⁴⁵, M. Baue^{73a,73b}, F. Bauer¹⁴⁴, P. Bauer²⁴, H.S. Bawa³¹, A. Bayirli^{12c}, J.B. Beacham⁴⁹, T. Beau¹³⁵, P.H. Beauchemin¹⁷⁰, F. Becherer⁵², P. Bechtel²⁴, H.C. Beck⁵³, H.P. Beck^{20,q}, K. Becker¹⁷⁸, C. Becot⁴⁶, A. Beddall^{12d}, A.J. Beddall^{12a}, V.A. Bednyakov⁸⁰, M. Bedognetti¹²⁰, C.P. Bee¹⁵⁵, T.A. Beermann¹⁸², M. Begalli^{81b}, M. Begel²⁹, A. Behera¹⁵⁵, J.K. Behr⁴⁶, F. Beisiegel²⁴, M. Belfkir⁵, A.S. Bell⁹⁵, G. Bella¹⁶¹, L. Bellagamba^{23b}, A. Bellerive³⁴, P. Bellos⁹, K. Beloborodov^{122b,122a}, K. Belotskiy¹¹², N.L. Belyaev¹¹², D. Bencheikroun^{35a}, N. Benekos¹⁰, Y. Benhammou¹⁶¹, D.P. Benjamin⁶, M. Benoit²⁹, J.R. Bensinger²⁶, S. Bentvelsen¹²⁰, L. Beresford¹³⁴, M. Beretta⁵¹, D. Berge¹⁹, E. Bergeaas Kuutmann¹⁷², N. Berger⁵, B. Bergmann¹⁴¹, L.J. Bergsten²⁶, J. Beringer¹⁸, S. Berlendis⁷, G. Bernardi¹³⁵, C. Bernius¹⁵³, F.U. Bernlochner²⁴, T. Berry⁹⁴, P. Berta¹⁰⁰, A. Berthold⁴⁸, I.A. Bertram⁹⁰, O. Bessidskaia Bylund¹⁸², N. Besson¹⁴⁴, S. Bethke¹¹⁵, A. Betti⁴², A.J. Bevan⁹³, J. Beyer¹¹⁵, S. Bhatta¹⁵⁵, D.S. Bhattacharya¹⁷⁷, P. Bhattarai²⁶, V.S. Bhopatkar⁶, R. Bi¹³⁸, R.M. Bianchi¹³⁸, O. Biebel¹¹⁴, D. Biedermann¹⁹, R. Bielski³⁶, K. Bierwagen¹⁰⁰, N.V. Biesuz^{72a,72b}, M. Biglietti^{75a}, T.R.V. Billoud¹⁴¹, M. Bindi⁵³, A. Bingul^{12d},

C. Bini^{73a,73b}, S. Biondi^{23b,23a}, C.J. Birch-sykes¹⁰¹, M. Birman¹⁸⁰, T. Bisanz³⁶, J.P. Biswal³, D. Biswas^{181j}, A. Bitadze¹⁰¹, C. Bittrich⁴⁸, K. Bjørke¹³³, T. Blazek^{28a}, I. Bloch⁴⁶, C. Blocker²⁶, A. Blue⁵⁷, U. Blumenschein⁹³, G.J. Bobbink¹²⁰, V.S. Bobrovnikov^{122b,122a}, S.S. Bocchetta⁹⁷, D. Bogavac¹⁴, A.G. Bogdanchikov^{122b,122a}, C. Bohm^{45a}, V. Boisvert⁹⁴, P. Bokan^{172,53}, T. Bold^{84a}, A.E. Bolz^{61b}, M. Bomben¹³⁵, M. Bona⁹³, J.S. Bonilla¹³¹, M. Boonekamp¹⁴⁴, C.D. Booth⁹⁴, A.G. Borbély⁵⁷, H.M. Borecka-Bielska⁹¹, L.S. Borgna⁹⁵, A. Borisov¹²³, G. Borissov⁹⁰, D. Bortoletto¹³⁴, D. Boscherini^{23b}, M. Bosman¹⁴, J.D. Bossio Sola¹⁰⁴, K. Bouaouda^{35a}, J. Boudreau¹³⁸, E.V. Bouhova-Thacker⁹⁰, D. Boumediene³⁸, A. Boveia¹²⁷, J. Boyd³⁶, D. Boye^{33c}, I.R. Boyko⁸⁰, A.J. Bozson⁹⁴, J. Bracinik²¹, N. Brahim^{60d}, G. Brandt¹⁸², O. Brandt³², F. Braren⁴⁶, B. Brau¹⁰³, J.E. Brau¹³¹, W.D. Breaden Madden⁵⁷, K. Brendlinger⁴⁶, R. Brenner¹⁶⁰, L. Brenner³⁶, R. Brenner¹⁷², S. Bressler¹⁸⁰, B. Brickwedde¹⁰⁰, D.L. Briglin²¹, D. Britton⁵⁷, D. Britzger¹¹⁵, I. Brock²⁴, R. Brock¹⁰⁷, G. Brooijmans³⁹, W.K. Brooks^{146d}, E. Brost²⁹, P.A. Bruckman de Renstrom⁸⁵, B. Brüers⁴⁶, D. Bruncko^{28b}, A. Bruni^{23b}, G. Bruni^{23b}, M. Bruschi^{23b}, N. Brusino^{73a,73b}, L. Bryngemark¹⁵³, T. Buanes¹⁷, Q. Buat¹⁵⁵, P. Buchholz¹⁵¹, A.G. Buckley⁵⁷, I.A. Budagov⁸⁰, M.K. Bugge¹³³, O. Bulekov¹¹², B.A. Bullard⁵⁹, T.J. Burch¹²¹, S. Burdin⁹¹, C.D. Burgard¹²⁰, A.M. Burger¹²⁹, B. Burghgrave⁸, J.T.P. Burr⁴⁶, C.D. Burton¹¹, J.C. Burzynski¹⁰³, V. Büscher¹⁰⁰, E. Buschmann⁵³, P.J. Bussey⁵⁷, J.M. Butler²⁵, C.M. Buttar⁵⁷, J.M. Butterworth⁹⁵, P. Butti³⁶, W. Buttinger¹⁴³, C.J. Buxo Vazquez¹⁰⁷, A. Buzatu¹⁵⁸, A.R. Buzykaev^{122b,122a}, G. Cabras^{23b,23a}, S. Cabrera Urbán¹⁷⁴, D. Caforio⁵⁶, H. Cai¹³⁸, V.M.M. Cairo¹⁵³, O. Cakir^{4a}, N. Calace³⁶, P. Calafiura¹⁸, G. Calderini¹³⁵, P. Calfayan⁶⁶, G. Callea⁵⁷, L.P. Caloba^{81b}, A. Caltabiano^{74a,74b}, S. Calvente Lopez⁹⁹, D. Calvet³⁸, S. Calvet³⁸, T.P. Calvet¹⁰², M. Calvetti^{72a,72b}, R. Camacho Toro¹³⁵, S. Camarda³⁶, D. Camarero Munoz⁹⁹, P. Camarri^{74a,74b}, M.T. Camerlingo^{75a,75b}, D. Cameron¹³³, C. Camincher³⁶, S. Campana³⁶, M. Campanelli⁹⁵, A. Camplani⁴⁰, V. Canale^{70a,70b}, A. Canesse¹⁰⁴, M. Cano Bret⁷⁸, J. Cantero¹²⁹, T. Cao¹⁶¹, Y. Cao¹⁷³, M. Capua^{41b,41a}, R. Cardarelli^{74a}, F. Cardillo¹⁷⁴, G. Carducci^{41b,41a}, I. Carli¹⁴², T. Carli³⁶, G. Carlino^{70a}, B.T. Carlson¹³⁸, E.M. Carlson^{176,168a}, L. Carminati^{69a,69b}, R.M.D. Carney¹⁵³, S. Caron¹¹⁹, E. Carquin^{146d}, S. Carrá⁴⁶, G. Carratta^{23b,23a}, J.W.S. Carter¹⁶⁷, T.M. Carter⁵⁰, M.P. Casado^{14g}, A.F. Casha¹⁶⁷, E.G. Castiglia¹⁸³, F.L. Castillo¹⁷⁴, L. Castillo Garcia¹⁴, V. Castillo Gimenez¹⁷⁴, N.F. Castro^{139a,139e}, A. Catinaccio³⁶, J.R. Catmore¹³³, A. Cattai³⁶, V. Cavaliere²⁹, V. Cavasinni^{72a,72b}, E. Celebi^{12b}, F. Celli¹³⁴, K. Cerny¹³⁰, A.S. Cerqueira^{81a}, A. Cerri¹⁵⁶, L. Cerrito^{74a,74b}, F. Cerutti¹⁸, A. Cervelli^{23b,23a}, S.A. Cetin^{12b}, Z. Chadi^{35a}, D. Chakraborty¹²¹, J. Chan¹⁸¹, W.S. Chan¹²⁰, W.Y. Chan⁹¹, J.D. Chapman³², B. Chargeishvili^{159b}, D.G. Charlton²¹, T.P. Charman⁹³, M. Chatterjee²⁰, C.C. Chau³⁴, S. Che¹²⁷, S. Chekanov⁶, S.V. Chekulaev^{168a}, G.A. Chelkov^{80,ae}, B. Chen⁷⁹, C. Chen^{60a}, C.H. Chen⁷⁹, H. Chen^{15c}, H. Chen²⁹, J. Chen^{60a}, J. Chen³⁹, J. Chen²⁶, S. Chen¹³⁶, S.J. Chen^{15c}, X. Chen^{15b}, Y. Chen^{60a}, Y-H. Chen⁴⁶, H.C. Cheng^{63a}, H.J. Cheng^{15a}, A. Cheplakov⁸⁰, E. Cheremushkina¹²³, R. Cherkaoui El Moursli^{35e}, E. Cheu⁷, K. Cheung⁶⁴, T.J.A. Chevalérias¹⁴⁴, L. Chevalier¹⁴⁴, V. Chiarella⁵¹, G. Chiarelli^{72a}, G. Chiodini^{68a}, A.S. Chisholm²¹, A. Chitan^{27b}, I. Chiu¹⁶³, Y.H. Chiu¹⁷⁶, M.V. Chizhov⁸⁰, K. Choi¹¹, A.R. Chomont^{73a,73b}, Y. Chou¹⁰³, Y.S. Chow¹²⁰, L.D. Christopher^{33e}, M.C. Chu^{63a}, X. Chu^{15a,15d}, J. Chudoba¹⁴⁰, J.J. Chwastowski⁸⁵, L. Chytka¹³⁰, D. Cieri¹¹⁵, K.M. Ciesla⁸⁵, V. Cindro⁹², I.A. Cioară^{27b}, A. Ciocio¹⁸, F. Ciroto^{70a,70b}, Z.H. Citron^{180,k}, M. Citterio^{69a}, D.A. Ciubotaru^{27b}, B.M. Ciungu¹⁶⁷, A. Clark⁵⁴, P.J. Clark⁵⁰, S.E. Clawson¹⁰¹, C. Clement^{45a,45b}, L. Clissa^{23b,23a}, Y. Coadou¹⁰², M. Cobal^{67a,67c}, A. Coccaro^{55b}, J. Cochran⁷⁹, R. Coelho Lopes De Sa¹⁰³, H. Cohen¹⁶¹, A.E.C. Coimbra³⁶, B. Cole³⁹, A.P. Colijn¹²⁰, J. Collot⁵⁸, P. Conde Muiño^{139a,139h}, S.H. Connell^{33c}, I.A. Connelly⁵⁷, S. Constantinescu^{27b}, F. Conventi^{70a,ak}, A.M. Cooper-Sarkar¹³⁴, F. Cormier¹⁷⁵, K.J.R. Cormier¹⁶⁷, L.D. Corpe⁹⁵, M. Corradi^{73a,73b}, E.E. Corrigan⁹⁷, F. Corriveau^{104,aa}, M.J. Costa¹⁷⁴, F. Costanza⁵, D. Costanzo¹⁴⁹, G. Cowan⁹⁴, J.W. Cowley³², J. Crane¹⁰¹, K. Cranmer¹²⁵, R.A. Creager¹³⁶, S. Crépe-Renaudin⁵⁸, F. Crescioli¹³⁵, M. Cristinziani²⁴, V. Croft¹⁷⁰, G. Crosetti^{41b,41a}, A. Cueto⁵, T. Cuhadar Donszelmann¹⁷¹, H. Cui^{15a,15d}, A.R. Cukierman¹⁵³, W.R. Cunningham⁵⁷, S. Czekierda⁸⁵,

P. Czodrowski³⁶, M.M. Czurylo^{61b}, M.J. Da Cunha Sargedas De Sousa^{60b}, J.V. Da Fonseca Pinto^{81b},
 C. Da Via¹⁰¹, W. Dabrowski^{84a}, F. Dachs³⁶, T. Dado⁴⁷, S. Dahbi^{33e}, T. Dai¹⁰⁶, C. Dallapiccola¹⁰³,
 M. Dam⁴⁰, G. D'amen²⁹, V. D'Amico^{75a,75b}, J. Damp¹⁰⁰, J.R. Dandoy¹³⁶, M.F. Daneri³⁰, M. Danning¹⁵²,
 V. Dao³⁶, G. Darbo^{55b}, O. Dartsis⁵, A. Dattagupta¹³¹, T. Daubney⁴⁶, S. D'Auria^{69a,69b}, C. David^{168b},
 T. Davidek¹⁴², D.R. Davis⁴⁹, I. Dawson¹⁴⁹, K. De⁸, R. De Asmundis^{70a}, M. De Beurs¹²⁰,
 S. De Castro^{23b,23a}, N. De Groot¹¹⁹, P. de Jong¹²⁰, H. De la Torre¹⁰⁷, A. De Maria^{15c}, D. De Pedis^{73a},
 A. De Salvo^{73a}, U. De Sanctis^{74a,74b}, M. De Santis^{74a,74b}, A. De Santo¹⁵⁶, J.B. De Vivie De Regie⁶⁵,
 D.V. Dedovich⁸⁰, A.M. Deiana⁴², J. Del Peso⁹⁹, Y. Delabat Diaz⁴⁶, D. Delgove⁶⁵, F. Deliot¹⁴⁴,
 C.M. Delitzsch⁷, M. Della Pietra^{70a,70b}, D. Della Volpe⁵⁴, A. Dell'Acqua³⁶, L. Dell'Asta^{74a,74b},
 M. Delmastro⁵, C. Delporte⁶⁵, P.A. Delsart⁵⁸, S. Demers¹⁸³, M. Demichev⁸⁰, G. Demontigny¹¹⁰,
 S.P. Denisov¹²³, L. D'Eramo¹²¹, D. Derendarz⁸⁵, J.E. Derkaoui^{35d}, F. Derue¹³⁵, P. Dervan⁹¹, K. Desch²⁴,
 K. Dette¹⁶⁷, C. Deutsch²⁴, M.R. Devesa³⁰, P.O. Deviveiros³⁶, F.A. Di Bello^{73a,73b}, A. Di Ciaccio^{74a,74b},
 L. Di Ciaccio⁵, C. Di Donato^{70a,70b}, A. Di Girolamo³⁶, G. Di Gregorio^{72a,72b}, A. Di Luca^{76a,76b},
 B. Di Micco^{75a,75b}, R. Di Nardo^{75a,75b}, K.F. Di Petrillo⁵⁹, R. Di Sipio¹⁶⁷, C. Diaconu¹⁰², F.A. Dias¹²⁰,
 T. Dias Do Vale^{139a}, M.A. Diaz^{146a}, F.G. Diaz Capriles²⁴, J. Dickinson¹⁸, M. Didenko¹⁶⁶, E.B. Diehl¹⁰⁶,
 J. Dietrich¹⁹, S. Díez Cornell⁴⁶, C. Diez Pardos¹⁵¹, A. Dimitrievska¹⁸, W. Ding^{15b}, J. Dingfelder²⁴,
 S.J. Dittmeier^{61b}, F. Dittus³⁶, F. Djama¹⁰², T. Djobava^{159b}, J.I. Djuvslund¹⁷, M.A.B. Do Vale¹⁴⁷,
 M. Dobre^{27b}, D. Dodsworth²⁶, C. Doglioni⁹⁷, J. Dolejsi¹⁴², Z. Dolezal¹⁴², M. Donadelli^{81c}, B. Dong^{60c},
 J. Donini³⁸, A. D'onofrio^{15c}, M. D'Onofrio⁹¹, J. Dopke¹⁴³, A. Doria^{70a}, M.T. Dova⁸⁹, A.T. Doyle⁵⁷,
 E. Drechsler¹⁵², E. Dreyer¹⁵², T. Dreyer⁵³, A.S. Drobac¹⁷⁰, D. Du^{60b}, T.A. du Pree¹²⁰, Y. Duan^{60d},
 F. Dubinin¹¹¹, M. Dubovsky^{28a}, A. Dubreuil⁵⁴, E. Duchovni¹⁸⁰, G. Duckeck¹¹⁴, O.A. Ducu³⁶, D. Duda¹¹⁵,
 A. Dudarev³⁶, A.C. Dudder¹⁰⁰, E.M. Duffield¹⁸, M. D'uffizi¹⁰¹, L. Duflo⁶⁵, M. Dührssen³⁶, C. Dülsen¹⁸²,
 M. Dumancic¹⁸⁰, A.E. Dumitriu^{27b}, M. Dunford^{61a}, S. Dungs⁴⁷, A. Duperrin¹⁰², H. Duran Yildiz^{4a},
 M. Düren⁵⁶, A. Durglishvili^{159b}, D. Duschinger⁴⁸, B. Dutta⁴⁶, D. Duvnjak¹, G.I. Dyckes¹³⁶, M. Dyndal³⁶,
 S. Dysch¹⁰¹, B.S. Dziedzic⁸⁵, M.G. Eggleston⁴⁹, T. Eifert⁸, G. Eigen¹⁷, K. Einsweiler¹⁸, T. Ekelof¹⁷²,
 H. El Jarrari^{35e}, V. Ellajosyula¹⁷², M. Ellert¹⁷², F. Ellinghaus¹⁸², A.A. Elliot⁹³, N. Ellis³⁶, J. Elmsheuser²⁹,
 M. Elsing³⁶, D. Emelianov¹⁴³, A. Emerman³⁹, Y. Enari¹⁶³, M.B. Epland⁴⁹, J. Erdmann⁴⁷, A. Ereditato²⁰,
 P.A. Erland⁸⁵, M. Errenst¹⁸², M. Escalier⁶⁵, C. Escobar¹⁷⁴, O. Estrada Pastor¹⁷⁴, E. Etzion¹⁶¹,
 G.E. Evans^{139a}, H. Evans⁶⁶, M.O. Evans¹⁵⁶, A. Ezhilov¹³⁷, F. Fabbri⁵⁷, L. Fabbri^{23b,23a}, V. Fabiani¹¹⁹,
 G. Facini¹⁷⁸, R.M. Fakhruddinov¹²³, S. Falciano^{73a}, P.J. Falke²⁴, S. Falke³⁶, J. Faltova¹⁴², Y. Fang^{15a},
 Y. Fang^{15a}, G. Fanourakis⁴⁴, M. Fanti^{69a,69b}, M. Faraj^{67a,67c}, A. Farbin⁸, A. Farilla^{75a}, E.M. Farina^{71a,71b},
 T. Farooque¹⁰⁷, S.M. Farrington⁵⁰, P. Farthouat³⁶, F. Fassi^{35e}, P. Fassnacht³⁶, D. Fassouliotis⁹,
 M. Faucci Giannelli⁵⁰, W.J. Fawcett³², L. Fayard⁶⁵, O.L. Fedin^{137,p}, W. Fedorko¹⁷⁵, A. Fehr²⁰,
 M. Feickert¹⁷³, L. Feligioni¹⁰², A. Fell¹⁴⁹, C. Feng^{60b}, M. Feng⁴⁹, M.J. Fenton¹⁷¹, A.B. Fenyuk¹²³,
 S.W. Ferguson⁴³, J. Ferrando⁴⁶, A. Ferrari¹⁷², P. Ferrari¹²⁰, R. Ferrari^{71a}, D.E. Ferreira de Lima^{61b},
 A. Ferrer¹⁷⁴, D. Ferrere⁵⁴, C. Ferretti¹⁰⁶, F. Fiedler¹⁰⁰, A. Filipčić⁹², F. Filthaut¹¹⁹, K.D. Finelli²⁵,
 M.C.N. Fiolhais^{139a,139c,a}, L. Fiorini¹⁷⁴, F. Fischer¹¹⁴, J. Fischer¹⁰⁰, W.C. Fisher¹⁰⁷, T. Fitschen²¹,
 I. Fleck¹⁵¹, P. Fleischmann¹⁰⁶, T. Flick¹⁸², B.M. Flierl¹¹⁴, L. Flores¹³⁶, L.R. Flores Castillo^{63a},
 F.M. Follega^{76a,76b}, N. Fomin¹⁷, J.H. Foo¹⁶⁷, G.T. Forcolin^{76a,76b}, B.C. Forland⁶⁶, A. Formica¹⁴⁴,
 F.A. Förster¹⁴, A.C. Forti¹⁰¹, E. Fortin¹⁰², M.G. Foti¹³⁴, D. Fournier⁶⁵, H. Fox⁹⁰, P. Francavilla^{72a,72b},
 S. Francescato^{73a,73b}, M. Franchini^{23b,23a}, S. Franchino^{61a}, D. Francis³⁶, L. Franco⁵, L. Franconi²⁰,
 M. Franklin⁵⁹, G. Frattari^{73a,73b}, A.N. Fray⁹³, P.M. Freeman²¹, B. Freund¹¹⁰, W.S. Freund^{81b},
 E.M. Freundlich⁴⁷, D.C. Frizzell¹²⁸, D. Froidevaux³⁶, J.A. Frost¹³⁴, M. Fujimoto¹²⁶, C. Fukunaga¹⁶⁴,
 E. Fullana Torregrosa¹⁷⁴, T. Fusayasu¹¹⁶, J. Fuster¹⁷⁴, A. Gabrielli^{23b,23a}, A. Gabrielli³⁶, S. Gadatsch⁵⁴,
 P. Gadow¹¹⁵, G. Gagliardi^{55b,55a}, L.G. Gagnon¹¹⁰, G.E. Gallardo¹³⁴, E.J. Gallas¹³⁴, B.J. Gallop¹⁴³,
 R. Gamboa Goni⁹³, K.K. Gan¹²⁷, S. Ganguly¹⁸⁰, J. Gao^{60a}, Y. Gao⁵⁰, Y.S. Gao^{31,m}, F.M. Garay Walls^{146a},
 C. García¹⁷⁴, J.E. García Navarro¹⁷⁴, J.A. García Pascual^{15a}, C. Garcia-Argos⁵², M. Garcia-Sciveres¹⁸,

R.W. Gardner³⁷, N. Garelli¹⁵³, S. Gargiulo⁵², C.A. Garner¹⁶⁷, V. Garonne¹³³, S.J. Gasiorowski¹⁴⁸, P. Gaspar^{81b}, A. Gaudiello^{55b,55a}, G. Gaudio^{71a}, P. Gauzzi^{73a,73b}, I.L. Gavrilenko¹¹¹, A. Gavriluk¹²⁴, C. Gay¹⁷⁵, G. Gaycken⁴⁶, E.N. Gazis¹⁰, A.A. Geanta^{27b}, C.M. Gee¹⁴⁵, C.N.P. Gee¹⁴³, J. Geisen⁹⁷, M. Geisen¹⁰⁰, C. Gemme^{55b}, M.H. Genest⁵⁸, C. Geng¹⁰⁶, S. Gentile^{73a,73b}, S. George⁹⁴, T. Geralis⁴⁴, L.O. Gerlach⁵³, P. Gessinger-Befurt¹⁰⁰, G. Gessner⁴⁷, M. Ghasemi Bostanabad¹⁷⁶, M. Ghneimat¹⁵¹, A. Ghosh⁶⁵, A. Ghosh⁷⁸, B. Giacobbe^{23b}, S. Giagu^{73a,73b}, N. Giangiacomi¹⁶⁷, P. Giannetti^{72a}, A. Giannini^{70a,70b}, G. Giannini¹⁴, S.M. Gibson⁹⁴, M. Gignac¹⁴⁵, D.T. Gil^{84b}, B.J. Gilbert³⁹, D. Gillberg³⁴, G. Gilles¹⁸², N.E.K. Gillwald⁴⁶, D.M. Gingrich^{3,aj}, M.P. Giordani^{67a,67c}, P.F. Giraud¹⁴⁴, G. Giugliarelli^{67a,67c}, D. Giugni^{69a}, F. Giuli^{74a,74b}, S. Gkaitatzis¹⁶², I. Gkialas^{9,h}, E.L. Gkoukousis¹⁴, P. Gkoutoumis¹⁰, L.K. Gladilin¹¹³, C. Glasman⁹⁹, J. Glatzer¹⁴, P.C.F. Glaysher⁴⁶, A. Glazov⁴⁶, G.R. Gledhill¹³¹, I. Gnesi^{41b,c}, M. Goblirsch-Kolb²⁶, D. Godin¹¹⁰, S. Goldfarb¹⁰⁵, T. Golling⁵⁴, D. Golubkov¹²³, A. Gomes^{139a,139b}, R. Goncalves Gama⁵³, R. Gonçalo^{139a,139c}, G. Gonella¹³¹, L. Gonella²¹, A. Gongadze⁸⁰, F. Gonnella²¹, J.L. Gonski³⁹, S. González de la Hoz¹⁷⁴, S. Gonzalez Fernandez¹⁴, R. Gonzalez Lopez⁹¹, C. Gonzalez Renteria¹⁸, R. Gonzalez Suarez¹⁷², S. Gonzalez-Sevilla⁵⁴, G.R. Gonzalvo Rodriguez¹⁷⁴, L. Goossens³⁶, N.A. Gorasia²¹, P.A. Gorbounov¹²⁴, H.A. Gordon²⁹, B. Gorini³⁶, E. Gorini^{68a,68b}, A. Gorišek⁹², A.T. Goshaw⁴⁹, M.I. Gostkin⁸⁰, C.A. Gottardo¹¹⁹, M. Gouighri^{35b}, A.G. Goussiou¹⁴⁸, N. Govender^{33c}, C. Goy⁵, I. Grabowska-Bold^{84a}, E.C. Graham⁹¹, J. Gramling¹⁷¹, E. Gramstad¹³³, S. Grancagnolo¹⁹, M. Grandi¹⁵⁶, V. Gratchev¹³⁷, P.M. Gravila^{27f}, F.G. Gravili^{68a,68b}, C. Gray⁵⁷, H.M. Gray¹⁸, C. Grefe²⁴, K. Gregersen⁹⁷, I.M. Gregor⁴⁶, P. Grenier¹⁵³, K. Grevtsov⁴⁶, C. Grieco¹⁴, N.A. Grieser¹²⁸, A.A. Grillo¹⁴⁵, K. Grimm^{31,l}, S. Grinstein^{14,w}, J.-F. Grivaz⁶⁵, S. Groh¹⁰⁰, E. Gross¹⁸⁰, J. Grosse-Knetter⁵³, Z.J. Grout⁹⁵, C. Grud¹⁰⁶, A. Grummer¹¹⁸, J.C. Grundy¹³⁴, L. Guan¹⁰⁶, W. Guan¹⁸¹, C. Gubbels¹⁷⁵, J. Guenther⁷⁷, A. Guerguichon⁶⁵, J.G.R. Guerrero Rojas¹⁷⁴, F. Guescini¹¹⁵, D. Guest⁷⁷, R. Gugel¹⁰⁰, A. Guida⁴⁶, T. Guillemin⁵, S. Guindon³⁶, J. Guo^{60c}, W. Guo¹⁰⁶, Y. Guo^{60a}, Z. Guo¹⁰², R. Gupta⁴⁶, S. Gurbuz^{12c}, G. Gustavino¹²⁸, M. Guth⁵², P. Gutierrez¹²⁸, C. Gutschow⁹⁵, C. Guyot¹⁴⁴, C. Gwenlan¹³⁴, C.B. Gwilliam⁹¹, E.S. Haaland¹³³, A. Haas¹²⁵, C. Haber¹⁸, H.K. Hadavand⁸, A. Hadei¹⁰⁰, M. Haleem¹⁷⁷, J. Haley¹²⁹, J.J. Hall¹⁴⁹, G. Halladjian¹⁰⁷, G.D. Hallewell¹⁰², K. Hamano¹⁷⁶, H. Hamdaoui^{35e}, M. Hamer²⁴, G.N. Hamity⁵⁰, K. Han^{60a}, L. Han^{15c}, L. Han^{60a}, S. Han¹⁸, Y.F. Han¹⁶⁷, K. Hanagaki^{82,u}, M. Hance¹⁴⁵, D.M. Handl¹¹⁴, M.D. Hank³⁷, R. Hankache¹³⁵, E. Hansen⁹⁷, J.B. Hansen⁴⁰, J.D. Hansen⁴⁰, M.C. Hansen²⁴, P.H. Hansen⁴⁰, E.C. Hanson¹⁰¹, K. Hara¹⁶⁹, T. Harenberg¹⁸², S. Harkusha¹⁰⁸, P.F. Harrison¹⁷⁸, N.M. Hartman¹⁵³, N.M. Hartmann¹¹⁴, Y. Hasegawa¹⁵⁰, A. Hasib⁵⁰, S. Hassani¹⁴⁴, S. Haug²⁰, R. Hauser¹⁰⁷, M. Havranek¹⁴¹, C.M. Hawkes²¹, R.J. Hawkings³⁶, S. Hayashida¹¹⁷, D. Hayden¹⁰⁷, C. Hayes¹⁰⁶, R.L. Hayes¹⁷⁵, C.P. Hays¹³⁴, J.M. Hays⁹³, H.S. Hayward⁹¹, S.J. Haywood¹⁴³, F. He^{60a}, Y. He¹⁶⁵, M.P. Heath⁵⁰, V. Hedberg⁹⁷, A.L. Heggelund¹³³, N.D. Hehir⁹³, C. Heidegger⁵², K.K. Heidegger⁵², W.D. Heidorn⁷⁹, J. Heilman³⁴, S. Heim⁴⁶, T. Heim¹⁸, B. Heinemann^{46,ah}, J.G. Heinlein¹³⁶, J.J. Heinrich¹³¹, L. Heinrich³⁶, J. Hejbal¹⁴⁰, L. Helary⁴⁶, A. Held¹²⁵, S. Hellesund¹³³, C.M. Helling¹⁴⁵, S. Hellman^{45a,45b}, C. Helsens³⁶, R.C.W. Henderson⁹⁰, L. Henkelmann³², A.M. Henriques Correia³⁶, H. Herde²⁶, Y. Hernández Jiménez^{33e}, H. Herr¹⁰⁰, M.G. Herrmann¹¹⁴, T. Herrmann⁴⁸, G. Herten⁵², R. Hertenberger¹¹⁴, L. Hervas³⁶, G.G. Hesketh⁹⁵, N.P. Hessey^{168a}, H. Hibi⁸³, S. Higashino⁸², E. Higón-Rodríguez¹⁷⁴, K. Hildebrand³⁷, J.C. Hill³², K.K. Hill²⁹, K.H. Hiller⁴⁶, S.J. Hillier²¹, M. Hils⁴⁸, I. Hinchliffe¹⁸, F. Hinterkeuser²⁴, M. Hirose¹³², S. Hirose¹⁶⁹, D. Hirschbuehl¹⁸², B. Hiti⁹², O. Hladik¹⁴⁰, J. Hobbs¹⁵⁵, R. Hobincu^{27e}, N. Hod¹⁸⁰, M.C. Hodgkinson¹⁴⁹, A. Hoecker³⁶, D. Hohn⁵², D. Hohov⁶⁵, T. Holm²⁴, T.R. Holmes³⁷, M. Holzbock¹¹⁵, L.B.A.H. Hommels³², T.M. Hong¹³⁸, J.C. Honig⁵², A. Hönle¹¹⁵, B.H. Hooberman¹⁷³, W.H. Hopkins⁶, Y. Horii¹¹⁷, P. Horn⁴⁸, L.A. Horyn³⁷, S. Hou¹⁵⁸, A. Hoummada^{35a}, J. Howarth⁵⁷, J. Hoya⁸⁹, M. Hrabovsky¹³⁰, J. Hrivnac⁶⁵, A. Hrynevich¹⁰⁹, T. Hryn'ova⁵, P.J. Hsu⁶⁴, S.-C. Hsu¹⁴⁸, Q. Hu³⁹, S. Hu^{60c}, Y.F. Hu^{15a,15d,al}, D.P. Huang⁹⁵, X. Huang^{15c}, Y. Huang^{60a}, Y. Huang^{15a}, Z. Hubacek¹⁴¹, F. Hubaut¹⁰², M. Huebner²⁴, F. Huegging²⁴, T.B. Huffman¹³⁴,

M. Huhtinen³⁶, R. Hulsken⁵⁸, R.F.H. Hunter³⁴, N. Huseynov^{80,ab}, J. Huston¹⁰⁷, J. Huth⁵⁹, R. Hyneman¹⁵³,
S. Hyrych^{28a}, G. Iacobucci⁵⁴, G. Iakovidis²⁹, I. Ibragimov¹⁵¹, L. Iconomidou-Fayard⁶⁵, P. Iengo³⁶,
R. Ignazzi⁴⁰, R. Iguchi¹⁶³, T. Iizawa⁵⁴, Y. Ikegami⁸², M. Ikeno⁸², N. Ilic^{119,167,aa}, F. Iltzsche⁴⁸, H. Imam^{35a},
G. Introzzi^{71a,71b}, M. Iodice^{75a}, K. Iordanidou^{168a}, V. Ippolito^{73a,73b}, M.F. Isacson¹⁷², M. Ishino¹⁶³,
W. Islam¹²⁹, C. Issever^{19,46}, S. Istin¹⁶⁰, J.M. Iturbe Ponce^{63a}, R. Iuppa^{76a,76b}, A. Ivina¹⁸⁰, J.M. Izen⁴³,
V. Izzo^{70a}, P. Jacka¹⁴⁰, P. Jackson¹, R.M. Jacobs⁴⁶, B.P. Jaeger¹⁵², V. Jain², G. Jäkel¹⁸², K.B. Jakobi¹⁰⁰,
K. Jakobs⁵², T. Jakoubek¹⁸⁰, J. Jamieson⁵⁷, K.W. Janas^{84a}, R. Jansky⁵⁴, M. Janus⁵³, P.A. Janus^{84a},
G. Jarlskog⁹⁷, A.E. Jaspán⁹¹, N. Javadov^{80,ab}, T. Javůrek³⁶, M. Javurkova¹⁰³, F. Jeanneau¹⁴⁴, L. Jeanty¹³¹,
J. Jejelava^{159a}, P. Jenni^{52,d}, N. Jeong⁴⁶, S. Jézéquel⁵, J. Jia¹⁵⁵, Z. Jia^{15c}, H. Jiang⁷⁹, Y. Jiang^{60a}, Z. Jiang¹⁵³,
S. Jiggins⁵², F.A. Jimenez Morales³⁸, J. Jimenez Pena¹¹⁵, S. Jin^{15c}, A. Jinaru^{27b}, O. Jinnouchi¹⁶⁵,
H. Jivan^{33e}, P. Johansson¹⁴⁹, K.A. Johns⁷, C.A. Johnson⁶⁶, E. Jones¹⁷⁸, R.W.L. Jones⁹⁰, S.D. Jones¹⁵⁶,
T.J. Jones⁹¹, J. Jovicevic³⁶, X. Ju¹⁸, J.J. Junggeburch¹¹⁵, A. Juste Rozas^{14,w}, A. Kaczmarska⁸⁵,
M. Kado^{73a,73b}, H. Kagan¹²⁷, M. Kagan¹⁵³, A. Kahn³⁹, C. Kahra¹⁰⁰, T. Kaji¹⁷⁹, E. Kajomovitz¹⁶⁰,
C.W. Kalderon²⁹, A. Kaluza¹⁰⁰, A. Kamenshchikov¹²³, M. Kaneda¹⁶³, N.J. Kang¹⁴⁵, S. Kang⁷⁹,
Y. Kano¹¹⁷, J. Kanzaki⁸², L.S. Kaplan¹⁸¹, D. Kar^{33e}, K. Karava¹³⁴, M.J. Kareem^{168b}, I. Karkanas¹⁶²,
S.N. Karpov⁸⁰, Z.M. Karpova⁸⁰, V. Kartvelishvili⁹⁰, A.N. Karyukhin¹²³, E. Kasimi¹⁶², A. Kastanas^{45a,45b},
C. Kato^{60d}, J. Katzy⁴⁶, K. Kawade¹⁵⁰, K. Kawagoe⁸⁸, T. Kawaguchi¹¹⁷, T. Kawamoto¹⁴⁴, G. Kawamura⁵³,
E.F. Kay¹⁷⁶, F.I. Kaya¹⁷⁰, S. Kazakos¹⁴, V.F. Kazanin^{122b,122a}, J.M. Keaveney^{33a}, R. Keeler¹⁷⁶,
J.S. Keller³⁴, E. Kellermann⁹⁷, D. Kelsey¹⁵⁶, J.J. Kempster²¹, J. Kendrick²¹, K.E. Kennedy³⁹, O. Kepka¹⁴⁰,
S. Kersten¹⁸², B.P. Kerševan⁹², S. Ketabchi Haghghat¹⁶⁷, F. Khalil-Zada¹³, M. Khandoga¹⁴⁴,
A. Khanov¹²⁹, A.G. Kharlamov^{122b,122a}, T. Kharlamova^{122b,122a}, E.E. Khoda¹⁷⁵, T.J. Khoo⁷⁷,
G. Khoriauli¹⁷⁷, E. Khramov⁸⁰, J. Khubua^{159b}, S. Kido⁸³, M. Kiehn³⁶, E. Kim¹⁶⁵, Y.K. Kim³⁷,
N. Kimura⁹⁵, A. Kirchhoff⁵³, D. Kirchmeier⁴⁸, J. Kirk¹⁴³, A.E. Kiryunin¹¹⁵, T. Kishimoto¹⁶³,
D.P. Kisliuk¹⁶⁷, V. Kitali⁴⁶, C. Kitsaki¹⁰, O. Kivernyk²⁴, T. Klapdor-Kleingrothaus⁵², M. Klassen^{61a},
C. Klein³⁴, M.H. Klein¹⁰⁶, M. Klein⁹¹, U. Klein⁹¹, K. Kleinknecht¹⁰⁰, P. Klimek³⁶, A. Klimentov²⁹,
F. Klimpel³⁶, T. Klingl²⁴, T. Klioutchnikova³⁶, F.F. Klitzner¹¹⁴, P. Kluit¹²⁰, S. Kluth¹¹⁵, E. Kneringer⁷⁷,
E.B.F.G. Knoop¹⁰², A. Knue⁵², D. Kobayashi⁸⁸, M. Kobel⁴⁸, M. Kocian¹⁵³, T. Kodama¹⁶³, P. Kodys¹⁴²,
D.M. Koeck¹⁵⁶, P.T. Koenig²⁴, T. Koffas³⁴, N.M. Köhler³⁶, M. Kolb¹⁴⁴, I. Koletsou⁵, T. Komarek¹³⁰,
T. Kondo⁸², K. Köneke⁵², A.X.Y. Kong¹, A.C. König¹¹⁹, T. Kono¹²⁶, V. Konstantinides⁹⁵,
N. Konstantinidis⁹⁵, B. Konya⁹⁷, R. Kopeliansky⁶⁶, S. Koperny^{84a}, K. Korcyl⁸⁵, K. Kordas¹⁶²,
G. Koren¹⁶¹, A. Korn⁹⁵, I. Korolkov¹⁴, E.V. Korolkova¹⁴⁹, N. Korotkova¹¹³, O. Kortner¹¹⁵, S. Kortner¹¹⁵,
V.V. Kostyukhin^{149,166}, A. Kotsokchagia⁶⁵, A. Kotwal⁴⁹, A. Koulouris¹⁰,
A. Kourkoumeli-Charalampidi^{71a,71b}, C. Kourkoumelis⁹, E. Kourlitis⁶, V. Kouskoura²⁹, R. Kowalewski¹⁷⁶,
W. Kozanecki¹⁰¹, A.S. Kozhin¹²³, V.A. Kramarenko¹¹³, G. Kramberger⁹², D. Krasnopevtsev^{60a},
M.W. Krasny¹³⁵, A. Krasznahorkay³⁶, D. Krauss¹¹⁵, J.A. Kremer¹⁰⁰, J. Kretzschmar⁹¹, K. Kreul¹⁹,
P. Krieger¹⁶⁷, F. Krieter¹¹⁴, S. Krishnamurthy¹⁰³, A. Krishnan^{61b}, M. Krivos¹⁴², K. Krizka¹⁸,
K. Kroeninger⁴⁷, H. Kroha¹¹⁵, J. Kroll¹⁴⁰, J. Kroll¹³⁶, K.S. Krowpman¹⁰⁷, U. Kruchonak⁸⁰, H. Krüger²⁴,
N. Krumnack⁷⁹, M.C. Kruse⁴⁹, J.A. Krzysiak⁸⁵, A. Kubota¹⁶⁵, O. Kuchinskaia¹⁶⁶, S. Kuday^{4b},
D. Kuechler⁴⁶, J.T. Kuechler⁴⁶, S. Kuehn³⁶, T. Kuhl⁴⁶, V. Kukhtin⁸⁰, Y. Kulchitsky^{108,ad}, S. Kuleshov^{146b},
Y.P. Kulinich¹⁷³, M. Kuna⁵⁸, A. Kupco¹⁴⁰, T. Kupfer⁴⁷, O. Kuprash⁵², H. Kurashige⁸³,
L.L. Kurchaninov^{168a}, Y.A. Kurochkin¹⁰⁸, A. Kurova¹¹², M.G. Kurth^{15a,15d}, E.S. Kuwertz³⁶, M. Kuze¹⁶⁵,
A.K. Kvam¹⁴⁸, J. Kvita¹³⁰, T. Kwan¹⁰⁴, C. Lacasta¹⁷⁴, F. Lacava^{73a,73b}, D.P.J. Lack¹⁰¹, H. Lacker¹⁹,
D. Lacour¹³⁵, E. Ladygin⁸⁰, R. Lafaye⁵, B. Laforge¹³⁵, T. Lagouri^{146c}, S. Lai⁵³, I.K. Lakomic^{84a},
J.E. Lambert¹²⁸, S. Lammers⁶⁶, W. Lampl⁷, C. Lampoudis¹⁶², E. Lançon²⁹, U. Landgraf⁵²,
M.P.J. Landon⁹³, V.S. Lang⁵², J.C. Lange⁵³, R.J. Langenberg¹⁰³, A.J. Lankford¹⁷¹, F. Lanni²⁹,
K. Lantzsche²⁴, A. Lanza^{71a}, A. Lapertosa^{55b,55a}, J.F. Laporte¹⁴⁴, T. Lari^{69a}, F. Lasagni Manghi^{23b,23a},
M. Lassnig³⁶, V. Latonova¹⁴⁰, T.S. Lau^{63a}, A. Laudrain¹⁰⁰, A. Laurier³⁴, M. Lavorgna^{70a,70b},

S.D. Lawlor⁹⁴, M. Lazzaroni^{69a,69b}, B. Le¹⁰¹, E. Le Guirriec¹⁰², A. Lebedev⁷⁹, M. LeBlanc⁷, T. LeCompte⁶, F. Ledroit-Guillon⁵⁸, A.C.A. Lee⁹⁵, C.A. Lee²⁹, G.R. Lee¹⁷, L. Lee⁵⁹, S.C. Lee¹⁵⁸, S. Lee⁷⁹, B. Lefebvre^{168a}, H.P. Lefebvre⁹⁴, M. Lefebvre¹⁷⁶, C. Leggett¹⁸, K. Lehmann¹⁵², N. Lehmann²⁰, G. Lehmann Miotto³⁶, W.A. Leight⁴⁶, A. Leisos^{162,v}, M.A.L. Leite^{81c}, C.E. Leitgeb¹¹⁴, R. Leitner¹⁴², K.J.C. Leney⁴², T. Lenz²⁴, S. Leone^{72a}, C. Leonidopoulos⁵⁰, A. Leopold¹³⁵, C. Leroy¹¹⁰, R. Les¹⁰⁷, C.G. Lester³², M. Levchenko¹³⁷, J. Levêque⁵, D. Levin¹⁰⁶, L.J. Levinson¹⁸⁰, D.J. Lewis²¹, B. Li^{15b}, B. Li¹⁰⁶, C-Q. Li^{60c,60d}, F. Li^{60c}, H. Li^{60a}, H. Li^{60b}, J. Li^{60c}, K. Li¹⁴⁸, L. Li^{60c}, M. Li^{15a,15d}, Q.Y. Li^{60a}, S. Li^{60d,60c,b}, X. Li⁴⁶, Y. Li⁴⁶, Z. Li^{60b}, Z. Li¹³⁴, Z. Li¹⁰⁴, Z. Li⁹¹, Z. Liang^{15a}, M. Liberatore⁴⁶, B. Liberti^{74a}, K. Lie^{63c}, S. Lim²⁹, C.Y. Lin³², K. Lin¹⁰⁷, R.A. Linck⁶⁶, R.E. Lindley⁷, J.H. Lindon²¹, A. Linss⁴⁶, A.L. Lioni⁵⁴, E. Lipeles¹³⁶, A. Lipniacka¹⁷, T.M. Liss^{173,ai}, A. Lister¹⁷⁵, J.D. Little⁸, B. Liu⁷⁹, B.X. Liu¹⁵², H.B. Liu²⁹, J.B. Liu^{60a}, J.K.K. Liu³⁷, K. Liu^{60d}, M. Liu^{60a}, M.Y. Liu^{60a}, P. Liu^{15a}, X. Liu^{60a}, Y. Liu⁴⁶, Y. Liu^{15a,15d}, Y.L. Liu¹⁰⁶, Y.W. Liu^{60a}, M. Livan^{71a,71b}, A. Lleres⁵⁸, J. Llorente Merino¹⁵², S.L. Lloyd⁹³, C.Y. Lo^{63b}, E.M. Lobodzinska⁴⁶, P. Loch⁷, S. Loffredo^{74a,74b}, T. Lohse¹⁹, K. Lohwasser¹⁴⁹, M. Lokajicek¹⁴⁰, J.D. Long¹⁷³, R.E. Long⁹⁰, I. Longarini^{73a,73b}, L. Longo³⁶, I. Lopez Paz¹⁰¹, A. Lopez Solis¹⁴⁹, J. Lorenz¹¹⁴, N. Lorenzo Martinez⁵, A.M. Lory¹¹⁴, A. Lösle⁵², X. Lou^{45a,45b}, X. Lou^{15a}, A. Lounis⁶⁵, J. Love⁶, P.A. Love⁹⁰, J.J. Lozano Bahilo¹⁷⁴, M. Lu^{60a}, Y.J. Lu⁶⁴, H.J. Lubatti¹⁴⁸, C. Luci^{73a,73b}, F.L. Lucio Alves^{15c}, A. Lucotte⁵⁸, F. Luehring⁶⁶, I. Luise¹⁵⁵, L. Luminari^{73a}, B. Lund-Jensen¹⁵⁴, N.A. Luongo¹³¹, M.S. Lutz¹⁶¹, D. Lynn²⁹, H. Lyons⁹¹, R. Lysak¹⁴⁰, E. Lytken⁹⁷, F. Lyu^{15a}, V. Lyubushkin⁸⁰, T. Lyubushkina⁸⁰, H. Ma²⁹, L.L. Ma^{60b}, Y. Ma⁹⁵, D.M. Mac Donell¹⁷⁶, G. Maccarrone⁵¹, C.M. Macdonald¹⁴⁹, J.C. MacDonald¹⁴⁹, J. Machado Miguens¹³⁶, R. Madar³⁸, W.F. Mader⁴⁸, M. Madugoda Ralalage Don¹²⁹, N. Madysa⁴⁸, J. Maeda⁸³, T. Maeno²⁹, M. Maerker⁴⁸, V. Magerl⁵², N. Magini⁷⁹, J. Magro^{67a,67c,r}, D.J. Mahon³⁹, C. Maidantchik^{81b}, A. Maio^{139a,139b,139d}, K. Maj^{84a}, O. Majersky^{28a}, S. Majewski¹³¹, Y. Makida⁸², N. Makovec⁶⁵, B. Malaescu¹³⁵, Pa. Malecki⁸⁵, V.P. Maleev¹³⁷, F. Malek⁵⁸, D. Malito^{41b,41a}, U. Mallik⁷⁸, C. Malone³², S. Maltezos¹⁰, S. Malyukov⁸⁰, J. Mamuzic¹⁷⁴, G. Mancini⁵¹, J.P. Mandalia⁹³, I. Mandić⁹², L. Manhaes de Andrade Filho^{81a}, I.M. Maniatis¹⁶², J. Manjarres Ramos⁴⁸, K.H. Mankinen⁹⁷, A. Mann¹¹⁴, A. Manousos⁷⁷, B. Mansoulie¹⁴⁴, I. Mantos¹⁶², S. Manzoni¹²⁰, A. Marantis¹⁶², G. Marceca³⁰, L. Marchese¹³⁴, G. Marchiori¹³⁵, M. Marcisovsky¹⁴⁰, L. Marcoccia^{74a,74b}, C. Marcon⁹⁷, M. Marjanovic¹²⁸, Z. Marshall¹⁸, M.U.F. Martensson¹⁷², S. Marti-Garcia¹⁷⁴, C.B. Martin¹²⁷, T.A. Martin¹⁷⁸, V.J. Martin⁵⁰, B. Martin dit Latour¹⁷, L. Martinelli^{75a,75b}, M. Martinez^{14,w}, P. Martinez Agullo¹⁷⁴, V.I. Martinez Outschoorn¹⁰³, S. Martin-Haugh¹⁴³, V.S. Martoiu^{27b}, A.C. Martyniuk⁹⁵, A. Marzin³⁶, S.R. Maschek¹¹⁵, L. Masetti¹⁰⁰, T. Mashimo¹⁶³, R. Mashinistov¹¹¹, J. Masik¹⁰¹, A.L. Maslennikov^{122b,122a}, L. Massa^{23b,23a}, P. Massarotti^{70a,70b}, P. Mastrandrea^{72a,72b}, A. Mastroberardino^{41b,41a}, T. Masubuchi¹⁶³, D. Matakias²⁹, A. Matic¹¹⁴, N. Matsuzawa¹⁶³, P. Mättig²⁴, J. Maurer^{27b}, B. Maček⁹², D.A. Maximov^{122b,122a}, R. Mazini¹⁵⁸, I. Maznas¹⁶², S.M. Mazza¹⁴⁵, J.P. Mc Gowan¹⁰⁴, S.P. Mc Kee¹⁰⁶, T.G. McCarthy¹¹⁵, W.P. McCormack¹⁸, E.F. McDonald¹⁰⁵, A.E. McDougall¹²⁰, J.A. Mcfayden¹⁸, G. Mchedlidze^{159b}, M.A. McKay⁴², K.D. McLean¹⁷⁶, S.J. McMahon¹⁴³, P.C. McNamara¹⁰⁵, C.J. McNicol¹⁷⁸, R.A. McPherson^{176,aa}, J.E. Mdhului^{33e}, Z.A. Meadows¹⁰³, S. Meehan³⁶, T. Megy³⁸, S. Mehlhase¹¹⁴, A. Mehta⁹¹, B. Meirose⁴³, D. Melini¹⁶⁰, B.R. Mellado Garcia^{33e}, J.D. Mellenthin⁵³, M. Melo^{28a}, F. Meloni⁴⁶, A. Melzer²⁴, E.D. Mendes Gouveia^{139a,139e}, A.M. Mendes Jacques Da Costa²¹, H.Y. Meng¹⁶⁷, L. Meng³⁶, X.T. Meng¹⁰⁶, S. Menke¹¹⁵, E. Meoni^{41b,41a}, S. Mergelmeyer¹⁹, S.A.M. Merkt¹³⁸, C. Merlassino¹³⁴, P. Mermod⁵⁴, L. Merola^{70a,70b}, C. Meroni^{69a}, G. Merz¹⁰⁶, O. Meshkov^{113,111}, J.K.R. Meshreki¹⁵¹, J. Metcalfe⁶, A.S. Mete⁶, C. Meyer⁶⁶, J-P. Meyer¹⁴⁴, M. Michetti¹⁹, R.P. Middleton¹⁴³, L. Mijović⁵⁰, G. Mikenberg¹⁸⁰, M. Mikestikova¹⁴⁰, M. Mikuž⁹², H. Mildner¹⁴⁹, A. Milic¹⁶⁷, C.D. Milke⁴², D.W. Miller³⁷, L.S. Miller³⁴, A. Milov¹⁸⁰, D.A. Milstead^{45a,45b}, A.A. Minaenko¹²³, I.A. Minashvili^{159b}, L. Mince⁵⁷, A.I. Mincer¹²⁵, B. Mindur^{84a}, M. Mineev⁸⁰, Y. Minegishi¹⁶³, Y. Mino⁸⁶, L.M. Mir¹⁴, M. Mironova¹³⁴, T. Mitani¹⁷⁹, J. Mitrevski¹¹⁴, V.A. Mitsou¹⁷⁴,

M. Mittal^{60c}, O. Miu¹⁶⁷, A. Miucci²⁰, P.S. Miyagawa⁹³, A. Mizukami⁸², J.U. Mjörnmark⁹⁷,
T. Mkrtchyan^{61a}, M. Mlynarikova¹²¹, T. Moa^{45a,45b}, S. Mobius⁵³, K. Mochizuki¹¹⁰, P. Moder⁴⁶,
P. Mogg¹¹⁴, S. Mohapatra³⁹, R. Moles-Valls²⁴, K. Mönig⁴⁶, E. Monnier¹⁰², A. Montalbano¹⁵²,
J. Montejo Berlingen³⁶, M. Montella⁹⁵, F. Monticelli⁸⁹, S. Monzani^{69a}, N. Morange⁶⁵,
A.L. Moreira De Carvalho^{139a}, D. Moreno^{22a}, M. Moreno Llácer¹⁷⁴, C. Moreno Martinez¹⁴,
P. Morettini^{55b}, M. Morgenstern¹⁶⁰, S. Morgenstern⁴⁸, D. Mori¹⁵², M. Morii⁵⁹, M. Morinaga¹⁷⁹,
V. Morisbak¹³³, A.K. Morley³⁶, G. Mornacchi³⁶, A.P. Morris⁹⁵, L. Morvaj³⁶, P. Moschovakos³⁶,
B. Moser¹²⁰, M. Mosidze^{159b}, T. Moskalets¹⁴⁴, P. Moskvitina¹¹⁹, J. Moss^{31,n}, E.J.W. Moyse¹⁰³,
S. Muanza¹⁰², J. Mueller¹³⁸, R.S.P. Mueller¹¹⁴, D. Muenstermann⁹⁰, G.A. Mullier⁹⁷, D.P. Mungo^{69a,69b},
J.L. Munoz Martinez¹⁴, F.J. Munoz Sanchez¹⁰¹, P. Murin^{28b}, W.J. Murray^{178,143}, A. Murrone^{69a,69b},
J.M. Muse¹²⁸, M. Muškinja¹⁸, C. Mwewa^{33a}, A.G. Myagkov^{123,ae}, A.A. Myers¹³⁸, G. Myers⁶⁶, J. Myers¹³¹,
M. Myska¹⁴¹, B.P. Nachman¹⁸, O. Nackenhorst⁴⁷, A.Nag Nag⁴⁸, K. Nagai¹³⁴, K. Nagano⁸², Y. Nagasaka⁶²,
J.L. Nagle²⁹, E. Nagy¹⁰², A.M. Nairz³⁶, Y. Nakahama¹¹⁷, K. Nakamura⁸², T. Nakamura¹⁶³, H. Nanjo¹³²,
F. Napolitano^{61a}, R.F. Naranjo Garcia⁴⁶, R. Narayan⁴², I. Naryshkin¹³⁷, M. Naseri³⁴, T. Naumann⁴⁶,
G. Navarro^{22a}, P.Y. Nechaeva¹¹¹, F. Nechansky⁴⁶, T.J. Neep²¹, A. Negri^{71a,71b}, M. Negrini^{23b}, C. Nellist¹¹⁹,
C. Nelson¹⁰⁴, M.E. Nelson^{45a,45b}, S. Nemecek¹⁴⁰, M. Nessi^{36,f}, M.S. Neubauer¹⁷³, F. Neuhaus¹⁰⁰,
M. Neumann¹⁸², R. Newhouse¹⁷⁵, P.R. Newman²¹, C.W. Ng¹³⁸, Y.S. Ng¹⁹, Y.W.Y. Ng¹⁷¹, B. Ngair^{35e},
H.D.N. Nguyen¹⁰², T. Nguyen Manh¹¹⁰, E. Nibigira³⁸, R.B. Nickerson¹³⁴, R. Nicolaidou¹⁴⁴,
D.S. Nielsen⁴⁰, J. Nielsen¹⁴⁵, M. Niemeyer⁵³, N. Nikiforou¹¹, V. Nikolaenko^{123,ae}, I. Nikolic-Audit¹³⁵,
K. Nikolopoulos²¹, P. Nilsson²⁹, H.R. Nindhito⁵⁴, A. Nisati^{73a}, N. Nishu^{60c}, R. Nisius¹¹⁵, I. Nitsche⁴⁷,
T. Nitta¹⁷⁹, T. Nobe¹⁶³, D.L. Noel³², Y. Noguchi⁸⁶, I. Nomidis¹³⁵, M.A. Nomura²⁹, M. Nordberg³⁶,
J. Novak⁹², T. Novak⁹², O. Novgorodova⁴⁸, R. Novotny¹¹⁸, L. Nozka¹³⁰, K. Ntekas¹⁷¹, E. Nurse⁹⁵,
F.G. Oakham^{34,aj}, J. Ocariz¹³⁵, A. Ochi⁸³, I. Ochoa^{139a}, J.P. Ochoa-Ricoux^{146a}, K. O'Connor²⁶, S. Oda⁸⁸,
S. Odaka⁸², S. Oerdek⁵³, A. Ogrodnik^{84a}, A. Oh¹⁰¹, C.C. Ohm¹⁵⁴, H. Oide¹⁶⁵, R. Oishi¹⁶³, M.L. Ojeda¹⁶⁷,
H. Okawa¹⁶⁹, Y. Okazaki⁸⁶, M.W. O'Keefe⁹¹, Y. Okumura¹⁶³, A. Olariu^{27b}, L.F. Oleiro Seabra^{139a},
S.A. Olivares Pino^{146a}, D. Oliveira Damazio²⁹, J.L. Oliver¹, M.J.R. Olsson¹⁷¹, A. Olszewski⁸⁵,
J. Olszowska⁸⁵, Ö.O. Öncel²⁴, D.C. O'Neil¹⁵², A.P. O'Neill¹³⁴, A. Onofre^{139a,139e}, P.U.E. Onyisi¹¹,
H. Oppen¹³³, R.G. Oreamuno Madriz¹²¹, M.J. Oreglia³⁷, G.E. Orellana⁸⁹, D. Orestano^{75a,75b},
N. Orlando¹⁴, R.S. Orr¹⁶⁷, V. O'Shea⁵⁷, R. Ospanov^{60a}, G. Otero y Garzon³⁰, H. Otono⁸⁸, P.S. Ott^{61a},
G.J. Ottino¹⁸, M. Ouchrif^{35d}, J. Ouellette²⁹, F. Ould-Saada¹³³, A. Ouraou^{144,*}, Q. Ouyang^{15a}, M. Owen⁵⁷,
R.E. Owen¹⁴³, V.E. Ozcan^{12c}, N. Ozturk⁸, J. Pacalt¹³⁰, H.A. Pacey³², K. Pachal⁴⁹, A. Pacheco Pages¹⁴,
C. Padilla Aranda¹⁴, S. Pagan Griso¹⁸, G. Palacino⁶⁶, S. Palazzo⁵⁰, S. Palestini³⁶, M. Palka^{84b}, P. Palni^{84a},
C.E. Pandini⁵⁴, J.G. Panduro Vazquez⁹⁴, P. Pani⁴⁶, G. Panizzo^{67a,67c}, L. Paolozzi⁵⁴, C. Papadatos¹¹⁰,
K. Papageorgiou^{9,h}, S. Parajuli⁴², A. Paramonov⁶, C. Paraskevopoulos¹⁰, D. Paredes Hernandez^{63b},
S.R. Paredes Saenz¹³⁴, B. Parida¹⁸⁰, T.H. Park¹⁶⁷, A.J. Parker³¹, M.A. Parker³², F. Parodi^{55b,55a},
E.W. Parrish¹²¹, J.A. Parsons³⁹, U. Parzefall⁵², L. Pascual Dominguez¹³⁵, V.R. Pascuzzi¹⁸,
J.M.P. Pasner¹⁴⁵, F. Pasquali¹²⁰, E. Pasqualucci^{73a}, S. Passaggio^{55b}, F. Pastore⁹⁴, P. Pasuwan^{45a,45b},
S. Patariaia¹⁰⁰, J.R. Pater¹⁰¹, A. Pathak^{181,j}, J. Patton⁹¹, T. Pauly³⁶, J. Parkes¹⁵³, M. Pedersen¹³³,
L. Pedraza Diaz¹¹⁹, R. Pedro^{139a}, T. Peiffer⁵³, S.V. Peleganchuk^{122b,122a}, O. Penc¹⁴⁰, C. Peng^{63b},
H. Peng^{60a}, B.S. Peralva^{81a}, M.M. Perego⁶⁵, A.P. Pereira Peixoto^{139a}, L. Pereira Sanchez^{45a,45b},
D.V. Perepelitsa²⁹, E. Perez Codina^{168a}, L. Perini^{69a,69b}, H. Pernegger³⁶, S. Perrella³⁶, A. Perrevoort¹²⁰,
K. Peters⁴⁶, R.F.Y. Peters¹⁰¹, B.A. Petersen³⁶, T.C. Petersen⁴⁰, E. Petit¹⁰², V. Petousis¹⁴¹, C. Petridou¹⁶²,
F. Petrucci^{75a,75b}, M. Pettee¹⁸³, N.E. Pettersson¹⁰³, K. Petukhova¹⁴², A. Peyaud¹⁴⁴, R. Pezoa^{146d},
L. Pezzotti^{71a,71b}, T. Pham¹⁰⁵, P.W. Phillips¹⁴³, M.W. Phipps¹⁷³, G. Piacquadio¹⁵⁵, E. Pianori¹⁸,
A. Picazio¹⁰³, R.H. Pickles¹⁰¹, R. Piegaia³⁰, D. Pietreanu^{27b}, J.E. Pilcher³⁷, A.D. Pilkington¹⁰¹,
M. Pinamonti^{67a,67c}, J.L. Pinfold³, C. Pitman Donaldson⁹⁵, M. Pitt¹⁶¹, L. Pizzimento^{74a,74b}, A. Pizzini¹²⁰,
M.-A. Pleier²⁹, V. Plesanovs⁵², V. Pleskot¹⁴², E. Plotnikova⁸⁰, P. Podberezko^{122b,122a}, R. Poettgen⁹⁷,

R. Poggi⁵⁴, L. Poggioli¹³⁵, I. Pogrebnyak¹⁰⁷, D. Pohl²⁴, I. Pokharel⁵³, G. Polesello^{71a}, A. Poley^{152,168a}, A. Policicchio^{73a,73b}, R. Polifka¹⁴², A. Polini^{23b}, C.S. Pollard⁴⁶, V. Polychronakos²⁹, D. Ponomarenko¹¹², L. Pontecorvo³⁶, S. Popa^{27a}, G.A. Popeneciu^{27d}, L. Portales⁵, D.M. Portillo Quintero⁵⁸, S. Pospisil¹⁴¹, K. Potamianos⁴⁶, I.N. Potrap⁸⁰, C.J. Potter³², H. Potti¹¹, T. Poulsen⁹⁷, J. Poveda¹⁷⁴, T.D. Powell¹⁴⁹, G. Pownall⁴⁶, M.E. Pozo Astigarraga³⁶, A. Prades Ibanez¹⁷⁴, P. Pralavorio¹⁰², M.M. Prapa⁴⁴, S. Prell⁷⁹, D. Price¹⁰¹, M. Primavera^{68a}, M.L. Proffitt¹⁴⁸, N. Proklova¹¹², K. Prokofiev^{63c}, F. Prokoshin⁸⁰, S. Protopopescu²⁹, J. Proudfoot⁶, M. Przybycien^{84a}, D. Pudzha¹³⁷, A. Puri¹⁷³, P. Puzo⁶⁵, D. Pyatiizbyantseva¹¹², J. Qian¹⁰⁶, Y. Qin¹⁰¹, A. Quadt⁵³, M. Queitsch-Maitland³⁶, G. Rabanal Bolanos⁵⁹, M. Racko^{28a}, F. Ragusa^{69a,69b}, G. Rahal⁹⁸, J.A. Raine⁵⁴, S. Rajagopalan²⁹, A. Ramirez Morales⁹³, K. Ran^{15a,15d}, D.F. Rassloff^{61a}, D.M. Rauch⁴⁶, F. Rauscher¹¹⁴, S. Rave¹⁰⁰, B. Ravina⁵⁷, I. Ravinovich¹⁸⁰, J.H. Rawling¹⁰¹, M. Raymond³⁶, A.L. Read¹³³, N.P. Readoff¹⁴⁹, M. Reale^{68a,68b}, D.M. Rebuffi^{71a,71b}, G. Redlinger²⁹, K. Reeves⁴³, D. Reikher¹⁶¹, A. Reiss¹⁰⁰, A. Rej¹⁵¹, C. Rembser³⁶, A. Renardi⁴⁶, M. Renda^{27b}, M.B. Rendel¹¹⁵, A.G. Rennie⁵⁷, S. Resconi^{69a}, E.D. Resseguie¹⁸, S. Rettie⁹⁵, B. Reynolds¹²⁷, E. Reynolds²¹, O.L. Rezanova^{122b,122a}, P. Reznicek¹⁴², E. Ricci^{76a,76b}, R. Richter¹¹⁵, S. Richter⁴⁶, E. Richter-Was^{84b}, M. Ridel¹³⁵, P. Rieck¹¹⁵, O. Rifki⁴⁶, M. Rijssenbeek¹⁵⁵, A. Rimoldi^{71a,71b}, M. Rimoldi⁴⁶, L. Rinaldi^{23b}, T.T. Rinn¹⁷³, G. Ripellino¹⁵⁴, I. Riu¹⁴, P. Rivadeneira⁴⁶, J.C. Rivera Vergara¹⁷⁶, F. Rizatdinova¹²⁹, E. Rizvi⁹³, C. Rizzi³⁶, S.H. Robertson^{104,aa}, M. Robin⁴⁶, D. Robinson³², C.M. Robles Gajardo^{146d}, M. Robles Manzano¹⁰⁰, A. Robson⁵⁷, A. Rocchi^{74a,74b}, C. Roda^{72a,72b}, S. Rodriguez Bosca¹⁷⁴, A. Rodriguez Rodriguez⁵², A.M. Rodríguez Vera^{168b}, S. Roe³⁶, J. Roggel¹⁸², O. Røhne¹³³, R. Röhrig¹¹⁵, R.A. Rojas^{146d}, B. Roland⁵², C.P.A. Roland⁶⁶, J. Roloff²⁹, A. Romaniouk¹¹², M. Romano^{23b,23a}, N. Rompotis⁹¹, M. Ronzani¹²⁵, L. Roos¹³⁵, S. Rosati^{73a}, G. Rosin¹⁰³, B.J. Rosser¹³⁶, E. Rossi⁴⁶, E. Rossi^{75a,75b}, E. Rossi^{70a,70b}, L.P. Rossi^{55b}, L. Rossini⁴⁶, R. Rosten¹⁴, M. Rotaru^{27b}, B. Rottler⁵², D. Rousseau⁶⁵, G. Rovelli^{71a,71b}, A. Roy¹¹, D. Roy^{33e}, A. Rozanov¹⁰², Y. Rozen¹⁶⁰, X. Ruan^{33e}, T.A. Ruggeri¹, F. Rühr⁵², A. Ruiz-Martinez¹⁷⁴, A. Rummeler³⁶, Z. Rurikova⁵², N.A. Rusakovich⁸⁰, H.L. Russell¹⁰⁴, L. Rustige^{38,47}, J.P. Rutherford⁷, E.M. Rüttinger¹⁴⁹, M. Rybar¹⁴², G. Rybkin⁶⁵, E.B. Rye¹³³, A. Ryzhov¹²³, J.A. Sabater Iglesias⁴⁶, P. Sabatini¹⁷⁴, L. Sabetta^{73a,73b}, S. Sacerdoti⁶⁵, H.F.W. Sadrozinski¹⁴⁵, R. Sadykov⁸⁰, F. Safai Tehrani^{73a}, B. Safarzadeh Samani¹⁵⁶, M. Safdari¹⁵³, P. Saha¹²¹, S. Saha¹⁰⁴, M. Sahinsoy¹¹⁵, A. Sahu¹⁸², M. Saimpert³⁶, M. Saito¹⁶³, T. Saito¹⁶³, H. Sakamoto¹⁶³, D. Salamani⁵⁴, G. Salamanna^{75a,75b}, A. Salnikov¹⁵³, J. Salt¹⁷⁴, A. Salvador Salas¹⁴, D. Salvatore^{41b,41a}, F. Salvatore¹⁵⁶, A. Salvucci^{63a}, A. Salzburger³⁶, J. Samarati³⁶, D. Sammel⁵², D. Sampsonidis¹⁶², D. Sampsonidou^{60d,60c}, J. Sánchez¹⁷⁴, A. Sanchez Pineda^{67a,36,67c}, H. Sandaker¹³³, C.O. Sander⁴⁶, I.G. Sanderswood⁹⁰, M. Sandhoff¹⁸², C. Sandoval^{22b}, D.P.C. Sankey¹⁴³, M. Sannino^{55b,55a}, Y. Sano¹¹⁷, A. Sansoni⁵¹, C. Santoni³⁸, H. Santos^{139a,139b}, S.N. Santpur¹⁸, A. Santra¹⁷⁴, K.A. Saoucha¹⁴⁹, A. Sapronov⁸⁰, J.G. Saraiva^{139a,139d}, O. Sasaki⁸², K. Sato¹⁶⁹, F. Sauerburger⁵², E. Sauvan⁵, P. Savard^{167,aj}, R. Sawada¹⁶³, C. Sawyer¹⁴³, L. Sawyer⁹⁶, I. Sayago Galvan¹⁷⁴, C. Sbarra^{23b}, A. Sbrizzi^{67a,67c}, T. Scanlon⁹⁵, J. Schaarschmidt¹⁴⁸, P. Schacht¹¹⁵, D. Schaefer³⁷, L. Schaefer¹³⁶, U. Schäfer¹⁰⁰, A.C. Schaffer⁶⁵, D. Schaile¹¹⁴, R.D. Schamberger¹⁵⁵, E. Schanet¹¹⁴, C. Scharf¹⁹, N. Scharmberg¹⁰¹, V.A. Schegelsky¹³⁷, D. Scheirich¹⁴², F. Schenck¹⁹, M. Schernau¹⁷¹, C. Schiavi^{55b,55a}, L.K. Schildgen²⁴, Z.M. Schillaci²⁶, E.J. Schioppa^{68a,68b}, M. Schioppa^{41b,41a}, K.E. Schleicher⁵², S. Schlenker³⁶, K.R. Schmidt-Sommerfeld¹¹⁵, K. Schmieden¹⁰⁰, C. Schmitt¹⁰⁰, S. Schmitt⁴⁶, L. Schoeffel¹⁴⁴, A. Schoening^{61b}, P.G. Scholer⁵², E. Schopf¹³⁴, M. Schott¹⁰⁰, J.F.P. Schouwenberg¹¹⁹, J. Schovancova³⁶, S. Schramm⁵⁴, F. Schroeder¹⁸², A. Schulte¹⁰⁰, H-C. Schultz-Coulon^{61a}, M. Schumacher⁵², B.A. Schumm¹⁴⁵, Ph. Schune¹⁴⁴, A. Schwartzman¹⁵³, T.A. Schwarz¹⁰⁶, Ph. Schwemling¹⁴⁴, R. Schwienhorst¹⁰⁷, A. Sciandra¹⁴⁵, G. Sciolla²⁶, F. Scuri^{72a}, F. Scutti¹⁰⁵, L.M. Scyboz¹¹⁵, C.D. Sebastiani⁹¹, K. Sedlaczek⁴⁷, P. Seema¹⁹, S.C. Seidel¹¹⁸, A. Seiden¹⁴⁵, B.D. Seidlitz²⁹, T. Seiss³⁷, C. Seitz⁴⁶, J.M. Seixas^{81b}, G. Sekhniaidze^{70a}, S.J. Sekula⁴², N. Semprini-Cesari^{23b,23a}, S. Sen⁴⁹, C. Serfon²⁹, L. Serin⁶⁵, L. Serkin^{67a,67b}, M. Sessa^{60a}, H. Severini¹²⁸, S. Sevova¹⁵³, F. Sforza^{55b,55a},

A. Sfyra⁵⁴, E. Shabalina⁵³, J.D. Shahinian¹³⁶, N.W. Shaikh^{45a,45b}, D. Shaked Renous¹⁸⁰, L.Y. Shan^{15a},
 M. Shapiro¹⁸, A. Sharma³⁶, A.S. Sharma¹, P.B. Shatalov¹²⁴, K. Shaw¹⁵⁶, S.M. Shaw¹⁰¹, M. Shehade¹⁸⁰,
 Y. Shen¹²⁸, A.D. Sherman²⁵, P. Sherwood⁹⁵, L. Shi⁹⁵, C.O. Shimmin¹⁸³, Y. Shimogama¹⁷⁹,
 M. Shimojima¹¹⁶, J.D. Shinner⁹⁴, I.P.J. Shipsey¹³⁴, S. Shirabe¹⁶⁵, M. Shiyakova^{80,y}, J. Shlomi¹⁸⁰,
 A. Shmeleva¹¹¹, M.J. Shochet³⁷, J. Shojaii¹⁰⁵, D.R. Shope¹⁵⁴, S. Shrestha¹²⁷, E.M. Shrif^{33e}, M.J. Shroff¹⁷⁶,
 E. Shulga¹⁸⁰, P. Sicho¹⁴⁰, A.M. Sickles¹⁷³, E. Sideras Haddad^{33e}, O. Sidiropoulou³⁶, A. Sidoti^{23b,23a},
 F. Siegert⁴⁸, Dj. Sijacki¹⁶, M.Jr. Silva¹⁸¹, M.V. Silva Oliveira³⁶, S.B. Silverstein^{45a}, S. Simion⁶⁵,
 R. Simoniello¹⁰⁰, C.J. Simpson-allsoy²¹, S. Simsek^{12b}, P. Sinervo¹⁶⁷, V. Sinetckii¹¹³, S. Singh¹⁵²,
 S. Sinha^{33e}, M. Sioli^{23b,23a}, I. Siral¹³¹, S.Yu. Sivoklov¹¹³, J. Sjölin^{45a,45b}, A. Skaf⁵³, E. Skorda⁹⁷,
 P. Skubic¹²⁸, M. Slawinska⁸⁵, K. Sliwa¹⁷⁰, V. Smakhtin¹⁸⁰, B.H. Smart¹⁴³, J. Smiesko^{28b}, N. Smirnov¹¹²,
 S.Yu. Smirnov¹¹², Y. Smirnov¹¹², L.N. Smirnova^{113,s}, O. Smirnova⁹⁷, E.A. Smith³⁷, H.A. Smith¹³⁴,
 M. Smizanska⁹⁰, K. Smolek¹⁴¹, A. Smykiewicz⁸⁵, A.A. Snesarev¹¹¹, H.L. Snoek¹²⁰, I.M. Snyder¹³¹,
 S. Snyder²⁹, R. Sobie^{176,aa}, A. Soffer¹⁶¹, A. Sjøgaard⁵⁰, F. Sohns⁵³, C.A. Solans Sanchez³⁶,
 E.Yu. Soldatov¹¹², U. Soldevila¹⁷⁴, A.A. Solodkov¹²³, A. Soloshenko⁸⁰, O.V. Solovyanov¹²³,
 V. Solovyev¹³⁷, P. Sommer¹⁴⁹, H. Son¹⁷⁰, A. Sonay¹⁴, W. Song¹⁴³, W.Y. Song^{168b}, A. Sopczak¹⁴¹,
 A.L. Sopio⁹⁵, F. Sopkova^{28b}, S. Sottocornola^{71a,71b}, R. Soualah^{67a,67c}, A.M. Soukharev^{122b,122a}, D. South⁴⁶,
 S. Spagnolo^{68a,68b}, M. Spalla¹¹⁵, M. Spangenberg¹⁷⁸, F. Spanò⁹⁴, D. Sperlich⁵², T.M. Spieker^{61a},
 G. Spigo³⁶, M. Spina¹⁵⁶, D.P. Spiteri⁵⁷, M. Spousta¹⁴², A. Stabile^{69a,69b}, B.L. Stamas¹²¹, R. Stamen^{61a},
 M. Stamenkovic¹²⁰, A. Stampekis²¹, E. Stanecka⁸⁵, B. Stanislaus¹³⁴, M.M. Stanitzki⁴⁶, M. Stankaityte¹³⁴,
 B. Stapf¹²⁰, E.A. Starchenko¹²³, G.H. Stark¹⁴⁵, J. Stark⁵⁸, P. Staroba¹⁴⁰, P. Starovoitov^{61a}, S. Stärz¹⁰⁴,
 R. Staszewski⁸⁵, G. Stavropoulos⁴⁴, M. Stegler⁴⁶, P. Steinberg²⁹, A.L. Steinhebel¹³¹, B. Stelzer^{152,168a},
 H.J. Stelzer¹³⁸, O. Stelzer-Chilton^{168a}, H. Stenzel⁵⁶, T.J. Stevenson¹⁵⁶, G.A. Stewart³⁶, M.C. Stockton³⁶,
 G. Stoicea^{27b}, M. Stolarski^{139a}, S. Stonjek¹¹⁵, A. Straessner⁴⁸, J. Strandberg¹⁵⁴, S. Strandberg^{45a,45b},
 M. Strauss¹²⁸, T. Strebler¹⁰², P. Strizenec^{28b}, R. Ströhmer¹⁷⁷, D.M. Strom¹³¹, L.R. Strom⁴⁶,
 R. Stroynowski⁴², A. Strubig^{45a,45b}, S.A. Stucci²⁹, B. Stugu¹⁷, J. Stupak¹²⁸, N.A. Styles⁴⁶, D. Su¹⁵³,
 W. Su^{60d,148,60c}, X. Su^{60a}, N.B. Suarez¹³⁸, V.V. Sulin¹¹¹, M.J. Sullivan⁹¹, D.M.S. Sultan⁵⁴, S. Sultansoy^{4c},
 T. Sumida⁸⁶, S. Sun¹⁰⁶, X. Sun¹⁰¹, C.J.E. Suster¹⁵⁷, M.R. Sutton¹⁵⁶, S. Suzuki⁸², M. Svatos¹⁴⁰,
 M. Swiatlowski^{168a}, S.P. Swift², T. Swirski¹⁷⁷, A. Sydorenko¹⁰⁰, I. Sykora^{28a}, M. Sykora¹⁴², T. Sykora¹⁴²,
 D. Ta¹⁰⁰, K. Tackmann^{46,x}, J. Taenzer¹⁶¹, A. Taffard¹⁷¹, R. Tafirout^{168a}, E. Tagiev¹²³, R.H.M. Taibah¹³⁵,
 R. Takashima⁸⁷, K. Takeda⁸³, T. Takeshita¹⁵⁰, E.P. Takeva⁵⁰, Y. Takubo⁸², M. Talby¹⁰²,
 A.A. Talyshev^{122b,122a}, K.C. Tam^{63b}, N.M. Tamir¹⁶¹, J. Tanaka¹⁶³, R. Tanaka⁶⁵, S. Tapia Araya¹⁷³,
 S. Tapprogge¹⁰⁰, A. Tarek Abouelfadl Mohamed¹⁰⁷, S. Tarem¹⁶⁰, K. Tariq^{60b}, G. Tarna^{27b,e},
 G.F. Tartarelli^{69a}, P. Tas¹⁴², M. Tasevsky¹⁴⁰, E. Tassi^{41b,41a}, G. Tateno¹⁶³, A. Tavares Delgado^{139a},
 Y. Tayalati^{35e}, A.J. Taylor⁵⁰, G.N. Taylor¹⁰⁵, W. Taylor^{168b}, H. Teagle⁹¹, A.S. Tee⁹⁰,
 R. Teixeira De Lima¹⁵³, P. Teixeira-Dias⁹⁴, H. Ten Kate³⁶, J.J. Teoh¹²⁰, K. Terashi¹⁶³, J. Terron⁹⁹,
 S. Terzo¹⁴, M. Testa⁵¹, R.J. Teuscher^{167,aa}, N. Themistokleous⁵⁰, T. Thevenaux-Pelzer¹⁹, D.W. Thomas⁹⁴,
 J.P. Thomas²¹, E.A. Thompson⁴⁶, P.D. Thompson²¹, E. Thomson¹³⁶, E.J. Thorpe⁹³, V.O. Tikhomirov^{111,af},
 Yu.A. Tikhonov^{122b,122a}, S. Timoshenko¹¹², P. Tipton¹⁸³, S. Tisserant¹⁰², K. Todome^{23b,23a},
 S. Todorova-Nova¹⁴², S. Todt⁴⁸, J. Tojo⁸⁸, S. Tokár^{28a}, K. Tokushuku⁸², E. Tolley¹²⁷, R. Tombs³²,
 K.G. Tomiwa^{33e}, M. Tomoto^{82,117}, L. Tompkins¹⁵³, P. Tornambe¹⁰³, E. Torrence¹³¹, H. Torres⁴⁸,
 E. Torró Pastor¹⁷⁴, M. Toscani³⁰, C. Toscirì¹³⁴, J. Toth^{102,z}, D.R. Tovey¹⁴⁹, A. Traeet¹⁷, C.J. Treado¹²⁵,
 T. Trefzger¹⁷⁷, F. Tresoldi¹⁵⁶, A. Tricoli²⁹, I.M. Trigger^{168a}, S. Trincaz-Duvoid¹³⁵, D.A. Trischuk¹⁷⁵,
 W. Trischuk¹⁶⁷, B. Trocme⁵⁸, A. Trofymov⁶⁵, C. Troncon^{69a}, F. Trovato¹⁵⁶, L. Truong^{33c}, M. Trzebinski⁸⁵,
 A. Trzupek⁸⁵, F. Tsai⁴⁶, P.V. Tsiarehka^{108,ad}, A. Tsirigotis^{162,v}, V. Tsiskaridze¹⁵⁵, E.G. Tskhadadze^{159a},
 M. Tsopoulou¹⁶², I.I. Tsukerman¹²⁴, V. Tsulaia¹⁸, S. Tsuno⁸², D. Tsybychev¹⁵⁵, Y. Tu^{63b}, A. Tudorache^{27b},
 V. Tudorache^{27b}, A.N. Tuna³⁶, S. Turchikhin⁸⁰, D. Turgeman¹⁸⁰, I. Turk Cakir^{4b,t}, R.J. Turner²¹,
 R. Turra^{69a}, P.M. Tuts³⁹, S. Tzamarias¹⁶², E. Tzovara¹⁰⁰, K. Uchida¹⁶³, F. Ukegawa¹⁶⁹, G. Unal³⁶,

M. Unal¹¹, A. Undrus²⁹, G. Unel¹⁷¹, F.C. Ungaro¹⁰⁵, Y. Unno⁸², K. Uno¹⁶³, J. Urban^{28b}, P. Urquijo¹⁰⁵,
G. Usai⁸, Z. Uysal^{12d}, V. Vacek¹⁴¹, B. Vachon¹⁰⁴, K.O.H. Vadla¹³³, T. Vafeiadis³⁶, A. Vaidya⁹⁵,
C. Valderanis¹¹⁴, E. Valdes Santurio^{45a,45b}, M. Valente^{168a}, S. Valentinetti^{23b,23a}, A. Valero¹⁷⁴, L. Valéry⁴⁶,
R.A. Vallance²¹, A. Vallier³⁶, J.A. Valls Ferrer¹⁷⁴, T.R. Van Daalen¹⁴, P. Van Gemmeren⁶, S. Van Stroud⁹⁵,
I. Van Vulpen¹²⁰, M. Vanadia^{74a,74b}, W. Vandelli³⁶, M. Vandenbroucke¹⁴⁴, E.R. Vandewall¹²⁹,
D. Vannicola^{73a,73b}, R. Vari^{73a}, E.W. Varnes⁷, C. Varni^{55b,55a}, T. Varol¹⁵⁸, D. Varouchas⁶⁵, K.E. Varvell¹⁵⁷,
M.E. Vasile^{27b}, G.A. Vasquez¹⁷⁶, F. Vazeille³⁸, D. Vazquez Furelos¹⁴, T. Vazquez Schroeder³⁶, J. Veatch⁵³,
V. Vecchio¹⁰¹, M.J. Veen¹²⁰, L.M. Veloce¹⁶⁷, F. Veloso^{139a,139c}, S. Veneziano^{73a}, A. Ventura^{68a,68b},
A. Verbytskyi¹¹⁵, V. Vercesi^{71a}, M. Verducci^{72a,72b}, C.M. Vergel Infante⁷⁹, C. Vergis²⁴, W. Verkerke¹²⁰,
A.T. Vermeulen¹²⁰, J.C. Vermeulen¹²⁰, C. Vernieri¹⁵³, P.J. Verschuuren⁹⁴, M.C. Vetterli^{152,aj},
N. Viaux Maira^{146d}, T. Vickey¹⁴⁹, O.E. Vickey Boeriu¹⁴⁹, G.H.A. Viehhauser¹³⁴, L. Vigani^{61b},
M. Villa^{23b,23a}, M. Villaplana Perez¹⁷⁴, E.M. Villhauer⁵⁰, E. Vilucchi⁵¹, M.G. Vinciter³⁴, G.S. Virdee²¹,
A. Vishwakarma⁵⁰, C. Vittori^{23b,23a}, I. Vivarelli¹⁵⁶, M. Vogel¹⁸², P. Vokac¹⁴¹, J. Von Ahnen⁴⁶,
S.E. von Buddenbrock^{33e}, E. Von Toerne²⁴, V. Vorobel¹⁴², K. Vorobev¹¹², M. Vos¹⁷⁴, J.H. Vosseveld⁹¹,
M. Vozak¹⁰¹, N. Vranjes¹⁶, M. Vranjes Milosavljevic¹⁶, V. Vrba¹⁴¹, M. Vreeswijk¹²⁰, N.K. Vu¹⁰²,
R. Vuillermet³⁶, I. Vukotic³⁷, S. Wada¹⁶⁹, P. Wagner²⁴, W. Wagner¹⁸², J. Wagner-Kuhr¹¹⁴, S. Wahdan¹⁸²,
H. Wahlberg⁸⁹, R. Wakasa¹⁶⁹, V.M. Walbrecht¹¹⁵, J. Walder¹⁴³, R. Walker¹¹⁴, S.D. Walker⁹⁴,
W. Walkowiak¹⁵¹, V. Wallangen^{45a,45b}, A.M. Wang⁵⁹, A.Z. Wang¹⁸¹, C. Wang^{60a}, C. Wang^{60c}, H. Wang¹⁸,
H. Wang³, J. Wang^{63a}, P. Wang⁴², Q. Wang¹²⁸, R.-J. Wang¹⁰⁰, R. Wang^{60a}, R. Wang⁶, S.M. Wang¹⁵⁸,
W.T. Wang^{60a}, W. Wang^{15c}, W.X. Wang^{60a}, Y. Wang^{60a}, Z. Wang¹⁰⁶, C. Wanotayaroj⁴⁶, A. Warburton¹⁰⁴,
C.P. Ward³², R.J. Ward²¹, N. Warrack⁵⁷, A.T. Watson²¹, M.F. Watson²¹, G. Watts¹⁴⁸, B.M. Waugh⁹⁵,
A.F. Webb¹¹, C. Weber²⁹, M.S. Weber²⁰, S.A. Weber³⁴, S.M. Weber^{61a}, Y. Wei¹³⁴, A.R. Weidberg¹³⁴,
J. Weingarten⁴⁷, M. Weirich¹⁰⁰, C. Weiser⁵², P.S. Wells³⁶, T. Wenaus²⁹, B. Wendland⁴⁷, T. Wengler³⁶,
S. Wenig³⁶, N. Wermes²⁴, M. Wessels^{61a}, T.D. Weston²⁰, K. Whalen¹³¹, A.M. Wharton⁹⁰, A.S. White¹⁰⁶,
A. White⁸, M.J. White¹, D. Whiteson¹⁷¹, B.W. Whitmore⁹⁰, W. Wiedenmann¹⁸¹, C. Wiel⁴⁸, M. Wielers¹⁴³,
N. Wieseotte¹⁰⁰, C. Wiglesworth⁴⁰, L.A.M. Wiik-Fuchs⁵², H.G. Wilkens³⁶, L.J. Wilkins⁹⁴,
D.M. Williams³⁹, H.H. Williams¹³⁶, S. Williams³², S. Willocq¹⁰³, P.J. Windischhofer¹³⁴,
I. Wingerter-Seez⁵, E. Winkels¹⁵⁶, F. Winklmeier¹³¹, B.T. Winter⁵², M. Wittgen¹⁵³, M. Wobisch⁹⁶,
A. Wolf¹⁰⁰, R. Wölker¹³⁴, J. Wollrath⁵², M.W. Wolter⁸⁵, H. Wolters^{139a,139c}, V.W.S. Wong¹⁷⁵,
A.F. Wongel⁴⁶, N.L. Woods¹⁴⁵, S.D. Worm⁴⁶, B.K. Wosiek⁸⁵, K.W. Woźniak⁸⁵, K. Wraight⁵⁷, S.L. Wu¹⁸¹,
X. Wu⁵⁴, Y. Wu^{60a}, J. Wuerzinger¹³⁴, T.R. Wyatt¹⁰¹, B.M. Wynne⁵⁰, S. Xella⁴⁰, L. Xia¹⁷⁸, J. Xiang^{63c},
X. Xiao¹⁰⁶, X. Xie^{60a}, I. Xiotidis¹⁵⁶, D. Xu^{15a}, H. Xu^{60a}, H. Xu^{60a}, L. Xu²⁹, R. Xu¹³⁶, T. Xu¹⁴⁴, W. Xu¹⁰⁶,
Y. Xu^{15b}, Z. Xu^{60b}, Z. Xu¹⁵³, B. Yabsley¹⁵⁷, S. Yacoob^{33a}, D.P. Yallup⁹⁵, N. Yamaguchi⁸⁸,
Y. Yamaguchi¹⁶⁵, A. Yamamoto⁸², M. Yamatani¹⁶³, T. Yamazaki¹⁶³, Y. Yamazaki⁸³, J. Yan^{60c}, Z. Yan²⁵,
H.J. Yang^{60c,60d}, H.T. Yang¹⁸, S. Yang^{60a}, T. Yang^{63c}, X. Yang^{60a}, X. Yang^{60b,58}, Y. Yang¹⁶³, Z. Yang^{60a},
W-M. Yao¹⁸, Y.C. Yap⁴⁶, H. Ye^{15c}, J. Ye⁴², S. Ye²⁹, I. Yeletskikh⁸⁰, M.R. Yexley⁹⁰, E. Yigitbasi²⁵,
P. Yin³⁹, K. Yorita¹⁷⁹, K. Yoshihara⁷⁹, C.J.S. Young³⁶, C. Young¹⁵³, J. Yu⁷⁹, R. Yuan^{60b,i}, X. Yue^{61a},
M. Zaazoua^{35e}, B. Zabinski⁸⁵, G. Zacharis¹⁰, E. Zaffaroni⁵⁴, J. Zahreddine¹³⁵, A.M. Zaitsev^{123,ae},
T. Zakareishvili^{159b}, N. Zakharchuk³⁴, S. Zambito³⁶, D. Zanzi³⁶, S.V. Zeißner⁴⁷, C. Zeitnitz¹⁸²,
G. Zemaityte¹³⁴, J.C. Zeng¹⁷³, O. Zenin¹²³, T. Ženiš^{28a}, D. Zerwas⁶⁵, M. Zgubić¹³⁴, B. Zhang^{15c},
D.F. Zhang^{15b}, G. Zhang^{15b}, J. Zhang⁶, Kaili. Zhang^{15a}, L. Zhang^{15c}, L. Zhang^{60a}, M. Zhang¹⁷³,
R. Zhang¹⁸¹, S. Zhang¹⁰⁶, X. Zhang^{60c}, X. Zhang^{60b}, Y. Zhang^{15a,15d}, Z. Zhang^{63a}, Z. Zhang⁶⁵, P. Zhao⁴⁹,
Y. Zhao¹⁴⁵, Z. Zhao^{60a}, A. Zhemchugov⁸⁰, Z. Zheng¹⁰⁶, D. Zhong¹⁷³, B. Zhou¹⁰⁶, C. Zhou¹⁸¹, H. Zhou⁷,
M. Zhou¹⁵⁵, N. Zhou^{60c}, Y. Zhou⁷, C.G. Zhu^{60b}, C. Zhu^{15a,15d}, H.L. Zhu^{60a}, H. Zhu^{15a}, J. Zhu¹⁰⁶,
Y. Zhu^{60a}, X. Zhuang^{15a}, K. Zhukov¹¹¹, V. Zhulanov^{122b,122a}, D. Zieminska⁶⁶, N.I. Zimine⁸⁰,
S. Zimmermann⁵², Z. Zinonos¹¹⁵, M. Ziolkowski¹⁵¹, L. Živković¹⁶, G. Zobernig¹⁸¹, A. Zoccoli^{23b,23a},
K. Zoch⁵³, T.G. Zorbass¹⁴⁹, R. Zou³⁷, L. Zwalinski³⁶.

- ¹Department of Physics, University of Adelaide, Adelaide; Australia.
- ²Physics Department, SUNY Albany, Albany NY; United States of America.
- ³Department of Physics, University of Alberta, Edmonton AB; Canada.
- ⁴(^a)Department of Physics, Ankara University, Ankara; (^b)Istanbul Aydin University, Application and Research Center for Advanced Studies, Istanbul; (^c)Division of Physics, TOBB University of Economics and Technology, Ankara; Turkey.
- ⁵LAPP, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.
- ⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.
- ⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.
- ⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.
- ⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.
- ¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.
- ¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.
- ¹²(^a)Bahcesehir University, Faculty of Engineering and Natural Sciences, Istanbul; (^b)Istanbul Bilgi University, Faculty of Engineering and Natural Sciences, Istanbul; (^c)Department of Physics, Bogazici University, Istanbul; (^d)Department of Physics Engineering, Gaziantep University, Gaziantep; Turkey.
- ¹³Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- ¹⁴Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.
- ¹⁵(^a)Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (^b)Physics Department, Tsinghua University, Beijing; (^c)Department of Physics, Nanjing University, Nanjing; (^d)University of Chinese Academy of Science (UCAS), Beijing; China.
- ¹⁶Institute of Physics, University of Belgrade, Belgrade; Serbia.
- ¹⁷Department for Physics and Technology, University of Bergen, Bergen; Norway.
- ¹⁸Physics Division, Lawrence Berkeley National Laboratory and University of California, Berkeley CA; United States of America.
- ¹⁹Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.
- ²⁰Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.
- ²¹School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.
- ²²(^a)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (^b)Departamento de Física, Universidad Nacional de Colombia, Bogotá, Colombia; Colombia.
- ²³(^a)INFN Bologna and Università di Bologna, Dipartimento di Fisica; (^b)INFN Sezione di Bologna; Italy.
- ²⁴Physikalisches Institut, Universität Bonn, Bonn; Germany.
- ²⁵Department of Physics, Boston University, Boston MA; United States of America.
- ²⁶Department of Physics, Brandeis University, Waltham MA; United States of America.
- ²⁷(^a)Transilvania University of Brasov, Brasov; (^b)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (^c)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (^d)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (^e)University Politehnica Bucharest, Bucharest; (^f)West University in Timisoara, Timisoara; Romania.
- ²⁸(^a)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (^b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.
- ²⁹Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.
- ³⁰Departamento de Física, Universidad de Buenos Aires, Buenos Aires; Argentina.
- ³¹California State University, CA; United States of America.

- ³²Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.
- ³³(^a)Department of Physics, University of Cape Town, Cape Town; (^b)iThemba Labs, Western Cape; (^c)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (^d)University of South Africa, Department of Physics, Pretoria; (^e)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.
- ³⁴Department of Physics, Carleton University, Ottawa ON; Canada.
- ³⁵(^a)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (^b)Faculté des Sciences, Université Ibn-Tofail, Kénitra; (^c)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (^d)Faculté des Sciences, Université Mohamed Premier and LPTPM, Oujda; (^e)Faculté des sciences, Université Mohammed V, Rabat; Morocco.
- ³⁶CERN, Geneva; Switzerland.
- ³⁷Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.
- ³⁸LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.
- ³⁹Nevis Laboratory, Columbia University, Irvington NY; United States of America.
- ⁴⁰Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.
- ⁴¹(^a)Dipartimento di Fisica, Università della Calabria, Rende; (^b)INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy.
- ⁴²Physics Department, Southern Methodist University, Dallas TX; United States of America.
- ⁴³Physics Department, University of Texas at Dallas, Richardson TX; United States of America.
- ⁴⁴National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.
- ⁴⁵(^a)Department of Physics, Stockholm University; (^b)Oskar Klein Centre, Stockholm; Sweden.
- ⁴⁶Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.
- ⁴⁷Lehrstuhl für Experimentelle Physik IV, Technische Universität Dortmund, Dortmund; Germany.
- ⁴⁸Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.
- ⁴⁹Department of Physics, Duke University, Durham NC; United States of America.
- ⁵⁰SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.
- ⁵¹INFN e Laboratori Nazionali di Frascati, Frascati; Italy.
- ⁵²Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.
- ⁵³II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.
- ⁵⁴Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ⁵⁵(^a)Dipartimento di Fisica, Università di Genova, Genova; (^b)INFN Sezione di Genova; Italy.
- ⁵⁶II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.
- ⁵⁷SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.
- ⁵⁸LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.
- ⁵⁹Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.
- ⁶⁰(^a)Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; (^b)Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; (^c)School of Physics and Astronomy, Shanghai Jiao Tong University, KLPPAC-MoE, SKLPPC, Shanghai; (^d)Tsung-Dao Lee Institute, Shanghai; China.
- ⁶¹(^a)Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; (^b)Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.
- ⁶²Faculty of Applied Information Science, Hiroshima Institute of Technology, Hiroshima; Japan.
- ⁶³(^a)Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; (^b)Department of Physics, University of Hong Kong, Hong Kong; (^c)Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.

- ⁶⁴Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.
- ⁶⁵IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.
- ⁶⁶Department of Physics, Indiana University, Bloomington IN; United States of America.
- ⁶⁷(^a)INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; (^b)ICTP, Trieste; (^c)Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.
- ⁶⁸(^a)INFN Sezione di Lecce; (^b)Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.
- ⁶⁹(^a)INFN Sezione di Milano; (^b)Dipartimento di Fisica, Università di Milano, Milano; Italy.
- ⁷⁰(^a)INFN Sezione di Napoli; (^b)Dipartimento di Fisica, Università di Napoli, Napoli; Italy.
- ⁷¹(^a)INFN Sezione di Pavia; (^b)Dipartimento di Fisica, Università di Pavia, Pavia; Italy.
- ⁷²(^a)INFN Sezione di Pisa; (^b)Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.
- ⁷³(^a)INFN Sezione di Roma; (^b)Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.
- ⁷⁴(^a)INFN Sezione di Roma Tor Vergata; (^b)Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.
- ⁷⁵(^a)INFN Sezione di Roma Tre; (^b)Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.
- ⁷⁶(^a)INFN-TIFPA; (^b)Università degli Studi di Trento, Trento; Italy.
- ⁷⁷Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität, Innsbruck; Austria.
- ⁷⁸University of Iowa, Iowa City IA; United States of America.
- ⁷⁹Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.
- ⁸⁰Joint Institute for Nuclear Research, Dubna; Russia.
- ⁸¹(^a)Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; (^b)Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; (^c)Instituto de Física, Universidade de São Paulo, São Paulo; Brazil.
- ⁸²KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.
- ⁸³Graduate School of Science, Kobe University, Kobe; Japan.
- ⁸⁴(^a)AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; (^b)Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.
- ⁸⁵Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.
- ⁸⁶Faculty of Science, Kyoto University, Kyoto; Japan.
- ⁸⁷Kyoto University of Education, Kyoto; Japan.
- ⁸⁸Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.
- ⁸⁹Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.
- ⁹⁰Physics Department, Lancaster University, Lancaster; United Kingdom.
- ⁹¹Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.
- ⁹²Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.
- ⁹³School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.
- ⁹⁴Department of Physics, Royal Holloway University of London, Egham; United Kingdom.
- ⁹⁵Department of Physics and Astronomy, University College London, London; United Kingdom.
- ⁹⁶Louisiana Tech University, Ruston LA; United States of America.
- ⁹⁷Fysiska institutionen, Lunds universitet, Lund; Sweden.
- ⁹⁸Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules (IN2P3), Villeurbanne; France.
- ⁹⁹Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.
- ¹⁰⁰Institut für Physik, Universität Mainz, Mainz; Germany.
- ¹⁰¹School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.

- ¹⁰²CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.
- ¹⁰³Department of Physics, University of Massachusetts, Amherst MA; United States of America.
- ¹⁰⁴Department of Physics, McGill University, Montreal QC; Canada.
- ¹⁰⁵School of Physics, University of Melbourne, Victoria; Australia.
- ¹⁰⁶Department of Physics, University of Michigan, Ann Arbor MI; United States of America.
- ¹⁰⁷Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ¹⁰⁸B.I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk; Belarus.
- ¹⁰⁹Research Institute for Nuclear Problems of Byelorussian State University, Minsk; Belarus.
- ¹¹⁰Group of Particle Physics, University of Montreal, Montreal QC; Canada.
- ¹¹¹P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow; Russia.
- ¹¹²National Research Nuclear University MEPhI, Moscow; Russia.
- ¹¹³D.V. Skobel'syn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University, Moscow; Russia.
- ¹¹⁴Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.
- ¹¹⁵Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.
- ¹¹⁶Nagasaki Institute of Applied Science, Nagasaki; Japan.
- ¹¹⁷Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.
- ¹¹⁸Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.
- ¹¹⁹Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen; Netherlands.
- ¹²⁰Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.
- ¹²¹Department of Physics, Northern Illinois University, DeKalb IL; United States of America.
- ¹²²^(a)Budker Institute of Nuclear Physics and NSU, SB RAS, Novosibirsk; ^(b)Novosibirsk State University Novosibirsk; Russia.
- ¹²³Institute for High Energy Physics of the National Research Centre Kurchatov Institute, Protvino; Russia.
- ¹²⁴Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of National Research Centre "Kurchatov Institute", Moscow; Russia.
- ¹²⁵Department of Physics, New York University, New York NY; United States of America.
- ¹²⁶Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.
- ¹²⁷Ohio State University, Columbus OH; United States of America.
- ¹²⁸Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.
- ¹²⁹Department of Physics, Oklahoma State University, Stillwater OK; United States of America.
- ¹³⁰Palacký University, RCPTM, Joint Laboratory of Optics, Olomouc; Czech Republic.
- ¹³¹Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.
- ¹³²Graduate School of Science, Osaka University, Osaka; Japan.
- ¹³³Department of Physics, University of Oslo, Oslo; Norway.
- ¹³⁴Department of Physics, Oxford University, Oxford; United Kingdom.
- ¹³⁵LPNHE, Sorbonne Université, Université de Paris, CNRS/IN2P3, Paris; France.
- ¹³⁶Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.
- ¹³⁷Konstantinov Nuclear Physics Institute of National Research Centre "Kurchatov Institute", PNPI, St. Petersburg; Russia.
- ¹³⁸Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.

- ¹³⁹(*a*) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; (*b*) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; (*c*) Departamento de Física, Universidade de Coimbra, Coimbra; (*d*) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; (*e*) Departamento de Física, Universidade do Minho, Braga; (*f*) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); (*g*) Dep Física and CEFITEC of Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Caparica; (*h*) Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.
- ¹⁴⁰Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.
- ¹⁴¹Czech Technical University in Prague, Prague; Czech Republic.
- ¹⁴²Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.
- ¹⁴³Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.
- ¹⁴⁴IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.
- ¹⁴⁵Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.
- ¹⁴⁶(*a*) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; (*b*) Universidad Andres Bello, Department of Physics, Santiago; (*c*) Instituto de Alta Investigación, Universidad de Tarapacá; (*d*) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.
- ¹⁴⁷Universidade Federal de São João del Rei (UFSJ), São João del Rei; Brazil.
- ¹⁴⁸Department of Physics, University of Washington, Seattle WA; United States of America.
- ¹⁴⁹Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.
- ¹⁵⁰Department of Physics, Shinshu University, Nagano; Japan.
- ¹⁵¹Department Physik, Universität Siegen, Siegen; Germany.
- ¹⁵²Department of Physics, Simon Fraser University, Burnaby BC; Canada.
- ¹⁵³SLAC National Accelerator Laboratory, Stanford CA; United States of America.
- ¹⁵⁴Physics Department, Royal Institute of Technology, Stockholm; Sweden.
- ¹⁵⁵Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- ¹⁵⁶Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- ¹⁵⁷School of Physics, University of Sydney, Sydney; Australia.
- ¹⁵⁸Institute of Physics, Academia Sinica, Taipei; Taiwan.
- ¹⁵⁹(*a*) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; (*b*) High Energy Physics Institute, Tbilisi State University, Tbilisi; Georgia.
- ¹⁶⁰Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- ¹⁶¹Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- ¹⁶²Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- ¹⁶³International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- ¹⁶⁴Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo; Japan.
- ¹⁶⁵Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- ¹⁶⁶Tomsk State University, Tomsk; Russia.
- ¹⁶⁷Department of Physics, University of Toronto, Toronto ON; Canada.
- ¹⁶⁸(*a*) TRIUMF, Vancouver BC; (*b*) Department of Physics and Astronomy, York University, Toronto ON; Canada.
- ¹⁶⁹Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- ¹⁷⁰Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- ¹⁷¹Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of

America.

¹⁷²Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.

¹⁷³Department of Physics, University of Illinois, Urbana IL; United States of America.

¹⁷⁴Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.

¹⁷⁵Department of Physics, University of British Columbia, Vancouver BC; Canada.

¹⁷⁶Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.

¹⁷⁷Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.

¹⁷⁸Department of Physics, University of Warwick, Coventry; United Kingdom.

¹⁷⁹Waseda University, Tokyo; Japan.

¹⁸⁰Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.

¹⁸¹Department of Physics, University of Wisconsin, Madison WI; United States of America.

¹⁸²Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.

¹⁸³Department of Physics, Yale University, New Haven CT; United States of America.

^a Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.

^b Also at Center for High Energy Physics, Peking University; China.

^c Also at Centro Studi e Ricerche Enrico Fermi; Italy.

^d Also at CERN, Geneva; Switzerland.

^e Also at CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.

^f Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

^g Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain.

^h Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.

ⁱ Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.

^j Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America.

^k Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel.

^l Also at Department of Physics, California State University, East Bay; United States of America.

^m Also at Department of Physics, California State University, Fresno; United States of America.

ⁿ Also at Department of Physics, California State University, Sacramento; United States of America.

^o Also at Department of Physics, King's College London, London; United Kingdom.

^p Also at Department of Physics, St. Petersburg State Polytechnical University, St. Petersburg; Russia.

^q Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.

^r Also at Dipartimento di Matematica, Informatica e Fisica, Università di Udine, Udine; Italy.

^s Also at Faculty of Physics, M.V. Lomonosov Moscow State University, Moscow; Russia.

^t Also at Giresun University, Faculty of Engineering, Giresun; Turkey.

^u Also at Graduate School of Science, Osaka University, Osaka; Japan.

^v Also at Hellenic Open University, Patras; Greece.

^w Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.

^x Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.

^y Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria.

^z Also at Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Budapest; Hungary.

^{aa} Also at Institute of Particle Physics (IPP); Canada.

- ab* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- ac* Also at Instituto de Fisica Teorica, IFT-UAM/CSIC, Madrid; Spain.
- ad* Also at Joint Institute for Nuclear Research, Dubna; Russia.
- ae* Also at Moscow Institute of Physics and Technology State University, Dolgoprudny; Russia.
- af* Also at National Research Nuclear University MPhI, Moscow; Russia.
- ag* Also at Physics Department, An-Najah National University, Nablus; Palestine.
- ah* Also at Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.
- ai* Also at The City College of New York, New York NY; United States of America.
- aj* Also at TRIUMF, Vancouver BC; Canada.
- ak* Also at Università di Napoli Parthenope, Napoli; Italy.
- al* Also at University of Chinese Academy of Sciences (UCAS), Beijing; China.
- * Deceased