

JA

CONSIGLIO NAZIONALE DELLE RICERCHE

ISTITUTO DI COSMO-GEOFISICA

TORINO

**Observation of electromagnetic
interactions of high energy muons
deep underground**

C.Castagnoli, A.Castellina, O.Saavedra,
T.M.Kirina, R.P. Kokoulin, A.A.Petrukhin

Rap. int. n.310/1994

CERN LIBRARIES, GENEVA



SCAN-9502040

529506

Submitted to: *Physical Review D*

Observation of electromagnetic interactions of high energy muons deep underground

C.Castagnoli, A.Castellina, O.Saavedra
*Istituto di Cosmogeofisica del CNR, cs.Fiume 4, 10133 Torino
e Istituto di Fisica Generale dell'Universita' di Torino, Italy.*

T.M.Kirina, R.P.Kokoulin, A.A.Petrukhin,
Moscow Engineering Physics Institute, 115409 Moscow, Russia.

The first experimental measurement of the spectrum of cascade showers produced by cosmic ray muon interactions deep underground in the transferred energy range 0.4 - 200 GeV has been performed by means of the NUSEX calorimeter, at a depth of about 5000 hg/cm², the average muon energy being close to 360 GeV. The spectrum of energy transfers is in good agreement with calculations based on the commonly used theoretical formulae for muon interaction cross sections and on a conservative picture of the formation of muon energy spectrum at great depths. The present data do not confirm the deficit of the cross section at low energy transfers recently reported by the CCFR collaboration [1]. A possible explanation of the difference between the observed and the expected stochastic energy loss distributions presented by the CCFR group is discussed.

PACS number(s): 95.55.Vj, 95.85.Ry, 13.10+q

I. INTRODUCTION

A correct knowledge of the muon interaction cross section and energy loss rate is important for the interpretation of experimental results and the planning of new experiments both at accelerators and in cosmic rays. The attention to this problem is also raised by some evidences for a deviation of the cross section from the commonly used theoretical formulae. For example, in the recently published results of the CCFR collaboration [1], the observed stochastic energy loss distribution for high muon energies (around 1 TeV) at low energy transfers appears considerably lower than the expected one, indicating a 30% deficit of the cross section in comparison with the theoretical prediction at relative

transfers $\nu \simeq 10^{-3}$. Only few measurements of the electromagnetic interaction cross sections for high energy muons exist; because of the absence of high energy muon beams at accelerators and of a rather slow progress in their creation, most of these experiments have been performed by studying muons in the cosmic ray flux.

A comprehensive survey of the different techniques used for muon cross section investigations in cosmic rays and a critical analysis of the earlier experimental results may be found in [2]. Two methods are most widely applied to check the energy dependence of the muon interaction cross section. One of them is based on the use of combined arrangements including a target calorimeter for the detection of the interactions and a magnetic spectrometer for the muon momentum evaluation. However, this approach requires the creation of powerful spectrometers, which necessary sizes grow too fast with increasing muon energy. Because of this, the second method is most often used, requiring only detectors for muon interactions. In this case, the transferred energy spectrum, which is the convolution of the differential cross section $\sigma(E, \epsilon)$ with the muon energy spectrum at the detector site, is measured. Conclusions about the cross section behaviour are obtained by comparing the observed and the expected interaction rates, presuming a sufficiently accurate knowledge of the local muon spectrum, which is either taken from independent measurements or calculated. Earlier experiments of this kind were performed at shallow depths underground corresponding to average muon energies below 10^2 GeV.

Deep underground investigations of muon interactions are of special interest. Owing to the features of the energy dependence of cosmic ray muon flux and muon interaction in rock, the differential muon spectrum at large depths is essentially flat at low energies (up to several tens of GeV) and is largely enriched by high energy muons, the average local muon energy approaching 300 - 400 GeV. Besides a high value of the average energy, the deep underground muon spectrum is characterized by a rather weak dependence of the spectrum shape on the depth, which greatly simplifies calculations of the expected spectrum of muon induced cascades. No direct measurements of the muon energy spectrum at such depths exist; therefore, in the muon interaction analysis one has to use the spectrum obtained by calculating the cosmic ray muon flux penetration through a large thickness of rock assuming an *a priori* knowledge of the cross sections. Because of this, some logical contradiction appears here: to estimate the cross section, we need to know it in advance. Luckily, the shape of the local muon energy spectrum is mainly determined by the muon spectrum slope at the surface, by the total muon energy loss and (to a lower extent, through the fluctuation effect) by the cross section energy dependence at large energy transfers ($\nu \simeq 1$), but it does practically not depend on the details of the differential cross section behaviour at low relative energy

transfers ($v \ll 1$), thus leaving us the possibility to consistently check the interaction cross section in this domain.

In the present paper, the results on the first deep underground study of the transferred energy spectrum are reported. Electromagnetic interactions of high energy muons in the NUSEX detector [3] were analysed. The apparatus was located at a depth of about 5000 hg/cm² underground, the average muon energy being close to 360 GeV. The fine structure of the detector allowed to detect cascades with energies as low as 0.4 GeV. The experimental data collected during 1982 - 1988 have been used in the analysis.

II. EXPERIMENTAL ARRANGEMENT

The NUSEX detector is a digital tracking calorimeter, consisting of 134 horizontal iron plates 3.5 m x 3.5 m x 1 cm thick, interleaved by 3.5 m long plastic streamer tubes with 9 x 9 mm² inner cross section. The detector height is 3.7 m and the total mass ~150 t. The average mass composition of the calorimeter is given by 90% iron and 10% PVC. The average layer thickness (including chamber material and spacers) equals to 0.62 radiation lengths (r.l.). The tube chamber planes are equipped with a two-dimensional external strip readout system. X-coordinate pickup strips are parallel to the tubes, 0.4 cm wide and positioned in correspondence of each tube wire (1 cm pitch). Y-strips (orthogonal to wires) are 1 cm wide and have 1.2 cm pitch. The detection efficiency for charged particles crossing a tube is greater than 98 %. The multihit probability (i.e. the probability to detect an induced signal on a strip adjacent to the hit tube) was estimated by muon track analysis and amounted to approximately 7% for X-view, while for Y-strips typically 2 or 3 hits per plane for a single muon track were detected. Because of this, only the X-strips information has been considered in the present analysis for the quantitative description of the showers, while the Y-strips data were only used for the geometrical reconstruction of muon tracks. The detector was operated with very soft trigger conditions, thus providing a registration efficiency for cosmic ray muons close to 100%. A test module of the NUSEX detector [3], having almost identical structure, was calibrated at the CERN-PS accelerator and the response of the calorimeter for electromagnetic cascade showers was measured up to incident electron energies 1.2 GeV.

The NUSEX detector was installed in the Mont Blanc tunnel; the rock thickness exceeded 4800 hg/cm² in every direction. The average composition of the surrounding rock is characterised by the following parameters: $\langle Z/A \rangle = 0.494$, $\langle Z^2/A \rangle = 5.12$, $\rho = 2.68$ g/cm³. NUSEX was a multipurpose detector: while its main goal was the study of nucleon stability, important results have been reached in the investigations of primary cosmic ray composition and spectrum and detection of muon

signals from celestial sources; neutrino oscillation, ν_e/ν_μ - ratio and dark matter studies were also performed [4-8].

III. MUON SELECTION

In a preliminary step, all events with at least one straight penetrating track crossing any 50 planes of the calorimeter have been selected from raw data. These data have been used in order to study the detector performances in operation. The time intervals of improper operation for some parts of the detector and the channels often producing false signals could be recognised and excluded on the basis of these considerations. Typically, the number of such channels did not exceed 1% of the total. Examples of muon events (X-view) registered by the NUSEX detector are given in Fig.1. Only muons crossing at least 90 consecutive planes (2/3 of the detector height) with projected zenith angles $|\theta_{x,y}| \leq 40^\circ$ have been used for the physical analysis. Muon tracks appreciably distorted by multiple scattering have been rejected (as an example, an end-of-range track is shown in Fig.1a). For this purpose, the linear fit of the entire track has been compared with the partial fits of two halves of the track. The track was excluded if the maximum deviation between these fits exceeded 0.8 cm in the X-view or 2.0 cm in the Y-view. The Monte Carlo simulation of multiple scattering of muons in the detector showed that this selection corresponded to an effective cutoff energy $E_{\min} \sim 4$ GeV. The total amount of rejected scattered tracks (about 2 %) is in reasonable agreement with the expected muon spectrum at the observation depth. Besides single muons, muon bundle events have been observed (Fig.1b). Individual muons from these bundles, satisfying the above selection criteria, have also been included into the analysis. In all, about 1.7×10^4 muon tracks have been selected for the present muon interaction study.

IV. SELECTION OF CASCADE SHOWERS

An automatic scanning procedure has been applied to find cascade showers initiated by muon interactions. Only hits detected at projected distances $D_x > 1.0$ cm from the track fit have been used for cascade recognition and energy estimation. The average background caused by induced signals from muon track and by secondary particles produced in low energy interactions was estimated from experimental data and amounted to 0.07 hits per plane for $D_x > 1.0$ cm. An example of the operation of the scanning routine (for the event presented in Fig.1c) is given in Fig.2. First, the number of hits in each plane within a road $D_x = 1-15$ cm along the track is calculated (upper histogram in the figure). Then, the running sum of the numbers of hits in 7 consecutive planes is

computed (lower histogram), thus producing a smoothed longitudinal profile of the event with an averaging window of about 4 radiation lengths. Two showers are considered separated if the minimum observed in the smoothed profile is less than a half of each of the corresponding maxima. Only showers with total number of hits outside the muon track $n_h \geq 9$ have been considered. To suppress the background, only structures with more than 2 hit planes have been taken into account.

Besides cascades due to purely electromagnetic interaction processes (knock-on electrons, direct electron-positron pair production, bremsstrahlung process), nuclear - electromagnetic showers have been recorded during the experiment (Fig.1d). Nuclear showers show a different lateral structure in comparison with electromagnetic cascades and a correct estimation of their energies is not simple. On the other hand, the expected rate of nuclear interactions is small at low transferred energies: less than 2% of the total at < 5 GeV. Furthermore, nuclear showers with energies exceeding several GeV are easily recognised due to the presence of penetrating tracks (shower hadrons - Fig.1d). In the present analysis, we chose to exclude nuclear showers on the basis of a visual scanning of the events. Similar problems with the energy estimation arise in the case of cascades originated outside the detector (Fig.1e) or near its edges. To exclude these events, only showers centred within the inner volume of the calorimeter (more than 20 cm from the sides, and planes from 14 to 119) have been selected, so defining a fiducial target mass of about 90 ton. Finally, about 9200 reconstructed showers with $n_h \geq 9$ have been included in the analysis.

V. CALCULATION OF THE EXPECTED DISTRIBUTION

The calculation of the expected shower distribution is simplified by the following circumstances. Firstly, the muon energy practically does not change in the calorimeter and separate interactions may be considered as independent, the total number of interactions being proportional to the muon path in the fiducial volume. Secondly, at great depths the shape of the muon energy spectrum only slightly varies with depth, which allows us to use a single energy distribution function averaged in the corresponding zenith angle interval. In these approximations the interaction rate per muon and per unit thickness is

$$N_{\text{int}}(\epsilon) = \int_{E_{\text{min}}}^{\infty} F_{\mu}(E) \sigma(E, \epsilon) dE , \quad (1)$$

where $F_{\mu}(E)$ is the normalised muon energy distribution, and $\sigma(E,\epsilon)$ is the interaction cross section. The expected shower distribution in the number of hits is given by

$$N_{\text{exp}}(n_h) = X_{\text{tot}} \int_0^{\infty} P(\epsilon, n_h) N_{\text{int}}(\epsilon) d\epsilon . \quad (2)$$

Here $X_{\text{tot}} = 4.4 \times 10^6$ cm (1.48×10^7 g/cm² in NUSEX material) is the total track length of selected muons within the fiducial volume, which is determined directly from the geometry reconstruction of the observed events; $P(\epsilon, n_h)$ is the probability to observe n_h hits in a shower with energy ϵ , which is assumed to follow a Poisson distribution with an average value $\langle n_h \rangle$ defined by the corresponding calibration curve (see below).

VI. CALIBRATION CURVES FOR MUON INITIATED CASCADE SHOWERS

As we mentioned above, the test module of the NUSEX detector was calibrated at the accelerator electron beam [3]. However, some difficulties appear in the muon interaction study in a digital calorimeter when the influence of muon track on the shower pattern near the axis has to be taken into account. Because of this, only hits detected at some distances from the track may be used for the shower energy estimation, while the electron calibration gives only the total number of hits corresponding to a given shower energy. Moreover, the detector calibration was performed only in the energy region interesting for proton decay searches, that is up to electron energies 1.2 GeV. To extrapolate the calibration results to higher energies and to various distances from the muon track, we use here a simple analytical model [9] (Appendix A) describing the average behaviour of a shower in a digital calorimeter. The model contains two free fitting parameters (the effective values of the Moliere unit r'_M and of the electron critical energy β') which characterise the shower development in the real detector structure and may be directly estimated from the experimental data (electron beam calibration results and lateral distributions of hits in the registered cascades).

The experimental projected lateral distribution functions (average number of hits as a function of distance from the shower axis) have been compared with the best fit model calculations for 4 intervals in the number of hits in reconstructed cascades, corresponding to shower energies from 1 to 8 GeV; they are shown in Fig.3. The agreement between experimental data and calculations seems good, with the exception of large distances at the lowest energy. This region ($D_x \geq 10$ cm) gives however only a minor contribution to the total number of hits.

The results of the accelerator electron beam calibration experiment are given in Fig.4 (squares) together with the best-fit model curve. The lower region in the figure represents the model prediction for the number of hits in the distance range $D_x = 1 - 15$ cm, the one used in the present analysis. The uncertainty in the latter calibration curve is mainly due to variations in the functions used in the analytical model and to ambiguities in the parameters, arising from the not full identity of the measurement conditions in the NUSEX detector and in the test module exposed to the accelerator beam (see Appendix A).

VII. LOCAL MUON ENERGY SPECTRUM

The underground muon energy spectrum has been calculated both analytically and numerically by various authors. The main features of the underground spectrum may be illustrated on the basis of the analytical solution assuming the constancy of the coefficients in the muon energy loss relation

$$\langle dE/dx \rangle = a + bE \quad (3)$$

and neglecting energy loss fluctuations. For a power type differential muon energy spectrum at the surface

$$N_\mu(E,0) = B E^{-(\gamma+1)} \quad (4)$$

one can easily derive the energy spectrum at the depth h :

$$N_\mu(E,h) = B \exp(-\gamma bh) \left\{ E + \frac{a}{b} [1 - \exp(-bh)] \right\}^{-(\gamma+1)}. \quad (5)$$

Here, the first exponent describes the attenuation of the high energy muon flux, whereas the factor in braces determines the shape of the local energy spectrum. The average underground muon energy is given by

$$E_{av} = \frac{a}{b(\gamma-1)} [1 - \exp(-bh)]. \quad (6)$$

The energy spectrum is essentially flat at energies $E \ll E_{av}$, and reproduces the slope of the surface spectrum at $E \gg E_{av}$. At shallow depths ($bh \ll 1$), the average muon energy is nearly proportional to h ; at great depths, which are usually defined by the condition $bh > 1$ (or $h > 2 \times 10^5$ g/cm² in rock), the exponent in the square brackets of eqs.(5) and (6) becomes small, and the spectrum shape only slowly

varies with the rock thickness, the average energy approaching a limit $a/(b(\gamma-1))$.

A more rigorous approach to the muon transport problem must allow for fluctuations in the energy losses due to rare interactions with high relative energy transfers, mostly in the processes of muon bremsstrahlung and nuclear interaction. In fact, the muon flux at great depths is formed mainly by particles evading these catastrophic collisions. Zatsepin and Michal'chi [10] found that the influence of the fluctuations may be taken into account by the introduction of some effective energy loss coefficients ($b_{ef}, b'_{ef} < b$) instead of the value determining the average energy loss rate (3). In the first approximation, the underground spectrum may be expressed as

$$N_{\mu}(E, h) = B \exp(-\gamma b_{ef} h) \left\{ E + \frac{a}{b'_{ef}} \left[1 - \exp(-b'_{ef} h) \right] \right\}^{-(\gamma+1)}. \quad (7)$$

The comparison between (5) and (7) shows that the energy loss fluctuations lead to a slower absorption of the muon flux and to an increase of the average muon energy deep underground.

The role of different relative energy transfer regions in the fluctuation effect on the muon flux penetration may be clarified if we confront the expressions for b and b_{ef} . Following Zatsepin and Michal'chi approach, we may write

$$b = \int_0^1 \sigma(v) v dv ; \quad b_{ef} = \int_0^1 \sigma(v) \left[\frac{1 - (1-v)^\gamma}{\gamma} \right] dv. \quad (8)$$

Here $\sigma(v)$ is the cross section (in cm^2/g) for muon interactions in the processes of muon bremsstrahlung, nuclear interaction and pair production. At high muon energies, this cross section depends only on the relative energy transfer v . Comparison of the formulae for b and b_{ef} shows that only the cross section dependence near $v \simeq 1$ is important for the fluctuation problem, whereas in the low relative transfer region $v \ll 1$ only the integral energy loss is essential. Considerations about the value of b'_{ef} are somewhat more complicated but lead to the same conclusion.

Quantitatively, here we rely on the results of local muon spectrum calculations performed by means of two independent Monte Carlo codes. In one of them [11], direct pair production and ionisation (including high energy knock-on electron production) are treated as continuous energy losses, while polynomial approximations for muon bremsstrahlung and nuclear interaction cross sections based on the accurate expressions [11] (and refs. therein) are used. The second simulation is based on the code

described in [12]. In this case, all muon interactions with energy transfers above $\varepsilon_0 = 1$ GeV are simulated following the energy dependence of the corresponding cross sections, while the lowest energy transfers ($\varepsilon < \varepsilon_0$) are included in the continuous energy losses. The average muon energy losses calculated for standard rock ($Z^2/A = 5.5$, $Z/A = 0.5$) for both programs agree with the commonly used values [13] in the muon energy range 5 - 10000 GeV. The muon energies at the surface have been sampled from a power type spectrum with integral power index $\gamma=2.70$. The average muon energies at different depth underground as calculated by means of the two programs are given in Fig.5 by full and open circles, respectively. The results of calculations based on the semi-analytical solution of the muon transport problem [14] are given for comparison (squares). The differential energy spectra of muons for different depths of standard rock are shown by histograms in Fig.6.

The present Monte Carlo results are well approximated by the analytical expression (7) but with different values of the coefficients (solid curves in Figs.5 and 6). From the least square fit of the calculated muon intensities and average energies for standard rock, in the depth range $(4-10) \times 10^5$ g/cm², we get for the parameters in (7)

$$b_{ef} = 3.81 \times 10^{-6} \text{ cm}^2/\text{g}; \quad b'_{ef} = 3.74 \times 10^{-6} \text{ cm}^2/\text{g}; \quad \frac{a}{b'_{ef}} = 647 \text{ GeV}. \quad (9)$$

The transition to a different rock composition may be performed by accounting for the difference in the average energy losses; approximately, the a-term and the b-term in the energy loss relation are proportional to $\langle Z/A \rangle$ and to $\langle Z(Z+1)/A \rangle$, respectively. Thus, for the Mont Blanc rock we found

$$b_{ef} = 3.59 \times 10^{-6} \text{ cm}^2/\text{g}; \quad b'_{ef} = 3.52 \times 10^{-6} \text{ cm}^2/\text{g}; \quad \frac{a}{b'_{ef}} = 680 \text{ GeV}. \quad (10)$$

Other corrections to the calculated muon spectrum included the averaging over zenith angles (a $\sec\Theta$ -enhancement factor for the surface intensity was assumed) and the correction for the non-asymptotic slope of the muon spectrum at the surface (a more accurate expression for muons originating from π - and K-decays was used). Taking into account the low energy cut-off E_{\min} (Sect. III), the estimated average muon energy for the present experiment equals to 360 GeV. Variations of the parameters used in the muon energy spectrum calculation (5% in depth, 5% and 2% in the values of b- and a-terms in the average energy loss, 0.1 in the spectrum slope around 4 TeV at the surface) correspond to changes in the average muon energy by 2, 3, 2, and 5%, respectively.

VIII. CORRECTIONS

Three kinds of corrections have been introduced in the experimental distributions of the reconstructed showers and/or in calculations.

A. Imitation of cascades by the background

Occasional condensations of background hits caused by induced signals from the muon itself or by low energy secondaries along the track may imitate false cascades in the shower reconstruction procedure. The number of such cases has been estimated via a Monte Carlo simulation based on the experimental distribution of the number of hits (at distances 1 - 15 cm from the track) in isolated planes with "clean environment" (no hits outside the track in 5 nearest planes above and below). The estimated fraction of false cascades is about 10% at $n_h = 9$ and falls off rapidly with n_h , being negligible at $n_h > 15$. The estimated spectrum of these false cascades has been subtracted from the experimental distribution.

B. Overestimation of cascade shower energy

Background hits which occur near the muon track close to the true shower before or after it may be added to the cascade by the shower reconstruction procedure. The average number of such hits per shower may be estimated as a double width of the averaging window multiplied by the number of background hits per plane. Thus, on average, the number of hits in a real shower is overestimated by $\Delta n_h = 2 \times 7 \times 0.07 \simeq 1.0$. The corresponding correction is relatively large owing to the falling spectrum of cascades. It has been taken into account in the calculations of the expected spectrum (2) by substituting the average number of hits $\langle n_h \rangle$ determined by the calibration curve with the corrected increased value $(\langle n_h \rangle + \Delta n_h)$.

C. Correction for shower superposition

At high muon energies, the mean free path for muon interactions is not large in comparison with the typical longitudinal size of the shower. If the distance between two interactions is less than the length of the shower, two cascades overlap each other and cannot be distinguished.

The superposition of the cascades leads to two competing effects ([15], Appendix B): firstly, the loss of some interactions with small energy

transfers, the effect being dominant at relative energy transfers $v < 3 \times 10^{-3}$; secondly, an overestimation of the energy of one of the interactions. In comparison with analog calorimeters (with amplitude measurements), the latter effect is enhanced for digital ones because of the non-linearity of their response, since two overlapping (but shifted in depth) showers on the average produce a larger number of hits than one shower with the same total energy. For the present analysis, the correction for cascade shower superposition has been calculated by means of Monte Carlo technique. Muons have been sampled from the local energy spectrum, and interactions with energy transfers greater than 0.1 GeV have been simulated. The lateral and longitudinal distributions in the number of cascade particles have been calculated using the analytical model (see Appendix A) and then converted into event longitudinal profiles in the number of hits. Finally, the simulated events have been analysed by the standard shower reconstruction routine. The correction factor has been estimated as the ratio of the reconstructed spectra with and without shower superposition. The value of the corresponding correction to the expected cascade distribution is of about -15% at $n_h = 10$, crosses 0 near $n_h = 20 - 25$ and becomes positive at larger n_h (about 16% above $n_h = 80$).

IX. RESULTS AND DISCUSSION

The ratio of the distribution of the reconstructed electromagnetic showers in the number of hits n_h to the expected one is given in Fig.7. All corrections are included. The last point contains 18 showers and corresponds to $n_h > 420$ (cascade shower energies $\epsilon > 220$ GeV). The commonly used theoretical formulae for muon electromagnetic interaction cross sections have been adopted in calculating the expected spectrum: Bhabha expression [16] for knock-on electron production, Petrukhin and Shestakov formulae [17] for the bremsstrahlung process, and Kokoulin and Petrukhin cross section [18] for direct electron pair production. The solid curves in the figure outline the estimated region of systematic uncertainties, which are mainly related to the uncertainty in the energy calibration curve. The observed distribution is in good agreement with the expectation in the entire range of measurements. In Fig.8, the event distribution in the number of hits has been turned into the transferred energy distribution (1). The weighted average energy ϵ corresponding to a certain number of hits was calculated similarly to relation (2). The experimental points in the figure are plotted as deviations from the expectation taking into account all corrections. Horizontal bars indicate the estimated systematic uncertainty in the energy calibration.

The derived transferred energy spectrum agrees well with calculations at $\epsilon \geq 0.4$ GeV. It is important to note that the present data

provide a check of the direct pair production cross section, which contribution to the total generation spectrum of the electromagnetic cascades in the range $\varepsilon = 0.4 - 20$ GeV is greater than 60% (Fig.9). At energy transfers $\varepsilon > 60$ GeV, the muon bremsstrahlung dominates. The observation does not contradict the expected shower rate in this region either (last 4 points in Figs.7 and 8), though statistical errors and systematic uncertainties here are relatively large.

The present results do not confirm the existence of a deficit in the muon interaction rate at low relative energy transfers. At first sight, the comparison of our data with CCFR results [1] seems impossible, since in the present experiment the average muon energy was only 360 GeV, whereas a serious deviation from the theory in [1] (about 30%) was observed only at $\langle E \rangle = 1130$ GeV, the effect being small at $\langle E \rangle = 545$ GeV. However, our data set contains 1.7×10^4 muons above 4 GeV, and, owing to the deep underground location of the setup, about 20% of these muons (or about 3400) have energies exceeding 500 GeV. The CCFR results [1] are based on 9411 cosmic ray muons with energies above 40 GeV, 393 of which having energies greater than 500 GeV (see [1], p.3043). Taking into account the difference in the target thickness (7 times larger in [1]), we can conclude that the statistical power of the two experiments for high energy muon interactions is quite comparable. If we assume the deviation in the ν -dependence of the cross section from the theory similar to that presented in Fig.9 of the CCFR publication [1], the total muon energy loss coefficient b in (3) would be reduced by about 5%, the effective values b_{ef} and b'_{ef} would be lower by 6%. In accordance with Sect. VII, this reduction would lead to an increase of the average muon energy at the NUSEX depth by only 4%, whereas the spectrum of the transferred energies (1) in the interval 0.4 - 2 GeV should appear lower by 13 - 7% (dashed curve in Fig.8) and if the deviation reported in the CCFR paper really exists, our experimental data (in spite of the integration of the effect over the muon energy spectrum) would be sensitive to it. Clearly, our data contradict such a hypothesis. The most probable explanation of the difference between the observed and expected stochastic energy loss distributions in [1] may be a serious underestimation of the distorting influence of cascade shower superposition. The value of the corresponding correction mentioned in that paper was about 5% for 1 TeV muons and 1 GeV energy transfer (see Ref.15 in [1]). Recently performed calculations [15] (Appendix B) gave five times larger effect for these energies, which is in quantitative agreement with the above difference.

X. SUMMARY

The energy spectrum of cascade showers produced by cosmic ray muons in the deep underground detector was measured. In the energy transfer range 0.4 - 200 GeV the observed spectrum of electromagnetic

interactions is in good agreement with calculations based on a conservative picture of the formation of the muon energy spectrum at great depths and of muon interactions in iron. The general consistency of the present results implies that energy transfer spectra for various deep underground and underwater detectors may be calculated on the basis of widely used theoretical formulae for muon interaction cross sections.

Present data do not confirm the deficit of the cross section in the region of low energy transfers reported by the CCFR group [1]. A possible reason for the difference between the observed and the expected distributions of reconstructed cascades in their analysis is the underestimation of the distortions arising from the superposition of showers produced in individual muon interactions.

APPENDIX A: ANALYTICAL MODEL FOR CASCADE SHOWER DESCRIPTION IN A DIGITAL CALORIMETER

A model for calculating the average response of a digital calorimeter for electromagnetic cascade showers has been suggested in [9]. In deriving its basic relations, it is assumed that the counter is discharged every time at least one charged particle crosses its volume. Let $n_e(\epsilon, t, r)$ be the average lateral density of charged particles (electrons and positrons) in a shower with energy ϵ at depth t and distance r from the shower axis. Then the average number of particles crossing the counter at depth t_j and distance x_j may be calculated as

$$n_c(\epsilon, t_j, x_j) = \int_{x_j - \Delta}^{x_j + \Delta} dx \int_{-\infty}^{+\infty} n_e(\epsilon, t_j, r) dy, \quad r = \sqrt{x^2 + y^2}. \quad (\text{A1})$$

Here 2Δ is the counter width. Infinite limits of integration over y (along the counter) may be used if the counter length is large in comparison with the transverse shower size. Assuming that the number of particles crossing a small counter volume obeys the Poisson distribution, we find for the probability that at least one particle hits the counter

$$P_h(\epsilon, t_j, x_j) = 1 - \exp[-n_c(\epsilon, t_j, x_j)]. \quad (\text{A2})$$

Formula (A2) gives the average number of hits detected in the counter at a given location (t_j, x_j) . The summation of P_h over depth, width and the whole volume of the detector gives the average projected lateral distribution of hits, the average longitudinal profile, and the average total number of hits in a shower with energy ϵ , respectively.

We have tested three different parameterizations of the particle density $n_e(\epsilon, t, r)$. One is based on the well-known formulae given by Greisen [19] for the longitudinal profile of a shower and on the NKG-function for the lateral distribution of electrons:

$$n_e(\epsilon, t, r) = n_G(\epsilon, t) f_{\text{NKG}}(S, r), \quad (\text{A3})$$

where

$$n_G(\epsilon, t) = 0.32 \exp\left[t \left(1 - \frac{3}{2} \ln S\right)\right] / \sqrt{\ln \epsilon}; \quad (\text{A4})$$

$$S = 3t / (t + 2 \ln \epsilon); \quad 0 < S \leq 2; \quad (\text{A5})$$

$$f_{\text{NKG}}(S, r) = C(S) r^{S-2} (r + 1)^{S-4.5}; \quad (\text{A6})$$

$$C(S) = \Gamma(4.5 - S) / [2\pi \Gamma(S) \Gamma(4.5 - 2S)]; \quad (\text{A7})$$

S is the shower age; $C(S)$ is the normalisation factor; $\Gamma(z)$ is the Gamma-function. Hereafter, the shower energy ϵ is measured in effective critical energies β' , and the distance from the axis in effective Moliere units r'_M . These effective parameters may significantly differ from the values introduced in the cascade theory and may be estimated for a specific calorimeter structure from the observed characteristics of cascade showers and the available calibration results.

The second version of the formulae for particle density differs from (A3) - (A7) by a modified expression for a longitudinal cascade curve

$$n_{\text{GM}}(\epsilon, t) = 0.29 \exp\left[t \left(1 - \frac{7}{4} \ln S'\right)\right] / \sqrt{\ln \epsilon}; \quad (\text{A8})$$

$$S' = 7t / (3t + 4 \ln \epsilon). \quad (\text{A9})$$

This expression describes the shower absorption in iron after the shower maximum considerably better than (A4). Finally, the third version includes a modification of the depth dependence for the lateral distribution function. According to the results of the numerical integration of cascade equations [20], a better description is reached by introducing the S - dependence of the effective Moliere unit

$$r'_M(S) = r'_M(1) [1 - 0.37 (S - 1)] \quad (\text{A10})$$

instead of $r'_M = \text{const.}$ We estimated the parameters r'_M, β' for the NUSEX detector from the fit of the experimental lateral distributions of hits (Fig.3) and from the electron beam calibration data of the test module (squares in Fig.4). The effective Moliere unit was derived from the lateral distributions which only slightly depend on the β' value used in the calculations, whereas the effective critical energy defining the absolute energy calibration was determined for the corresponding r'_M from the electron calibration point at 1 GeV shower energy. The estimates of r'_M and β' for different versions of the parameterization of particle density are given in Table A1.

Table A1. Best-fit values of the parameters for different parameterizations of cascade particle density.

Formulae used	Best-fit parameters
(A3 - A7)	$b' = 26.8 \text{ MeV}; r'_M = 1.87 \text{ cm}$
(A3), (A8-A9),(A5-A7)	$b' = 27.4 \text{ MeV}; r'_M = 1.67 \text{ cm}$
(A3), (A8-A9),(A5-A7),(A10)	$b' = 29.0 \text{ MeV}; r'_M(1) = 2.11 \text{ cm}$

Owing to the normalisation at 1 GeV, all three fits give practically the same total number of hits in the range 0.4 - 4 GeV, but predict slightly different calibration curves out of this energy range and for different distances D_x from the shower axis. Since the available data do not allow a clear choice among the three versions of the analytical formulae, we consider the difference between these fits as a systematic uncertainty of the calibration curves used in the present analysis. An additional ambiguity (about 5% in the energy scale) arises from the not full identity of the measurement conditions in the NUSEX detector and in the test module exposed at the accelerator beam (multihit probability, orientation of the shower axes, influence of spacers, etc.) The overall region of calibration curve uncertainties is given in Fig.4.

APPENDIX B: DISTORTION OF THE ENERGY SPECTRUM OF RECONSTRUCTED CASCADES DUE TO SHOWER SUPERPOSITION

The superposition of showers originated from individual muon interactions leads to a distortion of the energy distribution of reconstructed cascades in comparison with the true one, the value of the distortion depending on the reconstruction procedure and on the sampling thickness (if the latter is not negligibly small in comparison with the longitudinal shower size). A quantitative consideration of this effect has been given in [15].

A Monte Carlo simulation of the response of a sampling calorimeter for high energy muons has been performed. Interaction points and energy transfers above the threshold $\varepsilon_0 = 0.05$ GeV have been selected according to the muon interaction cross section. In order to obtain the pure effect of shower superposition, the following simplifications have been adopted: the muon energy was not modified after the interaction (i.e., it was constant); cascade fluctuations and measurement errors were not taken into account, and the contribution of each interaction into the calorimeter response $n(t_i)$ was calculated following the average transition curve; showers from photonuclear muon interaction were treated as usual electromagnetic ones. The sampling thickness of the setup was taken equal to 2 cm of iron (1.14 r.l.).

Showers in simulated events were reconstructed using the following procedure. Two showers were considered to be separated if the minimum between them was at least twice less than each of the corresponding maxima. In order to exclude a systematic distortion of the shower energy, the value of the signal in this minimum was in this case equally shared between the two neighbouring reconstructed cascades. The energy transfer in the reconstructed interaction was estimated by means of the usual sampling calorimeter formula:

$$\varepsilon = \beta\tau \sum n(t_i). \quad (\text{B1})$$

Here β is the track length constant, and τ is the sampling thickness (in r.l.). The summation was performed within the reconstructed shower boundaries.

For every muon energy, two distributions were confronted: the distribution of the energies transferred in simulated interactions $N_{\text{sim}}(\Delta\varepsilon)$, and that of the reconstructed showers $N_{\text{rec}}(\Delta\varepsilon)$. Note that the first of these distributions equals the product of the cross section for muon interactions in the $\Delta\varepsilon$ interval and the total muon path in the target. Then the distortion of the energy spectrum of reconstructed interactions may be represented by the ratio

$$\eta(\Delta\varepsilon) = N_{\text{rec}}(\Delta\varepsilon)/N_{\text{sim}}(\Delta\varepsilon). \quad (\text{B2})$$

The results of calculations of the distortion function (B2) for several muon energies are given by solid curves in Fig.10. Because of the scaling behaviour of the muon interaction cross section at high energies, the regions where the cascade superposition is important are determined mainly by the relative energy transfer. Thus, the loss of separate interactions is appreciable when they are frequent, namely, for $v \leq 3 \times 10^{-3}$. On the other hand, the overestimation of the reconstructed energies due to superposition of two or several cascade showers is significant for $10^{-2} \leq v \leq 10^{-1}$, where the cross section falls off rapidly

with the increase of ν . For $\nu > 10^{-1}$ interactions are rare and the effects discussed here are small. As we have pointed out above, the value of the distortion depends on the reconstruction procedure details and on the sampling thickness. The dashed curve in Fig.10 was calculated for conditions close to those of the CCFR experiment [1]. Similarly to the procedure adopted there, in order to separate two neighbouring showers a dip between them below a fixed discrimination level (about 2 equivalent particles above the single muon signal) was required. The calculated distortion (about 30% around 1 GeV energy transfer) is in quantitative agreement with the difference between the measured and expected stochastic energy loss distributions reported by the CCFR group.

Obviously, the overlap of separate cascades in a usual (analog) calorimeter does not influence the measurements of the total muon energy loss rate $\langle dE/dx \rangle$ and the CCFR results concerning the latter value are not affected by this effect.

References

- [1] W.K.Sakumoto et al., Phys. Rev. D45, 3042 (1992).
- [2] C.Grupen, Fortschritte der Physik 23, 127 (1976).
- [3] G.Battistoni et al., Nucl. Instrum. Methods A245, 277 (1986).
- [4] G.Battistoni et al., Phys. Lett. 133B, 454 (1983).
- [5] M.Aglietta et al., Nucl. Phys. B14, 193 (1990).
- [6] M.Aglietta et al., Europhys. Lett. 8, 611 (1989).
- [7] G.Battistoni et al., Nucl. Phys. B14,211 (1990).
- [8] M.Aglietta et al., Europhys. Lett.15, 559 (1991).
- [9] R.P.Kokoulin, Moscow Engineering Physics Institute, Preprint No.019-93, 1993.
- [10] G.T.Zatsepin and E.D.Mikhail'chi, Izv. Akad. Nauk SSSR, ser. fiz., 30, 1679 (1966); V.I.Gurentsov, G.T.Zatsepin and E.D.Mikhail'chi, Sov. J. Nucl. Phys. 23, 527 (1976).
- [11] H.Bilokon et al., Nucl. Instrum. Methods A303, 381 (1991).
- [12] R.P.Kokoulin, Institute for Cosmic Ray Research, University of Tokyo, Report No.102-82-5, 1982.
- [13] W.Lohmann, R.Kopp and R.Voss, CERN Report 85-03, Geneva, 1985.
- [14] V.I.Gurentsov, Institute for Nuclear Research, Preprint No.0380, Moscow, 1984 (in Russian).
- [15] R.P.Kokoulin and A.A.Petrukhin, Moscow Engineering Physics Institute, Preprint No.020-93, 1993.
- [16] H.J.Bhabha, Proc. Roy. Soc. A164, 257 (1938).
- [17] A.A.Petrukhin and V.V.Shestakov, Can. J. Phys. 46, S 377 (1968).
- [18] R.P.Kokoulin and A.A.Petrukhin, in Proceedings of the 11th International Conference on Cosmic Rays, Budapest, Hungary, 1969 [Acta Phys. Acad. Sci. Hung. 29, Suppl.4, 277 (1970)]; in Proceedings of the 12th International Conference on Cosmic Rays, Hobart, Australia, 1971, vol.6, p.2436.
- [19] K.Greisen, in Progress in Cosmic Ray Physics, vol.3, edited by J.G.Wilson, Amsterdam, 1956, p.1.
- [20] A.A.Lagutin, Thesis, Moscow Engineering Physics Institute, 1979 (unpublished).

Figure Captions

Fig.1. Examples of muon events in the NUSEX detector: (a) scattered low energy muon; (b) muon bundle; (c) multiple interactions of a high energy muon; (d) nuclear interaction; (e) interaction outside the detector volume. Solid vertical lines correspond to the positions of iron spacers; dotted lines indicate the 16-tube chamber boundaries.

Fig.2. Operation of the scanning routine for the event plotted in Fig.1(c). Upper histogram: number of hits in each plane at distances 1 - 15 cm from the muon track. Lower histogram: sum of the numbers of hits in 7 consecutive planes. Vertical bars indicate the reconstructed shower boundaries.

Fig.3. Projected lateral distribution functions for showers observed in the NUSEX detector. Points: experimental data corresponding to $n_H = 11 - 20$, $21 - 36$, $37 - 61$, and $62 - 100$ (from the bottom to the top, respectively). Curves: results of analytical model calculations with best-fit parameters.

Fig.4. Average number of hits versus cascade shower energy. Squares: accelerator electron beam calibration [3] of the test module. Upper curve: analytical model calculations (best-fit parameters) for the test module, $D_x \geq 0$. The lower region represents the model prediction for the distance range $D_x = 1 - 15$ cm in the NUSEX detector.

Fig.5. Average energy of vertical muons versus depth in standard rock. Full and open circles: calculations with two independent Monte Carlo codes (see text). Squares: results of the semi-analytical solution of the muon transport problem [14]. The curve represents the fit based on the analytical expression (7) for the muon spectrum with the parameters given by (9).

Fig.6. Differential energy spectra of vertical muons at different depths of standard rock (4×10^5 , 5×10^5 and 7×10^5 g/cm² respectively, from the top to the bottom). Histograms: Monte Carlo calculations. Curves: analytical approximation (7) with the parameters given by (9).

Fig.7. Ratio of the observed shower distribution in the number of hits to the expected one. All corrections are included. The curves outline the estimated region of systematic uncertainties.

Fig.8. Derived transferred energy spectrum in the NUSEX detector. Circles: experimental data (with the corrections discussed in the text). Vertical bars correspond to statistical errors; horizontal bars indicate the uncertainties in the energy estimate. The solid curve represents the

expected spectrum of muon electromagnetic interactions in the NUSEX material at the observation depth (average muon energy 360 GeV). Dashed curve: spectrum calculated with a decreased cross section in the region of low relative transfers (according to deviations in Fig.9 of [1]).

Fig.9. Relative contribution of various muon interaction processes to the generation spectrum of cascades in the NUSEX detector.

Fig.10. Distortion function (B2). Solid curves 1, 2, and 3: sampling thickness 2 cm of iron, energies equal to 100, 300, and 1000 GeV, respectively. Dashed curve: sampling thickness 10 cm of iron, muon energy 1130 GeV, shower separation procedure similar to that described in [1].




FIGURE 1





FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10