

Two-parton scattering in the high-energy limit: climbing two- and three-Reggeon ladders

G. Falcioni¹, E. Gardi¹, N. Maher¹, C. Milloy² and L. Vernazza^{2,3*}

¹ Higgs Centre for Theoretical Physics, School of Physics and Astronomy,
The University of Edinburgh, Edinburgh EH9 3FD, Scotland, UK

² Dipartimento di Fisica and Arnold-Regge Center, Università di Torino,
and INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy

³ Theoretical Physics Department, CERN, Geneva, Switzerland

* Leonardo.Vernazza@to.infn.it



15th International Symposium on Radiative Corrections:
Applications of Quantum Field Theory to Phenomenology,
FSU, Tallahassee, FL, USA, 17-21 May 2021
doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)

Abstract

We review recent progress on the calculation of scattering amplitudes in the high-energy limit. We start by illustrating the shockwave formalism, which allows one to calculate amplitudes as iterated solutions of rapidity evolution equations. We then focus on our recent results regarding $2 \rightarrow 2$ parton scattering. We present the calculation of the imaginary part of the amplitude, at next-to-leading logarithmic accuracy in the high-energy logarithms, formally to all orders, and in practice to 13 loops. We then discuss the computation of the real part of the amplitude at next-to-next-to-leading logarithmic accuracy and through four loops. Both computations are carried in full colour, and provide new insight into the analytic structure of scattering amplitudes and their infrared singularity structure.



Copyright L. Vernazza *et al.*

This work is licensed under the Creative Commons
[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 18-11-2021

Accepted 02-02-2022

Published 20-06-2022

doi:[10.21468/ciPostPhysProc.7.007](https://doi.org/10.21468/ciPostPhysProc.7.007)



Check for
updates

Contents

1 Introduction	1
2 Methodology	4
3 Two-Reggeon cut	5
4 Three-Reggeon cut	10
5 Conclusion	15
References	17

1 Introduction

The high-energy, or Regge limit of gauge-theory scattering amplitudes offers a unique insight into gauge dynamics. In this limit amplitudes simplify and factorize in rapidity, revealing new degrees of freedom that propagate in two transverse dimensions. Rapidity evolution equations, such as BFKL [1, 2] and its non-linear generalization [3–7], allow us to translate concepts from Regge theory [8] into calculation tools, leading to concrete predictions for partonic amplitudes [9–17]. The study of the Regge limit has been crucial in determining multi-leg planar $\mathcal{N} = 4$ super Yang-Mills (SYM) amplitudes in general kinematics to unprecedented accuracy, see e.g. [10, 16, 18, 19]. In parallel, computation of the Regge limit in $2 \rightarrow 2$ scattering for general colour [11–14] have been shown to provide powerful constraints on soft singularities of amplitudes in general kinematics [15, 20–22].

In this talk we focus on $2 \rightarrow 2$ gauge-theory scattering amplitudes, and review recent results in [13, 14, 23, 24], obtained within a framework developed in [11, 12]. In a scattering configuration $1 + 2 \rightarrow 3 + 4$, with Mandelstam invariants $s \equiv (p_1 + p_2)^2$, $t \equiv (p_1 - p_4)^2$ and $u \equiv (p_1 - p_3)^2 = -s - t$, the high-energy limit is defined by the condition $s \gg -t$. Assuming that the momentum transfer $-t$ is large compared to the QCD scale, these amplitudes can be calculated as an expansion in the strong coupling, and notoriously develop large logarithms in the ratio $\frac{s}{-t}$. It has long been known that the Leading Logarithms (LLs) are resummed to all orders [25, 26] via

$$\mathcal{M}_{ij \rightarrow ij}^{\text{LL}}(s, t) = \left(\frac{s}{-t}\right)^{C_A \alpha_g(t)} \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad \text{with} \quad \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} = g_s^2 \frac{2s}{t} T_i \cdot T_j, \quad (1)$$

where the colour generator T_i is in the representation of parton i , and

$$\alpha_g = \frac{\alpha_s}{\pi} \frac{r_\Gamma}{2\epsilon} + \mathcal{O}(\alpha_s^2); \quad r_\Gamma = e^{\epsilon\gamma_E} \frac{\Gamma^2(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(1-2\epsilon)} \quad (2)$$

is the gluon Regge trajectory, written in dimensional regularization with $D = 4 - 2\epsilon$. Higher-order corrections, not written in eq. (2), contribute beyond LL. The simple exponentiation property in eq. (1) can be interpreted as due to the exchange of an effective degree of freedom, which is identified as a single *Reggeized gluon*, or *Reggeon*. At higher logarithmic accuracy the scattering amplitude develops a more complex analytic structure, which can be understood in term of the exchange of multiple Reggeons [27–29] in the transverse plane. These involve evolution equations which are integrable in the planar limit [30–33], but are difficult to solve in general. However, in the perturbative regime they can be integrated iteratively [11–14], to obtain perturbative high-energy amplitudes order-by-order in α_s .

The factorization structure of the amplitude beyond LLs becomes clearer by exploiting the symmetry under the exchange $s \leftrightarrow u$, also known as “signature” symmetry. To this end, the scattering amplitude is split into odd (–) and even (+) component with respect to $s \leftrightarrow u$:

$$\mathcal{M} = \mathcal{M}^{(-)} + \mathcal{M}^{(+)}. \quad (3)$$

It is also convenient to expand the amplitude into a signature-symmetric definition for the large logarithm:

$$\frac{1}{2} \left(\log \frac{-s - i0}{-t} + \log \frac{-u - i0}{-t} \right) = \log \left| \frac{s}{t} \right| - i \frac{\pi}{2} \equiv L, \quad (4)$$

such that one has

$$\mathcal{M}_{ij \rightarrow ij}^{(\pm)} = \sum_{n=0}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \sum_{m=0}^n L^m \mathcal{M}_{ij \rightarrow ij}^{(\pm, n, m)}, \quad \text{with} \quad \mathcal{M}_{ij \rightarrow ij}^{(-, 0, 0)} \equiv \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}. \quad (5)$$




Figure 1: Schematic relation between logarithmic counting and Reggeon exchanges for two-parton amplitudes in the high-energy limit. The grey blobs represent impact factors, while the red ones are corrections to the Regge trajectory.

Splitting the amplitudes into (\pm) components offers several advantages: first of all, signature is preserved under rapidity evolution, which greatly simplifies the computation of these amplitudes [11–14]. Furthermore, given that a single Reggeon has the quantum numbers of a gluon exchanged in the t channel, which is odd under the signature symmetry, one finds that $\mathcal{M}^{(-)}$ consists of an odd number of Reggeons, while $\mathcal{M}^{(+)}$ involves an even number of them [11]. In addition, there are other useful properties: for instance, it is possible to show [12] that the odd amplitude coefficients $\mathcal{M}^{(-,n,m)}$ are purely *real*, while the even ones $\mathcal{M}^{(+,n,m)}$ are purely *imaginary*, when expanded in powers of the symmetric logarithm L . Moreover, for gluon-gluon and quark-gluon scattering Bose symmetry links the kinematic dependence to that of colour, therefore $\mathcal{M}^{(+)}$ and $\mathcal{M}^{(-)}$ are governed by t -channel exchange of colour representations which are respectively even and odd under $1 \leftrightarrow 4$ (or $2 \leftrightarrow 3$) interchange.

One obtains a schematic representation for the amplitude, which connects the logarithmic accuracy with the number of Reggeon exchanges, as represented in fig. 1. One has

$$\mathcal{M}^{(-)} = \mathcal{M}^{(-),\text{SRS}} + \mathcal{M}^{(-),\text{MRS}}, \quad \mathcal{M}^{(+)} = \mathcal{M}^{(+),\text{MRS}} \quad (6)$$

where the superscripts label amplitudes involving single-Reggeon states (SRS) and multi-Reggeon states (MRS), respectively. As shown in fig. 1, the odd component $\mathcal{M}^{(-)}$ is given up to NLL accuracy by a single Reggeon exchange [34, 35], with $\mathcal{O}(\alpha_s^2)$ corrections to the trajectory in (2), along with s -independent impact factors [36–39]:

$$\mathcal{M}_{ij \rightarrow ij}^{(-),\text{LL+NLL}} = \mathcal{M}_{ij \rightarrow ij}^{(-),\text{SRS}} \Big|_{\text{NLL}} = e^{C_A \alpha_s(t)L} C_i(t) C_j(t) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}. \quad (7)$$

The factors $C_i(t)$ are expanded in perturbation theory, with coefficients $C_i^{(n)}$. The $n = 1$ term contributes at NLL, while the higher-order terms give rise to further subleading logarithms.

A proper description of the scattering amplitude beyond eq. (7) requires to take into account the t -channel exchange of multi-Reggeon states. The signature even amplitude $\mathcal{M}^{(+)}$, which starts at NLL accuracy, is governed at this logarithmic order by two-Reggeon exchange [11, 13]. Furthermore, starting with the odd amplitude $\mathcal{M}^{(-)}$ at NNLL accuracy, one needs to take into account the exchange of three Reggeons and their mixing with one-Reggeon states [12, 40]. In this talk we review recent results obtained in [13, 14] and [23, 24], where the two towers of coefficients $\mathcal{M}^{(+,n,n-1)}$ and $\mathcal{M}^{(-,n,n-2)}$ have been expressed to all orders as iterated solution of the BFKL equation and its generalization, the Balitsky-JIMWLK equation.

Thanks to the development of new methods, the $\mathcal{M}^{(+,n,n-1)}$ tower has been evaluated explicitly to 13 loops, [14], while the tower $\mathcal{M}^{(-,n,n-2)}$ has been computed to 4 loops in [23, 24].

The calculation of scattering amplitudes in terms of Reggeon exchanges is particularly interesting, because it allows one to connect modern perturbation theory with concepts such as Regge poles and cuts, obtaining a deeper understanding of the analytic structure of scattering amplitudes. In general, an amplitude that can be brought to the form of eq. (7) is said to admit *Regge-pole factorization*. It is only admitted by the signature odd amplitude, to NLL accuracy. The signature even amplitude is described in turn by a Regge cut associated with a two-Reggeon exchange. Starting at NNLL accuracy, also the signature odd amplitude receives contributions due to multi-Reggeon states, giving rise to Regge cuts, such that one has

$$\mathcal{M}_{ij \rightarrow ij}^{(-), \text{LL+NLL+NNLL}} = e^{C_A \alpha_g(t)L} C_i(t) C_j(t) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} + \mathcal{M}_{ij \rightarrow ij}^{(-), \text{MRS}}. \quad (8)$$

The second term in the r.h.s. first arises at NNLL at two loops [41, 42], and it has been interpreted as being due to three-Reggeon exchange in refs. [12, 40, 43], where the three-loop NNLL amplitude was determined. As a consequence of eq. (8), at NNLL accuracy $C_{i/j}(t)$ and $\alpha_g(t)$ start to depend on how the separation in eq. (6) is precisely defined, which is referred to below as a scheme choice. Following [12], we adopt the scheme in which the Regge cut contribution is then identified with the MRS, which is explicitly computed using the Balitsky-JIMWLK rapidity evolution equation. The Regge-pole contribution is identified with the SRS contribution, determined by matching the r.h.s of eq. (8) to the full amplitude. Other schemes may be possible, and we refer to [24] for a more detailed discussion on this topic.

An additional motivation to investigate multi-loop corrections in the high-energy limit is that it provides information on the infrared singularity structure of amplitudes. Further discussion of this aspect can be found in the original papers [11–13, 20, 23, 24] and in a talk given at this conference [44], to which we refer for further details.

2 Methodology

Following [11, 12], we describe two parton scattering in the high-energy limit within the shock-wave formalism: fast particles moving in the (+) lightcone direction, defined as *projectile*, scatter against fast particles moving in the (−) lightcone direction, or *target*. The fast projectile $|\psi_i\rangle$ appears as a set of infinite Wilson lines $U(z_1) \otimes \dots \otimes U(z_n)$ [45, 46] at transverse position $z_k \equiv x_{k\perp}$, crossing the target $\langle\psi_j|$ (seen as a “shockwave”) at $x^- = 0$, with

$$U(z) = \mathcal{P} \exp \left[i g_s \mathbf{T}^a \int_{-\infty}^{+\infty} dx^+ A_+^a(x^+, x^- = 0, z) \right]. \quad (9)$$

In perturbation theory the unitary matrices $U(z)$ are close to the identity, therefore they can be parameterized in terms of a colour-adjoint field W^a :

$$U(z) = e^{i g_s \mathbf{T}^a W^a(z)}, \quad (10)$$

which is identified as a source for Reggeized gluons. The projectile and target are thus expanded in Reggeon fields: schematically,

$$|\psi_i\rangle = \sum_{n=1}^{\infty} |\psi_{i,n}\rangle \sim g_s [\mathbf{T}W]_i(p) - g_s^2 [\mathbf{T}W \otimes \mathbf{T}W]_i(p) - g_s^3 [\mathbf{T}W \otimes \mathbf{T}W \otimes \mathbf{T}W]_i(p) + \dots, \quad (11)$$

where $|\psi_{i,n}\rangle$ represents a state of n Reggeons, which depends on the transverse momentum p , but not on the center-of-mass energy. The energy dependence enters through the fact that infinite Wilson lines develop rapidity divergences, regulated introducing a rapidity cutoff $\eta = L$,

leading to a rapidity evolution equation for the projectile (and the target)

$$-\frac{d}{d\eta} |\psi_i\rangle = H |\psi_i\rangle, \tag{12}$$

where H is the Balitsky-JIMWLK Hamiltonian [3–7]. A key feature of eq. (12) is the non-linearity of H : evolution of the full projectile $|\psi_i\rangle$ in rapidity generates an increasing number of Wilson lines $U(z_j)$, eventually leading to the phenomenon of gluon saturation. However, in applications to partonic scattering processes it is appropriate to take the limit of dilute projectile and target [11, 12], in which case H acts on states with a given number of Reggeon fields W . In this perturbative regime, H is diagonal to leading order in g_s^2 ; the non-linearity of H manifests at higher orders in the coupling, producing transitions between states with different number of Reggeon fields: a transition $k \rightarrow k + 2n$ is of order $g_s^{2(1+n)}$. Also, note that, as a consequence of the signature symmetry, only transitions $k \rightarrow k \pm 2$ are allowed, while transitions $k \rightarrow k \pm 1$ are forbidden. In short, the r.h.s of eq. (12) has the form

$$H \begin{pmatrix} W \\ WW \\ WWW \\ \dots \end{pmatrix} \equiv \begin{pmatrix} H_{1 \rightarrow 1} & 0 & H_{3 \rightarrow 1} & \dots \\ 0 & H_{2 \rightarrow 2} & 0 & \dots \\ H_{1 \rightarrow 3} & 0 & H_{3 \rightarrow 3} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} W \\ WW \\ WWW \\ \dots \end{pmatrix} \simeq \begin{pmatrix} g_s^2 & 0 & g_s^4 & \dots \\ 0 & g_s^2 & 0 & \dots \\ g_s^4 & 0 & g_s^2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} W \\ WW \\ WWW \\ \dots \end{pmatrix}, \tag{13}$$

where non-vanishing entries on the r.h.s. display the perturbative orders at which the various transition Hamiltonian start contributing. While off-diagonal entries are obtained from expansion of the Leading Order (LO) Balitsky-JIMWLK equation, diagonal entries are given by the Regge trajectory for $H_{1 \rightarrow 1}$, or the LO BFKL equation for $H_{n \rightarrow n}$, with $n \neq 1$.

A scattering amplitude in the high-energy limit is then given as an expectation value between states with fixed number of Reggeon fields W evolved to equal rapidity:

$$\frac{i(Z_i Z_j)^{-1}}{2s} \mathcal{M}_{ij \rightarrow ij} = \langle \psi_j | e^{-HL} | \psi_i \rangle, \quad \text{with} \quad Z_i(t) = \exp \left\{ -\frac{1}{2} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \Gamma_i(\alpha_s(\lambda^2), \lambda^2) \right\}. \tag{14}$$

Notice that, for simplicity, we define the expectation value in terms of states $|\psi_i\rangle$ where collinear divergences have been removed. These are given by the factors $Z_i(t)$, defined as an integral over the scale of Γ_i , where, to three loops, $\Gamma_i = \frac{1}{2} \gamma_K(\alpha_s(\lambda^2)) C_i \log \frac{\mu^2}{\lambda^2} + 2\gamma_i$. Here γ_K is the component of the cusp anomalous dimension [47–51] proportional to the quadratic Casimir C_i , in the representation of parton i , and γ_i are anomalous dimensions associated with on-shell form factors [42, 52].

Our aim is to calculate transitions which involve multi-Regge states, therefore, it proves useful to introduce a *reduced amplitude*, obtained from the original amplitude by removing single Reggeon transitions:

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij} = e^{-T_t^2 \alpha_g(t)L} \langle \psi_j | e^{-HL} | \psi_i \rangle \equiv \langle \psi_j | e^{-\hat{H}L} | \psi_i \rangle, \quad \hat{H}_{k \rightarrow k+2n} = H_{k \rightarrow k+2n} + \delta_{n0} \mathbf{T}_t^2 \alpha_g(t), \tag{15}$$

which we have expressed in terms of a *reduced Hamiltonian* \hat{H} . After evolution has been performed, the contraction of Reggeons of equal rapidity is evaluated in terms of free propagators $\langle W^a(p) | W^b(q) \rangle = \frac{i}{p^2} \delta^{ab} \delta^{2-2\epsilon}(p-q) + \mathcal{O}(g_s^2)$, see [11]. We are now ready to discuss the calculation of the amplitudes $\hat{\mathcal{M}}^{(+,n,n-1)}$ and $\hat{\mathcal{M}}^{(-,n,n-2)}$.

3 Two-Reggeon cut

According to eq. (15) and fig. 1, the NLL contribution to the even amplitude is given by

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(+),\text{NLL}} = \langle \psi_{j,2} | e^{-\hat{H}L} | \psi_{i,2} \rangle, \tag{16}$$




Figure 2: Representation of the two Reggeon amplitude and the corresponding BFKL wavefunction. A single application of the BFKL Hamiltonian adds a rung in the ladder. The amplitude at ℓ loops, $\mathcal{M}^{(\ell)}(p)$, is obtained by integrating over the wavefunction $\Omega^{(\ell-1)}(p, k)$, which itself consists of $\ell - 1$ rungs. Figure taken from [15] under the Creative Commons Attribution 4.0 International License.

where the subscript of $|\psi_{i,2}\rangle$ identifies the two-Reggeon component of the projectile i (and target j). Upon expansion one obtains [1, 2, 11, 13, 53]:

$$\hat{\mathcal{M}}_{ij \rightarrow ij}^{(+, \ell, \ell-1)} = -i\pi \frac{r_\Gamma^\ell}{(\ell-1)!} \int [Dk] \frac{p^2}{k^2(p-k)^2} \Omega^{(\ell-1)}(p, k) \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad (17)$$

where $[Dk] \equiv \frac{\pi}{r_\Gamma} \left(\frac{\mu^2}{4\pi e^{-\gamma_E}} \right)^\epsilon \frac{d^{2-2\epsilon}k}{(2\pi)^{2-2\epsilon}}$, and r_Γ has been defined in eq. (2). We refer to $\Omega^{(\ell-1)}(p, k)$ as the two Reggeized gluon wavefunction at $\ell - 1$ loop order, as shown in fig. 2. The wavefunction admits the BFKL evolution equation

$$\frac{d}{dL} \Omega(p, k) = \frac{\alpha_s r_\Gamma}{\pi} \hat{H} \Omega(p, k), \quad (18)$$

where the Hamiltonian has two components $\hat{H} = (2C_A - \mathbf{T}_t^2) \hat{H}_i + (C_A - \mathbf{T}_t^2) \hat{H}_m$ with

$$\hat{H}_i \Psi(p, k) = \int [Dk'] f(p, k, k') [\Psi(p, k') - \Psi(p, k)], \quad \hat{H}_m \Psi(p, k) = J(p, k) \Psi(p, k), \quad (19a)$$

where

$$f(p, k, k') \equiv \frac{k^2}{k'^2(k-k')^2} + \frac{(p-k)^2}{(p-k')^2(k-k')^2} - \frac{p^2}{k'^2(p-k')^2}, \quad (20)$$

and

$$J(p, k) = \frac{1}{2\epsilon} + \int [Dk'] f(p, k, k') = \frac{1}{2\epsilon} \left[2 - \left(\frac{p^2}{k^2} \right)^\epsilon - \left(\frac{p^2}{(p-k)^2} \right)^\epsilon \right]. \quad (21)$$

While the exact solution is unknown, the equation can be solved iteratively

$$\Omega(p, k) = \sum_{\ell=1}^{\infty} \left(\frac{\alpha_s r_\Gamma}{\pi} \right)^\ell \frac{L^{\ell-1}}{(\ell-1)!} \Omega^{(\ell-1)}(p, k), \quad \Omega^{(\ell-1)}(p, k) = \hat{H} \Omega^{(\ell-2)}(p, k), \quad (22)$$

and we get

$$\Omega^{(0)}(p, k) = 1, \quad \Omega^{(1)}(p, k) = (C_A - \mathbf{T}_t^2) J(p, k), \quad (23)$$

$$\Omega^{(2)}(p, k) = (C_A - \mathbf{T}_t^2)^2 J^2(p, k) + (2C_A - \mathbf{T}_t^2)(C_A - \mathbf{T}_t^2) \int [Dk'] f(p, k, k') [J(p, k') - J(p, k)],$$

and so on. While for the first few orders it is possible to integrate directly these wavefunctions to get the dimensionally-regularized amplitude, this task becomes quickly intricate at higher orders. It is thus necessary to resort to more sophisticated techniques [13, 14].

There are two crucial observations that allows one to carry the integration in eqs. (17) and (19) to high order: the first is that the wavefunction $\Omega^{(\ell)}(p, k)$ is finite, to any order, which follows from the form of the Hamiltonian (19). As a consequence, all infrared singularities in the NLL amplitude must be generated in the final integration over k in eq. (17), from the soft regions where $k \ll p$ or where $p - k \ll p$. The symmetry of the problem implies that these regions give the same contribution, hence it is sufficient to focus on the former, and multiply the result by two. The second observation is that BFKL evolution closes in the soft approximation, $k \ll p$. This is easily seen by considering a single evolution step of the wavefunction generated by the Hamiltonian (19), and observing that $f(p, k, k')$ vanishes in regions other than the soft region $k \sim k' \ll p$. Applying these arguments iteratively, we deduce that the soft limit of the wavefunction (and thus the complete singular part of the amplitude) is fully determined by the configuration where the entire side rail of the ladder is soft. It is thus possible to expand eq. (19) and solve it for $k \sim k' \ll p$. This is much easier, because in this approximation the original two-scale problem, k^2 and $(p - k)^2$, is reduced to a single-scale problem. The integrals are then simple to perform, and the solution for the wavefunction at any order is given as a polynomial in $\xi \equiv (p^2/k^2)^\epsilon$:

$$\Omega^{(\ell-1)}(p, k) = \frac{(C_A - \mathbf{T}_t)^{\ell-1}}{(2\epsilon)^{\ell-1}} \sum_{n=0}^{\ell-1} (-1)^n \binom{\ell-1}{n} \left(\frac{p^2}{k^2}\right)^{n\epsilon} \prod_{m=0}^{n-1} \left\{ 1 - \hat{B}_m(\epsilon) \frac{2C_A - \mathbf{T}_t}{C_A - \mathbf{T}_t} \right\}, \quad (24)$$

where $\hat{B}_n(\epsilon) = 2n(2+n)\zeta_3\epsilon^3 + 3n(2+n)\zeta_4\epsilon^4 + \dots$. Performing the final integration over k according to (17) we obtain all the infrared-singular contributions to the NLL amplitude at any loop order ℓ . Remarkably, the result can be resummed into a closed-form expression:

$$\hat{\mathcal{M}}_{\text{NLL}}^{(+)} \Big|_s = \frac{i\pi}{L(C_A - \mathbf{T}_t^2)} \left(1 - R(\epsilon) \frac{C_A}{C_A - \mathbf{T}_t^2} \right)^{-1} \left[\exp \left\{ \frac{r_\Gamma}{2\epsilon} \frac{\alpha_s}{\pi} L(C_A - \mathbf{T}_t) \right\} - 1 \right] \mathbf{T}_{s-u}^2 \mathcal{M}^{(\text{tree})} \quad (25)$$

up to $\mathcal{O}(\epsilon^0)$ terms, where

$$R(\epsilon) = \frac{\Gamma^3(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(1-2\epsilon)} - 1 = -2\zeta_3\epsilon^3 - 3\zeta_4\epsilon^4 - 6\zeta_5\epsilon^5 - (10\zeta_6 - 2\zeta_3^2)\epsilon^6 + \mathcal{O}(\epsilon^7). \quad (26)$$

Now that the singularities have been determined, the remaining task it to compute the $\mathcal{O}(\epsilon^0)$ finite terms in the amplitude. Once again, the fact that the wavefunction is finite plays a central role [14]: it implies that the wavefunction can be computed consistently in two transverse dimensions, setting $\epsilon = 0$. This may seem to pose a problem for the subsequent calculation of the amplitude, given that the integral in eq. (17) requires (dimensional) regularization. However, this issue can be overcome by carefully combining the soft limit discussed above with the two-dimensional calculation [14].

The calculation of the wavefunction in exactly two transverse dimension is rather technical, and we refer to [14] for a detailed discussion. Let us mention here that this computation is done by representing the two-dimensional momentum k , k' and p in the BFKL kernel of eq. (19) as complex parameters, $k = k_x + ik_y$, $k' = k'_x + ik'_y$ and $p = p_x + ip_y$, and implementing a variable change in the BFKL equation:

$$\frac{k_x + ik_y}{p_x + ip_y} = \frac{z}{z-1} \quad \text{and} \quad \frac{k'_x + ik'_y}{p_x + ip_y} = \frac{w}{w-1}. \quad (27)$$

It is possible to show that in these variables, the 2-dimensional wavefunction can be expressed in terms of single-valued harmonic polylogarithms (SVHPLs, [9, 54–58]) at any order. Single

valuedness is to be expected here, because branch cuts are physically inadmissible in the Euclidean two-dimensional transverse space. Moreover, the structure of the Hamiltonian guarantees that the wavefunction at order ℓ is a pure function of uniform weight ℓ . It is then possible to express the action of the 2-dimensional Hamiltonian on any linear combination of SVHPLs $\mathcal{L}_w(z, \bar{z})$ (where the word w corresponds to a set of 0 and 1 indices) as a set of differential equations, which can be integrated iteratively, using the soft limit $z, \bar{z} \rightarrow 0$ as boundary data. The procedure has been automated in [14]. For instance, the first two orders reads:

$$\begin{aligned} \Omega_{2d}^{(1)} &= \frac{1}{2} C_2 (\mathcal{L}_0 + 2\mathcal{L}_1), \\ \Omega_{2d}^{(2)} &= \frac{1}{2} C_2^2 (\mathcal{L}_{0,0} + 2\mathcal{L}_{0,1} + 2\mathcal{L}_{1,0} + 4\mathcal{L}_{1,1}) + \frac{1}{4} C_1 C_2 (-\mathcal{L}_{0,1} - \mathcal{L}_{1,0} - 2\mathcal{L}_{1,1}). \end{aligned} \tag{28}$$

With the amplitude determined to all orders in the soft approximation, see eq. (25), and the wavefunction in two dimensions available to any required order, we return to the question of how to combine this information to obtain the full amplitude. This is achieved [14] by writing the full wavefunction as the sum of a soft and an hard components: $\Omega(p, k) = \Omega_{\text{soft}}(p, k) + \Omega_{\text{hard}}(p, k)$, where the two components are defined as follows: $\Omega_{\text{soft}}(p, k)$ is given as the symmetric version of the dimensionally-regularized solution in eq. (24), where $(p^2/k^2)^\epsilon$ is replaced by $\left(\frac{(p^2)^2}{k^2(p-k)^2}\right)^{\epsilon/2}$. This guarantees that both soft limits are taken into account; moreover, $\Omega_{\text{soft}}(p, k)$ admits an ϵ expansion in terms of SVHPLs, and hence its two-dimensional limit reproduces exactly all non-vanishing terms in the $z \rightarrow 0$ and $z \rightarrow \infty$ limits of the two-dimensional solution $\Omega^{(2d)}(z, \bar{z})$ of (28). As a consequence, one can isolate the two-dimensional limit of the hard wavefunction as follows:

$$\Omega_{\text{hard}}^{(2d)}(z, \bar{z}) \equiv \lim_{\epsilon \rightarrow 0} \Omega_{\text{hard}} = \Omega^{(2d)}(z, \bar{z}) - \Omega_{\text{soft}}^{(2d)}(z, \bar{z}). \tag{29}$$

Given that all the singularities in the amplitude arise from the soft component of the wavefunction, the full NLL amplitude (17) to $\mathcal{O}(\epsilon)$ is given by

$$\hat{\mathcal{M}}_{ij \rightarrow ij}^{(+), \text{NLL}} \left(\frac{s}{-t} \right) = -i\pi \left[\int [Dk] \frac{p^2 \Omega_{\text{soft}}(p, k)}{k^2(p-k)^2} + \frac{1}{4\pi} \int \frac{d^2z}{z\bar{z}} \Omega_{\text{hard}}^{(2d)}(z, \bar{z}) \right] \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{(\text{tree})}. \tag{30}$$

The soft wavefunction is integrated in dimensional regularization in $d = 2 - 2\epsilon$ dimensions, while the hard wavefunction, which by construction vanishes in the soft limits, can be integrated in $d = 2$. We thus obtain the full NLL $2 \rightarrow 2$ amplitude, i.e. the leading tower of logarithms in the imaginary part of the amplitude. For the first few orders one has

$$\begin{aligned} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(+,1,0)} &= i\pi \frac{1}{2\epsilon} \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{(\text{tree})}, & \hat{\mathcal{M}}_{ij \rightarrow ij}^{(+,2,1)} &= i\pi C_2 \left[\frac{1}{8\epsilon^2} - \frac{\zeta(2)}{8} \right] \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{(\text{tree})}, \\ \hat{\mathcal{M}}_{ij \rightarrow ij}^{(+,3,2)} &= i\pi C_2^2 \left[\frac{1}{48\epsilon^3} - \frac{\zeta(2)}{32\epsilon} - \frac{29}{48} \zeta(3) \right] \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{(\text{tree})}, \\ \hat{\mathcal{M}}_{ij \rightarrow ij}^{(+,4,3)} &= i\pi C_2^2 \left[\frac{C_2}{384\epsilon^4} - \frac{C_2 \zeta(2)}{192\epsilon^2} - \left(\frac{7C_2}{288} + \frac{C_A}{192} \right) \frac{\zeta(3)}{\epsilon} - \frac{C_2 \zeta(4)}{48} - \frac{C_A \zeta(4)}{128} \right] \mathbf{T}_{s-u}^2 \mathcal{M}_{ij \rightarrow ij}^{(\text{tree})}. \end{aligned} \tag{31}$$

The wavefunction is given up to $\ell = 13$ loops in [14]; higher orders can be obtained with the same algorithm. The amplitude coefficients $\hat{\mathcal{M}}^{(+, \ell, \ell-1)}$ have uniform transcendental weight, and are expressed in terms of single- and multiple-zeta values: the latter appear starting at 11 loops, with $\zeta_{5,3,3}$ being the first one to emerge. Such multiple zeta values originate from the hard part of the amplitude in eq. (30), and are thus only of the type that originates in SVHPLs.

The singularities of $\hat{\mathcal{M}}^{(+, \ell, \ell-1)}$ are part of the soft contribution, and are thus resummed by eq. (25). Combining this information with the factorization theorem for infrared singularities

allows one to extract the corresponding soft anomalous dimension and sum it to all orders, as obtained in [13]. In short, it is well known that infrared divergences in amplitudes factorize and exponentiate [48–50, 59–62] according to

$$\mathcal{M} = \mathbf{Z} \cdot \mathcal{H}, \quad \mathbf{Z} = \mathcal{P} \exp \left\{ -\frac{1}{2} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \Gamma \right\}, \quad (32)$$

where \mathcal{H} is an infrared-renormalized hard amplitude, which is finite, and Γ is the soft anomalous dimension. In the high-energy limit the latter takes the form [20]

$$\Gamma = \frac{\gamma_K}{2} [L\mathbf{T}_t^2 + i\pi\mathbf{T}_{s-u}^2] + \Gamma_i + \Gamma_j + \Delta, \quad (33)$$

where γ_{K_i} has been defined after eq. (14) and Δ represents non-dipole corrections starting at three loops [48–50, 63]. The dipole contribution is well known [20, 42]: the $L\mathbf{T}_t^2$ term contributes starting at LL, while the $i\pi\mathbf{T}_{s-u}^2$ and Γ_i terms start at NLL. Here we focus on Δ , which we expand as

$$\Delta = \sum_{n=3}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^n \sum_{m=0}^{n-1} L^m \Delta^{(n,m)}. \quad (34)$$

The calculation of the tower $\hat{\mathcal{M}}^{(+,\ell,\ell-1)}$ allows us to determine the coefficients $\Delta^{(-,\ell,\ell-1)}$ for all ℓ . We find [13]

$$\Delta^{(-,\ell,\ell-1)} = i\pi G^{(\ell)} \mathbf{T}_{s-u}^2, \quad \text{with} \quad G^{(\ell)} \equiv \frac{1}{(\ell-1)!} \left[\frac{(C_A - \mathbf{T}_t^2)}{2} \right]^{\ell-1} \left(1 - R(\epsilon) \frac{C_A}{C_A - \mathbf{T}_t^2} \right)^{-1} \Big|_{\epsilon^{\ell-1}}, \quad (35)$$

where $R(\epsilon)$ is given in eq. (26), and the subscript $|\epsilon^{\ell-1}$ indicates that one should extract the coefficient of $\epsilon^{\ell-1}$. The result in eq. (35) can be resummed to all orders: one finds

$$\Delta_{\text{NLL}}^{(-)} = i\pi \frac{\alpha_s}{\pi} G \left(\frac{\alpha_s}{\pi} L \right) \mathbf{T}_{s-u}^2, \quad (36)$$

where $G(x) = \sum_{\ell=1}^{\infty} x^{\ell-1} G^{(\ell)}$. It is interesting to note [13] that $G(x)$ is an entire function, thus it admits an infinite radius of convergence, which implies that one can compute the NLL soft anomalous dimension at any value of the effective coupling, including $\frac{\alpha_s}{\pi} L \gg 1$.

The finite, $\mathcal{O}(\epsilon^0)$ contribution to the NLL amplitude in eq. (31) can be written as

$$\hat{\mathcal{M}}_{\text{NLL}}^{(+)} = \frac{i\pi}{L} \Xi_{\text{NLL}}^{(+)} \mathbf{T}_{s-u}^2 \mathcal{M}_{\text{tree}}. \quad (37)$$

It is not yet known how to resum $\Xi_{\text{NLL}}^{(+)}$. In any case, it is clear that resummation would not involve Γ functions only, because $\Xi_{\text{NLL}}^{(+)}$ contains multiple zeta values. However, it is possible to study numerically the convergence of the series given by the coefficients $\Xi_{\text{NLL}}^{(+,\ell,\ell-1)}$, for the relevant representations for $N_c = 3$, the singlet ($\mathbf{T}_t^2 \mathcal{M}^{[1]} = 0$) and the 27 representation ($\mathbf{T}_t^2 \mathcal{M}^{[27]} = 2(N_c + 1) \mathcal{M}^{[27]}$). One finds:

$$\Xi_{\text{NLL}}^{(+)[1]} = -0.6169 x^2 - 6.536 x^3 - 0.8371 x^4 - 8.483 x^5 - 1.529 x^6 - 12.67 x^7 + 1.610 x^8 - 20.62 x^9 + 16.48 x^{10} - 35.98 x^{11} + 46.07 x^{12} - 74.04 x^{13} + \mathcal{O}(x^{14}), \quad (38)$$

$$\Xi_{\text{NLL}}^{(+)[27]} = 1.028 x^2 - 18.16 x^3 + 2.184 x^4 - 196.0 x^5 + 372.3 x^6 - 2821 x^7 + 9382 x^8 - 46494 x^9 + 180397 x^{10} - 797524 x^{11} + 3.239 \times 10^6 x^{12} - 1.374 \times 10^7 x^{13} + \mathcal{O}(x^{14}). \quad (39)$$

By using Padé approximants, it is possible to conclude [14] that the series has a finite radius of convergence, as is illustrated in fig. 3. We find asymptotic geometric progression with the powers of $-\frac{1}{2} C_2 \frac{\alpha_s}{\pi} L = -\frac{3}{2} x$ and $(C_2 - \frac{3}{8} C_1) \frac{\alpha_s}{\pi} L = -\frac{17}{4} x$ for the singlet and the 27 representations, respectively. In both cases the series displays sign-oscillations, indicating that once resummed, it could be extrapolated beyond its radius of convergence.




Figure 3: Partial sums of the amplitude coefficients $\Xi_{\text{NLL}}^{(+,\ell)}$, up to 13th order, for the singlet (left plot) and the 27 colour representation (right plot). The horizontal axis x represents $\frac{\alpha_s}{\pi}L$. The dashed vertical line represents the radius of convergence, R , determined by the pole closest to $x = 0$, using Padé approximants. Figure taken from [15] under the Creative Commons Attribution 4.0 International License.

4 Three-Reggeon cut

We turn now to considering the odd amplitude at NNLL accuracy, i.e., the tower of coefficients $\mathcal{M}^{(-,n,n-2)}$. This part of the amplitude is particularly interesting from the point of view of Regge theory, because it involves a Regge cut in the real part of the amplitude, and it has quite different factorization properties compared to the two-Reggeon cut discussed so far, which contributes to the imaginary part of the amplitude. In this respect, an aspect of particular interest is the mixing between the single- and triple-Reggeon exchange, as discussed at the end of section 1, for which we refer the reader to [24]. In this talk we focus on the calculation of $\mathcal{M}^{(-,n,n-2)}$, which follows from the expansion to NNLL accuracy of

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-)} = \langle \psi_j^{(-)} | e^{-\hat{H}L} | \psi_i^{(-)} \rangle = D_i(t) D_j(t) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} + \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-),\text{MRS}}, \quad (40)$$

where the second equality follows from eq. (8) and the definition of reduced amplitude in eq. (15). As shown in [23, 24], this matrix element can be written to all orders as the sum of four contributions. In general, the NNLL contribution to the odd amplitude at n -loop is proportional to L^{n-2} , which is obtained from $n - 2$ repeated actions of the Balitsky-JIMWLK Hamiltonian H . The diagonal transitions $H_{k \rightarrow k}$ in eq. (13) are $\mathcal{O}(\alpha_s)$, while next-to-diagonal transitions $H_{k \rightarrow k \pm 2n}$ with $n = 1$ are $\mathcal{O}(\alpha_s^2)$. Noting that $|\psi_{i,n}\rangle$ is $\mathcal{O}(g_s^{n-1})$, there are four different types of contributions $\mathcal{O}(\alpha_s^n L^{n-2})$:

$$\begin{aligned} & \langle \psi_{j,3} | \hat{H}_{3 \rightarrow 3}^k | \psi_{i,3} \rangle, & \langle \psi_{j,1} | \hat{H}_{3 \rightarrow 1} \hat{H}_{3 \rightarrow 3}^{k-2} \hat{H}_{1 \rightarrow 3} | \psi_{i,1} \rangle, \\ & \langle \psi_{j,1} | \hat{H}_{3 \rightarrow 1} \hat{H}_{3 \rightarrow 3}^{k-1} | \psi_{i,3} \rangle, & \langle \psi_{j,3} | \hat{H}_{3 \rightarrow 3}^{k-1} \hat{H}_{1 \rightarrow 3} | \psi_{i,1} \rangle, \end{aligned} \quad (41)$$

where the two transitions in the second line of eq. (41) are related by target-projectile symmetry. These contributions can be expressed in a compact form by making the power-counting in the strong coupling manifest. To this end we introduce normalized Reggeon projectile states $|i_n\rangle$ (and similarly $|j\rangle$ for the target) and a normalized reduced Hamiltonian \tilde{H} , defined as

$$(r_\Gamma \alpha_s)^{(n-1)/2} |i_n\rangle \equiv |\psi_{i,n}\rangle, \quad \left(\frac{\alpha_s}{\pi} r_\Gamma\right)^{1+n} \pi^n \tilde{H}_{k \rightarrow k+2n} \equiv \hat{H}_{k \rightarrow k+2n}, \quad (42)$$

where we introduce rescalings by the constant r_Γ defined in eq. (2), and in turn we expand $|i_n\rangle = |i_n\rangle^{\text{LO}} + \sum_{k=1}^\infty \left(\frac{\alpha_s}{\pi}\right)^k |i_n\rangle^{\text{N}^k\text{LO}}$. With this notation, the four terms in eq. (41) give rise to

the NNLL odd amplitude as follows:

$$\begin{aligned} \frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-), \text{NNLL}} &= \pi^2 \left[\sum_{\ell=0}^{\infty} \frac{(-L)^\ell}{\ell!} \left(\frac{\alpha_s}{\pi} r_\Gamma \right)^{\ell+2} \langle j_3 | \tilde{H}_{3 \rightarrow 3}^\ell | i_3 \rangle \right. \\ &+ \sum_{\ell=1}^{\infty} \frac{(-L)^\ell}{\ell!} \left(\frac{\alpha_s}{\pi} r_\Gamma \right)^{\ell+2} \left[\langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{3 \rightarrow 3}^{\ell-1} | i_3 \rangle + \langle j_3 | \tilde{H}_{3 \rightarrow 3}^{\ell-1} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle \right] \\ &\left. + \sum_{\ell=2}^{\infty} \frac{(-L)^\ell}{\ell!} \left(\frac{\alpha_s}{\pi} r_\Gamma \right)^{\ell+2} \langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{3 \rightarrow 3}^{\ell-2} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle \right]^{\text{LO}} + \left(\frac{\alpha_s}{\pi} \right)^2 \langle j_1 | i_1 \rangle^{\text{NNLO}}. \end{aligned} \quad (43)$$

Notice that all but the last term are described within the leading order (LO) formalism, where the matter content of the theory is irrelevant. The matrix elements are given in terms of LO projectile/target states with either one or three Reggeons, which in turn are evolved by means of the LO Balitsky-JIMWLK Hamiltonian. The two-loop contribution $\langle j_1 | i_1 \rangle^{\text{NNLO}}$, instead, needs to be extracted from two-loop amplitudes, see ref. [12], but it does not enter the calculation of reduced amplitudes to higher orders. Apart from $\langle j_1 | i_1 \rangle^{\text{NNLO}}$, eq. (43) is therefore universal, in that it applies in any gauge theory, fully governed by the LO Balitsky-JIMWLK evolution of infinite Wilson lines in eq. (12).

Eq. (43) has characteristic analytic properties. First of all, it is interesting to notice that the NNLL amplitude is proportional, to all orders, to a factor of π^2 , if we exclude the two-loop single-Reggeon contribution. This is a characteristic feature of the three-Reggeon cut, playing a role similar to the universal factor of $i\pi$ appearing in case of the two-Reggeon cut in eqs. (17) and (31). Furthermore, as in case of the two-Reggeon cut eq. (31), based on form of the LO Balitsky-JIMWLK Hamiltonian, we expect the coefficients $\mathcal{M}^{(-,n,n-2)}$ to have maximal, uniform transcendent weight, when the dimensional regularization parameter ϵ is counted as having weight -1 .

Eq. (43) can be calculated iteratively. Compared to the computation of the two-Reggeon cut discussed in section 3, the integrals involved in the calculation of eq. (43) are of similar complexity; however, the colour structure is more involved, and it is non trivial to express it in terms of colour operators acting on the tree level. For this purpose, specific techniques and new colour identities need to be developed. This task has been carried out in [23, 24] where the odd amplitude has been determined explicitly up to four loops. For a detailed discussion of these techniques we refer the reader to these works. In the following we focus on presenting the result for the odd amplitude up to four loops.

The two and three-loops amplitude coefficients were already obtained in [12]. Using the notation of eq. (43), at the two-loop amplitude has two contributions:

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-,2,0)} = \langle j_1 | i_1 \rangle^{\text{NNLO}} + \pi^2 r_\Gamma^2 \langle j_3 | i_3 \rangle. \quad (44)$$

The single-Reggeon transition, comparing with eq. (40), reads

$$\langle j_1 | i_1 \rangle^{\text{NNLO}} = \left[D_i^{(2)} + D_j^{(2)} + D_i^{(1)} D_j^{(1)} \right] \langle j_1 | i_1 \rangle. \quad (45)$$

The three-reggeon transition gives

$$\langle j_3 | i_3 \rangle = -\frac{1}{8} \left(\frac{1}{\epsilon^2} - 6\epsilon \hat{\zeta}_3 + \mathcal{O}(\epsilon^3) \right) \left[\left(\mathbf{T}_{s-u}^2 \right)^2 - \frac{C_A^2}{12} \right] \langle j_1 | i_1 \rangle, \quad (46)$$

where the tree-level factor $\langle j_1 | i_1 \rangle$ reads, according to the normalization of the states in eq. (42), $\langle j_1 | i_1 \rangle = \frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{\text{tree}}$. Furthermore, $\hat{\zeta}_3 = \zeta_3 + \frac{3}{2}\epsilon \zeta_4 - \frac{5}{2}\epsilon^3 \zeta_6 + \mathcal{O}(\epsilon^5)$. Summing the two contributions, one has

$$\hat{\mathcal{M}}^{(-,2,0)} = \left[D_i^{(2)} + D_j^{(2)} + D_i^{(1)} D_j^{(1)} + \pi^2 r_\Gamma^2 S^{(2)}(\epsilon) \left(\left(\mathbf{T}_{s-u}^2 \right)^2 - \frac{1}{12} C_A^2 \right) \right] \mathcal{M}^{\text{tree}}, \quad (47)$$




Figure 4: Representation of transitions involving Multi-Reggeon States (MRS), which contribute to the odd-signature amplitude at four loops. Figure taken from [23] under the Creative Commons Attribution 4.0 International License.

where

$$S^{(2)}(\epsilon) = -\frac{1}{8\epsilon^2} + \frac{3}{4}\epsilon\zeta_3 + \frac{9}{8}\epsilon^2\zeta_4 + \mathcal{O}(\epsilon^3). \quad (48)$$

At three loops the action of the Hamiltonian kicks-in, and one has to take into account three MRS contributions:

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-,3,1)} = -\pi^2 r_\Gamma^3 \left[\langle j_3 | \tilde{H}_{3 \rightarrow 3} | i_3 \rangle + (\langle j_1 | \tilde{H}_{3 \rightarrow 1} | i_3 \rangle + \langle j_3 | \tilde{H}_{1 \rightarrow 3} | i_1 \rangle) \right]. \quad (49)$$

The various matrix elements can be evaluated to give

$$\begin{aligned} \langle j_3 | \tilde{H}_{3 \rightarrow 3} | i_3 \rangle &= -24 S_C^{(3)}(\epsilon) \left(\frac{d_{AR_i}}{N_{R_i} C_i} + \frac{d_{AR_j}}{N_{R_j} C_j} \right) \langle j_1 | i_1 \rangle \\ &+ \left[S_A^{(3)}(\epsilon) \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] + S_B^{(3)}(\epsilon) [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_{s-u}^2 + S_C^{(3)}(\epsilon) C_A^3 \right] \langle j_1 | i_1 \rangle, \end{aligned} \quad (50)$$

and

$$\langle j_3 | \tilde{H}_{1 \rightarrow 3} | i_3 \rangle + \langle j_1 | \tilde{H}_{3 \rightarrow 1} | i_3 \rangle = 24 S_C^{(3)}(\epsilon) \left(\frac{d_{AR_i}}{N_{R_i} C_i} + \frac{d_{AR_j}}{N_{R_j} C_j} \right) \langle j_1 | i_1 \rangle, \quad (51)$$

where d_{AR_i} and d_{AR_j} are quartic Casimir associated to the projectile and target:

$$d_{AR_i} = \frac{1}{4!} \sum_{\sigma \in S_4} \text{Tr} [F^{\sigma(a)} F^{\sigma(b)} F^{\sigma(c)} F^{\sigma(d)}] \text{Tr} [\mathbf{T}_i^a \mathbf{T}_i^b \mathbf{T}_i^c \mathbf{T}_i^d], \quad (52)$$

and the functions $S_i^{(3)}(\epsilon)$ arise from the evaluation of loop integrals in $2 - 2\epsilon$ dimensions:

$$S_A^{(3)}(\epsilon) = \frac{1}{48\epsilon^3} + \frac{37\hat{\zeta}_3}{24} + \mathcal{O}(\epsilon^2), \quad (53a)$$

$$S_B^{(3)}(\epsilon) = \frac{1}{24\epsilon^3} + \frac{\hat{\zeta}_3}{12} + \mathcal{O}(\epsilon^2), \quad (53b)$$

$$S_C^{(3)}(\epsilon) = -\frac{1}{432} \left(\frac{1}{2\epsilon^3} - 35\hat{\zeta}_3 + \mathcal{O}(\epsilon^2) \right). \quad (53c)$$

It is interesting to note that the sum of the transitions $1 \rightarrow 3$ and $3 \rightarrow 1$ does not involve a matrix in colour space, but is simply proportional to $\langle j_1 | i_1 \rangle$. This is due to the fact that colour is carried by a single Reggeon on either the target or projectile sides. This property holds for all terms in the second and third lines of (43) at any order. Another intriguing feature [23, 24] is that the quartic Casimir contributions of eq. (51), exactly cancel an identical contribution in the $3 \rightarrow 3$ transition. One then finds

$$\hat{\mathcal{M}}^{(-,3,1)} = -\pi^2 r_\Gamma^3 \left[S_A^{(3)}(\epsilon) \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] + S_B^{(3)}(\epsilon) [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_{s-u}^2 + S_C^{(3)}(\epsilon) C_A^3 \right] \mathcal{M}^{\text{tree}}. \quad (54)$$

Next, we consider the four-loop amplitude, determined for the first time in [23]. In this case we need to take into account two consecutive applications of the Balitsky-JIMWLK Hamiltonian, which gives rise for the first time to all four MRS terms in eq. (46). One has

$$\frac{i}{2s} \hat{\mathcal{M}}_{ij \rightarrow ij}^{(-,4,2)} = \frac{\pi^2 r_\Gamma^4}{2} \left[\langle j_3 | \tilde{H}_{3 \rightarrow 3}^2 | i_3 \rangle + (\langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{3 \rightarrow 3} | i_3 \rangle + \langle j_3 | \tilde{H}_{3 \rightarrow 3} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle) + \langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle \right], \quad (55)$$

The first term, i.e. the double insertion of $\tilde{H}_{3 \rightarrow 3}$ in the three-Reggeon ladder, gives the most complicated contribution, which is schematically represented by the two diagrams on the left in fig. 4. The first diagrams involve BFKL evolution applied twice to the same pair of Reggeons, while in the second diagram evolution is applied at each stage to a different pair. One obtains

$$\langle j_3 | \tilde{H}_{3 \rightarrow 3}^2 | i_3 \rangle = \left[\frac{\mathbf{C}^{(4,-4)}}{\epsilon^4} + \frac{\hat{\zeta}_3}{\epsilon} \mathbf{C}^{(4,-1)} + \mathcal{O}(\epsilon) \right] \langle j_1 | i_1 \rangle, \quad (56)$$

where we have defined the following operators in colour space

$$\begin{aligned} \mathbf{C}^{(4,-4)} = & \frac{1}{432} \left[\frac{d_{AA}}{N_A} - 3C_A \left(\frac{d_{AR_i}}{N_{R_i} C_i} + \frac{d_{AR_j}}{N_{R_j} C_j} \right) + \frac{C_A^4}{12} \right] - \frac{1}{192} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 \\ & + \frac{1}{96} [\mathbf{T}_{s-u}^2, [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2]] \mathbf{T}_t^2 + \frac{7}{576} \mathbf{T}_t^2 [(\mathbf{T}_{s-u}^2)^2, \mathbf{T}_t^2] - \frac{5}{192} \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2, \end{aligned} \quad (57)$$

$$\begin{aligned} \mathbf{C}^{(4,-1)} = & -\frac{101}{216} \left[\frac{d_{AA}}{N_A} - \frac{312}{101} C_A \left(\frac{d_{AR_i}}{N_{R_i} C_i} + \frac{d_{AR_j}}{N_{R_j} C_j} \right) + \frac{211 C_A^4}{2424} \right] + \frac{101}{96} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 \\ & + \frac{49}{48} [\mathbf{T}_{s-u}^2, [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2]] \mathbf{T}_t^2 - \frac{47}{288} \mathbf{T}_t^2 [(\mathbf{T}_{s-u}^2)^2, \mathbf{T}_t^2] - \frac{49}{48} \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2. \end{aligned} \quad (58)$$

The terms in square brackets in eqs. (57) and (58) are leading in the planar limit, while all other terms, involving commutators of \mathbf{T}_{s-u}^2 and \mathbf{T}_t^2 , are suppressed in N_c , and result in mixing of the octet exchange with other colour.

The third and fourth diagrams in fig. 4 correspond to the $3 \rightarrow 1$ and $1 \rightarrow 3$ transitions given by the second and third terms in eq. (55). The last diagram correspond to the $1 \rightarrow 1$ transition mediated by three Reggeons, i.e. the last term in eq. (55). These contributions are all proportional to the unit matrix in colour space. For the $3 \rightarrow 1$ and $1 \rightarrow 3$ transitions we obtain [23, 24]

$$\begin{aligned} \langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{3 \rightarrow 3} | i_3 \rangle + \langle j_3 | \tilde{H}_{3 \rightarrow 3} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle = \\ = \frac{C_A}{144} \left(\frac{d_{AR_i}}{N_{R_i} C_i} + \frac{d_{AR_j}}{N_{R_j} C_j} \right) \left(\frac{1}{\epsilon^4} - \frac{208 \hat{\zeta}_3}{\epsilon} + \mathcal{O}(\epsilon) \right) \langle j_1 | i_1 \rangle, \end{aligned} \quad (59)$$

and for the $1 \rightarrow 1$ transition

$$\langle j_1 | \tilde{H}_{3 \rightarrow 1} \tilde{H}_{1 \rightarrow 3} | i_1 \rangle = \left[-\frac{1}{432 \epsilon^4} \left(\frac{d_{AA}}{N_A} + \frac{C_A^4}{12} \right) + \frac{55 \hat{\zeta}_3}{108 \epsilon} \left(\frac{d_{AA}}{N_A} + \frac{101 C_A^4}{1320} \right) \right] \langle j_1 | i_1 \rangle. \quad (60)$$

Summing the contributions in eqs. (56), (59) and (60) we get

$$\hat{\mathcal{M}}^{(-,4,2)} = \frac{r_\Gamma^4 \pi^2}{2} \left[\frac{1}{\epsilon^4} \mathbf{K}^{(4)} + \left(\frac{1}{\epsilon} \zeta_3 + \frac{3}{2} \zeta_4 \right) \mathbf{K}^{(1)} + \mathcal{O}(\epsilon) \right] \mathcal{M}^{\text{tree}}, \quad (61)$$

where the colour structures are defined by

$$\begin{aligned} \mathbf{K}^{(4)} = & \frac{1}{96} [\mathbf{T}_{s-u}^2, [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2]] \mathbf{T}_t^2 + \frac{7}{576} \mathbf{T}_t^2 [(\mathbf{T}_{s-u}^2)^2, \mathbf{T}_t^2] \\ & - \frac{1}{192} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 - \frac{5}{192} \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2, \end{aligned} \quad (62a)$$

$$\begin{aligned} \mathbf{K}^{(1)} = & \frac{49}{48} [\mathbf{T}_{s-u}^2, [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2]] \mathbf{T}_t^2 - \frac{47}{288} \mathbf{T}_t^2 [(\mathbf{T}_{s-u}^2)^2, \mathbf{T}_t^2] \\ & + \frac{101}{96} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 - \frac{49}{48} \mathbf{T}_{s-u}^2 [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 + \frac{1}{24} \left(\frac{d_{AA}}{N_A} - \frac{C_A^4}{24} \right). \end{aligned} \quad (62b)$$

Eq. (62b) depends on the quartic Casimir d_{AA} , irrespective of the representation associated to the target and to the projectile. The expressions in eq. (62a) and (62b) imply that the reduced amplitude $\hat{\mathcal{M}}^{(-,4,2)}$ is non-planar, which is expected on general grounds, [24], based on its direct connection to the Regge cut. This is because eq. (62a) is written in terms of the commutator $[\mathbf{T}_t^2, \mathbf{T}_{s-u}^2]$, which is subleading in N_c , and the only non-commutator term is also subleading, given that

$$\frac{d_{AA}}{N_A} - \frac{C_A^4}{24} = 0 \cdot N_c^4 + \frac{3}{2} N_c^2. \quad (63)$$

We note that, as at three loops, all terms arising from $3 \rightarrow 1$ and $1 \rightarrow 3$ transitions, which are proportional to the quartic Casimir invariants d_{AR_i} and d_{AR_j} , cancel in the sum with the $3 \rightarrow 3$ amplitude, to all orders in ϵ . These contributions are separately leading in the planar limit, thus their cancellation is essential for the complete $\hat{\mathcal{M}}^{(-,4,2)}$ to be non-planar. Based on this observation, we conjecture that transition amplitudes connecting three-Reggeon states with a single Reggeon cancel from the reduced amplitude $\hat{\mathcal{M}}^{(-,NNLL)}$ to all perturbative orders.

With the odd amplitude coefficients $\mathcal{M}^{(-,n,n-2)}$ evaluated up to four-loop, several aspects concerning the analytic structure of two-parton amplitudes can be investigated. We refer to [24] for a thorough analysis. Here we limit ourselves to one application, namely, the extraction of the soft anomalous dimension, (see refs. [24, 44]) which can be obtained by combining the knowledge of the amplitude in the high-energy limit with the infrared factorization theorem, as discussed for the even amplitude at NLL accuracy. In this case, the calculation of the tower $\mathcal{M}^{(-,\ell,\ell-2)}$ gives access to $\Delta^{(+,\ell,\ell-2)}$. Here we have determined $\mathcal{M}^{(-,\ell,\ell-2)}$ up to $\ell = 4$, therefore we obtain [12, 23, 24]:

$$\Delta^{(+,3,1)} = 0, \quad (64)$$

$$\Delta^{(+,4,2)} = \zeta_2 \zeta_3 \left(\frac{d_{AA}}{N_A} - \frac{C_A^4}{24} - \frac{1}{4} \mathbf{T}_t^2 [(\mathbf{T}_{s-u}^2)^2, \mathbf{T}_t^2] + \frac{3}{4} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 \right). \quad (65)$$

Thanks to the calculation of the four-loop amplitude, we have obtained the first non-trivial real contribution to the tower $\Delta^{(+,\ell,\ell-2)}$. For completeness, we recall that also the imaginary contribution $\Delta^{(-,3,1)}$ has been determined from the three-loop exact calculation in general kinematic, [63], and reads $\Delta^{(3,1)} = i\pi \frac{\zeta_3}{4} [\mathbf{T}_t^2, [\mathbf{T}_t^2, \mathbf{T}_{s-u}^2]]$ [12]. We refer to [24] for a complete list of known coefficients $\Delta^{(\pm,\ell,m)}$ in the high-energy limit.

With the result for the soft anomalous dimension, it is easy to invert eq. (32) and obtain a prediction for the finite remainder of the amplitude at four loop. To this end, given the reduced amplitude in eq. (61), one needs first to restore the single Reggeon contribution which had been removed according to eq. (15). The finite remainder then reads

$$\mathcal{H} = Z_i Z_j Z^{-1} e^{\alpha_s \mathbf{T}_i^2 L} \hat{\mathcal{M}}. \quad (66)$$

Upon expansion the four-loop contribution at $\mathcal{O}(\epsilon^0)$ we get

$$\mathcal{H}^{(-,4,2)} = \left\{ \frac{C_A^2}{2} \left(\hat{\alpha}_g^{(2,0)} \right)^2 + \frac{3}{16} \zeta_4 \zeta_2 C_{\Delta}^{(+,4,2)} + \mathcal{O}(\epsilon) \right\} \mathcal{M}^{\text{tree}}, \quad (67)$$

where $\hat{\alpha}_g^{(2,0)} = C_A \left(\frac{101}{108} - \frac{\zeta_3}{8} \right) - \frac{7}{27} T_R n_f$ is the $\mathcal{O}(\epsilon^0)$ term of the two-loop Regge trajectory, and

$$C_{\Delta}^{(+,4,2)} \equiv \frac{d_{AA}}{N_A} - \frac{C_A^4}{24} + \frac{1}{4} \mathbf{T}_t^2 [\mathbf{T}_t^2, (\mathbf{T}_{s-u}^2)^2] + \frac{3}{4} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2. \quad (68)$$

Eq. (67) conveniently displays the theory dependence of $\mathcal{H}^{(-,4,2)}$, which is in fact restricted to $\hat{\alpha}_g^{(2,0)}$. We can thus obtain explicit result in QCD:

$$\mathcal{H}_{\text{QCD}}^{(-,4,2)} = \left\{ C_A^2 T_R^2 n_f^2 \frac{49}{1458} + C_A^3 T_R n_f \left(\frac{7\zeta_3}{216} - \frac{707}{2916} \right) + C_A^4 \left(\frac{\zeta_3^2}{128} - \frac{101\zeta_3}{864} + \frac{10201}{23328} \right) + \frac{3}{16} \zeta_4 \zeta_2 C_{\Delta}^{(+,4,2)} + \mathcal{O}(\epsilon) \right\} \mathcal{M}^{\text{tree}}, \quad (69)$$

which is a new result. Furthermore, according to the principle of maximum transcendentality, in $\mathcal{N} = 4$ SYM one has:

$$\mathcal{H}_{\text{SYM}}^{(-,4,2)} = \left\{ \frac{C_A^4}{128} \zeta_3^2 + \frac{3}{16} \zeta_4 \zeta_2 C_{\Delta}^{(+,4,2)} + \mathcal{O}(\epsilon) \right\} \mathcal{M}^{\text{tree}}. \quad (70)$$

While for $\mathcal{N} = 4$ SYM the planar limit, $N_c \rightarrow \infty$, is already known [64–66], the non-planar correction, i.e. the second term in eq. (70) is new.

5 Conclusion

In this talk we have summarized recent progress concerning the calculation of amplitudes in the high-energy limit. Within the shockwave formalism, scattering amplitudes can be calculated as expectation values of Wilson lines associated to the external partons [3, 67]. In this framework one can identify an effective degree of freedom, identified as a ‘‘Reggeon’’ [11], which plays a central role in determining the factorization properties of the amplitude in the high-energy limit. Upon expansion in perturbation theory, the amplitude can be described in terms of transition amplitudes between states labelled by the number Reggeons, which needs to be evolved to equal rapidity by means of the BFKL equation and its generalization, the Balitsky-JIMWLK equation [11, 12]. While it is not known how to solve these equations exactly, it is possible to develop iterative solutions, which allows one to calculate the amplitude to high order in perturbation theory.

In this context, we have discussed recent calculations of $2 \rightarrow 2$ scattering amplitudes in the high-energy limit. Refs. [13, 14] focus on the imaginary part of the amplitude at NLL accuracy in the high-energy logarithm. Within the shockwave formalism this can be expressed as a class of ladder graphs involving two Reggeons with any number of rungs, which arise from the iterative solution of the BFKL equation. We have shown that the singular part [13] can be associated to a kinematic configuration in which one of the two Reggeons become soft. Within this approximation the evolution equation simplifies, such that the singular part of the amplitude can be calculated to all order and resummed in a closed form involving only gamma functions. We have then discussed the calculation of the finite contribution, for which a general algorithm has been set up, based on an iterative solution of the BFKL equation in two transverse dimensions, where the calculation is greatly simplified by the fact that the two-dimensional two-Reggeon wavefunction consists of single-valued HPLs [14]. Explicit results

have been presented up to 13 loops order. Finite corrections to the amplitude have a more complicated pattern, involving multiple zeta values, and hence they cannot be resummed in terms of gamma functions.

We moved then to discuss the calculation of the real part of the two-parton scattering amplitude at NNLL accuracy, recently investigated in [23, 24]. We have shown that the amplitude can be expressed to all order as an iterated solution of the Balitsky-JIMWLK equation, with a simple diagrammatic interpretation in terms of ladder graphs involving three Reggions, with any number of rungs. Remarkably, the calculation of the entire tower requires only the LO Balitsky-JIMWLK formalism, while the complete odd amplitude at NNLL requires only two additional inputs, given by the single-Reggeon impact factors at two loops and Regge trajectory at three loops, which can be determined by matching with the full amplitude. In [23, 24] the signature-odd amplitude has been calculated for the first time to four loops, where it was expressed in a form valid for scattering of particles in general colour representations. The four-loop result, eq. (61), is entirely non-planar, a property that can be understood based on its direct connection to the Regge cut [24], and it involves a purely adjoint quartic Casimir.

Finally, we have briefly discussed one application of these calculations, which exploit the fact that it is possible to apply infrared factorization to the scattering amplitude calculated in the high-energy limit, to extract the soft anomalous dimension within the same limit. In this context, the calculation of the even NLL amplitude discussed above made it possible to extract the soft anomalous dimension to all orders, which, at this logarithmic accuracy, is found to have an infinite radius of convergence. From the computation of the odd amplitude at NNLL accuracy we determine the soft anomalous dimension at this logarithmic accuracy to four loop. Ref. [51] analyzed the colour structure of the soft anomalous dimension, incorporating the recently computed four-loop cusp anomalous dimension [68, 69], which introduces quartic Casimirs. Our results show that the soft anomalous dimension contains additional quartic Casimirs, beyond those associated with the cusp.

The knowledge of the soft anomalous dimension in the high-energy limit is extremely useful for bootstrap approaches, see for instance [22]. In practice, the information obtained in particular kinematic limits, such as the Regge limit, can be used to reconstruct the structure of infrared divergences in full kinematic. Preliminary steps in this direction, using the new information obtained in [13, 14, 23], have been carried out in [24], and have been discussed in another talk in this conference, [44], to which we refer for further details.

Acknowledgements

We would like to thank Simon Caron-Huot and Josha Reichel for stimulating collaborations on some of the topics discussed in this talk.

Funding information EG, GF and NM are supported by the STFC Consolidated Grant “Particle Physics at the Higgs Centre”. GF is supported by the ERC Starting Grant 715049 “QCD-forfuture” with Principal Investigator Jennifer Smillie. CM’s work is supported by the Italian Ministry of University and Research (MIUR), grant PRIN 20172LNEEZ. LV is supported by Fellini, Fellowship for Innovation at INFN, funded by the European Union’s Horizon 2020 research programme under the Marie Skłodowska-Curie Cofund Action, grant agreement no. 754496.

References

- [1] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, *The Pomeron Singularity in Nonabelian Gauge Theories*, Sov. Phys. JETP **45**, 199 (1977), [Zh. Eksp. Teor. Fiz. **72**, 377 (1977)].
- [2] I. I. Balitsky and L. N. Lipatov, *The Pomeron Singularity in Quantum Chromodynamics*, Sov. J. Nucl. Phys. **28**, 822 (1978), [Yad. Fiz. **28**, 1597 (1978)].
- [3] I. Balitsky, *Operator expansion for high-energy scattering*, Nucl. Phys. B **463**, 99 (1996), doi:[10.1016/0550-3213\(95\)00638-9](https://doi.org/10.1016/0550-3213(95)00638-9).
- [4] Y. V. Kovchegov, *Small- x F_2 structure function of a nucleus including multiple Pomeron exchanges*, Phys. Rev. D **60**, 034008 (1999), doi:[10.1103/PhysRevD.60.034008](https://doi.org/10.1103/PhysRevD.60.034008).
- [5] J. Jalilian-Marian, A. Kovner, L. McLerran and H. Weigert, *Intrinsic glue distribution at very small x* , Phys. Rev. D **55**, 5414 (1997), doi:[10.1103/PhysRevD.55.5414](https://doi.org/10.1103/PhysRevD.55.5414).
- [6] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *Wilson renormalization group for low- x physics: Towards the high density regime*, Phys. Rev. D **59**, 014014 (1998), doi:[10.1103/PhysRevD.59.014014](https://doi.org/10.1103/PhysRevD.59.014014).
- [7] E. Iancu, A. Leonidov and L. McLerran, *The renormalization group equation for the color glass condensate*, Phys. Lett. B **510**, 133 (2001), doi:[10.1016/S0370-2693\(01\)00524-X](https://doi.org/10.1016/S0370-2693(01)00524-X).
- [8] P. D. B. Collins, *An Introduction to Regge Theory and High-Energy Physics*, Cambridge Monographs on Mathematical Physics, Cambridge Univ. Press, Cambridge, UK, ISBN 9780521110358 (2009).
- [9] L. J. Dixon, C. Duhr and J. Pennington, *Single-valued harmonic polylogarithms and the multi-Regge limit*, J. High Energy Phys. **10**, 074 (2012), doi:[10.1007/JHEP10\(2012\)074](https://doi.org/10.1007/JHEP10(2012)074).
- [10] L. J. Dixon, J. M. Drummond, C. Duhr and J. Pennington, *The four-loop remainder function and multi-Regge behavior at NNLLA in planar $\mathcal{N} = 4$ super-Yang-Mills theory*, J. High Energy Phys. **06**, 116 (2014), doi:[10.1007/JHEP06\(2014\)116](https://doi.org/10.1007/JHEP06(2014)116).
- [11] S. Caron-Huot, *When does the gluon reggeize?*, J. High Energy Phys. **05**, 093 (2015), doi:[10.1007/JHEP05\(2015\)093](https://doi.org/10.1007/JHEP05(2015)093).
- [12] S. Caron-Huot, E. Gardi and L. Vernazza, *Two-parton scattering in the high-energy limit*, J. High Energy Phys. **06**, 016 (2017), doi:[10.1007/JHEP06\(2017\)016](https://doi.org/10.1007/JHEP06(2017)016).
- [13] S. Caron-Huot, E. Gardi, J. Reichel and L. Vernazza, *Infrared singularities of QCD scattering amplitudes in the Regge limit to all orders*, J. High Energy Phys. **03**, 098 (2018), doi:[10.1007/JHEP03\(2018\)098](https://doi.org/10.1007/JHEP03(2018)098).
- [14] S. Caron-Huot, E. Gardi, J. Reichel and L. Vernazza, *Two-parton scattering amplitudes in the Regge limit to high loop orders*, J. High Energy Phys. **08**, 116 (2020), doi:[10.1007/JHEP08\(2020\)116](https://doi.org/10.1007/JHEP08(2020)116).
- [15] E. Gardi, S. Caron-Huot, J. Reichel and L. Vernazza, *The High-Energy Limit of 2-to-2 Partonic Scattering Amplitudes*, Proc. Sci. **375**, 050 (2019), doi:[10.22323/1.375.0050](https://doi.org/10.22323/1.375.0050).
- [16] V. Del Duca, S. Druc, J. M. Drummond, C. Duhr, F. Dulat, R. Marzucca, G. Papathanasiou and B. Verbeek, *All-Order Amplitudes at any Multiplicity in the Multi-Regge Limit*, Phys. Rev. Lett. **124**, 161602 (2020), doi:[10.1103/PhysRevLett.124.161602](https://doi.org/10.1103/PhysRevLett.124.161602).

- [17] J. Bartels, *Analytic structure of the 8-point scattering amplitude in multi-Regge kinematics in $N=4$ SYM: conformal Regge pole and Regge cut contributions*, [arXiv:2005.08818](https://arxiv.org/abs/2005.08818).
- [18] S. Caron-Huot, L. J. Dixon, A. McLeod and M. von Hippel, *Bootstrapping a Five-Loop Amplitude Using Steinmann Relations*, *Phys. Rev. Lett.* **117**, 241601 (2016), doi:[10.1103/PhysRevLett.117.241601](https://doi.org/10.1103/PhysRevLett.117.241601).
- [19] S. Caron-Huot, L. J. Dixon, F. Dulat, M. von Hippel, A. J. McLeod and G. Papathanasiou, *Six-Gluon amplitudes in planar $\mathcal{N} = 4$ super-Yang-Mills theory at six and seven loops*, *J. High Energy Phys.* **08**, 016 (2019), doi:[10.1007/JHEP08\(2019\)016](https://doi.org/10.1007/JHEP08(2019)016).
- [20] V. Del Duca, C. Duhr, E. Gardi, L. Magnea and C. D. White, *The infrared structure of gauge theory amplitudes in the high-energy limit*, *J. High Energy Phys.* **12**, 021 (2011), doi:[10.1007/JHEP12\(2011\)021](https://doi.org/10.1007/JHEP12(2011)021).
- [21] J. M. Henn and B. Mistlberger, *Four-Gluon Scattering at Three Loops, Infrared Structure, and the Regge Limit*, *Phys. Rev. Lett.* **117**, 171601 (2016), doi:[10.1103/PhysRevLett.117.171601](https://doi.org/10.1103/PhysRevLett.117.171601).
- [22] Ø. Almelid, C. Duhr, E. Gardi, A. McLeod and C. D. White, *Bootstrapping the QCD soft anomalous dimension*, *J. High Energy Phys.* **09**, 073 (2017), doi:[10.1007/JHEP09\(2017\)073](https://doi.org/10.1007/JHEP09(2017)073).
- [23] G. Falcioni, E. Gardi, C. Milloy and L. Vernazza, *Climbing three-Reggeon ladders: Four-loop amplitudes in the high-energy limit in full color*, *Phys. Rev. D* **103**, L111501 (2021), doi:[10.1103/PhysRevD.103.L111501](https://doi.org/10.1103/PhysRevD.103.L111501).
- [24] G. Falcioni, E. Gardi, N. Maher, C. Milloy and L. Vernazza, *Scattering amplitudes in the Regge limit and the soft anomalous dimension through four loops*, to appear, (2021).
- [25] L. Lipatov, *Reggeization of the Vector Meson and the Vacuum Singularity in Nonabelian Gauge Theories*, *Sov. J. Nucl. Phys.* **23**, 338 (1976).
- [26] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, *Multi - Reggeon Processes in the Yang-Mills Theory*, *Sov. Phys. JETP* **44**, 443 (1976).
- [27] F. E. Low, *Model of the bare Pomeron*, *Phys. Rev. D* **12**, 163 (1975), doi:[10.1103/PhysRevD.12.163](https://doi.org/10.1103/PhysRevD.12.163).
- [28] S. Nussinov, *Colored-Quark Version of Some Hadronic Puzzles*, *Phys. Rev. Lett.* **34**, 1286 (1975), doi:[10.1103/PhysRevLett.34.1286](https://doi.org/10.1103/PhysRevLett.34.1286).
- [29] J. F. Gunion and D. E. Soper, *Quark-counting and hadron-size effects for total cross sections*, *Phys. Rev. D* **15**, 2617 (1977), doi:[10.1103/PhysRevD.15.2617](https://doi.org/10.1103/PhysRevD.15.2617).
- [30] L. N. Lipatov, *High Energy Asymptotics of Multi-Colour QCD and Exactly Solvable Lattice Models*, *JETP Lett.* **59**, 596 (1994), [arXiv:hep-th/9311037](https://arxiv.org/abs/hep-th/9311037).
- [31] L. D. Faddeev and G. P. Korchemsky, *High energy QCD as a completely integrable model*, *Phys. Lett. B* **342**, 311 (1995), doi:[10.1016/0370-2693\(94\)01363-H](https://doi.org/10.1016/0370-2693(94)01363-H).
- [32] S. É. Derkachov, G. P. Korchemsky and A. N. Manashov, *Noncompact Heisenberg spin magnets from high-energy QCD*, *Nucl. Phys. B* **617**, 375 (2001), doi:[10.1016/S0550-3213\(01\)00457-6](https://doi.org/10.1016/S0550-3213(01)00457-6).

- [33] S. É. Derkachov, G. P. Korchemsky, J. Kotański and A. N. Manashov, *Noncompact Heisenberg spin magnets from high-energy QCD II. Quantization conditions and energy spectrum*, Nucl. Phys. B **645**, 237 (2002), doi:[10.1016/S0550-3213\(02\)00842-8](https://doi.org/10.1016/S0550-3213(02)00842-8).
- [34] V. S. Fadin, R. Fiore, M. G. Kozlov and A. V. Reznichenko, *Proof of the multi-Regge form of QCD amplitudes with gluon exchanges in the NLA*, Phys. Lett. B **639**, 74 (2006), doi:[10.1016/j.physletb.2006.03.031](https://doi.org/10.1016/j.physletb.2006.03.031).
- [35] V. S. Fadin, M. G. Kozlov and A. V. Reznichenko, *Gluon Reggeization in Yang-Mills theories*, Phys. Rev. D **92**, 085044 (2015), doi:[10.1103/PhysRevD.92.085044](https://doi.org/10.1103/PhysRevD.92.085044).
- [36] V. S. Fadin, R. Fiore and M. I. Kotsky, *Gluon Reggeization in QCD in the next to leading order*, Phys. Lett. B **359**, 181 (1995), doi:[10.1016/0370-2693\(95\)01016-J](https://doi.org/10.1016/0370-2693(95)01016-J).
- [37] V. S. Fadin, R. Fiore and M. I. Kotsky, *Gluon Regge trajectory in the two-loop approximation*, Phys. Lett. B **387**, 593 (1996), doi:[10.1016/0370-2693\(96\)01054-4](https://doi.org/10.1016/0370-2693(96)01054-4).
- [38] V. S. Fadin, R. Fiore and A. Quartarolo, *Reggeization of the quark-quark scattering amplitude in QCD*, Phys. Rev. D **53**, 2729 (1996), doi:[10.1103/PhysRevD.53.2729](https://doi.org/10.1103/PhysRevD.53.2729).
- [39] J. Blümlein, V. Ravindran and W. L. van Neerven, *Gluon Regge trajectory in $O(\alpha_s^2)$* , Phys. Rev. D **58**, 091502 (1998), doi:[10.1103/PhysRevD.58.091502](https://doi.org/10.1103/PhysRevD.58.091502).
- [40] V. S. Fadin, *Particularities of the NNLLA BFKL*, AIP Conf. Proc. **1819**, 060003 (2017), doi:[10.1063/1.4977159](https://doi.org/10.1063/1.4977159).
- [41] V. Del Duca and E. W. Nigel Glover, *The high energy limit of QCD at two loops*, J. High Energy Phys. **10**, 035 (2001), doi:[10.1088/1126-6708/2001/10/035](https://doi.org/10.1088/1126-6708/2001/10/035).
- [42] V. Del Duca, G. Falcioni, L. Magnea and L. Vernazza, *Analyzing high-energy factorization beyond next-to-leading logarithmic accuracy*, J. High Energy Phys. **02**, 029 (2015), doi:[10.1007/JHEP02\(2015\)029](https://doi.org/10.1007/JHEP02(2015)029).
- [43] V. S. Fadin and L. N. Lipatov, *Reggeon cuts in QCD amplitudes with negative signature*, Eur. Phys. J. C **78**, 439 (2018), doi:[10.1140/epjc/s10052-018-5910-1](https://doi.org/10.1140/epjc/s10052-018-5910-1).
- [44] N. Maher, G. Falcioni, E. Gardi, C. Milloy and L. Vernazza, *The Soft Anomalous Dimension at four loops in the Regge Limit*, [arXiv:2111.01517](https://arxiv.org/abs/2111.01517).
- [45] I. A. Korchemskaya and G. P. Korchemsky, *High-energy scattering in QCD and cross singularities of Wilson loops*, Nucl. Phys. B **437**, 127 (1995), doi:[10.1016/0550-3213\(94\)00553-Q](https://doi.org/10.1016/0550-3213(94)00553-Q).
- [46] I. A. Korchemskaya and G. P. Korchemsky, *Evolution equation for gluon Regge trajectory*, Phys. Lett. B **387**, 346 (1996), doi:[10.1016/0370-2693\(96\)01016-7](https://doi.org/10.1016/0370-2693(96)01016-7).
- [47] G. P. Korchemsky and A. V. Radyushkin, *Renormalization of the Wilson loops beyond the leading order*, Nucl. Phys. B **283**, 342 (1987), doi:[10.1016/0550-3213\(87\)90277-X](https://doi.org/10.1016/0550-3213(87)90277-X).
- [48] E. Gardi and L. Magnea, *Factorization constraints for soft anomalous dimensions in QCD scattering amplitudes*, J. High Energy Phys. **03**, 079 (2009), doi:[10.1088/1126-6708/2009/03/079](https://doi.org/10.1088/1126-6708/2009/03/079).
- [49] E. Gardi and L. Magnea, *Infrared singularities in QCD amplitudes*, Il Nuovo Cim. **5-6**, 137 (2010), doi:[10.1393/ncc/i2010-10528-x](https://doi.org/10.1393/ncc/i2010-10528-x).

- [50] T. Becher and M. Neubert, *Infrared Singularities of Scattering Amplitudes in Perturbative QCD*, Phys. Rev. Lett. **102**, 162001 (2009), doi:[10.1103/PhysRevLett.102.162001](https://doi.org/10.1103/PhysRevLett.102.162001).
- [51] T. Becher and M. Neubert, *Infrared singularities of scattering amplitudes and N^3LL resummation for n -jet processes*, J. High Energy Phys. **01**, 025 (2020), doi:[10.1007/JHEP01\(2020\)025](https://doi.org/10.1007/JHEP01(2020)025).
- [52] S. Moch, J. A. M. Vermaseren and A. Vogt, *Three-loop results for quark and gluon form factors*, Phys. Lett. B **625**, 245 (2005), doi:[10.1016/j.physletb.2005.08.067](https://doi.org/10.1016/j.physletb.2005.08.067).
- [53] L. N. Lipatov, *The Bare Pomeron in Quantum Chromodynamics*, Sov. Phys. JETP **63**, 904 (1986).
- [54] F. C.S. Brown, *Polylogarithmes multiples uniformes en une variable*, Compt. Rend. Math. **338**, 527 (2004), doi:[10.1016/j.crma.2004.02.001](https://doi.org/10.1016/j.crma.2004.02.001).
- [55] F. Brown, *Single-valued Motivic Periods and Multiple Zeta Values*, SIGMA **2**, e25 (2014), doi:[10.1017/fms.2014.18](https://doi.org/10.1017/fms.2014.18).
- [56] O. Schnetz, *Graphical functions and single-valued multiple polylogarithms*, Commun. Num. Theor. Phys. **8**, 589 (2014), doi:[10.4310/CNTP2014.v8.n4.a1](https://doi.org/10.4310/CNTP2014.v8.n4.a1).
- [57] J. Pennington, *The six-point remainder function to all loop orders in the multi-Regge limit*, J. High Energy Phys. **01**, 059 (2013), doi:[10.1007/JHEP01\(2013\)059](https://doi.org/10.1007/JHEP01(2013)059).
- [58] V. Del Duca, L. J. Dixon, C. Duhr and J. Pennington, *The BFKL equation, Mueller-Navelet jets and single-valued harmonic polylogarithms*, J. High Energy Phys. **02**, 086 (2014), doi:[10.1007/JHEP02\(2014\)086](https://doi.org/10.1007/JHEP02(2014)086).
- [59] S. Catani, *The singular behaviour of QCD amplitudes at two-loop order*, Phys. Lett. B **427**, 161 (1998), doi:[10.1016/S0370-2693\(98\)00332-3](https://doi.org/10.1016/S0370-2693(98)00332-3).
- [60] G. Sterman and M. E. Tejeda-Yeomans, *Multi-loop amplitudes and resummation*, Phys. Lett. B **552**, 48 (2003), doi:[10.1016/S0370-2693\(02\)03100-3](https://doi.org/10.1016/S0370-2693(02)03100-3).
- [61] S. Aybat, L. J. Dixon and G. Sterman, *Two-Loop Anomalous-Dimension Matrix for Soft-Gluon Exchange*, Phys. Rev. Lett. **97**, 072001 (2006), doi:[10.1103/PhysRevLett.97.072001](https://doi.org/10.1103/PhysRevLett.97.072001).
- [62] S. Aybat, L. J. Dixon and G. Sterman, *Two-loop soft anomalous dimension matrix and resummation at next-to-next-to-leading poles*, Phys. Rev. D **74**, 074004 (2006), doi:[10.1103/PhysRevD.74.074004](https://doi.org/10.1103/PhysRevD.74.074004).
- [63] Ø. Almelid, C. Duhr and E. Gardi, *Three-Loop Corrections to the Soft Anomalous Dimension in Multileg Scattering*, Phys. Rev. Lett. **117**, 172002 (2016), doi:[10.1103/PhysRevLett.117.172002](https://doi.org/10.1103/PhysRevLett.117.172002).
- [64] C. Anastasiou, L. Dixon, Z. Bern and D. A. Kosower, *Planar Amplitudes in Maximally Supersymmetric Yang-Mills Theory*, Phys. Rev. Lett. **91**, 251602 (2003), doi:[10.1103/PhysRevLett.91.251602](https://doi.org/10.1103/PhysRevLett.91.251602).
- [65] Z. Bern, L. J. Dixon and V. A. Smirnov, *Iteration of planar amplitudes in maximally supersymmetric Yang-Mills theory at three loops and beyond*, Phys. Rev. D **72**, 085001 (2005), doi:[10.1103/PhysRevD.72.085001](https://doi.org/10.1103/PhysRevD.72.085001).

- [66] J. M. Drummond, G. P. Korchemsky and E. Sokatchev, *Conformal properties of four-gluon planar amplitudes and Wilson loops*, Nucl. Phys. B **795**, 385 (2008), doi:[10.1016/j.nuclphysb.2007.11.041](https://doi.org/10.1016/j.nuclphysb.2007.11.041).
- [67] I. I. Balitsky and A. V. Belitsky, *Nonlinear evolution in high-density QCD*, Nucl. Phys. B **629**, 290 (2002), doi:[10.1016/S0550-3213\(02\)00149-9](https://doi.org/10.1016/S0550-3213(02)00149-9).
- [68] J. M. Henn, G. P. Korchemsky and B. Mistlberger, *The full four-loop cusp anomalous dimension in $\mathcal{N} = 4$ super Yang-Mills and QCD*, J. High Energy Phys. **04**, 018 (2020), doi:[10.1007/JHEP04\(2020\)018](https://doi.org/10.1007/JHEP04(2020)018).
- [69] A. von Manteuffel, E. Panzer and R. M. Schabinger, *Cusp and Collinear Anomalous Dimensions in Four-Loop QCD from Form Factors*, Phys. Rev. Lett. **124**, 162001 (2020), doi:[10.1103/PhysRevLett.124.162001](https://doi.org/10.1103/PhysRevLett.124.162001).