

Renormalization & improvement of the tensor operator for $N_f = 3$ QCD in a χ SF setup

ALPHA
Collaboration

Isabel Campos Plasencia,^a Mattia Dalla Brida,^b Giulia Maria de Divitiis,^{c,d,*}
Andrew Lytle,^e Mauro Papinutto,^f Ludovica Pirelli^{c,d} and Anastassios Vladikas^d

^a*Instituto de Física de Cantabria IFCA-CSIC,
Avda. de los Castros s/n, E-39005 Santander, Spain*

^b*Theoretical Physics Department, CERN,
CH-1211 Geneva 23, Switzerland*

^c*Dipartimento di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy*

^d*INFN, Sezione di Tor Vergata, c/o Dipartimento di Fisica, Università di Roma “Tor Vergata”,
Via della Ricerca Scientifica 1, I-00133 Rome, Italy*

^e*Department of Physics, University of Illinois at Urbana-Champaign,
Urbana, IL 61801, USA*

^f*Dipartimento di Fisica, Università di Roma La Sapienza and INFN, Sezione di Roma,
Piazzale A. Moro 2, Roma, I-00185, Italy*

*E-mail: isabel.campos@csic.es, mattia.dalla.brida@cern.ch,
giulia.dedivitiis@roma2.infn.it, atlytle@illinois.edu,
mauro.papinutto@roma1.infn.it, ludovica.pirelli@students.uniroma2.eu,
tassos.vladikas@roma2.infn.it*

We present preliminary results of the non-perturbative renormalization group (RG) running of the flavor non-singlet tensor operator. We employ the χ SF scheme for $N_f = 3$ QCD using ensembles generated by the ALPHA collaboration for the computation of the quark mass running. The χ SF property of automatic $O(a)$ improvement prevents the $O(a)$ mixing of the correlation functions.

*The 38th International Symposium on Lattice Field Theory, LATTICE2021 26th-30th July, 2021
Zoom/Gather@Massachusetts Institute of Technology*

*Speaker

1. Flavor non-singlet tensor operator

A non-perturbative determination of renormalisation group running between hadronic and electroweak scales for the flavor non-singlet tensor operator

$$T_{\mu\nu}^a(x) = i\bar{\psi}(x) \sigma_{\mu\nu} \frac{1}{2}\tau^a \psi(x) \quad (1)$$

$$\sigma_{\mu\nu} \equiv \frac{i}{2}[\gamma_\mu, \gamma_\nu] \quad (2)$$

is very interesting from both phenomenological and theoretical points of view. The tensor enters the amplitudes of effective Hamiltonians, which describe, for example, rare heavy meson decays, neutron beta decays and possible Beyond Standard Model effects:

$$\mathcal{A} = \langle f | \mathcal{H}_{eff} | i \rangle = C_W(\mu) \langle f | \mathcal{O}(\mu) | i \rangle \quad (3)$$

$$\mathcal{O} \sim (\bar{l}\sigma_{\mu\nu}e)(\bar{q}_i\sigma_{\mu\nu}q_j), G_{\mu\nu}(\bar{q}_i\sigma_{\mu\nu}q_j) \dots \quad (4)$$

Moreover, the computation of the scale dependence of the renormalization factor completes the ALPHA renormalization and improvement programme of the bilinear operators. For $N_f = 0, 2$ such a study has appeared in ref. [1]. For $N_f = 3$, preliminary results of the RG-running in the relatively high energy range $2 \text{ GeV} \lesssim \mu \lesssim 128 \text{ GeV}$ have been reported in ref. [2]. $N_f = 3$ renormalisation factors at different scales are also presented in ref. [3].

2. RG flow

We employ a χ SF setup (see [4–8]), which is a mass-independent renormalization scheme. Such schemes are characterized by RG equations of the following form:

$$\mu \frac{\partial}{\partial \mu} T_R(\mu) = \gamma(g_R(\mu)) T_R(\mu), \quad T_R(\mu) = Z_T(\mu) T, \quad (5)$$

where g_R is the running coupling. The anomalous dimension γ has the perturbative expansion

$$\gamma(g_R) \stackrel{g_R \rightarrow 0}{\sim} -g_R^2(\gamma_0 + \gamma_1 g_R^2 + \gamma_2 g_R^4 + \mathcal{O}(g_R^6)), \quad (6)$$

with a universal coefficient γ_0 . The solution $T_R(\mu)$ is expressed in terms of an integration constant T_{RGI} , which is renormalization group invariant (RGI):

$$T_{\text{RGI}} = T_R(\mu) \left[\frac{g_R^2(\mu)}{4\pi} \right]^{-\frac{\gamma_0}{2b_0}} \exp \left\{ - \int_0^{g_R(\mu)} dg \left[\frac{\gamma(g)}{\beta(g)} - \frac{\gamma_0}{b_0 g} \right] \right\}. \quad (7)$$

It is possible to factorize the running in many evolutions between two scales:

$$T_R(\mu) = \frac{T_R(\mu)}{T_R(\mu_n)} \dots \frac{T_R(\mu_2)}{T_R(\mu_1)} \frac{T_R(\mu_1)}{T_{\text{RGI}}} T_{\text{RGI}}, \quad (8)$$

leading naturally to the definition of the step scaling function:

$$\sigma_T(s, u) = \frac{T_R(\mu_2)}{T_R(\mu_1)} = \frac{Z_T(\mu_2)}{Z_T(\mu_1)}, \quad (9)$$

where $s \equiv \frac{\mu_1}{\mu_2}$ and $u \equiv g_R^2(\mu_1)$. A common and convenient choice is to take successive scales at fixed ratio $s = 2$:

$$\sigma_T(u) \equiv \sigma_T(2, u) = \exp \left\{ \int_{g_R(\mu)}^{g_R(\mu/2)} dg \frac{\gamma(g)}{\beta(g)} \right\}. \quad (10)$$

On the lattice, the scale evolution can be studied non-perturbatively as a finite size scaling, with the renormalization scale identified as the inverse of the lattice size $\mu = \frac{1}{L}$:

$$\mu = \frac{1}{L}, \quad u \equiv g_R^2(L) \quad (11)$$

$$\sigma(u) = \lim_{a \rightarrow 0} \Sigma(u, a/L) \quad \Sigma(u, a/L) = g_R^2(2L) \quad (12)$$

$$\sigma_T(u) = \lim_{a \rightarrow 0} \Sigma_T(u, a/L) \quad \Sigma_T(u, a/L) = \frac{Z_T(g_0^2, a/2L)}{Z_T(g_0^2, a/L)}, \quad (13)$$

where a is the lattice spacing. The renormalization constants $Z_T(g_0^2, a/L)$ are defined imposing renormalization conditions on the correlation functions, as shown in eqs. (22,23) of the next section.

Our actual RG flow materializes in a sequence of many lattices. We used the same gauge configurations generated by the ALPHA collaboration for the determination of the quark mass running (see [9] for details of the simulations). They refer to $N_f = 3$ massless Wilson-clover fermions with Schrödinger Functional (SF) boundary conditions. The simulation parameters correspond to a RG evolution from an hadronic scale μ_{had} of about 200 MeV to a perturbative scale μ_{pt} around 128 GeV. The peculiarity of this flow is the change of schemes at the intermediate scale $\mu_0/2 \sim 2$ GeV: in the high energy region the running coupling is defined in the SF scheme ($g_R = g_{SF}$) [10–12], while in the low energy region it is defined in the gradient flow (GF) scheme ($g_R = g_{GF}$) [13, 14]:

$$\begin{array}{ccccccc} & \mu_{had} & \text{GF scheme} & \mu_0/2 & \text{SF scheme} & \mu_{pt} & \\ & \approx 200\text{MeV} & & \approx 2\text{GeV} & & \approx 128\text{GeV} & \\ & \leftarrow & & \leftarrow & & \leftarrow & \mu \end{array}$$

We impose the same definition of $Z_T(g_0^2, a/L)$ at all scales, which implies that the anomalous dimension, which is a different function of g_{SF}^2 and of g_{GF}^2 , has the same value at a given renormalisation scale μ :

$$\gamma(\mu) = \gamma_{SF}(g_{SF}^2(\mu)) = \gamma_{GF}(g_{GF}^2(\mu)). \quad (14)$$

3. χ SF Chirally Rotated Schrödinger Functional

In our study we adopt a mixed action approach (see also [15, 16]): while the sea quarks obey the standard SF boundary conditions, for the valence quarks we impose χ SF boundary conditions. In the continuum and chiral limit, the SF and χ SF setups are equivalent, being connected by a chiral flavor transformation [4]:

$$R = \exp \left(i \frac{\alpha}{2} \gamma_5 \tau^3 \right) \Big|_{\alpha=\pi/2} \quad \begin{cases} \psi & \rightarrow \psi' = R\psi \\ \bar{\psi} & \rightarrow \bar{\psi}' = \bar{\psi}R \end{cases} \quad (15)$$

$$P_{\pm} \equiv \frac{1}{2}(1 \pm \gamma_0) \rightarrow Q_{\pm} \equiv \frac{1}{2}(1 \pm i\gamma_0\gamma_5\tau^3), \quad (16)$$

where P_{\pm} and Q_{\pm} are the SF and the χ SF projectors acting on fermionic fields at the boundaries.

At finite lattice spacing however, χ SF breaks the parity-flavor symmetry $\mathcal{P}_5 = i\gamma_0\gamma_5\tau^3$, which is recovered by introducing an extra boundary counterterm of dimension 3 with coefficient z_f . The parameters z_f and the bare mass m_0 must be tuned non-perturbatively to their critical values in order to restore parity-flavor and chiral symmetries up to discretisation effects. In practice the two tunings can be done independently, so we inherited the value of the critical hopping parameter κ from the SF simulations [9], while we fixed z_f imposing the vanishing of g_A^{ud} , a \mathcal{P}_5 -odd correlation function:

$$\begin{cases} m = \frac{\tilde{\delta}_0 f_{A,I}^{ud}(x_0)}{2f_P^{ud}(x_0)} \Big|_{x_0=L/2} = 0 & m_{cr} \text{ tuning} \\ g_A^{ud}(x_0) \Big|_{x_0=L/2} = 0 & z_f \text{ tuning.} \end{cases} \quad (17)$$

Here m stands for the SF-PCAC quark mass, $f_{A,I}$ and f_P are the usual SF correlation functions of the improved axial current and the pseudoscalar density, while g_A^{ud} is the χ SF correlation function involving the axial current with flavors u, d . See Table 1 for a brief overview of the correlation functions. Some details of our tuning procedure can be found in ref. [16]

A boundary improvement counterterm proportional to the coefficients d_s is also needed in order to cancel $O(a)$ discretisation effects originating at the time borders.

Once these requirements are fulfilled, the argument of automatic $O(a)$ improvement is achieved in χ SF [4, 5]: the \mathcal{P}_5 -even correlation functions receive corrections only at second order in the lattice spacing, whereas the \mathcal{P}_5 -odd ones are pure lattice artefacts:

$$g_{even} = g_{even}^{\text{continuum}} + O(a^2) \quad (18)$$

$$g_{odd} = O(a). \quad (19)$$

This property turns out to be particularly advantageous for the tensor operator, because $O(a)$ improvement does not require mixing with bulk counterterms in the correlation functions. For l_T with flavor combination ud , for example, the improvement coefficient c_T is irrelevant, since the vector correlation function l_V , being \mathcal{P}_5 -odd, is $O(a)$. Therefore the Symanzik correction, being $O(a^2)$, may be dropped:

$$T_{\mu\nu}^I = T_{\mu\nu} + c_T(g_0^2) a (\tilde{\delta}_\mu V_\nu - \tilde{\delta}_\nu V_\mu), \quad (20)$$

$$l_T^{ud,I} = l_T^{ud} + c_T(g_0^2) a \tilde{\delta}_0 l_V^{ud}. \quad (21)$$

The rich variety of correlation functions is an interesting feature of χ SF, offering the possibility of several definitions of Z_T , for example through the renormalization condition on the electric tensors as well as on the magnetic ones:

$$Z_T(g_0, a/L) \frac{l_T^{ud}(L/2)_{\text{re}}}{\sqrt{l_1^{ud}}} = \frac{l_T^{ud}(L/2)_{\text{re}}}{\sqrt{l_1^{ud}}} \Big|_{\text{Tree Level}} \quad \text{“electric” tensor } T_{0k} \quad (22)$$

$$Z_T(g_0, a/L) \frac{l_T^{uu'}(L/2)_{\text{im}}}{\sqrt{l_1^{uu'}}} = \frac{l_T^{uu'}(L/2)_{\text{im}}}{\sqrt{l_1^{uu'}}} \Big|_{\text{Tree Level}} \quad \text{“magnetic” tensor } \tilde{T}_{0k} = -\frac{1}{2}\varepsilon_{0kij}T_{ij}. \quad (23)$$

flavors	$f_1, f_2 = u, d, u', d'$	
bulk operators	$X = V_0, A_0, S, P$	$Y_k = V_k, A_k, T_{k0}, \bar{T}_{k0}$
SF	χSF	
$f_X(x_0) = -\frac{1}{2} \langle X^{f_1 f_2}(x) O_5^{f_2 f_1} \rangle$	$g_X(x_0) = -\frac{1}{2} \langle X^{f_1 f_2}(x) Q_5^{f_2 f_1} \rangle$	
$k_Y(x_0) = -\frac{1}{6} \sum_{k=1}^3 \langle Y_k^{f_1 f_2}(x) O_k^{f_2 f_1} \rangle$	$l_Y(x_0) = -\frac{1}{6} \sum_{k=1}^3 \langle Y_k^{f_1 f_2}(x) Q_k^{f_2 f_1} \rangle$	
$f_1 = -\frac{1}{2} \langle O_5^{f_1 f_2} O_5'^{f_2 f_1} \rangle$	$g_1 = -\frac{1}{2} \langle Q_5^{f_1 f_2} Q_5'^{f_2 f_1} \rangle$	
$k_1 = -\frac{1}{6} \sum_{k=1}^3 \langle O_k^{f_1 f_2} O_k'^{f_2 f_1} \rangle$	$l_1 = -\frac{1}{6} \sum_{k=1}^3 \langle Q_k^{f_1 f_2} Q_k'^{f_2 f_1} \rangle$	
$O_5^{f_1 f_2} = a^6 \sum_{y,z} \bar{\zeta}_{f_1}(y) P_+ \gamma_5 \zeta_{f_2}(z)$	$Q_5^{uu'} = a^6 \sum_{y,z} \bar{\zeta}_u(y) \gamma_0 \gamma_5 Q_- \zeta_{u'}(z)$	
$O_k^{f_1 f_2} = a^6 \sum_{y,z} \bar{\zeta}_{f_1}(y) P_+ \gamma_k \zeta_{f_2}(z)$	$Q_k^{uu'} = a^6 \sum_{y,z} \bar{\zeta}_u(y) \gamma_k Q_- \zeta_{u'}(z)$	
\vdots	\vdots	
$P_{\pm} \equiv \frac{1}{2}(1 \pm \gamma_0)$	$Q_{\pm} \equiv \frac{1}{2}(1 \pm i\gamma_0 \gamma_5)$	

Table 1: Brief overview of correlation functions in SF and χ SF setups.

The subscripts $_{\text{re}}$, $_{\text{im}}$ here denote the real and the imaginary parts. Both relations correspond to the same SF renormalization condition in the continuum. In fact, the continuum equivalence of the SF and χ SF setups implies universality relations among correlation functions:

$$k_T = i l_{\bar{T}}^{uu'} = -i l_{\bar{T}}^{dd'} = l_T^{ud} = l_T^{du}, \quad (24)$$

$$k_1 = l_1^{uu'} = l_1^{dd'} = l_1^{ud} = l_1^{du}. \quad (25)$$

4. Results

Our preliminary results are based on the determination of Z_T from eq. (22). We obtain the continuum limit of the tensor lattice step scaling function $\Sigma_T(u, a/L)$ performing global fits of the data at different couplings and lattice spacings:

$$\Sigma_T(u, a/L) = \frac{Z_T(g_0^2, a/2L)}{Z_T(g_0^2, a/L)} \quad (26)$$

$$= \sigma_T(u) + \rho_T(u) \left(\frac{a}{L}\right)^2, \quad (27)$$

with $\sigma_T(u)$ and $\rho_T(u)$ parameterised by polynomials. Fig. 1 shows Σ_T as a function of $(a/L)^2$, and parametrized by the coupling u (in different colours) for the high energy region. The relative continuum limit is illustrated in the top panel of Fig. 2. Our data (red circles) are in agreement with perturbation theory at two loops (gray line) [1] and with the data obtained in a purely SF setup (circles in black) [2, 17]. This demonstrates the continuum SF- χ SF universality. The results at

low energies are shown in the bottom panel of the same figure. We have extracted the anomalous dimension γ , relying on the formula:

$$\sigma_T(u) = \exp \left\{ \int_{\sqrt{u}}^{\sqrt{\sigma(u)}} dg \frac{\gamma(g)}{\beta(g)} \right\}. \quad (28)$$

In Fig. 3 we then finally show our preliminary results for γ_{SF} and γ_{GF} , over the full range of couplings available.




Figure 1: The step scaling function $\Sigma_T(u, a/L)$ in the high energy region

5. Conclusions

We have presented preliminary results for the RG running of flavor non-singlet tensor operator in $N_f = 3$ QCD, using the gauge configurations generated by the ALPHA collaboration [9]. The data span, in a fully non-perturbative way, a range of energies of about three orders of magnitude, going from hadronic to electro-weak scales. We obtained the anomalous dimension of the tensor operator, aiming to complete the computation of the non-perturbative RG running of all dimension 3 bilinear operators.




Figure 2: The continuum step scaling function $\sigma_T(u)$ at high energies (top panel) and at low energies (bottom panel). The two energy regions correspond to different definitions of the running coupling: $u = g_{SF}^2$ [10–12] and $u = g_{GF}^2$ [13, 14].

6. Acknowledgements

We wish to thank Patrick Fritzsche, Carlos Pena, David Preti, and Alberto Ramos for their help. This work is partially supported by INFN and CINECA, as part of research project of the QCDLAT INFN-initiative. We acknowledge the Santander Supercomputacion support group at the University of Cantabria which provided access to the Altamira Supercomputer at the Institute of Physics of Cantabria (IFCA-CSIC). We also acknowledge support by the Poznan Supercomputing and Networking Center (PSNC) under the project with grant number 466. AL acknowledges support by the U.S. Department of Energy under grant number DE-SC0015655.




Figure 3: The anomalous dimensions $\gamma_{SF}(u)$ and $\gamma_{GF}(u)$. The two schemes SF and GF correspond to different definitions of the running coupling: $u = g_{SF}^2$ [10–12] and $u = g_{GF}^2$ [13, 14].

References

- [1] C. Pena and D. Preti, *Eur. Phys. J. C* **78** (2018) no.7, 575 doi:10.1140/epjc/s10052-018-6022-7 [arXiv:1706.06674 [hep-lat]].
- [2] L. Chimirri *et al.* [ALPHA], *PoS LATTICE2019* (2020), 212 doi:10.22323/1.363.0212 [arXiv:1910.06759 [hep-lat]].
- [3] G. S. Bali, S. Bürger, S. Collins, M. Göckeler, M. Gruber, S. Piemonte, A. Schäfer, A. Sternbeck and P. Wein, *Phys. Rev. D* **103** (2021) no.9, 094511 doi:10.1103/PhysRevD.103.094511 [arXiv:2012.06284 [hep-lat]].
- [4] S. Sint, *Nucl. Phys. B* **847** (2011), 491-531 doi:10.1016/j.nuclphysb.2011.02.002 [arXiv:1008.4857 [hep-lat]].
- [5] S. Sint and B. Leder, *PoS LATTICE2010* (2010), 265 doi:10.22323/1.105.0265 [arXiv:1012.2500 [hep-lat]].
- [6] M. Dalla Brida, S. Sint and P. Vilaseca, *JHEP* **08** (2016), 102 doi:10.1007/JHEP08(2016)102 [arXiv:1603.00046 [hep-lat]].
- [7] P. V. Mainar, M. Dalla Brida and M. Papinutto, *PoS LATTICE2015* (2016), 252 doi:10.22323/1.251.0252
- [8] M. Dalla Brida, T. Korzec, S. Sint and P. Vilaseca, *Eur. Phys. J. C* **79** (2019) no.1, 23 doi:10.1140/epjc/s10052-018-6514-5 [arXiv:1808.09236 [hep-lat]].

- [9] I. Campos *et al.* [ALPHA], Eur. Phys. J. C **78** (2018) no.5, 387 doi:10.1140/epjc/s10052-018-5870-5 [arXiv:1802.05243 [hep-lat]].
- [10] M. Lüscher, R. Narayanan, P. Weisz and U. Wolff, Nucl. Phys. B **384** (1992), 168-228 doi:10.1016/0550-3213(92)90466-O [arXiv:hep-lat/9207009 [hep-lat]].
- [11] S. Sint, Nucl. Phys. B **421** (1994), 135-158 doi:10.1016/0550-3213(94)90228-3 [arXiv:hep-lat/9312079 [hep-lat]].
- [12] M. Dalla Brida *et al.* [ALPHA], Phys. Rev. Lett. **117** (2016) no.18, 182001 doi:10.1103/PhysRevLett.117.182001 [arXiv:1604.06193 [hep-ph]].
- [13] M. Dalla Brida *et al.* [ALPHA], Phys. Rev. D **95** (2017) no.1, 014507 doi:10.1103/PhysRevD.95.014507 [arXiv:1607.06423 [hep-lat]].
- [14] P. Fritzsche and A. Ramos, JHEP **10** (2013), 008 doi:10.1007/JHEP10(2013)008 [arXiv:1301.4388 [hep-lat]].
- [15] I. Campos, M. Dalla Brida, G. M. de Divitiis, A. Lytle, M. Papinutto and A. Vladikas, PoS **LATTICE2019** (2019), 202 doi:10.22323/1.363.0202 [arXiv:1910.01898 [hep-lat]].
- [16] I. Campos Plasencia, M. Dalla Brida, G. M. de Divitiis, A. Lytle, M. Papinutto, L. Pirelli and A. Vladikas, “Quark mass RG-running for $N_f = 3$ QCD in a χSF setup,” [arXiv:2111.15384 [hep-lat]].
- [17] Fabian Joswig, “Non-perturbative renormalization of $O(a)$ improved tensor currents,” [talk slides] **LATTICE2019** (2019)

https://indico.cern.ch/event/764552/contributions/3420570/attachments/1863864/3064496/fabian_joswig_LATTICE19.pdf