

# MINIMALISTIC MUSINGS ABOUT THE STANDARD MODEL\*

CHRIS P. KORTHALS ALTES

NIKHEF Theory Group, NIKHEF Amsterdam  
Science Park 105, 1098 XG Amsterdam, The Netherlands  
and  
Centre de Physique Théorique, Campus de Luminy  
163 Avenue de Luminy, 13009 Marseille, France

*(Received May 14, 2021; accepted May 17, 2021)*

Martinus (“Tini”) Veltman’s early contributions to the Standard Model were essential for its success. After some nostalgic reminiscences, I turn to the Standard Model with a minimalistic attitude, the point of view that beyond the SM there is only the Planck scale. Known since long, the gravitational force can be obtained as the gauge theory of local Poincaré symmetry, called gauge gravity. This gauge theory of gravity embodies *per se* a Palatini formulation. This causes the potential of non-minimally coupled Higgs inflation to have an intriguing improved large field behaviour. Some of its effects are experimentally accessible or refutable. The question of quantum corrections is discussed.

DOI:10.5506/APhysPolB.52.607

## 1. Introduction

In writing a tribute to Prof. Tini Veltman, it occurred to me that he was, like many, a physicist prey to periods of elation ranging to bouts of despair about perceived lack of understanding of his work. He had certainly some reasons to have despaired, when in the late sixties he was analysing gauge theories for weak interactions, which were, so he was told by colleagues, “one of those funny corners of physics”.

But in Tini’s case, elation had always the last word, amongst more as a recipient at the 1999 Nobel banquet in Stockholm’s town hall. His elation did flow over to the audience in his after-dinner speech; he likened the anguish of potential Nobel candidates all over the world waiting for the call from Stockholm, to the terror spread by the Viking ancestors of the Nobel committee a millennium ago amongst innocents on European shores.

---

\* Funded by SCOAP<sup>3</sup> under Creative Commons License, CC-BY 4.0.

His Burgundian after-dinner speeches were matched by caustic diatribes against anything he considered pretentious.

Indeed, it is with due circumspection that I write these lines. Tini would have despaired about the title and abstract.

Actually, he had, as far as I know, four famous encounters with gravity that went into print. The first one was in a paper with Henk van Dam [1] on the lack of a continuous limit from massive vector gauge theory and Einstein gravity theories to their massless version, and concluded that the latter should be strictly massless. The second one on the one-loop gravity action [2], with Gerard 't Hooft. The third one was a Letter on the effect of the Higgs condensate as a source of the gravitational field brought up by Linde [3] a few months before. The fourth one in his lecture course on gravity at the Les Houches School [4].

The first and second papers are quite influential till the present day.

His lectures are typical for the particle physicist he was, in heart and soul. He did forego the geometrical approach, and presented gravity as the theory of gravitons. The lectures are reflecting the point of view taken in the CERN report DIAGRAMMAR [5]. That is, for him the quantum theory of gravity was the collection of diagrams. This mantra had served him well in the elucidation of radiative corrections to weak and electromagnetic interactions.

Curiously, he did not appreciate the usefulness of solitons or instantons in theories like QCD. Not only in QCD. I was present when he explained to a well-known cosmologist in the CERN coffee room why black holes cannot possibly form — although he may well have played the devil's advocate, something he loved to do in order to challenge people. Not only solitons but also certain disciplines in physics did not carry his approval. At lunch with a group of condensed matter physicists in a restaurant in the Quartier Latin, during a meeting at the Ecole Normale, he aired the opinion that Statistical Mechanics was the art of finding an interesting physics property and then averaging it out.

I had a brush with him in the corridor of CERN just after the second gravity paper appeared, because I asked him why gravity had anything to do with particle physics. Why was gravity quantized? After all, gravitons had not been seen. After starting of course a muscular response, to my astonishment he acquiesced. Nowadays, astrophysical black holes have been detected, but the discussion how to detect effects from the quantum nature of gravity [8] and detect gravitons [9] is very interesting, but still at an academic stage. Tini knew very well that quantum gravity, with his approach or any other, was a formidable opponent and he did not engage it anymore after the mid-seventies, at least not in published form.

Tini entered my horizon when he gave a seminar in the mid-sixties in Amsterdam where I studied theory. He wore sandals, a pullover and displayed a very caustic sense of humour. This in stark contrast to the professors in Amsterdam who wore — at that time at least — a conventional outfit.

A little later, involved myself in current algebra I profited from his PRL letter on the divergence equations [6]. It explained wonderfully well how sum rules of current algebra could be obtained from low energy theorems and dispersion relations, avoiding any discussion of Schwinger terms. This paper signalled the start of his conviction that non-Abelian vector fields were the key to understanding weak interactions.

Tini was a regular guest at the CPT in Marseille, from 1968 on, partly because the man in charge there, Toni Visconti, and his students were early birds in computer generation of Feynman graphs and wanted to profit from Tini's *SCHOONSCHIP* expertise. I had arrived there as a postdoc after the 1971 revolution at the Amsterdam HEP conference, where Tini had Gerard 't Hooft presented part of his thesis work [10].

At the 1971 Orsay Summer Institute Tini told me that he worried about the chiral anomalies in what was then still the purely leptonic version of the SM. Michel Perrottet and I wrote a paper belabouring this worry [12]. In 1972, both Tini and Gerard played a crucial role at the Marseille conference on Yang–Mills fields (at that time the Standard Model including quarks was two months old [11]), and on a similar occasion in 1974.

Of course, the famous Summer Institutes at the Orsay, Ecole Normale and Triangular Meetings were hotbeds for developing Particle Theory, and Tini and Gerard were faithful participants, with crucial contributions.

In 1974, Tini had invited me to spend some months in Utrecht, which happened to coincide with the advent of the 1974 October revolution. He discussed his paper on the Linde effect extensively before he sent it off. However, the ongoing revolution posed more pressing subjects . . .

From autumn 1978 till spring 1981, I spent more than two years in Utrecht. I had decided to work on numerical lattice gauge theory although Tini certainly did not encourage me to do so. He was busy with his work on radiative corrections in the electroweak sector and gave his well-known talk (“Quarks and Leptons: what next?”) at the 1979 Lepton Photon Conference at Fermi Lab. It makes interesting reading still now. Bottom was then just discovered. Tini discussed the expected top–bottom mass difference in connection with custodial symmetry and radiative corrections to the  $\rho$  parameter. He was playing with a Higgs heavier than the unitarity limit of vector boson scattering, creating a strongly interacting sector. Nevertheless, his talk ends with the admonition “We should keep checking electron muon universality”.

Well, his advice was heeded, according to the latest news from the LHCb [7].

Some years later, he spent long periods at the Theory Department of the Universita Autonoma in Madrid, working with Francisco (“Paco”) Indurain. I happened to have an ongoing collaboration at that very same place and time with Antonio Gonzalez Arroyo. Apart from enjoying a marvellous hospitality, I had ample time to observe the flamboyant encounters that Tini had with the quite picturesque population of the Residencia de Estudiantes, a revered Madrileño landmark where we had the honour to stay. It had traditionally harboured many intellectuals and scientists from in and outside Spain.

After these recollections, I would like to come back to the minimalistic view of the Standard Model, the fact that gravity is just such as the electroweak and strong force a gauge force, with as gauge group the local Poincaré group, and how it may embody inflation.

Of course, this idea of “gauge gravity” is an old idea, starting with a series of papers precisely sixty years ago [13].

Kibble’s conclusion<sup>1</sup> was that the theory was for all practical purposes indistinguishable from Einstein’s theory. More precisely, in absence of matter, they are identical, but in the presence of fermionic matter, they are differing by a cosmological term

$$\sim \sqrt{-g}M_{\text{P}}^{-4} \left( \sum_f \bar{\psi}(x)\gamma^a\gamma^b\gamma^c\psi(x) \right)^2. \quad (1)$$

$M_{\text{P}} = \sqrt{(8\pi G_{\text{N}})^{-1}}$  is the reduced Planck mass, and the indices  $a, b, c$  on the Dirac matrices are fully anti-symmetric. The sum is over all fermions in the SM, provided the right-handed neutrinos are the Dirac type. The square of Newtons constant renders all effects of this term unobservably small.

However, if one introduces a non-minimal coupling of the Standard Model Higgs to the curvature scalar  $R$

$$-\xi H^\dagger H R, \quad (2)$$

then the behaviour of the Higgs potential for large field values is crucially changed, with measurable consequences in inflation. This idea, very much in the spirit of minimalism, is due to Bezrukov and Shaposhnikov [22] in the context of Einstein gravity. Subsequently, Bauer and Demir [24] realized that the Palatini formulation [19] of Einstein gravity improved parametrically (in terms of  $\xi$ ) this inflation potential for high values of the Higgs field. Recall that the Palatini method starts from positing the metric and the connection to be independent variables in the Einstein Hilbert action. Moreover, gauge

---

<sup>1</sup> For an identical view, see also Weyl [14].

gravity has a Palatini method built-in, essentially because the Lorentz group and the translations generate independent connections, as will be amply clear from Section 2.

Controversy sets in when quantum effects are taken into account. The traditional approach is to look for symmetries that are respected on the quantum level. That tells you the general structure of the effective action. The crucial symmetry is the approximate translation invariance of the flat Higgs potential.

I was ruminating on this idea in late summer last year, thinking that gauge gravity was a fairly remote and quiet corner. Till I discussed with Jan Smit who soon brought to my attention a couple of quite recent papers by Shaposhnikov and coworkers [23]. So what follows is mostly history and I do apologise for incomplete referencing.

How gauge gravity emerges from local Poincaré invariance is narrated in Section 2. I could not withstand the temptation to do it in the form of a pastiche using two well known personalities. Both came into existence practically on the CERN site, in Ferney–Voltaire two and a half centuries ago<sup>2</sup>. They are the legendary Candide and his companion, Dr. Pangloss. They are supposed to have learnt how the forces in the Standard Model are obtained by the gauge principle but never learnt about any theoretical description of gravity, in particular, they are blissfully unaware of the use of geometrical ideas. However, they are supposed to be acquainted with the experiments that infer that all bodies fall equally fast *in vacuo*. This permits them to almost deduce that the force associated with the local Poincaré group is that of gravity. The pastiche serves a dual purpose, to realize how simply the SM begets gauge gravity and how much we are preconditioned by the geometric point of view.

In Section 3, I discuss Higgs inflation in the light of gauge gravity.

## 2. In which Candide contemplates gauge gravity potentials

As already mentioned in the introduction, the strict believer in the gauge principle of the SM is Dr. Pangloss. He is very intent on finding out what physical force is associated with the gauging of Poincaré symmetry. Unfortunately, he is prone to pedantry and condescendence. Candide has his reservations and is waiting to punch on weaknesses in the exposé of his colleague.

The two decide to write a Poincaré transformation as

$$x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu} + a^{\mu}, \quad \Lambda^{\mu}_{\nu} \eta^{\nu\rho} \Lambda_{\rho}^{\sigma} = \eta^{\mu\sigma}, \quad \eta = (1, -1, -1, -1). \quad (3)$$

---

<sup>2</sup> Due to Voltaire's fulminations [15] against Leibniz' idea of "All is for the best in the best of all possible worlds".

Dr. Pangloss' aim is to introduce a gauge covariant derivative for the Poincaré symmetry made local. This asks for gauge potentials. Candide concurs to take the potentials  $t_a^\mu$  for the translations with  $\partial_\mu$  as the translation charge. The index  $a = 0, 1, 2, 3$ . The local Lorentz symmetry needs a potential  $A_{ab\mu}$  with the normalized Lorentz group generators  $M^{ab}$  as charges. The latter will depend on whether one has a quark ( $M^{ab} = \gamma^{[a}\gamma^{b]}/2i$ ) or a vector boson ( $(M^{ab})_{cd} = \delta_c^{[a}\delta_d^{b]}$ ). They should appear in covariant derivatives.

They decide to find the covariant derivative for the Higgs field  $H$  first, as that should be the simplest case. If the transformation is global  $H'(x') = H(x)$  and

$$\partial'_\mu H'(x') = \Lambda_\mu^\nu \partial_\nu H(x)$$

they want a derivative  $D_a$ , such that in the case of  $x'^\mu = \Lambda_\nu^\mu(x)x^\nu + a^\mu(x)$ ,

$$D'_a H'(x') = \Lambda_a^b(x) D_b H(x). \quad (4)$$

First, they note that the local Poincaré transformation is a general coordinate transformation, and  $H'(x') = H(x)$ . Hence,

$$\partial'_\mu H'(x') = \frac{\partial x^\nu}{\partial x'^\mu} \partial_\nu H(x).$$

However, this is not yet what they want, see Eq. (4). But if they put  $D_a = t_a^\mu(x)\partial_\mu$ , then assuming for the translation potential the rule

$$t'^\mu_a(x') = \frac{\partial x^\nu}{\partial x'^\mu} \Lambda_a^b t_b^\nu(x), \quad (5)$$

their Eq. (4) results.

The Higgs kinetic term becomes  $\int d^4x t (D_a H)^\dagger D_b H \eta^{ab}$ ,  $t$  being the determinant of the inverse  $t_a^\mu$  of  $t_b^\nu(x)$ <sup>3</sup>.

Candide makes the point that the transformation law (5) does not have an inhomogeneous term, whereas the transformation law for the local Lorentz potential has one, by construction. He suggests another name to avoid confusion and he proposes the name Vierbein for  $t_\mu^a$ .

Dr. Pangloss agrees and proposes to carry on with the less simple fermion case with  $\psi'(x') = S(\Lambda)\psi(x)$ .

Were it not for the transformed coordinate  $x'$  on the left-hand side, they would be concerned with an internal symmetry operation as in the Standard Model with an  $SO(3, 1)$  potential  $A(x)_{ab\mu}$ , and a familiar covariant derivative

$$D_\mu(A) = \partial_\mu + iA_\mu, \quad A_\mu = \frac{1}{2} A_{ab\mu} M^{ab}. \quad (6)$$

<sup>3</sup> Plugging in the definition of  $D_a$  gives the minimal coupling of a tensor  $g^{\mu\nu} \equiv \eta^{ab} t_a^\mu(x) t_b^\nu(x)$  to the usual derivatives. From this, identification follows that the translation potential is nothing but the Vierbein. Remember, however, that our friends are unaware of metric, connections *etc.*

This derivative transforms under a local Lorentz transformation as they are used to in the Standard Model to obtain a covariant derivative. After a local Poincaré transformation  $x'(x)$  in  $\psi(x')$ , it behaves as a contravariant vector

$$D'_\mu(A') \equiv \partial'_\mu + iA'(x')_\mu = \frac{\partial x^\nu}{\partial x'^\mu} D_\nu(A). \tag{7}$$

So they obtain for the combination

$$D'_\mu(A') \psi'(x') = \frac{\partial x^\nu}{\partial x'^\mu} S(A) D_\nu(A) \psi(x).$$

Like for the scalar case, this is not yet their desired result. They multiply the gauge covariant derivative (6) with the Vierbein  $t_a^\mu$  to get for the local Poincaré covariant derivative  $D_a(A) = t_a^\mu \partial_\mu + it_a^\mu A_\mu$

$$D'_a(A') \psi'(x') = \Lambda_a^b(x) S(A) D_b(A) \psi(x).$$

Dr. Pangloss notes that  $D_a(A)$  is the sum of a translation term  $t_a^\mu \partial_\mu$  and a local Lorentz transform term  $t_a^\mu A_\mu$ . This is, as expected from a local Poincaré, invariant derivative. However, as brought up by Candide already in the scalar case, the translational “potential” has no inhomogeneous term.

The kinetic term for the fermion field becomes in terms of the new covariant derivative

$$L_f(\psi, D_a \psi) = \bar{\psi} i \gamma^a D_a(A) \psi.$$

However, Candide demurs. He repeats his observation made in the scalar case. Only the local Lorentz potential has an inhomogeneous part. He then goes on reproaching Dr. Pangloss that he has produced a nice mathematical framework to obtain a local Poincaré-invariant action for scalar and fermion fields but no physical interpretation. Where is the potential with an inhomogeneous term under a general coordinate transformation?

The two gentlemen agree to disagree and to reconvene quickly.

*2.1. In which Candide discovers what describes the gravitational force*

The discussion is taken up again with Dr. Pangloss announcing he has found an answer to the question of Candide. He claims to have found a combination of the translation and local Lorentz potentials that has the property Candide desires.

In what follows, Dr. Pangloss is explaining himself within the quotation marks.

“Like in the Standard Model, the commutators of our derivatives give the field strengths. In our case, I have two. One is the field strength formed from

$$[D_\mu(A) D_\nu(A)] \equiv \frac{1}{2} \hat{R}_{\mu\nu}(A). \tag{8}$$

This is the SO(3, 1) gauge field strength and can be written in component form  $\hat{R}_{\mu\nu} = \hat{R}^{bc}_{\mu\nu} M_{bc}$ . I can multiply this tensor with the Vierbeins  $t_b^\kappa t_c^\lambda$  and the result is the tensor  $\hat{R}^{\kappa\lambda}_{\mu\nu}$ .

On the other hand, I can form the commutator from the derivatives  $D_a(A) = t_a^\mu D_\mu(A)$ . The result is *different* from  $\hat{R}_{\mu\nu} t_a^\mu t_b^\nu \equiv \hat{R}_{ab}$

$$\begin{aligned} [D_a D_b] \psi &= \frac{1}{2} \hat{R}_{ab} \psi - T_{ab}^c D_c \psi, \\ T_{ab}^c &= - (t_a^\mu t_b^\nu - t_b^\mu t_a^\nu) D_\mu(A) t_\nu^c. \end{aligned} \quad (9)$$

The torsion  $T_{ab}^c$  is the new element, obviously antisymmetric in the last two indices<sup>4</sup>. Let me project it on Einstein indices  $T_{\sigma\tau}^\rho$  with the Vierbeins. It is then the antisymmetric part of a new quantity, a composite of Vierbeins and SO(3, 1) gauge potential

$$\Gamma_{\sigma\tau}^\rho(x) \equiv -t^b_\sigma D_\tau(A) t_b^\rho. \quad (10)$$

At this point, Candide gets impatient and asks what all this is good for. Dr. Pangloss implores him to let him finish.

He shows how  $\Gamma$  transforms under general coordinate transformations by using in (10) the transformation properties (7) and (5). He comes up with the following formula for the transformation law:

$$\Gamma'^\rho_{\sigma\tau}(x') = \frac{\partial x'^\rho}{\partial x^\lambda} \frac{\partial x^\mu}{\partial x'^\sigma} \frac{\partial x^\nu}{\partial x'^\tau} \Gamma^\lambda_{\mu\nu}(x) + \frac{\partial x'^\rho}{\partial x^\mu} \frac{\partial^2 x^\mu}{\partial x'^\sigma \partial x'^\tau}.$$

Candide is delighted by this expression: “Dr. Pangloss, this is a very interesting finding. Pray, let us try to interpret it.”

He starts by noting that this law permits the symmetric part of the new quantity  $\Gamma$  to be transformed away, at least locally, where the first derivatives can be set to unity. Dr. Pangloss concurs in this view. The latter then remembers the experiments of his colleagues Philoponus [16] and Galileo [17] that indicate all bodies fall equally fast *in vacuo*. Candide argues that this is consistent with the property just established. “Alas, my friend,” answers the inexorably pedantic Dr. Pangloss “you may be too quick. The experiment shows all gravitating bodies feel the same acceleration. But a uniform acceleration of the system may undo the gravitational one only if the inertial mass and the gravitational mass are the same.” Candide has to give in, and says they will have to wait for the outcome of the relevant experiment.

<sup>4</sup> The action of  $D(A)_\mu$  on the Vierbein is defined as

$$D_\mu(A) t_\nu^c = \partial_\mu t_\nu^c + A^c_{d\mu} t_\nu^d.$$



Nevertheless, the two gentlemen are quite satisfied with their result. Pending a positive experimental outcome, the gravitational force can be described by this composite gauge potential  $F_{\lambda\mu}^\kappa$  of their original  $SO(3,1)$  potentials.

They embark now on a natural question. What is the field strength related to this gauge potential? Dr. Pangloss quickly comes up with the answer

$$R_{\lambda\mu\nu}^\kappa(\Gamma) = \partial_\mu \Gamma_{\lambda\nu}^\kappa + \Gamma_{\alpha\mu}^\kappa \Gamma_{\lambda\nu}^\alpha - \mu \leftrightarrow \nu.$$

They do check that after a general coordinate transformation indeed the inhomogeneous part cancels as behooves a field strength.

As usual, Candide gets uneasy. “Dr. Pangloss, should this gravitational field strength not be related to our local Lorentz field strength (8)?”

The latter does agree that this may be true. The answer he gives is obtained by plugging his original definition of  $\Gamma$  in terms of translation and local Lorentz potential, Eq. (10), into the new field strength.

And indeed, Candide’s expectation was justified. The field strength of gravity turns out to be a contraction of the  $SO(3,1)$  field tensor with two Vierbeins

$$\hat{R}_{ab\mu\nu}(A)t^{a\kappa}t^b_{\lambda} = R_{\lambda\mu\nu}^\kappa(\Gamma). \tag{11}$$

The two are quite happy with their findings. The field strength of gravity  $R(\Gamma)$  is identical to the  $SO(3,1)$  field strength contracted with two Vierbeins. They conclude that the gravitational force is described by a gauge potential, a composite of Vierbein fields and the  $SO(3,1)$  gauge potential. They decide for a good dinner and exit the narrative.

**2.1.1. Epilogue to the dialogue**

Clearly, had the hands of the protagonists not been tied up on their back by their lack of knowledge of geometric concepts, the whole conversation could have ended at Eq. (10). The latter becomes after some trivial rearrangement the condition of metricity on the Vierbein field.

The geometry for an affine connexion with torsion was developed by Cartan [18]. In the seventies, Cartan’s geometric viewpoint was taken up again by Trautman and Hehl [18].

Still, we have to choose the pure gravity term in the SM action. Of course, there is the curvature  $R(\Gamma)$ . However, with torsion, the Riemann tensor is no longer symmetric in the exchange of the first pair of indices with the last pair. Hence, contracting the Riemann tensor with the epsilon tensor gives a non-vanishing pseudoscalar  $\tilde{R}(\Gamma)$ . There is one more independent scalar that can be taken as the Nieh–Yan invariant [23]. In this note, we will stick to the curvature  $R(\Gamma)$ .

What is certainly not academic is that this gauge gravity has a Palatini-type formalism built-in through the local Lorentz potentials. They are independent of the Vierbein fields. Just like in Einstein's gravity, the Palatini point of view is the independence of the affine connection of the metric.

In conclusion, the Standard Model produces gauge gravity, along with the strong and em forces. By constraining the torsion in Eq. (9) to vanish one retrieves Einstein's gravity.

And luckily, the original hypercharge gauge anomaly cancellations of the strong, electroweak and hypercharge forces [11] do include as well that of gravity [21]! They fix the hypercharge assignments, once the electric charge, say of the electron, is fixed.

Evidently, there is no clue why the electroweak and gravitational mass scale are so far apart. That is, scale invariance is badly broken.

This brings us to Weyl transformations. They do replace the scaling of coordinates and become an internal symmetry when the Vierbein fields are available. They do act multiplicatively on Vierbeins.

However, they do leave the Lorentz potential invariant, in analogy with the invariance of the affine connection in the Palatini version of Einstein gravity. Hence, the covariant derivative  $\partial_\mu + iA_\mu$  is invariant. And so is their commutator  $\hat{R}_{\mu\nu}(A)$  in (8). Hence, the Riemann curvature transforms according to Eq. (11) like the product of two Vierbein fields<sup>5</sup>. There is a good use for this in Section 3 when discussing the high field behaviour of the Higgs field in Higgs inflation.

### 3. Does the Standard Model accommodate the inflationary universe?

Inflation refers to a period in the early universe where the energy of the universe was all stored in the potential energy of the inflaton field. In order to have sufficient inflation, the field is supposed to roll slowly, *i.e.* the potential energy of the field at these high values should be flat, *i.e.* approximately translation invariant.

This flatness obviously excludes the Higgs potential  $V_{\text{ew}} = \frac{\lambda}{4} \left( H^\dagger H - \frac{v^2}{2} \right)^2$  with  $v \sim 246$  GeV as a candidate for inflation.

Bezrukov *et al.* [22] exploited the fact that Newton's constant decreases or rather the Planck mass increases when they added the Higgs coupled to curvature

$$M_{\text{P}}^2 \rightarrow M_{\text{P}}^2 + 2\xi|H|^2 .$$

---

<sup>5</sup> On matter fields, they act with a power fixed by the engineering dimension. Symmetry under Weyl transformations admits only dimensionless couplings.

To see the consequences at high field values, the traditional approach is to make a Weyl transformation on the gravitational field such that in the new, ‘‘Einstein’’ frame, the constant in front of the curvature is again  $M_{\text{P}}^2$  and the Higgs field in that frame has a canonical kinetic term<sup>6</sup>. The Higgs potential in the Einstein frame then becomes due to the decrease of Newton’s constant

$$\hat{V}(H^\dagger H) = \frac{V_{\text{ew}}(H^\dagger H)}{\left(1 + 2\xi \frac{H^\dagger H}{M_{\text{P}}^2}\right)^2}. \quad (12)$$

Now, the potential is flat at high field values  $|H| \gg \frac{M_{\text{P}}}{\sqrt{\xi}}$ , whereas the electroweak breaking at  $|H| \sim v$  is left unchanged unless  $\xi \sim 10^{17}$ . If we want the inflationary plateau large enough to create enough e-foldings  $\xi$  must be large, of the order of  $10^{7-8}$ . The original *raison d’être* of the  $|H|^2 R$  term was that it enhances an internal symmetry, Weyl symmetry, for a fixed numerical value of the coupling [20],  $\xi = -1/12$ . A huge value of the coupling seems unnatural. However, we want to describe inflation, an event where scale invariance was broken in an unprecedented fashion. It is then somewhat of a miracle that a numerical window of couplings describing successful inflation exists at all.

To see in more detail how the flattening of the potential comes about, we write down the relevant part of the action in terms of the radial Higgs field  $h$  with  $h^2 = 2H^\dagger H$  and drop its couplings to the gauge fields

$$S_{\text{J}} = \int d^4x \sqrt{-g} \left( -\frac{1}{2} (M_{\text{P}}^2 + \xi h^2) R + \frac{1}{2} g^{\mu\nu} \partial_\mu h \partial_\nu h - V_{\text{ew}}(h) \right).$$

This is the action in the Jordan frame. It has a Higgs-field-dependent gravitational coupling and the canonical kinetic term for the Higgs and the canonical potential causing electroweak breaking. The question is what the effect of the Higgs dependence of the gravity coupling will be on the potential. This can be computed directly [26] in the Jordan frame.

Alternatively, a Weyl transformation  $\exp(-\omega)$  on the gravitational field can restore the coupling in front of the curvature by the physical coupling  $M_{\text{P}}^2$ <sup>7</sup>. This Weyl transformation acts only on the Vierbein fields, and using the relation between the curvature tensor and the local Lorentz field strength, we see that the curvature  $R$  and the metric transform both like  $\exp(-2\omega)$  and the determinant like  $\exp(4\omega)$

<sup>6</sup> For a justification, see reference [26].

<sup>7</sup> The discussion here is in the context of gauge gravity, but the mathematics is identical to that in Ref. [24].

$$S_E = \int d^4x \sqrt{-g} \left( -\frac{1}{2} (M_P^2 + \xi h^2) \exp(2\omega) R + \frac{1}{2} g^{\mu\nu} \exp(2\omega) \partial_\mu h \partial_\nu h - \exp(4\omega) V(h) \right).$$

This is the action in the Einstein frame. Requiring that the curvature coupling is the Planck mass fixes  $\omega$  in terms of  $h$

$$\exp(-2\omega) = 1 + \xi \frac{h^2}{M_P^2}.$$

The prefactor  $\exp(4\omega)$  of the Higgs potential is thereby determined as in Eq. (12).

The Higgs kinetic term is normalized in the Einstein frame by introducing the renormalized Higgs  $\phi$ . It becomes quite simply related to  $h$  by

$$h = \frac{M_P}{\sqrt{\xi}} \sinh \left( \sqrt{\xi} \frac{\phi}{M_P} \right). \quad (13)$$

Thus, in the Einstein frame, the field with a canonical kinetic term is very non-linear function of the original Higgs field. Moreover, the potential  $U(\phi) = \hat{V}(h(\phi))$  becomes

$$U_{\text{Pal}}(\phi) = U_\infty \tanh^4 \left( \frac{\xi \phi}{M_P^2} \right). \quad (14)$$

The potential has an asymptotic translation symmetry in the Einstein frame. This is what we need for inflation. In the Jordan frame, this becomes an asymptotic scale symmetry.

So far for gauge gravity.

For the Einstein gravity, the torsion in Eq. (9) is vanishing. Thus, the Lorentz potential depends now on the Vierbein and so does the affine connection in (10). Hence, we have to do with metric Einstein gravity, not Palatini–Einstein gravity.

As a consequence, the renormalization of the Higgs kinetic term gets replaced by

$$\exp(2\omega) \rightarrow \exp(2\omega) + \exp(4\omega) \xi^2 h^2 / 6M_P^2. \quad (15)$$

Note that for small  $h/M_P$ , the new term is a dimension-6 operator suppressed by a factor of  $\Lambda_m = \sqrt{6}M_P/\xi$ .

The relation of  $h$  to the renormalized Higgs field  $\phi$  is for large fields and large value of  $\xi$

$$h = \frac{M_P}{\sqrt{\xi}} \exp \left( \frac{\phi}{\sqrt{6}M_P} \right), \quad \text{metric Einstein gravity}$$

and the potential in the Einstein frame becomes under the same conditions

$$U_m(\phi) = U_\infty \left( 1 - 2 \exp \left( -2\phi / \left( \sqrt{6} M_{\text{P}} \right) \right) \right). \quad (16)$$

For slow roll, the inflationary Hubble parameter equals

$$H_{\text{I}}^2 = \frac{U_\infty}{3M_{\text{P}}^2} = \frac{\lambda}{24} \left( \frac{M_{\text{P}}}{\xi} \right)^2, \quad (17)$$

and this will be the benchmark scale for the inflationary regime.

With the input of potentials (14) and (16), we can extract the values of  $\xi$  and other inflationary parameters, and compare them in both cases.

Inflation in the Einstein frame is taking place between  $\phi_i$  at time  $t_i$  and  $\phi_f$  at  $t_f$  and the number  $N$  of e-foldings in that frame is given by the integral

$$N = \frac{1}{M_{\text{P}}^2} \int_{\phi_f}^{\phi_i} U(\phi) / (\partial U(\phi) / \partial \phi) d\phi.$$

We skip further details on the traditional comparison to data and give the results. The upshot is that the number of e-folds and the scalar tilt are numerically pretty much the same in the gauge gravity case and the Einstein gravity case [22]. However, the  $\xi$  value for gauge gravity is  $10^7$ , while  $10^4$  for Einstein gravity. It follows that for gauge gravity the ratio tensor to scalar tilt is very small,  $\sim 10^{-11}$  for gauge gravity, seven orders of magnitude smaller than its Einstein gravity value.

Unfortunately, the measurement of this ratio will be very hard below  $r \sim 10^{-5}$ . Of course, it suffices to find an experimental value for  $r$  above this barrier and the Palatini case is falsified.

The current upper bound [25] is  $r < 0.064$  and together with the Hubble rate during inflation  $H_{\text{I}} = 1.06 \times 10^{-4} r^{1/2} M_{\text{P}}$  it follows that  $H_{\text{I}} < 6 \times 10^{13}$  GeV.

So far, for the classical description of both metric and Palatini inflation there is a striking parametric difference between the two cases, Eqs. (16) and (14). The plateau begins parametrically earlier in the Palatini case and, as a consequence, inflation takes place well below the Planck scale. Asymptotically, the height of the plateau is the same.

### 3.1. The issue of quantum corrections

The coupling  $\xi$  of the non-minimal term is quite large. This raises two concerns. The first is ( $h \ll M_{\text{P}}$ ) how big the unitarity violations in scattering amplitudes are for small Higgs field values. The second and most important is whether quantum corrections affect slow roll.

Elastic Higgs–Higgs scattering is the least complicated case to demonstrate this. We take the metric approach. From reference [28], one finds for the  $J = 0$  elastic amplitude with one graviton exchange in the Jordan frame that the maximum energy in the center-of-mass frame is

$$E_{\text{CM}}^2 = \frac{\pi}{2} \frac{M_{\text{P}}^2}{(1 + \xi/12)^2}, \quad h \ll M_{\text{P}}. \quad (18)$$

If  $\xi = -1/12$ , there is no bound. The reason is that the amplitude vanishes in the Weyl-invariant theory [20]. For large  $\xi$ , the typical scale  $\Lambda_m \equiv \frac{M_{\text{P}}}{\xi}$  emerges for  $E_{\text{CM}}$ .

This result indicates that for energies well below  $M_{\text{P}}$ , perturbative unitarity breaks down. Does this mean the minimalistic view of the Standard Model is not tenable in the presence of this large coupling? Not necessarily, because a strong coupling regime of the SM could set in.

The scale  $\Lambda_m$  is again retrieved in the transition from the Jordan frame to the Einstein frame. Equation (15) tells us that the dimension six operator characteristic for the metric approach is

$$\frac{1}{\Lambda_m^2} h^2 \partial_\mu h \partial^\mu h \quad \text{with} \quad \Lambda_m = \frac{\sqrt{6} M_{\text{P}}}{\xi}.$$

This is the cut-off scale for small  $h/M_{\text{P}}$  and is of the same order of magnitude as the unitarity limit.

A different way to come to the same conclusion is to look at how a given interaction term involving the Higgs in the Jordan frame looks in the Einstein frame. Then there are two regimes, small Higgs values ( $\xi h \ll M_{\text{P}}$ ) and large Higgs values. For the first regime, the answer is what we found already for the cut-off. For the latter regime, the cut-off  $\Lambda_m = \sqrt{6} M_{\text{P}}$ . This is parametrically higher than the value of the inflationary Hubble scale (17). This was the point of the authors of reference [27]: the cut-off is in principle background dependent. However, some authors have expressed dissenting opinions [29].

**3.1.1. The Palatini case**

Let us now contrast the Palatini case with the metric case. We follow the same approach mentioned at the end of the last section but now with the relation between the Higgs fields in Jordan and Einstein frames

$$h = \frac{M_P}{\sqrt{\xi}} \sinh\left(\frac{\sqrt{\xi}\phi}{M_P}\right). \tag{19}$$

Take a simple term in the SM action, the Yukawa term  $h\bar{\psi}\psi$ . In the Einstein frame,  $h$  is written in terms of the  $\phi$  field and the fermion fields transform according to their engineering dimension  $\psi_E = \exp(-\frac{3}{2}\omega)\psi$ . Taking this into account we find for the Yukawa term in the Einstein frame

$$\exp(\omega)h\bar{\psi}_E\psi_E = \frac{M_P}{\sqrt{\xi}} \tanh\left(\frac{\sqrt{\xi}\phi}{M_P}\right) \bar{\psi}_E\psi_E.$$

In the small  $\phi$  regime the expansion gives

$$\exp(\omega)h\bar{\psi}_E\psi_E = \phi\bar{\psi}_E\psi_E + \frac{1}{A_{\text{Pal}}^2}\phi^3\bar{\psi}_E\psi_E + \dots$$

where  $A_{\text{Pal}} = \frac{M_P}{\sqrt{\xi}}$  is the scale where new physics or a strong coupling regime of the SM sets in. This is parametrically less conservative than the metric case.

**3.1.2. Quantum corrections to the inflationary plateau**

This question is really answered by knowing how to compute the effective potential for the Higgs field.

In the Jordan frame, one can set up the effective potential in the conventional way by a constraint in the path integral. The constraint is of the invariant form

$$\delta\left(\overline{H^\dagger H} - \frac{h^2}{2}\right). \tag{20}$$

The bar means a convenient space-time average of  $H^\dagger H$ . The action is  $S_J + S_m$ ,  $S_m$  being the SM action without the Higgs sector.

The use of an invariant variable seems obligatory due to the way one extracts the e-foldings from the potential.

In the Einstein frame, the shift invariance is perhaps better to exploit. However, it is a shift invariance in a variable that is a non-linear function of the original Higgs field. Clearly, a reliable approximation is needed [27].

Recapitulating, we have found that for the Palatini case the cut-off is certainly parametrically larger than the inflationary Hubble constant Eq. (17)

$$\Lambda_{\text{Pal}} \geq \xi H_I. \quad (21)$$

This is certainly an encouraging result. It means that in the inflationary phase one has not to appeal to new degrees of freedom beyond the SM.

For a recent quite positive conclusion on the Palatini case see [30]. Gauge gravity has as mentioned more than only scalar curvature for the gravity sector, namely the pseudo-scalar curvature and the Nieh–Han invariant [23]. In that context, Higgs inflation seems to be a generic phenomenon.

#### 4. Epilogue

It was in Ann Arbor that we celebrated Tini’s sixtieth birthday. One of the contributions to the after-dinner speeches was a very apt and witty rendition of his personality in the form of a poem. The title [31] was

“The Volcano”.

We have to learn to live without this force of nature.

#### REFERENCES

- [1] H. van Dam, M.J.G. Veltman, «Massive and mass-less Yang–Mills and gravitational fields», *Nucl. Phys. B* **22**, 397 (1970).
- [2] G. ’t Hooft, M.J.G. Veltman, «One loop divergencies in the theory of gravitation», *Ann. Inst. H. Poincaré Phys. Theor. A* **20**, 69 (1974).
- [3] M.J.G. Veltman, «Cosmology and the Higgs Mass», *Phys. Rev. Lett.* **34**, 777 (1975); A.D. Linde, «Is the Lee constant a cosmological constant?», *Pis’ma Zh. Eksp. Toer. Fiz.* **19**, 320 (1974).
- [4] M.J.G. Veltman, «Quantum Theory of Gravitation», *Conf. Proc. C* **7507281**, 265 (1975).
- [5] G. ’t Hooft, M.J.G. Veltman, «DIAGRAMMAR», CERN Yellow Report, CERN-73-09.
- [6] M.J.G. Veltman, «Divergence Conditions and Sum Rules», *Phys. Rev. Lett.* **17**, 553 (1966).
- [7] LHCb Collaboration (R. Aaij *et al.*), «Test of lepton universality in beauty-quark decays», [arXiv:2103.11769](https://arxiv.org/abs/2103.11769) [hep-ex].
- [8] Sougato Bose *et al.*, «Spin Entanglement Witness for Quantum Gravity», *Phys. Rev. Lett.* **119**, 240401 (2017), [arXiv:1707.06050](https://arxiv.org/abs/1707.06050) [quant-ph]; Ch. Marletto, V. Vedral, «Gravitationally-induced entanglement between two massive particles is sufficient evidence of quantum effects in gravity», *Phys. Rev. Lett.* **119**, 240402 (2017), [arXiv:1707.06036](https://arxiv.org/abs/1707.06036) [quant-ph].



- [9] F. Dyson, «Is the graviton detectable?», *Int. J. Mod. Phys. A* **28**, 1330041 (2012); T. Rothman, S. Boughn, «Can gravitons be detected?», *Found. Phys.* **36**, 1801 (2006), [arXiv:gr-qc/0601043](https://arxiv.org/abs/gr-qc/0601043).
- [10] G. 't Hooft, «Renormalizable Lagrangians for massive Yang–Mills fields», *Nucl. Phys. B* **35**, 167 (1971).
- [11] C. Bouchiat, J. Iliopoulos, P. Meyer, «An anomaly-free version of Weinberg's model», *Phys. Lett. B* **38**, 519 (1972).
- [12] C.P. Korthals Altes, M. Perrottet, «Anomalous Ward-identities, gauge-variance and appearance of ghosts in Higgs–Kibble type theories», *Phys. Lett. B* **39**, 546 (1972).
- [13] T.W.B. Kibble, «Lorentz Invariance and the Gravitational Field», *J. Math. Phys.* **2**, 212 (1961); R. Utiyama, «Invariant Theoretical Interpretation of Interaction», *Phys. Rev.* **101**, 1597 (1956); D.W. Sciama, «On the analogy between charge and spin in General Relativity», in: «Recent Developments in General Relativity, Festschrift for Infeld Press», *Pergamon Press*, Oxford 1962, p. 415.
- [14] H. Weyl, «A remark on the coupling of gravitation and electron», *Phys. Rev.* **77**, 699 (1950).
- [15] Voltaire, «Candide, ou l'Optimisme», traduit de l'Allemand de Mr le Docteur Ralph, Genève 1759, Ed. Cramer.
- [16] M.R. Cohen, I.E. Drabkin, «A Source Book in Greek Science», *McGraw Hill*, New York 1948.
- [17] Galileo Galilei, «Dialogues Concerning Two New Sciences», *MacMillan*, New York 1914, pp. 62–64, translation by H. Crew, A. Favaro.
- [18] E. Cartan, «Sur une généralisation de la notion de courbure de Riemann et les espaces de torsion», *C.R. Acad. Sci. (Paris)* **174**, 593 (1922); E. Cartan, «Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie)», *Annales scientifiques de l'École Normale Supérieure, Série 3* **40**, 325 (1923); «Sur les variétés à connexion affine, et la théorie de la relativité généralisée (première partie) (Suite)», *Annales scientifiques de l'École Normale Supérieure, Série 3* **41**, 1 (1924); «Sur les variétés à connexion affine, et la théorie de la relativité généralisée (deuxième partie)», *Annales scientifiques de l'École Normale Supérieure, Série 3* **42**, 17 (1925); A. Trautman, «Einstein–Cartan Theory», in: J.-P. Francoise, G.L. Naber, S.T. Tsou (Eds.) «Encyclopedia of Mathematical Physics», *Elsevier*, 2006, pp. 189–195; F.W. Hehl, P. von der Heyde, G.D. Kerlick, J.M. Nester, «General relativity with spin and torsion: Foundations and prospects», *Rev. Mod. Phys.* **48**, 393 (1976); I.L. Shapiro, «Physical aspects of the space–time torsion», *Phys. Rep.* **357**, 113 (2002), [arXiv:hep-th/0103093](https://arxiv.org/abs/hep-th/0103093).
- [19] A. Palatini, «Deduzione invariante delle equazioni gravitazionali dal principio di Hamilton», *Rend. Circ. Mat. Palermo* **43**, 203 (1919); A. Einstein, «Einheitliche Feld-theorie von Gravitation und Elektrizität», *Sitzungber. Preuss. Akad. Wiss.* **1925**(XXII), 414 (1925).
- [20] C.G. Callan, S. Coleman, R. Jackiw, «A New Improved Energy-Momentum Tensor», *Ann. Phys.* **59**, 42 (1970); S. Deser, «Scale Invariance and Gravitational Coupling», *Ann. Phys.* **59**, 248 (1970).

- [21] C.Q. Geng, R.E. Marshak, «Uniqueness of quark and lepton representations in the Standard Model from the anomalies viewpoint», *Phys. Rev. D* **39**, 693 (1989); J.A. Minahan, P. Ramond, R.C. Warner, «Comment on anomaly cancellation in the Standard Model», *Phys. Rev. D* **41**, 715 (1990); C.Q. Geng, R.E. Marshak, «Reply to “Comment on anomaly cancellation in the Standard Model”», *Phys. Rev. D* **41**, 717 (1990); P.H. Frampton, R.N. Mohapatra, «Partial derivation of transformation properties of quarks and leptons», *Phys. Rev. D* **50**, 3569 (1994).
- [22] F.L. Bezrukov, M. Shaposhnikov, «The Standard Model Higgs boson as the inflaton», *Phys. Lett. B* **659**, 703 (2008), [arXiv:0710.3755 \[hep-th\]](#).
- [23] M. Shaposhnikov, A. Shkerin, I. Timiryasov, S. Zell, «Higgs inflation in Einstein–Cartan gravity», *J. Cosmol. Astropart. Phys.* **2021**, 008 (2021), [arXiv:2007.14978 \[hep-ph\]](#). See also: M. Langvik, J.-M. Ojanpera, S. Raatikainen, S. Rasanen, «Higgs Inflation with Holst and Nieh-Han terms», *Phys. Rev. D* **103**, 083514 (2021), [arXiv:2007.12595 \[astro-ph.CO\]](#).
- [24] F. Bauer, D.A. Demir, «Inflation with non-minimal coupling: Metric *vs.* Palatini formulations», *Phys. Lett. B* **665**, 222 (2008), [arXiv:0803.2664 \[hep-ph\]](#); F. Bauer, D.A. Demir, «Higgs–Palatini inflation and unitarity», *Phys. Lett. B* **698**, 425 (2011), [arXiv:1012.2900 \[hep-ph\]](#).
- [25] Planck Collaboration (Y. Akrami *et al.*), «Planck 2018 results. I. Overview and the cosmological legacy of Planck», *Astron. Astrophys.* **641**, A1 (2020), [arXiv:1807.06205 \[astro-ph.CO\]](#).
- [26] N. Makino, M. Sasaki, «The Density Perturbation in the Chaotic Inflation with Non-Minimal Coupling», *Prog. Theor. Phys.* **86**, 103 (1991); R. Fakir, S. Habib, W. Unruh, «Cosmological density perturbations with modified gravity», *Astrophys. J.* **394**, 396 (1992).
- [27] F. Bezrukov, A. Magnin, M. Shaposhnikov, S. Sibiryakov, «Higgs inflation: consistency and generalisations», *J. High Energy Phys.* **1101**, 016 (2011), [arXiv:1008.5157 \[hep-ph\]](#).
- [28] T. Han, S. Willenbrock, «Scale of Quantum Gravity», *Phys. Lett. B* **616**, 215 (2005), [arXiv:hep-ph/0404182](#).
- [29] J.L.F. Barbon, J.R. Espinosa, «On the Naturalness of Higgs Inflation», *Phys. Rev. D* **79**, 081302 (2009), [arXiv:0903.0355 \[hep-ph\]](#); J.L.F. Barbon, J.A. Casas, J. Elias-Miro, J.R. Espinosa, «Higgs Inflation as a Mirage», *J. High Energy Phys.* **1509**, 027 (2015), [arXiv:1501.02231 \[hep-ph\]](#); C.P. Burgess, H.M. Lee, M. Trott, «Power-counting and the validity of the classical approximation during inflation», *J. High Energy Phys.* **0909**, 103 (2009), [arXiv:0902.4465 \[hep-ph\]](#); C.P. Burgess, H.M. Lee, M. Trott, «On Higgs inflation and naturalness», *J. High Energy Phys.* **1007**, 007 (2010), [arXiv:1002.2730 \[hep-ph\]](#); G.F. Giudice, H.M. Lee, «Unitarizing Higgs Inflation», *Phys. Lett. B* **694**, 294 (2011), [arXiv:1010.1417 \[hep-ph\]](#).

- [30] J. McDonald, «Does Palatini Higgs Inflation Conserve Unitarity?», *J. Cosmol. Astropart. Phys.* **04**, 069 (2021), [arXiv:2007.04111 \[hep-ph\]](#).
- [31] D. Maison, «Der Vulkan (TINICATEPETL)», in: R. Akhoury, B. de Wit, P. van Nieuwenhuizen, H. Veltman (Eds.) «Gauge Theories — Past and Future. In Commemoration of the 60<sup>th</sup> Birthday of M. Veltman. World Scientific Series in 20<sup>th</sup> Century Physics: Volume 1», *World Scientific*, Singapore 1992, p. 1.