

CERN-EP-2021-271
2022/01/22

CMS-BPH-18-004

Observation of $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ and $B_s^0 \rightarrow \psi(2S)K_S^0$ decays

The CMS Collaboration*

Abstract

Using a data sample of $\sqrt{s} = 13$ TeV proton-proton collisions collected by the CMS experiment at the LHC in 2017 and 2018 with an integrated luminosity of 103 fb^{-1} , the $B_s^0 \rightarrow \psi(2S)K_S^0$ and $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decays are observed with significances exceeding 5 standard deviations. The resulting branching fraction ratios, measured for the first time, correspond to $\mathcal{B}(B_s^0 \rightarrow \psi(2S)K_S^0)/\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0) = (3.33 \pm 0.69 \text{ (stat)} \pm 0.11 \text{ (syst)} \pm 0.34 (f_s/f_d)) \times 10^{-2}$ and $\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)/\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0) = 0.480 \pm 0.013 \text{ (stat)} \pm 0.032 \text{ (syst)}$, where the last uncertainty in the first ratio is related to the uncertainty in the ratio of production cross sections of B_s^0 and B^0 mesons, f_s/f_d .

Submitted to the European Physical Journal C

1 Introduction

Decays of neutral B mesons into charmonium resonances (J/ψ , $\psi(2S)$, etc.) are well suited to study the flavour sector of the standard model (SM) and to search for indications of new physics beyond the SM. In the last decade, interest in b hadron decays to final states containing a charmonium resonance has increased after several exotic hadrons have been observed as intermediate resonances in multibody decays. Starting from the observation of $X(3872)$ [1], many new charmonium-like states have been observed, such as $X(4140)$ [2–5], $Y(4260)$ [6, 7], and others, with properties (mass, width and decay pattern) not fitting into the landscape of traditional charmonium states. The first charged tetraquark candidate, $Z(4430)^+$, was discovered in the $B \rightarrow \psi(2S)K\pi^+$ decay as a peak in the $\psi(2S)\pi^+$ mass spectrum [8–11]. Many other exotic hadrons have been observed in the last 15 years [12, 13], and the nature of most of them is still unclear. Moreover, channels whose final state is accessible both from B and \bar{B} can be used to measure time-dependent CP asymmetry [14–27] as well.

This paper presents the first measurement of the $B_s^0 \rightarrow \psi(2S)K_S^0$ and $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decays, using a data sample of proton-proton collisions at $\sqrt{s} = 13$ TeV collected by the CMS experiment at the CERN LHC in 2017 and 2018 with an integrated luminosity of 103 fb^{-1} [28, 29]. Both decays can potentially be used for CP asymmetry measurements, and, in addition, the second one can also be used to search for intermediate exotic resonances. The $\psi(2S)$ and K_S^0 mesons are reconstructed using their decays into $\mu^+\mu^-$ and $\pi^+\pi^-$, respectively. The $B^0 \rightarrow \psi(2S)K_S^0$ decay is chosen as the normalization channel for the measurement of the branching fractions, since its probability is precisely known [13], and its topology and kinematic properties are similar to those of the $B_s^0 \rightarrow \psi(2S)K_S^0$ or $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decays. Therefore, using this normalization reduces the systematic uncertainties related to muon and track reconstruction. The relative branching fractions are measured using the relations

$$R_s \equiv \frac{\mathcal{B}(B_s^0 \rightarrow \psi(2S)K_S^0)}{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0)} = \frac{f_d}{f_s} \frac{\epsilon(B^0 \rightarrow \psi(2S)K_S^0)}{\epsilon(B_s^0 \rightarrow \psi(2S)K_S^0)} \frac{N(B_s^0 \rightarrow \psi(2S)K_S^0)}{N(B^0 \rightarrow \psi(2S)K_S^0)},$$

$$R_{\pi^+\pi^-} \equiv \frac{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)}{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0)} = \frac{\epsilon(B^0 \rightarrow \psi(2S)K_S^0)}{\epsilon(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)} \frac{N(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)}{N(B^0 \rightarrow \psi(2S)K_S^0)}, \quad (1)$$

where \mathcal{B} is the branching fraction, N is the number of reconstructed events in data, ϵ is the total reconstruction efficiency, and f_d/f_s is the ratio of production cross sections of B^0 and B_s^0 mesons (also called fragmentation fraction ratio). Charge-conjugate states are implied to be included throughout the paper.

Tabulated results are provided in the HEPData record for this analysis [30].

2 The CMS detector and simulated event samples

The central feature of the CMS apparatus [31] is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a brass and scintillator hadron calorimeter. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid.

Events of interest are selected using a two-tiered trigger system [32]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a time interval of less than $4 \mu\text{s}$ [33]. The first-level trigger used in this analysis requires at least two muons. The second level, known

as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing that reduces the event rate to around 1 kHz before data storage. The high-level trigger algorithm used in the analysis requires two opposite-sign muons compatible with the dimuon decay of a $\psi(2S)$ meson with transverse momentum (p_T) larger than 18 GeV.

Simulated Monte Carlo samples for the decays of interest are generated for the analysis. The PYTHIA 8.230 package [34] with the CP5 tune [35] is used to simulate the production of the B^0 and B_s^0 mesons, whose subsequent decays are performed by EVTGEN 1.6.0 [36], where final-state photon radiation is included using PHOTOS 3.61 [37, 38]. The generated events are passed to a detailed GEANT4-based simulation [39] of the CMS detector, and are then processed using the same trigger and reconstruction as used for the collision data. The simulation includes effects from multiple proton-proton interactions in the same or nearby bunch crossings (pileup) with the multiplicity distribution tuned to match those of the data.

3 Event reconstruction and selection

The reconstruction procedure starts with finding two muons of opposite charges, that must match those that triggered the event readout. The muon candidates are required to have $p_T(\mu^\pm) > 3$ GeV, a pseudorapidity $|\eta(\mu^\pm)| < 2.4$, and to satisfy general identification (soft-muon) criteria [40]. The two muons with a two-prong vertex fit probability $P_{\text{vtx}}(\mu^+\mu^-) > 1\%$ are paired to form the $\psi(2S)$ candidate, which must have $p_T(\psi(2S)) > 18$ GeV and an invariant mass $3500 < m(\mu^+\mu^-) < 3950$ MeV (the world average $\psi(2S)$ meson mass is $m_{\psi(2S)}^{\text{PDG}} = 3686.10$ MeV [13]).

The $K_S^0 \rightarrow \pi^+\pi^-$ candidates are formed from displaced two-prong vertices, as described in Ref. [41]. The $\pi^+\pi^-$ invariant mass is required to be within ± 20 MeV of the world average value $m_{K^0}^{\text{PDG}} = 497.611$ MeV [13], which corresponds to approximately three times the mass resolution. Selected π^+ and π^- tracks are then refitted with their invariant mass constrained to $m_{K^0}^{\text{PDG}}$, and the obtained K_S^0 candidate is required to have $P_{\text{vtx}}(\pi^+\pi^-) > 1\%$.

The $B \rightarrow \psi(2S)K_S^0$ candidates are obtained through a kinematic vertex fit on the $\mu^+\mu^-K_S^0$ system which constrains the dimuon mass to $m_{\psi(2S)}^{\text{PDG}}$. The K_S^0 candidates are required to have $p_T(K_S^0) > 1$ GeV, a 3D pointing angle between $\vec{D}(K_S^0)$ and $\vec{p}(K_S^0)$ to satisfy $\cos(\vec{D}(K_S^0), \vec{p}(K_S^0)) > 0.99$, and a transverse displacement significance for K_S^0 of $D_{xy} > 5\sigma_{D_{xy}}$. Here $\vec{D}(K_S^0)$ denotes the vector from the K_S^0 production vertex to the K_S^0 decay vertex, while D_{xy} and $\sigma_{D_{xy}}$ correspond to the length of \vec{D}_{xy} , the transverse component of \vec{D} , and its uncertainty. To suppress the combinatorial background, additional requirements are applied: $P_{\text{vtx}}(\mu^+\mu^-K_S^0) > 5\%$, $\cos(\vec{D}_{xy}(B), \vec{p}_T(B)) > 0.99$, and $D_{xy}(B) > 5\sigma_{D_{xy}(B)}$, where the B meson transverse displacement $D_{xy}(B)$ is calculated with respect to the primary vertex (PV). From all reconstructed proton-proton collision points, the PV is chosen as the one with the smallest B pointing angle, as in Refs. [42–44]. The pointing angle is the angle formed by the B candidate momentum and the vector from the PV to the reconstructed B candidate vertex. Furthermore, if in this procedure any of the tracks used in the B candidate reconstruction is included in the fit of the chosen PV, the track is removed, and the PV is refitted.

For the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ candidates, two additional, oppositely charged, high-purity [45] tracks, assumed to be pions and having $p_T > 0.9$ GeV, are included in the B meson vertex fit, while the rest of the selection criteria are the same.

4 Observation of the $B_s^0 \rightarrow \psi(2S)K_S^0$ decay

The measured $\psi(2S)K_S^0$ invariant mass distribution is presented in Fig. 1 (left). The B^0 signal (left peak) is described with a double Gaussian function with common mean, whose parameters are free to vary in an unbinned maximum-likelihood fit. It is found in simulation that the $B_s^0 \rightarrow \psi(2S)K_S^0$ signal (right peak) has the same shape as the $B^0 \rightarrow \psi(2S)K_S^0$ signal, but it is about 10% wider, because of the larger energy release in the decay. Therefore, the B_s^0 signal is modelled with a double Gaussian function of the same shape as the B^0 signal, with the resolution parameters scaled by the ratio of the widths found in the simulation. The background is modelled with an exponential function. The good quality of the fit is verified by calculating the χ^2 between the binned distribution and the fit function, resulting in $\chi^2 = 83$ for 91 degrees of freedom.

The ratio of signal yields $N(B_s^0 \rightarrow \psi(2S)K_S^0)/N(B^0 \rightarrow \psi(2S)K_S^0) = (6.8 \pm 1.4) \times 10^{-3}$ is extracted from the fit. Its uncertainty is calculated by taking into account the correlation between the uncertainties in B_s^0 and B^0 yields, which are found to be 113 ± 23 and 16660 ± 140 , respectively, where the uncertainties are statistical only.

The statistical significance of the $B_s^0 \rightarrow \psi(2S)K_S^0$ signal is evaluated with the likelihood ratio technique, comparing the background-only and signal-plus-background hypotheses, with the standard asymptotic formula [46], assuming that the conditions to apply Wilks' theorem [47] are satisfied. For a significance estimation, the mass difference between the B_s^0 and B^0 signals is fixed to the known value of 83.78 MeV [13]. The obtained significance is 5.2 standard deviations and varies in the range 5.1–5.4 standard deviations when accounting for the systematic uncertainties due to the choice of the fit model, discussed in Section 7.

5 Observation of the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay

As shown in Fig. 1 (right), the measured $\psi(2S)K_S^0\pi^+\pi^-$ mass distribution presents a clear $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ signal peak on top of a relatively small background. The B^0 signal is modelled with a double Gaussian function with common mean with all parameters free to vary, and the combinatorial background is described by an exponential function.




Figure 1: Measured invariant mass distributions of $\psi(2S)K_S^0$ (left) and $\psi(2S)K_S^0\pi^+\pi^-$ (right) candidates. The overlaid results from the fit are described in the text.

Studies of simulated events show that the $B_s^0 \rightarrow \psi(2S)K_S^0K^-\pi^+$ decay contributes to the reconstructed $\psi(2S)K_S^0\pi^+\pi^-$ mass distribution when the charged kaon is reconstructed as a pion.

This relevant background contribution is accounted for in the fit to data by including a dedicated component with a freely varying normalization and a fixed shape that is obtained from simulation (Fig. 1, right).

The signal yield $N(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)$ is found to be 3498 ± 87 , where the uncertainty is statistical only. The χ^2 between the binned distribution and the fit function is 75 for 92 degrees of freedom, demonstrating the good quality of the fit. The significance of the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ signal, evaluated as described in Section 4, exceeds 30 standard deviations.

The intermediate invariant mass distributions, corresponding to the four-body $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay, are produced using the $_s\mathcal{P}$ lot [48] technique to subtract the non- B^0 background, using the $m(\psi(2S)K_S^0\pi^+\pi^-)$ distribution fit described above. The correlations between the intermediate invariant masses and $m(\psi(2S)K_S^0\pi^+\pi^-)$ have been checked to be below 10%. Figures 2 and 3 show the 2- and 3-body invariant mass distributions. Overlaid are the predictions of the 4-body phase space simulations, which provide poor description of the data since the simulations do not account for the intermediate resonance structure. The simulation after application of the reweighting procedure described in Section 7 is also shown. The mass distributions of $\psi(2S)$ and one or two light mesons ($\psi(2S)K_S^0$, $\psi(2S)\pi^\pm$, $\psi(2S)K_S^0\pi^\pm$, $\psi(2S)\pi^+\pi^-$) do not present any significant narrow peak that could indicate a contribution from an exotic charmonium state. The small excess at about 4.3 GeV in the $m(\psi(2S)\pi^+)$ distribution (Fig. 2, bottom left) is not significant, and there is no similar excess in the $m(\psi(2S)\pi^-)$ distribution (Fig. 2, middle left). Moreover, exotic states previously found in this mass range are known to have large natural widths [12, 13]. Signs of the $K^*(892)^\pm$ (Fig. 2, middle and bottom right), $\rho(770)^0$ (Fig. 2, top left), and $K_1(1270)^0$ (Fig. 3, top right) resonances are seen in the mass distributions of $K_S^0\pi^\pm$, $\pi^+\pi^-$, and $K_S^0\pi^+\pi^-$, respectively.

6 Efficiencies

The total reconstruction efficiency for each decay channel is evaluated using samples of simulated events. It is calculated as the number of reconstructed events divided by the number of generated events, and includes the detector acceptance, trigger, and candidate reconstruction efficiencies. Only the ratios of such efficiencies are needed to measure the ratios R_s and $R_{\pi^+\pi^-}$, thus reducing the systematic uncertainties associated with muon and track reconstruction.

The obtained efficiency ratios are found to be

$$\frac{\epsilon(B^0 \rightarrow \psi(2S)K_S^0)}{\epsilon(B_s^0 \rightarrow \psi(2S)K_S^0)} = 1.019 \pm 0.013 \quad \text{and} \quad \frac{\epsilon(B^0 \rightarrow \psi(2S)K_S^0)}{\epsilon(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)} = 2.288 \pm 0.026,$$

where the uncertainties are statistical only and are related to the size of the simulated event samples. The first ratio is close to unity, as expected, while the second ratio is significantly greater than unity because of the presence of two additional tracks in the denominator.

The validation of Monte Carlo samples is performed by comparing distributions of variables used in the event selection between simulation and background-subtracted data. No significant deviation is found, and thus no systematic uncertainties in the efficiency ratio are assigned related to data-simulation discrepancies in those variables.

7 Systematic uncertainties

Many systematic uncertainties, related to the efficiency of the trigger as well as the reconstruction and identification of the muons, cancel out in the measured ratios R_s and $R_{\pi^+\pi^-}$. Since




Figure 2: Distributions of 2-body intermediate invariant masses from the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay. The data distributions (black dots) are background subtracted. Overlaid are the predictions of phase space simulations (red triangles), as well as the predictions after applying the reweighting procedure described in Section 7 (grey squares).

the $B_s^0 \rightarrow \psi(2S)K_S^0$ and $B^0 \rightarrow \psi(2S)K_S^0$ decays have the same number of tracks in the final state, uncertainties related to the track reconstruction are of the same size and correlated, and therefore cancel out when propagated to the measured ratio R_s . For the ratio $R_{\pi^+\pi^-}$, we consider an additional uncertainty of 4.2% from the uncertainty in the tracking efficiency of two additional pions [49].

The systematic uncertainty related to the choice of the fit model is evaluated by testing different models. The largest deviation in the measured ratio from its baseline value is taken as a systematic uncertainty, separately for the variations of the signal and background models. Several alternative signal models were considered. One is a double Gaussian function for B^0 and B_s^0 signals with the resolution shape fixed to the expectations taken from simulation with only the resolution scaling parameter being free in the fit. Another signal model is a Student's t -distribution [50] with the value of the n parameter fixed to the one measured in simulation.




Figure 3: Distributions of 3-body intermediate invariant masses from the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay. Data distributions (black dots) are background subtracted. Overlaid are the predictions of phase space simulations (red triangles), as well as the predictions after applying the reweighting procedure described in Section 7 (grey squares).

Alternative background models include polynomials of the second and third degrees, an exponential multiplied by a polynomial, and a power function multiplied by an exponential, where in all cases the background shape parameters are free to vary in the fits.




Figure 4: Background-subtracted $p_T(B^0)$ distribution in data for the $B^0 \rightarrow \psi(2S)K_S^0$ signal. The last bin includes the overflow.

The uncertainty related to the finite size of the simulation samples (used to measure the efficiencies in Section 6) is also considered as a systematic uncertainty.

The uncertainty associated with the shape of the $B_s^0 \rightarrow \psi(2S)K_S^0K^-\pi^+$ contribution to the $\psi(2S)K_S^0\pi^+\pi^-$ invariant mass distribution is estimated by varying the shape parameters within their uncertainties. The largest deviation of $N(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)$ from the base-

line value is 0.5% which is taken as a systematic uncertainty.

As discussed in Section 5, the simulation for the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay does not take into account the intermediate resonance structure, leading to a significant disagreement between data and simulation in the 2- and 3-body mass distributions. This results in a potential bias in the efficiency reported in Section 6. To estimate the corresponding systematic uncertainty, the simulated sample is reweighted to be consistent with the data, and the difference between the baseline efficiency and the efficiency obtained on the weighted sample is taken as a systematic uncertainty. Due to the limited number of events, it is impossible to assign weights taking the ratio of data to simulation in bins of multi-dimensional phase space of the considered 4-body decay. An iterative procedure has been developed that operates with one-dimensional weights corresponding to each 2- and 3-body invariant mass, gradually making the mass distributions on weighted simulation sample closer and closer to data, until a satisfactory agreement in all intermediate invariant mass distributions is achieved. The distributions of invariant masses obtained with the weighted simulation sample are presented in Figs. 2 and 3. The efficiency obtained on the weighted simulation sample deviates from the baseline value by 5%, which is taken as a systematic uncertainty due to the intermediate resonance structure. This efficiency correction procedure with iterative reweighting is verified using a dedicated simulation sample instead of data, in which the contributions from $K^*(892)^\pm$ and $\rho(770)^0$ resonances are included with arbitrary magnitudes.

All uncertainties described above, excluding the one related to f_s/f_d for the ratio R_s , are summarized in Table 1 together with a total systematic uncertainty, calculated as a sum in quadrature of the individual sources.

Table 1: Systematic uncertainties (in %) of the measured branching fraction ratios.

Source	R_s	$R_{\pi^+\pi^-}$
Background model	2.5	0.8
Signal model	1.5	0.8
Shape of $B_s^0 \rightarrow \psi(2S)K_S^0K^-\pi^+$ contribution	—	0.5
Finite size of simulation samples	1.3	1.1
Intermediate resonances	—	5.0
Tracking efficiency	—	4.2
Total	3.2	6.7

A measurement of the ratio of the B_s^0 and B^0 fragmentation fractions, f_s/f_d , in proton-proton collisions at the LHC has been recently reported by the LHCb Collaboration [51]: $f_s/f_d = (0.263 \pm 0.008) - (17.6 \pm 2.1) \times 10^{-4} p_T(B)$, where $p_T(B)$ in GeV is the transverse momentum of a B meson produced in 13 TeV proton-proton collisions. The ratio was found to be independent of the rapidity of the B meson, but with a significant dependence on the transverse momentum of the B candidate. The $p_T(B)$ distribution used in this analysis is shown in Fig. 4, where the background is subtracted using s Plot. Using the LHCb result and the average $p_T(B)$ in our events of 31.2 GeV, the f_s/f_d value for the kinematic range of this analysis is obtained to be $f_s/f_d = 0.208 \pm 0.007$. The LHCb f_s/f_d measurement is mostly dependent on the events with $p_T < 20$ GeV, while the majority of the events in this analysis have $p_T(B) > 20$ GeV. Therefore, we assign an additional systematic uncertainty on f_s/f_d as the difference between 0.208 and the value obtained under the assumption that f_s/f_d becomes constant (0.2278) in the region $p_T > 20$ GeV. This additional uncertainty is estimated to be 0.020, and the total uncertainty on f_s/f_d is obtained by summing it in quadrature with the uncertainty of 0.007 obtained above. The resulting fragmentation fraction ratio used in the R_s measurement is $f_s/f_d = 0.208 \pm 0.021$,

with a relative uncertainty of 10%.

8 Measured branching fraction ratios

The branching fraction ratio of the $B_s^0 \rightarrow \psi(2S)K_S^0$ decay relative to the $B^0 \rightarrow \psi(2S)K_S^0$ one is measured using Eq. (1) to be

$$R_s = \frac{\mathcal{B}(B_s^0 \rightarrow \psi(2S)K_S^0)}{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0)} = (3.33 \pm 0.69 \text{ (stat)} \pm 0.11 \text{ (syst)} \pm 0.34 (f_s/f_d)) \times 10^{-2},$$

where the last uncertainty is related to the used value $f_s/f_d = 0.208 \pm 0.021$. Since the knowledge of f_s/f_d at large p_T can be updated with future measurements, allowing to improve the R_s evaluation, we also provide the measurement of the product

$$R_s \frac{f_s}{f_d} = \frac{f_s}{f_d} \frac{\mathcal{B}(B_s^0 \rightarrow \psi(2S)K_S^0)}{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0)} = (0.69 \pm 0.14 \text{ (stat)} \pm 0.02 \text{ (syst)}) \times 10^{-2}.$$

In addition, the transverse momentum distribution of the measured B candidates is presented in Fig. 4 and in the HEPData record for this analysis [30].

The branching fraction ratio of the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay with respect to the $B^0 \rightarrow \psi(2S)K_S^0$ one is measured to be

$$R_{\pi^+\pi^-} = \frac{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)}{\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0)} = 0.480 \pm 0.013 \text{ (stat)} \pm 0.032 \text{ (syst)}.$$

9 Summary

The $B_s^0 \rightarrow \psi(2S)K_S^0$ and $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decays are observed using proton-proton collision data collected by the CMS experiment at 13 TeV with an integrated luminosity of 103 fb^{-1} . Their branching fractions are measured with respect to the $B^0 \rightarrow \psi(2S)K_S^0$ decay to be $\mathcal{B}(B_s^0 \rightarrow \psi(2S)K_S^0)/\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0) = (3.33 \pm 0.69 \text{ (stat)} \pm 0.11 \text{ (syst)} \pm 0.34 (f_s/f_d)) \times 10^{-2}$, and $\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-)/\mathcal{B}(B^0 \rightarrow \psi(2S)K_S^0) = 0.480 \pm 0.013 \text{ (stat)} \pm 0.032 \text{ (syst)}$, where the last uncertainty in the first ratio corresponds to the uncertainty in the ratio of production cross sections of B_s^0 and B^0 mesons. The 2- and 3-body invariant mass distributions of the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay products do not show significant exotic narrow structures in addition to the known light meson resonances. Further studies with more data will be needed to investigate more precisely the internal dynamics of the $B^0 \rightarrow \psi(2S)K_S^0\pi^+\pi^-$ decay, and to perform CP asymmetry measurements in the two observed decays in the future.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the

LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NK-FIA (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS, RFBR, and NRC KI (Russia); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR, and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 884104, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010; the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Lendület ("Momentum") Programme and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, the New National Excellence Program ÚNKP, the NKFI research grants 123842, 123959, 124845, 124850, 125105, 128713, 128786, and 129058 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Science and Higher Education and the National Science Center, contracts Opus 2014/15/B/ST2/03998 and 2015/19/B/ST2/02861 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; the Ministry of Science and Higher Education, projects no. 0723-2020-0041 and no. FSWW-2020-0008, and the Russian Foundation for Basic Research, project No.19-42-703014 (Russia); MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Stavros Niarchos Foundation (Greece); the Rachadapisek Sompot Fund for Postdoctoral Fellowship, Chulalongkorn University and the Chulalongkorn Academic into Its 2nd Century Project Advancement Project (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] Belle Collaboration, "Observation of a narrow charmoniumlike state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays", *Phys. Rev. Lett.* **91** (2003) 262001, doi:10.1103/PhysRevLett.91.262001, arXiv:hep-ex/0309032.

-
- [2] CDF Collaboration, “Evidence for a narrow near-threshold structure in the $J/\psi\phi$ mass spectrum in $B^+ \rightarrow J/\psi\phi K^+$ decays”, *Phys. Rev. Lett.* **102** (2009) 242002, doi:10.1103/PhysRevLett.102.242002, arXiv:0903.2229.
- [3] CDF Collaboration, “Observation of the $Y(4140)$ structure in the $J/\psi\phi$ mass spectrum in $B^\pm \rightarrow J/\psi\phi K^\pm$ decays”, *Mod. Phys. Lett. A* **32** (2017) 1750139, doi:10.1142/S0217732317501395, arXiv:1101.6058.
- [4] CMS Collaboration, “Observation of a peaking structure in the $J/\psi\phi$ mass spectrum from $B^\pm \rightarrow J/\psi\phi K^\pm$ decays”, *Phys. Lett. B* **734** (2014) 261, doi:10.1016/j.physletb.2014.05.055, arXiv:1309.6920.
- [5] LHCb Collaboration, “Amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays”, *Phys. Rev. D* **95** (2017) 012002, doi:10.1103/PhysRevD.95.012002, arXiv:1606.07898.
- [6] BaBar Collaboration, “Observation of a broad structure in the $\pi^+\pi^-J/\psi$ mass spectrum around $4.26 \text{ GeV}/c^2$ ”, *Phys. Rev. Lett.* **95** (2005) 142001, doi:10.1103/PhysRevLett.95.142001, arXiv:hep-ex/0506081.
- [7] BESIII Collaboration, “Precise measurement of the $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ cross section at center-of-mass energies from 3.77 to 4.60 GeV”, *Phys. Rev. Lett.* **118** (2017) 092001, doi:10.1103/PhysRevLett.118.092001, arXiv:1611.01317.
- [8] Belle Collaboration, “Observation of a resonancelike structure in the $\pi^\pm\psi'$ mass distribution in exclusive $B \rightarrow K\pi^\pm\psi'$ decays”, *Phys. Rev. Lett.* **100** (2008) 142001, doi:10.1103/PhysRevLett.100.142001, arXiv:0708.1790.
- [9] Belle Collaboration, “Dalitz analysis of $B \rightarrow K\pi^+\psi'$ decays and the $Z(4430)^+$ ”, *Phys. Rev. D* **80** (2009) 031104, doi:10.1103/PhysRevD.80.031104, arXiv:0905.2869.
- [10] Belle Collaboration, “Experimental constraints on the spin and parity of the $Z(4430)^+$ ”, *Phys. Rev. D* **88** (2013) 074026, doi:10.1103/PhysRevD.88.074026, arXiv:1306.4894.
- [11] LHCb Collaboration, “Observation of the resonant character of the $Z(4430)^-$ state”, *Phys. Rev. Lett.* **112** (2014) 222002, doi:10.1103/PhysRevLett.112.222002, arXiv:1404.1903.
- [12] A. Hosaka et al., “Exotic hadrons with heavy flavors: X, Y, Z, and related states”, *Prog. Theor. Exp. Phys.* **2016** (2016) 062C01, doi:10.1093/ptep/ptw045, arXiv:1603.09229.
- [13] Particle Data Group, P. A. Zyla et al., “Review of particle physics”, *Prog. Theor. Exp. Phys.* **2020** (2020) 083C01, doi:10.1093/ptep/ptaa104.
- [14] N. Cabibbo, “Unitary symmetry and leptonic decays”, *Phys. Rev. Lett.* **10** (1963) 531, doi:10.1103/PhysRevLett.10.531.
- [15] Belle Collaboration, “Observation of large CP violation in the neutral B meson system”, *Phys. Rev. Lett.* **87** (2001) 091802, doi:10.1103/PhysRevLett.87.091802, arXiv:hep-ex/0107061.
- [16] BaBar Collaboration, “Observation of CP violation in the B^0 meson system”, *Phys. Rev. Lett.* **87** (2001) 091801, doi:10.1103/PhysRevLett.87.091801, arXiv:hep-ex/0107013.

- [17] BaBar Collaboration, “Evidence for CP violation in $B^0 \rightarrow J/\psi\pi^0$ decays”, *Phys. Rev. Lett.* **101** (2008) 021801, doi:10.1103/PhysRevLett.101.021801, arXiv:0804.0896.
- [18] BaBar Collaboration, “Measurement of time-dependent CP asymmetry in $B^0 \rightarrow c\bar{c}K^{(*)0}$ decays”, *Phys. Rev. D* **79** (2009) 072009, doi:10.1103/PhysRevD.79.072009, arXiv:0902.1708.
- [19] Belle Collaboration, “Precise measurement of the CP violation parameter $\sin 2\phi_1$ in $B^0 \rightarrow (c\bar{c})K^0$ decays”, *Phys. Rev. Lett.* **108** (2012) 171802, doi:10.1103/PhysRevLett.108.171802, arXiv:1201.4643.
- [20] LHCb Collaboration, “Measurement of the CP -violating phase β in $B^0 \rightarrow J/\psi\pi^+\pi^-$ decays and limits on penguin effects”, *Phys. Lett. B* **742** (2015) 38, doi:10.1016/j.physletb.2015.01.008, arXiv:1411.1634.
- [21] LHCb Collaboration, “Measurement of CP violation in $B^0 \rightarrow J/\psi K_S^0$ decays”, *Phys. Rev. Lett.* **115** (2015) 031601, doi:10.1103/PhysRevLett.115.031601, arXiv:1503.07089.
- [22] LHCb Collaboration, “Measurement of the time-dependent CP asymmetries in $B_s^0 \rightarrow J/\psi K_S^0$ ”, *JHEP* **06** (2015) 131, doi:10.1007/JHEP06(2015)131, arXiv:1503.07055.
- [23] LHCb Collaboration, “Measurement of CP violation in $B^0 \rightarrow J/\psi K_S^0$ and $B^0 \rightarrow \psi(2S)K_S^0$ decays”, *JHEP* **11** (2017) 170, doi:10.1007/JHEP11(2017)170, arXiv:1709.03944.
- [24] Belle Collaboration, “Measurement of the branching fraction and time-dependent CP asymmetry for $B^0 \rightarrow J/\psi\pi^0$ decays”, *Phys. Rev. D* **98** (2018) 112008, doi:10.1103/PhysRevD.98.112008, arXiv:1810.01356.
- [25] LHCb Collaboration, “Updated measurement of time-dependent CP -violating observables in $B_s^0 \rightarrow J/\psi K^+K^-$ decays”, *Eur. Phys. J. C* **79** (2019) 706, doi:10.1140/epjc/s10052-019-7159-8, arXiv:1906.08356. [Erratum: doi:10.1140/epjc/s10052-020-7875-0].
- [26] CMS Collaboration, “Measurement of the CP -violating phase ϕ_s in the $B_s^0 \rightarrow J/\psi\phi(1020) \rightarrow \mu^+\mu^-K^+K^-$ channel in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **816** (2021) 136188, doi:10.1016/j.physletb.2021.136188, arXiv:2007.02434.
- [27] ATLAS Collaboration, “Measurement of the CP -violating phase ϕ_s in $B_s^0 \rightarrow J/\psi\phi$ decays in ATLAS at 13 TeV”, *Eur. Phys. J. C* **81** (2021) 342, doi:10.1140/epjc/s10052-021-09011-0, arXiv:2001.07115.
- [28] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [29] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [30] “HEPData record for this analysis”, 2021. doi:10.17182/hepdata.114370.
- [31] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.

-
- [32] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [33] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [34] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [35] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [36] D. J. Lange, “The EvtGen particle decay simulation package”, *Nucl. Instrum. Meth. A* **462** (2001) 152, doi:10.1016/S0168-9002(01)00089-4.
- [37] E. Barberio, B. van Eijk, and Z. Waş, “PHOTOS — a universal Monte Carlo for QED radiative corrections in decays”, *Comput. Phys. Commun.* **66** (1991) 115, doi:10.1016/0010-4655(91)90012-A.
- [38] E. Barberio and Z. Waş, “PHOTOS – a universal Monte Carlo for QED radiative corrections: version 2.0”, *Comput. Phys. Commun.* **79** (1994) 291, doi:10.1016/0010-4655(94)90074-4.
- [39] GEANT4 Collaboration, “GEANT4—A simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [40] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [41] CMS Collaboration, “CMS tracking performance results from early LHC operation”, *Eur. Phys. J. C* **70** (2010) 1165, doi:10.1140/epjc/s10052-010-1491-3, arXiv:1007.1988.
- [42] CMS Collaboration, “Search for the X(5568) state decaying into $B_s^0 \pi^\pm$ in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. Lett.* **120** (2018) 202005, doi:10.1103/PhysRevLett.120.202005, arXiv:1712.06144.
- [43] CMS Collaboration, “Study of the $B^+ \rightarrow J/\psi \bar{\Lambda} p$ decay in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **12** (2019) 100, doi:10.1007/JHEP12(2019)100, arXiv:1907.05461.
- [44] CMS Collaboration, “Observation of the $\Lambda_b^0 \rightarrow J/\psi \Lambda \phi$ decay in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **802** (2020) 135203, doi:10.1016/j.physletb.2020.135203, arXiv:1911.03789.
- [45] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [46] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].













- [47] S. S. Wilks, "The large-sample distribution of the likelihood ratio for testing composite hypotheses", *Annals Math. Statist.* **9** (1938) 60, doi:10.1214/aoms/1177732360.
- [48] M. Pivk and F. R. Le Diberder, " \mathcal{P} lot: a statistical tool to unfold data distributions", *Nucl. Instrum. Meth. A* **555** (2005) 356, doi:10.1016/j.nima.2005.08.106, arXiv:physics/0402083.
- [49] CMS Collaboration, "Tracking POG results for pion efficiency with the D^* meson using data from 2016 and 2017", CMS Detector Performance Note CMS-DP-2018-050, 2018.
- [50] S. Jackman, "Bayesian analysis for the social sciences". John Wiley & Sons, New Jersey, USA, 2009. doi:10.1002/9780470686621.
- [51] LHCb Collaboration, "Precise measurement of the f_s/f_d ratio of fragmentation fractions and of B_s^0 decay branching fractions", *Phys. Rev. D* **104** (2021) 032005, doi:10.1103/PhysRevD.104.032005, arXiv:2103.06810.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Tumasyan

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis, M. Dragicevic , A. Escalante Del Valle , R. Frühwirth¹, M. Jeitler¹ , N. Krammer, L. Lechner , D. Liko, I. Mikulec, P. Paulitsch, F.M. Pitters, J. Schieck¹ , R. Schöfbeck , D. Schwarz, S. Templ , W. Waltenberger , C.-E. Wulz¹ 











Institute for Nuclear Problems, Minsk, Belarus

V. Chekhovsky, A. Litomin, V. Makarenko 





Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish², E.A. De Wolf, T. Janssen , T. Kello³, A. Lelek , H. Rejeb Sfar, P. Van Mechelen , S. Van Putte, N. Van Remortel 

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , A. De Moor, M. Delcourt, H. El Faham , S. Lowette , S. Moortgat , A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier , W. Van Doninck, D. Vannerom 













Université Libre de Bruxelles, Bruxelles, Belgium

D. Beghin, B. Bilin , B. Clerbaux , G. De Lentdecker, L. Favart , A.K. Kalsi , K. Lee, M. Mahdavihorrami, I. Makarenko , L. Moureaux , S. Paredes , L. Pétré, A. Popov , N. Postiau, E. Starling , L. Thomas , M. Vanden Bemden, C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

T. Cornelis , D. Dobur, J. Knolle , L. Lambrecht, G. Mestdach, M. Niedziela , C. Rendón, C. Roskas, A. Samalan, K. Skovpen , M. Tytgat , B. Vermassen, L. Wezenbeek














Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke, A. Bethani , G. Bruno, F. Bury , C. Caputo , P. David , C. Delaere , I.S. Donertas , A. Giammanco , K. Jaffel, Sa. Jain , V. Lemaitre, K. Mondal , J. Prisciandaro, A. Taliercio, M. Teklishyn , T.T. Tran, P. Vischia , S. Wertz 

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , C. Hensel, A. Moraes , P. Rebello Teles 


Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho, H. Brandao Malbouisson, W. Carvalho , J. Chinellato⁴, E.M. Da Costa , G.G. Da Silveira⁵ , D. De Jesus Damiao , V. Dos Santos Sousa, S. Fonseca De Souza , C. Mora Herrera , K. Mota Amarilo, L. Mundim , H. Nogima, A. Santoro, S.M. Silva Do Amaral , A. Sznajder , M. Thiel, F. Torres Da Silva De Araujo⁶ , A. Vilela Pereira 

Universidade Estadual Paulista (a), Universidade Federal do ABC (b), São Paulo, Brazil

C.A. Bernardes⁵ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , D.S. Lemos , P.G. Mercadante , S.F. Novaes , Sandra S. Padula 


Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov, G. Antchev , R. Hadjiiska, P. Iaydjiev, M. Misheva, M. Rodozov, M. Shopova, G. Sultanov





University of Sofia, Sofia, Bulgaria

A. Dimitrov, T. Ivanov, L. Litov , B. Pavlov, P. Petkov, A. Petrov













Beihang University, Beijing, China

T. Cheng , T. Javaid⁷, M. Mittal, L. Yuan

Department of Physics, Tsinghua University, Beijing, China

M. Ahmad , G. Bauer, C. Dozen⁸ , Z. Hu , J. Martins⁹ , Y. Wang, K. Yi^{10,11}


Institute of High Energy Physics, Beijing, China

E. Chapon , G.M. Chen⁷ , H.S. Chen⁷ , M. Chen , F. Iemmi, A. Kapoor , D. Leggat, H. Liao, Z.-A. Liu⁷ , V. Milosevic , F. Monti , R. Sharma , J. Tao , J. Thomas-Wilsker, J. Wang , H. Zhang , J. Zhao 



State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos, Y. An, Y. Ban, C. Chen, A. Levin , Q. Li , X. Lyu, Y. Mao, S.J. Qian, D. Wang , J. Xiao, H. Yang

Sun Yat-Sen University, Guangzhou, China

M. Lu, Z. You 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao³, H. Okawa , Y. Zhang 


Zhejiang University, Hangzhou, China, Zhejiang, China

Z. Lin , M. Xiao 

Universidad de Los Andes, Bogota, Colombia

C. Avila , A. Cabrera , C. Florez , J. Fraga


Universidad de Antioquia, Medellin, Colombia

J. Mejia Guisao, F. Ramirez, J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic, N. Godinovic , D. Lelas , I. Puljak 

University of Split, Faculty of Science, Split, Croatia

Z. Antunovic, M. Kovac, T. Sculac 


Institute Rudjer Boskovic, Zagreb, Croatia

V. Brigljevic , D. Ferencek , D. Majumder , M. Roguljic, A. Starodumov¹² , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou, G. Kole , M. Kolosova, S. Konstantinou, J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis, H. Rykaczewski, H. Saka 


Charles University, Prague, Czech Republic

M. Finger¹³, M. Finger Jr.¹³ , A. Kveton





















































Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala

Universidad San Francisco de Quito, Quito, Ecuador






E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egypt



































tian Network of High Energy Physics, Cairo, EgyptA.A. Abdelalim^{14,15} , E. Salama^{16,17}**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M.A. Mahmoud , Y. Mohammed **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**S. Bhowmik , R.K. Dewanjee , K. Ehataht, M. Kadastik, S. Nandan, C. Nielsen, J. Pata, M. Raidal , L. Tani, C. Veelken**Department of Physics, University of Helsinki, Helsinki, Finland**P. Eerola , H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar, E. Brücken , F. Garcia , J. Havukainen , M.S. Kim , R. Kinnunen, T. Lampén, K. Lassila-Perini , S. Lehti , T. Lindén, M. Lotti, L. Martikainen, M. Myllymäki, J. Ott , M.m. Rantanen, H. Siikonen, E. Tuominen , J. Tuominiemi**Lappeenranta University of Technology, Lappeenranta, Finland**P. Luukka , H. Petrow, T. Tuuva**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**C. Amendola , M. Besancon, F. Couderc , M. Dejardin, D. Denegri, J.L. Faure, F. Ferri , S. Ganjour, P. Gras, G. Hamel de Monchenault , P. Jarry, B. Lenzi , J. Malcles, J. Rander, A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁸, M. Titov , G.B. Yu **Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**S. Ahuja , F. Beaudette , M. Bonanomi , A. Buchot Perraguin, P. Busson, A. Cappati, C. Charlot, O. Davignon, B. Diab, G. Falmagne , S. Ghosh, R. Granier de Cassagnac , A. Hakimi, I. Kucher , J. Motta, M. Nguyen , C. Ochando , P. Paganini , J. Rembser, R. Salerno , U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini, A. Zabi, A. Zghiche **Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**J.-L. Agram¹⁹ , J. Andrea, D. Apparù, D. Bloch , G. Bourgatte, J.-M. Brom, E.C. Chabert, C. Collard , D. Darej, J.-C. Fontaine¹⁹, U. Goerlach, C. Grimault, A.-C. Le Bihan, E. Nibigira , P. Van Hove **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**E. Asilar , S. Beauceron , C. Bernet , G. Boudoul, C. Camen, A. Carle, N. Chanon , D. Contardo, P. Depasse , H. El Mamouni, J. Fay, S. Gascon , M. Gouzevitch , B. Ille, I.B. Laktineh, H. Lattaud , A. Lesauvage , M. Lethuillier , L. Mirabito, S. Perries, K. Shchablo, V. Sordini , L. Torterotot , G. Touquet, M. Vander Donckt, S. Viret**Georgian Technical University, Tbilisi, Georgia**I. Lomidze, T. Toriashvili²⁰, Z. Tsamalaidze¹³**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**V. Botta, L. Feld , K. Klein, M. Lipinski, D. Meuser, A. Pauls, N. Röwert, J. Schulz, M. Teroerde **RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**A. Dodonova, D. Eliseev, M. Erdmann , P. Fackeldey , B. Fischer, T. Hebbeker , K. Hoepfner, F. Ivone, L. Mastrolorenzo, M. Merschmeyer , A. Meyer , G. Mocellin, S. Mondal, S. Mukherjee , D. Noll , A. Novak, A. Pozdnyakov , Y. Rath, H. Reithler,

A. Schmidt , S.C. Schuler, A. Sharma , L. Vigilante, S. Wiedenbeck, S. Zaleski





















RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok, G. Flügge, W. Haj Ahmad²¹ , O. Hlushchenko, T. Kress, A. Nowack , O. Pooth, D. Roy , A. Stahl²² , T. Ziemons , A. Zotz














Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen, M. Aldaya Martin, P. Asmuss, S. Baxter, M. Bayatmakou, O. Behnke, A. Bermúdez Martínez, S. Bhattacharya, A.A. Bin Anuar , F. Blekman , K. Borras²³, D. Brunner, A. Campbell , A. Cardini , C. Cheng, F. Colombina, S. Consuegra Rodríguez , G. Correia Silva, M. De Silva, L. Didukh, G. Eckerlin, D. Eckstein, L.I. Estevez Banos , O. Filatov , E. Gallo²⁴, A. Geiser, A. Giraldi, A. Grohsjean , M. Guthoff, A. Jafari²⁵ , N.Z. Jomhari , H. Jung , A. Kasem²³ , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , D. Krücker , W. Lange, K. Lipka, W. Lohmann²⁶, R. Mankel, I.-A. Melzer-Pellmann , M. Mendizabal Morentin, J. Metwally, A.B. Meyer , M. Meyer , J. Mnich , A. Mussgiller, A. Nürnberg, Y. Otariid, D. Pérez Adán , D. Pitzl, A. Raspereza, B. Ribeiro Lopes, J. Rübenach, A. Saggio , A. Saibel , M. Savitskyi , M. Scham²⁷, V. Scheurer, S. Schnake, P. Schütze, C. Schwanenberger²⁴ , M. Shchedrolosiev, R.E. Sosa Ricardo , D. Stafford, N. Tonon , M. Van De Klundert , F. Vazzoler , R. Walsh , D. Walter, Q. Wang , Y. Wen , K. Wichmann, L. Wiens, C. Wissing, S. Wuchterl 

University of Hamburg, Hamburg, Germany

R. Aggleton, S. Albrecht , S. Bein , L. Benato , P. Connor , K. De Leo , M. Eich, K. El Morabit, F. Feindt, A. Fröhlich, C. Garbers , E. Garutti , P. Gunnellini, M. Hajheidari, J. Haller , A. Hinzmann , G. Kasieczka, R. Klanner , T. Kramer, V. Kutzner, J. Lange , T. Lange , A. Lobanov , A. Malara , C. Matthies, A. Mehta , A. Nigamova, K.J. Pena Rodriguez, M. Rieger , O. Rieger, P. Schleper, M. Schröder , J. Schwandt , J. Sonneveld , H. Stadie, G. Steinbrück, A. Tews, I. Zoi 





Karlsruher Institut fuer Technologie, Karlsruhe, Germany

J. Bechtel , S. Brommer, M. Burkart, E. Butz , R. Caspart , T. Chwalek, W. De Boer[†], A. Dierlamm, A. Droll, N. Faltermann , M. Giffels, J.O. Gosewisch, A. Gottmann, F. Hartmann²² , C. Heidecker, U. Husemann , P. Keicher, R. Koppenhöfer, S. Maier, S. Mitra , Th. Müller, M. Neukum, G. Quast , K. Rabbertz , J. Rauser, D. Savoia , M. Schnepf, D. Seith, I. Shvetsov, H.J. Simonis, R. Ulrich , J. Van Der Linden, R.F. Von Cube, M. Wassmer, M. Weber , S. Wieland, R. Wolf , S. Wozniewski, S. Wunsch


Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis, A. Kyriakis, D. Loukas, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

M. Diamantopoulou, D. Karasavvas, P. Kontaxakis , C.K. Koraka, A. Manousakis-Katsikakis, A. Panagiotou, I. Papavergou, N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis, E. Vourliotis

National Technical University of Athens, Athens, Greece

G. Bakas, K. Kousouris , I. Papakrivopoulos, G. Tsipolitis, A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece






K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios, P. Katsoulis, P. Kokkas, N. Manthos, I. Papadopoulos , J. Strogas 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University,

Budapest, Hungary

M. Csanad , K. Farkas, M.M.A. Gadallah²⁸ , S. Lökös²⁹ , P. Major, K. Mandal , G. Pasztor , A.J. Rádl, O. Surányi, G.I. Veres 


Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók³⁰ , G. Bencze, C. Hajdu , D. Horvath^{31,32} , F. Sikler , V. Veszpremi 


Institute of Nuclear Research ATOMKI, Debrecen, Hungary

S. Czellar, D. Fasanella , F. Fienga , J. Karancsi³⁰ , J. Molnar, Z. Szillasi, D. Teyssier






Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, Z.L. Trocsanyi³³ , B. Ujvari³⁴











Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³⁵ , F. Nemes³⁵, T. Novak

National Institute of Science Education and Research, HBNI, Bhubaneswar, India

S. Bahinipati³⁶ , C. Kar , P. Mal, T. Mishra , V.K. Muraleedharan Nair Bindhu³⁷, A. Nayak³⁷ , P. Saha, N. Sur , S.K. Swain, D. Vats³⁷






Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁸ , R. Gupta, A. Kaur, H. Kaur, M. Kaur , P. Kumari , M. Meena, K. Sandeep , J.B. Singh³⁹ , A.K. Viridi 







University of Delhi, Delhi, India

A. Ahmed, A. Bhardwaj , B.C. Choudhary , M. Gola, S. Keshri , A. Kumar , M. Naimuddin , P. Priyanka , K. Ranjan, S. Saumya, A. Shah 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

M. Bharti⁴⁰, R. Bhattacharya, S. Bhattacharya , D. Bhowmik, S. Dutta, S. Dutta, B. Gomber⁴¹ , M. Maity⁴², P. Palit , P.K. Rout , G. Saha, B. Sahu , S. Sarkar, M. Sharan

Indian Institute of Technology Madras, Madras, India

P.K. Behera , S.C. Behera, P. Kalbhor , J.R. Komaragiri⁴³ , D. Kumar⁴³, A. Muhammad, L. Panwar⁴³ , R. Pradhan, P.R. Pujahari, A. Sharma , A.K. Sikdar, P.C. Tiwari⁴³ 



Bhabha Atomic Research Centre, Mumbai, India

K. Naskar⁴⁴





Tata Institute of Fundamental Research-A, Mumbai, India

T. Aziz, S. Dugad, M. Kumar

Tata Institute of Fundamental Research-B, Mumbai, India

S. Banerjee , R. Chudasama, M. Guchait, S. Karmakar, S. Kumar, G. Majumder, K. Mazumdar, S. Mukherjee 




Indian Institute of Science Education and Research (IISER), Pune, India

A. Alpana, S. Dube , B. Kansal, A. Laha, S. Pandey , A. Rastogi , S. Sharma 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi^{45,46} , E. Khazaie⁴⁶, M. Zeinali⁴⁷



























Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani⁴⁸, S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi 

























University College Dublin, Dublin, Ireland

M. Grunewald 






INFN Sezione di Bari ^a, Bari, Italy, Università di Bari ^b, Bari, Italy, Politecnico di Bari ^c, Bari, Italy

M. Abbrescia^{a,b} , R. Aly^{a,b,49} , C. Aruta^{a,b}, A. Colaleo^a , D. Creanza^{a,c} , N. De Filippis^{a,c} , M. De Palma^{a,b} , A. Di Florio^{a,b}, A. Di Pilato^{a,b} , W. Elmetenawee^{a,b} , F. Errico^{a,b} , L. Fiore^a , G. Iaselli^{a,c} , M. Ince^{a,b} , S. Lezki^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^{a,b}, V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b}, A. Pompili^{a,b} , G. Pugliese^{a,c} , D. Ramos^a, A. Ranieri^a , G. Selvaggi^{a,b} , L. Silvestris^a , F.M. Simone^{a,b} , Ü. Sözbilir^a, R. Venditti^a , P. Verwilligen^a 













INFN Sezione di Bologna ^a, Bologna, Italy, Università di Bologna ^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , L. Borgonovi^a, L. Brigliadori^a, R. Campanini^{a,b} , P. Capiluppi^{a,b} , A. Castro^{a,b} , F.R. Cavallo^a , C. Ciocca^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b}, S. Lo Meo^{a,50}, L. Lunerti^{a,b}, S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania ^a, Catania, Italy, Università di Catania ^b, Catania, Italy

S. Albergò^{a,b,51} , S. Costa^{a,b,51} , A. Di Mattia^a , R. Potenza^{a,b}, A. Tricomi^{a,b,51} , C. Tuve^{a,b} 

INFN Sezione di Firenze ^a, Firenze, Italy, Università di Firenze ^b, Firenze, Italy

G. Barbagli^a , A. Cassese^a , R. Ceccarelli^{a,b}, V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^{a,b}, M. Meschini^a , S. Paoletti^a , R. Seidita^{a,b}, G. Sguazzoni^a , L. Viliani^a 





















INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , D. Piccolo 











INFN Sezione di Genova ^a, Genova, Italy, Università di Genova ^b, Genova, Italy

M. Bozzo^{a,b} , F. Ferro^a , R. Mulargia^a, E. Robutti^a , S. Tosi^{a,b} 


















INFN Sezione di Milano-Bicocca ^a, Milano, Italy, Università di Milano-Bicocca ^b, Milano, Italy








A. Benaglia^a , G. Boldrini , F. Brivio^{a,b}, F. Cetorelli^{a,b}, F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a, S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , D. Pedrini^a , B.S. Pinolini, S. Ragazzi^{a,b} , N. Redaelli^a , T. Tabarelli de Fatis^{a,b} , D. Valsecchi^{a,b,22}, D. Zuolo^{a,b} 

INFN Sezione di Napoli ^a, Napoli, Italy, Università di Napoli 'Federico II' ^b, Napoli, Italy, Università della Basilicata ^c, Potenza, Italy, Università G. Marconi ^d, Roma, Italy












S. Buontempo^a , F. Carnevali^{a,b}, N. Cavallo^{a,c} , A. De Iorio^{a,b} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,52} , S. Meola^{a,d,22} , P. Paolucci^{a,22} , B. Rossi^a , C. Sciacca^{a,b} 

INFN Sezione di Padova ^a, Padova, Italy, Università di Padova ^b, Padova, Italy, Università di Trento ^c, Trento, Italy
















P. Azzi^a , N. Bacchetta^a , D. Bisello^{a,b} , P. Bortignon^a , A. Bragagnolo^{a,b} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^a , U. Dosselli^a , F. Gasparini^{a,b} , U. Gasparini^{a,b} , G. Grosso, L. Layer^{a,53}, E. Lusiani , M. Margoni^{a,b} , F. Marini, A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b}, F. Simonetto^{a,b} 

G. Strong^a , M. Tosi^{a,b} , H. Yarar^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} ,
G. Zumerle^{a,b} 

































INFN Sezione di Pavia ^a, Pavia, Italy, Università di Pavia ^b, Pavia, Italy

C. Aimè^{a,b} , A. Braghieri^a , S. Calzaferri^{a,b} , D. Fiorina^{a,b} , P. Montagna^{a,b} , S.P. Ratti^{a,b} ,
V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^a , P. Vitulo^{a,b} 


















INFN Sezione di Perugia ^a, Perugia, Italy, Università di Perugia ^b, Perugia, Italy

P. Asenov^{a,54} , G.M. Bilei^a , D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^b ,
G. Mantovani^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,54} , A. Piccinelli^{a,b} ,
M. Presilla^{a,b} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 







































INFN Sezione di Pisa ^a, Pisa, Italy, Università di Pisa ^b, Pisa, Italy, Scuola Normale Superiore di Pisa ^c, Pisa, Italy, Università di Siena ^d, Siena, Italy

P. Azzurri^a , G. Bagliesi^a , V. Bertacchi^{a,c} , L. Bianchini^a , T. Boccali^a , E. Bossini^{a,b} ,
R. Castaldi^a , M.A. Ciocci^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , M.R. Di Domenico^{a,d} ,
S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , E. Manca^{a,c} , G. Mandorli^{a,c} ,
D. Matos Figueiredo, A. Messineo^{a,b} , M. Musich^a , F. Palla^a , S. Parolia^{a,b} , G. Ramirez-Sanchez^{a,c} ,
A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,c} , A. Scribano^a , N. Shafiei^{a,b} ,
P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , A. Venturi^a , P.G. Verdini^a 








INFN Sezione di Roma ^a, Rome, Italy, Sapienza Università di Roma ^b, Rome, Italy

P. Barria^a , M. Campana^{a,b} , F. Cavallari^a , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a ,
E. Longo^{a,b} , P. Meridiani^a , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} ,
C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a ,
R. Tramontano^{a,b} 









INFN Sezione di Torino ^a, Torino, Italy, Università di Torino ^b, Torino, Italy, Università del Piemonte Orientale ^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a ,
R. Bellan^{a,b} , A. Bellora^{a,b} , J. Berenguer Antequera^{a,b} , C. Biino^a , N. Cartiglia^a ,
M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a ,
C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , E. Migliore^{a,b} , E. Monteil^{a,b} , M. Monteno^a ,
M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Pelliccioni^a ,
M. Ruspa^{a,c} , K. Shchelina^a , F. Siviero^{a,b} , V. Sola^a , A. Solano^{a,b} , D. Soldi^{a,b} ,
A. Staiano^a , M. Tornago^{a,b} , D. Trocino^a , G. Umoret^{a,b} , A. Vagnerini^{a,b} 

INFN Sezione di Trieste ^a, Trieste, Italy, Università di Trieste ^b, Trieste, Italy

S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , A. Da Rold^{a,b} ,
G. Della Ricca^{a,b} , G. Sorrentino^{a,b} 




Kyungpook National University, Daegu, Korea

S. Dogra , C. Huh , B. Kim, D.H. Kim , G.N. Kim , J. Kim, J. Lee, S.W. Lee ,
C.S. Moon , Y.D. Oh , S.I. Pak, S. Sekmen , Y.C. Yang

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

B. Francois , T.J. Kim , J. Park 

Korea University, Seoul, Korea

S. Cho, S. Choi , B. Hong , K. Lee, K.S. Lee , J. Lim, J. Park, S.K. Park, J. Yoo





Kyung Hee University, Department of Physics, Seoul, Republic of Korea, Seoul, Korea

J. Goh , A. Gurtu



Sejong University, Seoul, Korea

H.S. Kim , Y. Kim

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi, S. Jeon, J. Kim, J.S. Kim, S. Ko, H. Kwon, H. Lee , S. Lee, B.H. Oh, M. Oh , S.B. Oh, H. Seo , U.K. Yang, I. Yoon 


University of Seoul, Seoul, Korea

W. Jang, D.Y. Kang, Y. Kang, S. Kim, B. Ko, J.S.H. Lee , Y. Lee, J.A. Merlin, I.C. Park, Y. Roh, M.S. Ryu, D. Song, I.J. Watson , S. Yang

Yonsei University, Department of Physics, Seoul, Korea

S. Ha, H.D. Yoo

Sungkyunkwan University, Suwon, Korea

M. Choi, H. Lee, Y. Lee, I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Egaila, Kuwait, Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi


Riga Technical University, Riga, Latvia

K. Dreimanis , V. Veckalns⁵⁵ 





Vilnius University, Vilnius, Lithuania

M. Ambrozas, A. Carvalho Antunes De Oliveira , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 





National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , Z. Zolkapli


Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello, J.A. Murillo Quijada , A. Sehrawat, L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala, H. Castilla-Valdez, E. De La Cruz-Burelo , I. Heredia-De La Cruz⁵⁶ , R. Lopez-Fernandez, C.A. Mondragon Herrera, D.A. Perez Navarro, R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

S. Carrillo Moreno, C. Oropeza Barrera , F. Vazquez Valencia


Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Pedraza, H.A. Salazar Ibarguen, C. Uribe Estrada

University of Montenegro, Podgorica, Montenegro

J. Mijuskovic⁵⁷, N. Raicevic




University of Auckland, Auckland, New Zealand

D. Krofcheck 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 


National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad, M.I. Asghar, A. Awais, M.I.M. Awan, M. Gul , H.R. Hoorani, W.A. Khan, M.A. Shah, M. Shoaib , M. Waqas 




AGH University of Science and Technology Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

V. Avati, L. Grzanka, M. Malawski

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska, M. Bluj , B. Boimska , M. Górski, M. Kazana, M. Szleper , P. Zalewski



Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski, K. Doroba, A. Kalinowski , M. Konecki , J. Krolikowski 




Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo, P. Bargassa , D. Bastos, A. Boletti , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad, M. Pisano, J. Seixas , O. Toldaiev , J. Varela 

Joint Institute for Nuclear Research, Dubna, Russia

S. Afanasiev, D. Budkouski, I. Golutvin, I. Gorbunov , V. Karjavine, V. Korenkov , A. Lanev, A. Malakhov, V. Matveev^{58,59}, V. Palichik, V. Perelygin, M. Savina, V. Shalaev, S. Shmatov, S. Shulha, V. Smirnov, O. Teryaev, N. Voytishin, B.S. Yuldashev⁶⁰, A. Zarubin, I. Zhizhin


Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia

G. Gavrilo , V. Golovtsov, Y. Ivanov, V. Kim⁶¹ , E. Kuznetsova⁶², V. Murzin, V. Oreshkin, I. Smirnov, D. Sosnov , V. Sulimov, L. Uvarov, S. Volkov, A. Vorobyev

Institute for Nuclear Research, Moscow, Russia

Yu. Andreev , A. Dermenev, S. Gninenko , N. Golubev, A. Karneyeu , D. Kirpichnikov , M. Kirsanov, N. Krasnikov, A. Pashenkov, G. Pivovarov , A. Toropin




Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of NRC 'Kurchatov Institute', Moscow, Russia

V. Epshteyn, V. Gavrilo, N. Lychkovskaya, A. Nikitenko⁶³, V. Popov, A. Stepenov, M. Toms, E. Vlasov , A. Zhokin


Moscow Institute of Physics and Technology, Moscow, Russia

T. Aushev

National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia

O. Bychkova, R. Chistov⁶⁴ , M. Danilov⁶⁴ , A. Oskin, P. Parygin, S. Polikarpov⁶⁴ , A. Tulupov


P.N. Lebedev Physical Institute, Moscow, Russia

V. Andreev, M. Azarkin, I. Dremin , M. Kirakosyan, A. Terkulov






Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia

A. Belyaev, E. Boos , M. Dubinin⁶⁵ , L. Dudko , A. Ershov, A. Gribushin, V. Klyukhin , O. Kodolova , I. Lokhtin , S. Obraztsov, S. Petrushanko, V. Savrin, A. Snigirev 

Novosibirsk State University (NSU), Novosibirsk, Russia

V. Blinov⁶⁶, T. Dimova⁶⁶, L. Kardapoltsev⁶⁶, A. Kozyrev⁶⁶, I. Ovtin⁶⁶, O. Radchenko⁶⁶, Y. Skovpen⁶⁶ 



**Institute for High Energy Physics of National Research Centre ‘Kurchatov Institute’,
Protvino, Russia**

I. Azhgirey , I. Bayshev, D. Elumakhov, V. Kachanov, D. Konstantinov , P. Mandrik ,
V. Petrov, R. Ryutin, S. Slabospitskii , A. Sobol, S. Troshin , N. Tyurin, A. Uzunian, A. Volkov

National Research Tomsk Polytechnic University, Tomsk, Russia

A. Babaev, V. Okhotnikov




















Tomsk State University, Tomsk, Russia

V. Borshch, V. Ivanchenko , E. Tcherniaev 

**University of Belgrade: Faculty of Physics and VINCA Institute of Nuclear Sciences,
Belgrade, Serbia**

P. Adzic⁶⁷ , M. Dordevic , P. Milenovic , J. Milosevic 

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT),
Madrid, Spain**

M. Aguilar-Benitez, J. Alcaraz Maestre , A. Álvarez Fernández, I. Bachiller, M. Barrio Luna,
Cristina F. Bedoya , C.A. Carrillo Montoya , M. Cepeda , M. Cerrada, N. Colino ,
B. De La Cruz, A. Delgado Peris , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gon-
zalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran,
Á. Navarro Tobar , C. Perez Dengra, A. Pérez-Calero Yzquierdo , J. Puerta Pelayo ,
I. Redondo , L. Romero, S. Sánchez Navas, L. Urda Gómez , C. Willmott














Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz

**Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de As-
turias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , C. Erice , J. Fernandez Menendez , S. Folgueras ,
I. Gonzalez Caballero , J.R. González Fernández, E. Palencia Cortezon , C. Ramón Álvarez,
V. Rodríguez Bouza , A. Soto Rodríguez, A. Trapote, N. Trevisani , C. Vico Villalba

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

J.A. Brochero Cifuentes , I.J. Cabrillo, A. Calderon , J. Duarte Campderros , M. Fernan-
dez , C. Fernandez Madrazo , P.J. Fernández Manteca , A. García Alonso, G. Gomez,
C. Martinez Rivero, P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas ,
J. Piedra Gomez , C. Prieels, A. Ruiz-Jimeno , L. Scodellaro , I. Vila, J.M. Vizán García 































University of Colombo, Colombo, Sri Lanka



























M.K. Jayananda, B. Kailasapathy⁶⁸, D.U.J. Sonnadara, D.D.C. Wickramarathna

University of Ruhuna, Department of Physics, Matara, Sri Lanka




W.G.D. Dharmaratna , K. Liyanage, N. Perera, N. Wickramage

CERN, European Organization for Nuclear Research, Geneva, Switzerland

















T.K. Aarrestad , D. Abbaneo, J. Alimena , E. Auffray, G. Auzinger, J. Baechler,
P. Baillon[†], D. Barney , J. Bendavid, M. Bianco , A. Bocci , C. Caillol, T. Camporesi,
M. Capeans Garrido , G. Cerminara, N. Chernyavskaya , S.S. Chhibra , S. Choudhury,
M. Cipriani , L. Cristella , D. d’Enterria , A. Dabrowski , A. David , A. De Roeck ,
M.M. Defranchis , M. Deile , M. Dobson, M. Dünser , N. Dupont, A. Elliott-Peisert,
F. Fallavollita⁶⁹, A. Florent , L. Forthomme , G. Franzoni , W. Funk, S. Ghosh , S. Giani,
D. Gigi, K. Gill, F. Glege, L. Gouskos , E. Govorkova , M. Haranko , J. Hegeman ,
V. Innocente , T. James, P. Janot , J. Kaspar , J. Kieseler , M. Komm , N. Kratochwil,

C. Lange , S. Laurila, P. Lecoq , A. Lintuluoto, K. Long , C. Lourenço , B. Maier, L. Malgeri , S. Mallios, M. Mannelli, A.C. Marini , F. Meijers, S. Mersi , E. Meschi , F. Moortgat , M. Mulders , S. Orfanelli, L. Orsini, F. Pantaleo , E. Perez, M. Peruzzi , A. Petrilli, G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo, M. Pitt , H. Qu , T. Quast, D. Rabady , A. Racz, G. Reales Gutiérrez, M. Rovere, H. Sakulin, J. Salfeld-Nebgen , S. Scarfi, C. Schwick, M. Selvaggi , A. Sharma, P. Silva , W. Snoeys , P. Sphicas⁷⁰ , S. Summers , K. Tatar , V.R. Tavolaro , D. Treille, P. Tropea, A. Tsiro, J. Wanczyk⁷¹, K.A. Wozniak, W.D. Zeuner














Paul Scherrer Institut, Villigen, Switzerland

L. Caminada⁷² , A. Ebrahimi , W. Erdmann, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, U. Langenegger, M. Missiroli⁷² , L. Noehte⁷², T. Rohe



ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

K. Androsov⁷¹ , M. Backhaus , P. Berger, A. Calandri , A. De Cosa, G. Dissertori , M. Dittmar, M. Donegà, C. Dorfer , F. Eble, K. Gedia, F. Glessgen, T.A. Gómez Espinosa , C. Grab , D. Hits, W. Lustermann, A.-M. Lyon, R.A. Manzoni , L. Marchese , C. Martin Perez, M.T. Meinhard, F. Nessi-Tedaldi, J. Niedziela , F. Pauss, V. Perovic, S. Pigazzini , M.G. Ratti , M. Reichmann, C. Reissel, T. Reitenspiess, B. Ristic , D. Ruini, D.A. Sanz Becerra , V. Stampf, J. Steggemann⁷¹ , R. Wallny 







Universität Zürich, Zurich, Switzerland

C. AMSLER⁷³ , P. Bärtschi, C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier, A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler, W. Jin, A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti, A. Macchiolo , P. Meiring, V.M. Mikuni , U. Molinatti, I. Neutelings, A. Reimers, P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger, Y. Takahashi 




National Central University, Chung-Li, Taiwan

C. Adloff⁷⁴, C.M. Kuo, W. Lin, A. Roy , T. Sarkar⁴² , S.S. Yu











National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao, K.F. Chen , P.H. Chen , P.s. Chen, H. Cheng , W.-S. Hou , Y.y. Li, R.-S. Lu, E. Paganis , A. Psallidas, A. Steen, H.y. Wu, E. Yazgan , P.r. Yu

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

B. Asavapibhop , C. Asawatangtrakuldee , N. Srimanobhas 





Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

F. Boran , S. Damarseckin⁷⁵, Z.S. Demiroglu , F. Dolek , I. Dumanoglu⁷⁶ , E. Eskut, Y. Guler⁷⁷ , E. Gurpinar Guler⁷⁷ , C. Isik, O. Kara, A. Kayis Topaksu, U. Kiminsu , G. Onengut, K. Ozdemir⁷⁸, A. Polatoz, A.E. Simsek , B. Tali⁷⁹, U.G. Tok , S. Turkcapar, I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Karapinar, K. Ocalan⁸⁰ , M. Yalvac⁸¹ 



Bogazici University, Istanbul, Turkey

B. Akgun, I.O. Atakisi , E. Gulmez , M. Kaya⁸² , O. Kaya⁸³, Ö. Özçelik, S. Tekten⁸⁴, E.A. Yetkin⁸⁵ 

Istanbul Technical University, Istanbul, Turkey


A. Cakir , K. Cankocak⁷⁶ , Y. Komurcu, S. Sen⁸⁶ 

Istanbul University, Istanbul, Turkey












S. Cerci⁷⁹, I. Hos⁸⁷, B. Isildak⁸⁸, B. Kaynak, S. Ozkorucuklu, H. Sert , C. Simsek, D. Sunar Cerci⁷⁹ , C. Zorbilmez

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkov, Ukraine







B. Grynyov

National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine
L. Levchuk 



















University of Bristol, Bristol, United Kingdom

D. Anthony, E. Bhal , S. Bologna, J.J. Brooke , A. Bundock , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , G.P. Heath, H.F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran, S. Seif El Nasr-Storey, V.J. Smith, N. Stylianou⁸⁹ , K. Walkingshaw Pass, R. White





Rutherford Appleton Laboratory, Didcot, United Kingdom

K.W. Bell, A. Belyaev⁹⁰ , C. Brew , R.M. Brown, D.J.A. Cockerill, C. Cooke, K.V. Ellis, K. Harder, S. Harper, M.-L. Holmberg⁹¹, J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt, T. Reis , T. Schuh, C.H. Shepherd-Themistocleous, I.R. Tomalin, T. Williams 

Imperial College, London, United Kingdom

R. Bainbridge , P. Bloch , S. Bonomally, J. Borg , S. Breeze, O. Buchmuller, V. Cepaitis , G.S. Chahal⁹² , D. Colling, P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi, G. Iles, J. Langford, L. Lyons, A.-M. Magnan, S. Malik, A. Martelli , D.G. Monk, J. Nash⁹³ , M. Pesaresi, B.C. Radburn-Smith, D.M. Raymond, A. Richards, A. Rose, E. Scott , C. Seez, A. Shtipliyski, A. Tapper , K. Uchida, T. Virdee²² , M. Vojinovic , N. Wardle , S.N. Webb , D. Winterbottom

Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid , L. Teodorescu, S. Zahid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , A.R. Kanuganti, B. McMaster , M. Saunders , S. Sawant, C. Sutantawibul, J. Wilson 











Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , R. Uniyal , A.M. Vargas Hernandez














The University of Alabama, Tuscaloosa, Alabama, USA

A. Buccilli , S.I. Cooper , D. Di Croce , S.V. Gleyzer , C. Henderson , C.U. Perez , P. Rumerio⁹⁴ , C. West 











Boston University, Boston, Massachusetts, USA

A. Akpinar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , E. Fontanesi, D. Gastler, S. May , J. Rohlf , K. Salyer , D. Sperka, D. Spitzbart , I. Suarez , A. Tsatsos, S. Yuan, D. Zou

Brown University, Providence, Rhode Island, USA

G. Benelli , B. Burkle , X. Coubez²³, D. Cutts , M. Hadley , U. Heintz , J.M. Hogan⁹⁵ , T. Kwon, G. Landsberg , K.T. Lau , D. Li, M. Lukasik, J. Luo , M. Narain, N. Pervan, S. Sagir⁹⁶ , F. Simpson, E. Usai , W.Y. Wong, X. Yan , D. Yu , W. Zhang







University of California, Davis, Davis, California, USA

J. Bonilla , C. Brainerd , R. Breedon, M. Calderon De La Barca Sanchez, M. Chertok , J. Conway , P.T. Cox, R. Erbacher, G. Haza, F. Jensen , O. Kukral, R. Lander, M. Mulhearn , D. Pellett, B. Regnery , D. Taylor , Y. Yao , F. Zhang 




University of California, Los Angeles, California, USA

M. Bachtis , R. Cousins , A. Datta , D. Hamilton, J. Hauser , M. Ignatenko, M.A. Iqbal, T. Lam, W.A. Nash, S. Regnard , D. Saltzberg , B. Stone, V. Valuev 











University of California, Riverside, Riverside, California, USA

Y. Chen, R. Clare , J.W. Gary , M. Gordon, G. Hanson , G. Karapostoli , O.R. Long , N. Manganeli, W. Si , S. Wimpenny, Y. Zhang













University of California, San Diego, La Jolla, California, USA

J.G. Branson, P. Chang , S. Cittolin, S. Cooperstein , D. Diaz , J. Duarte , R. Gerosa , L. Giannini , J. Guiang, R. Kansal , V. Krutelyov , R. Lee, J. Letts , M. Masciovecchio , F. Mokhtar, M. Pieri , B.V. Sathia Narayanan , V. Sharma , M. Tadel, F. Würthwein , Y. Xiang , A. Yagil 







University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

N. Amin, C. Campagnari , M. Citron , G. Collura , A. Dorsett, V. Dutta , J. Incandela , M. Kilpatrick , J. Kim , B. Marsh, H. Mei, M. Oshiro, M. Quinnan , J. Richman, U. Sarica , F. Setti, J. Sheplock, P. Siddireddy, D. Stuart, S. Wang 


California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, I. Dutta , J.M. Lawhorn , N. Lu , J. Mao, H.B. Newman , T.Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , Z. Zhang , R.Y. Zhu 















Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M.B. Andrews, P. Bryant , T. Ferguson , A. Harilal, C. Liu, T. Mudholkar , M. Paulini , A. Sanchez, W. Terrill












































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hassani, G. Karathanasis, E. MacDonald, R. Patel, A. Perloff , C. Savard, N. Schonbeck, K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper











Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , Y. Cheng , D.J. Cranshaw , S. Hogan, J. Monroy , J.R. Patterson , D. Quach , J. Reichert , M. Reid , A. Ryd, W. Sun , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , G. Apollinari, A. Apresyan , A. Apyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat, K. Burkett , J.N. Butler, A. Canepa, G.B. Cerati , H.W.K. Cheung , F. Chlebana, K.F. Di Petrillo , J. Dickinson , V.D. Elvira , Y. Feng, J. Freeman, Z. Gece, L. Gray, D. Green, S. Grünendahl , O. Gutsche , R.M. Harris , R. Heller, T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani, M. Johnson, U. Joshi, T. Klijnsma , B. Klima , K.H.M. Kwok, S. Lammel , D. Lincoln , R. Lipton, T. Liu, C. Madrid, K. Maeshima, C. Mantilla , D. Mason, P. McBride , P. Merkel, S. Mrenna , S. Nahn , J. Ngadiuba , V. Papadimitriou, N. Pastika, K. Pedro , C. Pena⁶⁵ , F. Ravera , A. Reinsvold Hall⁹⁷ , L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel, S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk, N.V. Tran , L. Uplegger , E.W. Vaandering , H.A. Weber 







University of Florida, Gainesville, Florida, USA

P. Avery, D. Bourilkov , L. Cadamuro , V. Cherepanov, R.D. Field, D. Guerrero, M. Kim, E. Koenig, J. Konigsberg , A. Korytov, K.H. Lo, K. Matchev , N. Menendez , G. Mitselmakher , A. Muthirakalayil Madhu, N. Rawal, D. Rosenzweig, S. Rosenzweig, K. Shi , J. Wang , Z. Wu , E. Yigitbasi , X. Zuo












Florida State University, Tallahassee, Florida, USA

T. Adams , A. Askew , R. Habibullah , V. Hagopian, K.F. Johnson, R. Khurana, T. Kolberg , G. Martinez, H. Prosper , C. Schiber, O. Viazlo , R. Yohay , J. Zhang





Florida Institute of Technology, Melbourne, Florida, USA

M.M. Baarmand , S. Butalla, T. Elkafrawy¹⁷ , M. Hohlmann , R. Kumar Verma , D. Noonan , M. Rahmani, F. Yumiceva 








University of Illinois at Chicago (UIC), Chicago, Illinois, USA

M.R. Adams, H. Becerril Gonzalez , R. Cavanaugh , S. Dittmer, O. Evdokimov , C.E. Gerber , D.J. Hofman , A.H. Merrit, C. Mills , G. Oh , T. Roy, S. Rudrabhatla, M.B. Tonjes , N. Varelas , J. Viinikainen , X. Wang, Z. Ye 















The University of Iowa, Iowa City, Iowa, USA

M. Alhousseini , K. Dilsiz⁹⁸ , L. Emediato, R.P. Gandrajula , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili⁹⁹, J. Nachtman, H. Ogul¹⁰⁰ , Y. Onel , A. Penzo, C. Snyder, E. Tiras¹⁰¹ 




Johns Hopkins University, Baltimore, Maryland, USA

O. Amram , B. Blumenfeld , L. Corcodilos , J. Davis, A.V. Gritsan , S. Kyriacou, P. Maksimovic , J. Roskes , M. Swartz, T.Á. Vámi 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu, J. Anguiano, C. Baldenegro Barrera , P. Baringer , A. Bean , Z. Flowers, T. Isidori, S. Khalil , J. King, G. Krintiras , A. Kropivnitskaya , M. Lazarovits, C. Le Mahieu, C. Lindsey, J. Marquez, N. Minafra , M. Murray , M. Nickel, C. Rogan , C. Royon, R. Salvatico , S. Sanders, E. Schmitz, C. Smith , Q. Wang , Z. Warner, J. Williams , G. Wilson 









Kansas State University, Manhattan, Kansas, USA

S. Duric, A. Ivanov , K. Kaadze , D. Kim, Y. Maravin , T. Mitchell, A. Modak, K. Nam
















Lawrence Livermore National Laboratory, Livermore, California, USA

F. Rebassoo, D. Wright








University of Maryland, College Park, Maryland, USA

E. Adams, A. Baden, O. Baron, A. Belloni , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg, T. Koeth, Y. Lai, S. Lascio, A.C. Mignerey, S. Nabili, C. Palmer , M. Seidel , A. Skuja , L. Wang, K. Wong 






Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

D. Abercrombie, G. Andreassi, R. Bi, W. Busza , I.A. Cali, Y. Chen , M. D'Alfonso , J. Eysermans, C. Freer , G. Gomez Ceballos, M. Goncharov, P. Harris, M. Hu, M. Klute , D. Kovalskyi , J. Krupa, Y.-J. Lee , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland, Z. Shi , G.S.F. Stephans , J. Wang, Z. Wang , B. Wyslouch 

University of Minnesota, Minneapolis, Minnesota, USA

R.M. Chatterjee, A. Evans , J. Hiltbrand, Sh. Jain , B.M. Joshi , M. Krohn, Y. Kubota, J. Mans , M. Revering, R. Rusack , R. Saradhy, N. Schroeder , N. Strobbe , M.A. Wadud











University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , M. Bryson, S. Chauhan , D.R. Claes, C. Fangmeier, L. Finco , F. Golf , C. Joo, I. Kravchenko , I. Reed, J.E. Siado, G.R. Snow[†], W. Tabb, A. Wightman, F. Yan, A.G. Zecchinelli






State University of New York at Buffalo, Buffalo, New York, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava, C. McLean , D. Nguyen, J. Pekkanen , S. Rappoccio , A. Williams 











Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis, Y. Haddad , Y. Han, A. Hortiangtham, A. Krishna, J. Li , J. Lidrych , G. Madigan, B. Marzocchi , D.M. Morse , V. Nguyen, T. Orimoto , A. Parker, L. Skinnari , A. Tishelman-Charny, T. Wamorkar, B. Wang , A. Wisecarver, D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Bhattacharya , J. Bueghly, Z. Chen , A. Gilbert , T. Gunter , K.A. Hahn, Y. Liu, N. Odell, M.H. Schmitt , M. Velasco














University of Notre Dame, Notre Dame, Indiana, USA

R. Band , R. Bucci, M. Cremonesi, A. Das , N. Dev , R. Goldouzian , M. Hildreth, K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence, N. Loukas , D. Lutton, J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady, K. Mohrman, C. Moore, Y. Musienko⁵⁸, R. Ruchti, A. Townsend, M. Wayne, M. Zarucki , L. Zygala

The Ohio State University, Columbus, Ohio, USA

B. Bylsma, L.S. Durkin , B. Francis , C. Hill , M. Nunez Ornelas , K. Wei, B.L. Winer, B.R. Yates 







Princeton University, Princeton, New Jersey, USA

F.M. Addesa , B. Bonham , P. Das , G. Dezoort, P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich, S. Higginbotham, A. Kalogeropoulos , G. Kopp, S. Kwan , D. Lange, D. Marlow , K. Mei , I. Ojalvo, J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , S. Norberg












Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi, V.E. Barnes , R. Chawla , S. Das , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy, M. Liu, G. Negro, N. Neumeister , G. Paspalaki, S. Piperov , A. Purohit, J.F. Schulte , M. Stojanovic¹⁸, J. Thieman , F. Wang , R. Xiao , W. Xie 








Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar

Rice University, Houston, Texas, USA

D. Acosta , A. Baty , T. Carnahan, M. Decaro, S. Dildick , K.M. Ecklund , S. Freed, P. Gardner, F.J.M. Geurts , A. Kumar , W. Li, B.P. Padley , R. Redjimi, J. Rotter, W. Shi , A.G. Stahl Leiton , S. Yang , L. Zhang¹⁰², Y. Zhang 












University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro, R. Demina , J.L. Dulemba , C. Fallon, T. Ferbel , M. Galanti, A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili, E. Ranken, R. Taus, G.P. Van Onsem 

The Rockefeller University, New York, New York, USA

K. Goulios








Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , A. Gandrakota , Y. Gershtein , E. Halkiadakis , A. Hart, M. Heindl , O. Karacheban²⁶ , I. Laflotte, A. Lath , R. Montalvo, K. Nash, M. Osherson, S. Salur , S. Schnetzer, S. Somalwar , R. Stone, S.A. Thayil , S. Thomas, H. Wang 




University of Tennessee, Knoxville, Tennessee, USA

H. Acharya, A.G. Delannoy , S. Fiorendi , S. Spanier 







Texas A&M University, College Station, Texas, USA

O. Bouhali¹⁰³ , M. Dalchenko , A. Delgado , R. Eusebi, J. Gilmore, T. Huang, T. Kamon¹⁰⁴, H. Kim , S. Luo , S. Malhotra, R. Mueller, D. Overton, D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin, J. Damgov, V. Hegde, K. Lamichhane, S.W. Lee , T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev, Z. Wang, A. Whitbeck


Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , S. Greene, A. Gurrola , W. Johns, A. Melo, K. Padeken , F. Romeo , P. Sheldon , S. Tuo, J. Velkovska 











University of Virginia, Charlottesville, Virginia, USA

M.W. Arenton , B. Cardwell, B. Cox , G. Cummings , J. Hakala , R. Hirosky , M. Joyce , A. Ledovskoy , A. Li, C. Neu , C.E. Perez Lara , B. Tannenwald , S. White 

Wayne State University, Detroit, Michigan, USA

N. Poudyal 

University of Wisconsin - Madison, Madison, WI, Wisconsin, USA

S. Banerjee, K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He, M. Herndon , A. Herve, U. Hussain, A. Lanaro, A. Loeliger, R. Loveless, J. Madhusudanan Sreekala , A. Mallampalli, A. Mohammadi, D. Pinna, A. Savin, V. Shang, V. Sharma , W.H. Smith , D. Teague, S. Trembath-Reichert, W. Vetens 

†: Deceased

1: Also at TU Wien, Wien, Austria

2: Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

3: Also at Université Libre de Bruxelles, Bruxelles, Belgium

4: Also at Universidade Estadual de Campinas, Campinas, Brazil

5: Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

6: Also at The University of the State of Amazonas, Manaus, Brazil

7: Also at University of Chinese Academy of Sciences, Beijing, China

8: Also at Department of Physics, Tsinghua University, Beijing, China

9: Also at UFMS, Nova Andradina, Brazil

10: Also at Nanjing Normal University Department of Physics, Nanjing, China

11: Now at The University of Iowa, Iowa City, Iowa, USA

12: Also at Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of NRC 'Kurchatov Institute', Moscow, Russia

13: Also at Joint Institute for Nuclear Research, Dubna, Russia

14: Also at Helwan University, Cairo, Egypt

15: Now at Zewail City of Science and Technology, Zewail, Egypt

16: Also at British University in Egypt, Cairo, Egypt

17: Now at Ain Shams University, Cairo, Egypt

-
- 18: Also at Purdue University, West Lafayette, Indiana, USA
 - 19: Also at Université de Haute Alsace, Mulhouse, France
 - 20: Also at Tbilisi State University, Tbilisi, Georgia
 - 21: Also at Erzincan Binali Yildirim University, Erzincan, Turkey
 - 22: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
 - 23: Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
 - 24: Also at University of Hamburg, Hamburg, Germany
 - 25: Also at Isfahan University of Technology, Isfahan, Iran
 - 26: Also at Brandenburg University of Technology, Cottbus, Germany
 - 27: Also at Forschungszentrum Jülich, Juelich, Germany
 - 28: Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
 - 29: Also at Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary
 - 30: Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
 - 31: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
 - 32: Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
 - 33: Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
 - 34: Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
 - 35: Also at Wigner Research Centre for Physics, Budapest, Hungary
 - 36: Also at IIT Bhubaneswar, Bhubaneswar, India
 - 37: Also at Institute of Physics, Bhubaneswar, India
 - 38: Also at Punjab Agricultural University, Ludhiana, India
 - 39: Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
 - 40: Also at Shoolini University, Solan, India
 - 41: Also at University of Hyderabad, Hyderabad, India
 - 42: Also at University of Visva-Bharati, Santiniketan, India
 - 43: Also at Indian Institute of Science (IISc), Bangalore, India
 - 44: Also at Indian Institute of Technology (IIT), Mumbai, India
 - 45: Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
 - 46: Now at Department of Physics, Isfahan University of Technology, Isfahan, Iran
 - 47: Also at Sharif University of Technology, Tehran, Iran
 - 48: Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
 - 49: Now at INFN Sezione di Bari, Università di Bari, Politecnico di Bari, Bari, Italy
 - 50: Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
 - 51: Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
 - 52: Also at Scuola Superiore Meridionale, Università di Napoli Federico II, Napoli, Italy
 - 53: Also at Università di Napoli 'Federico II', Napoli, Italy
 - 54: Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
 - 55: Also at Riga Technical University, Riga, Latvia
 - 56: Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
 - 57: Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
 - 58: Also at Institute for Nuclear Research, Moscow, Russia
 - 59: Now at National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia
 - 60: Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan
 - 61: Also at St. Petersburg Polytechnic University, St. Petersburg, Russia

- 62: Also at University of Florida, Gainesville, Florida, USA
- 63: Also at Imperial College, London, United Kingdom
- 64: Also at P.N. Lebedev Physical Institute, Moscow, Russia
- 65: Also at California Institute of Technology, Pasadena, California, USA
- 66: Also at Budker Institute of Nuclear Physics, Novosibirsk, Russia
- 67: Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia
- 68: Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- 69: Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
- 70: Also at National and Kapodistrian University of Athens, Athens, Greece
- 71: Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- 72: Also at Universität Zürich, Zurich, Switzerland
- 73: Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- 74: Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- 75: Also at Şirnak University, Sirnak, Turkey
- 76: Also at Near East University, Research Center of Experimental Health Science, Nicosia, Turkey
- 77: Also at Konya Technical University, Konya, Turkey
- 78: Also at Piri Reis University, Istanbul, Turkey
- 79: Also at Adiyaman University, Adiyaman, Turkey
- 80: Also at Necmettin Erbakan University, Konya, Turkey
- 81: Also at Bozok Universititesi Rektörlüğü, Yozgat, Turkey
- 82: Also at Marmara University, Istanbul, Turkey
- 83: Also at Milli Savunma University, Istanbul, Turkey
- 84: Also at Kafkas University, Kars, Turkey
- 85: Also at Istanbul Bilgi University, Istanbul, Turkey
- 86: Also at Hacettepe University, Ankara, Turkey
- 87: Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- 88: Also at Ozyegin University, Istanbul, Turkey
- 89: Also at Vrije Universiteit Brussel, Brussel, Belgium
- 90: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- 91: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- 92: Also at IPPP Durham University, Durham, United Kingdom
- 93: Also at Monash University, Faculty of Science, Clayton, Australia
- 94: Also at Università di Torino, Torino, Italy
- 95: Also at Bethel University, St. Paul, Minneapolis, USA
- 96: Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- 97: Also at United States Naval Academy, Annapolis, N/A, USA
- 98: Also at Bingol University, Bingol, Turkey
- 99: Also at Georgian Technical University, Tbilisi, Georgia
- 100: Also at Sinop University, Sinop, Turkey
- 101: Also at Erciyes University, Kayseri, Turkey
- 102: Also at Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China
- 103: Also at Texas A&M University at Qatar, Doha, Qatar
- 104: Also at Kyungpook National University, Daegu, Korea