

Observation of short range three-particle correlations in e^+e^- annihilations at LEP energies

DELPHI Collaboration

Abstract

Measurements are presented of short range three-particle correlations in e^+e^- annihilations at LEP using data collected by the DELPHI detector. At small values of the four-momentum difference, strong three-particle correlations are observed for like-sign (+++ and ---) and for unlike-sign (+ + - and + - -) pion combinations which are not a consequence of two-particle correlations. A possible explanation of the observed effects in like-sign combinations is the existence of higher order Bose-Einstein interference, which significantly changes the particle distributions in jets.

(To be submitted to Phys. Lett. B)

P.Abreu²¹, W.Adam⁵⁰, T.Adye³⁷, E.Agasi³¹, I.Ajinenko⁴², R.Aleksan³⁹, G.D.Alekseev¹⁶, P.P.Allport²², S.Almehed²⁴, S.J.Alvsvaag⁴, U.Amaldi⁹, S.Amato⁴⁷, A.Andreazza²⁸, M.L.Andrieux¹⁴, P.Antilogus²⁵, W-D.Apel¹⁷, Y.Arnoud³⁹, B.Åsman⁴⁴, J-E.Augustin¹⁹, A.Augustinus³¹, P.Baillon⁹, P.Bambade¹⁹, F.Barao²¹, R.Barate¹⁴, G.Barbiellini⁴⁶, D.Y.Bardin¹⁶, G.J.Barker³⁵, A.Baroncelli⁴⁰, O.Barring²⁴, J.A.Barrio²⁶, W.Bartl⁵⁰, M.J.Bates³⁷, M.Battaglia¹⁵, M.Baubillier²³, J.Baudot³⁹, K-H.Becks⁵², M.Begalli⁶, P.Beilliere⁸, Yu.Belokopytov⁹, K.Belous⁴², A.C.Benvenuti⁵, M.Berggren⁴¹, D.Bertrand², F.Bianchi⁴⁵, M.Bigi⁴⁵, M.S.Bilenky¹⁶, P.Billoir²³, D.Bloch¹⁰, M.Blume⁵², S.Blyth³⁵, V.Bocci³⁸, T.Bolognese³⁹, M.Bonesini²⁸, W.Bonivento²⁸, P.S.L.Booth²², G.Borisov⁴², C.Bosio⁴⁰, S.Bosworth³⁵, O.Botner⁴⁸, B.Bouquet¹⁹, C.Bourdarios⁹, T.J.V.Bowcock²², M.Bozzo¹³, P.Branchini⁴⁰, K.D.Brand³⁶, R.A.Brenner¹⁵, C.Bricman², L.Brillault²³, R.C.A.Brown⁹, P.Bruckman¹⁸, J-M.Brunet⁸, L.Bugge³³, T.Buran³³, A.Buys⁹, M.Caccia²⁸, M.Calvi²⁸, A.J.Camacho Rozas⁴¹, T.Camporesi⁹, V.Canale³⁸, M.Canepa¹³, K.Cankocak⁴⁴, F.Cao², F.Carena⁹, P.Carrillo⁴⁷, L.Carroll²², C.Caso¹³, M.V.Castillo Gimenez⁴⁹, A.Cattai⁹, F.R.Cavallo⁵, L.Cerrito³⁸, V.Chabaud⁹, Ph.Charpentier⁹, L.Chaussard²⁵, J.Chauveau²³, P.Checchia³⁶, G.A.Chelkov¹⁶, R.Chierici⁴⁵, P.Chochula⁷, V.Chorowicz⁹, V.Cindro⁴³, P.Collins⁹, J.L.Contreras¹⁹, R.Contri¹³, E.Cortina⁴⁹, G.Cosme¹⁹, F.Cossutti⁴⁶, H.B.Crawley¹, D.Crennell³⁷, G.Crosetti¹³, J.Cuevas Maestro³⁴, S.Czellar¹⁵, E.Dahl-Jensen²⁹, J.Dahm⁵², B.Dalmagne¹⁹, M.Dam³³, G.Damgaard²⁹, A.Daum¹⁷, P.D.Dauncey³⁷, M.Davenport⁹, W.Da Silva²³, C.Defoix⁸, G.Della Ricca⁴⁶, P.Delpierre²⁷, N.Demaria³⁵, A.De Angelis⁹, H.De Boeck², W.De Boer¹⁷, S.De Brabandere², C.De Clercq², C.De La Vaissiere²³, B.De Lotto⁴⁶, A.De Min²⁸, L.De Paula⁴⁷, C.De Saint-Jean³⁹, H.Dijkstra⁹, L.Di Ciaccio³⁸, F.Djama¹⁰, J.Dolbeau⁸, M.Donszelmann⁹, K.Doroba⁵¹, M.Dracos¹⁰, J.Drees⁵², K.-A.Drees⁵², M.Dris³², Y.Dufour⁸, F.Dupont¹⁴, D.Edsall¹, R.Ehret¹⁷, G.Eigen⁴, T.Ekelof⁴⁸, G.Ekspong⁴⁴, M.Elsing⁵², J-P.Engel²³, N.Ershaidat²³, B.Erzen⁴³, M.Espirito Santo²¹, E.Falk²⁴, D.Fassouliotis³², M.Feindt⁹, A.Fenyuk⁴², A.Ferrer⁴⁹, T.A.Filippas³², A.Firestone¹, P.-A.Fischer¹⁰, H.Foeth⁹, E.Fokitis³², F.Fontanelli¹³, F.Formenti⁹, B.Franek³⁷, P.Frenkiel⁸, D.C.Fries¹⁷, A.G.Frodesen⁴, R.Fruhworth⁵⁰, F.Fulda-Quenzer¹⁹, H.Furstenau⁹, J.Fuster⁴⁹, A.Galloni²², D.Gamba⁴⁵, M.Gandelman⁶, C.Garcia⁴⁹, J.Garcia⁴¹, C.Gaspar⁹, U.Gasparini³⁶, Ph.Gavillet⁹, E.N.Gazis³², D.Gele¹⁰, J-P.Gerber¹⁰, M.Gibbs²², D.Gillespie⁹, R.Gokiel⁵¹, B.Golob⁴³, G.Gopal³⁷, L.Gorn¹, M.Gorski⁵¹, Yu.Gouz⁴², V.Gracco¹³, E.Graziani⁴⁰, G.Grosdidier¹⁹, P.Gunnarsson⁴⁴, M.Gunther⁴⁸, J.Guy³⁷, U.Haedinger¹⁷, F.Hahn⁵², M.Hahn¹⁷, S.Hahn⁵², Z.Hajduk¹⁸, A.Hallgren⁴⁸, K.Hamacher⁵², W.Hao³¹, F.J.Harris³⁵, V.Hedberg²⁴, R.Henriques²¹, J.J.Hernandez⁴⁹, P.Herquet², H.Herr⁹, T.L.Hessing⁹, E.Higon⁴⁹, H.J.Hilke⁹, T.S.Hill¹, S.-O.Holmgren⁴⁴, P.J.Holt³⁵, D.Holthuizen³¹, M.Houlden²², K.Huet², K.Hultqvist⁴⁴, P.Ioannou³, J.N.Jackson²², R.Jacobsson⁴⁴, P.Jalocha¹⁸, R.Janik⁷, G.Jarlskog²⁴, P.Jarry³⁹, B.Jean-Marie¹⁹, E.K.Johansson⁴⁴, L.Jonsson²⁴, P.Jonsson²⁴, C.Joram⁹, P.Juillot¹⁰, M.Kaiser¹⁷, G.Kalmus³⁷, F.Kapusta²³, M.Karlsson⁴⁴, E.Karvelas¹¹, A.Katargin⁴², S.Katsanevas³, E.C.Katsoufis³², R.Keranen¹⁵, B.A.Khomenko¹⁶, N.N.Khovanski¹⁶, B.King²², N.J.Kjaer²⁹, H.Klein⁹, A.Kloving⁴, P.Kluit³¹, J.H.Koehne¹⁷, B.Koene³¹, P.Kokkinias¹¹, M.Koratzinos⁹, V.Kostioukhine⁴², C.Kourkoumelis³, O.Kouznetsov¹³, P.-H.Kramer⁵², M.Krammer⁵⁰, C.Kreuter¹⁷, J.Krolikowski⁵¹, I.Kronkvist²⁴, Z.Krumstein¹⁶, W.Krupinski¹⁸, P.Kubinec⁷, W.Kucewicz¹⁸, K.Kurvinen¹⁵, C.Lacasta⁴⁹, I.Laktineh²⁵, S.Lamblot²³, J.W.Lamsa¹, L.Lanceri⁴⁶, D.W.Lane¹, P.Langefeld⁵², V.Lapin⁴², I.Last²², J-P.Laugier³⁹, R.Lauhakangas¹⁵, G.Leder⁵⁰, F.Ledroit¹⁴, V.Lefebure², C.K.Legan¹, R.Leitner³⁰, Y.Lemoigne³⁹, J.Lemonne², G.Lenzen⁵², V.Lepeltier¹⁹, T.Lesiak³⁶, D.Liko⁵⁰, R.Lindner⁵², A.Lipniacka¹⁹, I.Lippi³⁶, B.Loerstad²⁴, M.Lokajicek¹², J.G.Loken³⁵, J.M.Lopez⁴¹, A.Lopez-Fernandez⁹, M.A.Lopez Aguera⁴¹, D.Loukas¹¹, P.Lutz³⁹, L.Lyons³⁵, J.MacNaughton⁵⁰, G.Maehlum¹⁷, A.Maio²¹, V.Malychev¹⁶, F.Mand⁵⁰, J.Marco⁴¹, B.Marechal⁴⁷, M.Margoni³⁶, J.-C.Marin⁹, C.Mariotti⁴⁰, A.Markou¹¹, T.Maron⁵², C.Martinez-Rivero⁴¹, F.Martinez-Vidal⁴⁹, S.Marti i Garcia⁴⁹, F.Matorras⁴¹, C.Matteuzzi²⁸, G.Matthiae³⁸, M.Mazzucato³⁶, M.Mc Cubbin⁹, R.Mc Kay¹, R.Mc Nulty²², J.Medbo⁴⁸, C.Meroni²⁸, W.T.Meyer¹, A.Miagkov⁴², M.Michelotto³⁶, E.Migliore⁴⁵, L.Mirabito²⁵, W.A.Mitaroff⁵⁰, U.Mjoernmark²⁴, T.Moa⁴⁴, R.Moeller²⁹, K.Moenig⁹, M.R.Monge¹³, P.Moretini¹³, H.Mueller¹⁷, L.M.Mundim⁶, W.J.Murray³⁷, B.Muryn¹⁸, G.Myatt³⁵, F.Naraghi¹⁴, F.L.Navarria⁵, S.Navas⁴⁹, P.Negri²⁸, S.Nemecek¹², W.Neumann⁵², N.Neumeister⁵⁰, R.Nicolaidou³, B.S.Nielsen²⁹, M.Nieuwenhuizen³¹, V.Nikolaenko¹⁰, P.Niss⁴⁴, A.Nomerotski³⁶, A.Normand³⁵, W.Oberschulte-Beckmann¹⁷, V.Obraztsov⁴², A.G.Olshevski¹⁶, A.Onofre²¹, R.Orava¹⁵, K.Osterberg¹⁵, A.Ouraou³⁹, P.Paganini¹⁹, M.Paganoni²⁸, P.Pages¹⁰, H.Palka¹⁸, Th.D.Papadopoulou³², L.Pape⁹, C.Parkes³⁵, F.Parodi¹³, A.Passeri⁴⁰, M.Pegoraro³⁶, L.Peralta²¹, H.Pernegger⁵⁰, M.Pernicka⁵⁰, A.Perrotta⁵, C.Petridou⁴⁶, A.Petrolini¹³, H.T.Phillips³⁷, G.Piana¹³, F.Pierre³⁹, M.Pimenta²¹, S.Plaszczynski¹⁹, O.Podobrin¹⁷, M.E.Pol⁶, G.Polok¹⁸, P.Poropat⁴⁶, V.Pozdniakov¹⁶, M.Prest⁴⁶, P.Privitera³⁸, N.Pukhaeva¹⁶, A.Pullia²⁸, D.Radojicic³⁵, S.Ragazzi²⁸, H.Rahmani³², J.Rames¹², P.N.Ratoff²⁰, A.L.Read³³, M.Reale⁵², P.Rebecchi¹⁹, N.G.Redaeli²⁸, M.Regler⁵⁰, D.Reid⁹, P.B.Renton³⁵, L.K.Resvanis³, F.Richard¹⁹, J.Richardson²², J.Ridky¹², G.Rinaudo⁴⁵, I.Ripp³⁹, A.Romero⁴⁵, I.Roncagliolo¹³, P.Ronchese³⁶, L.Roos¹⁴, E.I.Rosenberg¹, E.Rosso⁹, P.Roudeau¹⁹, T.Rovelli⁵, W.Ruckstuhl³¹, V.Ruhlmann-Kleider³⁹, A.Ruiz⁴¹, H.Saarikko¹⁵, Y.Sacquin³⁹, A.Sadovsky¹⁶, G.Sajot¹⁴, J.Salt⁴⁹, J.Sanchez²⁶, M.Sannino¹³, H.Schneider¹⁷, M.A.E.Schyns⁵², G.Sciolla⁴⁵, F.Scuri⁴⁶, Y.Sedykh¹⁶, A.M.Segar³⁵, A.Seitz¹⁷, R.Sekulin³⁷, R.C.Shellard⁶, I.Siccama³¹, P.Siegrist³⁹, S.Simonetti³⁹, F.Simonetto³⁶, A.N.Sisakian¹⁶, B.Sitar⁷, T.B.Skaali³³, G.Smadja²⁵, N.Smirnov⁴², O.Smirnova¹⁶, G.R.Smith³⁷, A.Sokolov⁴², R.Sosnowski⁵¹, D.Souza-Santos⁶, T.Spaso²¹, E.Spiriti⁴⁰, S.Squarcia¹³, H.Staek⁵², C.Stanescu⁴⁰, S.Stapnes³³, I.Stavitski³⁶, K.Stepaniak⁵¹, F.Stichelbaut⁹, A.Stocchi¹⁹, J.Strauss⁵⁰, R.Strub¹⁰, B.Stugu⁴, M.Szczekowski⁵¹, M.Szeptycka⁵¹, T.Tabarelli²⁸, J.P.Tavernet²³, A.Tilquin²⁷, J.Timmermans³¹, L.G.Tkatchev¹⁶, T.Todorov¹⁰, D.Z.Toet³¹, A.Tomaradze²

B.Tome²¹, L.Tortora⁴⁰, G.Transtromer²⁴, D.Treille⁹, W.Trischuk⁹, G.Tristram⁸, A.Trombini¹⁹, C.Troncon²⁸, A.Tsirou⁹, M-L.Turluer³⁹, I.A.Tyapkin¹⁶, M.Tyndel³⁷, S.Tzamaras²², B.Ueberschaer⁵², S.Ueberschaer⁵², O.Ullaland⁹, G.Valenti⁵, E.Vallazza⁹, C.Vander Velde², G.W.Van Apeldoorn³¹, P.Van Dam³¹, J.Van Eldik³¹, N.Vassilopoulos³⁵, G.Vegni²⁸, L.Ventura³⁶, W.Venus³⁷, F.Verbeure², M.Verlato³⁶, L.S.Vertogradov¹⁶, D.Vilanova³⁹, P.Vincent²⁵, L.Vitale⁴⁶, E.Vlasov⁴², A.S.Vodopyanov¹⁶, V.Vrba¹², H.Wahlen⁵², C.Walck⁴⁴, A.Wehr⁵², M.Weierstall⁵², P.Weilhammer⁹, A.M.Wetherell⁹, D.Wicke⁵², J.H.Wickens², M.Wielers¹⁷, G.R.Wilkinson³⁵, W.S.C.Williams³⁵, M.Winter¹⁰, M.Witek⁹, K.Woschnagg⁴⁸, K.Yip³⁵, O.Yushchenko⁴², F.Zach²⁵, C.Zacharatou²⁴, A.Zaitsev⁴², A.Zalewska¹⁸, P.Zalewski⁵¹, D.Zavrtanik⁴³, E.Zevgolatakos¹¹, N.I.Zimin¹⁶, M.Zito³⁹, D.Zontar⁴³, R.Zuberi³⁵, G.C.Zucchelli⁴⁴, G.Zumerle³⁶

¹Ames Laboratory and Department of Physics, Iowa State University, Ames IA 50011, USA

²Physics Department, Univ. Instelling Antwerpen, Universiteitsplein 1, B-2610 Wilrijk, Belgium and IIHE, ULB-VUB, Pleinlaan 2, B-1050 Brussels, Belgium

and Faculté des Sciences, Univ. de l'Etat Mons, Av. Maistriau 19, B-7000 Mons, Belgium

³Physics Laboratory, University of Athens, Solonos Str. 104, GR-10680 Athens, Greece

⁴Department of Physics, University of Bergen, Allégaten 55, N-5007 Bergen, Norway

⁵Dipartimento di Fisica, Università di Bologna and INFN, Via Irnerio 46, I-40126 Bologna, Italy

⁶Centro Brasileiro de Pesquisas Físicas, rua Xavier Sigaud 150, RJ-22290 Rio de Janeiro, Brazil and Depto. de Física, Pont. Univ. Católica, C.P. 38071 RJ-22453 Rio de Janeiro, Brazil

and Inst. de Física, Univ. Estadual do Rio de Janeiro, rua São Francisco Xavier 524, Rio de Janeiro, Brazil

⁷Comenius University, Faculty of Mathematics and Physics, Mlynska Dolina, SK-84215 Bratislava, Slovakia

⁸Collège de France, Lab. de Physique Corpusculaire, IN2P3-CNRS, F-75231 Paris Cedex 05, France

⁹CERN, CH-1211 Geneva 23, Switzerland

¹⁰Centre de Recherche Nucléaire, IN2P3 - CNRS/ULP - BP20, F-67037 Strasbourg Cedex, France

¹¹Institute of Nuclear Physics, N.C.S.R. Demokritos, P.O. Box 60228, GR-15310 Athens, Greece

¹²FZU, Inst. of Physics of the C.A.S. High Energy Physics Division, Na Slovance 2, 180 40, Praha 8, Czech Republic

¹³Dipartimento di Fisica, Università di Genova and INFN, Via Dodecaneso 33, I-16146 Genova, Italy

¹⁴Institut des Sciences Nucléaires, IN2P3-CNRS, Université de Grenoble 1, F-38026 Grenoble Cedex, France

¹⁵Research Institute for High Energy Physics, SEFT, P.O. Box 9, FIN-00014 Helsinki, Finland

¹⁶Joint Institute for Nuclear Research, Dubna, Head Post Office, P.O. Box 79, 101 000 Moscow, Russian Federation

¹⁷Institut für Experimentelle Kernphysik, Universität Karlsruhe, Postfach 6980, D-76128 Karlsruhe, Germany

¹⁸High Energy Physics Laboratory, Institute of Nuclear Physics, Ul. Kawioru 26a, PL-30055 Krakow 30, Poland

¹⁹Université de Paris-Sud, Lab. de l'Accélérateur Linéaire, IN2P3-CNRS, Bât. 200, F-91405 Orsay Cedex, France

²⁰School of Physics and Materials, University of Lancaster, Lancaster LA1 4YB, UK

²¹LIP, IST, FCUL - Av. Elias Garcia, 14-1º, P-1000 Lisboa Codex, Portugal

²²Department of Physics, University of Liverpool, P.O. Box 147, Liverpool L69 3BX, UK

²³LPNHE, IN2P3-CNRS, Universités Paris VI et VII, Tour 33 (RdC), 4 place Jussieu, F-75252 Paris Cedex 05, France

²⁴Department of Physics, University of Lund, Sölvegatan 14, S-22363 Lund, Sweden

²⁵Université Claude Bernard de Lyon, IPNL, IN2P3-CNRS, F-69622 Villeurbanne Cedex, France

²⁶Universidad Complutense, Avda. Complutense s/n, E-28040 Madrid, Spain

²⁷Univ. d'Aix - Marseille II - CPP, IN2P3-CNRS, F-13288 Marseille Cedex 09, France

²⁸Dipartimento di Fisica, Università di Milano and INFN, Via Celoria 16, I-20133 Milan, Italy

²⁹Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen 0, Denmark

³⁰NC, Nuclear Centre of MFF, Charles University, Areal MFF, V Holesovickach 2, 180 00, Praha 8, Czech Republic

³¹NIKHEF-H, Postbus 41882, NL-1009 DB Amsterdam, The Netherlands

³²National Technical University, Physics Department, Zografou Campus, GR-15773 Athens, Greece

³³Physics Department, University of Oslo, Blindern, N-1000 Oslo 3, Norway

³⁴Dpto. Física, Univ. Oviedo, C/P. Pérez Casas, S/N-33006 Oviedo, Spain

³⁵Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK

³⁶Dipartimento di Fisica, Università di Padova and INFN, Via Marzolo 8, I-35131 Padua, Italy

³⁷Rutherford Appleton Laboratory, Chilton, Didcot OX11 0QX, UK

³⁸Dipartimento di Fisica, Università di Roma II and INFN, Tor Vergata, I-00173 Rome, Italy

³⁹Centre d'Etudes de Saclay, DSM/DAPNIA, F-91191 Gif-sur-Yvette Cedex, France

⁴⁰Istituto Superiore di Sanità, Ist. Naz. di Fisica Nucl. (INFN), Viale Regina Elena 299, I-00161 Rome, Italy

⁴¹C.E.A.F.M., C.S.I.C. - Univ. Cantabria, Avda. los Castros, S/N-39006 Santander, Spain, (CICYT-AEN93-0832)

⁴²Inst. for High Energy Physics, Serpukov P.O. Box 35, Protvino, (Moscow Region), Russian Federation

⁴³J. Stefan Institute and Department of Physics, University of Ljubljana, Jamova 39, SI-61000 Ljubljana, Slovenia

⁴⁴Fysikum, Stockholm University, Box 6730, S-113 85 Stockholm, Sweden

⁴⁵Dipartimento di Fisica Sperimentale, Università di Torino and INFN, Via P. Giuria 1, I-10125 Turin, Italy

⁴⁶Dipartimento di Fisica, Università di Trieste and INFN, Via A. Valerio 2, I-34127 Trieste, Italy

and Istituto di Fisica, Università di Udine, I-33100 Udine, Italy

⁴⁷Univ. Federal do Rio de Janeiro, C.P. 68528 Cidade Univ., Ilha do Fundão BR-21945-970 Rio de Janeiro, Brazil

⁴⁸Department of Radiation Sciences, University of Uppsala, P.O. Box 535, S-751 21 Uppsala, Sweden

⁴⁹IFIC, Valencia-CSIC, and D.F.A.M.N., U. de Valencia, Avda. Dr. Moliner 50, E-46100 Burjassot (Valencia), Spain

⁵⁰Institut für Hochenergiephysik, Österr. Akad. d. Wissensch., Nikolsdorfergasse 18, A-1050 Vienna, Austria

⁵¹Inst. Nuclear Studies and University of Warsaw, Ul. Hoza 69, PL-00681 Warsaw, Poland

⁵²Fachbereich Physik, University of Wuppertal, Postfach 100 127, D-42097 Wuppertal 1, Germany

1 Introduction

Studies of the properties of hadronic events in e^+e^- annihilations and tests of QCD generators are usually performed using global event shape variables and single particle distributions. As the multiplicity of particles increases with increasing initial energy, correlations between particles change their distributions in jets and an understanding of these correlations becomes important. Studies of correlations at high energy are thus related to the composition of jets and tests of QCD models. A clear example of the influence of correlations on particle distributions is the distortion of the Breit-Wigner shape for oppositely charged pions from the decay of broad resonances by residual Bose-Einstein (BE) correlations, which was observed at LEP energies [1](see also [2]). Recently, it was also shown that the observed W mass at LEP200 is likely to be affected by BE correlations [3]. A better understanding of these correlations, as well as comparisons of BE algorithms with experimental data at high energies, becomes necessary for high precision measurements of the W mass.

Two-particle correlations have been studied extensively in hadron-hadron interactions [4,5] and in e^+e^- annihilations at LEP energies [6]. However, very few experimental results are available on three-particle correlations which can be generated either by two-particle correlations or by genuine three-particle correlations. Identifying the latter is complicated because of the need for high statistics of large multiplicity events and the problem of subtracting the direct consequences of two-particle correlations. Obvious $\pi^+\pi^+\pi^-$ and $\pi^+\pi^-\pi^-$ correlations are generated in the decays of many resonances. The most prominent example at low masses is the η' decay into $\eta\pi^+\pi^-$ with subsequent decay of the η into $\pi^+\pi^-\pi^0$ or $\pi^+\pi^-\gamma$. At higher three-pion masses there are many contributions from the decays of charm and beauty particles.

Short-range rapidity correlations are observed for $(++-)$ and $(+--)$ combinations in several experiments [7–10]. In contrast to the unlike-sign combinations, no genuine correlations were found for the like-sign $(+++)$ and $(---)$ configuration. Furthermore, in the three-particle Bose-Einstein effect studied in [8,10,11], no evidence was found for three-particle correlations beyond those generated from pairs in the triplet. Only one hadron-proton experiment recently reported the observation of positive short-range higher order like-sign correlations [12].

Experimental evidence for genuine like-sign three-particle correlations has still to be reported; it constitutes an important theoretical issue for the understanding of BE correlations [13].

In this paper, the first evidence is presented for genuine three particle correlations at LEP using data collected by the DELPHI detector. We discuss like-sign and unlike-sign three-particle correlations after subtraction of the two-particle ones. It is demonstrated that positive short range correlations exist for both like-sign and unlike-sign configurations. Both these correlations can be understood, at least partly, as following from higher order Bose-Einstein correlations, which change the event shape at small relative momenta. Experimental data are compared with various modifications of the JETSET PS model [14].

2 Analysis

The single-particle density is defined as

$$\rho_1(q_a) = \frac{E_a}{N_{ev}} \frac{d^3 n}{d^3 p_a}, \quad (1)$$

the two-particle density as

$$\rho_2(q_a, q_b) = \frac{E_a E_b}{N_{ev}} \frac{d^6 n}{d^3 p_a d^3 p_b} \quad (2)$$

and the three-particle density as

$$\rho_3(q_a, q_b, q_c) = \frac{E_a E_b E_c}{N_{ev}} \frac{d^9 n}{d^3 p_a d^3 p_b d^3 p_c}, \quad (3)$$

where q_i and p_i are the four vector and three vector momentum, respectively, and E_i the energy of the particle; n is the number of particles (in Eq. 1), of doublets (in Eq. 2) or triplets (in Eq. 3) and N_{ev} is the number of events. For n identical particles, these densities are normalized to $\langle n \rangle$, $\langle n(n-1) \rangle$ and $\langle n(n-1)(n-2) \rangle$, respectively. We study the normalized three-body correlation function written as [15]

$$R_3 = R_3'' - R_3' + 1 \quad (4)$$

where the quantity

$$R_3'' = [\rho_3(q_a, q_b, q_c) - \rho_1(q_a)\rho_1(q_b)\rho_1(q_c)] / \rho_1(q_a)\rho_1(q_b)\rho_1(q_c) \quad (5)$$

is the full three-particle correlation function and

$$R_3' = \frac{[\rho_2(q_a, q_b)\rho_1(q_c) + \rho_2(q_a, q_c)\rho_1(q_b) + \rho_2(q_b, q_c)\rho_1(q_a) - 3\rho_1(q_a)\rho_1(q_b)\rho_1(q_c)]}{\rho_1(q_a)\rho_1(q_b)\rho_1(q_c)} \quad (6)$$

is the three-particle correlation arising from correlations in the pairs of the triplet. The $\rho_1(q_a)\rho_1(q_b)\rho_1(q_c)$ term represents uncorrelated production of the three particles, each to be taken with its density defined in Eq. 1. The $\rho_2(q_a, q_b)\rho_1(q_c)$ term, as well as the next two terms in Eq. 6, contain the two-body correlations. The genuine three-particle correlation function R_3 is then given by

$$R_3 = [\rho_3 + 3\rho_1\rho_1\rho_1 - 3\rho_2\rho_1] / \rho_1\rho_1\rho_1 \quad (7)$$

where the q -variables have been omitted for clarity and $3\rho_2\rho_1$ is the sum of the three $\rho_2\rho_1$ terms in Eq. 6. The quantity R_3 therefore does not contain three-particle correlations which arise from two-body correlations nor is it sensitive to possible bias in two-particle combinations due to Dalitz pairs, γ -conversions, or particle or resonance decays.

The kinematical variable used in this analysis is

$$Q^2 = q_{12}^2 + q_{23}^2 + q_{31}^2 \quad (8)$$

with $q_{ij}^2 = -(q_i - q_j)^2$. For three-pion systems, $Q^2 = M_{123}^2 - 9m_\pi^2$, with M_{123} the invariant mass of the pion triplet. It has been shown [5] that this Lorentz invariant variable is more sensitive to correlations than the rapidity variable.

3 Experimental procedure

This study is based on a sample of about 1.5 million hadronic events collected with the DELPHI detector during the 1992 and 1993 running periods, at a center of mass energy at and around the Z^0 peak. The detector has been described in [16]. The analysis uses the same methodology as in [17]. All charged particles were considered to be pions and only charged particles satisfying the following requirements were used:

- polar angle θ with respect to the beam axis between 25° and 155° ;
- momentum greater than $0.2 \text{ GeV}/c$ and less than $50 \text{ GeV}/c$;
- measured track length in the TPC, the main tracking chamber, greater than 50 cm ;
- measured impact parameter with respect to the event vertex less than 5 cm in the transverse plane and 10 cm along the beam direction.

Hadronic events were selected by requiring that:

- there were at least 5 charged particles in the event;
- the total energy of the charged particles exceeded 3 GeV in each of the two hemispheres defined with respect to the beam direction;
- the total energy of all charged particles was larger than 15 GeV ;
- the total momentum imbalance was less than $30 \text{ GeV}/c$;
- the polar angle of the thrust axis θ_{th} , defined with respect to the beam direction, satisfied $|\cos \theta_{\text{th}}| < 0.75$.

The contamination from events due to beam-gas scattering, $\gamma\gamma$ interactions and $\tau^+\tau^-$ events was estimated to be less than 0.3% of the selected events. Only charged particles with an impact parameter less than 0.1 cm in the transverse plane and 1 cm in the beam direction were used in the calculation of the Q distribution. Thus only particles originating from the primary vertex were included. These strict cuts on both transverse and longitudinal impact parameters removed most tracks from K_S^0 and Λ decays. To select clean two-jet events, a cut on the thrust variable $T > 0.97$ was imposed. A sample of 327,162 events was available after all cuts.

In order to calculate the terms $\rho_2\rho_1$ and $\rho_1\rho_1\rho_1$ in Eq. 5-7, the following mixing technique was used. A pool of particle four-vectors with momentum components calculated with respect to the thrust axis, was constructed from a large number of events. To calculate $\rho_1\rho_1\rho_1$, three tracks were randomly drawn from the pool and the term was calculated in the same way as in real events; to calculate $\rho_2\rho_1$, two particles were taken from the same event and the third one from another event.

The analysis was limited to $Q > 0.15 \text{ GeV}/c$ in order to avoid the region where the three tracks could be geometrically confused. This value for triplets corresponds, on average, to $q_{ij} > 0.09 \text{ GeV}/c$ for pairs. Moreover, if false two-body correlations are introduced they are subtracted in the quantity R_3 . The correction for geometrical acceptance, kinematical cuts, particle interactions with the detector material and other detector imperfections, was made by using the Monte Carlo simulation program DELSIM [18]. A sample of Z^0 hadronic events was generated with JETSET and all tracks were followed through the full detector. After application of the same cuts as on the data, including the thrust cut, about 1,000,000 events remained. Bin by bin correction factors were calculated for ρ_3 , $\rho_2\rho_1$, and $\rho_1\rho_1\rho_1$. In the following, only corrected distributions will be presented. However, it was checked that the same features are present at the level of the uncorrected distributions.

4 Results and discussion

Figure 1a shows R_3'' , the full three-particle correlation function for like-sign triplets using the normalization of the density functions as given above. A strong correlation is observed at small Q -values. However, R_3' , the background three-particle correlation arising from two-particle correlations between particles in the triplet, also shows a strong rise towards $Q = 0$. Figure 1a demonstrates that most of the three-particle correlations are due to two-particle correlations, but also that some excess of R_3'' over R_3' is present at small Q . The three-particle correlation functions R_3'' and R_3' for unlike-sign triplets are presented in Figure 1b. A difference between the R_3'' and R_3' functions at small Q is also present for unlike-sign triplets.

The genuine three-particle correlations for like-sign triplets, R_3 , are shown in Figure 2a. A clear signal is observed at $Q < 0.5$ GeV/ c , corresponding to a short-range three-particle correlation. A reference sample such as the one of unlike-sign pion combinations is inadequate because of the presence of correlations in this sample. This is clearly seen from Figure 2b where the function $R_3(Q)$ is plotted for unlike-sign triplets.

The predictions of JETSET 7.3 PS [14] with no BE correlations included, are shown in Figure 2 (full lines). Here and in the following, the JETSET events are subject to the same cuts and contributions to ρ_3 calculated from single events and the other terms with the same mixing technique as the experimental data. The statistical errors on the JETSET are about three times smaller than those on the data points. At small Q -values, strong differences are observed between JETSET and the data for both like-sign and unlike-sign triplets. For like-sign triplets the model does not show any increase of the correlation function for $Q \rightarrow 0$, compatible with no like-sign three-particle correlations in the model. A steep increase of the correlation function at small Q -values is observed in the model without BE correlations for $(+ + -)$ and $(+ - -)$ combinations (full line in Figure 2b). In the model, the η' production rate was reduced by a factor 4, based on a measurement of η' production at LEP [19] which is about 4 times smaller than the one predicted by the original version of JETSET. It was checked that this modification has little effect on the $(+ + +)$ and $(- - -)$ correlations but changes the $(+ + -)$ and $(+ - -)$ correlations (dashed line in Figure 2b). The larger Q value of the first data point for $(+ + -)$ and $(+ - -)$ correlations (0.25 GeV/ c as compared to 0.15 GeV/ c for like-sign triplets) and the larger errors on the the first two data points, are due to the large uncertainty on η' production in JETSET and consequent uncertainties in the correction factors. Simulations based on JETSET show that R_3 for like-sign combinations is only slightly affected if $b\bar{b}$ events are removed from the sample whereas the change in R_3 for unlike-sign combinations is comparable to the effect of reducing η' production. All modifications of JETSET yield expected results, supporting the method used.

Higher-order BE correlations are a natural source of like-sign three-particle correlations. The shape of $R_3(Q)$ for $(+ + +)$ and $(- - -)$ combinations is reminiscent of the shape of the two-body BE correlations [20–22]. In analogy to the latter it can be parametrized by [23]

$$R_3(Q) \sim [1 + 2\lambda_3 \exp(-r_3^2 Q^2)]. \quad (9)$$

The factor two before the λ_3 parameter arises from the presence of two possible diagrams with exchange of identical pions within a triplet [13]. The total three-particle BE correlation is built up from three diagrams with interchange of two particles in the triplet, plus two diagrams of interchange of the three particles. The latter correspond to the quantity R_3 mentioned above, and the former to the “background” term R_3' .

N	0.92 ± 0.01
δ	$0.11 \pm 0.01 \text{ (GeV}/c\text{)}^{-1}$
λ_3	$0.28 \pm 0.05 \text{ (stat)} \pm 0.07 \text{ (syst)}$
r_3	$3.33 \pm 0.20 \text{ (stat)} \pm 0.16 \text{ (syst)} \text{ (GeV}/c\text{)}^{-1}$

Table 1: Results of the fit of the three-particle correlation function $R_3(Q)$ for $(+++)$ and $(---)$ combinations with Eq. 10.

	λ_3	$r_3 \text{ (GeV}/c\text{)}^{-1}$
$T > 0.95$	0.34 ± 0.05	3.29 ± 0.19
$\theta > 2^\circ$	0.35 ± 0.06	3.48 ± 0.20
Data/JETSET ($T > 0.97$)	0.24 ± 0.04	3.32 ± 0.25
Data/JETSET ($T > 0.95$)	0.26 ± 0.04	3.17 ± 0.22
Uncorrected data	0.25 ± 0.04	4.16 ± 0.30
Data/DELSIM	0.19 ± 0.03	3.55 ± 0.24

Table 2: Values of the parameters λ_3 and r_3 with different analysis methods (see text).

A modification of Eq. 9 is often used to parametrize the experimental data:

$$R_3(Q) = N(1 + \delta Q)[1 + 2\lambda_3 \exp(-r_3^2 Q^2)], \quad (10)$$

where N is a normalization factor and the term $(1 + \delta Q)$ is an empirical term which takes into account the rise of the correlation function at large Q -values. A fit of Eq. 10 to the data with $Q < 1.35 \text{ GeV}/c$ yielded the parameter values given in Table 1 ($\chi^2/\text{DF}=5.2/8$). The fitted form is shown in Figure 2a as a dot-dashed curve. The experimental resolution for Q at $Q < 0.5 \text{ GeV}/c$ is less than $25 \text{ MeV}/c$, i.e. four times smaller than the bin size, and was neglected.

To estimate the systematic errors on the parameters λ_3 and r_3 :

- the cut on the thrust variable was chosen to be $T > 0.95$; this increased the data sample by 60%;
- a cut was made of minimum 2° on the opening angle θ between pairs of tracks in the triplet;
- R_3 for the corrected data was divided by R_3 of JETSET in order to reduce possible residual correlations which are not due to BE correlations.

The values obtained are given in Table 2. The maximal deviation of the λ_3 and r_3 values was used as an estimate of the systematic errors given in Table 1. The results were checked by relaxing the impact parameter requirements to 5 and 10 cm respectively. The conclusions of this analysis are not affected by these cuts. In order to check detector influences Eq. 10 was also fit to the uncorrected data of R_3 and to the uncorrected data of R_3 divided by R_3 of simulated events, fully tracked through the detector by means of DELSIM. The values of λ_3 and R_3 obtained with this procedure are presented in Table 2.

It should be remarked that calculation with JETSET shows that the fraction of pure $\pi\pi\pi$ triplets in all charge combinations is 72% for the interval $0.15 < Q < 0.5 \text{ GeV}/c$. The values of λ_3 and r_3 in Table 1 are not corrected for the contamination by triplets where (at least) one of the particles is not a pion. Correcting for this effect would increase the value of λ_3 by 39%, in the hypothesis that non- $\pi\pi\pi$ triplets are uncorrelated.

Bose-Einstein correlations can be included in JETSET where they are introduced as a final state interaction. Practically, after the generation of the pion momenta the generated values of the momenta of all identical pions are modified in such a way that their momentum vector differences are reduced with a quantity determined by the chosen form of parametrization and given parameters of correlation strength λ and radius r [14] in order to describe the like-sign two-pion distribution for $R(Q)$. In [17], DELPHI demonstrated that JETSET reproduces well the two-particle correlations if BE correlations are switched on in the model with Gaussian parametrisation for pions which are produced promptly or are decay products of short-lived resonances[†]. The maximal correlation strength $\lambda = 1$ and a radius $r = 0.50$ fm were used. This method of including Bose-Einstein correlations in the model moves *all* direct like-sign pions closer together in momentum space and therefore also yields genuine higher-order correlations with, however, some arbitrary form and correlation strength [3]. Another weakness of the model for BE correlations is that JETSET does not take into account the lifetime of resonances. Consequently, the model can only give a qualitative picture of these final state effects and provide evidence that genuine higher order Bose-Einstein correlations are present at small Q for $R_3(Q)$.

In Figure 3, the three-particle correlation functions R_3 are shown with the predictions of JETSET with and without BE correlations included. The parameters used to include the BE correlations are the same as in the two-particle correlation study of DELPHI [17]. The model is in reasonable agreement with the data for the $(+ + -)$ and $(+ - -)$ configurations and gives an enhancement for the $(+ + +)$ and $(- - -)$ correlations. Even if the enhancement in R_3 at low Q values would be partly due to two-particle correlations, arising from the method used, the λ_3 value obtained from the ratio R_3 of the data to R_3 of JETSET with BE effects included would give a lower limit of λ_3 . The value obtained by a fit to Eq. 10 is 0.14 ± 0.03 .

Bose-Einstein interference in JETSET not only changes the distribution of like-sign correlations (Figure 3a), but also the unlike-sign ones (Figure 3b) and leads to better agreement with the data. Due to BE correlations, particle distributions and invariant masses of jets are changed [3]. This observation is important for studies at high energies, particularly for the W mass measurement at LEP200.

5 Conclusions

The first observation is reported of genuine three-like-sign particle correlations in e^+e^- annihilations at the Z^0 mass. These correlations can be explained as a higher order Bose-Einstein enhancement. The JETSET model with BE correlations included, with the same λ and r parameters which are used for describing the two-particle correlations, yields reasonable agreement for the $(+ + -)$ and $(+ - -)$ correlations and gives an enhancement for $(+ + +)$ and $(- - -)$ correlations. One possible explanation of the inadequacy to describe the $(+ + +)$ and $(- - -)$ configurations could be an incomplete treatment of Bose-Einstein correlations in the model.

Acknowledgements — We thank T. Sjöstrand for very useful discussions on Bose-Einstein correlations in JETSET. We are greatly indebted to our technical collaborators and to the funding agencies for their support in building and operating the DELPHI detector, and to the members of the CERN-SL Division for the excellent performance of the LEP collider.

[†]All resonances with lifetime larger than the $K^*(890)$ were considered as long-lived.

References

- [1] P.D. Acton et al. (OPAL Coll.), *Z. Phys.* C56 (1992) 521; P. Abreu et al. (DELPHI Coll.), *Z. Phys.* C65 (1995) 587.
- [2] G. Lafferty, *Z. Phys.* C60 (1993) 659; S. Haywood, *Where are we going with Bose-Einstein – A mini review*, RAL Report 94-074.
- [3] L. Lönnblad and T. Sjöstrand, *Bose-Einstein effects and W mass determinations*, CERN-TH/95-17.
- [4] W.A. Zajc in *Hadronic Multiparticle production*, Ed. P. Carruthers, World Scientific, 1988, p. 235-288; E.A. De Wolf, *Bose-Einstein correlations* Proc. XXVII Int. Conf. on High Energy Physics, Eds. P.J. Bussey and I.G. Knowles, Inst. of Physics Publ., 1995, p. 1281.
- [5] E.A. De Wolf, I.M. Dremin, W.K. Kittel, *Scaling laws for density correlations and fluctuations in Multiparticle Dynamics*, Nijmegen preprint HEN-362 (1993), Brussels preprint IIHE-ULB-VUB-93-01; to be published in *Phys. Reports*.
- [6] P.D. Acton et al. (OPAL Coll.), *Phys. Lett.* B267 (1991) 143; P. Abreu et al. (DELPHI Coll.), *Phys. Lett.* B286 (1992) 201; D. Decamp et al. (ALEPH Coll.), *Z. Phys.* C54 (1992) 75; P.D. Acton et al., (OPAL Coll.), *Z. Phys.* C58 (1993) 207; P.D. Acton et al. (OPAL Coll.), *Phys. Lett.* B287 (1992) 401.
- [7] V.P. Kenney et al., *Nucl. Phys.* B144 (1978) 312.
- [8] J.L. Bailly et al. (NA23 Coll.), *Z. Phys.* C43 (1989) 341.
- [9] A. Breakstone et al. (SFM Coll.), *Mod. Phys. Lett.* A6 (1991) 2785.
- [10] M. Althoff et al. (TASSO Coll.), *Z. Phys.* C30 (1986) 355.
- [11] T. Akesson et al. (AFS Coll.), *Z. Phys.* C36 (1987) 517.
- [12] N.M. Agababyan et al. (NA22 Coll.), *Phys. Lett.* B332 (1994) 458.
- [13] M. Biyajima et al., *Progr. Theor. Phys.* 84 (1990) 931.
- [14] T. Sjöstrand, *Comp. Phys. Comm.* 27 (1982) 243; *ibid.* 28 (1983) 229; T. Sjöstrand, M. Bengtsson, *Comp. Phys. Comm.* 43 (1987) 367.
- [15] A.H. Mueller, *Phys. Rev.* D4 (1971) 150.
- [16] P. Aarnio et al. (DELPHI Coll.), *Nucl. Instr. Methods* A303 (1991) 233.
- [17] P. Abreu et al. (DELPHI Coll.), *Z. Phys.* C63 (1994) 17.
- [18] DELSIM User Manual, DELPHI 87-96 PROG-99; DELSIM Reference Manual, DELPHI 87-97 PROG-100.
- [19] D. Buskulic et al. (ALEPH Coll.), *Phys. Lett.* B292 (1992) 210.
- [20] D. Decamp et al. (ALEPH Coll.), *Z. Phys.* C54 (1992) 75
- [21] P. Abreu et al. (DELPHI Coll.), *Phys. Lett.* B286 (1992) 201
- [22] P.D. Acton et al. (OPAL Coll.), *Phys. Lett.* B267 (1991) 143.
- [23] I. Juricic et al., *Phys. Rev.* D39 (1989) 1.

DELPHI




Figure 1: Q -dependence of R''_3 , the total three-particle correlation, and R'_3 , the three-particle correlation arising from two-body correlations, (a) for like-sign and (b) for unlike-sign triplets.

DELPHI




Fig.2

Figure 2: The three-particle correlation function R_3 for (a) like-sign triplets and (b) for unlike-sign triplets, compared to the JETSET prediction with no Bose-Einstein correlations included. The dot-dashed curve in (a) is the result of a fit with Eq. 10.

DELPHI




Fig.3

Figure 3: The function $R_3(Q)$ for **(a)** like-sign triplets and **(b)** unlike-sign triplets. The predictions of JETSET without BE (dashed line) and with BE correlations included for direct pions (full line) are also shown, using parameters $\lambda = 1$ and $r = 0.50$ fm in the Gaussian parametrization. In both versions of the model η' production was reduced by a factor four.