



Probing small Bjorken- x nuclear gluonic structure via coherent J/ψ photoproduction in ultraperipheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

The CMS Collaboration*

Abstract

Quasi-real photons exchanged in relativistic heavy ion interactions are powerful probes of the gluonic structure of nuclei. The first measurement of the coherent J/ψ photoproduction cross section in ultraperipheral lead-lead collisions is presented over a wide range of photon-nucleus center-of-mass energies per nucleon ($W_{\gamma N}^{\text{Pb}}$). Results are obtained using data at the nucleon-nucleon center-of-mass energy of 5.02 TeV collected by the CMS experiment at the CERN LHC, corresponding to an integrated luminosity of 1.52 nb^{-1} . The cross section is observed to rise rapidly at low $W_{\gamma N}^{\text{Pb}}$, and plateau above $W_{\gamma N}^{\text{Pb}} \approx 40 \text{ GeV}$, up to 400 GeV, center a new regime of small Bjorken- x ($\approx 6 \times 10^{-5}$) gluons being probed in a heavy nucleus. The observed energy dependence is not predicted by current quantum chromodynamic models.

Submitted to Physical Review Letters

The electromagnetic fields surrounding relativistic heavy ions are highly Lorentz contracted and can be effectively treated as linearly polarized quasi-real photons [1, 2]. These photons are powerful and clean probes of the gluonic structure of nuclei [3]. At high energy colliders, when two heavy ions pass each other at an impact parameter greater than twice the nuclear radius, it is referred to as an “ultraperipheral collision” or UPC. In UPCs, a projectile photon from one ion can fluctuate into a virtual quark-antiquark pair and interact with the other target ion via gluon exchanges before turning exclusively into a vector meson (VM) final state [4–17]. If the photon interacts with the nucleus as a whole, the process is known as a coherent interaction, while the incoherent process involves interactions of photons with the nuclear constituents of the target nucleus. At energies accessible at the CERN LHC, studies of VM photoproduction in UPCs open the possibility of probing the poorly known nuclear gluonic structure and dynamics in the low Bjorken- x region ($10^{-5} < x < 10^{-2}$). For instance, the increasingly large gluon density at small x may ultimately reach the unitarity limit, leading to the saturation of the gluon density [18, 19] and thus to a saturation of the observed VM cross section. It has also been suggested that the black-disk limit (BDL) [20–25] of strong absorption may be reached at high energies, where the gluon density itself may not have saturated, but the scattering probability by multiple gluons becomes large enough that the cross section may also saturate at the geometric limit of the nucleus.

The photoproduction of heavy-flavor VMs is of particular interest since it can be studied in the perturbative quantum chromodynamic (QCD) framework with a large energy scale defined by the quark mass. In the leading-order QCD contribution, which is sketched in Fig. 1, the coherent J/ψ photoproduction cross section is proportional to the square of the gluon density function. In photon-proton interactions, the J/ψ production cross section has been extensively measured and is observed to significantly increase with the photon-proton center-of-mass energy [7, 26–30]. This is a consequence of the rapidly rising gluon density in the proton at small x [23, 31]. For photons scattering off heavy-nuclear targets, while significant nuclear suppression effects have been observed [11–17, 32], the underlying mechanism remains poorly understood. Little is known regarding the behavior of the interaction towards the high energy limit, which probes gluons at small x (i.e., $x < 10^{-4}$). The major limitation is the experimental difficulty in distinguishing the photon-emitter nucleus and the target nucleus in a symmetric collision system [33]. For VM photoproduction off nuclei at a given rapidity, there is always a mixture of low and high energy photon contributions, as shown by the opposing sides of Fig. 1. This two-way ambiguity limits the measurements at LHC energies to below 125 GeV in the photon-nucleus center-of-mass energy per nucleon ($W_{\gamma N}^A$).

A novel approach for resolving this two-way ambiguity has been proposed in Refs. [33, 34]. By controlling the UPC impact parameter using forward-emitted neutron multiplicities from the electromagnetic dissociation (EMD) of ions, the relative contributions of low and high energy photons to the measured J/ψ cross section can be separated without ambiguity. The energy exchange in an EMD process is of the order of 10 MeV, which is six orders of magnitude lower than the beam energy.

The results presented in this Letter use PbPb UPCs at a nucleon-nucleon center-of-mass energy $\sqrt{s_{NN}} = 5.02$ TeV, which were collected by the CMS experiment at the LHC in 2018. The data correspond to an integrated luminosity of 1.52 nb^{-1} . By applying the aforementioned forward neutron tagging technique, this Letter reports the cross section measurement of the coherent J/ψ photoproduction off Pb nuclei over a broad energy range of $40 < W_{\gamma N}^{\text{Pb}} < 400$ GeV, which includes previously unexplored small- x regions (i.e., $x < 10^{-4}$). Tabulated results are provided in the HEPData record for this analysis [35].




Figure 1: A pictorial representation of the J/ψ photoproduction process in PbPb at leading-order QCD, showing the origin of the two-way ambiguity. The blue wavy and black helical lines represent emitted photons and gluons, respectively.

The CMS apparatus [36] is a multipurpose, nearly hermetic detector, designed to trigger on [37, 38] and identify electrons, muons, photons, and hadrons [39–41]. A global “particle-flow” algorithm [42] aims to reconstruct all individual particles in an event, combining information provided by the all-silicon inner tracker and by the crystal electromagnetic and brass-scintillator hadron calorimeters, operating inside a 3.8 T superconducting solenoid, with data from the gas-ionization muon detectors embedded in the flux-return yoke outside the solenoid. Forward calorimeters [43], made of steel and quartz-fibres, extend the pseudorapidity coverage provided by the barrel and endcap detectors. Two zero-degree calorimeters (ZDCs) [44], made of quartz fibers and plates embedded in tungsten absorbers, are used to detect neutrons from nuclear dissociation events.

Events are selected using a hardware-based trigger system that requires at least one muon candidate with no explicit selection on its minimum transverse momentum (p_T), coincident with a PbPb bunch crossing [38]. At the trigger level, events with an energy deposit above the noise threshold in either of the two forward calorimeters are vetoed. For the offline analysis, events are required to have a primary interaction vertex, formed by using two or more tracks coming from the collision, that is within 20 cm along the beam axis and at a radius less than 2 cm in the transverse plane from the CMS detector center. To suppress hadronic PbPb collisions, the largest energy deposits in the forward calorimeters are required to be below 7.3 and 7.6 GeV in the positive and negative rapidity sides, respectively. These energy requirements are determined from empty bunch-crossing events [45], providing a purity of nearly 100% with negligible efficiency loss. In addition, events must contain exactly two muon candidates and no additional tracks in the range $|\eta| < 2.4$.

To select high-quality muons, both muons are required to be “soft muons”, as defined in Ref. [46], based on the combined information of the tracker and muon detector systems. Pairs of muons of opposite signs with an invariant mass in the range $2.6 < m_{\mu^+\mu^-} < 4.2$ GeV are taken as J/ψ and $\psi(2S)$ meson candidates. To reconstruct J/ψ mesons at low p_T , the dimuon rapidity must satisfy $1.6 < |y| < 2.4$ for both muons to reach the muon detector systems. For each muon pair, at least one of the muon candidates is required to be matched to the triggered muon. Combinatorial backgrounds are estimated using same-sign muon pairs, and found to be negligible after the event and muon requirements.

The detector reconstruction efficiency is estimated using a Monte Carlo (MC) sample of coher-

ent $J/\psi \rightarrow \mu^+\mu^-$ events generated with the STARLIGHT (v3.13) event generator [47]. These simulated events are further processed with the full CMS detector response simulation using GEANT4 [48].

The raw signal yields of J/ψ and $\psi(2S)$ mesons, $N_{J/\psi}$ and $N_{\psi(2S)}$, are extracted by fitting the dimuon invariant mass distribution for a given rapidity bin with two Crystal Ball (CB) functions [49] to describe the J/ψ and $\psi(2S)$ resonances, and a third-order polynomial function to parametrize the underlying background from quantum electrodynamic processes ($\gamma\gamma \rightarrow \mu^+\mu^-$) (see Appendix A). The raw number of J/ψ signal events contains contributions from several physics processes: coherent and incoherent (with or without nucleon dissociation) J/ψ production from direct photon-nucleus interactions, and also J/ψ mesons coming from the decay of coherent and incoherent $\psi(2S)$ production, referred to as “feed-down”. Coherent and incoherent productions each have a distinct characteristic p_T spectra, so their individual contributions can be disentangled using a template fit to the measured J/ψ p_T distribution, as done in Refs. [13, 15]. The feed-down contribution is estimated directly by measuring the $\psi(2S)$ signal yield from the fit to the dimuon invariant mass distributions. Details on the coherent J/ψ yield ($N_{J/\psi}^{\text{coh}}$) extraction for each rapidity bin, including corrections for the incoherent production and feed-down contribution, are provided in Appendix A.

The final corrected coherent J/ψ production differential cross section is determined using

$$\frac{d\sigma_{J/\psi}^{\text{coh}}}{dy} = \frac{N_{J/\psi}^{\text{coh}}}{\epsilon_{J/\psi} \mathcal{B}_{J/\psi \rightarrow \mu\mu} \mathcal{L}_{\text{int}} \epsilon_{\text{evtsel}} \Delta y'} \quad (1)$$

where the event selection efficiency of $\epsilon_{\text{evtsel}} = 94.2\%$ and the J/ψ reconstruction efficiency $\epsilon_{J/\psi}$, ranging from 2.6 to 11.9% for $1.6 < |y| < 2.4$, are evaluated using simulated samples. Additional corrections determined using the tag-and-probe technique [46, 50] are applied to account for any data-to-simulation discrepancy. The world-average value for the J/ψ dimuon branching fraction [51] $\mathcal{B}_{J/\psi \rightarrow \mu^+\mu^-}$ is used. The quantity \mathcal{L}_{int} is the total integrated luminosity of the data sample [52], and Δy is the rapidity binning used in the measurement.

Coherent J/ψ photoproduction cross sections are reported in classes of the forward-neutron multiplicity, which is determined by the energies deposited in the ZDCs, as done in Ref. [45]. Based on the neutron peaks observed in the total ZDC energy distribution, events are classified as having no neutrons (0n) or with at least one neutron (Xn, $X \geq 1$) in each ZDC. The purities of 0n and Xn classes are estimated using a multi-Gaussian function fit to the ZDC energy distribution, and are found to be better than 99.6% [45]. A total of three neutron multiplicity event classes are formed by combining the classifications of each ZDC side, labeled as 0n0n, 0nXn, and XnXn. Results without neutron multiplicity selection, labeled as AnAn, are also reported for comparison and complementarity with previous measurements. Except for the AnAn case, a correction is made for the effect of having an additional EMD in the event. This correction is obtained by evaluating the migration probability among neutron multiplicity classes, following the approach of Ref. [45]. An average change of +11, -30, and -29% in the raw J/ψ yields is estimated for the 0n0n, 0nXn, and XnXn event classes, respectively.

Because of the symmetry of the PbPb collision system, a J/ψ meson measured at rapidity y can result from two possible photon energies: $\omega_1 = (M_{J/\psi}/2) \exp(-y)$ and $\omega_2 = (M_{J/\psi}/2) \exp(+y)$. The x of the associated parton and $W_{\gamma N}^{\text{Pb}}$ with a photon of energy ω_1 can be found using $x = (M_{J/\psi}/\sqrt{s_{\text{NN}}}) \exp(+y)$ and $W_{\gamma N}^{\text{Pb}} = \sqrt{\sqrt{s_{\text{NN}}} M_{J/\psi} \exp(-y)}$. The corresponding values for a photon of energy ω_2 are obtained by changing the sign of y in each expression.

The measured differential cross section at a given rapidity y for a given neutron multiplicity

class, $injn$ ($= 0n0n, 0nXn, XnXn$), of UPC events is the sum of both photon energy contributions [33, 34]:

$$\frac{d\sigma_{J/\psi}^{injn}(y)}{dy} = n_{\gamma A}^{injn}(\omega_1) \sigma_{J/\psi}(\omega_1) + n_{\gamma A}^{injn}(\omega_2) \sigma_{J/\psi}(\omega_2), \quad (2)$$

where it should be kept in mind that ω_1 and ω_2 are themselves functions of y . Here, $n_{\gamma A}(\omega) = \omega dN_{\gamma A}(\omega)/d\omega$ is the photon flux and $\sigma_{J/\psi}(\omega)$ is the coherent J/ψ photoproduction cross section for a single γ -ion interaction, averaged over the given y range. The photon flux from a relativistic ion is determined with high precision using the equivalent approximation model as implemented with the STARLIGHT event generator [47]. The flux strongly depends on the ion-impact parameter; the high energy photon flux is enhanced relative to that of low energy photons at smaller impact parameters and for higher neutron multiplicities. Therefore, after simultaneously measuring the J/ψ differential coherent cross section as a function of rapidity (i.e., the left-hand side of Eq. (2)) and computing the photon fluxes in each of the three neutron classes, a combined χ^2 fit is performed to extract the differential cross sections for low and high energies, $\sigma_{J/\psi}(\omega_1)$ and $\sigma_{J/\psi}(\omega_2)$.

Systematic uncertainties are evaluated by taking the maximum variation from the nominal result for a given source. The uncertainties are estimated to be 1.5% in the integrated luminosity [52, 53], and 0.55% in $\mathcal{B}_{J/\psi \rightarrow \mu^+ \mu^-}$ [51]. Uncertainties in the tag-and-probe corrections are determined to be 3–8% from low to high $W_{\gamma N}^{Pb}$ values. The choice of the forward calorimeter threshold energy in rejecting hadronic contamination leads to 1–4% uncertainties found by modifying the limits and noting the change in the results. For the systematic uncertainty in the raw signal yield extraction, different functional forms for modeling the shapes of the signal and background are studied, including: (i) a sum of CB and Gaussian functions for the J/ψ signal shape; (ii) a second-order polynomial for the dimuon continuum background; (iii) fixing the CB shape parameters to the values determined from MC signal distributions; (iv) varying the range of the dimuon invariant mass distribution used in the fit; (v) changing the dimuon continuum p_T shape to the one determined from side-band mass regions. The results vary by less than 8% when these changes are made. The uncertainty in the neutron migration correction is estimated to be less than 2% by comparing to an alternative approach following Ref. [54], using the observed rate of single- and double-sided neutron emissions per bunch crossing. Individual sources of experimental uncertainties are added in quadrature to obtain the total systematic uncertainty of 6–12%. Theoretical uncertainties in the photon flux calculation are separately evaluated to be 3–16% by varying the Pb nuclei radius ($R = 6.67 \pm 0.03$ fm), the nuclear skin thickness ($a = 0.56 \pm 0.03$ fm) [55], and the EMD cross section [56] within the experimental uncertainties, where uncertainties at low and high $W_{\gamma N}^{Pb}$ are found to be strongly anticorrelated. Correlations between uncertainties are evaluated as a function of $W_{\gamma N}^{Pb}$ and included in the covariance matrix [35] of the fit.

The resulting measured coherent J/ψ photoproduction differential cross sections over the rapidity range $1.6 < |y| < 2.4$ are reported in Fig. 2 (left) in three neutron multiplicity classes. The results with no neutron selection are given in the right plot of Fig. 2. The measurements are performed as a function of $|y|$, but are shown at negative y values, following the ALICE convention [13, 15]. Predictions from several theoretical calculations are also shown in both plots for comparison. The leading twist approximation (LTA) [34] is a perturbative QCD calculation that takes into account nuclear shadowing effects from multi-nucleon interference. Both weak (WS) and strong (SS) shadowing scenarios are shown [34]. The color dipole (CD) models, with different model parameters (BGK, BGW, IIM), assume quark-antiquark dipole scattering from the nuclear targets [57].




Figure 2: The differential coherent J/ψ photoproduction cross section as a function of rapidity, in different neutron multiplicity classes (left): $0n0n$, $0nXn$ and $XnXn$; (right): $AnAn$. The small vertical bars and shaded boxes represent the statistical and systematic uncertainties, respectively. The horizontal bars show the bin widths. Theoretical predictions from LTA weak/strong shadowing [34], color dipole models (CD_BGK, CD_BGW, and CD_IIM) [57], and STARLIGHT [47] are shown by the curves. The right plot also displays the results from the ALICE [13, 15] and LHCb [17] experiments.

For the case of no neutron selection ($AnAn$), the data follow the trend of the forward-rapidity measurements from ALICE [13] over a new y region. None of the models describe the combined results over the full rapidity range. The color dipole models agree with the measurements in the forward-rapidity region, but fail to describe the data at $y \approx 0$. In each neutron multiplicity class, the LTA predictions tend to be lower than the CMS results, particularly for the strong shadowing scenario. These comparisons indicate that there are key ingredients missing from the theoretical understanding of high energy photon-nucleus scattering processes.

To gain further insight, the total measured J/ψ coherent photoproduction cross section as a function of $W_{\gamma N}^{Pb}$ up to ≈ 400 GeV is shown in Fig. 3, after decomposing the two-way ambiguity. Because the contributions of high energy photons are negligible at very forward rapidity (less than 5% for $-4.5 < y < -3.5$) [33, 34], and the fact that $y \approx 0$, $\omega_1 \approx \omega_2 \approx M_{J/\psi}/2$, the total cross section at lower $W_{\gamma N}^{Pb}$ values can be approximated using ALICE and LHCb measurements. These results are also shown in Fig. 3. The experimental and theoretical (from the photon flux) uncertainties are displayed separately in Fig. 3. Predictions from the LTA and CD models, as well as the gluon saturation models bBK [58], IPsat [59], and GG [60]), are compared to the experimental measurements. The naive prediction from the impulse approximation (IA) model [33] is also shown, providing a reference assuming the absence of any nuclear effects except for coherence.

The measured total cross section has an unexpected energy dependence, rising by more than a factor of 4 as $W_{\gamma N}^{Pb}$ goes from $W_{\gamma N}^{Pb} \approx 15$ to 40 GeV. This is consistent with the expectation of a fast-growing gluon density at low x (e.g., from the IA model). However, this trend vanishes for $W_{\gamma N}^{Pb} > 40$ GeV, and instead the total cross section begins a slow linear rise with a slope of $(3.0 \pm 0.4 \text{ (stat)} \pm 1.4 \text{ (syst)}) \times 10^{-5} \text{ mb/GeV}$ up to $W_{\gamma N}^{Pb} \approx 400$ GeV. Considering the experimental uncertainties across the measured $W_{\gamma N}^{Pb}$ range, none of the theoretical models are consistent with the measurements, with the CD-BGK model having the best p -value of 1.6×10^{-8} . This result could imply the onset of novel physics mechanisms in the coherent J/ψ photoproduction process starting at $W_{\gamma N}^{Pb} \approx 40$ GeV, for example, the saturation of the gluon density in the Pb nucleus at the corresponding x value. This picture may be less straightforward due to higher-order perturbative QCD corrections (e.g., quark-antiquark exchange) as suggested in




Figure 3: The total coherent J/ψ photoproduction cross section as a function of $W_{\gamma N}^{\text{Pb}}$ from the CMS measurement in PbPb UPCs at $\sqrt{s_{\text{NN}}} = 5.02$ TeV. Approximated results (implied by the asterisk) from the ALICE [13, 15] and LHCb [17] experiments are displayed for specific rapidity regions, where the two-way ambiguity effect is expected to be negligible. The $W_{\gamma N}^{\text{Pb}}$ values used correspond to the center of each experiment's rapidity range. The vertical bars and the shaded and open boxes represent the statistical, experimental, and theoretical (photon flux) uncertainties, respectively. The predictions from various theoretical calculations [33, 34, 57–60] are shown by the curves.

Refs. [61, 62]. On the other hand, the observed behavior is also expected when approaching the BDL, where major part of the target Pb nucleus becomes completely absorptive to projectile photons because of the large scattering probability in the presence of dense gluons at small x [21]. In this scenario, the photon-nucleus cross section approaches the the unitarity limit allowed by the geometric size of the nucleus. The slow rise in the measured cross section from about 40 to 400 GeV would then be due to periphery of the nucleus not becoming fully “black.”

To quantify the nuclear effects on the observed gluon density function in a Pb nucleus, a nuclear gluon suppression factor, $R_g^{\text{Pb}}(x, \mu^2 = 2.4 \text{ GeV}^2)$ as a function of x and with $\mu = M_{J/\psi}/2$, where μ is the energy scale, is defined as the square root of the ratio of the measured coherent J/ψ photoproduction cross section σ^{Meas} to the IA prediction σ^{IA} , $R_g^{\text{Pb}} = \sqrt{\sigma^{\text{Meas}}/\sigma^{\text{IA}}}$ [63]. The extracted R_g^{Pb} values are shown in Fig. 4 as a function of x . The suppression in the relatively high- x (low $W_{\gamma N}^{\text{Pb}}$) region of $x > 5 \times 10^{-3}$ is moderate, approximately 0.8–0.9. Moving toward smaller x values, R_g^{Pb} starts dropping rapidly to 0.4–0.5 for $x \approx 6 \times 10^{-5}$. Similar to the cross section measurement, no theoretical model predicts the observed values and x dependence of R_g^{Pb} over the wide x range reported. If the BDL of strong absorption is reached for $x < 5 \times 10^{-3}$,

corresponding to $W_{\gamma N}^{\text{Pb}} > 40 \text{ GeV}$, the direct connection between the measured cross section and the single gluon density function, as in the weak absorption limit, no longer holds. Instead, the cross section will reach the unitarity limit while the gluon density can continue increasing. Therefore, new theoretical approaches are needed to understand the structure and dynamics of the strong force for a heavy nucleus in this new domain of extreme gluon density.




Figure 4: The nuclear gluon suppression factor R_g^{Pb} as a function of Bjorken x extracted from the CMS measurement of the coherent J/ψ photoproduction in PbPb UPCs at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$. Approximated results (implied by the asterisk) from the ALICE [13, 15] and LHCb [17] experiments are displayed for specific rapidity regions, where the two-way ambiguity effect is expected to be negligible. A measurement of J/ψ photoproduction from the E691 experiment [64, 65] is also given. The x values are evaluated at the centers of their corresponding rapidity ranges. The vertical bars and shaded and open boxes represent the statistical, experimental systematic, and theoretical systematic uncertainties, respectively. The latter is due to the uncertainties in the photon flux and the IA. Predictions from various theoretical calculations [34, 57, 58, 60] are shown by the curves.

In summary, the first measurement of the coherent J/ψ photoproduction cross section off lead nuclei, as a function of the photon-nuclear center-of-mass energy per nucleon ($W_{\gamma N}^{\text{Pb}}$) has been presented over a broad energy range. In a coherent process, the J/ψ is produced by the photon interacting with the nucleus as a whole. Results are obtained with ultraperipheral PbPb collision data by applying the forward neutron tagging technique. The cross section is observed to rise rapidly at low $W_{\gamma N}^{\text{Pb}}$ but appears to plateau above $W_{\gamma N}^{\text{Pb}} \approx 40 \text{ GeV}$, up to 400 GeV , where a new regime of gluon momentum fraction (Bjorken $x \approx 6 \times 10^{-5}$) in a heavy nucleus is probed. This observed trend is not predicted by current theoretical models. This can be interpreted

either as the first direct evidence for the saturation of nuclear gluonic density or that the scattering cross section is near the black-disk limit.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); Minciencias (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

References

- [1] E. J. Williams, "Nature of the high energy particles of penetrating radiation and status of ionization and radiation formulae", *Phys. Rev.* **45** (1934) 729, doi:10.1103/PhysRev.45.729.
- [2] C. F. von Weizsacker, "Radiation emitted in collisions of very fast electrons", *Z. Phys.* **88** (1934) 612, doi:10.1007/BF01333110.
- [3] S. R. Klein and P. Steinberg, "Photonuclear and two-photon interactions at high-energy nuclear colliders", *Annu. Rev. Nucl. Part. Sci.* **70** (2020) 323, doi:10.1146/annurev-nucl-030320-033923, arXiv:2005.01872.
- [4] STAR Collaboration, "Coherent ρ^0 production in ultraperipheral heavy ion collisions", *Phys. Rev. Lett.* **89** (2002) 272302, doi:10.1103/PhysRevLett.89.272302, arXiv:nucl-ex/0206004.
- [5] STAR Collaboration, "Coherent diffractive photoproduction of ρ^0 mesons on gold nuclei at 200 GeV/nucleon-pair at the Relativistic Heavy Ion Collider", *Phys. Rev. C* **96** (2017) 054904, doi:10.1103/PhysRevC.96.054904, arXiv:1702.07705.
- [6] PHENIX Collaboration, "Photoproduction of J/ψ and of high mass e^+e^- in ultra-peripheral Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV", *Phys. Lett. B* **679** (2009) 321, doi:10.1016/j.physletb.2009.07.061, arXiv:0903.2041.

- [7] ALICE Collaboration, “Exclusive J/ψ photoproduction off protons in ultra-peripheral p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Phys. Rev. Lett.* **113** (2014) 232504, doi:10.1103/PhysRevLett.113.232504, arXiv:1406.7819.
- [8] CMS Collaboration, “Measurement of exclusive Y photoproduction from protons in pPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Eur. Phys. J. C* **79** (2019) 277, doi:10.1140/epjc/s10052-019-6774-8, arXiv:1809.11080.
- [9] CMS Collaboration, “Measurement of exclusive $\rho(770)^0$ photoproduction in ultraperipheral pPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Eur. Phys. J. C* **79** (2019) 702, doi:10.1140/epjc/s10052-019-7202-9, arXiv:1902.01339.
- [10] STAR Collaboration, “Tomography of ultrarelativistic nuclei with polarized photon-gluon collisions”, *Sci. Adv.* **9** (2023) eabq3903, doi:10.1126/sciadv.abq3903, arXiv:2204.01625.
- [11] ALICE Collaboration, “Coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV”, *Phys. Lett. B* **718** (2013) 1273, doi:10.1016/j.physletb.2012.11.059, arXiv:1209.3715.
- [12] ALICE Collaboration, “Charmonium and e^+e^- pair photoproduction at mid-rapidity in ultra-peripheral PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV”, *Eur. Phys. J. C* **73** (2013) 2617, doi:10.1140/epjc/s10052-013-2617-1, arXiv:1305.1467.
- [13] ALICE Collaboration, “Coherent J/ψ photoproduction at forward rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Phys. Lett. B* **798** (2019) 134926, doi:10.1016/j.physletb.2019.134926, arXiv:1904.06272.
- [14] CMS Collaboration, “Coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV with the CMS experiment”, *Phys. Lett. B* **772** (2017) 489, doi:10.1016/j.physletb.2017.07.001, arXiv:1605.06966.
- [15] ALICE Collaboration, “Coherent J/ψ and $\psi(2S)$ photoproduction at midrapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Eur. Phys. J. C* **81** (2021) 712, doi:10.1140/epjc/s10052-021-09437-6, arXiv:2101.04577.
- [16] LHCb Collaboration, “Study of coherent J/ψ production in lead-lead collisions at $\sqrt{s_{\text{NN}}} = 5$ TeV”, *JHEP* **07** (2022) 117, doi:10.1007/JHEP07(2022)117, arXiv:2107.03223.
- [17] LHCb Collaboration, “Study of coherent charmonium production in ultra-peripheral lead-lead collisions”, 2022. arXiv:2206.08221.
- [18] E. Iancu and R. Venugopalan, “The color glass condensate and high-energy scattering in QCD”, p. 249. 2003. arXiv:hep-ph/0303204. doi:10.1142/9789812795533_0005.
- [19] F. Gelis, E. Iancu, J. Jalilian-Marian, and R. Venugopalan, “The color glass condensate”, *Ann. Rev. Nucl. Part. Sci.* **60** (2010) 463, doi:10.1146/annurev.nucl.010909.083629, arXiv:1002.0333.
- [20] V. N. Gribov, “Interaction of gamma quanta and electrons with nuclei at high-energies”, *Zh. Eksp. Teor. Fiz.* **57** (1969) 1306.

-
- [21] L. Frankfurt, V. Guzey, M. McDermott, and M. Strikman, “Revealing the black body regime of small- x DIS through final state signals”, *Phys. Rev. Lett.* **87** (2001) 192301, doi:10.1103/PhysRevLett.87.192301, arXiv:hep-ph/0104154.
- [22] L. Frankfurt, M. Strikman, and M. Zhalov, “Signals for black body limit in coherent ultraperipheral heavy ion collisions”, *Phys. Lett. B* **537** (2002) 51, doi:10.1016/S0370-2693(02)01882-8, arXiv:hep-ph/0204175.
- [23] L. Frankfurt, M. Strikman, and C. Weiss, “Small- x physics: From HERA to LHC and beyond”, *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 403, doi:10.1146/annurev.nucl.53.041002.110615, arXiv:hep-ph/0507286.
- [24] T. Rogers, X. Zu, V. Guzey, and M. Strikman, “Determining the proximity of γN scattering to the black body limit using deep inelastic scattering and J/ψ production”, *Phys. Rev. D* **69** (2004) 074011, doi:10.1103/PhysRevD.69.074011, arXiv:hep-ph/0309099.
- [25] A. Alkin, E. Martynov, O. Kovalenko, and S. M. Troshin, “Impact-parameter analysis of TOTEM data at the LHC: Black disk limit exceeded”, *Phys. Rev. D* **89** (2014) 091501(R), doi:10.1103/PhysRevD.89.091501, arXiv:1403.8036.
- [26] ZEUS Collaboration, “Exclusive photoproduction of J/ψ mesons at HERA”, *Eur. Phys. J. C* **24** (2002) 345, doi:10.1007/s10052-002-0953-7, arXiv:hep-ex/0201043.
- [27] H1 Collaboration, “Elastic J/ψ production at HERA”, *Eur. Phys. J. C* **46** (2006) 585, doi:10.1140/epjc/s2006-02519-5, arXiv:hep-ex/0510016.
- [28] H1 Collaboration, “Elastic and proton-dissociative photoproduction of J/ψ mesons at HERA”, *Eur. Phys. J. C* **73** (2013) 2466, doi:10.1140/epjc/s10052-013-2466-y, arXiv:1304.5162.
- [29] LHCb Collaboration, “Exclusive J/ψ and $\psi(2S)$ production in pp collisions at $\sqrt{s} = 7$ TeV”, *J. Phys. G* **40** (2013) 045001, doi:10.1088/0954-3899/40/4/045001, arXiv:1301.7084.
- [30] ALICE Collaboration, “Energy dependence of exclusive J/ψ photoproduction off protons in ultra-peripheral p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Eur. Phys. J. C* **79** (2019) 402, doi:10.1140/epjc/s10052-019-6816-2, arXiv:1809.03235.
- [31] J. D. Bjorken, “Asymptotic sum rules at infinite momentum”, *Phys. Rev.* **179** (1969) 1547, doi:10.1103/PhysRev.179.1547.
- [32] L. Favart, M. Guidal, T. Horn, and P. Kroll, “Deeply virtual meson production on the nucleon”, *Eur. Phys. J. A* **52** (2016) 158, doi:10.1140/epja/i2016-16158-2, arXiv:1511.04535.
- [33] V. Guzey, M. Strikman, and M. Zhalov, “Disentangling coherent and incoherent quasielastic J/ψ photoproduction on nuclei by neutron tagging in ultraperipheral ion collisions at the LHC”, *Eur. Phys. J. C* **74** (2014) 2942, doi:10.1140/epjc/s10052-014-2942-z, arXiv:1312.6486.
- [34] V. Guzey, E. Kryshen, and M. Zhalov, “Coherent photoproduction of vector mesons in ultraperipheral heavy ion collisions: Update for run 2 at the CERN Large Hadron Collider”, *Phys. Rev. C* **93** (2016) 055206, doi:10.1103/PhysRevC.93.055206, arXiv:1602.01456.

- [35] CMS Collaboration, “HEPData record for this analysis”, 2023.
doi:10.17182/hepdata.138867.
- [36] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004,
doi:10.1088/1748-0221/3/08/S08004.
- [37] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [38] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020,
doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [39] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014,
doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [40] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015,
doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [41] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009,
doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [42] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [43] G. Bayatian et al., “Design, performance and calibration of the CMS forward calorimeter wedges”, *Eur. Phys. J. C* **53** (2008) 139, doi:10.1140/epjc/s10052-007-0459-4.
- [44] O. Surányi et al., “Performance of the CMS zero degree calorimeters in pPb collisions at the LHC”, *JINST* **16** (2021) P05008, doi:10.1088/1748-0221/16/05/P05008, arXiv:2102.06640.
- [45] CMS Collaboration, “Observation of forward neutron multiplicity dependence of dimuon acoplanarity in ultraperipheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Rev. Lett.* **127** (2021) 122001, doi:10.1103/PhysRevLett.127.122001, arXiv:2011.05239.
- [46] CMS Collaboration, “Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV”, *JINST* **7** (2012) P10002, doi:10.1088/1748-0221/7/10/P10002, arXiv:1206.4071.
- [47] S. R. Klein et al., “STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions”, *Comput. Phys. Commun.* **212** (2017) 258,
doi:10.1016/j.cpc.2016.10.016, arXiv:1607.03838.
- [48] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [49] Oreglia, “A study of the reaction $\psi(2S) \rightarrow \gamma\gamma/\psi$ ”, *Phys. Rev. D* **25** (1982) 2259,
doi:10.1103/PhysRevD.25.2259.

- [50] CMS Collaboration, “Measurement of the inclusive W and Z production cross sections in pp collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **10** (2011) 132, doi:10.1007/JHEP10(2011)132, arXiv:1107.4789.
- [51] Particle Data Group Collaboration, “Review of particle physics”, *Prog. Theor. Exp. Phys.* **2022** (2022) 083C01, doi:10.1093/ptep/ptac097.
- [52] CMS Collaboration, “CMS luminosity measurement using nucleus-nucleus collisions at $\sqrt{s_{NN}} = 5.02$ TeV in 2018”, Technical Report CMS-PAS-LUM-18-001, 2022. CMS Physics Analysis Summary.
- [53] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [54] ATLAS Collaboration, “Exclusive dimuon production in ultraperipheral Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ATLAS”, *Phys. Rev. C* **104** (2021) 024906, doi:10.1103/PhysRevC.104.024906, arXiv:2011.12211.
- [55] D. d’Enterria and C. Loizides, “Progress in the Glauber model at collider energies”, *Ann. Rev. Nucl. Part. Sci.* **71** (2021) 315, doi:10.1146/annurev-nucl-102419-060007, arXiv:2011.14909.
- [56] A. J. Baltz, M. J. Rhoades-Brown, and J. Weneser, “Heavy ion partial beam lifetimes due to coulomb induced processes”, *Phys. Rev. E* **54** (1996) 4233, doi:10.1103/PhysRevE.54.4233.
- [57] A. Łuszczak and W. Schäfer, “Coherent photoproduction of J/ψ in nucleus-nucleus collisions in the color dipole approach”, *Phys. Rev. C* **99** (2019) 044905, doi:10.1103/PhysRevC.99.044905, arXiv:1901.07989.
- [58] D. Bendova, J. Cepila, J. G. Contreras, and M. Matas, “Photonuclear J/ψ production at the LHC: proton-based versus nuclear dipole scattering amplitudes”, *Phys. Lett. B* **817** (2021) 136306, doi:10.1016/j.physletb.2021.136306, arXiv:2006.12980.
- [59] H. Mäntysaari and B. Schenke, “Probing subnucleon scale fluctuations in ultraperipheral heavy ion collisions”, *Phys. Lett. B* **772** (2017) 832, doi:10.1016/j.physletb.2017.07.063, arXiv:1703.09256.
- [60] J. Cepila, J. G. Contreras, and M. Krelina, “Coherent and incoherent J/ψ photonuclear production in an energy-dependent hot-spot model”, *Phys. Rev. C* **97** (2018) 024901, doi:10.1103/PhysRevC.97.024901, arXiv:1711.01855.
- [61] K. J. Eskola et al., “Exclusive J/ψ photoproduction in ultraperipheral Pb+Pb collisions at the CERN Large Hadron Collider calculated at next-to-leading order perturbative QCD”, *Phys. Rev. C* **106** (2022) 035202, doi:10.1103/PhysRevC.106.035202, arXiv:2203.11613.
- [62] K. J. Eskola et al., “Next-to-leading order perturbative QCD predictions for exclusive J/ψ photoproduction in oxygen-oxygen and lead-lead collisions at the LHC”, 10, 2022. arXiv:2210.16048.

- [63] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Evidence for nuclear gluon shadowing from the ALICE measurements of PbPb ultraperipheral exclusive J/ψ production”, *Phys. Lett. B* **726** (2013) 290, doi:10.1016/j.physletb.2013.08.043, arXiv:1305.1724.
- [64] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Nuclear suppression from coherent J/ψ photoproduction at the Large Hadron Collider”, *Phys. Lett. B* **816** (2021) 136202, doi:10.1016/j.physletb.2021.136202, arXiv:2008.10891.
- [65] Fermilab Tagged Photon Spectrometer Collaboration, “An experimental study of the A-dependence of J/ψ photoproduction”, *Phys. Rev. Lett.* **57** (1986) 3003, doi:10.1103/PhysRevLett.57.3003.

A Method of coherent J/ψ yield extraction

To extract the J/ψ yield from coherent photoproduction, a two-step analysis is performed by fitting first the dimuon invariant mass distribution at $p_T < 0.2$ GeV, and then the dimuon p_T distribution within the mass window of $2.95 < m_{\mu^+\mu^-} < 3.25$ GeV.




Figure 5: The invariant mass distribution of $\mu^+\mu^-$ pairs with dimuon $p_T < 0.2$ GeV from the signal events with no neutron multiplicity selection (AnAn) (left). The results of the fit are shown by the various curves, and the corresponding values of the χ^2 / degrees of freedom (ndf) and the fitted number of J/ψ and $\psi(2S)$ events are given. The values of the factors R_N , R , and f_D , defined below, are also shown. The transverse momentum distribution of $\mu^+\mu^-$ events with a dimuon invariant mass in the J/ψ mass window of ($2.95 < m_{\mu^+\mu^-} < 3.25$ GeV) (right). Again, the results of the fit are shown by the various curves. In both plots, the vertical bars on the data points give the statistical uncertainty.

An example of an invariant mass distribution for the reconstructed dimuon pairs is shown for events with a dimuon $p_T < 0.2$ GeV and no neutron multiplicity selection in Fig. 5 (left). The parameters of the CB functions left free in the fit include the pole mass $\mu_{J/\psi}$ and width $\Gamma_{J/\psi}$ of the J/ψ . The analogous parameters for the $\psi(2S)$ are constrained by demanding $\mu_{\psi(2S)} = \mu_{J/\psi} M_{\psi(2S)} / M_{J/\psi}$ and $\Gamma_{\psi(2S)} = \Gamma_{J/\psi} M_{\psi(2S)} / M_{J/\psi}$, respectively, where $M_{J/\psi}$ and $M_{\psi(2S)}$ are the world-average mass values [51]. The results of the fit are shown by the various curves, with the J/ψ and $\psi(2S)$ signals being clearly visible. The raw yields of J/ψ and $\psi(2S)$ events, $N_{J/\psi}$ and $N_{\psi(2S)}$, are given in the figure.

Figure 5 (right) shows an example of the $\mu^+\mu^-$ p_T spectrum for events with a dimuon invariant mass in the J/ψ mass window ($2.95 < m_{\mu^+\mu^-} < 3.25$ GeV). Because the different physics

processes involved have distinct shapes, their individual contributions can be measured via template fits. The template p_T spectra for coherent J/ψ production, feed-down J/ψ from coherently photoproduced $\psi(2S)$ decays, incoherent J/ψ production without nucleon dissociation, and the dimuon continuum from quantum electrodynamic processes ($\gamma\gamma \rightarrow \mu^+\mu^-$) are obtained from simulated events with the STARLIGHT event generator [47] and the GEANT4 [48] simulation of the CMS detector. The predicted slope of the p_T distribution for coherently photoproduced J/ψ particles from STARLIGHT is adjusted by varying the nuclear radius parameter to best describe the data p_T distribution.

To describe incoherent J/ψ p_T spectra with nucleon dissociation, an empirical function employed by the H1 [28] and ALICE [13, 15] experiments is used: $dN/dp_T \approx p_T [1 + (b_{\text{pd}}/n_{\text{pd}})p_T^2]^{-n_{\text{pd}}}$, where b_{pd} and n_{pd} are fit parameters. Normalizations of the coherent J/ψ , incoherent J/ψ without nucleon dissociation, and incoherent J/ψ with nucleon dissociation components are free parameters in the fit. The normalization of the dimuon continuum is fixed to the value determined from the invariant mass fitting, as shown in Fig. 5 (left). Normalizations of coherent and incoherent $\psi(2S)$ feed-down to J/ψ are constrained to those of the prompt coherent and incoherent J/ψ components according to the feed-down fractions (f_D) extracted from $\psi(2S)$ yields in the invariant mass distribution, as described below. The entire procedure is repeated for each neutron multiplicity class.

The measured p_T spectrum in the range $0.2 < p_T < 0.5 \text{ GeV}$ is not well described by the fit, as was also seen in previous ALICE studies [13, 15]. This discrepancy is not unexpected since the STARLIGHT event generator does not precisely describe all the physics processes involved. To estimate the systematic uncertainty due to this effect, the p_T upper limit is changed to $p_T < 0.3 \text{ GeV}$ and the extraction of the coherent cross section is repeated, leading to an uncertainty of less than 2%.

The final raw coherent J/ψ yield, $N_{J/\psi}^{\text{coh}}$, is evaluated by:

$$N_{J/\psi}^{\text{coh}} = \frac{N_{J/\psi}}{1 + f_I + f_D}, \quad (3)$$

where f_I and f_D are the fractional contamination of incoherent processes and coherent $\psi(2S)$ feed-down, respectively. The incoherent contamination, f_I , is obtained by fitting the p_T distribution and calculating the fraction of incoherent J/ψ events in the $p_T < 0.2 \text{ GeV}$ region. The feed-down contamination, f_D , is estimated based on the ratio of the number of $\psi(2S)$ to J/ψ events, $R_N = N_{\psi(2S)}/N_{J/\psi}$, found from the fit to the dimuon invariant mass distribution. This parameter can be expressed as

$$R_N = \frac{\sigma_{\psi(2S)} \mathcal{B}_{\psi(2S) \rightarrow \mu\mu} \epsilon_{\psi(2S)}}{(\sigma_{J/\psi} \epsilon_{J/\psi} + \sigma_{\psi(2S)} \mathcal{B}_{\psi(2S) \rightarrow J/\psi} \epsilon_{\psi(2S) \rightarrow J/\psi}) \mathcal{B}_{J/\psi \rightarrow \mu\mu}}. \quad (4)$$

The symbols ϵ and \mathcal{B} represent the two reconstruction efficiencies and decay branching fractions, respectively.

The ratio of the cross sections of coherent $\psi(2S)$ to coherent J/ψ production, $R = \sigma_{\psi(2S)}/\sigma_{J/\psi}$, can then be expressed in terms of R_N as

$$R = \frac{R_N \mathcal{B}_{J/\psi \rightarrow \mu\mu} \epsilon_{J/\psi}}{\mathcal{B}_{\psi(2S) \rightarrow \mu\mu} \epsilon_{\psi(2S)} - R_N \mathcal{B}_{\psi(2S) \rightarrow J/\psi} \epsilon_{\psi(2S) \rightarrow J/\psi} \mathcal{B}_{J/\psi \rightarrow \mu\mu}}. \quad (5)$$

Finally, the fraction of the coherent J/ψ yield that comes from the feed-down of coherently

photoproduced $\psi(2S)$ is found using the expression

$$f_D = R \frac{\epsilon_{\psi(2S) \rightarrow J/\psi}}{\epsilon_{J/\psi}} \mathcal{B}_{\psi(2S) \rightarrow J/\psi}. \quad (6)$$

The world-average values [51] for the J/ψ and $\psi(2S)$ branching fractions are used in these equations.


The resulting measured values of the incoherent and feed-down fractions and their uncertainties from the fits are given in Table 1 for each of the neutron multiplicity class.

Table 1: The measured values of the incoherent and feed-down fractions (f_I) and (f_D), respectively, and their uncertainties for each of the neutron multiplicity class.




















	0n0n	0nXn	XnXn	AnAn
f_I	0.005 ± 0.001	0.103 ± 0.005	0.090 ± 0.011	0.035 ± 0.003
f_D	0.028 ± 0.005	0.037 ± 0.007	0.050 ± 0.012	0.032 ± 0.004

B The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Tumasyan¹ 













Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , A. Escalante Del Valle , P.S. Hussain , M. Jeitler² , N. Krammer , L. Lechner , D. Liko , I. Mikulec , J. Schieck² , R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ , W. Waltenberger , C.-E. Wulz² 














Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish³ , T. Janssen , T. Kello⁴, P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

E.S. Bols , J. D'Hondt , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier , W. Van Doninck, S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Clerbaux , S. Dansana , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , K. Lee , M. Mahdavikhorrani , I. Makarenko , A. Malara , S. Paredes , L. Pétré, N. Postiau, L. Thomas , M. Vanden Bemden, C. Vander Velde , P. Vanlaer 






Ghent University, Ghent, Belgium

D. Dobur , J. Knolle , L. Lambrecht , G. Mestdach, C. Rendón, A. Samalan, K. Skovpen , M. Tytgat , N. Van Den Bossche , B. Vermassen, L. Wezenbeek 





















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , G. Bruno , F. Bury , C. Caputo , P. David , C. Delaere , I.S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaître, J. Lidrych , K. Mondal , T.T. Tran , P. Vischia , S. Wertz 








Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , A. Moraes , P. Rebello Teles 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira⁶ , D. De Jesus Damiao , V. Dos Santos Sousa , S. Fonseca De Souza , J. Martins⁷ , C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , S.M. Silva Do Amaral , A. Sznajder , M. Thiel , A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁶ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , P.G. Mercadante , S.F. Novaes , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Rodozov, M. Shopova , G. Sultanov 




University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov, E. Shumka 

Instituto de Alta Investigacion, Universidad de Tarapaca, Casilla 7 D, Arica, Chile

S. Keshri , S.Thakur 















Beihang University, Beijing, China

T. Cheng , Q. Guo, T. Javaid⁸, M. Mittal , L. Yuan 











Department of Physics, Tsinghua University, Beijing, China

G. Bauer⁹, Z. Hu , S. Lezki , K. Yi^{9,10}

Institute of High Energy Physics, Beijing, China

G.M. Chen⁸ , H.S. Chen⁸ , M. Chen⁸ , F. Iemmi , C.H. Jiang, A. Kapoor , H. Liao , Z.-A. Liu¹¹ , V. Milosevic , F. Monti , R. Sharma , J. Tao , J. Wang , H. Zhang , J. Zhao 


State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , A. Levin , C. Li , Q. Li , X. Lyu, Y. Mao, S.J. Qian , X. Sun , D. Wang , J. Xiao , H. Yang




Sun Yat-Sen University, Guangzhou, China

M. Lu , Z. You 

University of Science and Technology of China, Hefei, China

N. Lu 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao⁴ , D. Leggat, H. Okawa , Y. Zhang 





Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 





Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega



Universidad de Antioquia, Medellin, Colombia

J. Mejia Guisao , F. Ramirez , M. Rodriguez , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , I. Puljak 








University of Split, Faculty of Science, Split, Croatia

Z. Antunovic, M. Kovac , T. Sculac 




Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , S. Mishra , M. Roguljic , A. Starodumov¹² , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

Y. Assran^{13,14}, S. Elgammal¹⁴

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

A. Lotfy , M.A. Mahmoud 













National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

S. Bhowmik , R.K. Dewanjee , K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

P. Eerola , H. Kirschenmann , K. Osterberg , M. Voutilainen 
















Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , M.m. Rantanen , H. Siikonen , E. Tuominen , J. Tuominiemi 





























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow , T. Tuuva[†]

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

C. Amendola , M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , V. Lohezic , J. Malcles , J. Rander, A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁵ , P. Simkina , M. Titov 

















Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

C. Baldenegro Barrera , F. Beaudette , A. Buchot Perraguin , P. Busson , A. Cappati , C. Charlot , F. Damas , O. Davignon , B. Diab , G. Falmagne , B.A. Fontana Santos Alves , S. Ghosh , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , G. Liu , J. Motta , M. Nguyen , C. Ochando , L. Portales , R. Salerno , U. Sarkar , J.B. Sauvan , Y. Sirois , A. Tarabini , E. Vernazza , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram¹⁶ , J. Andrea, D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , U. Goerlach , C. Grimault, A.-C. Le Bihan , P. Van Hove 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , J. Choi , D. Contardo , P. Depasse , C. Dozen¹⁷ , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , G. Grenier , B. Ille , I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, M. Vander Donckt , P. Verdier , S. Viret

Georgian Technical University, Tbilisi, Georgia



















D. Chokheli , I. Lomidze , Z. Tsamalaidze¹² 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany










V. Botta , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany































































S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , M. Erdmann , P. Fackeldey 

B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , M.y. Lee , L. Mastrolorenzo, M. Merschmeyer , A. Meyer , S. Mondal , S. Mukherjee , D. Noll , A. Novak , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb, F. Rehm, H. Reithler , A. Schmidt , S.C. Schuler, A. Sharma , A. Stein , F. Torres Da Silva De Araujo¹⁸ , L. Vigilante, S. Wiedenbeck , S. Zaleski





































RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , W. Haj Ahmad¹⁹ , O. Hlushchenko, T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 




























Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen, M. Aldaya Martin , J. Alimena , Y. An , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke, S. Bhattacharya , F. Blekman²⁰ , K. Borrás²¹ , D. Brunner , A. Campbell , A. Cardini , C. Cheng, F. Colombina, S. Consuegra Rodríguez , G. Correia Silva , M. De Silva , G. Eckerlin, D. Eckstein, L.I. Estevez Banos , O. Filatov , E. Gallo²⁰ , A. Geiser , A. Giraldi , G. Greau, A. Grohsjean , V. Guglielmi , M. Guthoff , A. Jafari²² , N.Z. Jomhari , B. Kaech , M. Kasemann , H. Kaveh , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia, K. Lipka²³ , W. Lohmann²⁴ , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , J. Metwally, A.B. Meyer , G. Milella , M. Mormile , A. Mussgiller , A. Nürnberg , Y. Otariid, D. Pérez Adán , E. Ranken , A. Raspereza, B. Ribeiro Lopes , J. Rübenach, A. Saggio , M. Savitskyi , M. Scham^{25,21} , V. Scheurer, S. Schnake²¹ , P. Schütze , C. Schwanenberger²⁰ , M. Shchedrolosiev , R.E. Sosa Ricardo , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , Q. Wang , Y. Wen , K. Wichmann, L. Wiens²¹ , C. Wissing , S. Wuchterl , Y. Yang, A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , M. Bonanomi , P. Connor , K. De Leo , M. Eich, K. El Morabit , A. Fröhlich, C. Garbers , E. Garutti , M. Hajheidari, J. Haller , A. Hinzmann , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari, T. Kramer , V. Kutzner, F. Labe , J. Lange , A. Lobanov , C. Matthies , A. Mehta , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , M. Rieger , D. Savoio , J. Schindler, P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 






Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, N. Faltermann , M. Giffels , J.O. Gosewisch, A. Gottmann , F. Hartmann²⁶ , M. Horzela , U. Husemann , M. Klute , R. Koppenhöfer , M. Link, A. Lintuluoto , S. Maier , S. Mitra , Th. Müller , M. Neukum, M. Oh , G. Quast , K. Rabbertz , I. Shvetsov, H.J. Simonis, N. Trevisani , R. Ulrich , J. van der Linden , R.F. Von Cube , M. Wassmer , S. Wieland , R. Wolf , S. Wunsch, X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, P. Assiouras , G. Daskalakis , A. Kyriakis, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece







M. Diamantopoulou, D. Karasavvas, P. Kontaxakis , A. Manousakis-Katsikakis , G. Melachroinos, A. Panagiotou, I. Papavergou , N. Saoulidou , K. Theofilatos 

E. Tziaferi , K. Vellidis , I. Zisopoulos

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos ,
G. Tsipolitis, A. Zacharopoulou






University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , C. Kamtsikis,
P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos ,
J. Strologas 




MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , M.M.A. Gadallah²⁷ , P. Major , K. Mandal , G. Pásztor ,
A.J. Rádl²⁸ , O. Surányi , G.I. Veres 



Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók²⁹ , C. Hajdu , D. Horvath^{30,31} , F. Sikler , V. Veszpremi 




Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, N. Beni , S. Czellar, J. Karancsi²⁹ , J. Molnar, Z. Szillasi, D. Teyssier 

















Institute of Physics, University of Debrecen, Debrecen, Hungary

P. Raics, B. Ujvari³² , G. Zilizi 

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo²⁸ , F. Nemes²⁸ , T. Novak 









Panjab University, Chandigarh, India

J. Babbar , S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan ,
N. Dhingra³³ , R. Gupta, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar ,
P. Kumari , M. Meena , K. Sandeep , T. Sheokand, J.B. Singh³⁴ , A. Singla 














University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , M. Naimuddin ,
K. Ranjan , S. Saumya 


Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman³⁵ , S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta,
B. Gomber³⁶ , M. Maity³⁵, P. Palit , G. Saha , B. Sahu³⁶ , S. Sarkar

Indian Institute of Technology Madras, Madras, India

P.K. Behera , S.C. Behera , S. Chatterjee, P. Kalbhor , J.R. Komaragiri³⁷ , D. Kumar³⁷ ,
A. Muhammad , L. Panwar³⁷ , R. Pradhan , P.R. Pujahari , N.R. Saha , A. Sharma ,
A.K. Sikdar , S. Verma 









Bhabha Atomic Research Centre, Mumbai, India

K. Naskar³⁸ 

Tata Institute of Fundamental Research-A, Mumbai, India

T. Aziz, I. Das , S. Dugad, M. Kumar , G.B. Mohanty , P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India








A. Bala , S. Banerjee , M. Guchait , S. Karmakar , S. Kumar , G. Majumder ,
K. Mazumdar , S. Mukherjee , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National

Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁹ , A.K. Das, C. Kar , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu⁴⁰ , A. Nayak⁴⁰ , P. Saha , S.K. Swain, S. Varghese , D. Vats⁴⁰ 

Indian Institute of Science Education and Research (IISER), Pune, India

A. Alpana , S. Dube , B. Kansal , A. Laha , S. Pandey , A. Rastogi , S. Sharma 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴¹ , E. Khazaie , M. Zeinali⁴² 


































Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani⁴³ , S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi 

University College Dublin, Dublin, Ireland

M. Grunewald 






INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , R. Aly^{a,c,44} , C. Aruta^{a,b} , A. Colaleo^a , D. Creanza^{a,c} , L. Cristella^{a,b} , B. D' Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , A. Di Florio^{a,b} , W. Elmetenawee^{a,b} , F. Errico^{a,b} , L. Fiore^a , G. Iaselli^{a,c} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^{a,b} , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^a , D. Ramos^a , A. Ranieri^a , G. Selvaggi^{a,b} , L. Silvestris^a , F.M. Simone^{a,b} , Ü. Sözbilir^a , A. Stameria^a , R. Venditti^a , P. Verwilligen^a 

















INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , L. Borghonovi^a , L. Brigliadori^a , R. Campanini^{a,b} , P. Capiluppi^{a,b} , A. Castro^{a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^{a,b} , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,45} , L. Lunerti^{a,b} , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,46} , A. Di Mattia^a , R. Potenza^{a,b} , A. Tricomi^{a,b,46} , C. Tuve^{a,b} 






INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli^a , G. Bardelli^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^{a,b} , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^{a,b} , M. Meschini^a , S. Paoletti^a , G. Sguazzoni^a , L. Viliani^a 

























INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁷ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy












M. Bozzo^{a,b} , P. Chatagnon^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy




























A. Benaglia^a , G. Boldrini^a , F. Brivio^{a,b} , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , D. Pedrini^a , B.S. Pinolini^a , S. Ragazzi^{a,b} , N. Redaelli^a , T. Tabarelli de Fatis^{a,b} , D. Zuolo^{a,b} 

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della












Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b} , N. Cavallo^{a,c} , A. De Iorio^{a,b} ,
F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,48} , P. Paolucci^{a,26} , B. Rossi^a ,
C. Sciacca^{a,b} 
















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,49} , D. Bisello^{a,b} , P. Bortignon^a ,
A. Bragagnolo^{a,b} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^a , S. Fantinel^a ,
F. Gasparini^{a,b} , U. Gasparini^{a,b} , G. Grosso^a , L. Layer^{a,50} , E. Lusiani^a , M. Margoni^{a,b} ,
A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , G. Strong^a ,
M. Tosi^{a,b} , H. Yarar^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

































INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

S. Abu Zeid^{a,51} , C. Aimè^{a,b} , A. Braghieri^a , S. Calzaferri^{a,b} , D. Fiorina^{a,b} ,
P. Montagna^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 



















INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

P. Asenov^{a,52} , G.M. Bilei^a , D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} ,
G. Mantovani^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,52} , A. Piccinelli^{a,b} ,
M. Presilla^{a,b} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 










































INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Azzurri^a , G. Bagliesi^a , V. Bertacchi^{a,c} , R. Bhattacharya^a , L. Bianchini^{a,b} ,
T. Boccali^a , E. Bossini^{a,b} , D. Bruschini^{a,c} , R. Castaldi^a , M.A. Ciocci^{a,b} ,
V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} ,
D. Matos Figueiredo^a , A. Messineo^{a,b} , M. Musich^{a,b} , F. Palla^a , S. Parolia^a ,
G. Ramirez-Sanchez^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^a , T. Sarkar^a ,
A. Scribano^a , P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , A. Venturi^a ,
P.G. Verdini^a 







INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Barria^a , M. Campana^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} ,
E. Di Marco^a , M. Diemoz^a , E. Longo^{a,b} , P. Meridiani^a , G. Organtini^{a,b} ,
F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a ,
F. Santanastasio^{a,b} , L. Soffi^a , R. Tramontano^{a,b} 














INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a ,
R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} ,
N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} ,
C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , E. Migliore^{a,b} , M. Monteno^a ,
R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a ,
M. Pelliccioni^a , M. Ruspa^{a,c} , K. Shchelina^a , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} ,
D. Soldi^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , M. Tornago^{a,b} , D. Trocino^a ,
G. Umoret^{a,b} , A. Vagnerini^{a,b} , E. Vlasov^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , G. Della Ricca^{a,b} ,
G. Sorrentino^{a,b} 




Kyungpook National University, Daegu, Korea

S. Dogra , C. Huh , B. Kim , D.H. Kim , G.N. Kim , J. Kim, J. Lee , S.W. Lee , C.S. Moon , Y.D. Oh , S.I. Pak , M.S. Ryu , S. Sekmen , Y.C. Yang 





Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , T.J. Kim , J. Park 

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , J. Lim, J. Park, S.K. Park, J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh 

Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , S. Jeon , J. Kim , J.S. Kim, S. Ko , H. Kwon , H. Lee , S. Lee, B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , D. Kim , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , J.A. Merlin, I.C. Park , Y. Roh, D. Song, Watson, I.J. , S. Yang 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi 




Riga Technical University, Riga, Latvia

K. Dreimanis , G. Pikurs, A. Potrebko , M. Seidel , V. Veckalns 




Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , S.Y. Hoh⁵³ , I. Yusuff⁵³ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico






































































J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , I. Heredia-De La Cruz⁵⁴ , R. Lopez-Fernandez , C.A. Mondragon Herrera, D.A. Perez Navarro , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , F. Vazquez Valencia 

Benemerita Universidad Autonoma de Puebla, Puebla, MexicoI. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja, J. Mijuskovic⁵⁵, N. Raicevic **University of Auckland, Auckland, New Zealand**D. Krofcheck **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, M. Gul , H.R. Hoorani , W.A. Khan **AGH University of Science and Technology Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szeleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , A. Boletti , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , J. Varela **Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**M. Dordevic , J. Milosevic **Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**M. Aguilar-Benitez, J. Alcaraz Maestre , M. Barrio Luna, Cristina F. Bedoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , D.D. Redondo Ferrero , L. Romero, S. Sánchez Navas , J. Sastre , L. Urda Gómez , J. Vazquez Escobar , C. Willmott**Universidad Autónoma de Madrid, Madrid, Spain**J.F. de Trocóniz **Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba **Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , G. Gomez , C. Lasosa García , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , J. Piedra Gomez , C. Prieels, L. Scodellaro , I. Vila , J.M. Vizan Garcia 


























University of Colombo, Colombo, Sri Lanka

M.K. Jayananda , B. Kailasapathy⁵⁶ , D.U.J. Sonnadara , D.D.C. Wickramarathna 









University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna , K. Liyanage , N. Perera , N. Wickramage 













CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , B. Bilin , A.A. Bin Anuar , A. Bocci , E. Brondolin , C. Caillol , T. Camporesi , G. Cerminara , N. Chernyavskaya , S.S. Chhibra , S. Choudhury, M. Cipriani , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , M. Dünser , N. Dupont, F. Fallavollita⁵⁷, A. Florent , L. Forthomme , G. Franzoni , W. Funk , S. Ghosh⁵⁸ , S. Giani, D. Gigi, K. Gill, F. Glege , L. Gouskos , E. Govorkova , M. Haranko , J. Hegeman , V. Innocente , T. James , P. Janot , J. Kaspar , J. Kieseler , N. Kratochwil , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , B. Maier , L. Malgeri , M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi , F. Moortgat , M. Mulders , S. Orfanelli, L. Orsini, F. Pantaleo , E. Perez, M. Peruzzi , A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , M. Pitt , H. Qu , T. Quast, D. Rabady , A. Racz, G. Reales Gutiérrez, M. Rovere , H. Sakulin , J. Salfeld-Nebgen , S. Scarfi, M. Selvaggi , A. Sharma , P. Silva , P. Sphicas⁵⁹ , A.G. Stahl Leiton , A. Steen, S. Summers , K. Tatar , D. Treille , P. Tropea , A. Tsirou, D. Walter , J. Wanczyk⁶⁰ , K.A. Wozniak , W.D. Zeuner
























Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua⁶¹ , L. Caminada⁶¹ , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶¹ , L. Noehte⁶¹ , T. Rohe 




ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov⁶⁰ , M. Backhaus , A. Calandri , K. Datta , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , T.A. Gómez Espinosa , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , L. Marchese , C. Martin Perez , A. Mascellani⁶⁰ , F. Nessi-Tedaldi , J. Niedziela , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , F. Riti , D. Ruini, D.A. Sanz Becerra , R. Seidita , J. Steggemann⁶⁰ , D. Valsecchi , R. Wallny 











Universität Zürich, Zurich, Switzerland

C. Amsler⁶² , P. Bäertschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier , A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger , Y. Takahashi 

National Central University, Chung-Li, Taiwan

C. Adloff⁶³, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari³⁷ , S.S. Yu 

















National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.s. Chen, H. Cheng , W.-S. Hou , R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, J. Thomas-Wilsker , H.y. Wu, E. Yazgan 

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , Z.S. Demiroglu , F. Dolek , I. Dumanoglu⁶⁴ , E. Eskut, Y. Guler⁶⁵ , E. Gurpinar Guler⁶⁵ , C. Isik, O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir⁶⁶ , A. Polatoz, B. Tali⁶⁷ , U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Karapinar⁶⁸, K. Ocalan⁶⁹ , M. Yalvac⁷⁰ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁷¹ , O. Kaya⁷² , S. Tekten⁷³ 


Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak⁶⁴ , Y. Komurcu , S. Sen⁷⁴ 


Istanbul University, Istanbul, Turkey

O. Aydilek , S. Cerci⁶⁷ , V. Epshteyn , B. Hacisahinoglu , I. Hos⁷⁵ , B. Isildak⁷⁶ , B. Kaynak , S. Ozkorucuklu , C. Simsek , D. Sunar Cerci⁶⁷ 















Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

B. Grynyov 


















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 



























University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , L. Kreczko , B. Krikler , S. Paramesvaran , S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou⁷⁷ , K. Walkingshaw Pass, R. White 




Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁸ , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , K.V. Ellis, K. Harder , S. Harper , M.-L. Holmberg⁷⁹ , Sh. Jain , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin, T. Williams 














Imperial College, London, United Kingdom

R. Bainbridge , P. Bloch , J. Borg , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , V. Cepaitis , G.S. Chahal⁸⁰ , D. Colling , J.S. Dancu, P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , J. Langford , L. Lyons , A.-M. Magnan , S. Malik, A. Martelli , M. Mieskolainen , J. Nash⁸¹ , M. Pesaresi, B.C. Radburn-Smith , A. Richards, A. Rose , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee²⁶ , M. Vojinovic , N. Wardle , S.N. Webb , D. Winterbottom






Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , J. Hiltbrand , A.R. Kanuganti , B. McMaster , M. Saunders , S. Sawant , C. Sutantawibul , M. Toms , J. Wilson 

















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniyal , A.M. Vargas Hernandez 















The University of Alabama, Tuscaloosa, Alabama, USA

R. Chudasama , S.I. Cooper , D. Di Croce , S.V. Gleyzer , C.U. Perez , P. Rumerio⁸² , E. Usai , C. West 



















Boston University, Boston, Massachusetts, USA

A. Akpinar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , C. Erice , E. Fontanesi , D. Gastler , S. May , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan 











Brown University, Providence, Rhode Island, USA

G. Benelli , X. Coubez²¹, D. Cutts , M. Hadley , U. Heintz , J.M. Hogan⁸³ , T. Kwon , G. Landsberg , K.T. Lau , D. Li, J. Luo , M. Narain, N. Pervan , S. Sagir⁸⁴ , F. Simpson , W.Y. Wong, X. Yan , D. Yu , W. Zhang





University of California, Davis, Davis, California, USA

M.S. Abbott , J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , J. Conway , P.T. Cox , R. Erbacher , G. Haza , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , D. Pellett , B. Regnery , W. Wei, Y. Yao , F. Zhang 






















University of California, Los Angeles, California, USA

M. Bachtis , R. Cousins , A. Datta , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , E. Manca , W.A. Nash , D. Saltzberg , B. Stone , V. Valuev 



















University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , M. Gordon, G. Hanson , O.R. Long , W. Si , S. Wimpenny











University of California, San Diego, La Jolla, California, USA

J.G. Branson, S. Cittolin, S. Cooperstein , D. Diaz , J. Duarte , R. Gerosa , L. Giannini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 















University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

L. Brennan, C. Campagnari , M. Citron , G. Collura , A. Dorsett , J. Incandela , M. Kilpatrick , J. Kim , A.J. Li , P. Masterson , H. Mei , M. Oshiro , J. Richman , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , P. Siddireddy, D. Stuart , S. Wang 














California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, A. Latorre, J.M. Lawhorn , J. Mao , H.B. Newman , T. Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M.B. Andrews , P. Bryant , V. Dutta , T. Ferguson , A. Harilal , C. Liu , T. Mudholkar , S. Murthy , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 
















University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, N. Manganello , F. Marini , A. Perloff , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper 























Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , D. Gadkari , S. Hogan , J. Monroy , J.R. Patterson , J. Reichert , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 













Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , K.F. Di Petrillo , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , L. Horyn , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸⁵ , F. Ravera , A. Reinsvold Hall⁸⁶ , L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 








University of Florida, Gainesville, Florida, USA

P. Avery , D. Bourilkov , L. Cadamuro , P. Chang , V. Cherepanov , R.D. Field , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , E. Kuznetsova , K.H. Lo , K. Matchev , N. Menendez , G. Mitselmakher , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig , S. Rosenzweig , K. Shi , J. Wang , Z. Wu 


















Florida State University, Tallahassee, Florida, USA

T. Adams , A. Askew , N. Bower , R. Habibullah , V. Hagopian , T. Kolberg , G. Martinez , H. Prosper , O. Viazlo , M. Wulansatiti , R. Yohay , J. Zhang 















Florida Institute of Technology, Melbourne, Florida, USA

M.M. Baarmand , S. Butalla , T. Elkafrawy⁵¹ , M. Hohlmann , R. Kumar Verma , M. Rahmani , F. Yumiceva 

University of Illinois at Chicago (UIC), Chicago, Illinois, USA

M.R. Adams , R. Cavanaugh , S. Dittmer , O. Evdokimov , C.E. Gerber , D.J. Hofman , D. S. Lemos , A.H. Merrit , C. Mills , G. Oh , T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , X. Wang , Z. Ye , J. Yoo 























The University of Iowa, Iowa City, Iowa, USA

M. Alhousseini , K. Dilsiz⁸⁷ , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo , A. Mestvirishvili⁸⁸ , J. Nachtman , O. Neogi , H. Ogul⁸⁹ , Y. Onel , A. Penzo , C. Snyder , E. Tiras⁹⁰ 








Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , S. Kyriacou , P. Maksimovic , J. Roskes , S. Sekhar , M. Swartz , T.Á. Vámi 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , P. Baringer , A. Bean , Z. Flowers , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , C. Lindsey , J. Marquez , N. Minafra , M. Murray , M. Nickel , C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , Q. Wang , G. Wilson 



















Kansas State University, Manhattan, Kansas, USA

B. Allmond , S. Duric, A. Ivanov , K. Kaadze , A. Kalogeropoulos , D. Kim, Y. Maravin , T. Mitchell, A. Modak, K. Nam, J. Natoli , D. Roy 























Lawrence Livermore National Laboratory, Livermore, California, USA

F. Rebassoo , D. Wright 













University of Maryland, College Park, Maryland, USA

E. Adams , A. Baden , O. Baron, A. Belloni , A. Bethani , Y.m. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , Y. Lai , S. Lasocio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , L. Wang , K. Wong 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

J. Bendavid , W. Busza , I.A. Cali , Y. Chen , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , Y.-J. Lee , K. Long , C. Mironov , C. Paus , D. Rankin , C. Roland , G. Roland , Z. Shi , G.S.F. Stephans , J. Wang, Z. Wang , B. Wyslouch , T. J. Yang 












University of Minnesota, Minneapolis, Minnesota, USA

R.M. Chatterjee, B. Crossman, B.M. Joshi , C. Kapsiak , M. Krohn , Y. Kubota , D. Mahon , J. Mans , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M.A. Wadud 

University of Mississippi, Oxford, Mississippi, USA

L.M. Cremaldi 

















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , M. Bryson, D.R. Claes , C. Fangmeier , F. Golf , C. Joo , I. Kravchenko , I. Reed , J.E. Siado , G.R. Snow[†], W. Tabb , A. Wightman , F. Yan , A.G. Zecchinelli 










State University of New York at Buffalo, Buffalo, New York, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean , M. Morris, D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams 





















Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Krishna , J. Li , G. Madigan , B. Marzocchi , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , B. Wang , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Bhattacharya , J. Bueghly, Z. Chen , A. Gilbert , K.A. Hahn , Y. Liu , D.G. Monk , N. Odell , M.H. Schmitt , A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, USA

R. Band , R. Bucci, M. Cremonesi, A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , K. Mohrman , C. Moore , Y. Musienko¹² , R. Ruchti , A. Townsend , M. Wayne , H. Yockey, M. Zarucki , L. Zygalia 

The Ohio State University, Columbus, Ohio, USA

B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , A. Lesauvage , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

























F.M. Addesa , H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal 

B. Greenberg , N. Haubrich , S. Higginbotham , G. Kopp , S. Kwan , D. Lange ,
A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik 





















Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , V.E. Barnes , S. Chandra , R. Chawla , S. Das , A. Gu , L. Gutay , M. Jones ,
A.W. Jung , D. Kondratyev , A.M. Koshy , M. Liu , G. Negro , N. Neumeister ,
G. Paspalaki , S. Piperov , A. Purohit , J.F. Schulte , M. Stojanovic , J. Thieman ,
A. K. Viridi , F. Wang , R. Xiao , W. Xie 













Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar 

Rice University, Houston, Texas, USA

D. Acosta , A. Baty , T. Carnahan , S. Dildick , K.M. Ecklund , P.J. Fernández Man-
teca , S. Freed , P. Gardner , F.J.M. Geurts , A. Kumar , W. Li , J. Lin , O. Miguel Colin ,
B.P. Padley , R. Redjimi , J. Rotter , S. Yang⁹¹ , Z. Ye , E. Yigitbasi , Y. Zhang 























University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon , A. Garcia-
Bellido , O. Hindrichs , A. Khukhunaishvili , P. Parygin , E. Popova , R. Taus ,
G.P. Van Onsem 

The Rockefeller University, New York, New York, USA

K. Goulianos 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA
















B. Chiarito , J.P. Chou , Y. Gershtein , E. Halkiadakis , A. Hart , M. Heindl ,
D. Jaroslowski , O. Karacheban²⁴ , I. Laflotte , A. Lath , R. Montalvo , K. Nash ,
M. Osherson , H. Routray , S. Salur , S. Schnetzer , S. Somalwar , R. Stone ,
S.A. Thayil , S. Thomas , J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, USA

H. Acharya , A.G. Delannoy , S. Fiorendi , T. Holmes , E. Nibigira , S. Spanier 

Texas A

&M University, College Station, Texas, USA

M. Ahmad , O. Bouhali⁹² , M. Dalchenko , A. Delgado , R. Eusebi , J. Gilmore ,
T. Huang , T. Kamon⁹³ , H. Kim , S. Luo , S. Malhotra , R. Mueller , D. Overton ,
D. Rathjens , A. Safonov 










Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , V. Hegde , K. Lamichhane , S.W. Lee , T. Mengke ,
S. Muthumuni , T. Peltola , I. Volobouev , A. Whitbeck 


Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , S. Greene , A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo ,
E. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 






















University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , B. Cox , G. Cummings , J. Hakala , R. Hirosky , A. Ledovskoy ,
A. Li , C. Neu , C.E. Perez Lara 

Wayne State University, Detroit, Michigan, USA

P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev, V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev , V. Blinov⁹⁴, E. Boos , V. Borshch , D. Budkouski , V. Chekhovsky, R. Chistov⁹⁴ , M. Danilov⁹⁴ , A. Dermenev , T. Dimova⁹⁴ , I. Dremin , A. Ershov , G. Gavrilov , V. Gavrilov , S. Gninenko , V. Golovtsov , N. Golubev , I. Golutvin, I. Gorbunov , A. Gribushin , Y. Ivanov , V. Kachanov , A. Kaminskiy, L. Kardapoltsev⁹⁴ , V. Karjavine , A. Karneyev , L. Khein, V. Kim⁹⁴ , M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , O. Kodolova⁹⁵ , D. Konstantinov , V. Korenkov , V. Korotkikh, A. Kozyrev⁹⁴ , N. Krasnikov , A. Lanev , P. Levchenko⁹⁶ , A. Litomin, N. Lychkovskaya , V. Makarenko , A. Malakhov , V. Matveev⁹⁴ , V. Murzin , A. Nikitenko^{97,95} , S. Obraztsov , I. Ovtin⁹⁴ , V. Palichik , V. Perelygin , S. Petrushanko , S. Polikarpov⁹⁴ , V. Popov, O. Radchenko⁹⁴ , M. Savina , V. Savrin , D. Selivanova , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹⁴ , S. Slabospitskii , V. Smirnov , A. Snigirev , D. Sosnov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , I. Tlisova , A. Toropin , L. Uvarov , A. Uzunian , I. Vardanyan, A. Vorobyev[†], N. Voytishin , B.S. Yuldashev⁹⁸, A. Zarubin , I. Zhizhin , A. Zhokin 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

⁴Also at Université Libre de Bruxelles, Bruxelles, Belgium

⁵Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷Also at UFMS, Nova Andradina, Brazil

⁸Also at University of Chinese Academy of Sciences, Beijing, China

⁹Also at Nanjing Normal University Department of Physics, Nanjing, China

¹⁰Now at The University of Iowa, Iowa City, Iowa, USA

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹³Also at Suez University, Suez, Egypt

¹⁴Now at British University in Egypt, Cairo, Egypt

¹⁵Also at Purdue University, West Lafayette, Indiana, USA

¹⁶Also at Université de Haute Alsace, Mulhouse, France

¹⁷Also at Department of Physics, Tsinghua University, Beijing, China

¹⁸Also at The University of the State of Amazonas, Manaus, Brazil

¹⁹Also at Erzincan Binali Yildirim University, Erzincan, Turkey

²⁰Also at University of Hamburg, Hamburg, Germany

²¹Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

-
- ²²Also at Isfahan University of Technology, Isfahan, Iran
- ²³Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
- ²⁴Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁵Also at Forschungszentrum Jülich, Juelich, Germany
- ²⁶Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁷Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ²⁸Also at Wigner Research Centre for Physics, Budapest, Hungary
- ²⁹Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- ³⁰Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- ³¹Now at Universitatea Babeș-Bolyai - Facultatea de Fizică, Cluj-Napoca, Romania
- ³²Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary
- ³³Also at Punjab Agricultural University, Ludhiana, India
- ³⁴Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ³⁵Also at University of Visva-Bharati, Santiniketan, India
- ³⁶Also at University of Hyderabad, Hyderabad, India
- ³⁷Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁸Also at Indian Institute of Technology (IIT), Mumbai, India
- ³⁹Also at IIT Bhubaneswar, Bhubaneswar, India
- ⁴⁰Also at Institute of Physics, Bhubaneswar, India
- ⁴¹Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴²Also at Sharif University of Technology, Tehran, Iran
- ⁴³Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁴Also at Helwan University, Cairo, Egypt
- ⁴⁵Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁶Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁷Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁴⁸Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁴⁹Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵⁰Also at Università di Napoli 'Federico II', Napoli, Italy
- ⁵¹Also at Ain Shams University, Cairo, Egypt
- ⁵²Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵³Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁴Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁵Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
- ⁵⁶Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁷Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
- ⁵⁸Also at Indian Institute of Technology Hyderabad, Hyderabad, India
- ⁵⁹Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶⁰Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶¹Also at Universität Zürich, Zurich, Switzerland
- ⁶²Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶³Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁴Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁵Also at Konya Technical University, Konya, Turkey

- ⁶⁶Also at Piri Reis University, Istanbul, Turkey
- ⁶⁷Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁸Also at Istanbul Gedik University, Istanbul, Turkey
- ⁶⁹Also at Necmettin Erbakan University, Konya, Turkey
- ⁷⁰Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁷¹Also at Marmara University, Istanbul, Turkey
- ⁷²Also at Milli Savunma University, Istanbul, Turkey
- ⁷³Also at Kafkas University, Kars, Turkey
- ⁷⁴Also at Hacettepe University, Ankara, Turkey
- ⁷⁵Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁶Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁷Also at Vrije Universiteit Brussel, Brussel, Belgium
- ⁷⁸Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁹Also at University of Bristol, Bristol, United Kingdom
- ⁸⁰Also at IPPP Durham University, Durham, United Kingdom
- ⁸¹Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸²Also at Università di Torino, Torino, Italy
- ⁸³Also at Bethel University, St. Paul, Minnesota, USA
- ⁸⁴Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸⁵Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁶Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸⁷Also at Bingöl University, Bingöl, Turkey
- ⁸⁸Also at Georgian Technical University, Tbilisi, Georgia
- ⁸⁹Also at Sinop University, Sinop, Turkey
- ⁹⁰Also at Erciyes University, Kayseri, Turkey
- ⁹¹Also at Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China
- ⁹²Also at Texas A & M University at Qatar, Doha, Qatar
- ⁹³Also at Kyungpook National University, Daegu, Korea
- ⁹⁴Also at another institute or international laboratory covered by a cooperation agreement with CERN
- ⁹⁵Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁹⁶Also at Northeastern University, Boston, Massachusetts, USA
- ⁹⁷Also at Imperial College, London, United Kingdom
- ⁹⁸Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan