



Anomaly detection search for new resonances decaying into a Higgs boson and a generic new particle X in hadronic final states using $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector

The ATLAS Collaboration

A search is presented for a heavy resonance Y decaying into a Standard Model Higgs boson H and a new particle X in a fully hadronic final state. The full Large Hadron Collider Run 2 dataset of proton–proton collisions at $\sqrt{s} = 13$ TeV collected by the ATLAS detector from 2015 to 2018 is used, and corresponds to an integrated luminosity of 139 fb^{-1} . The search targets the high Y -mass region, where the H and X have a significant Lorentz boost in the laboratory frame. A novel signal region is implemented using anomaly detection, where events are selected solely because of their incompatibility with a learned background-only model. It is defined using a jet-level tagger for signal-model-independent selection of the boosted X particle, representing the first application of fully unsupervised machine learning to an ATLAS analysis. Two additional signal regions are implemented to target a benchmark X decay into two quarks, covering topologies where the X is reconstructed as either a single large-radius jet or two small-radius jets. The analysis selects Higgs boson decays into $b\bar{b}$, and a dedicated neural-network-based tagger provides sensitivity to the boosted heavy-flavor topology. No significant excess of data over the expected background is observed, and the results are presented as upper limits on the production cross section $\sigma(pp \rightarrow Y \rightarrow XH \rightarrow q\bar{q}b\bar{b})$ for signals with m_Y between 1.5 and 6 TeV and m_X between 65 and 3000 GeV.

1 Introduction

The Standard Model (SM) provides a framework for understanding fundamental particles and interactions that has been remarkably predictive of experimental results over several decades. The discovery of the Higgs boson in 2012 [1, 2] completed the set of particles predicted by the SM.

The sensitivity of the Higgs boson mass to radiative corrections implies either extreme fine-tuning in the SM or the existence of new particles at an energy scale not far above the Higgs boson mass. This theoretical motivation, coupled with the existing experimental mass reach of the Large Hadron Collider (LHC) at CERN, motivates searches for new particles with $\mathcal{O}(\text{TeV})$ masses. Because the Higgs boson couples more strongly to heavier particles, it is natural to expect that these new heavy particles may have decays to a Higgs boson.

A search is presented here for a new TeV-scale narrow-width particle Y , which decays into a Standard Model Higgs boson H and a new particle X with a mass near the weak scale. A fully hadronic final state is targeted for both particles. Tagging of the boosted Higgs decay into two b -quarks ($H \rightarrow b\bar{b}$ tagging) enhances a signal by using the highest branching fraction decay of the Higgs boson. A novel jet-level implementation of *anomaly detection* implemented via an unsupervised machine-learning architecture is used to select X particles solely because of their incompatibility with the expected SM background.

The application of anomaly detection to collider searches is a rapidly growing effort in the high-energy physics community (see Refs. [3–5] for overviews). The lack of evidence for new interactions and particles since the Higgs boson’s discovery has motivated the execution of generic searches to complement the existing rigorous, model-dependent analysis program. Machine learning provides an excellent framework for the construction of tools that can isolate events in data solely because of their incompatibility with a background-only hypothesis. Building a tool to perform model-independent classification of collision events involves training on data events, and therefore requires the ability to cope with a lack of labels indicating whether inputs are signal or background. This distinguishes the typical supervised classification problem, where all inputs are labeled with a known origin, from the anomaly detection approach, which makes use of unsupervised (no input labels) or weakly supervised (noisy labels) training. A selection tool that is constructed in this way has trade-offs: while it broadens sensitivity to many classes of new physics with a single classifier, it is expected that sensitivity to any specific model will be better with a dedicated supervised approach.

A Feynman diagram for the $Y \rightarrow XH$ signal process is shown in Figure 1, where the X can have a variety of hadronic decays. A nominal benchmark decay into two light quarks, $X \rightarrow q\bar{q}$, is generated to provide an interpretation framework for the results. The generic selection of the X is not strongly dependent on its mass, so the analysis is sensitive to X masses spanning several orders of magnitude, from $\mathcal{O}(10)$ GeV to $\mathcal{O}(1)$ TeV. The masses of the parent and daughter particles yield a kinematic scenario where the final-state particles are highly Lorentz-boosted in the laboratory frame, motivating a “boosted” reconstruction method using large-radius (large- R) jets and jet substructure observables to distinguish the boson decay products. An orthogonal “resolved” reconstruction method is used to recover sensitivity to topologies where the X is less boosted and reconstructed as separate small-radius (small- R) jets, significantly extending the sensitive phase-space region. Since the signal is resonant, it can be detected with a “bump hunt” on the invariant mass distribution of the reconstructed X and H candidates.

Although model-independent in nature, this search is motivated by several key extensions to the Standard Model which predict heavy diboson resonances. Examples of such theories include extended gauge symmetry models [6], warped extra dimensions [7–9], or two-Higgs-doublet models [10]. Signals are




Figure 1: Feynman diagram of the target signal process, where the Y is produced in the initial pp collision and decays to a fully hadronic final state via a SM Higgs boson $H \rightarrow b\bar{b}$ and a new particle X . The only assumption applied to the X decay is that it yields a hadronic final state.

generated with a simplified model based on spin-1 Heavy Vector Triplets (HVT) [11] which reproduces a large class of explicit beyond-the-SM (BSM) models. The HVT phenomenology provides a Lagrangian that fulfills SM symmetry constraints with an isospin $SU(2)$ triplet formed of a neutral Z' and two charged W'^{\pm} bosons.

This search uses the full LHC Run 2 $\sqrt{s} = 13$ TeV pp dataset collected by the ATLAS detector from 2015 to 2018 and corresponding to an integrated luminosity of 139 fb^{-1} . A search for the $Y \rightarrow XH$ process was previously performed by ATLAS using 36.1 fb^{-1} of data, assuming $X \rightarrow q\bar{q}$, and found no significant excess for Y masses from 1 to 4 TeV and X masses from 50 to 1000 GeV [12]. In addition to the much larger dataset, the present analysis includes several key improvements relative to the previous search, such as a neural-network-based tagger optimized for the boosted $H \rightarrow b\bar{b}$ topology, anomaly detection for enhanced signal model independence, and the use of two orthogonal regions to capture both boosted and resolved reconstruction of the nominal X decay into two quarks. The ATLAS Collaboration previously leveraged anomaly detection implemented with weakly supervised machine learning in a search for narrow new resonances in dijet events using the full Run 2 dataset [13]. The CMS experiment performed a multidimensional search for diboson resonances with their full Run 2 dataset of 138 fb^{-1} which includes sensitivity to the signature discussed here [14]. A maximum local (global) significance of 3.6 (2.3) standard deviations was observed for two mild excesses of events, at resonance masses of 2.1 and 2.9 TeV with X masses consistent with W , Z , or Higgs bosons.

The analysis regions in the present search are built by selecting two large- R jets, with additional criteria to reduce the background to events with Higgs and X particles in the search regions. Three overlapping analysis categories are defined with different X selections, namely the anomaly region and two orthogonal regions optimized for the benchmark $X \rightarrow q\bar{q}$ decay. The degree of model-independence of the X tagging is assessed using three simulated signals with different decays of the X particle that lead to differing large- R -jet topologies, in addition to the two-light-quark final state used to generate the (m_Y, m_X) signal grid. The background to the signal process consists primarily of multijet events from quantum chromodynamics (QCD) processes. It is estimated with a fully data-driven method that incorporates reweighting based on a deep neural network (DNN) to ensure good modeling. The final fits are made to the reconstructed invariant mass distribution of the Y candidates in overlapping ranges of X candidate mass to increase the sensitivity to a signal. Since no significant deviation from the estimated background is observed, results are presented as upper limits on the cross section times branching fraction for the generic HVT process.

2 ATLAS detector

The ATLAS detector [15] is a multipurpose particle detector with a forward-backward symmetric cylindrical geometry and a near 4π coverage in solid angle.¹ The inner-detector (ID) system is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$. The high-granularity silicon pixel detector covers the vertex region and typically provides four space-point measurements per track, the first hit normally being in the insertable B-layer installed before Run 2 [16, 17]. The next detector outward is the silicon microstrip tracker, which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker, which enables radially extended track reconstruction up to $|\eta| = 2.0$. Lead/liquid-argon (LAr) sampling calorimeters provide electromagnetic (EM) energy measurements with high granularity. A steel/scintillator-tile hadron calorimeter covers the central pseudorapidity range ($|\eta| < 1.7$). The endcap and forward regions are instrumented with LAr calorimeters for both the EM and hadronic energy measurements up to $|\eta| < 4.9$. The muon spectrometer surrounds the calorimeters and is based on three large superconducting air-core toroidal magnets with eight coils each. The field integral of the toroids ranges between 2.0 and 6.0 Tm across most of the detector. The muon spectrometer includes a system of precision chambers for tracking and fast detectors for triggering. A two-level trigger system is used to select events. The first-level trigger is implemented in hardware and uses a subset of the detector information to accept events at a rate below 100 kHz on average depending on the data-taking conditions. This is followed by a software-based high-level trigger (HLT) that reduces the rate of selected events to 1 kHz for offline storage. An extensive software suite [18] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

3 Data and Monte Carlo simulation

This search is performed with $\sqrt{s} = 13$ TeV pp collision data collected by the ATLAS detector during Run 2 of the LHC (2015–2018). After the application of data quality requirements [19] to ensure that all detector components are operating normally, the dataset corresponds to an integrated luminosity of $139.0 \pm 2.4 \text{ fb}^{-1}$ [20, 21]. Data events used in this analysis were triggered by the presence of a high- p_T large- R jet, built at trigger level using calorimeter-cell energy clusters calibrated to the hadronic scale utilizing the local cell-signal weighting method [22]. The unrescaled single-large- R -jet trigger with the lowest jet- p_T threshold was used in each year, corresponding to thresholds of 360 GeV in 2015, 420 GeV in 2016, and 440 GeV in 2017 and 2018. Further offline selection criteria are imposed (see Section 6) to ensure that selected events are all in the kinematic region where the trigger is fully efficient.

Monte Carlo (MC) event generators were used to simulate the signal targeted by this search. The simulated event samples include the effect of additional pp interactions (pileup) in the same or neighboring bunch crossings. The effect is assessed by overlaying simulated minimum-bias events on each hard-scatter event and accounting for how interactions in the previous or following bunch crossing affect the detector response. The simulated events are weighted to reproduce the pile-up distribution observed in each data-taking period.

¹ ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the center of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the center of the LHC ring, and y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$.

The detector response was simulated [23] using GEANT4 [24] and the events were reconstructed with the same software as used for data.

A generic HVT [11, 25] model with simulated qq scattering is used as a baseline for the $Y \rightarrow XH$ signal samples. The Y resonance is assumed to have a width that is narrow compared to the detector resolution. The generation of the $W' \rightarrow WH$ process was modified to replace the W boson with a new spin-1 boson X with a width of 2 GeV and a 100% branching fraction to $u\bar{d}$. The Standard Model Higgs boson was generated with decays into $b\bar{b}$ only. Samples with Y resonance masses of 1–6 TeV were generated using MADGRAPH5_AMC@NLO 2.7.2 and 2.7.3 [26] interfaced to PYTHIA 8.2 [27] for parton showering and hadronization with the NNPDF2.3LO parton distribution function (PDF) set [28], at leading order in QCD, and parameter values set according to the A14 tune [29]. A two-dimensional grid was generated for 195 signal points defined by Y and X mass values, with Y masses in the range 1.5–6 TeV and X masses in the range 65–3000 GeV. The Y and X mass values account for the kinematic constraint $m_Y > m_X$, and the X mass boundaries are dictated by achievable sensitivity. The signal acceptance times the efficiency has values ranging from a few percent to 40% for the various signal region selections.

In addition to the $Y \rightarrow XH$ signal grid, three signals with different jet topologies were used to craft and assess the degree of model independence in the X anomaly-tagging procedure. They were all generated with PYTHIA 8 using the same tune (A14) and PDFs (NNPDF2.3LO) as the $Y \rightarrow XH$ signal grid. The HVT $W' \rightarrow WZ$ configuration from Ref. [30] was used to create generic resonance signatures of the form $A \rightarrow BC$, where the daughter particles decay into either three light quarks or two heavy-flavor quarks. Choosing as benchmarks a parent A mass of 3 TeV and daughter masses of 200 and 400 GeV creates approximately the same kinematic scenario as the $Y \rightarrow XH$ phase space of interest, leading to boosted decay products that are captured as large- R jets. Depending on the daughter decay into either qqq or $b\bar{b}$, the large- R jet is produced with three-prong substructure or displaced vertices, respectively. Lastly, the PYTHIA 8 Hidden Valley Model A configuration was used to generate a “dark jet” signal that arises from the decay of a Z' dark sector mediator into two dark quarks, where the Z' has the same benchmark mass of 3 TeV [31]. These jets contain dark matter particles that do not interact with the detector, creating a hadronization pattern that contains both visible and invisible energy. Such jets are difficult to recognize using traditional substructure variables that focus on specific signal topologies, making them an ideal target for anomaly detection based on background-only characterization.

4 Object selection

Charged-particle tracks measured to have $p_T > 500$ MeV in the inner detector to are used to reconstruct pp collision vertices, and the one with the largest Σp_T^2 of such tracks is chosen as the event’s primary vertex.

Jets are built with the anti- k_t algorithm [32] in FASTJET [33], using two radius parameter values to cluster the constituent detector signals: $R = 1.0$ for large- R jets J , and $R = 0.4$ for small- R jets j . Large- R jets must satisfy $p_T > 200$ GeV and $|\eta| < 2.0$. They are constructed from combinations of tracks and calibrated calorimeter energy clusters known as Track-CaloClusters (TCCs) [34]. The TCCs take advantage of both the excellent energy resolution of the ATLAS calorimeter and the better angular resolution of the ID tracking system at very high energy, where the calorimeter is less able to discern jet substructure. This is particularly advantageous for measuring substructure in highly boosted jets, as is needed in this search to distinguish $Y \rightarrow XH$ signal from multijet background. To minimize the impact of pileup, large- R jets are trimmed [35] by removing any $R = 0.2$ subjets carrying less than 5% of the large- R jet’s p_T . The jet energy and mass is calibrated with a MC-based method [36].

Small- R jets are built from particle-flow objects [37], with improved accuracy of the jet’s charged-hadron energy component through measurements of the associated charged-particle momenta in the ID. Small- R jets must have $p_T > 20$ GeV and $|\eta| < 4.5$. An additional event-level veto is applied to reject events with a jet that is likely to have come from calorimeter noise, beam-induced background, or a cosmic ray [38].

Leptons, while excluded from the analysis region selections, participate in an overlap removal procedure that prevents double-counting of overlapping objects. Leptons must have $p_T > 7$ GeV, with $|\eta| < 2.47$ for electrons and $|\eta| < 2.7$ for muons. To ensure that each lepton’s track originates from the primary vertex, the transverse impact parameter d_0 , measured relative to the beam line, must have a significance $|d_0/\sigma(d_0)| < 3$ (5) for muons (electrons), and the longitudinal impact parameter z_0 , from the primary vertex to the point where d_0 is measured, must satisfy $|z_0 \sin \theta| < 0.5$ mm. No lepton isolation criteria are applied.

The overlap removal procedure is applied to all selected leptons and jets. If two electrons share the same track, or the separation between their two energy clusters satisfies $|\Delta\eta| < 0.075$ and $|\Delta\phi| < 0.125$, then the lower- p_T electron is discarded. Electrons that fall within $\Delta R = 0.02$ of a selected muon are also discarded. For electrons and nearby small- R jets, the jet is removed if the separation between the electron and jet satisfies $\Delta R < 0.2$; the electron is removed if the separation satisfies $0.2 < \Delta R < 0.4$. For muons and nearby small- R jets, the jet is removed if the separation between the muon and jet satisfies $\Delta R < 0.2$ and if the jet has less than three tracks or the energy and momentum differences between the muon and the jet are small; otherwise the muon is removed if the separation satisfies $\Delta R < 0.4$. To prevent double-counting of energy from an electron inside a large- R jet, the large- R jet is removed if its separation from the electron satisfies $\Delta R < 1.0$.

5 Variational recurrent neural network for anomalous-jet tagging

The anomaly detection in this search is performed with a jet-level Anomaly Score (S_A) [39]. The S_A value is given by a variational recurrent neural network (VRNN) [40], which consists of a variational autoencoder (VAE) whose latent space is updated at each time step of a recurrent neural network (RNN).

The class of models that are based on autoencoders are popular for the selection of rare or outlier events in high-energy physics [41–46]. Such an architecture is composed of two separate stages: an encoder that compresses the input into a lower-dimensional latent space through identification of its salient features, and a decoder that samples from the latent space and attempts to reproduce the input in its original dimensionality. Minimizing the reconstruction error between input and output is a key element of the training, and allows the autoencoder to implicitly learn the underlying distribution of features in its input dataset, and thus recognize anomalous data through its inability to reconstruct such events with low error.

VAEs extend this concept to perform Bayesian inference via a latent space that can accommodate a distribution of encoded information. The VRNN architecture combines the variational inference capabilities of a VAE with the sequence modeling provided by an RNN. The inclusion of a learned, time-dependent prior distribution is an essential feature of the VRNN, and allows the modeling of complex structured sequences with high variability.

The VRNN is trained on large- R TCC jets in the ATLAS Run 2 dataset satisfying the criteria in Section 6 for full trigger efficiency, the jet requirements described in Section 4, and $p_T(J) > 1.2$ TeV. Its architecture is the same as that described in Ref. [39]. Training is performed using the PyTorch deep-learning

library [47], and the network is updated using the Adam optimizer [48] with a learning rate parameter of 10^{-5} . No regularization via weight decay is applied, but gradient clipping is used with a clip value of 10.

The input jets are modeled as a sequence of up to 20 constituent four-vectors per jet, ordered in k_T -splitting starting from the highest- p_T constituent. The p_T selection is designed to chose input jets with highly boosted topologies, for which the VRNN is expected to most improve the sensitivity over the substructure-based approach, as these topologies are well-described by k_T -sorted sequence modeling and are also challenging to distinguish with substructure methods. The training is conditioned on four high-level variables, namely the energy-correlator substructure variable D_2 [49] for two-prong sensitivity, the N -subjettiness ratio τ_{32} [50] for three-prong sensitivity, and the two k_T -splitting-scale ratios d_{12} and d_{23} [51]. An alignment procedure applied to each jet rescales to the same p_T , boosts to the same energy, and rotates to the same orientation in η and ϕ , minimizing the ability of the VRNN to tag only on anomalous kinematic properties without considering internal constituent properties. This input modeling is designed to reveal correlations between constituents and substructure, allowing the VRNN to distinguish jets with anomalous energy-deposition patterns from the background of homogenous jets originating from QCD processes.

Since the input consists solely of jets from data, no labeling scheme is used in training, distinguishing this method of unsupervised learning from traditional supervised machine learning, where the input is labeled in signal or background categories. Of the data meeting the above criteria, 90% is used for training, and 10% for validation. No blinding criterion is applied to the training data, so events that are used in training may also enter the final analysis regions. Since the VRNN performance is consistent across a variety of signal contamination fractions in training [39], and the presence of signal in the training dataset would only lower the sensitivity and cannot produce a false excess, biasing the final result is not a concern.

The loss function of the VRNN is composed of two terms: a reconstruction error term, to minimize differences between the decoded result and the original input, and the Kullback–Leibler (KL) divergence D_{KL} [52] of the encoded approximate posterior distribution from the Gaussian prior distribution. The loss $\mathcal{L}(t)$ for each time step t is given by Eq. (1), where $\mathbf{x}(t)$ is the input constituent four-vector, $\mathbf{y}(t)$ is the output constituent four-vector, z is the approximate posterior, z_t is the learned prior, and λ is a constant parameter to weight the D_{KL} contribution:

$$\mathcal{L}(t) = |\mathbf{y}(t) - \mathbf{x}(t)|^2 + \lambda D_{\text{KL}}(z||z_t). \quad (1)$$

Here, the time step of the RNN refers to the location in the input sequence. The learned prior z_t is a function of the current time step’s hidden state, and the hidden state is updated via a recurrence relation after each time step. An overall loss \mathcal{L} over the sequence is then computed by averaging the individual time-step losses over the length of the sequence.

The KL divergence of the encoded posterior distribution from the true posterior distribution is jointly minimized along with the loss $\mathcal{L}(t)$, ensuring that the architecture accurately describes the underlying distribution of data [53]. The D_{KL} term has been shown to provide better discrimination between anomalous and standard jets than either the reconstruction error or the loss term as a whole. The Anomaly Score is therefore defined in terms of the D_{KL} of each constituent, averaged over the whole jet, and restricted to the range of (0, 1) via exponentiation, as shown in Eq. (2). The resulting variable S_A ensures that more-anomalous jets populate higher values and background populates lower values, and is defined as

$$S_A = 1 - e^{-D_{\text{KL}}}. \quad (2)$$

The performance of the VRNN can be assessed by applying Eq. (2) to both background and a variety of signal models, and studying the discrimination power of S_A . Its performance in this analysis, and the use of the Anomaly Score to select the X jet, are described in the following section.

6 Event selection

The experimental signature of the $Y \rightarrow XH$ signal contains at least two jets with high transverse momentum. Signal-like events are selected through event-level criteria, along with jet-tagging criteria for both the X and H boson. In total, the analysis is performed three times, once for each of the three signal regions (SRs). The three analysis categories are defined using the selection criteria for the X particle. Each signal region utilizes five background estimation regions, composed of three control regions (CRs) and two validation regions (VRs), which are defined using common selections on the Higgs boson. The definitions of these regions and motivation for the selection criteria are provided in this section.

The analysis uses data events passing a high- p_T -threshold single large- R -jet trigger. Selected events must have a primary vertex and at least two large- R jets. Only the two highest- p_T large- R jets in an event are considered further, and at least one of their masses must exceed 50 GeV to remove poorly reconstructed events. To ensure that the trigger is fully efficient, the invariant mass of the two large- R jets must satisfy $m_{JJ} > 1.3$ TeV and the leading large- R jet, J_1 , must have $p_T(J_1) > 500$ GeV. These requirements, along with a selection on the D_{Hbb} score defined in the next subsection, compose the analysis preselection.

The primary SR, based on anomaly detection, is referred to as the *anomaly SR*. The other two SRs target the benchmark $X \rightarrow q\bar{q}$ decay and are thus called the *two-prong SRs*, differing in reconstruction of the X particle as either a single large- R jet (*merged SR*) or two small- R jets (*resolved SR*). They are not required to be orthogonal to the anomaly SR, but do not overlap with each other. The analysis CRs include the high sideband (HSB) of the Higgs boson candidate mass m_H , used to generate the DNN reweighting for the background estimation procedure, and CRO, which provides the final background template for the SR. The VRs are defined in the low sideband (LSB) of the Higgs boson candidate mass, and validate the reweighting derived from the HSBs. The full analysis flow is shown in Figure 2, with selection details provided below.

6.1 X/H ambiguity resolution

Since the SRs are constructed by placing requirements on the different properties of the X and H large- R jets, an ambiguity must be resolved to determine which of the two J in the event is more likely to be the Higgs boson and thus subject to the Higgs boson selection criteria. This is done by using a classifier based on a neural network (NN) to separate bosons decaying into $b\bar{b}$ from top-quark and QCD jets [54]. The tagger is trained on the large- R jet p_T and η , along with the subjet flavor-tagging score DL1r [55] for up to three track-subjets constructed with a jet- p_T -dependent radius parameter [56]. The tagger version used here includes a reweighting of all training inputs to have the same p_T and η distributions, to minimize biasing of the tagger towards high- p_T or central jets respectively. The outputs of the NN are three classification scores corresponding to the probability for the jet to have originated from a Higgs boson (P_{Higgs}), top quark (P_{top}), or multijet process (P_{multijet}), and these are subsequently combined into the jet-level discriminant D_{Hbb} :

$$D_{Hbb} = \ln \frac{P_{\text{Higgs}}}{f_{\text{top}} \cdot P_{\text{top}} + (1 - f_{\text{top}}) \cdot P_{\text{multijet}}}. \quad (3)$$




Figure 2: Illustration of the selection flow, after preselection, and the analysis regions of the $Y \rightarrow XH$ search. The first step in resolving the ambiguity in assigning the two large- R jets to X and H candidates dictates which jet is used to make the first selection on either D_2^{trk} or the Anomaly Score of the X candidate. This sorts the preselection events into three separate categories, namely anomaly, two-prong merged, and two-prong resolved. Each region has its own background estimation and validation regions. The CRs (CR0, HSB0, HSB1), VRs (LSB0, LSB1), and SR are defined in the same way for all three analyses using the H candidate mass (m_H) and D_{Hbb} score. The large- R jet J_H is required to have $D_{Hbb} > -2$.

In Eq. (3), f_{top} determines the weight assigned to the top-quark background shape in the final discriminant, and its chosen value of 0.25 is based on signal-to-noise optimization studies.

To perform the large- R jet ambiguity resolution, D_{Hbb} is computed for both J in the event. The jet with the larger value of D_{Hbb} is labeled as the Higgs boson candidate (J_H), and the other J is by default the X candidate (J_X), thus determining which jet is subject to further H and X selection. This procedure has an accuracy of over 90% in the highly boosted region of the signal grid ($m_X/m_Y < 0.5$), decreasing to no less than 75% for less boosted mass points where the boosted $H \rightarrow b\bar{b}$ tagging is less efficient. The resulting distribution of D_{Hbb} for the J chosen as the Higgs boson candidate, for both data and representative $Y \rightarrow XH$ signals at preselection, is shown in Figure 3. Additionally, a preselection of $D_{Hbb} > -2$ is applied to remove events that are determined to be not $H \rightarrow b\bar{b}$ -like, thus ensuring the data-driven background estimation focuses on a phase-space region that is close to that of the signal.

6.2 X selection

After the X candidate's J is determined, a region-dependent selection is applied to enhance the anomalous-jet content in the anomaly, two-prong merged, and two-prong resolved regions. The anomaly region is defined using the Anomaly Score value S_A of the X jet as determined by the VRNN. Figure 4 shows the resulting distribution of X candidate S_A values after the analysis preselection is applied, comparing




Figure 3: Distributions of the J_H candidate D_{Hbb} score in data after preselection requirements are applied. Also shown are three $Y \rightarrow XH$ simulated signals, labeled by the masses of the Y and X particles. Requiring $D_{Hbb} > 2.44$ defines a working point that is 60% efficient for the selection of the boosted $H \rightarrow b\bar{b}$ topology across the full p_T range. All distributions are normalized to unity.

data with three different $Y \rightarrow XH$ baseline signal samples as well as the three signals with alternative X decays. The absence of signal information in the construction of the score means that it is not expected to outperform analytic or supervised methods for any single signal model. However, using the VRNN approach to characterize the background distribution provides broad sensitivity to a variety of signals that differ from jets originating from QCD processes. The most notable difference in S_A shape is found for dark jets, which have a substructure that is not well-characterized by existing variables. The dependence of the sensitivity on kinematics is also relevant to performance, as the S_A distribution for the highly resolved point ($m_Y = 5000$ GeV, $m_X = 2500$ GeV) populates lower, and thus less anomalous, values than the data, making this signal indistinguishable from background in the anomaly detection approach. A selection $S_A > 0.5$ is chosen for the J_X candidate, and provides modest but comparable discrimination power in data for the disparate signal-jet topologies considered in the full expected p_T range of the X candidate jets. This enhances the signal-to-background ratio for the two boosted points, ($m_Y = 2000$ GeV, $m_X = 300$ GeV) and ($m_Y = 3400$ GeV, $m_X = 110$ GeV), by approximately 25% relative to the inclusive selection.




Figure 4: Distributions of the J_X candidate Anomaly Score (S_A) in data after preselection requirements are applied. Also shown are three $Y \rightarrow XH$ simulated signals (left), labeled by the masses of the Y and X particles, and the three additional signals with alternative X decay hypotheses, namely heavy-flavor, three-prong, and dark jet (right). A selection $S_A > 0.5$ applied to J_X defines the anomaly analysis category. All distributions are normalized to unity.

Regions focusing on the benchmark $X \rightarrow q\bar{q}$ decay are used to supplement the anomaly detection analysis and interpret the results. The merged region is defined by applying a selection $D_2^{\text{trk}} < 1.2$, where D_2^{trk} is the same as D_2 but computed using only tracks associated with the jet. The choice of using only tracks in the D_2 calculation is motivated by the ability to propagate track-only uncertainties into the final signal efficiency. Detailed comparisons of D_2^{trk} with the standard D_2 variable at analysis preselection have shown D_2^{trk} and D_2 to be highly compatible, ensuring its sensitivity to two-prong signals with respect to multijet background. Because jets with two-prong substructure have lower values of D_2^{trk} , this selection enhances the presence of fully merged decays where the X decay products are well-contained by a single J . Since the efficiency of substructure variables is mainly driven by the boost of the large- R jet and therefore its p_T , the optimal cut value of 1.2 was chosen so as to maximize signal-vs-background discrimination across the p_T range of the X candidate, combining all mass points in the $Y \rightarrow XH$ grid. Distributions of D_2^{trk} for J_X are shown in Figure 5 for both data and representative $Y \rightarrow XH$ signals at preselection.

Due to kinematic constraints driven by the m_X/m_Y ratio, events that fail the merged selection typically also show poor reconstruction of the X mass by the large- R jet. To achieve better sensitivity in this regime, a resolved selection is defined by requiring the D_2^{trk} value of J_X to be greater than 1.2, corresponding to a less boosted X particle which is more appropriately reconstructed using two small- R jets. A dedicated algorithm matches two small- R jets to the X by requiring at least four j in the event, discarding the two with the smallest ΔR to the Higgs boson candidate J_H , and selecting the two remaining small- R jets with the highest p_T . The mass of the Y resonance is then computed using the Higgs boson candidate large- R jet and the two small- R jets matched to the X . Additional requirements of $|\Delta y_{j1,j2}| < 2.5$ and a p_T imbalance $p_T^{\text{bal}} < 0.8$ are applied to aid accurate resolved X reconstruction; here Δy refers to separation in rapidity and p_T^{bal} is defined as the difference of the X particle's small- R jet transverse momenta divided by their sum. The resolved region improves the signal efficiency for points with higher m_X/m_Y (corresponding to less boost for the X candidate decay) by a factor of approximately five relative to the boosted selection.




Figure 5: Distributions of J_X candidate D_2^{trk} in data after preselection requirements are applied. Also shown are three $Y \rightarrow XH$ simulated signals, labeled by the masses of the Y and X particles. A value of $D_2^{\text{trk}} = 1.2$ therefore separates the analysis events into merged (< 1.2) and resolved (> 1.2) two-prong categories. All distributions are normalized to unity.

6.3 H selection

The Higgs boson tagging is performed after the J_X selection is applied, and sorts the events into the three analysis categories, for which the background is estimated. For all signal regions, a requirement

$D_{Hbb} > 2.44$ providing a constant 60% efficiency across jet p_T is applied to the Higgs boson candidate J_H , along with a mass window requirement of $75 < m_H < 145$ GeV. The data-driven background estimation uses events from the HSB of the Higgs boson candidate mass ($145 < m_H < 200$ GeV), which are further separated into HSB0 and HSB1 depending on whether the Higgs boson candidate J_H fails or passes the D_{Hbb} selection. Validation is performed in the LSB, where the reconstructed Higgs boson mass must be between 65 and 75 GeV. LSB0 and LSB1 are similarly defined as having Higgs boson candidates that fail or pass the D_{Hbb} tagging criterion, respectively. CR0 is defined as the set of events where the H boson is in the SR mass window but fails the D_{Hbb} tagging. DNN-based reweighting is applied to CR0 to obtain the final background prediction in the SR. The contamination from signal events is found to be of the order of a few percent in the HSB, and no more than 10% in the LSB, for all mass points in the $Y \rightarrow XH$ grid.

In total, 3 SRs and 15 background estimation regions are used in the analysis. A summary of all region definitions is provided in Table 1.

Table 1: Preselection requirements, and optimized requirements defining the SRs and background estimation regions.

Parameter	Preselection requirements				
m_{JJ} [GeV]	> 1300				
$p_T(J_1)$ [GeV]	> 500				
m_J [GeV]	$m_{J_1} > 50 \parallel m_{J_2} > 50$				
D_{Hbb}	> -2				
	Signal regions				
	Merged	Resolved	Anomaly		
m_H [GeV]	(75, 145)				
D_{Hbb}	> 2.44				
D_2^{trk}	< 1.2	> 1.2	-		
$ \Delta y_{j_1, j_2} $	-	< 2.5	-		
p_T^{bal}	-	< 0.8	-		
Anomaly Score (S_A)	-	-	> 0.5		
	Background estimation regions				
	CR0	HSB0	HSB1	LSB0	LSB1
m_H [GeV]	(75, 145)	(145, 200)		(65, 75)	
D_{Hbb}	< 2.44	< 2.44	> 2.44	< 2.44	> 2.44

6.4 Binning optimization

The signal search in the two-dimensional space of m_Y vs. m_X applies a variable-width sliding window to the J_X candidate mass spectrum, dividing the data into a series of overlapping m_X ranges in which the m_{JJ} distribution is fitted. The chosen binning for m_Y and m_X is based on the ATLAS detector's mass resolution for a generic narrow-width resonance, given by the fitted width of the Gaussian core of a double-sided Crystal Ball function. Modifications are made to the binning to account for the limited number of data events. In the m_Y spectrum, bins are widened at higher values of m_Y to ensure that at least one event is present in each bin when using LSB1 data. For the m_X categories, an initial set of mass windows is generated from a linear fit of m_X resolution as a function of m_X . The first window is chosen to have a bin center at $m_X = 65$ GeV, the lowest generated signal X mass, and its width is chosen to be twice the resolution obtained from the fit. Subsequent windows are chosen in the same way, with their bin centers

higher than the previous window’s bin center by half of that window’s mass resolution. The widths of high-mass windows are expanded symmetrically, based on event counts in LSB1, until at least 10 events are present in each final m_X bin. In order to avoid having duplicate windows, all resulting windows which share at least one bin edge are replaced by a single window encompassing their union. Finally, to accommodate the highest signal mass point at $m_X = 3000$ GeV, the last bin’s upper edge is placed at 3200 GeV.

7 Background estimation

The background in the SRs arises mainly from high- p_T multijet events. Simulation of such QCD processes is known to suffer from mismodeling and is computationally expensive. Therefore, in this analysis the background estimation is fully data-driven and uses regions that are orthogonal to the SR because of their Higgs boson jet criteria.

The shape of the expected m_{JJ} distribution in the SR is obtained from data in CR0, and weights are derived that can be applied to HSB0 to reproduce the shape found in HSB1. This procedure is validated by applying the weights to data in LSB0 and comparing the resulting m_{JJ} spectrum with that observed in LSB1 data. The weights are generated inclusively in the X candidate selection, and applied separately to create three background predictions, one for each SR.

The reweighting function is defined as the ratio of the multidimensional probability distribution function (pdf) of data in HSB1 to that of data in HSB0. In this analysis, the statistical procedure of *direct importance estimation* [57, 58] is utilized, where the ratio is estimated directly from data without having to analytically compute each individual PDF. It is implemented via the training of a DNN, where a custom square-root loss function [59] is chosen to produce weights that can accurately reproduce the observed ratio in data. The DNN outputs event-level weights that are assumed to be approximately independent of m_H and can be applied to an untagged region to produce the m_{JJ} shape in the corresponding $H \rightarrow b\bar{b}$ -tagged region.

Events are considered for training the DNN if they pass the analysis preselection, satisfy $145 < m_H < 175$ GeV, and additionally have at least two track-jets associated with the Higgs boson candidate J . They are modeled as a set of variables, namely the transverse momentum, pseudorapidity, azimuthal angle ϕ and energy of the Higgs boson candidate; the number of tracks associated with the Higgs boson candidate; and the transverse momentum, pseudorapidity, ϕ and mass of the first two track-jets associated with the Higgs boson candidate, ordered in p_T . Each variable x is standardized with the transformation $x = (x - \mu)/\sigma$, where μ and σ are the mean and the standard deviation of the distribution of the x variable.

The DNN is built using a fully connected sequential model from KERAS [60] with 3 inner layers, each with 20 neurons and a rectified linear unit activation function. In order to reduce the problem of overfitting during training, 10% of connections among inner layers are randomly severed (called “dropout”). The last layer has a single output with a simple linear activation function. The model is trained using the Adam optimizer [61] in KERAS, with TENSORFLOW [62] as its backend. Training is performed using a batch size equal to the full dataset size for 1600 epochs, with early stopping at 100 epochs if the value of the loss calculated on the validation dataset does not decrease for 100 subsequent epochs.

The output weights of the DNN are validated using data in the LSB. Figure 6 shows the impact of the reweighting on distributions of several key analysis variables, using the two-prong merged LSB VR as an example. Three curves are shown for each variable, comparing the LSB0 data, before and after DNN reweighting is applied, with the target data distribution in LSB1. These variables are chosen so as to focus on kinematic variables of the Higgs boson, whose properties are used to distinguish the CR from

the SR, and variables involved in the final statistical treatment. Good agreement between the reweighted shape and the true tagged data is observed for each variable, suggesting a robust background model. Since the training is performed inclusively in the X -tagging, the same conclusion holds for the anomaly and two-prong resolved LSB regions. Also, since the reweighting is applied to m_{JJ} distributions after the X -tagging selection is applied, no extrapolation across D_2^{trk} or S_A is required, and the method shows similar background modeling across all three SRs. Any minor residual differences between predicted background and data in the SRs are covered by the nonclosure systematic uncertainty, described in Section 8.




Figure 6: Distribution of (a) p_T and (b) η for the H candidate J , along with (c) the mass of J_X and (d) m_{JJ} in the merged LSB validation region, overlaying data from LSB1 (solid black points) with the data in LSB0 shown before (orange histogram) and after (red circles) reweighting. The ratio of the LSB1 data to both the LSB0 data (orange) and the reweighted LSB0 data (red) is shown in the lower panel. Error bars indicate statistical uncertainties.

8 Systematic uncertainties

Systematic uncertainties affect both the data-driven background estimate and the simulated signal. All background uncertainties are estimated using an m_{JJ} shape that is inclusive in m_X , and applied to each m_X category fully correlated among m_{JJ} bins. Background shape variations are derived from three sources.

The first is from training the DNN in an alternative region of $165 < m_H < 200$ GeV to account for the way in which phase-space uncertainties may affect the obtained weights. This region has approximately the same number of events and tagging efficiency as the nominal training region, helping to isolate the way in which changing only the DNN model affects the weights. Upward and downward variations are

defined by symmetrizing the shape difference in m_{JJ} between the two different models, yielding an effect of $\mathcal{O}(1-10)\%$ across the distribution.

Another DNN variation accounts for the finite size of the training sample and the random initialization of the weights. It is estimated with a bootstrap procedure [63, 64] where a set of 100 bootstrap networks are trained. For each training, the training dataset is varied by resampling it with replacement, assigning to each event a new weight that comes from a Poisson variable with a mean of one. Two additional templates are formed to create a symmetric error band by taking the median weight for each event and adding or subtracting half of the interquartile range. This corresponds to a $\mathcal{O}(1)\%$ effect across m_Y .

Lastly, a nonclosure uncertainty is included to cover modeling discrepancies that may arise when weights derived from the NN training in the HSB are extrapolated to the LSB, and subsequently to the SR. It is defined by the symmetrized shape difference between the data and predicted background in the LSB. To avoid being sensitive to statistical fluctuations in the LSB, smoothing is applied to the variation, where it is rebinned to reduce the relative statistical uncertainty. The nonclosure uncertainty is negligible for low m_{JJ} and rises to $\mathcal{O}(10)\%$ in the m_{JJ} tails.

Normalization and shape uncertainties are applied to the simulated signals. An uncertainty of 1.7% is applied to the luminosity measured with the LUCID-2 detector [65]. The uncertainty from the trigger selection is negligible, and thus not included, since the requirement of $m_{JJ} > 1.3$ TeV ensures that the trigger is fully efficient.

Scale factors (SFs) computed to match the $H \rightarrow b\bar{b}$ tagger efficiency between data and simulation are applied to the signal template, and their uncertainties are similarly propagated to the signal normalization. These SFs are computed using the methodology described in Ref. [66], with the substitution of an updated D_{Hbb} that includes the η -reweighting of inputs as described in Section 6. They are binned in large- R -jet p_T , where the SF for the highest p_T bin is extrapolated to cover the upper end of the p_T region probed by the analysis selection. Agreement between data and MC simulation for these SFs is verified with $Z \rightarrow b\bar{b}$ events in calibration studies. The SFs are bounded by approximately 1.1 and 1.4 across all p_T bins, with uncertainties in the range of approximately $\pm(0.3-0.5)$.

Instrumental systematic uncertainties arise from uncertainty in the jet p_T scale and resolution, for both the large- R and small- R jets, and affect the shape of the m_{JJ} distribution. Uncertainty in the large- R -jet p_T scale is an important effect in the search for resonant structures in the presence of rapidly falling background spectra, as it can shift the peak of the resonance. It is evaluated using tracking-to-calorimetric p_T double ratios between data and simulation, where any observed differences are assigned as baseline systematic uncertainties [67]. Even though the analysis relies on jets built from TCCs, the total p_T of the jet is still solely derived from calorimeter information, keeping it independent of the track-based jet p_T . Past analyses have studied the possible impact of calorimeter vs. track-based p_T by cross-calibrating between per-jet TCC and calorimeter p_T and found it to be negligible [30]. The total jet- p_T -scale uncertainty varies with p_T and η and is typically around $\pm 5\%$. Uncertainties from the reconstruction and modeling of tracks are also taken into account; these cover track reconstruction efficiency, impact parameter resolution, tracking in dense environments, fake-track rates, and sagitta biases. The impact of the large- R -jet p_T resolution uncertainty is evaluated event-by-event by rerunning the analysis with an additional absolute 2% Gaussian smearing applied to the input jets' p_T to degrade the nominal resolution. Small- R -jet p_T scale and resolution uncertainties are similarly estimated by comparing data with simulation after applying in situ corrections [68].

Several sources of theoretical uncertainty affecting the signal models were considered. Uncertainties in the matrix element calculation are evaluated by varying the strong coupling constant (α_s), the renormalization

scale (μ_r), and the factorization scale (μ_f). The effects of uncertainties in the parton distribution functions are evaluated by comparing the m_{JJ} distributions obtained when using various alternative PDF sets and taking an envelope of these distributions, as prescribed by the PDF4LHC group [69]. Generator-level variations of the A14 tune’s parameter values are used to cover the initial- and final-state radiation (ISR and FSR) and multiple parton interaction (MPI) uncertainties. An overall conservative 3% normalization uncertainty is applied to all signals as a result of ISR/FSR/MPI modeling effects.

9 Statistical analysis

The statistical framework in the analysis is used to perform hypothesis tests in the SRs, testing the compatibility of the data with both the background-only and signal-plus-background hypotheses. The observable that is fitted is the m_Y distribution of the data in the SR. This fit is repeated several times in overlapping bins of the X candidate mass.

The parameter of interest in the statistical analysis is the signal strength μ , defined as a scale factor multiplying the nominal yield predicted by a 1 pb signal cross section so as to match the observed number of signal events. The background-only hypothesis corresponds to $\mu = 0$. The normalization of the data-driven background estimate is allowed to float, with each normalization factor being fitted independently because the m_X categories overlap. A test statistic based on the profile likelihood ratio calculated in the lowest-order asymptotic approximation is used to test the models corresponding to the signal grid. Systematic uncertainties are included in the fit as nuisance parameters (NPs) with Gaussian constraints. Both the signal strength and all signal systematic uncertainty NPs are correlated across the merged and resolved regions. The significance of an observed excess of data over the background prediction is quantified by the local p -value, which is the probability that the background-only model produces an excess at least as large as the one observed.

BUMPHUNTER [70] is used to find excesses in both the anomaly and two-prong SRs, with an algorithm that incorporates only the statistical uncertainty of the data and does not depend on a specific signal shape. It outputs a p -value that provides a goodness-of-fit metric, along with an interval of the invariant mass corresponding to the largest deviation of the data from the background prediction. It is thus used in all signal regions to search for significant deviations of the data from the background m_Y distribution. Dedicated studies indicate the fits are dominated by statistical uncertainties, further motivating the discounting of systematic uncertainties in the BUMPHUNTER fits. In the anomaly SR, no subsequent fits are performed using the $Y \rightarrow XH$ signal model, as this analysis is designed to keep signal-model dependence minimal and the two-prong regions are expected to have higher signal sensitivity across the $Y \rightarrow XH$ grid. However, given the absence of a significant excess in the two-prong regions, signal-plus-background fits are performed for each $Y \rightarrow XH$ signal model, and 95% confidence level (CL) upper limits on the signal cross section are set using a modified frequentist method (CL_s) [71].

The background estimation and statistical treatment are validated by comparison with data in the signal-depleted LSB VR for each of the three analysis categories. BUMPHUNTER p -values from all m_X -bin fits in the VR of the anomaly region approximate a flat distribution between 0 and 1, indicating good background modeling with no systematic biases across the phase space. Figure 7 shows the post-fit background prediction compared with the data passing the anomaly, merged, and resolved selections, in an example m_X window between 284.5 and 322.5 GeV where the whole kinematic phase space is well-populated. Good modeling is observed, and no significant pulls or constraints are observed in the background NPs. In the two-prong regions, checks are made for spurious signals in the LSB through a signal-plus-background fit

using each generated signal model. These checks indicate that no significant spurious signal is present beyond those likely to be produced by statistical fluctuations, with no systematic trend across the expected phase space.




Figure 7: Reconstructed m_γ distributions of the background, as determined by a background-only fit, and data in the LSB1 VR, for the anomaly, two-prong merged, and two-prong resolved selections, in the m_X bin [284.5, 322.5] GeV. The ratio of the observed data to the background is shown in the lower panel. The uncertainty band includes both the statistical and systematic effects.

10 Results

Results of background-only fits of the m_Y distribution across all m_X categories in the anomaly SR show the data to be compatible with the expected backgrounds given their uncertainties. Figure 8 shows the distribution of p -values across m_Y and m_X bins, where the background is the result of a background-only fit performed with all statistical and systematic uncertainties, and all uncertainties are included in the p -value calculation. The distribution of observed p -values is statistically compatible with the expectation, with a χ -squared value per degree of freedom of 1.32 between the distributions.




Figure 8: The distribution of observed p -values across all m_Y and m_X bins in the anomaly signal region, comparing data with the background estimates generated by a background-only fit, displayed in the two-dimensional (m_X , m_Y) grid. The p -value calculation is performed at the center of each m_X bin, and all statistical and background systematic uncertainties are considered. The lowest observed p -value corresponds to the bin with m_Y within [3608, 3805] GeV and m_X within [75.5, 95.5] GeV.

The largest observed excess in the anomaly SR is in the m_X window [75.5, 95.5] GeV. In this window, the greatest single-bin excess is found for m_Y between 3608 and 3805 GeV. The BUMPHUNTER interval with the greatest significance covers the m_Y range between bin edges of 3424 and 3805 GeV. The m_Y distribution corresponding to this m_X window is shown in Figure 9, along with the post-fit expected background. Studies of the individual jet mass distributions in this m_X category do not reveal any excesses in data that are consistent with a resonant particle decay. Given the number of individual search regions in this analysis, the impact of the trials factor is significant. A calculation is made to determine the global significance of the greatest single-bin deviation, accounting for all the overlapping m_X bins. First, the overlapping bin edges are used to define exclusive, nonoverlapping bins in m_X , and an integer is drawn from a Poisson distribution with a mean equal to the background expectation in each exclusive (m_Y , m_X) bin. Next, this yield is summed across exclusive bins to create a toy estimate for each overlapping bin in which the p -value is computed. Finally, this procedure is repeated N times, where N is the number of events, inclusively across all exclusive, nonoverlapping bins. This calculation yields a global significance of 1.43σ for the excess in the [3608, 3805] GeV bin of m_Y .

Results for the two-prong SRs are similarly derived by performing background-only fits and scanning with BUMPHUNTER for incompatibility with data. No significant deviations of data from the predicted




Figure 9: The m_Y distribution associated with the m_X bin [75.5, 95.5] GeV in the anomaly SR, where the background is determined by a background-only fit to data with both the statistical and systematic uncertainties included. The p -value and the lower panel of per-bin significances are computed with statistical uncertainties only. The single m_Y bin with the most significant excess is [3608, 3805] GeV. Incorporating the trials factor for all m_Y and overlapping m_X bins in the search, this excess has a global significance of 1.43σ .

background are observed beyond expected statistical fluctuations, in either the merged or resolved SR. An example m_Y distribution in both the merged and resolved SRs for the m_X bin [284.5, 322.5] GeV is shown in Figure 10, along with the background estimate that is determined from a background-only fit accounting for all uncertainties. Figure 11 shows a summary of the per-bin p -values in each m_Y bin for selected m_X bins that are centered on key X candidate mass hypotheses, namely at the W , Z , and H boson masses.

Given the absence of significant excesses in the data, signal-plus-background fits are performed to determine the 95% CL upper limit on the cross section of the $Y \rightarrow XH$ process. A summary of the expected and observed limits in the 2D grid of $Y \rightarrow XH$ signals is given in Figure 12, combining results from the merged and resolved regions. A bilinear interpolation procedure is applied to provide results between fully simulated signal points. The analysis is most sensitive in the very boosted regime, where the Y mass is approximately an order of magnitude larger than the X mass. Sensitivity is lowest in the highly resolved regime, where the required large- R J reconstruction of the Higgs boson sculpts the signal efficiency to favor high-momentum X particles. The observed limits range from cross sections of 0.341 fb for the signal point ($m_Y = 5000$ GeV, $m_X = 600$ GeV), to 1.22 pb for the signal point ($m_Y = 2500$ GeV, $m_X = 2000$ GeV).

The data in the anomaly and two-prong SRs can be used to provide a benchmark comparison of sensitivity across the set of large- R jet topologies considered for the X decay, thereby providing a metric for assessing the level of signal-model-dependence in both regions. The 95% CL upper limits on the production cross section of several benchmark signals are calculated for all three signal region selections, by injecting signal into the data until the BUMP HUNTER p -value exceeds a significance of 2σ . Seven signal points are considered in this study, including three highly boosted $Y \rightarrow XH$ points, one resolved $Y \rightarrow XH$ point, and the three alternative jet topologies. Because the systematic uncertainties in the signal efficiency of the Anomaly Score are not assessed, this comparison is performed using only statistical uncertainties and a post-fit background estimate in the limit calculation.




Figure 10: Reconstructed m_Y distributions of background and data in the SR, for the merged (left) and resolved (right) selections, in the m_X bin [284.5, 322.5] GeV. The background is determined by a background-only fit to the data with all statistical and systematic uncertainties included. The ratio of the observed data to the background is shown in the lower panel. The uncertainty band includes both the statistical and systematic effects.

Since the merged region uses D_2^{trk} and thus explicitly tags on the two-prong substructure of the X candidate's large- R jet in the generated $Y \rightarrow XH$ grid, it is possible that this region will outperform the fully unsupervised approach for the $Y \rightarrow XH$ signals. Figure 13 provides a summary of the limits obtained from the signal injection tests, comparing all three signal region selections for the seven benchmark signal points. For points where the X particle is highly boosted and thus the Anomaly Score is most sensitive, the upper limit on the cross section is approximately the same across the merged and anomaly SRs. The D_2^{trk} selection for this analysis is a general one; further optimization of the D_2^{trk} selection would be expected to provide better performance than the Anomaly Score across two-prong simulated signals. However, the signal-model-independent aspect of the anomaly detection used in the anomaly SR is more evident through improved limits on the alternative substructure signals. Notably, for the dark jets the cross-section upper limit is nearly an order of magnitude lower than that provided by the two-prong regions, which underlines the particular strength of the model-independent approach for signals that are challenging to characterize with existing high-level variables.

While these results cover regions of phase space that have not been studied directly by other searches, some analysis selections overlap with those of other recent ATLAS dijet resonance searches. The m_X bin of [75.5, 95.5] GeV would be sensitive to a VH resonance's hadronic final state, which is covered by a dedicated analysis using the same dataset [72]. The present approach differs in the tagging techniques used for both the vector boson and Higgs boson, but provides a similar 95% CL upper limit on the production cross section of a 3 TeV resonance. Due to its generality, the anomaly SR is expected to be sensitive to the same signatures as the weakly supervised dijet resonance search [13], but a direct comparison is not provided due to the assumptions made here about the mass and decay of the Higgs boson candidates.




Figure 11: The p -value per m_Y bin for both two-prong SRs, calculated using all systematic and statistical uncertainties of the background estimate. Two m_X bins are shown, $[75.5, 95.5]$ and $[113.0, 137.0]$ GeV, which corresponds to a window containing the W/Z and Higgs boson mass respectively. Events thus fall in (a) a merged W/Z window, (b) a merged Higgs window, (c) a resolved W/Z window, and (d) a resolved Higgs window. The background is determined by a background-only fit to the data with all statistical and systematic uncertainties included. In both m_X windows, the p -value has a high approximately constant value for the high Y -mass region of the resolved SR (bottom), as this phase-space region is far more likely to produce a highly boosted J_X that falls in the merged SR selection.




Figure 12: The expected (left) and observed (right) 95% CL limits on the cross section $\sigma(pp \rightarrow Y \rightarrow XH \rightarrow q\bar{q}b\bar{b})$ in pb in the two-dimensional space of m_Y versus m_X , obtained from a simultaneous fit of both the merged and resolved two-prong signal regions with all statistical and systematic uncertainties. A bilinear interpolation procedure is applied to provide results between fully simulated signal points. The observed limits range from 0.34 fb for the signal point ($m_Y = 5000$ GeV, $m_X = 600$ GeV) to 1.22 pb for the signal point ($m_Y = 2500$ GeV, $m_X = 2000$ GeV). The boundaries of the limit are defined by the simulated signal grid.




Figure 13: The 95% CL upper limit on the cross section for seven benchmark signal processes, comparing the anomaly, two-prong merged, and two-prong resolved signal region selections. The cross section is obtained by injecting signal into the observed data until 2σ sensitivity is achieved as determined by the `BUMPHUNTER` interval p -value. The background estimate is the result of a fit with all background uncertainties considered, while the `BUMPHUNTER` algorithm incorporates only statistical uncertainties. The uncertainty band reflects the standard deviation of obtained cross section values for 100 trials. All signals are labeled by the masses of the BSM particles except the $A \rightarrow BC$ signals, where $m_A = 3 \text{ TeV}$, $m_B = 200 \text{ GeV}$, and $m_C = 400 \text{ GeV}$. Compared to the D_2^{trk} -defined regions, the anomaly SR performs competitively for all signals considered, and considerably better for the dark jet signal, indicating a high degree of signal-model-agnostic sensitivity. The red upward-pointing arrow for $A \rightarrow BC \rightarrow qq qq qq$ indicates that the limit obtained from the two-pronged (merged) SR lies above the upper edge of the plot.

11 Conclusion

A search is performed for a heavy new boson Y decaying into a new particle X and a Standard Model Higgs boson in 139 fb^{-1} of 13 TeV pp collision data collected by the ATLAS detector during LHC Run 2. The analysis focuses on a fully hadronic final state, where the X particle and H boson are boosted such that their daughter particles are collimated. In the first application of fully unsupervised machine learning to an ATLAS search, a VRNN is trained on jets in data to define an anomaly detection SR, which selects the X particle based solely on its substructural incompatibility with background jets. Two supplementary SRs are designed to separately reconstruct merged and resolved decays of a nominal two-prong X benchmark signal. Sensitivity over the dominant multijet background is enhanced by additional machine-learning applications, namely a NN-based $H \rightarrow b\bar{b}$ tagger and a DNN-based reweighting to ensure good modeling. No significant deviations of the data from the predicted background are observed. The largest excess is found in the anomaly SR with a global significance of 1.43σ when considering all m_X and m_Y bins, and is not found to be compatible with the expected signal shape. Results are interpreted as 95% confidence-level upper limits on the cross section $\sigma(pp \rightarrow Y \rightarrow XH \rightarrow q\bar{q}b\bar{b})$, across the two-dimensional space where m_Y is between 1.5 and 6 TeV and m_X is between 65 and 3000 GeV. The most stringent upper limit of 0.341 fb is obtained in the merged SR for the signal point ($m_Y = 5000 \text{ GeV}$, $m_X = 600 \text{ GeV}$), while the weakest upper limit of 1.22 pb is obtained for the highly resolved point ($m_Y = 2500 \text{ GeV}$, $m_X = 2000 \text{ GeV}$).

Acknowledgments

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada and CRC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC

(Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [\[73\]](#).

References

- [1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1, arXiv: 1207.7214 [hep-ex].
- [2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30, arXiv: 1207.7235 [hep-ex].
- [3] G. Kasieczka et al., *The LHC Olympics 2020 a community challenge for anomaly detection in high energy physics*, *Rep. Prog. Phys.* **84** (2021) 124201, URL: <https://doi.org/10.1088%2F1361-6633%2Fac36b9>.
- [4] T. Aarrestad et al., *The Dark Machines Anomaly Score Challenge: Benchmark Data and Model Independent Event Classification for the Large Hadron Collider*, *SciPost Physics* **12** (2022), URL: <https://doi.org/10.21468%2Fscipostphys.12.1.043>.
- [5] G. Karagiorgi, G. Kasieczka, S. Kravitz, B. Nachman, and D. Shih, *Machine Learning in the Search for New Fundamental Physics*, 2021, URL: <https://arxiv.org/abs/2112.03769>.
- [6] G. Altarelli, B. Mele, and M. Ruiz-Altaba, *Searching for new heavy vector bosons in $p\bar{p}$ colliders*, *Z. Phys. C* **45** (1989) 109, [Erratum: *Z.Phys.C* 47, 676 (1990)].
- [7] K. S. Agashe et al., *LHC signals from cascade decays of warped vector resonances*, *JHEP* **05** (2017) 078, ISSN: 1029-8479, URL: [http://dx.doi.org/10.1007/JHEP05\(2017\)078](http://dx.doi.org/10.1007/JHEP05(2017)078).
- [8] L. Randall and R. Sundrum, *Large Mass Hierarchy from a Small Extra Dimension*, *Phys. Rev. Lett.* **83** (1999) 3370, ISSN: 1079-7114, URL: <http://dx.doi.org/10.1103/PhysRevLett.83.3370>.
- [9] T. Han, J. D. Lykken, and R.-J. Zhang, *Kaluza-Klein states from large extra dimensions*, *Phys. Rev. D* **59** (1999) 105006, ISSN: 1089-4918, URL: <http://dx.doi.org/10.1103/PhysRevD.59.105006>.
- [10] G. Branco et al., *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rep.* **516** (2012) 1, ISSN: 0370-1573, URL: <http://dx.doi.org/10.1016/j.physrep.2012.02.002>.
- [11] D. Pappadopulo, A. Thamm, R. Torre, and A. Wulzer, *Heavy vector triplets: bridging theory and data*, *JHEP* **9** (2014) 060, ISSN: 1029-8479, URL: [http://dx.doi.org/10.1007/JHEP09\(2014\)060](http://dx.doi.org/10.1007/JHEP09(2014)060).
- [12] ATLAS Collaboration, *A search for resonances decaying into a Higgs boson and a new particle X in the $XH \rightarrow qqbb$ final state with the ATLAS detector*, *Phys. Lett. B* **779** (2018) 24, ISSN: 0370-2693, URL: <http://dx.doi.org/10.1016/j.physletb.2018.01.042>.
- [13] ATLAS Collaboration, *Dijet Resonance Search with Weak Supervision Using $\sqrt{s} = 13$ TeV pp collisions in the ATLAS detector*, *Phys. Rev. Lett.* **125** (2020) 131801, arXiv: 2005.02983 [hep-ex].
- [14] CMS Collaboration, *Search for new heavy resonances decaying to WW , WZ , ZZ , WH , or ZH boson pairs in the all-jets final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2022, arXiv: 2210.00043 [hep-ex].

- [15] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, [JINST 3 \(2008\) S08003](#).
- [16] ATLAS Collaboration, *ATLAS Insertable B-Layer Technical Design Report*, ATLAS-TDR-19; CERN-LHCC-2010-013, 2010, URL: <https://cds.cern.ch/record/1291633>, Addendum: ATLAS-TDR-19-ADD-1; CERN-LHCC-2012-009, 2012, URL: <https://cds.cern.ch/record/1451888>.
- [17] B. Abbott et al., *Production and integration of the ATLAS Insertable B-Layer*, [JINST 13 \(2018\) T05008](#), arXiv: [1803.00844 \[physics.ins-det\]](#).
- [18] ATLAS Collaboration, *The ATLAS Collaboration Software and Firmware*, ATL-SOFT-PUB-2021-001, 2021, URL: <https://cds.cern.ch/record/2767187>.
- [19] ATLAS Collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, [JINST 15 \(2020\) P04003](#), arXiv: [1911.04632 \[physics.ins-det\]](#).
- [20] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC*, [Eur. Phys. J. C 76 \(2016\) 653](#), arXiv: [1608.03953 \[hep-ex\]](#).
- [21] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, [JINST 13 \(2018\) P07017](#).
- [22] ATLAS Collaboration, *Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1*, [Eur. Phys. J. C 77 \(2017\) 490](#), arXiv: [1603.02934 \[hep-ex\]](#).
- [23] ATLAS Collaboration, *The ATLAS Simulation Infrastructure*, [Eur. Phys. J. C 70 \(2010\) 823](#), arXiv: [1005.4568 \[physics.ins-det\]](#).
- [24] GEANT4 Collaboration, S. Agostinelli, et al., *GEANT4 – a simulation toolkit*, [Nucl. Instrum. Meth. A 506 \(2003\) 250](#).
- [25] J. de Blas, J. M. Lizana, and M. Pérez-Victoria, *Combining searches of Z' and W' bosons*, [JHEP 1 \(2013\) 166](#), URL: <https://doi.org/10.1007%2Fjhep01%282013%29166>.
- [26] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, [JHEP 07 \(2014\) 079](#), arXiv: [1405.0301 \[hep-ph\]](#).
- [27] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, [Comput. Phys. Commun. 178 \(2008\) 852](#), arXiv: [0710.3820 \[hep-ph\]](#).
- [28] R. D. Ball et al., *Impact of heavy quark masses on parton distributions and LHC phenomenology*, [Nucl. Phys. B 849 \(2011\) 296](#), arXiv: [1101.1300 \[hep-ph\]](#).
- [29] ATLAS Collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, tech. rep., ATL-PHYS-PUB-2014-021: CERN, 2014, URL: <https://cds.cern.ch/record/1966419>.
- [30] ATLAS Collaboration, *Search for diboson resonances in hadronic final states in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [JHEP 09 \(2019\) 091](#), [Erratum: [JHEP 06 \(2020\) 042](#)], arXiv: [1906.08589 \[hep-ex\]](#).
- [31] M. Park and M. Zhang, *Tagging a jet from a dark sector with jet substructure at colliders*, [Phys. Rev. D 100 \(2019\) 115009](#), arXiv: [1712.09279 \[hep-ph\]](#).
- [32] M. Cacciari, G. P. Salam, and G. Soyez, *The anti- k_t jet clustering algorithm*, [JHEP 04 \(2008\) 063](#), arXiv: [0802.1189 \[hep-ph\]](#).

- [33] M. Cacciari, G. P. Salam, and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896, arXiv: [1111.6097](https://arxiv.org/abs/1111.6097) [hep-ph].
- [34] ATLAS Collaboration, *Improving jet substructure performance in ATLAS using Track-CaloClusters*, tech. rep., ATL-PHYS-PUB-2017-015: CERN, 2017, URL: <https://cds.cern.ch/record/2275636>.
- [35] D. Krohn, J. Thaler, and L.-T. Wang, *Jet trimming*, *JHEP* **02** (2010) 084, arXiv: [0912.1342](https://arxiv.org/abs/0912.1342) [hep-ph].
- [36] ATLAS Collaboration, *Identification of boosted, hadronically-decaying W and Z bosons in $\sqrt{s} = 13$ TeV Monte Carlo Simulations for ATLAS*, tech. rep., ATL-PHYS-PUB-2015-033: CERN, 2015, URL: <https://cds.cern.ch/record/2041461>.
- [37] ATLAS Collaboration, *Jet reconstruction and performance using particle flow with the ATLAS Detector*, *Eur. Phys. J. C* **77** (2017) 466, arXiv: [1703.10485](https://arxiv.org/abs/1703.10485) [hep-ex].
- [38] ATLAS Collaboration, *Selection of jets produced in 13TeV proton-proton collisions with the ATLAS detector*, tech. rep., ATLAS-CONF-2015-029: CERN, 2015, URL: <https://cds.cern.ch/record/2037702>.
- [39] A. Kahn, J. Gonski, I. Ochoa, D. Williams, and G. Brooijmans, *Anomalous jet identification via sequence modeling*, *JINST* **16** (2021) P08012, URL: <https://doi.org/10.1088%2F1748-0221%2F16%2F08%2Fp08012>.
- [40] J. Chung et al., *A Recurrent Latent Variable Model for Sequential Data*, 2015, URL: <https://arxiv.org/abs/1506.02216>.
- [41] T. Heimel, G. Kasieczka, T. Plehn, and J. Thompson, *QCD or what?* *SciPost Physics* **6** (2019), URL: <https://doi.org/10.21468%2Fscipostphys.6.3.030>.
- [42] M. Farina, Y. Nakai, and D. Shih, *Searching for new physics with deep autoencoders*, *Phys. Rev. D* **101** (2020) 075021, URL: <https://doi.org/10.1103%2Fphysrevd.101.075021>.
- [43] B. Bortolato, A. Smolkovič, B. M. Dillon, and J. F. Kamenik, *Bump hunting in latent space*, *Phys. Rev. D* **105** (2022) 115009, URL: <https://doi.org/10.1103%2Fphysrevd.105.115009>.
- [44] T. Cheng, J.-F. Arguin, J. Leissner-Martin, J. Pilette, and T. Golling, *Variational autoencoders for anomalous jet tagging*, *Phys. Rev. D* **107** (2023) 016002, URL: <https://doi.org/10.1103%2Fphysrevd.107.016002>.
- [45] A. Blance, M. Spannowsky, and P. Waite, *Adversarially-trained autoencoders for robust unsupervised new physics searches*, *JHEP* **10** (2019) 047, arXiv: [1905.10384](https://arxiv.org/abs/1905.10384) [hep-ph].
- [46] O. Cerri, T. Q. Nguyen, M. Pierini, M. Spiropulu, and J.-R. Vlimant, *Variational autoencoders for new physics mining at the Large Hadron Collider*, *JHEP* **05** (2019) 036, arXiv: [1811.10276](https://arxiv.org/abs/1811.10276) [hep-ex].
- [47] A. Paszke et al., *PyTorch: An Imperative Style, High-Performance Deep Learning Library*, 2019, arXiv: [1912.01703](https://arxiv.org/abs/1912.01703) [cs.LG].
- [48] D. P. Kingma and J. Ba, *Adam: A Method for Stochastic Optimization*, 2017, arXiv: [1412.6980](https://arxiv.org/abs/1412.6980) [cs.LG].
- [49] A. J. Larkoski, G. P. Salam, and J. Thaler, *Energy correlation functions for jet substructure*, *JHEP* **6** (2013) 108, URL: <https://doi.org/10.1007%2Fjhep06%282013%29108>.

- [50] J. Thaler and K. Van Tilburg, *Identifying boosted objects with N -subjettiness*, *JHEP* **3** (2011) 015, ISSN: 1029-8479, URL: [http://dx.doi.org/10.1007/JHEP03\(2011\)015](http://dx.doi.org/10.1007/JHEP03(2011)015).
- [51] S. Marzani, G. Soyez, and M. Spannowsky, *Looking Inside Jets*, *Lecture Notes in Physics* (2019), ISSN: 1616-6361, URL: <http://dx.doi.org/10.1007/978-3-030-15709-8>.
- [52] S. Kullback, *Information Theory and Statistics*, New York: Wiley, 1959, ISBN: 978-0486696843.
- [53] D. P. Kingma and M. Welling, *Auto-Encoding Variational Bayes*, 2013, URL: <https://arxiv.org/abs/1312.6114>.
- [54] ATLAS Collaboration, *Identification of Boosted Higgs Bosons Decaying Into $b\bar{b}$ With Neural Networks and Variable Radius Subjets in ATLAS*, tech. rep., ATL-PHYS-PUB-2020-019: CERN, 2020, URL: <https://cds.cern.ch/record/2724739>.
- [55] ATLAS Collaboration, *ATLAS b -jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970, arXiv: [1907.05120](https://arxiv.org/abs/1907.05120) [hep-ex].
- [56] D. Krohn, J. Thaler, and L.-T. Wang, *Jets with variable R* , *JHEP* **06** (2009) 059, URL: <https://doi.org/10.1088/1126-6708/2009/06/059>.
- [57] T. Hastie, R. Tibshirani, and J. Friedman, *The Elements of Statistical Learning*, Springer Series in Statistics, New York, NY, USA: Springer New York Inc., 2001.
- [58] M. Sugiyama, T. Suzuki, and T. Kanamori, *Density Ratio Estimation in Machine Learning*, Cambridge University Press, 2012.
- [59] B. Nachman and J. Thaler, *Learning from many collider events at once*, *Phys. Rev. D* **103** (2021) 116003, URL: <https://doi.org/10.1103/PhysRevD.103.116013>.
- [60] F. Chollet et al., *Keras*, 2015, URL: <https://keras.io>.
- [61] D. P. Kingma and J. Ba, *Adam: A Method for Stochastic Optimization*, (2017), arXiv: [1412.6980](https://arxiv.org/abs/1412.6980) [cs.LG].
- [62] Martín Abadi et al, *TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems*, Software available from tensorflow.org, 2015, arXiv: [1603.04467](https://arxiv.org/abs/1603.04467) [cs.DC], URL: <https://www.tensorflow.org/>.
- [63] B. Efron, *Bootstrap Methods: Another Look at the Jackknife*, *Ann. Statist.* **7** (1979) 1, URL: <https://doi.org/10.1214/aos/1176344552>.
- [64] U. Michelucci and F. Venturini, *Estimating Neural Network's Performance with Bootstrap: A Tutorial*, *Mach. Learn. Knowl. Extr.* **3** (2021) 357, ISSN: 2504-4990, URL: <https://www.mdpi.com/2504-4990/3/2/18>.
- [65] ATLAS Collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, tech. rep., CERN, 2019, URL: <https://cds.cern.ch/record/2677054>.
- [66] ATLAS Collaboration, *Efficiency corrections for a tagger for boosted $H \rightarrow b\bar{b}$ decays in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, tech. rep., CERN, 2021, URL: <https://cds.cern.ch/record/2777811>.
- [67] ATLAS Collaboration, *Performance of jet substructure techniques for large- R jets in proton–proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector*, *JHEP* **09** (2013) 076, arXiv: [1306.4945](https://arxiv.org/abs/1306.4945) [hep-ex].

- [68] ATLAS Collaboration, *Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2020) 689, arXiv: [2007.02645](https://arxiv.org/abs/2007.02645) [[hep-ex](#)].
- [69] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys. G* **43** (2016) 023001, arXiv: [1510.03865](https://arxiv.org/abs/1510.03865) [[hep-ph](#)].
- [70] G. Choudalakis, *On hypothesis testing, trials factor, hypertexts and the BumpHunter*, 2011, URL: <https://arxiv.org/abs/1101.0390>.
- [71] A. L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693.
- [72] ATLAS Collaboration, *Search for resonances decaying into a weak vector boson and a Higgs boson in the fully hadronic final state produced in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **102** (2020) 112008, arXiv: [2007.05293](https://arxiv.org/abs/2007.05293) [[hep-ex](#)].
- [73] ATLAS Collaboration, *ATLAS Computing Acknowledgements*, ATL-SOFT-PUB-2021-003, 2021, URL: <https://cds.cern.ch/record/2776662>.

The ATLAS Collaboration

G. Aad ¹⁰¹, B. Abbott ¹¹⁹, D.C. Abbott ¹⁰², K. Abeling ⁵⁵, S.H. Abidi ²⁹, A. Aboulhorma ^{35e}, H. Abramowicz ¹⁵⁰, H. Abreu ¹⁴⁹, Y. Abulaiti ¹¹⁶, A.C. Abusleme Hoffman ^{136a}, B.S. Acharya ^{68a,68b,q}, B. Achkar ⁵⁵, C. Adam Bourdarios ⁴, L. Adamczyk ^{84a}, L. Adamek ¹⁵⁴, S.V. Addepalli ²⁶, J. Adelman ¹¹⁴, A. Adiguzel ^{21c}, S. Adorni ⁵⁶, T. Adye ¹³³, A.A. Affolder ¹³⁵, Y. Afik ³⁶, M.N. Agaras ¹³, J. Agarwala ^{72a,72b}, A. Aggarwal ⁹⁹, C. Agheorghiesei ^{27c}, J.A. Aguilar-Saavedra ^{129f}, A. Ahmad ³⁶, F. Ahmadov ^{38,z}, W.S. Ahmed ¹⁰³, S. Ahuja ⁹⁴, X. Ai ⁴⁸, G. Aielli ^{75a,75b}, I. Aizenberg ¹⁶⁸, M. Akbiyik ⁹⁹, T.P.A. Åkesson ⁹⁷, A.V. Akimov ³⁷, K. Al Khoury ⁴¹, G.L. Alberghi ^{23b}, J. Albert ¹⁶⁴, P. Albicocco ⁵³, S. Alderweireldt ⁵², M. Aleksa ³⁶, I.N. Aleksandrov ³⁸, C. Alexa ^{27b}, T. Alexopoulos ¹⁰, A. Alfonsi ¹¹³, F. Alfonsi ^{23b}, M. Alhroob ¹¹⁹, B. Ali ¹³¹, S. Ali ¹⁴⁷, M. Aliev ³⁷, G. Alimonti ^{70a}, W. Alkahi ⁵⁵, C. Allaire ⁶⁶, B.M.M. Allbrooke ¹⁴⁵, P.P. Allport ²⁰, A. Aloisio ^{71a,71b}, F. Alonso ⁸⁹, C. Alpigiani ¹³⁷, E. Alunno Camelia ^{75a,75b}, M. Alvarez Estevez ⁹⁸, M.G. Alvigi ^{71a,71b}, M. Aly ¹⁰⁰, Y. Amaral Coutinho ^{81b}, A. Ambler ¹⁰³, C. Amelung ³⁶, M. Amerl ¹, C.G. Ames ¹⁰⁸, D. Amidei ¹⁰⁵, S.P. Amor Dos Santos ^{129a}, S. Amoroso ⁴⁸, K.R. Amos ¹⁶², V. Ananiev ¹²⁴, C. Anastopoulos ¹³⁸, T. Andeen ¹¹, J.K. Anders ³⁶, S.Y. Andrean ^{47a,47b}, A. Andreatta ^{70a,70b}, S. Angelidakis ⁹, A. Angerami ^{41,ab}, A.V. Anisenkov ³⁷, A. Annovi ^{73a}, C. Antel ⁵⁶, M.T. Anthony ¹³⁸, E. Antipov ¹²⁰, M. Antonelli ⁵³, D.J.A. Antrim ^{17a}, F. Anulli ^{74a}, M. Aoki ⁸², T. Aoki ¹⁵², J.A. Aparisi Pozo ¹⁶², M.A. Aparo ¹⁴⁵, L. Aperio Bella ⁴⁸, C. Appelt ¹⁸, N. Aranzabal ³⁶, V. Araujo Ferraz ^{81a}, C. Arcangeletti ⁵³, A.T.H. Arce ⁵¹, E. Arena ⁹¹, J-F. Arguin ¹⁰⁷, S. Argyropoulos ⁵⁴, J.-H. Arling ⁴⁸, A.J. Armbruster ³⁶, O. Arnaez ¹⁵⁴, H. Arnold ¹¹³, Z.P. Arrubarrena Tame ¹⁰⁸, G. Artoni ^{74a,74b}, H. Asada ¹¹⁰, K. Asai ¹¹⁷, S. Asai ¹⁵², N.A. Asbah ⁶¹, J. Assahsah ^{35d}, K. Assamagan ²⁹, R. Astalos ^{28a}, R.J. Atkin ^{33a}, M. Atkinson ¹⁶¹, N.B. Atlay ¹⁸, H. Atmani ^{62b}, P.A. Atmasiddha ¹⁰⁵, K. Augsten ¹³¹, S. Auricchio ^{71a,71b}, A.D. Auriol ²⁰, V.A. Austrup ¹⁷⁰, G. Avner ¹⁴⁹, G. Avolio ³⁶, K. Axiotis ⁵⁶, M.K. Ayoub ^{14c}, G. Azuelos ^{107,ae}, D. Babal ^{28a}, H. Bachacou ¹³⁴, K. Bachas ^{151,s}, A. Bachiu ³⁴, F. Backman ^{47a,47b}, A. Badea ⁶¹, P. Bagnaia ^{74a,74b}, M. Bahmani ¹⁸, A.J. Bailey ¹⁶², V.R. Bailey ¹⁶¹, J.T. Baines ¹³³, C. Bakalis ¹⁰, O.K. Baker ¹⁷¹, P.J. Bakker ¹¹³, E. Bakos ¹⁵, D. Bakshi Gupta ⁸, S. Balaji ¹⁴⁶, R. Balasubramanian ¹¹³, E.M. Baldin ³⁷, P. Balek ¹³², E. Ballabene ^{70a,70b}, F. Balli ¹³⁴, L.M. Baltes ^{63a}, W.K. Balunas ³², J. Balz ⁹⁹, E. Banas ⁸⁵, M. Bandieramonte ¹²⁸, A. Bandyopadhyay ²⁴, S. Bansal ²⁴, L. Barak ¹⁵⁰, E.L. Barberio ¹⁰⁴, D. Barberis ^{57b,57a}, M. Barbero ¹⁰¹, G. Barbour ⁹⁵, K.N. Barends ^{33a}, T. Barillari ¹⁰⁹, M-S. Barisits ³⁶, T. Barklow ¹⁴², R.M. Barnett ^{17a}, P. Baron ¹²¹, D.A. Baron Moreno ¹⁰⁰, A. Baroncelli ^{62a}, G. Barone ²⁹, A.J. Barr ¹²⁵, L. Barranco Navarro ^{47a,47b}, F. Barreiro ⁹⁸, J. Barreiro Guimarães da Costa ^{14a}, U. Barron ¹⁵⁰, M.G. Barros Teixeira ^{129a}, S. Barsov ³⁷, F. Bartels ^{63a}, R. Bartoldus ¹⁴², A.E. Barton ⁹⁰, P. Bartos ^{28a}, A. Basalae ⁴⁸, A. Basan ⁹⁹, M. Baselga ⁴⁹, I. Bashta ^{76a,76b}, A. Bassalat ^{66,b}, M.J. Basso ¹⁵⁴, C.R. Basson ¹⁰⁰, R.L. Bates ⁵⁹, S. Batlamous ^{35e}, J.R. Batley ³², B. Batool ¹⁴⁰, M. Battaglia ¹³⁵, D. Battulga ¹⁸, M. Baucé ^{74a,74b}, P. Bauer ²⁴, A. Bayirli ^{21a}, J.B. Beacham ⁵¹, T. Beau ¹²⁶, P.H. Beauchemin ¹⁵⁷, F. Becherer ⁵⁴, P. Bechtel ²⁴, H.P. Beck ^{19,r}, K. Becker ¹⁶⁶, A.J. Beddall ^{21d}, V.A. Bednyakov ³⁸, C.P. Bee ¹⁴⁴, L.J. Beemster ¹⁵, T.A. Beermann ³⁶, M. Begalli ^{81d,81d}, M. Begel ²⁹, A. Behera ¹⁴⁴, J.K. Behr ⁴⁸, C. Beirao Da Cruz E Silva ³⁶, J.F. Beirer ^{55,36}, F. Beisiegel ²⁴, M. Belfkir ¹⁵⁸, G. Bella ¹⁵⁰, L. Bellagamba ^{23b}, A. Bellerive ³⁴, P. Bellos ²⁰, K. Beloborodov ³⁷, K. Belotskiy ³⁷, N.L. Belyaev ³⁷, D. Benckekroun ^{35a}, F. Bendebba ^{35a}, Y. Benhammou ¹⁵⁰, D.P. Benjamin ²⁹,

M. Benoit ²⁹, J.R. Bensinger ²⁶, S. Bentvelsen ¹¹³, L. Beresford ³⁶, M. Beretta ⁵³, D. Berge ¹⁸,
E. Bergeaas Kuutmann ¹⁶⁰, N. Berger ⁴, B. Bergmann ¹³¹, J. Beringer ^{17a}, S. Berlendis ⁷,
G. Bernardi ⁵, C. Bernius ¹⁴², F.U. Bernlochner ²⁴, T. Berry ⁹⁴, P. Berta ¹³², A. Berthold ⁵⁰,
I.A. Bertram ⁹⁰, S. Bethke ¹⁰⁹, A. Betti ^{74a,74b}, A.J. Bevan ⁹³, M. Bhamjee ^{33c}, S. Bhatta ¹⁴⁴,
D.S. Bhattacharya ¹⁶⁵, P. Bhattarai ²⁶, V.S. Bhopatkar ¹²⁰, R. Bi ^{29,ah}, R.M. Bianchi ¹²⁸,
O. Biebel ¹⁰⁸, R. Bielski ¹²², M. Biglietti ^{76a}, T.R.V. Billoud ¹³¹, M. Bindi ⁵⁵, A. Bingul ^{21b},
C. Bini ^{74a,74b}, S. Biondi ^{23b,23a}, A. Biondini ⁹¹, C.J. Birch-sykes ¹⁰⁰, G.A. Bird ^{20,133},
M. Birman ¹⁶⁸, T. Bisanz ³⁶, E. Bisceglie ^{43b,43a}, D. Biswas ^{169,m}, A. Bitadze ¹⁰⁰, K. Bjørke ¹²⁴,
I. Bloch ⁴⁸, C. Blocker ²⁶, A. Blue ⁵⁹, U. Blumenschein ⁹³, J. Blumenthal ⁹⁹, G.J. Bobbink ¹¹³,
V.S. Bobrovnikov ³⁷, M. Boehler ⁵⁴, D. Bogavac ³⁶, A.G. Bogdanchikov ³⁷, C. Bohm ^{47a},
V. Boisvert ⁹⁴, P. Bokan ⁴⁸, T. Bold ^{84a}, M. Bomben ⁵, M. Bona ⁹³, M. Boonekamp ¹³⁴,
C.D. Booth ⁹⁴, A.G. Borbély ⁵⁹, H.M. Borecka-Bielska ¹⁰⁷, L.S. Borgna ⁹⁵, G. Borissov ⁹⁰,
D. Bortoletto ¹²⁵, D. Boscherini ^{23b}, M. Bosman ¹³, J.D. Bossio Sola ³⁶, K. Bouaouda ^{35a},
N. Bouchhar ¹⁶², J. Boudreau ¹²⁸, E.V. Bouhova-Thacker ⁹⁰, D. Boumediene ⁴⁰, R. Bouquet ⁵,
A. Boveia ¹¹⁸, J. Boyd ³⁶, D. Boye ²⁹, I.R. Boyko ³⁸, J. Bracinik ²⁰, N. Brahimi ^{62d},
G. Brandt ¹⁷⁰, O. Brandt ³², F. Braren ⁴⁸, B. Brau ¹⁰², J.E. Brau ¹²², K. Brendlinger ⁴⁸,
R. Brenner ¹⁶⁸, L. Brenner ¹¹³, R. Brenner ¹⁶⁰, S. Bressler ¹⁶⁸, B. Brickwedde ⁹⁹, D. Britton ⁵⁹,
D. Britzger ¹⁰⁹, I. Brock ²⁴, G. Brooijmans ⁴¹, W.K. Brooks ^{136f}, E. Brost ²⁹, T.L. Bruckler ¹²⁵,
P.A. Bruckman de Renstrom ⁸⁵, B. Brüers ⁴⁸, D. Bruncko ^{28b,*}, A. Bruni ^{23b}, G. Bruni ^{23b},
M. Bruschi ^{23b}, N. Bruscinò ^{74a,74b}, L. Bryngemark ¹⁴², T. Buanes ¹⁶, Q. Buat ¹³⁷,
P. Buchholz ¹⁴⁰, A.G. Buckley ⁵⁹, I.A. Budagov ^{38,*}, M.K. Bugge ¹²⁴, O. Bulekov ³⁷,
B.A. Bullard ⁶¹, S. Burdin ⁹¹, C.D. Burgard ⁴⁸, A.M. Burger ⁴⁰, B. Burghgrave ⁸, J.T.P. Burr ³²,
C.D. Burton ¹¹, J.C. Burzynski ¹⁴¹, E.L. Busch ⁴¹, V. Büscher ⁹⁹, P.J. Bussey ⁵⁹, J.M. Butler ²⁵,
C.M. Buttar ⁵⁹, J.M. Butterworth ⁹⁵, W. Buttinger ¹³³, C.J. Buxo Vazquez ¹⁰⁶, A.R. Buzykaev ³⁷,
G. Cabras ^{23b}, S. Cabrera Urbán ¹⁶², D. Caforio ⁵⁸, H. Cai ¹²⁸, Y. Cai ^{14a,14d}, V.M.M. Cairo ³⁶,
O. Cakir ^{3a}, N. Calace ³⁶, P. Calafiura ^{17a}, G. Calderini ¹²⁶, P. Calfayan ⁶⁷, G. Callea ⁵⁹,
L.P. Caloba ^{81b}, D. Calvet ⁴⁰, S. Calvet ⁴⁰, T.P. Calvet ¹⁰¹, M. Calvetti ^{73a,73b},
R. Camacho Toro ¹²⁶, S. Camarda ³⁶, D. Camarero Munoz ²⁶, P. Camarri ^{75a,75b},
M.T. Camerlingo ^{76a,76b}, D. Cameron ¹²⁴, C. Camincher ¹⁶⁴, M. Campanelli ⁹⁵, A. Camplani ⁴²,
V. Canale ^{71a,71b}, A. Canesse ¹⁰³, M. Cano Bret ⁷⁹, J. Cantero ¹⁶², Y. Cao ¹⁶¹, F. Capocasa ²⁶,
M. Capua ^{43b,43a}, A. Carbone ^{70a,70b}, R. Cardarelli ^{75a}, J.C.J. Cardenas ⁸, F. Cardillo ¹⁶²,
T. Carli ³⁶, G. Carlino ^{71a}, J.I. Carlotto ¹³, B.T. Carlson ^{128,t}, E.M. Carlson ^{164,155a},
L. Carminati ^{70a,70b}, M. Carnesale ^{74a,74b}, S. Caron ¹¹², E. Carquin ^{136f}, S. Carrá ^{70a,70b},
G. Carratta ^{23b,23a}, F. Carri Argos ^{33g}, J.W.S. Carter ¹⁵⁴, T.M. Carter ⁵², M.P. Casado ^{13j},
A.F. Casha ¹⁵⁴, E.G. Castiglia ¹⁷¹, F.L. Castillo ^{63a}, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶²,
N.F. Castro ^{129a,129e}, A. Catinaccio ³⁶, J.R. Catmore ¹²⁴, V. Cavaliere ²⁹, N. Cavalli ^{23b,23a},
V. Cavasinni ^{73a,73b}, E. Celebi ^{21a}, F. Celli ¹²⁵, M.S. Centonze ^{69a,69b}, K. Cerny ¹²¹,
A.S. Cerqueira ^{81a}, A. Cerri ¹⁴⁵, L. Cerrito ^{75a,75b}, F. Cerutti ^{17a}, A. Cervelli ^{23b}, S.A. Cetin ^{21d},
Z. Chadi ^{35a}, D. Chakraborty ¹¹⁴, M. Chala ^{129f}, J. Chan ¹⁶⁹, W.Y. Chan ¹⁵², J.D. Chapman ³²,
B. Chargeishvili ^{148b}, D.G. Charlton ²⁰, T.P. Charman ⁹³, M. Chatterjee ¹⁹, S. Chekanov ⁶,
S.V. Chekulaev ^{155a}, G.A. Chelkov ^{38,a}, A. Chen ¹⁰⁵, B. Chen ¹⁵⁰, B. Chen ¹⁶⁴, H. Chen ^{14c},
H. Chen ²⁹, J. Chen ^{62c}, J. Chen ²⁶, S. Chen ¹⁵², S.J. Chen ^{14c}, X. Chen ^{62c}, X. Chen ^{14b,ad},
Y. Chen ^{62a}, C.L. Cheng ¹⁶⁹, H.C. Cheng ^{64a}, S. Cheong ¹⁴², A. Cheplakov ³⁸,
E. Cheremushkina ⁴⁸, E. Cherepanova ¹¹³, R. Cherkaoui El Moursli ^{35e}, E. Cheu ⁷, K. Cheung ⁶⁵,
L. Chevalier ¹³⁴, V. Chiarella ⁵³, G. Chiarelli ^{73a}, N. Chiedde ¹⁰¹, G. Chiodini ^{69a},
A.S. Chisholm ²⁰, A. Chitan ^{27b}, M. Chitishvili ¹⁶², Y.H. Chiu ¹⁶⁴, M.V. Chizhov ³⁸, K. Choi ¹¹,
A.R. Chomont ^{74a,74b}, Y. Chou ¹⁰², E.Y.S. Chow ¹¹³, T. Chowdhury ^{33g}, L.D. Christopher ^{33g},

K.L. Chu^{64a}, M.C. Chu^{64a}, X. Chu^{14a,14d}, J. Chudoba¹³⁰, J.J. Chwastowski⁸⁵, D. Cieri¹⁰⁹,
 K.M. Ciesla^{84a}, V. Cindro⁹², A. Ciocio^{17a}, F. Citroto^{71a,71b}, Z.H. Citron^{168,n}, M. Citterio^{70a},
 D.A. Ciubotaru^{27b}, B.M. Ciungu¹⁵⁴, A. Clark⁵⁶, P.J. Clark⁵², J.M. Clavijo Columbie⁴⁸,
 S.E. Clawson¹⁰⁰, C. Clement^{47a,47b}, J. Clercx⁴⁸, L. Clissa^{23b,23a}, Y. Coadou¹⁰¹,
 M. Cobal^{68a,68c}, A. Coccaro^{57b}, R.F. Coelho Barrue^{129a}, R. Coelho Lopes De Sa¹⁰²,
 S. Coelli^{70a}, H. Cohen¹⁵⁰, A.E.C. Coimbra^{70a,70b}, B. Cole⁴¹, J. Collot⁶⁰,
 P. Conde Muiño^{129a,129g}, M.P. Connell^{33c}, S.H. Connell^{33c}, I.A. Connelly⁵⁹, E.I. Conroy¹²⁵,
 F. Conventi^{71a,af}, H.G. Cooke²⁰, A.M. Cooper-Sarkar¹²⁵, F. Cormier¹⁶³, L.D. Corpe³⁶,
 M. Corradi^{74a,74b}, E.E. Corrigan⁹⁷, F. Corriveau^{103,x}, A. Cortes-Gonzalez¹⁸, M.J. Costa¹⁶²,
 F. Costanza⁴, D. Costanzo¹³⁸, B.M. Cote¹¹⁸, G. Cowan⁹⁴, J.W. Cowley³², K. Cranmer¹¹⁶,
 S. Crépe-Renaudin⁶⁰, F. Crescioli¹²⁶, M. Cristinziani¹⁴⁰, M. Cristoforetti^{77a,77b,d}, V. Croft¹⁵⁷,
 G. Crosetti^{43b,43a}, A. Cueto³⁶, T. Cuhadar Donszelmann¹⁵⁹, H. Cui^{14a,14d}, Z. Cui⁷,
 A.R. Cukierman¹⁴², W.R. Cunningham⁵⁹, F. Curcio^{43b,43a}, P. Czodrowski³⁶, M.M. Czurylo^{63b},
 M.J. Da Cunha Sargedas De Sousa^{62a}, J.V. Da Fonseca Pinto^{81b}, C. Da Via¹⁰⁰, W. Dabrowski^{84a},
 T. Dado⁴⁹, S. Dahbi^{33g}, T. Dai¹⁰⁵, C. Dallapiccola¹⁰², M. Dam⁴², G. D'amen²⁹,
 V. D'Amico¹⁰⁸, J. Damp⁹⁹, J.R. Dandoy¹²⁷, M.F. Daneri³⁰, M. Danninger¹⁴¹, V. Dao³⁶,
 G. Darbo^{57b}, S. Darmora⁶, S.J. Das²⁹, S. D'Auria^{70a,70b}, A. D'Avanzo^{71b}, C. David^{155b},
 T. Davidek¹³², D.R. Davis⁵¹, B. Davis-Purcell³⁴, I. Dawson⁹³, K. De⁸, R. De Asmundis^{71a},
 M. De Beurs¹¹³, N. De Biase⁴⁸, S. De Castro^{23b,23a}, N. De Groot¹¹², P. de Jong¹¹³,
 H. De la Torre¹⁰⁶, A. De Maria^{14c}, A. De Salvo^{74a}, U. De Sanctis^{75a,75b}, A. De Santo¹⁴⁵,
 J.B. De Vivie De Regie⁶⁰, D.V. Dedovich³⁸, J. Degens¹¹³, A.M. Deiana⁴⁴, F. Del Corso^{23b,23a},
 J. Del Peso⁹⁸, F. Del Rio^{63a}, F. Deliot¹³⁴, C.M. Delitzsch⁴⁹, M. Della Pietra^{71a,71b},
 D. Della Volpe⁵⁶, A. Dell'Acqua³⁶, L. Dell'Asta^{70a,70b}, M. Delmastro⁴, P.A. Delsart⁶⁰,
 S. Demers¹⁷¹, M. Demichev³⁸, S.P. Denisov³⁷, L. D'Eramo¹¹⁴, D. Derendarz⁸⁵,
 F. Derue¹²⁶, P. Dervan⁹¹, K. Desch²⁴, K. Dette¹⁵⁴, C. Deutsch²⁴, P.O. Deviveiros³⁶,
 F.A. Di Bello^{57b,57a}, A. Di Ciaccio^{75a,75b}, L. Di Ciaccio⁴, A. Di Domenico^{74a,74b},
 C. Di Donato^{71a,71b}, A. Di Girolamo³⁶, G. Di Gregorio⁵, A. Di Luca^{77a,77b}, B. Di Micco^{76a,76b},
 R. Di Nardo^{76a,76b}, C. Diaconu¹⁰¹, F.A. Dias¹¹³, T. Dias Do Vale¹⁴¹, M.A. Diaz^{136a,136b},
 F.G. Diaz Capriles²⁴, M. Didenko¹⁶², E.B. Diehl¹⁰⁵, L. Diehl⁵⁴, S. Díez Cornell⁴⁸,
 C. Diez Pardos¹⁴⁰, C. Dimitriadi^{24,160}, A. Dimitrievska^{17a}, W. Ding^{14b}, J. Dingfelder²⁴,
 I-M. Dinu^{27b}, S.J. Dittmeier^{63b}, F. Dittus³⁶, F. Djama¹⁰¹, T. Djobava^{148b}, J.I. Djuvsland¹⁶,
 C. Doglioni^{100,97}, J. Dolejsi¹³², Z. Dolezal¹³², M. Donadelli^{81c}, B. Dong^{62c}, J. Donini⁴⁰,
 A. D'Onofrio^{14c}, M. D'Onofrio⁹¹, J. Dopke¹³³, A. Doria^{71a}, M.T. Dova⁸⁹, A.T. Doyle⁵⁹,
 M.A. Draguet¹²⁵, E. Drechsler¹⁴¹, E. Dreyer¹⁶⁸, I. Drivas-koulouris¹⁰, A.S. Drobac¹⁵⁷,
 M. Drozdova⁵⁶, D. Du^{62a}, T.A. du Pree¹¹³, F. Dubinin³⁷, M. Dubovsky^{28a}, E. Duchovni¹⁶⁸,
 G. Duckeck¹⁰⁸, O.A. Ducu^{27b}, D. Duda¹⁰⁹, A. Dudarev³⁶, M. D'uffizi¹⁰⁰, L. Duflot⁶⁶,
 M. Dührssen³⁶, C. Dülsen¹⁷⁰, A.E. Dumitriu^{27b}, M. Dunford^{63a}, S. Dungs⁴⁹,
 K. Dunne^{47a,47b}, A. Duperrin¹⁰¹, H. Duran Yildiz^{3a}, M. Düren⁵⁸, A. Durglishvili^{148b},
 B.L. Dwyer¹¹⁴, G.I. Dyckes^{17a}, M. Dyndal^{84a}, S. Dysch¹⁰⁰, B.S. Dziedzic⁸⁵,
 Z.O. Earnshaw¹⁴⁵, B. Eckerova^{28a}, M.G. Eggleston⁵¹, E. Egidio Purcino De Souza^{81b},
 L.F. Ehrke⁵⁶, G. Eigen¹⁶, K. Einsweiler^{17a}, T. Ekelof¹⁶⁰, P.A. Ekman⁹⁷, Y. El Ghazali^{35b},
 H. El Jarrari^{35e,147}, A. El Moussaouy^{35a}, V. Ellajosyula¹⁶⁰, M. Ellert¹⁶⁰, F. Ellinghaus¹⁷⁰,
 A.A. Elliot⁹³, N. Ellis³⁶, J. Elmsheuser²⁹, M. Elsing³⁶, D. Emelianov¹³³, A. Emerman⁴¹,
 Y. Enari¹⁵², I. Ene^{17a}, S. Epari¹³, J. Erdmann^{49,ac}, A. Ereditato¹⁹, P.A. Erland⁸⁵,
 M. Errenst¹⁷⁰, M. Escalier⁶⁶, C. Escobar¹⁶², E. Etzion¹⁵⁰, G. Evans^{129a}, H. Evans⁶⁷,
 M.O. Evans¹⁴⁵, A. Ezhilov³⁷, S. Ezzarqtouni^{35a}, F. Fabbri⁵⁹, L. Fabbri^{23b,23a}, G. Facini⁹⁵,
 V. Fadeyev¹³⁵, R.M. Fakhrutdinov³⁷, S. Falciano^{74a}, P.J. Falke²⁴, S. Falke³⁶, J. Faltova¹³²,

Y. Fan [id](#)^{14a}, Y. Fang [id](#)^{14a,14d}, G. Fanourakis [id](#)⁴⁶, M. Fanti [id](#)^{70a,70b}, M. Faraj [id](#)^{68a,68b}, Z. Farazpay⁹⁶,
 A. Farbin [id](#)⁸, A. Farilla [id](#)^{76a}, T. Faroouque [id](#)¹⁰⁶, S.M. Farrington [id](#)⁵², F. Fassi [id](#)^{35e}, D. Fassouliotis [id](#)⁹,
 M. Faucci Giannelli [id](#)^{75a,75b}, W.J. Fawcett [id](#)³², L. Fayard [id](#)⁶⁶, P. Federicova [id](#)¹³⁰, O.L. Fedin [id](#)^{37a},
 G. Fedotov [id](#)³⁷, M. Feickert [id](#)¹⁶⁹, L. Feligioni [id](#)¹⁰¹, A. Fell [id](#)¹³⁸, D.E. Fellers [id](#)¹²², C. Feng [id](#)^{62b},
 M. Feng [id](#)^{14b}, Z. Feng [id](#)¹¹³, M.J. Fenton [id](#)¹⁵⁹, A.B. Fenyuk³⁷, L. Ferencz [id](#)⁴⁸, S.W. Ferguson [id](#)⁴⁵,
 J. Ferrando [id](#)⁴⁸, A. Ferrari [id](#)¹⁶⁰, P. Ferrari [id](#)^{113,112}, R. Ferrari [id](#)^{72a}, D. Ferrere [id](#)⁵⁶, C. Ferretti [id](#)¹⁰⁵,
 F. Fiedler [id](#)⁹⁹, A. Filipčič [id](#)⁹², E.K. Filmer [id](#)¹, F. Filthaut [id](#)¹¹², M.C.N. Fiolhais [id](#)^{129a,129c,c},
 L. Fiorini [id](#)¹⁶², F. Fischer [id](#)¹⁴⁰, W.C. Fisher [id](#)¹⁰⁶, T. Fitschen [id](#)¹⁰⁰, I. Fleck [id](#)¹⁴⁰, P. Fleischmann [id](#)¹⁰⁵,
 T. Flick [id](#)¹⁷⁰, L. Flores [id](#)¹²⁷, M. Flores [id](#)^{33d}, L.R. Flores Castillo [id](#)^{64a}, F.M. Follega [id](#)^{77a,77b},
 N. Fomin [id](#)¹⁶, J.H. Foo [id](#)¹⁵⁴, B.C. Forland⁶⁷, A. Formica [id](#)¹³⁴, A.C. Forti [id](#)¹⁰⁰, E. Fortin [id](#)¹⁰¹,
 A.W. Fortman [id](#)⁶¹, M.G. Foti [id](#)^{17a}, L. Fountas [id](#)⁹, D. Fournier [id](#)⁶⁶, H. Fox [id](#)⁹⁰, P. Francavilla [id](#)^{73a,73b},
 S. Francescato [id](#)⁶¹, S. Franchellucci [id](#)⁵⁶, M. Franchini [id](#)^{23b,23a}, S. Franchino [id](#)^{63a}, D. Francis³⁶,
 L. Franco [id](#)¹¹², L. Franconi [id](#)¹⁹, M. Franklin [id](#)⁶¹, G. Frattari [id](#)²⁶, A.C. Freegard [id](#)⁹³, P.M. Freeman²⁰,
 W.S. Freund [id](#)^{81b}, N. Fritzsche [id](#)⁵⁰, A. Froch [id](#)⁵⁴, D. Froidevaux [id](#)³⁶, J.A. Frost [id](#)¹²⁵, Y. Fu [id](#)^{62a},
 M. Fujimoto [id](#)¹¹⁷, E. Fullana Torregrosa [id](#)^{162,*}, J. Fuster [id](#)¹⁶², A. Gabrielli [id](#)^{23b,23a}, A. Gabrielli [id](#)¹⁵⁴,
 P. Gadow [id](#)⁴⁸, G. Gagliardi [id](#)^{57b,57a}, L.G. Gagnon [id](#)^{17a}, G.E. Gallardo [id](#)¹²⁵, E.J. Gallas [id](#)¹²⁵,
 B.J. Gallop [id](#)¹³³, R. Gamboa Goni [id](#)⁹³, K.K. Gan [id](#)¹¹⁸, S. Ganguly [id](#)¹⁵², J. Gao [id](#)^{62a}, Y. Gao [id](#)⁵²,
 F.M. Garay Walls [id](#)^{136a,136b}, B. Garcia^{29,ah}, C. García [id](#)¹⁶², J.E. García Navarro [id](#)¹⁶²,
 J.A. García Pascual [id](#)^{14a}, M. Garcia-Sciveres [id](#)^{17a}, R.W. Gardner [id](#)³⁹, D. Garg [id](#)⁷⁹, R.B. Garg [id](#)¹⁴²,
 S. Gargiulo [id](#)⁵⁴, C.A. Garner¹⁵⁴, V. Garonne [id](#)²⁹, S.J. Gasiorowski [id](#)¹³⁷, P. Gaspar [id](#)^{81b}, G. Gaudio [id](#)^{72a},
 V. Gautam¹³, P. Gauzzi [id](#)^{74a,74b}, I.L. Gavrilenko [id](#)³⁷, A. Gavrilyuk [id](#)³⁷, C. Gay [id](#)¹⁶³, G. Gaycken [id](#)⁴⁸,
 E.N. Gazis [id](#)¹⁰, A.A. Geanta [id](#)^{27b,27e}, C.M. Gee [id](#)¹³⁵, J. Geisen [id](#)⁹⁷, M. Geisen [id](#)⁹⁹, C. Gemme [id](#)^{57b},
 M.H. Genest [id](#)⁶⁰, S. Gentile [id](#)^{74a,74b}, S. George [id](#)⁹⁴, W.F. George [id](#)²⁰, T. Geralis [id](#)⁴⁶, L.O. Gerlach⁵⁵,
 P. Gessinger-Befurt [id](#)³⁶, M. Ghasemi Bostanabad [id](#)¹⁶⁴, M. Ghneimat [id](#)¹⁴⁰, K. Ghorbanian [id](#)⁹³,
 A. Ghosal [id](#)¹⁴⁰, A. Ghosh [id](#)¹⁵⁹, A. Ghosh [id](#)⁷, B. Giacobbe [id](#)^{23b}, S. Giagu [id](#)^{74a,74b},
 N. Giangiacomi [id](#)¹⁵⁴, P. Giannetti [id](#)^{73a}, A. Giannini [id](#)^{62a}, S.M. Gibson [id](#)⁹⁴, M. Gignac [id](#)¹³⁵,
 D.T. Gil [id](#)^{84b}, A.K. Gilbert [id](#)^{84a}, B.J. Gilbert [id](#)⁴¹, D. Gillberg [id](#)³⁴, G. Gilles [id](#)¹¹³, N.E.K. Gillwald [id](#)⁴⁸,
 L. Ginabat [id](#)¹²⁶, D.M. Gingrich [id](#)^{2,ae}, M.P. Giordani [id](#)^{68a,68c}, P.F. Giraud [id](#)¹³⁴, G. Giugliarelli [id](#)^{68a,68c},
 D. Giugni [id](#)^{70a}, F. Giuli [id](#)³⁶, I. Gkialas [id](#)^{9,k}, L.K. Gladilin [id](#)³⁷, C. Glasman [id](#)⁹⁸, G.R. Gledhill [id](#)¹²²,
 M. Glisic¹²², I. Gnesi [id](#)^{43b,g}, Y. Go [id](#)^{29,ah}, M. Goblirsch-Kolb [id](#)²⁶, B. Gocke [id](#)⁴⁹, D. Godin¹⁰⁷,
 S. Goldfarb [id](#)¹⁰⁴, T. Golling [id](#)⁵⁶, M.G.D. Gololo^{33g}, D. Golubkov [id](#)³⁷, J.P. Gombas [id](#)¹⁰⁶,
 A. Gomes [id](#)^{129a,129b}, G. Gomes Da Silva [id](#)¹⁴⁰, A.J. Gomez Delegido [id](#)¹⁶², R. Goncalves Gama [id](#)⁵⁵,
 R. Gonçalves [id](#)^{129a,129c}, G. Gonella [id](#)¹²², L. Gonella [id](#)²⁰, A. Gongadze [id](#)³⁸, F. Gonnella [id](#)²⁰,
 J.L. Gonski [id](#)⁴¹, R.Y. González Andana [id](#)⁵², S. González de la Hoz [id](#)¹⁶², S. Gonzalez Fernandez [id](#)¹³,
 R. Gonzalez Lopez [id](#)⁹¹, C. Gonzalez Renteria [id](#)^{17a}, R. Gonzalez Suarez [id](#)¹⁶⁰, S. Gonzalez-Sevilla [id](#)⁵⁶,
 G.R. Gonzalvo Rodriguez [id](#)¹⁶², L. Goossens [id](#)³⁶, N.A. Gorasia [id](#)²⁰, P.A. Gorbounov [id](#)³⁷, B. Gorini [id](#)³⁶,
 E. Gorini [id](#)^{69a,69b}, A. Gorišek [id](#)⁹², A.T. Goshaw [id](#)⁵¹, M.I. Gostkin [id](#)³⁸, C.A. Gottardo [id](#)³⁶,
 M. Goughri [id](#)^{35b}, V. Goumarre [id](#)⁴⁸, A.G. Goussiou [id](#)¹³⁷, N. Govender [id](#)^{33c}, C. Goy [id](#)⁴,
 I. Grabowska-Bold [id](#)^{84a}, K. Graham [id](#)³⁴, E. Gramstad [id](#)¹²⁴, S. Grancagnolo [id](#)¹⁸, M. Grandi [id](#)¹⁴⁵,
 V. Gratchev^{37,*}, P.M. Gravila [id](#)^{27f}, F.G. Gravili [id](#)^{69a,69b}, H.M. Gray [id](#)^{17a}, M. Greco [id](#)^{69a,69b},
 C. Grefe [id](#)²⁴, I.M. Gregor [id](#)⁴⁸, P. Grenier [id](#)¹⁴², C. Grieco [id](#)¹³, A.A. Grillo [id](#)¹³⁵, K. Grimm [id](#)^{31,o},
 S. Grinstein [id](#)^{13,v}, J.-F. Grivaz [id](#)⁶⁶, E. Gross [id](#)¹⁶⁸, J. Grosse-Knetter [id](#)⁵⁵, C. Grud¹⁰⁵, A. Grummer [id](#)¹¹¹,
 J.C. Grundy [id](#)¹²⁵, L. Guan [id](#)¹⁰⁵, W. Guan [id](#)¹⁶⁹, C. Gubbels [id](#)¹⁶³, J.G.R. Guerrero Rojas [id](#)¹⁶²,
 G. Guerrieri [id](#)^{68a,68b}, F. Guescini [id](#)¹⁰⁹, R. Gugel [id](#)⁹⁹, J.A.M. Guhit [id](#)¹⁰⁵, A. Guida [id](#)⁴⁸, T. Guillemain [id](#)⁴,
 E. Guilloton [id](#)^{166,133}, S. Guindon [id](#)³⁶, F. Guo [id](#)^{14a,14d}, J. Guo [id](#)^{62c}, L. Guo [id](#)⁶⁶, Y. Guo [id](#)¹⁰⁵,
 R. Gupta [id](#)⁴⁸, S. Gurbuz [id](#)²⁴, S.S. Gurdasani [id](#)⁵⁴, G. Gustavino [id](#)³⁶, M. Guth [id](#)⁵⁶, P. Gutierrez [id](#)¹¹⁹,
 L.F. Gutierrez Zagazeta [id](#)¹²⁷, C. Gutschow [id](#)⁹⁵, C. Guyot [id](#)¹³⁴, C. Gwenlan [id](#)¹²⁵, C.B. Gwilliam [id](#)⁹¹,

E.S. Haaland ¹²⁴, A. Haas ¹¹⁶, M. Habedank ⁴⁸, C. Haber ^{17a}, H.K. Hadavand ⁸, A. Hadeif ⁹⁹,
 S. Hadzic ¹⁰⁹, E.H. Haines ⁹⁵, M. Haleem ¹⁶⁵, J. Haley ¹²⁰, J.J. Hall ¹³⁸, G.D. Hallelwell ¹⁰¹,
 L. Halser ¹⁹, K. Hamano ¹⁶⁴, H. Hamdaoui ^{35e}, M. Hamer ²⁴, G.N. Hamity ⁵², J. Han ^{62b},
 K. Han ^{62a}, L. Han ^{14c}, L. Han ^{62a}, S. Han ^{17a}, Y.F. Han ¹⁵⁴, K. Hanagaki ⁸², M. Hance ¹³⁵,
 D.A. Hangal ^{41,ab}, H. Hanif ¹⁴¹, M.D. Hank ³⁹, R. Hankache ¹⁰⁰, J.B. Hansen ⁴²,
 J.D. Hansen ⁴², P.H. Hansen ⁴², K. Hara ¹⁵⁶, D. Harada ⁵⁶, T. Harenberg ¹⁷⁰, S. Harkusha ³⁷,
 Y.T. Harris ¹²⁵, N.M. Harrison ¹¹⁸, P.F. Harrison ¹⁶⁶, N.M. Hartman ¹⁴², N.M. Hartmann ¹⁰⁸,
 Y. Hasegawa ¹³⁹, A. Hasib ⁵², S. Haug ¹⁹, R. Hauser ¹⁰⁶, M. Havranek ¹³¹, C.M. Hawkes ²⁰,
 R.J. Hawkings ³⁶, S. Hayashida ¹¹⁰, D. Hayden ¹⁰⁶, C. Hayes ¹⁰⁵, R.L. Hayes ¹⁶³, C.P. Hays ¹²⁵,
 J.M. Hays ⁹³, H.S. Hayward ⁹¹, F. He ^{62a}, Y. He ¹⁵³, Y. He ¹²⁶, M.P. Heath ⁵², V. Hedberg ⁹⁷,
 A.L. Heggelund ¹²⁴, N.D. Hehir ⁹³, C. Heidegger ⁵⁴, K.K. Heidegger ⁵⁴, W.D. Heidorn ⁸⁰,
 J. Heilmann ³⁴, S. Heim ⁴⁸, T. Heim ^{17a}, J.G. Heinlein ¹²⁷, J.J. Heinrich ¹²², L. Heinrich ¹⁰⁹,
 J. Hejbal ¹³⁰, L. Helary ⁴⁸, A. Held ¹⁶⁹, S. Hellesund ¹²⁴, C.M. Helling ¹⁶³, S. Hellman ^{47a,47b},
 C. Helsens ³⁶, R.C.W. Henderson ⁹⁰, L. Henkelmann ³², A.M. Henriques Correia ³⁶, H. Herde ⁹⁷,
 Y. Hernández Jiménez ¹⁴⁴, L.M. Herrmann ²⁴, M.G. Herrmann ¹⁰⁸, T. Herrmann ⁵⁰, G. Herten ⁵⁴,
 R. Hertenberger ¹⁰⁸, L. Hervás ³⁶, N.P. Hesse ^{155a}, H. Hibi ⁸³, E. Higón-Rodríguez ¹⁶²,
 S.J. Hillier ²⁰, I. Hinchliffe ^{17a}, F. Hinterkeuser ²⁴, M. Hirose ¹²³, S. Hirose ¹⁵⁶,
 D. Hirschbuehl ¹⁷⁰, T.G. Hitchings ¹⁰⁰, B. Hiti ⁹², J. Hobbs ¹⁴⁴, R. Hobincu ^{27e}, N. Hod ¹⁶⁸,
 M.C. Hodgkinson ¹³⁸, B.H. Hodgkinson ³², A. Hoecker ³⁶, J. Hofer ⁴⁸, D. Hohn ⁵⁴, T. Holm ²⁴,
 M. Holzbock ¹⁰⁹, L.B.A.H. Hommels ³², B.P. Honan ¹⁰⁰, J. Hong ^{62c}, T.M. Hong ¹²⁸,
 J.C. Honig ⁵⁴, A. Hönle ¹⁰⁹, B.H. Hooberman ¹⁶¹, W.H. Hopkins ⁶, Y. Horii ¹¹⁰, S. Hou ¹⁴⁷,
 A.S. Howard ⁹², J. Howarth ⁵⁹, J. Hoya ⁶, M. Hrabovsky ¹²¹, A. Hrynevich ⁴⁸, T. Hryn'ova ⁴,
 P.J. Hsu ⁶⁵, S.-C. Hsu ¹³⁷, Q. Hu ⁴¹, Y.F. Hu ^{14a,14d,ag}, D.P. Huang ⁹⁵, S. Huang ^{64b},
 X. Huang ^{14c}, Y. Huang ^{62a}, Y. Huang ^{14a}, Z. Huang ¹⁰⁰, Z. Hubacek ¹³¹, M. Huebner ²⁴,
 F. Huegging ²⁴, T.B. Huffman ¹²⁵, M. Huhtinen ³⁶, S.K. Huiberts ¹⁶, R. Hulsken ¹⁰³,
 N. Huseynov ^{12,a}, J. Huston ¹⁰⁶, J. Huth ⁶¹, R. Hyneman ¹⁴², S. Hyrych ^{28a}, G. Iacobucci ⁵⁶,
 G. Iakovidis ²⁹, I. Ibragimov ¹⁴⁰, L. Iconomidou-Fayard ⁶⁶, P. Iengo ^{71a,71b}, R. Iguchi ¹⁵²,
 T. Iizawa ⁵⁶, Y. Ikegami ⁸², A. Ilg ¹⁹, N. Ilic ¹⁵⁴, H. Imam ^{35a}, T. Ingebretsen Carlson ^{47a,47b},
 G. Introzzi ^{72a,72b}, M. Iodice ^{76a}, V. Ippolito ^{74a,74b}, M. Ishino ¹⁵², W. Islam ¹⁶⁹, C. Issever ^{18,48},
 S. Istin ^{21a,aj}, H. Ito ¹⁶⁷, J.M. Iturbe Ponce ^{64a}, R. Iuppa ^{77a,77b}, A. Ivina ¹⁶⁸, J.M. Izen ⁴⁵,
 V. Izzo ^{71a}, P. Jacka ^{130,131}, P. Jackson ¹, R.M. Jacobs ⁴⁸, B.P. Jaeger ¹⁴¹, C.S. Jagfeld ¹⁰⁸,
 G. Jäkel ¹⁷⁰, K. Jakobs ⁵⁴, T. Jakoubek ¹⁶⁸, J. Jamieson ⁵⁹, K.W. Janas ^{84a}, G. Jarlskog ⁹⁷,
 A.E. Jaspan ⁹¹, M. Javurkova ¹⁰², F. Jeanneau ¹³⁴, L. Jeanty ¹²², J. Jejelava ^{148a,aa}, P. Jenni ^{54,h},
 C.E. Jessiman ³⁴, S. Jézéquel ⁴, J. Jia ¹⁴⁴, X. Jia ⁶¹, X. Jia ^{14a,14d}, Z. Jia ^{14c}, Y. Jiang ^{62a},
 S. Jiggins ⁵², J. Jimenez Pena ¹⁰⁹, S. Jin ^{14c}, A. Jinaru ^{27b}, O. Jinnouchi ¹⁵³, P. Johansson ¹³⁸,
 K.A. Johns ⁷, D.M. Jones ³², E. Jones ¹⁶⁶, P. Jones ³², R.W.L. Jones ⁹⁰, T.J. Jones ⁹¹,
 R. Joshi ¹¹⁸, J. Jovicevic ¹⁵, X. Ju ^{17a}, J.J. Junggeburth ³⁶, A. Juste Rozas ^{13,v}, S. Kabana ^{136e},
 A. Kaczmarzka ⁸⁵, M. Kado ^{74a,74b}, H. Kagan ¹¹⁸, M. Kagan ¹⁴², A. Kahn ⁴¹, A. Kahn ¹²⁷,
 C. Kahra ⁹⁹, T. Kaji ¹⁶⁷, E. Kajomovitz ¹⁴⁹, N. Kakati ¹⁶⁸, C.W. Kalderon ²⁹,
 A. Kamenshchikov ¹⁵⁴, S. Kanayama ¹⁵³, N.J. Kang ¹³⁵, Y. Kano ¹¹⁰, D. Kar ^{33g}, K. Karava ¹²⁵,
 M.J. Kareem ^{155b}, E. Karentzos ⁵⁴, I. Karkanas ^{151,f}, S.N. Karpov ³⁸, Z.M. Karpova ³⁸,
 V. Kartvelishvili ⁹⁰, A.N. Karyukhin ³⁷, E. Kasimi ^{151,f}, C. Kato ^{62d}, J. Katzy ⁴⁸, S. Kaur ³⁴,
 K. Kawade ¹³⁹, K. Kawagoe ⁸⁸, T. Kawamoto ¹³⁴, G. Kawamura ⁵⁵, E.F. Kay ¹⁶⁴, F.I. Kaya ¹⁵⁷,
 S. Kazakos ¹³, V.F. Kazanin ³⁷, Y. Ke ¹⁴⁴, J.M. Keaveney ^{33a}, R. Keeler ¹⁶⁴, G.V. Kehris ⁶¹,
 J.S. Keller ³⁴, A.S. Kelly ⁹⁵, D. Kelsey ¹⁴⁵, J.J. Kempster ²⁰, K.E. Kennedy ⁴¹, P.D. Kennedy ⁹⁹,
 O. Kepka ¹³⁰, B.P. Kerridge ¹⁶⁶, S. Kersten ¹⁷⁰, B.P. Kerševan ⁹², S. Keshri ⁶⁶,
 L. Keszeghova ^{28a}, S. Ketabchi Haghghat ¹⁵⁴, M. Khandoga ¹²⁶, A. Khanov ¹²⁰,

A.G. Kharlamov ³⁷, T. Kharlamova ³⁷, E.E. Khoda ¹³⁷, T.J. Khoo ¹⁸, G. Khorauli ¹⁶⁵,
 J. Khubua ^{148b}, Y.A.R. Khwaira ⁶⁶, M. Kiehn ³⁶, A. Kilgallon ¹²², D.W. Kim ^{47a,47b},
 E. Kim ¹⁵³, Y.K. Kim ³⁹, N. Kimura ⁹⁵, A. Kirchhoff ⁵⁵, D. Kirchmeier ⁵⁰, C. Kirfel ²⁴,
 J. Kirk ¹³³, A.E. Kiryunin ¹⁰⁹, T. Kishimoto ¹⁵², D.P. Kisliuk ¹⁵⁴, C. Kitsaki ¹⁰, O. Kivernyk ²⁴,
 M. Klassen ^{63a}, C. Klein ³⁴, L. Klein ¹⁶⁵, M.H. Klein ¹⁰⁵, M. Klein ⁹¹, S.B. Klein ⁵⁶,
 U. Klein ⁹¹, P. Klimek ³⁶, A. Klimentov ²⁹, F. Klimpel ¹⁰⁹, T. Klingl ²⁴, T. Klioutchnikova ³⁶,
 F.F. Klitzner ¹⁰⁸, P. Kluit ¹¹³, S. Kluth ¹⁰⁹, E. Kneringer ⁷⁸, T.M. Knight ¹⁵⁴, A. Knue ⁵⁴,
 D. Kobayashi ⁸⁸, R. Kobayashi ⁸⁶, M. Kocian ¹⁴², P. Kodyš ¹³², D.M. Koeck ¹⁴⁵, P.T. Koenig ²⁴,
 T. Koffas ³⁴, M. Kolb ¹³⁴, I. Koletsou ⁴, T. Komarek ¹²¹, K. Köneke ⁵⁴, A.X.Y. Kong ¹,
 T. Kono ¹¹⁷, N. Konstantinidis ⁹⁵, B. Konya ⁹⁷, R. Kopeliansky ⁶⁷, S. Koperny ^{84a}, K. Korcyl ⁸⁵,
 K. Kordas ^{151,f}, G. Koren ¹⁵⁰, A. Korn ⁹⁵, S. Korn ⁵⁵, I. Korolkov ¹³, N. Korotkova ³⁷,
 B. Kortman ¹¹³, O. Kortner ¹⁰⁹, S. Kortner ¹⁰⁹, W.H. KostECKA ¹¹⁴, V.V. Kostyukhin ¹⁴⁰,
 A. Kotsokchagia ¹³⁴, A. Kotwal ⁵¹, A. Koulouris ³⁶, A. Kourkoumeli-Charalampidi ^{72a,72b},
 C. Kourkoumelis ⁹, E. Kourlitis ⁶, O. Kovanda ¹⁴⁵, R. Kowalewski ¹⁶⁴, W. Kozanecki ¹³⁴,
 A.S. Kozhin ³⁷, V.A. Kramarenko ³⁷, G. Kramberger ⁹², P. Kramer ⁹⁹, M.W. Krasny ¹²⁶,
 A. Krasznahorkay ³⁶, J.A. Kremer ⁹⁹, T. Kresse ⁵⁰, J. Kretzschmar ⁹¹, K. Kreul ¹⁸,
 P. Krieger ¹⁵⁴, F. Krieter ¹⁰⁸, S. Krishnamurthy ¹⁰², A. Krishnan ^{63b}, M. Krivos ¹³²,
 K. Krizka ^{17a}, K. Kroeninger ⁴⁹, H. Kroha ¹⁰⁹, J. Kroll ¹³⁰, J. Kroll ¹²⁷, K.S. Krowpman ¹⁰⁶,
 U. Kruchonak ³⁸, H. Krüger ²⁴, N. Krumnack ⁸⁰, M.C. Kruse ⁵¹, J.A. Krzysiak ⁸⁵,
 O. Kuchinskaia ³⁷, S. Kuday ^{3a}, D. Kuechler ⁴⁸, J.T. Kuechler ⁴⁸, S. Kuehn ³⁶, T. Kuhl ⁴⁸,
 V. Kukhtin ³⁸, Y. Kulchitsky ^{37,a}, S. Kuleshov ^{136d,136b}, M. Kumar ^{33g}, N. Kumari ¹⁰¹,
 A. Kupco ¹³⁰, T. Kupfer ⁴⁹, A. Kupich ³⁷, O. Kuprash ⁵⁴, H. Kurashige ⁸³, L.L. Kurchaninov ^{155a},
 Y.A. Kurochkin ³⁷, A. Kurova ³⁷, M. Kuze ¹⁵³, A.K. Kvam ¹⁰², J. Kvita ¹²¹, T. Kwan ¹⁰³,
 K.W. Kwok ^{64a}, N.G. Kyriacou ¹⁰⁵, L.A.O. Laatu ¹⁰¹, C. Lacasta ¹⁶², F. Lacava ^{74a,74b},
 H. Lacker ¹⁸, D. Lacour ¹²⁶, N.N. Lad ⁹⁵, E. Ladygin ³⁸, B. Laforge ¹²⁶, T. Lagouri ^{136e},
 S. Lai ⁵⁵, I.K. Lakomic ^{84a}, N. Lalloue ⁶⁰, J.E. Lambert ¹¹⁹, S. Lammers ⁶⁷, W. Lampl ⁷,
 C. Lampoudis ^{151,f}, A.N. Lancaster ¹¹⁴, E. Lançon ²⁹, U. Landgraf ⁵⁴, M.P.J. Landon ⁹³,
 V.S. Lang ⁵⁴, R.J. Langenberg ¹⁰², A.J. Lankford ¹⁵⁹, F. Lanni ³⁶, K. Lantzsch ²⁴, A. Lanza ^{72a},
 A. Lapertosa ^{57b,57a}, J.F. Laporte ¹³⁴, T. Lari ^{70a}, F. Lasagni Manghi ^{23b}, M. Lassnig ³⁶,
 V. Latonova ¹³⁰, T.S. Lau ^{64a}, A. Laudrain ⁹⁹, A. Laurier ³⁴, S.D. Lawlor ⁹⁴, Z. Lawrence ¹⁰⁰,
 M. Lazzaroni ^{70a,70b}, B. Le ¹⁰⁰, B. Leban ⁹², A. Lebedev ⁸⁰, M. LeBlanc ³⁶, T. LeCompte ⁶,
 F. Ledroit-Guillon ⁶⁰, A.C.A. Lee ⁹⁵, G.R. Lee ¹⁶, L. Lee ⁶¹, S.C. Lee ¹⁴⁷, S. Lee ^{47a,47b},
 T.F. Lee ⁹¹, L.L. Leeuw ^{33c}, H.P. Lefebvre ⁹⁴, M. Lefebvre ¹⁶⁴, C. Leggett ^{17a}, K. Lehmann ¹⁴¹,
 G. Lehmann Miotto ³⁶, M. Leigh ⁵⁶, W.A. Leight ¹⁰², A. Leisos ^{151,u}, M.A.L. Leite ^{81c},
 C.E. Leitgeb ⁴⁸, R. Leitner ¹³², K.J.C. Leney ⁴⁴, T. Lenz ²⁴, S. Leone ^{73a}, C. Leonidopoulos ⁵²,
 A. Leopold ¹⁴³, C. Leroy ¹⁰⁷, R. Les ¹⁰⁶, C.G. Lester ³², M. Levchenko ³⁷, J. Levêque ⁴,
 D. Levin ¹⁰⁵, L.J. Levinson ¹⁶⁸, M.P. Lewicki ⁸⁵, D.J. Lewis ⁴, A. Li ⁵, B. Li ^{14b}, B. Li ^{62b},
 C. Li ^{62a}, C-Q. Li ^{62c}, H. Li ^{62a}, H. Li ^{62b}, H. Li ^{14c}, H. Li ^{62b}, J. Li ^{62c}, K. Li ¹³⁷, L. Li ^{62c},
 M. Li ^{14a,14d}, Q.Y. Li ^{62a}, S. Li ^{14a,14d}, S. Li ^{62d,62c,e}, T. Li ^{62b}, X. Li ¹⁰³, Z. Li ^{62b}, Z. Li ¹²⁵,
 Z. Li ¹⁰³, Z. Li ⁹¹, Z. Li ^{14a,14d}, Z. Liang ^{14a}, M. Liberatore ⁴⁸, B. Liberti ^{75a}, K. Lie ^{64c},
 J. Lieber Marin ^{81b}, K. Lin ¹⁰⁶, R.A. Linck ⁶⁷, R.E. Lindley ⁷, J.H. Lindon ², A. Linss ⁴⁸,
 E. Lipeles ¹²⁷, A. Lipniacka ¹⁶, A. Lister ¹⁶³, J.D. Little ⁴, B. Liu ^{14a}, B.X. Liu ¹⁴¹,
 D. Liu ^{62d,62c}, J.B. Liu ^{62a}, J.K.K. Liu ³², K. Liu ^{62d,62c}, M. Liu ^{62a}, M.Y. Liu ^{62a}, P. Liu ^{14a},
 Q. Liu ^{62d,137,62c}, X. Liu ^{62a}, Y. Liu ⁴⁸, Y. Liu ^{14c,14d}, Y.L. Liu ¹⁰⁵, Y.W. Liu ^{62a},
 M. Livan ^{72a,72b}, J. Llorente Merino ¹⁴¹, S.L. Lloyd ⁹³, E.M. Lobodzinska ⁴⁸, P. Loch ⁷,
 S. Loffredo ^{75a,75b}, T. Lohse ¹⁸, K. Lohwasser ¹³⁸, M. Lokajicek ¹³⁰, J.D. Long ¹⁶¹,
 I. Longarini ^{74a,74b}, L. Longo ^{69a,69b}, R. Longo ¹⁶¹, I. Lopez Paz ³⁶, A. Lopez Solis ⁴⁸,




























































































































































J. Lorenz ^{id108}, N. Lorenzo Martinez ^{id4}, A.M. Lory ^{id108}, A. Lösle ^{id54}, X. Lou ^{id47a,47b}, X. Lou ^{id14a,14d},
 A. Lounis ^{id66}, J. Love ^{id6}, P.A. Love ^{id90}, J.J. Lozano Bahilo ^{id162}, G. Lu ^{id14a,14d}, M. Lu ^{id79},
 S. Lu ^{id127}, Y.J. Lu ^{id65}, H.J. Lubatti ^{id137}, C. Luci ^{id74a,74b}, F.L. Lucio Alves ^{id14c}, A. Lucotte ^{id60},
 F. Luehring ^{id67}, I. Luise ^{id144}, O. Lukianchuk ^{id66}, O. Lundberg ^{id143}, B. Lund-Jensen ^{id143},
 N.A. Luongo ^{id122}, M.S. Lutz ^{id150}, D. Lynn ^{id29}, H. Lyons⁹¹, R. Lysak ^{id130}, E. Lytken ^{id97}, F. Lyu ^{id14a},
 V. Lyubushkin ^{id38}, T. Lyubushkina ^{id38}, H. Ma ^{id29}, L.L. Ma ^{id62b}, Y. Ma ^{id95}, D.M. Mac Donell ^{id164},
 G. Maccarrone ^{id53}, J.C. MacDonald ^{id138}, R. Madar ^{id40}, W.F. Mader ^{id50}, J. Maeda ^{id83}, T. Maeno ^{id29},
 M. Maerker ^{id50}, V. Magerl ^{id54}, H. Maguire ^{id138}, D.J. Mahon ^{id41}, C. Maidantchik ^{id81b},
 A. Maio ^{id129a,129b,129d}, K. Maj ^{id84a}, O. Majersky ^{id28a}, S. Majewski ^{id122}, N. Makovec ^{id66},
 V. Maksimovic ^{id15}, B. Malaescu ^{id126}, Pa. Malecki ^{id85}, V.P. Maleev ^{id37}, F. Malek ^{id60},
 D. Malito ^{id43b,43a}, U. Mallik ^{id79}, C. Malone ^{id32}, S. Maltezos¹⁰, S. Malyukov³⁸, J. Mamuzic ^{id13},
 G. Mancini ^{id53}, G. Manco ^{id72a,72b}, J.P. Mandalia ^{id93}, I. Mandić ^{id92},
 L. Manhaes de Andrade Filho ^{id81a}, I.M. Maniatis ^{id151,f}, M. Manisha ^{id134}, J. Manjarres Ramos ^{id50},
 D.C. Mankad ^{id168}, A. Mann ^{id108}, B. Mansoulie ^{id134}, S. Manzoni ^{id36}, A. Marantis ^{id151},
 G. Marchiori ^{id5}, M. Marcisovsky ^{id130}, L. Marcoccia ^{id75a,75b}, C. Marcon ^{id70a,70b}, M. Marinescu ^{id20},
 M. Marjanovic ^{id119}, E.J. Marshall ^{id90}, Z. Marshall ^{id17a}, S. Marti-Garcia ^{id162}, T.A. Martin ^{id166},
 V.J. Martin ^{id52}, B. Martin dit Latour ^{id16}, L. Martinelli ^{id74a,74b}, M. Martinez ^{id13,v},
 P. Martinez Agullo ^{id162}, V.I. Martinez Outschoorn ^{id102}, P. Martinez Suarez ^{id13}, S. Martin-Haugh ^{id133},
 V.S. Martoiu ^{id27b}, A.C. Martyniuk ^{id95}, A. Marzin ^{id36}, S.R. Maschek ^{id109}, L. Masetti ^{id99},
 T. Mashimo ^{id152}, J. Masik ^{id100}, A.L. Maslennikov ^{id37}, L. Massa ^{id23b}, P. Massarotti ^{id71a,71b},
 P. Mastrandrea ^{id73a,73b}, A. Mastroberardino ^{id43b,43a}, T. Masubuchi ^{id152}, T. Mathisen ^{id160},
 N. Matsuzawa¹⁵², J. Maurer ^{id27b}, B. Maček ^{id92}, D.A. Maximov ^{id37}, R. Mazini ^{id147}, I. Maznas ^{id151,f},
 M. Mazza ^{id106}, S.M. Mazza ^{id135}, C. Mc Ginn ^{id29,ah}, J.P. Mc Gowan ^{id103}, S.P. Mc Kee ^{id105},
 W.P. McCormack ^{id17a}, E.F. McDonald ^{id104}, A.E. McDougall ^{id113}, J.A. Mcfayden ^{id145},
 G. Mchedlidze ^{id148b}, R.P. Mckenzie ^{id33g}, T.C. Mclachlan ^{id48}, D.J. Mclaughlin ^{id95}, K.D. McLean ^{id164},
 S.J. McMahan ^{id133}, P.C. McNamara ^{id104}, C.M. Mccpartland ^{id91}, R.A. McPherson ^{id164,x}, T. Megy ^{id40},
 S. Mehlhase ^{id108}, A. Mehta ^{id91}, B. Meirose ^{id45}, D. Melini ^{id149}, B.R. Mellado Garcia ^{id33g},
 A.H. Melo ^{id55}, F. Meloni ^{id48}, E.D. Mendes Gouveia ^{id129a}, A.M. Mendes Jacques Da Costa ^{id20},
 H.Y. Meng ^{id154}, L. Meng ^{id90}, S. Menke ^{id109}, M. Mentink ^{id36}, E. Meoni ^{id43b,43a}, C. Merlassino ^{id125},
 L. Merola ^{id71a,71b}, C. Meroni ^{id70a}, G. Merz¹⁰⁵, O. Meshkov ^{id37}, J.K.R. Meshreki ^{id140}, J. Metcalfe ^{id6},
 A.S. Mete ^{id6}, C. Meyer ^{id67}, J-P. Meyer ^{id134}, M. Michetti ^{id18}, R.P. Middleton ^{id133}, L. Mijović ^{id52},
 G. Mikenberg ^{id168}, M. Mikesikova ^{id130}, M. Mikuž ^{id92}, H. Mildner ^{id138}, A. Milic ^{id36},
 C.D. Milke ^{id44}, D.W. Miller ^{id39}, L.S. Miller ^{id34}, A. Milov ^{id168}, D.A. Milstead^{47a,47b}, T. Min^{14c},
 A.A. Minaenko ^{id37}, I.A. Minashvili ^{id148b}, L. Mince ^{id59}, A.I. Mincer ^{id116}, B. Mindur ^{id84a},
 M. Mineev ^{id38}, Y. Mino ^{id86}, L.M. Mir ^{id13}, M. Miralles Lopez ^{id162}, M. Mironova ^{id125},
 M.C. Missio ^{id112}, T. Mitani ^{id167}, A. Mitra ^{id166}, V.A. Mitsou ^{id162}, O. Miu ^{id154}, P.S. Miyagawa ^{id93},
 Y. Miyazaki⁸⁸, A. Mizukami ^{id82}, J.U. Mjörnmark ^{id97}, T. Mkrtchyan ^{id63a}, T. Mlinarevic ^{id95},
 M. Mlynarikova ^{id36}, T. Moa ^{id47a,47b}, S. Mobius ^{id55}, K. Mochizuki ^{id107}, P. Moder ^{id48}, P. Mogg ^{id108},
 A.F. Mohammed ^{id14a,14d}, S. Mohapatra ^{id41}, G. Mokgatitwane ^{id33g}, B. Mondal ^{id140}, S. Mondal ^{id131},
 K. Mönig ^{id48}, E. Monnier ^{id101}, L. Monsonis Romero¹⁶², J. Montejo Berlingen ^{id36}, M. Montella ^{id118},
 F. Monticelli ^{id89}, N. Morange ^{id66}, A.L. Moreira De Carvalho ^{id129a}, M. Moreno Llácer ^{id162},
 C. Moreno Martinez ^{id56}, P. Morettini ^{id57b}, S. Morgenstern ^{id166}, M. Morii ^{id61}, M. Morinaga ^{id152},
 A.K. Morley ^{id36}, F. Morodei ^{id74a,74b}, L. Morvaj ^{id36}, P. Moschovakos ^{id36}, B. Moser ^{id36},
 M. Mosidze^{148b}, T. Moskalets ^{id54}, P. Moskvitina ^{id112}, J. Moss ^{id31,p}, E.J.W. Moyse ^{id102},
 O. Mtintsilana ^{id33g}, S. Muanza ^{id101}, J. Mueller ^{id128}, D. Muenstermann ^{id90}, R. Müller ^{id19},
 G.A. Mullier ^{id160}, J.J. Mullin¹²⁷, D.P. Mungo ^{id154}, J.L. Munoz Martinez ^{id13}, D. Munoz Perez ^{id162},
 F.J. Munoz Sanchez ^{id100}, M. Murin ^{id100}, W.J. Murray ^{id166,133}, A. Murrone ^{id70a,70b}, J.M. Muse ^{id119},

M. Muškinja ^{17a}, C. Mwewa ²⁹, A.G. Myagkov ^{37,a}, A.J. Myers ⁸, A.A. Myers ¹²⁸, G. Myers ⁶⁷,
M. Myska ¹³¹, B.P. Nachman ^{17a}, O. Nackenhorst ⁴⁹, A. Nag ⁵⁰, K. Nagai ¹²⁵, K. Nagano ⁸²,
J.L. Nagle ^{29,ah}, E. Nagy ¹⁰¹, A.M. Nairz ³⁶, Y. Nakahama ⁸², K. Nakamura ⁸², H. Nanjo ¹²³,
R. Narayan ⁴⁴, E.A. Narayanan ¹¹¹, I. Naryshkin ³⁷, M. Naseri ³⁴, C. Nass ²⁴, G. Navarro ^{22a},
J. Navarro-Gonzalez ¹⁶², R. Nayak ¹⁵⁰, A. Nayaz ¹⁸, P.Y. Nechaeva ³⁷, F. Nechansky ⁴⁸,
L. Nedic ¹²⁵, T.J. Neep ²⁰, A. Negri ^{72a,72b}, M. Negrini ^{23b}, C. Nellist ¹¹², C. Nelson ¹⁰³,
K. Nelson ¹⁰⁵, S. Nemecek ¹³⁰, M. Nessi ^{36,i}, M.S. Neubauer ¹⁶¹, F. Neuhaus ⁹⁹, J. Neundorf ⁴⁸,
R. Newhouse ¹⁶³, P.R. Newman ²⁰, C.W. Ng ¹²⁸, Y.S. Ng ¹⁸, Y.W.Y. Ng ⁴⁸, B. Ngair ^{35e},
H.D.N. Nguyen ¹⁰⁷, R.B. Nickerson ¹²⁵, R. Nicolaidou ¹³⁴, J. Nielsen ¹³⁵, M. Niemeyer ⁵⁵,
N. Nikiforou ³⁶, V. Nikolaenko ^{37,a}, I. Nikolic-Audit ¹²⁶, K. Nikolopoulos ²⁰, P. Nilsson ²⁹,
H.R. Nindhito ⁵⁶, A. Nisati ^{74a}, N. Nishu ², R. Nisius ¹⁰⁹, J-E. Nitschke ⁵⁰,
E.K. Nkadimeng ^{33g}, S.J. Noacco Rosende ⁸⁹, T. Nobe ¹⁵², D.L. Noel ³², Y. Noguchi ⁸⁶,
T. Nommensen ¹⁴⁶, M.A. Nomura ²⁹, M.B. Norfolk ¹³⁸, R.R.B. Norisam ⁹⁵, B.J. Norman ³⁴,
J. Novak ⁹², T. Novak ⁴⁸, O. Novgorodova ⁵⁰, L. Novotny ¹³¹, R. Novotny ¹¹¹, L. Nozka ¹²¹,
K. Ntekas ¹⁵⁹, N.M.J. Nunes De Moura Junior ^{81b}, E. Nurse ⁹⁵, F.G. Oakham ^{34,ae}, J. Ocariz ¹²⁶,
A. Ochi ⁸³, I. Ochoa ^{129a}, S. Oerdek ¹⁶⁰, A. Ogrodnik ^{84a}, A. Oh ¹⁰⁰, C.C. Ohm ¹⁴³,
H. Oide ⁸², R. Oishi ¹⁵², M.L. Ojeda ⁴⁸, Y. Okazaki ⁸⁶, M.W. O'Keefe ⁹¹, Y. Okumura ¹⁵²,
A. Olariu ^{27b}, L.F. Oleiro Seabra ^{129a}, S.A. Olivares Pino ^{136e}, D. Oliveira Damazio ²⁹,
D. Oliveira Goncalves ^{81a}, J.L. Oliver ¹⁵⁹, M.J.R. Olsson ¹⁵⁹, A. Olszewski ⁸⁵, J. Olszowska ^{85,*},
Ö.O. Öncel ⁵⁴, D.C. O'Neil ¹⁴¹, A.P. O'Neill ¹⁹, A. Onofre ^{129a,129e}, P.U.E. Onyisi ¹¹,
M.J. Oreglia ³⁹, G.E. Orellana ⁸⁹, D. Orestano ^{76a,76b}, N. Orlando ¹³, R.S. Orr ¹⁵⁴,
V. O'Shea ⁵⁹, R. Ospanov ^{62a}, G. Otero y Garzon ³⁰, H. Otono ⁸⁸, P.S. Ott ^{63a}, G.J. Ottino ^{17a},
M. Ouchrif ^{35d}, J. Ouellette ^{29,ah}, F. Ould-Saada ¹²⁴, M. Owen ⁵⁹, R.E. Owen ¹³³,
K.Y. Oyulmaz ^{21a}, V.E. Ozcan ^{21a}, N. Ozturk ⁸, S. Ozturk ^{21d}, J. Pacalt ¹²¹, H.A. Pacey ³²,
K. Pachal ⁵¹, A. Pacheco Pages ¹³, C. Padilla Aranda ¹³, G. Padovano ^{74a,74b}, S. Pagan Griso ^{17a},
G. Palacino ⁶⁷, A. Palazzo ^{69a,69b}, S. Palestini ³⁶, M. Palka ^{84b}, J. Pan ¹⁷¹, T. Pan ^{64a},
D.K. Panchal ¹¹, C.E. Pandini ¹¹³, J.G. Panduro Vazquez ⁹⁴, H. Pang ^{14b}, P. Pani ⁴⁸,
G. Panizzo ^{68a,68c}, L. Paolozzi ⁵⁶, C. Papadatos ¹⁰⁷, S. Parajuli ⁴⁴, A. Paramonov ⁶,
C. Paraskevopoulos ¹⁰, D. Paredes Hernandez ^{64b}, T.H. Park ¹⁵⁴, M.A. Parker ³², F. Parodi ^{57b,57a},
E.W. Parrish ¹¹⁴, V.A. Parrish ⁵², J.A. Parsons ⁴¹, U. Parzefall ⁵⁴, B. Pascual Dias ¹⁰⁷,
L. Pascual Dominguez ¹⁵⁰, V.R. Pascuzzi ^{17a}, F. Pasquali ¹¹³, E. Pasqualucci ^{74a}, S. Passaggio ^{57b},
F. Pastore ⁹⁴, P. Pasuwan ^{47a,47b}, P. Patel ⁸⁵, J.R. Pater ¹⁰⁰, T. Pauly ³⁶, J. Pearkes ¹⁴²,
M. Pedersen ¹²⁴, R. Pedro ^{129a}, S.V. Peleganchuk ³⁷, O. Penc ³⁶, E.A. Pender ⁵², C. Peng ^{64b},
H. Peng ^{62a}, K.E. Pensi ¹⁰⁸, M. Penzin ³⁷, B.S. Peralva ^{81d,81d}, A.P. Pereira Peixoto ⁶⁰,
L. Pereira Sanchez ^{47a,47b}, D.V. Perepelitsa ^{29,ah}, E. Perez Codina ^{155a}, M. Perganti ¹⁰,
L. Perini ^{70a,70b,*}, H. Pernegger ³⁶, S. Perrella ³⁶, A. Perrevoort ¹¹², O. Perrin ⁴⁰, K. Peters ⁴⁸,
R.F.Y. Peters ¹⁰⁰, B.A. Petersen ³⁶, T.C. Petersen ⁴², E. Petit ¹⁰¹, V. Petousis ¹³¹,
C. Petridou ^{151,f}, A. Petrukhin ¹⁴⁰, M. Pettee ^{17a}, N.E. Pettersson ³⁶, A. Petukhov ³⁷,
K. Petukhova ¹³², A. Peyaud ¹³⁴, R. Pezoa ^{136f}, L. Pezzotti ³⁶, G. Pezzullo ¹⁷¹, T.M. Pham ¹⁶⁹,
T. Pham ¹⁰⁴, P.W. Phillips ¹³³, M.W. Phipps ¹⁶¹, G. Piacquadio ¹⁴⁴, E. Pianori ^{17a},
F. Piazza ^{70a,70b}, R. Piegai ³⁰, D. Pietreanu ^{27b}, A.D. Pilkington ¹⁰⁰, M. Pinamonti ^{68a,68c},
J.L. Pinfeld ², G. Pinheiro Matos ⁴¹, B.C. Pinheiro Pereira ^{129a}, C. Pitman Donaldson ⁹⁵, D.A. Pizzi ³⁴,
L. Pizzimento ^{75a,75b}, A. Pizzini ¹¹³, M.-A. Pleier ²⁹, V. Plesanovs ⁵⁴, V. Pleskot ¹³²,
E. Plotnikova ³⁸, G. Poddar ⁴, R. Poettgen ⁹⁷, L. Poggioli ¹²⁶, I. Pogrebnyak ¹⁰⁶, D. Pohl ²⁴,
I. Pokharel ⁵⁵, S. Polacek ¹³², G. Polesello ^{72a}, A. Poley ^{141,155a}, R. Polifka ¹³¹, A. Polini ^{23b},
C.S. Pollard ¹²⁵, Z.B. Pollock ¹¹⁸, V. Polychronakos ²⁹, E. Pompa Pacchi ^{74a,74b},
D. Ponomarenko ³⁷, L. Pontecorvo ³⁶, S. Popa ^{27a}, G.A. Popeneciu ^{27d},

D.M. Portillo Quintero [ID155a](#), S. Pospisil [ID131](#), P. Postolache [ID27c](#), K. Potamianos [ID125](#), I.N. Potrap [ID38](#),
 C.J. Potter [ID32](#), H. Potti [ID1](#), T. Poulsen [ID48](#), J. Poveda [ID162](#), M.E. Pozo Astigarraga [ID36](#),
 A. Prades Ibanez [ID162](#), M.M. Prapa [ID46](#), J. Pretel [ID54](#), D. Price [ID100](#), M. Primavera [ID69a](#),
 M.A. Principe Martin [ID98](#), R. Privara [ID121](#), M.L. Proffitt [ID137](#), N. Proklova [ID127](#), K. Prokofiev [ID64c](#),
 G. Proto [ID75a,75b](#), S. Protopopescu [ID29](#), J. Proudfoot [ID6](#), M. Przybycien [ID84a](#), J.E. Puddefoot [ID138](#),
 D. Pudzha [ID37](#), P. Puzo [ID66](#), D. Pyatiizbyantseva [ID37](#), J. Qian [ID105](#), D. Qichen [ID100](#), Y. Qin [ID100](#),
 T. Qiu [ID93](#), A. Quadt [ID55](#), M. Queitsch-Maitland [ID100](#), G. Quetant [ID56](#), G. Rabanal Bolanos [ID61](#),
 D. Rafanoharana [ID54](#), F. Ragusa [ID70a,70b](#), J.L. Rainbolt [ID39](#), J.A. Raine [ID56](#), S. Rajagopalan [ID29](#),
 E. Ramakoti [ID37](#), K. Ran [ID48,14d](#), N.P. Rapheeha [ID33g](#), V. Raskina [ID126](#), D.F. Rassloff [ID63a](#), S. Rave [ID99](#),
 B. Ravina [ID55](#), I. Ravinovich [ID168](#), M. Raymond [ID36](#), A.L. Read [ID124](#), N.P. Readioff [ID138](#),
 D.M. Rebuzzi [ID72a,72b](#), G. Redlinger [ID29](#), K. Reeves [ID45](#), J.A. Reidelsturz [ID170](#), D. Reikher [ID150](#),
 A. Reiss [ID99](#), A. Rej [ID140](#), C. Rembser [ID36](#), A. Renardi [ID48](#), M. Renda [ID27b](#), M.B. Rendel [ID109](#), F. Renner [ID48](#),
 A.G. Rennie [ID59](#), S. Resconi [ID70a](#), M. Ressegotti [ID57b,57a](#), E.D. Resseguie [ID17a](#), S. Rettie [ID36](#),
 J.G. Reyes Rivera [ID106](#), B. Reynolds [ID118](#), E. Reynolds [ID17a](#), M. Rezaei Estabragh [ID170](#), O.L. Rezanova [ID37](#),
 P. Reznicek [ID132](#), E. Ricci [ID77a,77b](#), R. Richter [ID109](#), S. Richter [ID47a,47b](#), E. Richter-Was [ID84b](#),
 M. Ridel [ID126](#), P. Rieck [ID116](#), P. Riedler [ID36](#), M. Rijssenbeek [ID144](#), A. Rimoldi [ID72a,72b](#), M. Rimoldi [ID48](#),
 L. Rinaldi [ID23b,23a](#), T.T. Rinn [ID29](#), M.P. Rinnagel [ID108](#), G. Ripellino [ID143](#), I. Riu [ID13](#), P. Rivadeneira [ID48](#),
 J.C. Rivera Vergara [ID164](#), F. Rizatdinova [ID120](#), E. Rizvi [ID93](#), C. Rizzi [ID56](#), B.A. Roberts [ID166](#),
 B.R. Roberts [ID17a](#), S.H. Robertson [ID103,x](#), M. Robin [ID48](#), D. Robinson [ID32](#), C.M. Robles Gajardo [ID136f](#),
 M. Robles Manzano [ID99](#), A. Robson [ID59](#), A. Rocchi [ID75a,75b](#), C. Roda [ID73a,73b](#), S. Rodriguez Bosca [ID63a](#),
 Y. Rodriguez Garcia [ID22a](#), A. Rodriguez Rodriguez [ID54](#), A.M. Rodríguez Vera [ID155b](#), S. Roe [ID36](#),
 J.T. Roemer [ID159](#), A.R. Roepe-Gier [ID119](#), J. Roggel [ID170](#), O. Røhne [ID124](#), R.A. Rojas [ID164](#), B. Roland [ID54](#),
 C.P.A. Roland [ID67](#), J. Roloff [ID29](#), A. Romaniouk [ID37](#), E. Romano [ID72a,72b](#), M. Romano [ID23b](#),
 A.C. Romero Hernandez [ID161](#), N. Rompotis [ID91](#), L. Roos [ID126](#), S. Rosati [ID74a](#), B.J. Rosser [ID39](#),
 E. Rossi [ID4](#), E. Rossi [ID71a,71b](#), L.P. Rossi [ID57b](#), L. Rossini [ID48](#), R. Rosten [ID118](#), M. Rotaru [ID27b](#),
 B. Rottler [ID54](#), D. Rousseau [ID66](#), D. Rousso [ID32](#), G. Rovelli [ID72a,72b](#), A. Roy [ID161](#), A. Rozanov [ID101](#),
 Y. Rozen [ID149](#), X. Ruan [ID33g](#), A. Rubio Jimenez [ID162](#), A.J. Ruby [ID91](#), V.H. Ruelas Rivera [ID18](#),
 T.A. Ruggeri [ID1](#), F. Rühr [ID54](#), A. Ruiz-Martinez [ID162](#), A. Rummler [ID36](#), Z. Rurikova [ID54](#),
 N.A. Rusakovich [ID38](#), H.L. Russell [ID164](#), J.P. Rutherford [ID7](#), K. Rybacki [ID90](#), M. Rybar [ID132](#),
 E.B. Rye [ID124](#), A. Ryzhov [ID37](#), J.A. Sabater Iglesias [ID56](#), P. Sabatini [ID162](#), L. Sabetta [ID74a,74b](#),
 H.F-W. Sadrozinski [ID135](#), F. Safai Tehrani [ID74a](#), B. Safarzadeh Samani [ID145](#), M. Safdari [ID142](#),
 S. Saha [ID103](#), M. Sahinsoy [ID109](#), M. Saimpert [ID134](#), M. Saito [ID152](#), T. Saito [ID152](#), D. Salamani [ID36](#),
 G. Salamanna [ID76a,76b](#), A. Salnikov [ID142](#), J. Salt [ID162](#), A. Salvador Salas [ID13](#), D. Salvatore [ID43b,43a](#),
 F. Salvatore [ID145](#), A. Salzburger [ID36](#), D. Sammel [ID54](#), D. Sampsonidis [ID151,f](#), D. Sampsonidou [ID62d,62c](#),
 J. Sánchez [ID162](#), A. Sanchez Pineda [ID4](#), V. Sanchez Sebastian [ID162](#), H. Sandaker [ID124](#), C.O. Sander [ID48](#),
 J.A. Sandesara [ID102](#), M. Sandhoff [ID170](#), C. Sandoval [ID22b](#), D.P.C. Sankey [ID133](#), A. Sansoni [ID53](#),
 L. Santi [ID74a,74b](#), C. Santoni [ID40](#), H. Santos [ID129a,129b](#), S.N. Santpur [ID17a](#), A. Santra [ID168](#),
 K.A. Saoucha [ID138](#), J.G. Saraiva [ID129a,129d](#), J. Sardain [ID7](#), O. Sasaki [ID82](#), K. Sato [ID156](#), C. Sauer [ID63b](#),
 F. Sauerburger [ID54](#), E. Sauvan [ID4](#), P. Savard [ID154,ae](#), R. Sawada [ID152](#), C. Sawyer [ID133](#), L. Sawyer [ID96](#),
 I. Sayago Galvan [ID162](#), C. Sbarra [ID23b](#), A. Sbrizzi [ID23b,23a](#), T. Scanlon [ID95](#), J. Schaarschmidt [ID137](#),
 P. Schacht [ID109](#), D. Schaefer [ID39](#), U. Schäfer [ID99](#), A.C. Schaffer [ID66](#), D. Schaile [ID108](#),
 R.D. Schamberger [ID144](#), E. Schanet [ID108](#), C. Scharf [ID18](#), M.M. Schefer [ID19](#), V.A. Schegelsky [ID37](#),
 D. Scheirich [ID132](#), F. Schenck [ID18](#), M. Schernau [ID159](#), C. Scheulen [ID55](#), C. Schiavi [ID57b,57a](#),
 Z.M. Schillaci [ID26](#), E.J. Schioppa [ID69a,69b](#), M. Schioppa [ID43b,43a](#), B. Schlag [ID99](#), K.E. Schleicher [ID54](#),
 S. Schlenker [ID36](#), J. Schmeing [ID170](#), M.A. Schmidt [ID170](#), K. Schmieden [ID99](#), C. Schmitt [ID99](#),
 S. Schmitt [ID48](#), L. Schoeffel [ID134](#), A. Schoening [ID63b](#), P.G. Scholer [ID54](#), E. Schopf [ID125](#), M. Schott [ID99](#),
 J. Schovancova [ID36](#), S. Schramm [ID56](#), F. Schroeder [ID170](#), H-C. Schultz-Coulon [ID63a](#), M. Schumacher [ID54](#),

B.A. Schumm [ID135](#), Ph. Schune [ID134](#), A. Schwartzman [ID142](#), T.A. Schwarz [ID105](#), Ph. Schwemling [ID134](#),
 R. Schwienhorst [ID106](#), A. Sciandra [ID135](#), G. Sciolla [ID26](#), F. Scuri [ID73a](#), F. Scutti [ID104](#), C.D. Sebastiani [ID91](#),
 K. Sedlaczek [ID49](#), P. Seema [ID18](#), S.C. Seidel [ID111](#), A. Seiden [ID135](#), B.D. Seidlitz [ID41](#), T. Seiss [ID39](#),
 C. Seitz [ID48](#), J.M. Seixas [ID81b](#), G. Sekhniaidze [ID71a](#), S.J. Sekula [ID44](#), L. Selem [ID4](#),
 N. Semprini-Cesari [ID23b,23a](#), S. Sen [ID51](#), D. Sengupta [ID56](#), V. Senthilkumar [ID162](#), L. Serin [ID66](#),
 L. Serkin [ID68a,68b](#), M. Sessa [ID76a,76b](#), H. Severini [ID119](#), S. Sevova [ID142](#), F. Sforza [ID57b,57a](#), A. Sfyrla [ID56](#),
 E. Shabalina [ID55](#), R. Shaheen [ID143](#), J.D. Shahinian [ID127](#), D. Shaked Renous [ID168](#), L.Y. Shan [ID14a](#),
 M. Shapiro [ID17a](#), A. Sharma [ID36](#), A.S. Sharma [ID163](#), P. Sharma [ID79](#), S. Sharma [ID48](#), P.B. Shatalov [ID37](#),
 K. Shaw [ID145](#), S.M. Shaw [ID100](#), Q. Shen [ID62c,5](#), P. Sherwood [ID95](#), L. Shi [ID95](#), C.O. Shimmin [ID171](#),
 Y. Shimogama [ID167](#), J.D. Shinner [ID94](#), I.P.J. Shipsey [ID125](#), S. Shirabe [ID60](#), M. Shiyakova [ID38](#),
 J. Shlomi [ID168](#), M.J. Shochet [ID39](#), J. Shojaii [ID104](#), D.R. Shope [ID124](#), S. Shrestha [ID118,ai](#), E.M. Shrif [ID33g](#),
 M.J. Shroff [ID164](#), P. Sicho [ID130](#), A.M. Sickles [ID161](#), E. Sideras Haddad [ID33g](#), A. Sidoti [ID23b](#),
 F. Siegert [ID50](#), Dj. Sijacki [ID15](#), R. Sikora [ID84a](#), F. Sili [ID89](#), J.M. Silva [ID20](#), M.V. Silva Oliveira [ID36](#),
 S.B. Silverstein [ID47a](#), S. Simion [ID66](#), R. Simoniello [ID36](#), E.L. Simpson [ID59](#), N.D. Simpson [ID97](#),
 S. Simsek [ID21d](#), S. Sindhu [ID55](#), P. Sinervo [ID154](#), V. Sinetckii [ID37](#), S. Singh [ID141](#), S. Singh [ID154](#),
 S. Sinha [ID48](#), S. Sinha [ID33g](#), M. Sioli [ID23b,23a](#), I. Siral [ID36](#), S.Yu. Sivoklov [ID37,*](#), J. Sjölin [ID47a,47b](#),
 A. Skaf [ID55](#), E. Skorda [ID97](#), P. Skubic [ID119](#), M. Slawinska [ID85](#), V. Smakhtin [ID168](#), B.H. Smart [ID133](#),
 J. Smiesko [ID36](#), S.Yu. Smirnov [ID37](#), Y. Smirnov [ID37](#), L.N. Smirnova [ID37,a](#), O. Smirnova [ID97](#),
 A.C. Smith [ID41](#), E.A. Smith [ID39](#), H.A. Smith [ID125](#), J.L. Smith [ID91](#), R. Smith [ID142](#), M. Smizanska [ID90](#),
 K. Smolek [ID131](#), A. Smykiewicz [ID85](#), A.A. Snesarev [ID37](#), H.L. Snoek [ID113](#), S. Snyder [ID29](#),
 R. Sobie [ID164,x](#), A. Soffer [ID150](#), C.A. Solans Sanchez [ID36](#), E.Yu. Soldatov [ID37](#), U. Soldevila [ID162](#),
 A.A. Solodkov [ID37](#), S. Solomon [ID54](#), A. Soloshenko [ID38](#), K. Solovieva [ID54](#), O.V. Solovyanov [ID37](#),
 V. Solovyev [ID37](#), P. Sommer [ID36](#), A. Sonay [ID13](#), W.Y. Song [ID155b](#), A. Sopczak [ID131](#), A.L. Sopio [ID95](#),
 F. Sopkova [ID28b](#), V. Sothilingam [ID63a](#), S. Sottocornola [ID72a,72b](#), R. Soualah [ID115b](#), Z. Soumami [ID35e](#),
 D. South [ID48](#), S. Spagnolo [ID69a,69b](#), M. Spalla [ID109](#), F. Spanò [ID94](#), D. Sperlich [ID54](#), G. Spigo [ID36](#),
 M. Spina [ID145](#), S. Spinali [ID90](#), D.P. Spiteri [ID59](#), M. Spousta [ID132](#), E.J. Staats [ID34](#), A. Stabile [ID70a,70b](#),
 R. Stamen [ID63a](#), M. Stamenkovic [ID113](#), A. Stampekis [ID20](#), M. Standke [ID24](#), E. Stanecka [ID85](#),
 M.V. Stange [ID50](#), B. Stanislaus [ID17a](#), M.M. Stanitzki [ID48](#), M. Stankaityte [ID125](#), B. Stapf [ID48](#),
 E.A. Starchenko [ID37](#), G.H. Stark [ID135](#), J. Stark [ID101](#), D.M. Starke [ID155b](#), P. Staroba [ID130](#),
 P. Starovoitov [ID63a](#), S. Stärz [ID103](#), R. Staszewski [ID85](#), G. Stavropoulos [ID46](#), J. Steentoft [ID160](#),
 P. Steinberg [ID29](#), A.L. Steinhebel [ID122](#), B. Stelzer [ID141,155a](#), H.J. Stelzer [ID128](#), O. Stelzer-Chilton [ID155a](#),
 H. Stenzel [ID58](#), T.J. Stevenson [ID145](#), G.A. Stewart [ID36](#), M.C. Stockton [ID36](#), G. Stoicea [ID27b](#),
 M. Stolarski [ID129a](#), S. Stonjek [ID109](#), A. Straessner [ID50](#), J. Strandberg [ID143](#), S. Strandberg [ID47a,47b](#),
 M. Strauss [ID119](#), T. Strebler [ID101](#), P. Strizeneč [ID28b](#), R. Ströhmer [ID165](#), D.M. Strom [ID122](#), L.R. Strom [ID48](#),
 R. Stroynowski [ID44](#), A. Strubig [ID47a,47b](#), S.A. Stucci [ID29](#), B. Stugu [ID16](#), J. Stupak [ID119](#), N.A. Styles [ID48](#),
 D. Su [ID142](#), S. Su [ID62a](#), W. Su [ID62d,137,62c](#), X. Su [ID62a,66](#), K. Sugizaki [ID152](#), V.V. Sulin [ID37](#),
 M.J. Sullivan [ID91](#), D.M.S. Sultan [ID77a,77b](#), L. Sultanaliyeva [ID37](#), S. Sultansoy [ID3b](#), T. Sumida [ID86](#),
 S. Sun [ID105](#), S. Sun [ID169](#), O. Sunneborn Gudnadottir [ID160](#), M.R. Sutton [ID145](#), M. Svatos [ID130](#),
 M. Swiatlowski [ID155a](#), T. Swirski [ID165](#), I. Sykora [ID28a](#), M. Sykora [ID132](#), T. Sykora [ID132](#), D. Ta [ID99](#),
 K. Tackmann [ID48,w](#), A. Taffard [ID159](#), R. Tafirout [ID155a](#), J.S. Tafuya Vargas [ID66](#), R.H.M. Taibah [ID126](#),
 R. Takashima [ID87](#), K. Takeda [ID83](#), E.P. Takeva [ID52](#), Y. Takubo [ID82](#), M. Talby [ID101](#), A.A. Talyshev [ID37](#),
 K.C. Tam [ID64b](#), N.M. Tamir [ID150](#), A. Tanaka [ID152](#), J. Tanaka [ID152](#), R. Tanaka [ID66](#), M. Tanasini [ID57b,57a](#),
 J. Tang [ID62c](#), Z. Tao [ID163](#), S. Tapia Araya [ID80](#), S. Tapprogge [ID99](#), A. Tarek Abouelfadl Mohamed [ID106](#),
 S. Tarem [ID149](#), K. Tariq [ID62b](#), G. Tarna [ID101,27b](#), G.F. Tartarelli [ID70a](#), P. Tas [ID132](#), M. Tasevsky [ID130](#),
 E. Tassi [ID43b,43a](#), A.C. Tate [ID161](#), G. Tateno [ID152](#), Y. Tayalati [ID35e](#), G.N. Taylor [ID104](#), W. Taylor [ID155b](#),
 H. Teagle [ID91](#), A.S. Tee [ID169](#), R. Teixeira De Lima [ID142](#), P. Teixeira-Dias [ID94](#), J.J. Teoh [ID154](#),
 K. Terashi [ID152](#), J. Terron [ID98](#), S. Terzo [ID13](#), M. Testa [ID53](#), R.J. Teuscher [ID154,x](#), A. Thaler [ID78](#),

O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Thevenaux-Pelzer ¹⁸, O. Thielmann ¹⁷⁰, D.W. Thomas ⁹⁴, J.P. Thomas ²⁰, E.A. Thompson ⁴⁸, P.D. Thompson ²⁰, E. Thomson ¹²⁷, E.J. Thorpe ⁹³, Y. Tian ⁵⁵, V. Tikhomirov ^{37,a}, Yu.A. Tikhonov ³⁷, S. Timoshenko ³⁷, E.X.L. Ting ¹, P. Tipton ¹⁷¹, S. Tisserant ¹⁰¹, S.H. Tlou ^{33g}, A. Tnourji ⁴⁰, K. Todome ^{23b,23a}, S. Todorova-Nova ¹³², S. Todt ⁵⁰, M. Togawa ⁸², J. Tojo ⁸⁸, S. Tokár ^{28a}, K. Tokushuku ⁸², R. Tombs ³², M. Tomoto ^{82,110}, L. Tompkins ¹⁴², K.W. Topolnicki ^{84b}, P. Tornambe ¹⁰², E. Torrence ¹²², H. Torres ⁵⁰, E. Torró Pastor ¹⁶², M. Toscani ³⁰, C. Tosciri ³⁹, M. Tost ¹¹, D.R. Tovey ¹³⁸, A. Traeet ¹⁶, I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁵, A. Tricoli ²⁹, I.M. Trigger ^{155a}, S. Trincaz-Duvoid ¹²⁶, D.A. Trischuk ²⁶, B. Trocmé ⁶⁰, A. Trofymov ⁶⁶, C. Troncon ^{70a}, L. Truong ^{33c}, M. Trzebinski ⁸⁵, A. Trzupiek ⁸⁵, F. Tsai ¹⁴⁴, M. Tsai ¹⁰⁵, A. Tsiamis ^{151,f}, P.V. Tsiareshka ³⁷, S. Tsigaridas ^{155a}, A. Tsirigotis ^{151,u}, V. Tsiskaridze ¹⁴⁴, E.G. Tskhadadze ^{148a}, M. Tsopoulou ^{151,f}, Y. Tsujikawa ⁸⁶, I.I. Tsukerman ³⁷, V. Tsulaia ^{17a}, S. Tsuno ⁸², O. Tsur ¹⁴⁹, D. Tsybychev ¹⁴⁴, Y. Tu ^{64b}, A. Tudorache ^{27b}, V. Tudorache ^{27b}, A.N. Tuna ³⁶, S. Turchikhin ³⁸, I. Turk Cakir ^{3a}, R. Turra ^{70a}, T. Turtuvshin ^{38,y}, P.M. Tuts ⁴¹, S. Tzamarias ^{151,f}, P. Tzanis ¹⁰, E. Tzovara ⁹⁹, K. Uchida ¹⁵², F. Ukegawa ¹⁵⁶, P.A. Ulloa Poblete ^{136c}, E.N. Umaka ⁸⁰, G. Unal ³⁶, M. Unal ¹¹, A. Undrus ²⁹, G. Unel ¹⁵⁹, J. Urban ^{28b}, P. Urquijo ¹⁰⁴, G. Usai ⁸, R. Ushioda ¹⁵³, M. Usman ¹⁰⁷, Z. Uysal ^{21b}, L. Vacavant ¹⁰¹, V. Vacek ¹³¹, B. Vachon ¹⁰³, K.O.H. Vadla ¹²⁴, T. Vafeiadis ³⁶, A. Vaitkus ⁹⁵, C. Valderanis ¹⁰⁸, E. Valdes Santurio ^{47a,47b}, M. Valente ^{155a}, S. Valentinetti ^{23b,23a}, A. Valero ¹⁶², A. Vallier ¹⁰¹, J.A. Valls Ferrer ¹⁶², T.R. Van Daalen ¹³⁷, P. Van Gemmeren ⁶, M. Van Rijnbach ^{124,36}, S. Van Stroud ⁹⁵, I. Van Vulpen ¹¹³, M. Vanadia ^{75a,75b}, W. Vandelli ³⁶, M. Vandenbroucke ¹³⁴, E.R. Vandewall ¹²⁰, D. Vannicola ¹⁵⁰, L. Vannoli ^{57b,57a}, R. Vari ^{74a}, E.W. Varnes ⁷, C. Varni ^{17a}, T. Varol ¹⁴⁷, D. Varouchas ⁶⁶, L. Varriale ¹⁶², K.E. Varvell ¹⁴⁶, M.E. Vasile ^{27b}, L. Vaslin ⁴⁰, G.A. Vasquez ¹⁶⁴, F. Vazeille ⁴⁰, T. Vazquez Schroeder ³⁶, J. Veatch ³¹, V. Vecchio ¹⁰⁰, M.J. Veen ¹⁰², I. Veliscek ¹²⁵, L.M. Veloce ¹⁵⁴, F. Veloso ^{129a,129c}, S. Veneziano ^{74a}, A. Ventura ^{69a,69b}, A. Verbytskyi ¹⁰⁹, M. Verducci ^{73a,73b}, C. Vergis ²⁴, M. Verissimo De Araujo ^{81b}, W. Verkerke ¹¹³, J.C. Vermeulen ¹¹³, C. Vernieri ¹⁴², P.J. Verschuuren ⁹⁴, M. Vessella ¹⁰², M.C. Vetterli ^{141,ae}, A. Vgenopoulos ^{151,f}, N. Viaux Maira ^{136f}, T. Vickey ¹³⁸, O.E. Vickey Boeriu ¹³⁸, G.H.A. Viehhauser ¹²⁵, L. Vigani ^{63b}, M. Villa ^{23b,23a}, M. Villaplana Perez ¹⁶², E.M. Villhauer ⁵², E. Vilucchi ⁵³, M.G. Vinciter ³⁴, G.S. Virdee ²⁰, A. Vishwakarma ⁵², C. Vittori ^{23b,23a}, I. Vivarelli ¹⁴⁵, V. Vladimirov ¹⁶⁶, E. Voevodina ¹⁰⁹, F. Vogel ¹⁰⁸, P. Vokac ¹³¹, J. Von Ahnen ⁴⁸, E. Von Toerne ²⁴, B. Vormwald ³⁶, V. Vorobel ¹³², K. Vorobev ³⁷, M. Vos ¹⁶², J.H. Vossebeld ⁹¹, M. Vozak ¹¹³, L. Vozdecky ⁹³, N. Vranjes ¹⁵, M. Vranjes Milosavljevic ¹⁵, M. Vreeswijk ¹¹³, R. Vuillermet ³⁶, O. Vujinovic ⁹⁹, I. Vukotic ³⁹, S. Wada ¹⁵⁶, C. Wagner ¹⁰², W. Wagner ¹⁷⁰, S. Wahdan ¹⁷⁰, H. Wahlberg ⁸⁹, R. Wakasa ¹⁵⁶, M. Wakida ¹¹⁰, V.M. Walbrecht ¹⁰⁹, J. Walder ¹³³, R. Walker ¹⁰⁸, W. Walkowiak ¹⁴⁰, A.M. Wang ⁶¹, A.Z. Wang ¹⁶⁹, C. Wang ^{62a}, C. Wang ^{62c}, H. Wang ^{17a}, J. Wang ^{64a}, R.-J. Wang ⁹⁹, R. Wang ⁶¹, R. Wang ⁶, S.M. Wang ¹⁴⁷, S. Wang ^{62b}, T. Wang ^{62a}, W.T. Wang ⁷⁹, X. Wang ^{14c}, X. Wang ¹⁶¹, X. Wang ^{62c}, Y. Wang ^{62d}, Y. Wang ^{14c}, Z. Wang ¹⁰⁵, Z. Wang ^{62d,51,62c}, Z. Wang ¹⁰⁵, A. Warburton ¹⁰³, R.J. Ward ²⁰, N. Warrack ⁵⁹, A.T. Watson ²⁰, H. Watson ⁵⁹, M.F. Watson ²⁰, G. Watts ¹³⁷, B.M. Waugh ⁹⁵, A.F. Webb ¹¹, C. Weber ²⁹, H.A. Weber ¹⁸, M.S. Weber ¹⁹, S.M. Weber ^{63a}, C. Wei ^{62a}, Y. Wei ¹²⁵, A.R. Weidberg ¹²⁵, J. Weingarten ⁴⁹, M. Weirich ⁹⁹, C. Weiser ⁵⁴, C.J. Wells ⁴⁸, T. Wenaus ²⁹, B. Wendland ⁴⁹, T. Wengler ³⁶, N.S. Wenke ¹⁰⁹, N. Wermes ²⁴, M. Wessels ^{63a}, K. Whalen ¹²², A.M. Wharton ⁹⁰, A.S. White ⁶¹, A. White ⁸, M.J. White ¹, D. Whiteson ¹⁵⁹, L. Wickremasinghe ¹²³, W. Wiedenmann ¹⁶⁹, C. Wiel ⁵⁰, M. Wielers ¹³³, N. Wieseotte ⁹⁹, C. Wiglesworth ⁴², L.A.M. Wiik-Fuchs ⁵⁴, D.J. Wilbern ¹¹⁹, H.G. Wilkens ³⁶, D.M. Williams ⁴¹, H.H. Williams ¹²⁷,

S. Williams ³², S. Willocq ¹⁰², P.J. Windischhofer ¹²⁵, F. Winklmeier ¹²², B.T. Winter ⁵⁴, J.K. Winter ¹⁰⁰, M. Wittgen ¹⁴², M. Wobisch ⁹⁶, R. Wölker ¹²⁵, J. Wollrath ¹⁵⁹, M.W. Wolter ⁸⁵, H. Wolters ^{129a,129c}, V.W.S. Wong ¹⁶³, A.F. Wongel ⁴⁸, S.D. Worm ⁴⁸, B.K. Wosiek ⁸⁵, K.W. Woźniak ⁸⁵, K. Wraight ⁵⁹, J. Wu ^{14a,14d}, M. Wu ^{64a}, M. Wu ¹¹², S.L. Wu ¹⁶⁹, X. Wu ⁵⁶, Y. Wu ^{62a}, Z. Wu ^{134,62a}, J. Wuerzinger ¹²⁵, T.R. Wyatt ¹⁰⁰, B.M. Wynne ⁵², S. Xella ⁴², L. Xia ^{14c}, M. Xia ^{14b}, J. Xiang ^{64c}, X. Xiao ¹⁰⁵, M. Xie ^{62a}, X. Xie ^{62a}, S. Xin ^{14a,14d}, J. Xiong ^{17a}, I. Xiotidis ¹⁴⁵, D. Xu ^{14a}, H. Xu ^{62a}, H. Xu ^{62a}, L. Xu ^{62a}, R. Xu ¹²⁷, T. Xu ¹⁰⁵, W. Xu ¹⁰⁵, Y. Xu ^{14b}, Z. Xu ^{62b}, Z. Xu ^{14a}, B. Yabsley ¹⁴⁶, S. Yacoob ^{33a}, N. Yamaguchi ⁸⁸, Y. Yamaguchi ¹⁵³, H. Yamauchi ¹⁵⁶, T. Yamazaki ^{17a}, Y. Yamazaki ⁸³, J. Yan ^{62c}, S. Yan ¹²⁵, Z. Yan ²⁵, H.J. Yang ^{62c,62d}, H.T. Yang ^{62a}, S. Yang ^{62a}, T. Yang ^{64c}, X. Yang ^{62a}, X. Yang ^{14a}, Y. Yang ⁴⁴, Z. Yang ^{62a,105}, W.-M. Yao ^{17a}, Y.C. Yap ⁴⁸, H. Ye ^{14c}, H. Ye ⁵⁵, J. Ye ⁴⁴, S. Ye ²⁹, X. Ye ^{62a}, Y. Yeh ⁹⁵, I. Yeletskikh ³⁸, B.K. Yeo ^{17a}, M.R. Yexley ⁹⁰, P. Yin ⁴¹, K. Yorita ¹⁶⁷, S. Younas ^{27b}, C.J.S. Young ⁵⁴, C. Young ¹⁴², M. Yuan ¹⁰⁵, R. Yuan ^{62b,1}, L. Yue ⁹⁵, X. Yue ^{63a}, M. Zaazoua ^{35e}, B. Zabinski ⁸⁵, E. Zaid ⁵², T. Zakareishvili ^{148b}, N. Zakharchuk ³⁴, S. Zambito ⁵⁶, J.A. Zamora Saa ^{136d,136b}, J. Zang ¹⁵², D. Zanzi ⁵⁴, O. Zaplatilek ¹³¹, S.V. Zeißner ⁴⁹, C. Zeitnitz ¹⁷⁰, J.C. Zeng ¹⁶¹, D.T. Zenger Jr ²⁶, O. Zenin ³⁷, T. Ženiš ^{28a}, S. Zenz ⁹³, S. Zerradi ^{35a}, D. Zerwas ⁶⁶, B. Zhang ^{14c}, D.F. Zhang ¹³⁸, G. Zhang ^{14b}, J. Zhang ^{62b}, J. Zhang ⁶, K. Zhang ^{14a,14d}, L. Zhang ^{14c}, P. Zhang ^{14a,14d}, R. Zhang ¹⁶⁹, S. Zhang ¹⁰⁵, T. Zhang ¹⁵², X. Zhang ^{62c}, X. Zhang ^{62b}, Y. Zhang ^{62c,5}, Z. Zhang ^{17a}, Z. Zhang ⁶⁶, H. Zhao ¹³⁷, P. Zhao ⁵¹, T. Zhao ^{62b}, Y. Zhao ¹³⁵, Z. Zhao ^{62a}, A. Zhemchugov ³⁸, X. Zheng ^{62a}, Z. Zheng ¹⁴², D. Zhong ¹⁶¹, B. Zhou ¹⁰⁵, C. Zhou ¹⁶⁹, H. Zhou ⁷, N. Zhou ^{62c}, Y. Zhou ⁷, C.G. Zhu ^{62b}, C. Zhu ^{14a,14d}, H.L. Zhu ^{62a}, H. Zhu ^{14a}, J. Zhu ¹⁰⁵, Y. Zhu ^{62c}, Y. Zhu ^{62a}, X. Zhuang ^{14a}, K. Zhukov ³⁷, V. Zhulanov ³⁷, N.I. Zimine ³⁸, J. Zinsser ^{63b}, M. Ziolkowski ¹⁴⁰, L. Živković ¹⁵, A. Zoccoli ^{23b,23a}, K. Zoch ⁵⁶, T.G. Zorbas ¹³⁸, O. Zormpa ⁴⁶, W. Zou ⁴¹, L. Zwalinski ³⁶.

¹Department of Physics, University of Adelaide, Adelaide; Australia.

²Department of Physics, University of Alberta, Edmonton AB; Canada.

³(^a)Department of Physics, Ankara University, Ankara; (^b)Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye.

⁴LAPP, Univ. Savoie Mont Blanc, CNRS/IN2P3, Annecy; France.

⁵APC, Université Paris Cité, CNRS/IN2P3, Paris; France.

⁶High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America.

⁷Department of Physics, University of Arizona, Tucson AZ; United States of America.

⁸Department of Physics, University of Texas at Arlington, Arlington TX; United States of America.

⁹Physics Department, National and Kapodistrian University of Athens, Athens; Greece.

¹⁰Physics Department, National Technical University of Athens, Zografou; Greece.

¹¹Department of Physics, University of Texas at Austin, Austin TX; United States of America.

¹²Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.

¹³Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain.

¹⁴(^a)Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (^b)Physics Department, Tsinghua University, Beijing; (^c)Department of Physics, Nanjing University, Nanjing; (^d)University of Chinese Academy of Science (UCAS), Beijing; China.

¹⁵Institute of Physics, University of Belgrade, Belgrade; Serbia.

¹⁶Department for Physics and Technology, University of Bergen, Bergen; Norway.

¹⁷(^a)Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (^b)University of California,

Berkeley CA; United States of America.

¹⁸Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany.

¹⁹Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland.

²⁰School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom.

²¹(^a)Department of Physics, Bogazici University, Istanbul; (^b)Department of Physics Engineering, Gaziantep University, Gaziantep; (^c)Department of Physics, Istanbul University, Istanbul; (^d)Istinye University, Sariyer, Istanbul; Türkiye.

²²(^a)Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (^b)Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia.

²³(^a)Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; (^b)INFN Sezione di Bologna; Italy.

²⁴Physikalisches Institut, Universität Bonn, Bonn; Germany.

²⁵Department of Physics, Boston University, Boston MA; United States of America.

²⁶Department of Physics, Brandeis University, Waltham MA; United States of America.

²⁷(^a)Transilvania University of Brasov, Brasov; (^b)Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (^c)Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (^d)National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (^e)University Politehnica Bucharest, Bucharest; (^f)West University in Timisoara, Timisoara; (^g)Faculty of Physics, University of Bucharest, Bucharest; Romania.

²⁸(^a)Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (^b)Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic.

²⁹Physics Department, Brookhaven National Laboratory, Upton NY; United States of America.

³⁰Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina.

³¹California State University, CA; United States of America.

³²Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom.

³³(^a)Department of Physics, University of Cape Town, Cape Town; (^b)iThemba Labs, Western Cape; (^c)Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (^d)National Institute of Physics, University of the Philippines Diliman (Philippines); (^e)University of South Africa, Department of Physics, Pretoria; (^f)University of Zululand, KwaDlangezwa; (^g)School of Physics, University of the Witwatersrand, Johannesburg; South Africa.

³⁴Department of Physics, Carleton University, Ottawa ON; Canada.

³⁵(^a)Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; (^b)Faculté des Sciences, Université Ibn-Tofail, Kénitra; (^c)Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; (^d)LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; (^e)Faculté des sciences, Université Mohammed V, Rabat; (^f)Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco.

³⁶CERN, Geneva; Switzerland.

³⁷Affiliated with an institute covered by a cooperation agreement with CERN.

³⁸Affiliated with an international laboratory covered by a cooperation agreement with CERN.

³⁹Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America.

⁴⁰LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France.

⁴¹Nevis Laboratory, Columbia University, Irvington NY; United States of America.

⁴²Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark.

⁴³(^a)Dipartimento di Fisica, Università della Calabria, Rende; (^b)INFN Gruppo Collegato di Cosenza,

Laboratori Nazionali di Frascati; Italy.

⁴⁴Physics Department, Southern Methodist University, Dallas TX; United States of America.

⁴⁵Physics Department, University of Texas at Dallas, Richardson TX; United States of America.

⁴⁶National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece.

⁴⁷(^a) Department of Physics, Stockholm University; (^b) Oskar Klein Centre, Stockholm; Sweden.

⁴⁸Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany.

⁴⁹Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany.

⁵⁰Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany.

⁵¹Department of Physics, Duke University, Durham NC; United States of America.

⁵²SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom.

⁵³INFN e Laboratori Nazionali di Frascati, Frascati; Italy.

⁵⁴Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany.

⁵⁵II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany.

⁵⁶Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.

⁵⁷(^a) Dipartimento di Fisica, Università di Genova, Genova; (^b) INFN Sezione di Genova; Italy.

⁵⁸II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany.

⁵⁹SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom.

⁶⁰LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France.

⁶¹Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America.

⁶²(^a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; (^b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; (^c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; (^d) Tsung-Dao Lee Institute, Shanghai; China.

⁶³(^a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; (^b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany.

⁶⁴(^a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; (^b) Department of Physics, University of Hong Kong, Hong Kong; (^c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China.

⁶⁵Department of Physics, National Tsing Hua University, Hsinchu; Taiwan.

⁶⁶IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France.

⁶⁷Department of Physics, Indiana University, Bloomington IN; United States of America.

⁶⁸(^a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; (^b) ICTP, Trieste; (^c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy.

⁶⁹(^a) INFN Sezione di Lecce; (^b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy.

⁷⁰(^a) INFN Sezione di Milano; (^b) Dipartimento di Fisica, Università di Milano, Milano; Italy.

⁷¹(^a) INFN Sezione di Napoli; (^b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy.

⁷²(^a) INFN Sezione di Pavia; (^b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy.

⁷³(^a) INFN Sezione di Pisa; (^b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy.

⁷⁴(^a) INFN Sezione di Roma; (^b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy.

⁷⁵(^a) INFN Sezione di Roma Tor Vergata; (^b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy.

⁷⁶(^a) INFN Sezione di Roma Tre; (^b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy.

⁷⁷(^a) INFN-TIFPA; (^b) Università degli Studi di Trento, Trento; Italy.

⁷⁸Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria.

- ⁷⁹University of Iowa, Iowa City IA; United States of America.
- ⁸⁰Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America.
- ⁸¹(^a)Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; (^b)Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; (^c)Instituto de Física, Universidade de São Paulo, São Paulo; (^d)Rio de Janeiro State University, Rio de Janeiro; Brazil.
- ⁸²KEK, High Energy Accelerator Research Organization, Tsukuba; Japan.
- ⁸³Graduate School of Science, Kobe University, Kobe; Japan.
- ⁸⁴(^a)AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; (^b)Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland.
- ⁸⁵Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland.
- ⁸⁶Faculty of Science, Kyoto University, Kyoto; Japan.
- ⁸⁷Kyoto University of Education, Kyoto; Japan.
- ⁸⁸Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan.
- ⁸⁹Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina.
- ⁹⁰Physics Department, Lancaster University, Lancaster; United Kingdom.
- ⁹¹Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom.
- ⁹²Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia.
- ⁹³School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom.
- ⁹⁴Department of Physics, Royal Holloway University of London, Egham; United Kingdom.
- ⁹⁵Department of Physics and Astronomy, University College London, London; United Kingdom.
- ⁹⁶Louisiana Tech University, Ruston LA; United States of America.
- ⁹⁷Fysiska institutionen, Lunds universitet, Lund; Sweden.
- ⁹⁸Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain.
- ⁹⁹Institut für Physik, Universität Mainz, Mainz; Germany.
- ¹⁰⁰School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom.
- ¹⁰¹CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France.
- ¹⁰²Department of Physics, University of Massachusetts, Amherst MA; United States of America.
- ¹⁰³Department of Physics, McGill University, Montreal QC; Canada.
- ¹⁰⁴School of Physics, University of Melbourne, Victoria; Australia.
- ¹⁰⁵Department of Physics, University of Michigan, Ann Arbor MI; United States of America.
- ¹⁰⁶Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ¹⁰⁷Group of Particle Physics, University of Montreal, Montreal QC; Canada.
- ¹⁰⁸Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany.
- ¹⁰⁹Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany.
- ¹¹⁰Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan.
- ¹¹¹Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America.
- ¹¹²Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands.
- ¹¹³Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands.
- ¹¹⁴Department of Physics, Northern Illinois University, DeKalb IL; United States of America.
- ¹¹⁵(^a)New York University Abu Dhabi, Abu Dhabi; (^b)University of Sharjah, Sharjah; United Arab Emirates.

- ¹¹⁶Department of Physics, New York University, New York NY; United States of America.
- ¹¹⁷Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan.
- ¹¹⁸Ohio State University, Columbus OH; United States of America.
- ¹¹⁹Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America.
- ¹²⁰Department of Physics, Oklahoma State University, Stillwater OK; United States of America.
- ¹²¹Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic.
- ¹²²Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America.
- ¹²³Graduate School of Science, Osaka University, Osaka; Japan.
- ¹²⁴Department of Physics, University of Oslo, Oslo; Norway.
- ¹²⁵Department of Physics, Oxford University, Oxford; United Kingdom.
- ¹²⁶LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France.
- ¹²⁷Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America.
- ¹²⁸Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America.
- ¹²⁹^(a)Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; ^(b)Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; ^(c)Departamento de Física, Universidade de Coimbra, Coimbra; ^(d)Centro de Física Nuclear da Universidade de Lisboa, Lisboa; ^(e)Departamento de Física, Universidade do Minho, Braga; ^(f)Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); ^(g)Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal.
- ¹³⁰Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic.
- ¹³¹Czech Technical University in Prague, Prague; Czech Republic.
- ¹³²Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic.
- ¹³³Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom.
- ¹³⁴IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France.
- ¹³⁵Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America.
- ¹³⁶^(a)Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b)Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; ^(c)Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; ^(d)Universidad Andres Bello, Department of Physics, Santiago; ^(e)Instituto de Alta Investigación, Universidad de Tarapacá, Arica; ^(f)Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile.
- ¹³⁷Department of Physics, University of Washington, Seattle WA; United States of America.
- ¹³⁸Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom.
- ¹³⁹Department of Physics, Shinshu University, Nagano; Japan.
- ¹⁴⁰Department Physik, Universität Siegen, Siegen; Germany.
- ¹⁴¹Department of Physics, Simon Fraser University, Burnaby BC; Canada.
- ¹⁴²SLAC National Accelerator Laboratory, Stanford CA; United States of America.
- ¹⁴³Department of Physics, Royal Institute of Technology, Stockholm; Sweden.
- ¹⁴⁴Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America.
- ¹⁴⁵Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom.
- ¹⁴⁶School of Physics, University of Sydney, Sydney; Australia.
- ¹⁴⁷Institute of Physics, Academia Sinica, Taipei; Taiwan.
- ¹⁴⁸^(a)E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; ^(b)High

- Energy Physics Institute, Tbilisi State University, Tbilisi;^(c) University of Georgia, Tbilisi; Georgia.
- ¹⁴⁹Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel.
- ¹⁵⁰Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel.
- ¹⁵¹Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece.
- ¹⁵²International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan.
- ¹⁵³Department of Physics, Tokyo Institute of Technology, Tokyo; Japan.
- ¹⁵⁴Department of Physics, University of Toronto, Toronto ON; Canada.
- ¹⁵⁵^(a)TRIUMF, Vancouver BC;^(b)Department of Physics and Astronomy, York University, Toronto ON; Canada.
- ¹⁵⁶Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan.
- ¹⁵⁷Department of Physics and Astronomy, Tufts University, Medford MA; United States of America.
- ¹⁵⁸United Arab Emirates University, Al Ain; United Arab Emirates.
- ¹⁵⁹Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America.
- ¹⁶⁰Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden.
- ¹⁶¹Department of Physics, University of Illinois, Urbana IL; United States of America.
- ¹⁶²Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain.
- ¹⁶³Department of Physics, University of British Columbia, Vancouver BC; Canada.
- ¹⁶⁴Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada.
- ¹⁶⁵Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany.
- ¹⁶⁶Department of Physics, University of Warwick, Coventry; United Kingdom.
- ¹⁶⁷Waseda University, Tokyo; Japan.
- ¹⁶⁸Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel.
- ¹⁶⁹Department of Physics, University of Wisconsin, Madison WI; United States of America.
- ¹⁷⁰Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany.
- ¹⁷¹Department of Physics, Yale University, New Haven CT; United States of America.
- ^a Also Affiliated with an institute covered by a cooperation agreement with CERN.
- ^b Also at An-Najah National University, Nablus; Palestine.
- ^c Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- ^d Also at Bruno Kessler Foundation, Trento; Italy.
- ^e Also at Center for High Energy Physics, Peking University; China.
- ^f Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki ; Greece.
- ^g Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- ^h Also at CERN, Geneva; Switzerland.
- ⁱ Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ^j Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- ^k Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.
- ^l Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ^m Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America.
- ⁿ Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel.
- ^o Also at Department of Physics, California State University, East Bay; United States of America.

- p* Also at Department of Physics, California State University, Sacramento; United States of America.
- q* Also at Department of Physics, King's College London, London; United Kingdom.
- r* Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- s* Also at Department of Physics, University of Thessaly; Greece.
- t* Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- u* Also at Hellenic Open University, Patras; Greece.
- v* Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- w* Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- x* Also at Institute of Particle Physics (IPP); Canada.
- y* Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia.
- z* Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- aa* Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- ab* Also at Lawrence Livermore National Laboratory, Livermore; United States of America.
- ac* Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen; Germany.
- ad* Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- ae* Also at TRIUMF, Vancouver BC; Canada.
- af* Also at Università di Napoli Parthenope, Napoli; Italy.
- ag* Also at University of Chinese Academy of Sciences (UCAS), Beijing; China.
- ah* Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- ai* Also at Washington College, Maryland; United States of America.
- aj* Also at Yeditepe University, Physics Department, Istanbul; Türkiye.
- * Deceased