

High-energy limit of $2 \rightarrow 2$ scattering amplitudes at NNLL

Calum Milloy,^{a,*} Giulio Falcioni,^b Einan Gardi,^b Niamh Maher^b and Leonardo Vernazza^{a,c}

^a*Dipartimento di Fisica and Arnold-Regge Center,*

Università di Torino, and INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy

^b*Higgs Centre for Theoretical Physics, School of Physics and Astronomy,*

The University of Edinburgh, Edinburgh EH9 3FD, Scotland, UK

^c*Theoretical Physics Department, CERN, Geneva 1211, Switzerland*

E-mail: calumwilliam.milloy@unito.it

The high-energy limit of $2 \rightarrow 2$ scattering amplitudes offers an excellent setting to explore the universal features of gauge theories. At Leading Logarithmic (LL) accuracy the partonic amplitude is governed by Regge poles in the complex angular momentum plane. Beyond LL, Regge cuts in this plane begin to play an important role. Specifically, the real part of the amplitude at Next-to-Next-to-Leading Logarithmic (NNLL) accuracy presents for the first time both a Regge pole and a Regge cut. Analysing this tower of logarithms and computing it explicitly through four loops we are able to systematically separate between the Regge pole and the Regge cut. The former involves two fundamental parameters, namely the gluon Regge trajectory and impact factors. We explain how to consistently define the impact factors at two loops and the Regge trajectory at three loops. We confirm that the singularities of the trajectory are given by the cusp anomalous dimension. We also show that the Regge-cut contribution at four loop is nonplanar.

*Loops and Legs in Quantum Field Theory - LL2022,
25-30 April, 2022
Ettal, Germany*

*Speaker

1. Introduction to the high-energy limit of $2 \rightarrow 2$ scattering amplitudes

The high-energy limit has long been an interesting laboratory to study gauge-theory amplitudes, which simplify significantly in this regime. The simplification in the limit means that multi-loop corrections are accessible, and they may even be resummed to all orders (at a fixed logarithmic accuracy), while still retaining a rich structure in both colour and kinematics.




Figure 1: A representation of the $2 \rightarrow 2$ scattering of eq. (1).

We consider the $2 \rightarrow 2$ scattering of massless partons in a generic gauge theory

$$i(p_1, a_1, \lambda_1) + j(p_2, a_2, \lambda_2) \rightarrow j(p_3, a_3, \lambda_3) + i(p_4, a_4, \lambda_4), \quad (1)$$

where momenta, colour and helicity are labelled by p_i , a_i and λ_i respectively. A pictorial representation is given in Figure 1. The process is described in terms of the Mandelstam invariants,

$$s = (p_1 + p_2)^2 > 0 \quad t = (p_1 - p_4)^2 < 0 \quad u = (p_1 - p_3)^2 < 0, \quad (2)$$

where the signs dictate the physical scattering region. Momentum conservation implies

$$s + t + u = 0. \quad (3)$$

In $2 \rightarrow 2$ scattering, the high-energy (Regge) limit is defined to be when the centre-of-mass energy is much larger than the momentum transfer, i.e. $s \gg -t$. At leading power the tree-level amplitude is

$$\mathcal{M}_{ij \rightarrow ij}^{\text{tree}} = 4\pi\alpha_s \frac{2s}{t} (\mathbf{T}_i^b)_{a_1 a_4} (\mathbf{T}_j^b)_{a_2 a_3} \delta_{\lambda_1 \lambda_4} \delta_{\lambda_2 \lambda_3}, \quad (4)$$

where we observe that the helicity of each of the scattered partons is conserved. We use colour operator notation to keep expressions general for any representation. In the Regge limit it is convenient to define colour-flow operators following the conventions of the Mandelstams [1–3],

$$\mathbf{T}_s = \mathbf{T}_1 + \mathbf{T}_2, \quad \mathbf{T}_t = \mathbf{T}_1 + \mathbf{T}_4, \quad \mathbf{T}_u = \mathbf{T}_1 + \mathbf{T}_3. \quad (5)$$

The quantity \mathbf{T}_t^2 measures the colour charge of the t -channel adjoint exchange,

$$\mathbf{T}_t^2 \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} = C_A \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}. \quad (6)$$

In the high-energy limit it can be shown that in the complex angular momentum plane, the amplitude can be written as a sum over poles and cuts, dubbed *Regge poles* and *Regge cuts*. Regge cuts were shown by Mandelstam to arise from only nonplanar diagrams [4]. Note that Mandelstam's

observation only involved the *diagrams* being nonplanar. In this work we connect this to gauge theories, where planarity relates to colour and corresponds to the large- N_c limit.

Calculating the amplitude in the perturbative expansion in α_s we would find that large logarithms will develop in the ratio $s/(-t)$. In the Leading Logarithmic (LL) approximation, these can be resummed by the famous Regge pole,

$$\mathcal{M}_{ij \rightarrow ij}^{\text{LL}} = \left(\frac{s}{-t} \right)^{C_A \alpha_g(t, \mu^2)} \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad (7)$$

where α_g is called the Regge trajectory. This quantity admits a perturbative expansion,

$$\alpha_g(t, \mu^2) = \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^n \alpha_g^{(n)}(t, \mu^2), \quad (8)$$

where the one-loop coefficient relevant at LL is

$$\alpha_g^{(1)}(t, \mu^2) = \frac{r_\Gamma}{2\epsilon} \left(\frac{\mu^2}{-t} \right)^\epsilon, \quad r_\Gamma = e^{\epsilon \gamma_E} \frac{\Gamma^2(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(1-2\epsilon)}, \quad (9)$$

with $\epsilon = (4-d)/2$ the dimensional regularisation parameter. We will set the renormalisation scale $\mu^2 = -t$ for simplicity. At this accuracy the amplitude has the same colour structure as the tree-level, the octet exchange in the t -channel.

Before moving on to Next-to-Leading Logarithms (NLL), we shall first introduce the concept of signature under $s \leftrightarrow u$, where we notice that the tree-level amplitude in eq. (4) is signature-odd. We define even and odd amplitudes through

$$\mathcal{M}^{(\pm)}(s, t) = \frac{1}{2} (\mathcal{M}(s, t) \pm \mathcal{M}(-s - t, t)). \quad (10)$$

Upon defining the signature-even logarithm,

$$L \equiv \log \left(\frac{s}{-t} \right) - \frac{i\pi}{2} = \frac{1}{2} \left[\log \left(\frac{-s - i0}{-t} \right) + \log \left(\frac{-u - i0}{-t} \right) \right], \quad (11)$$

and then expanding the amplitude in L , the coefficients of L in the odd (even) amplitudes are entirely real (imaginary) [5].

In this work, we focus solely on the odd amplitude $\mathcal{M}^{(-)}$, and we will remove the superscript $(-)$ for convenience. For even amplitudes at NLL see [6–8]. At NLL, the amplitudes obey Regge pole factorisation [9–11],

$$\mathcal{M}_{ij \rightarrow ij}^{\text{NLL}} = e^{C_A \alpha_g(t) L} C_i(t) C_j(t) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad (12)$$

where $C_i(t)$ are energy-independent impact factors for each of the scattered partons. They are expanded in powers of α_s , similar to α_g in eq. (8). The dependence on s is entirely controlled by the Regge trajectory. NLL accuracy is obtained by taking the Regge trajectory up to two loops, and the impact factors at one loop.

It is worthwhile to compare Regge factorisation to infrared factorisation. It is well-known that long-distance singularities factorise from the amplitude [12–17],

$$\mathcal{M}_{ij \rightarrow ij} = \mathbf{P} \exp \left\{ -\frac{1}{2} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \Gamma_{ij \rightarrow ij} \right\} \cdot \mathcal{H} \quad (13)$$

where $\Gamma_{ij \rightarrow ij}$ is dubbed the soft anomalous dimension. In the high-energy limit it is given by [5]

$$\Gamma_{ij \rightarrow ij} = \frac{1}{2} \gamma_K [L \mathbf{T}_t^2 + i\pi \mathbf{T}_{s-u}^2] + \Gamma_i + \Gamma_j + \Delta \quad (14)$$

where at LL, divergences are captured by the lightlike cusp anomalous dimension γ_K and Γ_i is the collinear anomalous dimension, which collects collinear divergences associated to scattered parton i . The operator \mathbf{T}_{s-u}^2 is defined as

$$\mathbf{T}_{s-u}^2 = \frac{1}{2} (\mathbf{T}_s^2 - \mathbf{T}_u^2). \quad (15)$$

The quantity Δ starts at three loops [18] but only contributes at NLL for even amplitudes from four loops [6] and NNLL for odd amplitudes also from four loops [19]. Performing the integral over the d -dimensional running coupling in eq. (13), reveals the explicit infrared poles of the amplitude.

It is natural to compare the two exponentiations in eqs. (12) and (13). The coefficient of the logarithm L in the exponent gives

$$\alpha_g(t) \stackrel{?}{=} -\frac{1}{4} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \gamma_K(\alpha_s(\lambda^2)) + \mathcal{O}(\epsilon^0). \quad (16)$$

The equivalency holds at two loops (i.e. NLL) [20]. We shall show that it also holds at three loops through a proper definition of the trajectory and it remains a conjecture that it holds beyond. The factor independent of the logarithm suggests that the infrared divergences of the impact factors are related to the collinear anomalous dimension Γ_i [21, 22].

2. Distinguishing pole and cut contributions

At NNLL the factorisation in eq. (12) fails. The colour structure of the amplitude is no longer given only by the exchange of an octet in the t -channel, as for the tree-level amplitude, and new colour channels open up. To describe also these contributions to the amplitudes, we generalise eq. (12) by introducing a non-factorising term, which originates from Regge cuts [5, 19, 21–29]

$$\mathcal{M}_{ij \rightarrow ij}^{\text{NNLL}} = e^{C_A \alpha_g(t) L} C_i(t) C_j(t) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} + \mathcal{M}_{ij \rightarrow ij}^{\text{NF}}. \quad (17)$$

The high-energy behaviour of the amplitude will only be uncovered when the perturbative series is resummed. As such, it is not clear how to properly define the Regge pole and Regge cut in order-by-order computations. To see this explicitly, we expand eq. (17) to two loops

$$\mathcal{M}_{ij \rightarrow ij}^{(2), \text{NNLL}} = \left(C_i^{(1)} C_j^{(1)} + C_i^{(2)} + C_j^{(2)} \right) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}} + \mathcal{M}_{ij \rightarrow ij}^{(2), \text{NF}}. \quad (18)$$

In the expansion above, we are a priori free to move any term whose colour factor is proportional to the tree-level from $\mathcal{M}_{ij \rightarrow ij}^{(2), \text{NF}}$ into the two-loop impact factors $C_i^{(2)}$ (the one-loop $C_i^{(1)}$ are fixed at NLL). There is an ambiguity in the definition of the two-loop impact factors. Therefore, we need a prescription to disentangle the two-loop impact factors from the contribution of Regge cuts. How to properly distinguish the pole and cut contributions is the key result in what follows [30].

We can access this non-factorising term by computing certain diagrams in the high-energy limit. In this regime, the fundamental degrees of freedom are Reggeized gluons, dubbed *Reggeons*,




Figure 2: Example MR diagrams. The black lines represent the external scattering partons and the double red lines are the Reggeon exchanges. There is a symmetric sum over all possible orderings of the Reggeon attachments to the external partons.

see [6, 28] and references therein. At tree-level, the diagram featuring the exchange of a single Reggeon in the t -channel has the analytic structure of a Regge pole, i.e. the first term in eq. (17).

We use the shockwave formalism of ref. [6] to compute diagrams with the exchange of Multiple Reggeons (MR), which start at two loops¹. Example diagrams at two, three and four loops are given in Figure 2. The two-loop diagram displayed in Figure 2a was computed in [5], giving

$$\mathcal{M}_{ij \rightarrow ij}^{(2),\text{MR}} = \pi^2 \left(-\frac{1}{8\epsilon^2} + \frac{1}{8}\zeta_2 + \frac{4}{3}\epsilon\zeta_3 + \mathcal{O}(\epsilon^2) \right) \left((\mathbf{T}_{s-u}^2)^2 - \frac{1}{12}C_A^2 \right) \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}. \quad (19)$$

A couple of comments are in order. The overall $\pi^2 = -(i\pi)^2$ highly suggests it originates from a cut. It has uniform transcendental weight as this quantity is independent on the underlying gauge theory: the $\mathcal{N} = 4$ super Yang-Mills (SYM) result is the same as QCD.

These features are maintained in the MR exchanges at three and four loops, which are computed in [5] and [19], respectively. It is possible to identify the entire contribution of MR diagrams with the Regge-cut term in eq. (17), as done in [5]. This defines a scheme to extract the two-loop impact factors $C_i^{(2)}$ and the three-loop Regge trajectory $\alpha_g^{(3)}$ by matching eq. (17), where $\mathcal{M}_{ij \rightarrow ij}^{\text{NF}}$ is identified with $\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}$, with fixed-order calculations [31–35].

Here, we take a different point of view following refs. [21, 30]. Indeed, according to the analysis of Mandelstam [4], the Regge cut contribution $\mathcal{M}_{ij \rightarrow ij}^{\text{NF}}$ arises from nonplanar diagrams. Therefore, its colour structure should be subleading in the limit of large N_c . In contrast with this conclusion, the $\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}$ has non-vanishing contributions at two and three loops for large N_c [19, 21],

$$\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}|_{\text{planar}} = \frac{\pi^2 r_\Gamma^2 N_c^2}{6} \left(\frac{\alpha_s}{\pi} \right)^2 \left\{ S^{(2)}(\epsilon) - \left(\frac{\alpha_s}{\pi} \right) r_\Gamma N_c L \left[S^{(3)}(\epsilon) - \frac{1}{2\epsilon} S^{(2)}(\epsilon) \right] + \mathcal{O}(\alpha_s^3) \right\} \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad (20)$$

$$S^{(2)}(\epsilon) = -\frac{1}{8\epsilon^2} + \frac{3}{4}\epsilon\zeta_3 + \frac{9}{8}\epsilon^2\zeta_4 + \mathcal{O}(\epsilon^3), \quad (21)$$

$$S^{(3)}(\epsilon) = -\frac{1}{144\epsilon^3} + \frac{35}{72}\zeta_3 + \frac{35}{48}\epsilon\zeta_4 + \mathcal{O}(\epsilon^3). \quad (22)$$

¹At one-loop, it is possible to exchange only two Reggeons, but these contribute only to the imaginary (even) part of the amplitude.

By looking at eq. (20), we notice that, for every scattering process, the planar part of $\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}$ is colour proportional to the tree-level amplitude. Because of this property, by following the argument below eq. (18), at two loops it is possible to absorb the planar part of $\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}$ in the definition of the impact factors [21, 30]. At three loops, the planar terms in $\mathcal{M}_{ij \rightarrow ij}^{\text{MR}}$ are again proportional to the tree-level amplitude and can be absorbed in the definition of the three-loop gluon Regge trajectory. This naturally gives a method to capture the true separation between the pole and the cut, where we shift all the planar corrections into a new pole term with new impact factors and a new Regge trajectory [30]

$$\mathcal{M}_{ij \rightarrow ij}^{\text{pole}} = \tilde{C}_i(t) \tilde{C}_j(t) e^{\tilde{\alpha}_g(t) C_{AL}} \mathcal{M}_{ij \rightarrow ij}^{\text{tree}}, \quad (23)$$

where we have added a tilde on the parameters to distinguish between the old and new pole terms. As the whole amplitude is a well-defined quantity, the shifts in the two-loop impact factors and three-loop Regge trajectory can be calculated

$$\tilde{C}_{i/j}^{(2)} = C_{i/j}^{(2)} + N_c^2 (r_\Gamma)^2 \frac{\pi^2}{12} S^{(2)}(\epsilon), \quad (24)$$

$$\tilde{\alpha}_g^{(3)} = \alpha_g^{(3)} - N_c^2 (r_\Gamma)^3 \frac{\pi^2}{6} S^{(3)}(\epsilon). \quad (25)$$

This exhausts all possible parameters at NNLL to absorb planar contributions into the pole term. This is consistent with the fact that at four loops the MR term is entirely nonplanar [19, 21] and that it is conjectured to be so at all higher loops [21, 30].

3. Applications

Having properly distinguished pole and cut contributions order-by-order in the perturbation expansion, we can now extract Regge parameters. Using state-of-the-art calculations for the relevant amplitudes, we first extract two-loop impact factors and then the three-loop Regge trajectory. Lastly, as a by-product of the four-loop computation of $\mathcal{M}_{ij \rightarrow ij}^{(4), \text{MR}}$ we derive constraints on the infrared structure of four-loop amplitudes.

3.1 Two-loop impact factors

Using eq. (24) we can extract the two-loop impact factors. To display results it is more convenient to define the pole term as a symmetric sum of exponentials, rather than a single exponentiated sum. In eq. (23) we make the replacement

$$\tilde{C}_i(t) \tilde{C}_j(t) e^{\tilde{\alpha}_g(t) C_{AL}} \rightarrow Z_i(t) \bar{D}_i(t) \left[\left(\frac{s}{-t} \right)^{\tilde{\alpha}_g(t)} + \left(\frac{-s}{-t} \right)^{\tilde{\alpha}_g(t)} \right] Z_j(t) \bar{D}_j(t), \quad (26)$$

where $Z_i(t)$ is the exponential of the integral over the collinear anomalous dimension Γ_i

$$Z_i(t) \equiv \exp \left\{ -\frac{1}{2} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \Gamma_i \right\}. \quad (27)$$

This replacement defines a finite collinear-subtracted impact factor $\bar{D}_i(t)$, i.e. the poles of the impact factor are given by the collinear anomalous dimension [22]. It is related to the $\tilde{C}_i(t)$ by

$$\tilde{C}_i(t) = Z_i(t)\bar{D}_i(t)\sqrt{\cos\left(\frac{\pi C_A \tilde{\alpha}_g(t)}{2}\right)}. \quad (28)$$

Upon matching to amplitudes calculated in [31] we find the results

$$\begin{aligned} \bar{D}_g^{(2)} &= C_A^2 \left(\frac{335}{288} \zeta_2 + \frac{11}{18} \zeta_3 - \frac{3}{32} \zeta_4 - \frac{26675}{10368} \right) + C_A n_f \left(\frac{49}{108} - \frac{25}{144} \zeta_2 + \frac{5}{36} \zeta_3 \right) \\ &\quad + C_F n_f \left(\frac{55}{192} - \frac{\zeta_3}{4} \right) - \frac{25}{2592} n_f^2 + \mathcal{O}(\epsilon), \end{aligned} \quad (29)$$

$$\begin{aligned} \bar{D}_q^{(2)} &= C_A^2 \left(\frac{73\zeta_2}{32} - \frac{43\zeta_3}{48} - \frac{53\zeta_4}{64} + \frac{13195}{3456} \right) + C_A C_F \left(-\frac{5\zeta_2}{2} + \frac{475\zeta_3}{144} + \frac{65\zeta_4}{32} - \frac{78229}{10368} \right) \\ &\quad + C_F^2 \left(\frac{21\zeta_2}{16} - \frac{15\zeta_3}{8} - \frac{83\zeta_4}{64} + \frac{255}{128} \right) + C_A n_f \left(-\frac{5\zeta_2}{16} - \frac{7\zeta_3}{24} - \frac{385}{432} \right), \\ &\quad + C_F n_f \left(\frac{\zeta_2}{8} + \frac{19\zeta_3}{72} + \frac{505}{648} \right) + \frac{25}{864} n_f^2 + \mathcal{O}(\epsilon) \end{aligned} \quad (30)$$

for gluons and quarks respectively. The values through to $\mathcal{O}(\epsilon^2)$ are given in ref. [30]. Upon taking the planar limit we find that $\bar{D}_g^{(2)}$ agrees with the recent calculation of ref. [36]. Upon extracting the terms of highest weight of eq. (29), we find the gluon impact factor in planar $\mathcal{N} = 4$ SYM [37, 38], which also coincides with the impact factor in full colour [5, 32] when using the cut scheme.

3.2 Three-loop Regge trajectory

We can also extract the Regge trajectory. We use eq. (25) and the state-of-the-art three-loop four quark amplitudes of ref. [33]. In doing so, we find the poles are given by the cusp anomalous dimension, thus confirming eq. (16) at three loops. If we had used the MR scheme, we would find that there are infrared poles not described by γ_K [5].

Upon defining the cusp-subtracted Regge trajectory $\hat{\alpha}_g$ through

$$\hat{\alpha}_g \equiv \tilde{\alpha}_g + \frac{1}{4} \int_0^{\mu^2} \frac{d\lambda^2}{\lambda^2} \gamma_K(\alpha_s(\lambda^2)) \quad (31)$$

we find the finite result at three loops [30, 34]

$$\begin{aligned} \hat{\alpha}_g^{(3)} &= C_A^2 \left(\frac{297029}{93312} - \frac{799\zeta_2}{1296} - \frac{833\zeta_3}{216} - \frac{77\zeta_4}{192} + \frac{5}{24} \zeta_2 \zeta_3 + \frac{\zeta_5}{4} \right) \\ &\quad + C_A n_f \left(\frac{103\zeta_2}{1296} + \frac{139\zeta_3}{144} - \frac{5\zeta_4}{96} - \frac{31313}{46656} \right) + C_F n_f \left(\frac{19\zeta_3}{72} + \frac{\zeta_4}{8} - \frac{1711}{3456} \right) \\ &\quad + n_f^2 \left(\frac{29}{1458} - \frac{2\zeta_3}{27} \right) + \mathcal{O}(\epsilon). \end{aligned} \quad (32)$$

Notice that when $n_f = 0$ there are no N_c -subleading terms, in agreement with the expectation that the Regge trajectory is maximally non-Abelian [20]. The result of eq. (32) agrees with the planar extraction given in [36].

3.3 Infrared Constraints

Another application of the calculation of the MR term is that we can find constraints on the infrared structure of gauge-theory amplitudes. The structure of infrared divergences for multi-leg massless scattering is known at three loops in general kinematics [39]. In addition, in the high-energy limit the entire NLL tower of corrections has been resummed [7, 18]. Given that the four-loop computation of the Regge-cut term is known, we can match the Regge decomposition in eq. (17) to the soft anomalous dimension in the high-energy limit, eq. (14). It can be shown that only the Regge-cut term contributes to Δ . Expanding Δ as

$$\Delta = \sum_{m=3}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^m \sum_{n=0}^{m-1} L^n \Delta^{(m,n)}, \quad (33)$$

we have the four-loop results at NLL [6] and NNLL [19] respectively,

$$\Delta^{(4,3)} = -i\pi \frac{\zeta_3}{24} [\mathbf{T}_t^2, [\mathbf{T}_t^2, \mathbf{T}_{s-u}^2]] \mathbf{T}_t^2, \quad (34)$$

$$\text{Re} \left[\Delta^{(4,2)} \right] = \zeta_2 \zeta_3, \left\{ \frac{1}{4} \mathbf{T}_t^2 [\mathbf{T}_t^2, (\mathbf{T}_{s-u})^2] + \frac{3}{4} [\mathbf{T}_{s-u}^2, \mathbf{T}_t^2] \mathbf{T}_t^2 \mathbf{T}_{s-u}^2 + \frac{d_{AA}}{N_A} - \frac{C_A^4}{24} \right\}, \quad (35)$$

where only the real part of $\Delta^{(4,2)}$ is known as we only have NNLL results for the odd amplitude. By matching the above to an ansatz for the four-loop $\Delta^{(4)}$ in general kinematics [40] we can find the asymptotic limits of the functions [21].

The three-loop soft anomalous dimension was shown to be bootstrapable from general considerations and matching to known collinear and high-energy limits [41]. The above results gives useful constraints for a potential four-loop bootstrap.

4. Conclusions

Beyond NLL, in the high-energy limit, amplitudes no longer factorise in terms of a Regge pole, as given in eq. (12). A non-factorising term, originating from Regge cuts, needs to be included as in eq. (17). This leads to an ambiguity in how to properly define the pole term order-by-order in perturbation theory. In this work, we have shown how to consistently do so with the high-energy properties of Regge poles and Regge cuts in the complex angular momentum plane. We remove all planar MR contributions from the non-factorising term and define a new pole term with new NNLL parameters, namely the two-loop impact factor $\tilde{C}_i^{(2)}$ and the three-loop Regge trajectory $\tilde{\alpha}_g^{(3)}$.

As an application, we find $\mathcal{O}(\epsilon^2)$ terms of the two-loop impact factors of eqs. (29) and (30) using state-of-the-art amplitude computations [31]. We also extract, for the first time, the three-loop Regge trajectory, eq. (32), using the amplitudes of ref. [33]. The poles of the Regge trajectory are given by the integrated cusp anomalous dimension when using the cut scheme, eq. (31).

Furthermore, as a by-product of four-loop computations [6, 19], we can derive constraints on the structure of infrared divergences in the high-energy limit [21], see eqs. (34) and (35).

References

- [1] Yu. L. Dokshitzer and G. Marchesini. “Soft gluons at large angles in hadron collisions”. In: *JHEP* 01 (2006), p. 007. doi: [10.1088/1126-6708/2006/01/007](https://doi.org/10.1088/1126-6708/2006/01/007). arXiv: [hep-ph/0509078](https://arxiv.org/abs/hep-ph/0509078).
- [2] Vittorio Del Duca et al. “An infrared approach to Reggeization”. In: *Phys. Rev. D* 85 (2012), p. 071104. doi: [10.1103/PhysRevD.85.071104](https://doi.org/10.1103/PhysRevD.85.071104). arXiv: [1108.5947](https://arxiv.org/abs/1108.5947) [[hep-ph](#)].
- [3] Vittorio Del Duca et al. “The Infrared structure of gauge theory amplitudes in the high-energy limit”. In: *JHEP* 12 (2011), p. 021. doi: [10.1007/JHEP12\(2011\)021](https://doi.org/10.1007/JHEP12(2011)021). arXiv: [1109.3581](https://arxiv.org/abs/1109.3581) [[hep-ph](#)].
- [4] S. Mandelstam. “Cuts in the Angular Momentum Plane. 2”. In: *Nuovo Cim.* 30 (1963), pp. 1148–1162. doi: [10.1007/BF02828822](https://doi.org/10.1007/BF02828822).
- [5] Simon Caron-Huot, Einan Gardi, and Leonardo Vernazza. “Two-parton scattering in the high-energy limit”. In: *JHEP* 06 (2017), p. 016. doi: [10.1007/JHEP06\(2017\)016](https://doi.org/10.1007/JHEP06(2017)016). arXiv: [1701.05241](https://arxiv.org/abs/1701.05241) [[hep-ph](#)].
- [6] Simon Caron-Huot. “When does the gluon reggeize?” In: *JHEP* 05 (2015), p. 093. doi: [10.1007/JHEP05\(2015\)093](https://doi.org/10.1007/JHEP05(2015)093). arXiv: [1309.6521](https://arxiv.org/abs/1309.6521) [[hep-th](#)].
- [7] Simon Caron-Huot et al. “Infrared singularities of QCD scattering amplitudes in the Regge limit to all orders”. In: *JHEP* 03 (2018), p. 098. doi: [10.1007/JHEP03\(2018\)098](https://doi.org/10.1007/JHEP03(2018)098). arXiv: [1711.04850](https://arxiv.org/abs/1711.04850) [[hep-ph](#)].
- [8] Simon Caron-Huot et al. “Two-parton scattering amplitudes in the Regge limit to high loop orders”. In: *JHEP* 08 (2020), p. 116. doi: [10.1007/JHEP08\(2020\)116](https://doi.org/10.1007/JHEP08(2020)116). arXiv: [2006.01267](https://arxiv.org/abs/2006.01267) [[hep-ph](#)].
- [9] V.S. Fadin et al. “Proof of the multi-Regge form of QCD amplitudes with gluon exchanges in the NLA”. In: *Phys. Lett. B* 639 (2006), pp. 74–81. doi: [10.1016/j.physletb.2006.03.031](https://doi.org/10.1016/j.physletb.2006.03.031). arXiv: [hep-ph/0602006](https://arxiv.org/abs/hep-ph/0602006).
- [10] Boris Lazarevich Ioffe, Victor Sergeevich Fadin, and Lev Nikolaevich Lipatov. *Quantum chromodynamics: Perturbative and nonperturbative aspects*. Cambridge Univ. Press, 2010. ISBN: 978-1-107-42475-3, 978-0-521-63148-8, 978-0-511-71744-4. doi: [10.1017/CB09780511711817](https://doi.org/10.1017/CB09780511711817).
- [11] V.S. Fadin, M.G. Kozlov, and A.V. Reznichenko. “Gluon Reggeization in Yang-Mills Theories”. In: *Phys. Rev. D* 92.8 (2015), p. 085044. doi: [10.1103/PhysRevD.92.085044](https://doi.org/10.1103/PhysRevD.92.085044). arXiv: [1507.00823](https://arxiv.org/abs/1507.00823) [[hep-th](#)].
- [12] Stefano Catani. “The Singular behavior of QCD amplitudes at two loop order”. In: *Phys. Lett. B* 427 (1998), pp. 161–171. doi: [10.1016/S0370-2693\(98\)00332-3](https://doi.org/10.1016/S0370-2693(98)00332-3). arXiv: [hep-ph/9802439](https://arxiv.org/abs/hep-ph/9802439).
- [13] George F. Sterman and Maria E. Tejeda-Yeomans. “Multiloop amplitudes and resummation”. In: *Phys. Lett. B* 552 (2003), pp. 48–56. doi: [10.1016/S0370-2693\(02\)03100-3](https://doi.org/10.1016/S0370-2693(02)03100-3). arXiv: [hep-ph/0210130](https://arxiv.org/abs/hep-ph/0210130).

- [14] S.Mert Aybat, Lance J. Dixon, and George F. Sterman. “The Two-loop soft anomalous dimension matrix and resummation at next-to-next-to leading pole”. In: *Phys. Rev. D* 74 (2006), p. 074004. DOI: [10.1103/PhysRevD.74.074004](https://doi.org/10.1103/PhysRevD.74.074004). arXiv: [hep-ph/0607309](https://arxiv.org/abs/hep-ph/0607309).
- [15] S.Mert Aybat, Lance J. Dixon, and George F. Sterman. “The Two-loop anomalous dimension matrix for soft gluon exchange”. In: *Phys. Rev. Lett.* 97 (2006), p. 072001. DOI: [10.1103/PhysRevLett.97.072001](https://doi.org/10.1103/PhysRevLett.97.072001). arXiv: [hep-ph/0606254](https://arxiv.org/abs/hep-ph/0606254).
- [16] Yao Ma. “A Forest Formula to Subtract Infrared Singularities in Amplitudes for Wide-angle Scattering”. In: *JHEP* 05 (2020), p. 012. DOI: [10.1007/JHEP05\(2020\)012](https://doi.org/10.1007/JHEP05(2020)012). arXiv: [1910.11304](https://arxiv.org/abs/1910.11304) [[hep-ph](#)].
- [17] Ilya Feige and Matthew D. Schwartz. “Hard-Soft-Collinear Factorization to All Orders”. In: *Phys. Rev. D* 90.10 (2014), p. 105020. DOI: [10.1103/PhysRevD.90.105020](https://doi.org/10.1103/PhysRevD.90.105020). arXiv: [1403.6472](https://arxiv.org/abs/1403.6472) [[hep-ph](#)].
- [18] Einan Gardi et al. “The High-Energy Limit of 2-to-2 Partonic Scattering Amplitudes”. In: *PoS RADCOR2019* (2019), p. 050. DOI: [10.22323/1.375.0050](https://doi.org/10.22323/1.375.0050). arXiv: [1912.10883](https://arxiv.org/abs/1912.10883) [[hep-ph](#)].
- [19] Giulio Falcioni et al. “Climbing three-Reggeon ladders: four-loop amplitudes in the high-energy limit in full colour”. In: *Phys. Rev. D* 103 (2021), p. L111501. DOI: [10.1103/PhysRevD.103.L111501](https://doi.org/10.1103/PhysRevD.103.L111501). arXiv: [2012.00613](https://arxiv.org/abs/2012.00613) [[hep-ph](#)].
- [20] I. A. Korchemskaya and G. P. Korchemsky. “Evolution equation for gluon Regge trajectory”. In: *Phys. Lett.* B387 (1996), pp. 346–354. DOI: [10.1016/0370-2693\(96\)01016-7](https://doi.org/10.1016/0370-2693(96)01016-7). arXiv: [hep-ph/9607229](https://arxiv.org/abs/hep-ph/9607229) [[hep-ph](#)].
- [21] Giulio Falcioni et al. “Scattering amplitudes in the Regge limit and the soft anomalous dimension through four loops”. In: *JHEP* 03 (2022), p. 053. DOI: [10.1007/JHEP03\(2022\)053](https://doi.org/10.1007/JHEP03(2022)053). arXiv: [2111.10664](https://arxiv.org/abs/2111.10664) [[hep-ph](#)].
- [22] Vittorio Del Duca et al. “Analyzing high-energy factorization beyond next-to-leading logarithmic accuracy”. In: *JHEP* 02 (2015), p. 029. DOI: [10.1007/JHEP02\(2015\)029](https://doi.org/10.1007/JHEP02(2015)029). arXiv: [1409.8330](https://arxiv.org/abs/1409.8330) [[hep-ph](#)].
- [23] Vittorio Del Duca and E.W.Nigel Glover. “The High-energy limit of QCD at two loops”. In: *JHEP* 10 (2001), p. 035. DOI: [10.1088/1126-6708/2001/10/035](https://doi.org/10.1088/1126-6708/2001/10/035). arXiv: [hep-ph/0109028](https://arxiv.org/abs/hep-ph/0109028).
- [24] Vittorio Del Duca et al. “High-energy QCD amplitudes at two loops and beyond”. In: *Phys. Lett. B* 732 (2014), pp. 233–240. DOI: [10.1016/j.physletb.2014.03.033](https://doi.org/10.1016/j.physletb.2014.03.033). arXiv: [1311.0304](https://arxiv.org/abs/1311.0304) [[hep-ph](#)].
- [25] Vittorio Del Duca et al. “Beyond Reggeization for two- and three-loop QCD amplitudes”. In: *PoS RADCOR2013* (2013), p. 046. DOI: [10.22323/1.197.0046](https://doi.org/10.22323/1.197.0046). arXiv: [1312.5098](https://arxiv.org/abs/1312.5098) [[hep-ph](#)].
- [26] V.S. Fadin. “Particularities of the NNLLA BFKL”. In: *AIP Conf. Proc.* 1819.1 (2017). Ed. by Marcella Capua et al., p. 060003. DOI: [10.1063/1.4977159](https://doi.org/10.1063/1.4977159). arXiv: [1612.04481](https://arxiv.org/abs/1612.04481) [[hep-ph](#)].

- [27] V.S. Fadin and L.N. Lipatov. “Reggeon cuts in QCD amplitudes with negative signature”. In: *Eur. Phys. J. C* 78.6 (2018), p. 439. DOI: [10.1140/epjc/s10052-018-5910-1](https://doi.org/10.1140/epjc/s10052-018-5910-1). arXiv: [1712.09805](https://arxiv.org/abs/1712.09805) [hep-ph].
- [28] Victor Fadin. “BFKL – past and future”. In: Dec. 2020. arXiv: [2012.11931](https://arxiv.org/abs/2012.11931) [hep-ph].
- [29] V. S. Fadin. “Three-Reggeon Cuts in QCD Amplitudes”. In: *Phys. Atom. Nucl.* 84.1 (2021), pp. 100–104. DOI: [10.1134/S1063778820060149](https://doi.org/10.1134/S1063778820060149).
- [30] Giulio Falcioni et al. “Disentangling the Regge Cut and Regge Pole in Perturbative QCD”. In: *Phys. Rev. Lett.* 128.13 (2022), p. 132001. DOI: [10.1103/PhysRevLett.128.132001](https://doi.org/10.1103/PhysRevLett.128.132001). arXiv: [2112.11098](https://arxiv.org/abs/2112.11098) [hep-ph].
- [31] Taushif Ahmed, Johannes Henn, and Bernhard Mistlberger. “Four-particle scattering amplitudes in QCD at NNLO to higher orders in the dimensional regulator”. In: *JHEP* 12 (2019), p. 177. DOI: [10.1007/JHEP12\(2019\)177](https://doi.org/10.1007/JHEP12(2019)177). arXiv: [1910.06684](https://arxiv.org/abs/1910.06684) [hep-ph].
- [32] Johannes M. Henn and Bernhard Mistlberger. “Four-Gluon Scattering at Three Loops, Infrared Structure, and the Regge Limit”. In: *Phys. Rev. Lett.* 117.17 (2016), p. 171601. DOI: [10.1103/PhysRevLett.117.171601](https://doi.org/10.1103/PhysRevLett.117.171601). arXiv: [1608.00850](https://arxiv.org/abs/1608.00850) [hep-th].
- [33] Fabrizio Caola et al. “Three-loop helicity amplitudes for four-quark scattering in massless QCD”. In: (July 2021). arXiv: [2108.00055](https://arxiv.org/abs/2108.00055) [hep-ph].
- [34] Fabrizio Caola et al. “Three-Loop Gluon Scattering in QCD and the Gluon Regge Trajectory”. In: *Phys. Rev. Lett.* 128.21 (2022), p. 212001. DOI: [10.1103/PhysRevLett.128.212001](https://doi.org/10.1103/PhysRevLett.128.212001). arXiv: [2112.11097](https://arxiv.org/abs/2112.11097) [hep-ph].
- [35] Fabrizio Caola et al. “Three-loop helicity amplitudes for quark-gluon scattering in QCD”. In: (July 2022). arXiv: [2207.03503](https://arxiv.org/abs/2207.03503) [hep-ph].
- [36] Vittorio Del Duca, Robin Marzucca, and Bram Verbeek. “The gluon Regge trajectory at three loops from planar Yang-Mills theory”. In: *JHEP* 01 (2022), p. 149. DOI: [10.1007/JHEP01\(2022\)149](https://doi.org/10.1007/JHEP01(2022)149). arXiv: [2111.14265](https://arxiv.org/abs/2111.14265) [hep-ph].
- [37] Vittorio Del Duca and E. W. N. Glover. “Testing high-energy factorization beyond the next-to-leading-logarithmic accuracy”. In: *JHEP* 05 (2008), p. 056. DOI: [10.1088/1126-6708/2008/05/056](https://doi.org/10.1088/1126-6708/2008/05/056). arXiv: [0802.4445](https://arxiv.org/abs/0802.4445) [hep-th].
- [38] V. Del Duca, C. Duhr, and E. W. N. Glover. “Iterated amplitudes in the high-energy limit”. In: *JHEP* 12 (2008), p. 097. DOI: [10.1088/1126-6708/2008/12/097](https://doi.org/10.1088/1126-6708/2008/12/097). arXiv: [0809.1822](https://arxiv.org/abs/0809.1822) [hep-th].
- [39] Øyvind Almeliid, Claude Duhr, and Einan Gardi. “Three-loop corrections to the soft anomalous dimension in multileg scattering”. In: *Phys. Rev. Lett.* 117.17 (2016), p. 172002. DOI: [10.1103/PhysRevLett.117.172002](https://doi.org/10.1103/PhysRevLett.117.172002). arXiv: [1507.00047](https://arxiv.org/abs/1507.00047) [hep-ph].
- [40] Thomas Becher and Matthias Neubert. “Infrared singularities of scattering amplitudes and N^3 LL resummation for n -jet processes”. In: *JHEP* 01 (2020), p. 025. DOI: [10.1007/JHEP01\(2020\)025](https://doi.org/10.1007/JHEP01(2020)025). arXiv: [1908.11379](https://arxiv.org/abs/1908.11379) [hep-ph].
- [41] Øyvind Almeliid et al. “Bootstrapping the QCD soft anomalous dimension”. In: *JHEP* 09 (2017), p. 073. DOI: [10.1007/JHEP09\(2017\)073](https://doi.org/10.1007/JHEP09(2017)073). arXiv: [1706.10162](https://arxiv.org/abs/1706.10162) [hep-ph].