

Emergence of Species Scale Black Hole Horizons

José Calderón-Infante¹, Matilda Delgado², Angel M. Uranga²

¹ Theoretical Physics Department, CERN,
CH-1211 Geneva 23, Switzerland

² Instituto de Física Teórica IFT-UAM/CSIC,
C/ Nicolás Cabrera 13-15, Campus de Cantoblanco, 28049 Madrid, Spain
jose.calderon-infante@cern.ch, matilda.delgado@uam.es, angel.uranga@csic.es

Abstract

The scale at which quantum gravity becomes manifest, the species scale Λ_s , has recently been argued to take values parametrically lower than the Planck scale. We use black holes of vanishing horizon area (small black holes) in effective field theories coupled to quantum gravity to shed light on how the three different physical manifestations of the species scale Λ_s relate to each other. (i) Near the small black hole core, a scalar field runs to infinite distance in moduli space, a regime in which the Swampland Distance Conjecture predicts a tower of exponentially light states, which lower Λ_s . (ii) We integrate out modes in the tower and generate via Emergence a set of higher derivative corrections, showing that Λ_s is the scale at which such terms become relevant. (iii) Finally, higher derivative terms modify the black hole solution and grant it a non-zero, species scale sized stretched horizon of radius Λ_s^{-1} , showcasing the species scale as the size of the smallest possible black hole describable in the effective theory.

We present explicit 4d examples of small black holes in 4d $\mathcal{N} = 2$ supergravity, and the 10d example of type IIA D0-branes. The emergence of the species scale horizon for D0-branes requires a non-trivial interplay of different 8-derivative terms in type IIA and M-theory, providing a highly non-trivial check of our unified description of the different phenomena associated to the species scale.

Contents

1	Introduction	2
2	Warmup Example: 4d Small Black Holes	6
2.1	4d $\mathcal{N} = 2$ supergravity, attractors and higher derivative corrections	7
2.2	Small black holes, species scale horizons and the SDC tower	10
3	D0-branes and the Species Scale Horizon	15
3.1	Microscopic derivation of the entropy and of the species scale	17
3.2	Hot D0-branes	19
4	Species Scale Horizon from Emergent Higher Derivative Terms	21
4.1	Entropy function analysis of general R^4 terms	22
4.2	The D0-brane SDC tower and Higher Derivative Emergence	29
4.3	Supersymmetric 10d R^4 terms are not enough	31
4.4	Including F_2 : Species scale D0-brane horizon from M-theory couplings	33
5	Conclusions	38
A	Tensor Structure of R^4 Corrections	41

1 Introduction

Recent activity in the Swampland Program (see [1–4] for reviews) is providing strong support that the cutoff of an effective field theory (EFT) consistent with quantum gravity is not the Planck scale, but potentially a much lower one, the species scale Λ_s [5–9]. This implies that objects which can be reliably described in the EFT have a lower bound in their size of order Λ_s^{-1} . This is a very profound statement since it leads to quantum gravity effects at length scales which can potentially be parametrically larger than the Planck scale. This happens for instance near infinite distance points in moduli spaces, where the (Swampland) Distance Conjecture (SDC) [10] predicts the appearance of an infinite tower of light particles. Additionally, bounds on how fast these towers and the species scale become light have been proposed and tested in [11–14].

A natural setup in which to explore and sharpen these ideas is in black hole physics. In particular, charged extremal black holes with scalar dependent gauge kinetic functions have an attractor mechanism [15–18] that fixes the vevs of such scalars at the horizon. By appropriately choosing the charges of the black hole, one can tune the point of moduli space that is probed by the horizon. A particular choice of charges leads to small black holes, which classically have zero horizon area (i.e. a singularity), and for which scalars run off to infinite distance as one approaches the black hole core (see [19–21] for other uses of small black holes to explore swampland constraints and [20–29] for other spacetime-dependent configurations probing infinite distances in field space). One would then expect that quantum gravity effects smooth out the singularity by generating a stretched horizon of small but finite size. According to our discussion above, this size should be given by the species scale Λ_s^{-1} (hence parametrically larger than the Planck length). Indeed, it was argued in [19] that in order for small black hole solutions not to violate entropy bounds, they should acquire a finite size, of the order of the UV cut-off.

Black holes have been used at multiple occasions in the literature to probe the species scale [30, 31]. And conversely, the species scale itself is often defined as being the size of the smallest possible black hole (SPBH) that one can reliably describe within an EFT description [5, 32]. One of our objectives is to study how quantum gravitational

effects at the species scale come to modify small black hole solutions and ultimately promote them to these non-singular, finite-sized SPBHs, in perfect agreement with the works of [19].

The appearance of stretched horizons for small black holes has long been explored in the context of string theory, e.g. for supersymmetric (and non-supersymmetric) charged black holes in 4d $\mathcal{N} = 4, 2$ supersymmetric compactifications, by the inclusion of higher derivative corrections, resulting in string scale stretched horizons.¹ Actually this corresponds in those examples to precisely the species scale in the infinite distance limit of the corresponding small black hole, in agreement with our discussion above (see [30, 31] for related works).

In the above discussion, the appearance of the species scale associated to the SDC tower seems accidental. We will however argue that there is a direct link, by showing that integrating out the SDC tower produces, in the spirit of Emergence [35–38] (see [39–43] for recent discussions), the higher curvature terms which ultimately lead to the stretched horizon, which is thus naturally set by the species scale. Hence, we provide an explicit microscopic mechanism for the emergence of species scale horizons for this class of small black holes.² Our result moreover matches the approach in [47], where the species scale has been argued to be the scale at which higher curvature terms become relevant.

Hence, our work allows us to bring together three definitions of the species scale: the original definition accounting for the number of species in the SDC tower, the scale defining the size of SPBHs, and the scale of higher curvature corrections. The central idea is that small black hole solutions get resolved as SPBHs at the species scale, by including the effect of higher derivative corrections emerging from integrating out the SDC tower. Given that our main tool is small black holes, our results hold in asymptotic regimes in moduli space; it would be interesting to extend our understanding to test recent proposal involving the species scale in the interior of moduli space [30, 47, 48].

We expect that the emergence of species scale stretched horizons should hold be-

¹For a complementary viewpoint, see [33, 34].

²It would be interesting to explore connections with other proposals to produce stretched horizons parametrically larger than the Planck scale for small black holes, such as the fuzzball proposal (see [44–46] for reviews).

yond 4d systems. In fact, the main point of this work is to explore this question for D0-brane solutions of 10d type IIA. These behave as 10d charged small black holes, of classically zero size, and at whose core the dilaton runs off to infinitely strong coupling. Swampland considerations imply that a species scale horizon should arise. However, despite the fact that systems of N D0-branes have been under scrutiny for decades from diverse perspectives, including supersymmetric quantum mechanics [49, 50], M(atrix) theory [51], holography [52], their combination [53], and MonteCarlo simulations (see [54] for a recent update), the only discussion about D0-brane horizons seems to have appeared in [55].

We will provide several arguments for the existence of this species scale horizon for D0-brane solutions; our arguments include microstate counting, finite temperature considerations and the analysis of higher curvature terms using (a 10d version of) the entropy function. On the other hand, we compute the familiar higher curvature R^4 terms arising from emergence by integrating out the SDC tower of BPS particles (which are D0-branes themselves) and show that it does not suffice to generate the horizon. Hence, the appearance of the species scale horizon implied by Swampland considerations demands the inclusion of further higher derivative corrections. Indeed, we show that one indeed gets a finite size species scale horizon once we include 8-derivative terms involving curvatures and RR 2-form field strength, a computation which we carry out using a lift to 11d. This also nicely dovetails with the fact that the species scale is the 11d Planck scale.

We regard this as impressive evidence of the non-trivial power of the species scale proposal, which provides a rationale for the presence of these extra terms (which are otherwise largely ignored in the literature), and hence probes deep UV properties of M-theory.

One general aspect in our analysis is that the limited knowledge of higher derivative corrections implies that the computations can be carried out including only the leading terms. However, the scales probed, and in particular the species scale, are such that in principle the whole set of higher derivative terms contribute in comparable amounts (hence, the use of *leading* is misleading). The key point however is that the main effect of the correction is to turn the singular behaviour of the two-derivative approximation into a smoothed out behaviour controlled by the species scale, and it suffices to truncate

the infinite set of higher derivative terms to a tractable finite subset which captures the essence of this change, namely the presence of the species scale horizon, with the expectation that the remaining terms modify only order 1 numerical factors, but not the parametric dependence producing the species scale. In fact, in specific setups, such as 4d small black holes, this has been substantiated using scaling and other arguments [56, 57]. We expect this lesson to apply to other setups as well, including the 10d type IIA D0-branes. For this case, we do not have a clear argument why a truncation of the higher derivative terms suffices to capture the existence and parametric dependence of the stretched horizon. It would be interesting to argue for this, for instance by looking for general scaling properties of the higher derivative terms in the type IIA effective action with the dilaton.

The paper is organized as follows. In Section 2 we consider small black holes in 4d $\mathcal{N} = 2$. Section 2.1 reviews the attractor mechanism, including higher curvature corrections, and in section 2.2 we study a class of 4d small black holes, the appearance of a species scale horizon from the higher curvature corrections, and the emergence of the latter from integrating out an SDC tower of KK gravitons of M-theory on \mathbf{S}^1 (times the CY_3). In Section 3 we turn to the case of 10d type IIA D0-branes, regarded as small black holes. In section 3.1 we carry out a microscopic computation of the entropy of the system of N D0-branes and of the appearance of the species scale scaling as $N^{1/2}$. Section 3.2 presents a complementary discussion of the computation of the entropy from the perspective of the finite temperature deformation of the D0-brane supergravity solution.

In Section 4 we study the appearance of the horizon from higher derivative terms in the action. In section 4.1 we perform an entropy function analysis of the spacetime solution, and show that general R^4 corrections can lead to the appearance of a species scale stretched horizon matching the microscopic results. In Section 4.2 we consider the emergence of certain supersymmetric R^4 terms from integrating out the SDC tower in the limit explored by the solution (which is a D0-brane tower, or equivalently M-theory KK gravitons). In section 4.3 we apply the entropy function to these terms and show they still do not lead to the appearance of the horizon. Finally, in section 4.4 we show that including terms involving the RR 2-form field strength one recovers the species scale horizon for the system. We offer some final thoughts and future prospects in

Section 5.

2 Warmup Example: 4d Small Black Holes

In this section, we illustrate our picture of the diverse appearances of the species scale using small black holes in 4d $\mathcal{N} = 2$ EFTs obtained from compactifications of type IIA theory on a Calabi-Yau threefold. As is well known, there are large classes of BPS black hole solutions displaying the attractor mechanism [15]: Vector multiple moduli vary along the radial direction, and, in the $\text{AdS}_2 \times \mathbf{S}^2$ near-horizon geometry, attain values which are fixed in terms of the black hole charges and are independent of the asymptotic value of the moduli (while hypermultiplets remain fixed all along the flow). Small black holes correspond to solutions where the attractor mechanism drives the scalars to infinite distance in field space, leading to a horizon of formally zero size and to a singularity. As has been studied extensively in the literature (see [57, 58] for reviews), higher curvature corrections generically lead to a stretched horizon and a finite entropy, matching microscopic computations in many classes of examples. In section 2.1, we summarize these developments and describe the case of a specific class of small black hole solutions in full detail.

Our new angle is discussed in section 2.2, where we argue that the Swampland Distance Conjecture infinite tower of states becoming light in the infinite distance limit corresponding to small black holes is precisely responsible, via emergence, for the appearance of higher derivative terms producing the stretched horizon. We thus link the appearance of the species scale as derived from the SDC tower, as controlling higher derivative corrections, and as setting the size of the smallest possible black hole in the effective field theory. It is amusing to see that swampland ideas play such a central role in the almost three decade-long success story of microstate explanation of black hole entropy.³

³For other approaches to the Distance Conjecture and Black Holes, see [19–21, 28, 30, 59, 60]. For the interplay of higher curvature corrections and swampland constraints, mainly the Weak Gravity Conjecture [61], see e.g. [62–64]

2.1 4d $\mathcal{N} = 2$ supergravity, attractors and higher derivative corrections

2.1.1 $\mathcal{N} = 2$ supergravity black holes

Type II string theory on a Calabi-Yau threefold gives 4d $\mathcal{N} = 2$ supergravity theories. We will focus on the physics in vector multiplet moduli space, as hypermultiplets are inert. Including the graviphoton, there are $n_V + 1$ gauge bosons, labelled by $I = 0, \dots, n_V$. The structure of the bosonic Lagrangian is

$$\mathcal{L} = R - 2g_{i\bar{j}} \partial_\mu z^i \partial^\mu \bar{z}^{\bar{j}} + \text{Im} \mathcal{N}_{IJ} F_{\mu\nu}^I F^{J\mu\nu} + \text{Re} \mathcal{N}_{IJ} F_{\mu\nu}^I \frac{\epsilon^{\mu\nu\rho\sigma}}{2\sqrt{-g}} F_{\rho\sigma}^J, \quad (2.1)$$

where the different quantities are defined using special geometry (see [65] for reference). The scalars are parametrized by a set of projective coordinates X^I , and the Kähler potential is determined by the prepotential $F(X)$, a holomorphic function of degree 2, as

$$\mathcal{K} = -\log i \left(\bar{X}^I F_I - X^I \bar{F}_I \right) \quad , \quad \text{with } F_I = \partial_I F. \quad (2.2)$$

The gauge kinetic functions \mathcal{N}_{IJ} are also determined by special geometry, but we will skip its detailed structure.

The microscopic description in type IIA theory is as follows. There are $h_{1,1}$ vector multiplets whose gauge bosons arise from the 10d RR 3-form integrated over 2-cycles ω_i , $i = 1, \dots, h_{1,1}$. The Kähler moduli, complexified with the integrals of the NSNS 2-form over the 2-cycles, give rise to the vector moduli; morally, the affine coordinates Z^i . The corresponding $F_i \equiv \partial_i F$ are associated to the dual 4-cycles.

The structure of the prepotential in the large volume regime is

$$F_0(X) = -\frac{1}{6} C_{ijk} \frac{X^i X^j X^k}{X^0}, \quad (2.3)$$

where C_{ijk} are the integer triple intersection numbers of the Calabi-Yau. Away from the large volume limit, the prepotential receives corrections from worldsheet instantons. Also, as discussed later, certain higher derivative corrections can also be encoded in a corrected prepotential.

The theory contains BPS black holes constructed out of D-branes wrapped on holomorphic cycles in the CY. There is a vector of electric and magnetic charges $\Gamma = (q_I, p^I)$. The central charge is

$$\mathcal{Z} = e^{\mathcal{K}/2} (q_I X^I - F_I p^I). \quad (2.4)$$

Regular black holes have a near-horizon $\text{AdS}_2 \times \mathbf{S}^2$ geometry. The values of the moduli at the horizon and the entropy are fixed by the attractor equations

$$\partial_i \mathcal{Z}(X^I, \bar{X}^{\bar{I}}, q_I, p^I)|_h = 0 \quad , \quad S = \pi |\mathcal{Z}(q_I, p^I)|_h \quad , \quad (2.5)$$

where the subindex h indicates the value at the horizon. The solution can be expressed as

$$\begin{aligned} p^I &= \text{Re}[C_h X_h^I] = C_h X_h^I + \bar{C}_h \bar{X}_h^{\bar{I}} \quad , \\ q_I &= \text{Re}[C_h F_{hI}] = C_h F_{hI} + \bar{C}_h \bar{F}_{h\bar{I}} \quad , \end{aligned} \quad (2.6)$$

where we have introduced $C = -2i\bar{\mathcal{Z}}e^{\mathcal{K}/2}$.

In the type IIA setup, the charges q_0 and q_i correspond⁴ to D0-branes, and D2-branes on 2-cycles, while the charges p^0 and p^i correspond to D6-branes on the CY and D4-branes on 4-cycles.

A celebrated class (given its simple lift to M-theory [66]) is obtained by considering sets of D4-branes wrapped p_i times on the i^{th} 4-cycle and a number q_0 of D0-branes, for $|q_0| \gg p^i \gg 1$. Its attractor behaviour can be easily analyzed in the large volume limit (2.3), see [30] for a recent application. The attractor values for moduli, and the entropy, are (we introduce $q \equiv -q_0$)

$$X^0 = -\frac{1}{2} \sqrt{\frac{\frac{1}{6} C_{ijk} p^i p^j p^k}{q}} \quad , \quad X^i = -\frac{i}{2} p^i \quad , \quad S = 2\pi \sqrt{\frac{1}{6} q C_{ijk} p^i p^j p^k} \quad . \quad (2.7)$$

The CY volume modulus at the horizon is thus

$$\mathcal{V}_h = \sqrt{\frac{q^3}{\frac{1}{6} C_{ijk} p^i p^j p^k}} \quad . \quad (2.8)$$

Note that we need $q \gg p^i$ for a reliable large volume expansion. In section 2.2 we exploit this class of models to build small black holes and discuss its corrections.

2.1.2 Higher derivative corrections and quantum attractors

The discussion of higher derivative corrections to the attractor mechanism has been studied extensively in the literature. In particular, there is a class of $\mathcal{N} = 2$ $R^2 F^{2g-2}$

⁴One should take into account induced D-brane charges, e.g. a single D4-brane wrapped on a K3 has -1 units of induced D0-brane charge due to Chern-Simons couplings to background curvature.

F-term corrections which are computed by the topological string, and whose effects on black holes can be included systematically [67–69], see also [58, 70] for reviews. Following these references, one uses the Weyl superfield $W_{\mu\nu} = F_{\mu\nu}^+ - R_{\mu\nu\lambda\rho}^+ \theta\sigma^{\lambda\rho}\theta + \dots$, whose lower component is the (self-dual piece of the) graviphoton, and defines $\Upsilon = W^2$. The F-term corrections can be encoded in a degree 2 generalized prepotential

$$F(X^I, \Upsilon) = \sum_{g=0}^{\infty} F_g(X^I) \Upsilon^g. \quad (2.9)$$

The lowest term F_0 corresponds to the usual prepotential, and higher F_g 's are the genus g topological string amplitude. In the presence of these corrections, the attractor equations and the entropy are modified so we have

$$\begin{aligned} p^I &= i(X_h^I - \bar{X}_h^I) \quad , \quad q_I = i(F_I(X, \Upsilon)_h - \bar{F}_I(\bar{X}, \bar{\Upsilon}_h)) , \\ \Upsilon|_h &= -64 \quad , \quad S = \pi(|\mathcal{Z}_h|^2 + 4\text{Im}(\Upsilon \partial_{\Upsilon} F)_h) . \end{aligned} \quad (2.10)$$

The structure of corrections near the large volume limit is

$$F(X, \Upsilon) = -\frac{1}{6} C_{ijk} \frac{X^i X^j X^k}{X^0} + d_i \frac{X^i \Upsilon}{X^0} , \quad (2.11)$$

where the C_{ijk} are the triple-intersection numbers of the Calabi-Yau and the d_i are given in terms of the second Chern classes of the Calabi-Yau by

$$d_i = -\frac{1}{24} \frac{1}{64} \int_{\mathbf{X}_6} c_2(T\mathbf{X}_6) \wedge \omega_i \quad (2.12)$$

with ω_i a basis of $H^{1,1}(\mathbf{X}_6)$. While these corrections are present for general black holes, for regular ones they are suppressed with respect to the two-derivative result. However, they are crucial for small black holes for which the classical piece vanishes, so that the extra pieces produce a non-zero entropy, namely a stretched horizon. In the following we consider a particular class of small black holes, and show that the leading correction F_1 , which contain 4-derivative R^2 corrections, produce a stretched horizon. In section 2.2 we will show that these corrections precisely follow from the SDC tower and that the horizon size is controlled by the species scale.

We would like to point out that the results can also be derived in the formalism of minimization of the entropy function in an $\text{AdS}_2 \times \mathbf{S}^2$ ansatz near horizon geometry [71].

The added terms in the Lagrangian encoded in the corrected prepotential contribute to a modified entropy function. For small black holes, there is a run away but once the corrections are included, the entropy function can actually be minimized, signaling the presence of a horizon. This approach will be exploited in the 10d setup in section 4.1.

2.2 Small black holes, species scale horizons and the SDC tower

2.2.1 An illustrative class of stretched small black holes

We now consider a specific class of models, based on a 2-modulus version of the large volume expansion (2.3). We take the prepotential

$$F = -\frac{X^1(X^2)^2}{X^0}, \quad (2.13)$$

where we have chosen $C_{122} = 6$ for simplicity. This is a simple template that can model an internal space \mathbf{X}_6 given by $\text{K3} \times \mathbf{T}^2$ (for which extensive studies of small black holes and stretched horizons have been carried out, see [57] for a review), or K3-fibered CY threefolds. The affine coordinates $Z^1 = X^1/X^0$ and $Z^2 = X^2/X^0$ correspond to the sizes of the \mathbf{T}^2 (or the base \mathbf{P}_1 of the K3 fibration) and of the K3 (fiber), respectively.

We can easily build small black holes with a choice of non-zero charges $p \equiv p^1$, $q \equiv -q_0$, and setting $q_i, p^0, p^2 = 0$. The central charge, in terms of $t_i = \text{Im}Z^i$, is

$$\mathcal{Z} = \frac{q + p(t_2)^2}{2\sqrt{2}(t_1)^{1/2}t_2}. \quad (2.14)$$

The attractor equations imply that t_2 is stabilized at the horizon at

$$t_2 = \sqrt{\frac{q}{p}}, \quad (2.15)$$

which is still in the large volume regime if $q \gg p$, as we assume in the following. The attractor mechanism also leads to $t_1 \rightarrow \infty$, so this scalar runs off to infinite distance in moduli space. Consequently the overall volume in string units at the horizon also diverges:

$$e^{-\mathcal{K}_h} |X_h^0|^{-2} = 8\mathcal{V}_h = 8t_1 t_2^2 \rightarrow \infty, \quad (2.16)$$

where the $|X_h^0|^{-2}$ factor accounts for the choice of projective coordinates on the special Kähler manifold such that \mathcal{V} is independent of the Kähler gauge. Using the formulas

(2.5), the area of the horizon and therefore the entropy vanish:

$$S = \lim_{\substack{t_1 \rightarrow \infty \\ t_2 = \sqrt{\frac{q}{p}}} \pi |\mathcal{Z}| \rightarrow 0. \quad (2.17)$$

We now show that the small black hole acquires a stretched horizon upon the inclusion of the correction of the kind (2.11). In particular, we focus on the correction d_1 , so the prepotential is

$$F = -\frac{X^1(X^2)^2}{X^0} + d_1 \frac{X^1 \Upsilon}{X^0}, \quad (2.18)$$

where the last two terms correspond to F_1 , the genus 1 topological string amplitude. This corresponds to the class of solutions considered in [30] with $p^i = 0$, $i \neq 1$ (see also [57] for a related class).

The moduli at the horizon and entropy are given by

$$Z^1 = i \frac{\sqrt{pq}}{2\sqrt{d_1 \Upsilon_h}} \quad , \quad Z^2 = 0 \quad , \quad S \sim 4\pi \sqrt{d_1 \Upsilon_h p q}. \quad (2.19)$$

We see that the horizon no longer explores an infinite distance in moduli space and that the entropy is non-vanishing. Hence the solution develops a stretched horizon that cloaks up the singularity. The overall volume at the stretched horizon is modified by the curvature corrections (see [58] and references therein) and is given by:

$$e^{-\kappa_h} |X_h^0|^{-2} = 8\mathcal{V}_h \sim \frac{q(t_1)_h}{p} \sim \frac{q^{3/2}}{\sqrt{p d_1 \Upsilon_h}}, \quad (2.20)$$

which is finite, and still in the large volume regime if $q \gg p$.

Recall that the above result could have been obtained using the entropy function formalism, but we skip its explicit discussion. Rather we turn to the discussion of the appearance of the higher derivative correction from the SDC tower.

2.2.2 The species scale horizon

In a d -dimensional EFT weakly coupled to Einstein gravity, the species scale is given by:

$$\Lambda_s \sim \frac{M_d}{N_s^{\frac{1}{d-2}}}, \quad (2.21)$$

where N_s is the number of species below the species scale and M_d is the d -dimensional Planck mass. In EFTs obtained from string theory (or compactifications thereof), the number of light species changes as one moves along the moduli space. Thus, the species scale is dependent on the region of moduli-space under consideration. In particular, in infinite distance limits in moduli space where infinite towers of states are becoming exponentially light [10], the species scale falls as the number of light species increases drastically. In order to determine the species scale, we must therefore study what infinite distance limit the (formerly) small black hole was probing.

Recall that the small black holes described in the previous section probes the infinite distance limit $t_1 \rightarrow \infty$, with t_2 fixed. Note that in this case, the overall volume modulus (2.16) $\mathcal{V} \sim t_1(t_2)^2$ also goes to infinity at the horizon, where \mathcal{V} is the volume of the CY \mathbf{X}_6 in string units. At fixed 4d Planck scale $M_p^2 = M_s^2 \mathcal{V}/g_s^2$, the large \mathcal{V} limit corresponds to a large g_s limit in string variables (a similar conclusion follows from the fact that the 4d dilaton lies in a hypermultiplet, so it is constant along the attractor flow, hence \mathcal{V}/g_s^2 remains fixed). Hence, the limit of our interest correspond to a large g_s regime. There is therefore a tower of D0-brane particles becoming light; this also follows from the fact that the central charge for k D0-branes is given by $Z_{D0} = k(t_1)^{-1/2}(t_2)^{-1}$, which goes to zero in the limit. Actually, there may be other towers of particles becoming light, but they correspond to branes wrapped on some internal cycles, so they are suppressed in the regime $q \gg p$. The D0-brane particles thus encode the leading correction. Incidentally, we also note that the infinite distance limit explored by the singular two-derivative solution is a standard decompactification limit to M-theory compactified on \mathbf{X}_6 , rather than an emergent string limit [72, 73]. Indeed, one can show that the volume of the Calabi-Yau in M-theory units is related to the 4d dilaton and is therefore constant, signalling that there is no decompactification to 11d M-theory.

The computation of the species scale in this infinite distance limit can be done in different ways. We choose to focus on the approach of [48], where it was argued that higher derivative corrections to the EFT could be used as a proxy for determining the species scale. In this framework, the species scale acts as the energy scale that suppresses higher derivative corrections to the EFT. This recipe was applied to the case of Calabi-Yau compactifications of type II string theories to find that the species

scales is given by:

$$\Lambda_s = \frac{M_4}{\sqrt{F_1}}, \quad (2.22)$$

where once more, F_1 is the genus 1 topological free energy. Taking (2.19) and $F_1 \sim Z^1 d_1 \Upsilon$, we see that at the horizon of the stretched black hole, we have:

$$S \sim \left(\frac{\Lambda_s}{M_4} \right)^{-2}. \quad (2.23)$$

The radius of the stretched horizon is therefore given exactly by the species scale.

Evaluating the species scale can also be done by counting the number of D0 states that are light in the EFT, in the spirit of [39–43], using (2.21). In the next section we will bridge these two interpretations of the species scale by showing how the higher derivative corrections to the EFT Lagrangian originate from integrating out the tower of D0 states, in the spirit of Emergence.

2.2.3 The SDC tower and Higher Derivative Emergence

The SDC implies that the description of infinite distance limits in moduli space require the inclusion of microscopic physics, in particular an infinite tower of states whose masses decrease exponentially with the field distance [10]. On the other hand, in the previous section we have argued that the singular behaviour of the solution is cloaked by a stretched horizon if suitable higher derivative corrections are included. In this section we show that the two UV ingredients are related, and in fact the distance conjecture *implies* the appearance of higher derivative corrections of the necessary kind to generate the stretched horizon. Hence, the two explanations are two sides of the same phenomenon and link the different physical interpretations of the species scale.

The key idea is that the R^2 terms can be seen as generated by an infinite tower of 4d BPS D0-brane states, namely KK gravitons running on an \mathbf{S}^1 compactification of the 5d theory given by M-theory on \mathbf{X}_6 . In fact, they correspond to the leading term in the Gopakumar-Vafa expansion [74, 75] in the large volume limit (where the contribution from D2-branes is subleading).

Indeed, M-theory on \mathbf{X}_6 contains 5d BPS states which correspond to 11d gravitons on the groundstate of the \mathbf{X}_6 compactification. These are given by the K3 cohomology classes, giving an overall multiplicity of 24, times a linear dependence on the Kähler

modulus of Z^1 . These 5d BPS states can run with KK momentum in the \mathbf{S}^1 compactification to 4d, leading to the 4d tower of D0-brane states.⁵

Hence the problem is just the computation of 1-loop diagram, see [76] for computations in this spirit (see also [77–82] for related computations in other setups). For later convenience, we present the general n graviton scattering amplitude in d dimensions in the worldline formalism:

$$\mathcal{A}_{d,n} = \frac{1}{n! \pi^{d/2}} \sum_k \int d^d \mathbf{p} \int_0^\infty \frac{d\tau}{\tau} \tau^n e^{-\tau(\mathbf{p}^2 + \frac{k^2}{R^2})}. \quad (2.24)$$

Here τ is the worldline parameter, k is the KK momentum, and R is the \mathbf{S}^1 radius.

We want to obtain the R^2 corrections to the effective action, which are encoded in two graviton scattering. For pedagogical reasons, it is easier to start from the 6d perspective of M-theory on $\text{K3} \times \mathbf{S}^1$ and postpone compactifying on the \mathbf{T}^2 . Thus, we particularize to $d = 6$, $n = 2$. The above expression corresponds to the contribution of a single K3 ground state, and diverges in two ways: because of the integral and because of the infinite sum. Since the sum over momentum modes is infinite, we can perform a Poisson resummation, so that both are nicely combined. We get

$$\frac{1}{\pi^{3/2}} \mathcal{A}_{6,2} = \pi^{3/2} \tilde{K} \int_0^\infty d\hat{\tau} \hat{\tau}^{1/2} \sum_l e^{-\pi \hat{\tau} R^2 l^2} = C \tilde{K} + \frac{\zeta(3)}{\pi R^3} \tilde{K}, \quad (2.25)$$

where $\hat{\tau} = \tau^{-1}$. This Poisson summation trades the KK momentum k for the winding number l of the worldline along the \mathbf{S}^1 . The only divergence is now in the $l = 0$ piece, which has been isolated as the first terms in the second equality; C is an unknown coefficient regularizing the divergence. Including the K3 ground state multiplicity of 24 and the \mathbf{T}^2 modulus dependence upon compactification to 4d, we have

$$\frac{1}{\pi^{3/2}} \mathcal{A}_{4,2} = 24 Z^1 \left[C \tilde{K} + \frac{\zeta(3)}{\pi R^3} \tilde{K} \right], \quad (2.26)$$

where \tilde{K} is a 4d kinematical matrix. This is of the form introduced in the prepotential (2.18), hence it is precisely of the form required to produce a stretched horizon. We note that the same 4d result (2.26) could have been obtained via a 1-loop computation

⁵The D0-brane tower is actually analogous to that we will consider in the 10d context in section 4.2, as they are simply related by compactification.

in the 10d theory compactified directly on $K3 \times \mathbf{T}^2$, with the Z^1 prefactor arising from the momentum integral over the \mathbf{T}^2 . Hence, the 4d coupling of interest arises from higher derivative emergence.

This nicely illustrates the links among the appearance of the species scale from the SDC tower, the higher-curvature corrections and the stretching of a small black hole into the SPBH.

We finish with a comment in hindsight. It is a well-known fact that the R^2 corrections encoded in the corrected prepotential can be seen to arise from the compactification of suitable 10d R^4 terms [83]. For concreteness, focusing on the particularly simple case of compactification on $\mathbf{X}_6 = K3 \times \mathbf{T}^2$, we start from the familiar 10d supersymmetric R^4 terms, and put two curvature insertions in K3 to get a 6d R^2 term, with a prefactor of $\chi(K3) = 24$; upon further integration over \mathbf{T}^2 (which provides no background curvature) we obtain the 4d R^2 term with the Z^1 prefactor. It is easy to generalize this argument to general CY_3 compactifications and obtain the corrections⁶ (2.11) with d_i essentially determined by the second Chern classes of \mathbf{X}_6

$$d_i \sim c_{2i} = \int_{\mathbf{X}_6} R \wedge R \wedge \omega_i, \quad (2.27)$$

where ω_i are the Poincaré dual 2-forms of the basis 2-cycles. In fact, the above is the simplest way to derive the 4d R^2 corrections [83] (see [30] for a recent application). The dimensional reduction just described underlies the fact that the 4d R^2 terms can be obtained from integrating out a tower of 5d KK gravitons is related to a similar derivation of 10d R^4 terms from 11d M-theory gravitons, as we describe in section 4.2.

3 D0-branes and the Species Scale Horizon

We now turn to the discussion of the system of N D0-branes in 10d type IIA theory. We will argue for the existence of a stretched horizon controlled by the species scale, and explore its appearance from higher derivative couplings and their interplay with emergence upon integrating out the states in the corresponding SDC tower.

⁶Actually, there is one further constant correction to the prepotential, proportional to the Euler characteristic $\chi(\mathbf{X}_6)$, arising from integrating R^3 over the space \mathbf{X}_6 , and reabsorbing the resulting correction to the 4d Einstein term via a change of frame [83]. This will not be relevant for our discussion.

We start by describing D0-brane systems as small black holes in the theory. The relevant part of the string-frame type IIA action is

$$S = \frac{1}{2\kappa^2} \int d^{10}x \sqrt{-g} \left[e^{-2\phi} (R + 4\partial_\mu \phi \partial^\mu \phi) - \frac{1}{2} |F_2|^2 \right], \quad (3.1)$$

where $2\kappa^2 = (2\pi)^7 \alpha'^4$. The D0-brane solution is

$$\begin{aligned} ds^2 &= f(r)^{-1/2} (-dt^2) + f(r)^{1/2} (dx_\perp^2), \\ e^{(\phi - \phi_\infty)} &= f(r)^{3/4}, \\ f(r) &= 1 + \frac{\rho^7 g_s^\infty N}{r^7}, \quad \text{with } \rho^7 = (4\pi)^{\frac{5}{2}} \alpha'^{\frac{7}{2}} \Gamma\left(\frac{7}{2}\right). \end{aligned} \quad (3.2)$$

We will be more interested in the Einstein frame action

$$S_{\text{IIA}} = \frac{1}{2} \int d^{10}x \sqrt{-g} \left(R - \frac{1}{2} (\partial\phi)^2 - \frac{1}{2} e^{\frac{3}{2}\phi} |F_2|^2 \right), \quad (3.3)$$

where, from now on, we use 10d Planck units. The D0-brane solution reads

$$\begin{aligned} ds^2 &= f(r)^{-7/8} (-dt^2) + f(r)^{1/8} (dx_\perp^2), \\ e^{(\phi - \phi_\infty)} &= f(r)^{3/4}, \\ f(r) &= 1 + \frac{\rho^7 g_s^\infty N}{r^7}, \quad \text{with } \rho^7 = (4\pi)^{\frac{5}{2}} \alpha'^{\frac{7}{2}} \Gamma\left(\frac{7}{2}\right). \end{aligned} \quad (3.4)$$

Although not usually described from this viewpoint, this solution provides an interesting analogy with small black holes in lower dimensional theories. It has an \mathbf{S}^8 horizon of zero size at its core, where a scalar (the dilaton) goes off to infinite distance in moduli space, resulting in a singularity. The latter is not worrisome, because it simply reflects the presence of a source, whose microscopic description is beyond the two-derivative supergravity approximation.

On the other hand, swampland arguments would imply that, as we are driven to the core of the solution, namely near infinite distance limit in the dilaton moduli space, the effective field theory should include additional degrees of freedom, corresponding to an SDC tower of states. Inclusion of these in the effective field theory leads to a lowering of the cutoff scale and hence a minimal size for the system, which would naturally be associated to a species scale horizon. In the following sections we provide several arguments that such a horizon exists and is given by the species scale.

3.1 Microscopic derivation of the entropy and of the species scale

The most direct way to argue that the D0-brane system develops a finite size horizon is to show that it has a large number of microstates, to which one can associate an entropy. This provides a lower bound on the effective size of the system, by holographic bounds.⁷ In this section we perform the microscopic computation of the entropy of a system of D0-branes with total charge N (which is familiar from similar combinatorics problems in string theory), and also explain the microscopic derivation of the species scale (which is a novel result).

Recall that consistency of the description of 10d type IIA as M-theory on \mathbf{S}^1 requires that, for each value of k , there is exactly one threshold bound state in the k D0-brane system (corresponding to the 11d graviton with k units of KK momentum) [84]. Hence, for a total D0-brane charge N , the number of microstates is given by the number of partitions $p(N)$. The asymptotic behaviour for large N is given by the celebrated Hardy-Ramanujan formula

$$p(N) \sim \frac{1}{4\sqrt{3}N} \exp\left(\pi\sqrt{\frac{2}{3}}\sqrt{N}\right). \quad (3.5)$$

The entropy of this thermodynamic ensemble for large charge is thus

$$S \sim \sqrt{N}, \quad (3.6)$$

up to $\log N$ corrections. It is tantalizing to speculate the latter may be related to similar log corrections to the species scale discussed in [85].

The above expression implies that the corresponding horizon size in a gravitational description of the system is parametrically larger than the Planck scale, as we will explain in later sections. In fact, the expectation that it corresponds to the species scale can already be supported from the viewpoint of microscopic combinatorics, as we show next.

One may have expected that, in order for a charge N black hole to be describable in an effective field theory, the latter should include at least N species (i.e. different D0-brane bound states). However, this is not correct and highly overestimates the

⁷Note that in general the Bekenstein-Hawking area law gets corrected when the action includes higher curvature terms; hence, this relation will be made more precise in section 4.1.

number of species. In fact, rephrasing the definition in [48], the number of species N_s should be the minimum necessary to explain the entropy of black hole entropy. In other words,⁸ if we consider a theory with the number of species given by k and want to build a charge N black hole, the number of ways to do so is given by $p_k(N)$, the number of partitions of N with each part not larger than k . This is equivalent to the number of partitions of N into at most k parts.⁹ A classic result in [88] gives the asymptotic behaviour

$$\lim_{N \rightarrow \infty} \frac{p_k(N)}{p(N)} = \exp\left(-\frac{2}{C} e^{-\frac{1}{2}Cx}\right) \quad (3.7)$$

for

$$k = C^{-1} N^{\frac{1}{2}} \log N + x N^{\frac{1}{2}}, \quad \text{with } C = \pi \sqrt{\frac{2}{3}}. \quad (3.8)$$

Summing up, on the one hand the entropy of a black hole made out of N D0 branes in a theory with k species is given by $p_k(N)$ in (3.7). On the other, we know the entropy of such a black hole is given by $p(N) \sim \sqrt{N}$ by microstate counting. We now show that this scaling can be also recovered if and only if the number of species in the theory is at least of order \sqrt{N} . Indeed, equation (3.7) gives an order one number as $N \rightarrow \infty$ (so that both $p(N)$ and $p_k(N)$ are of the same order) unless $x \rightarrow -\infty$. Imposing that x should be bounded from below, (3.8) automatically implies that $k \gtrsim \sqrt{N}$ asymptotically, where we are ignoring a multiplicative $\log N$ correction. Hence, the number of species $N_s = \sqrt{N}$ suffices to explain the black hole entropy scaling. Equivalently, most microscopic configurations can be built using blocks of bound states of at most $N_s = \sqrt{N}$ D0-branes.

In fact, we incidentally note that the distribution is dominated by states containing at least one bound state of *exactly* $N_s = \sqrt{N}$ D0-branes. Following [89], we introduce

⁸Similar counting problems arise in many other setups, for instance characterizing 1/2 BPS states with potentially large R-charges in AdS/CFT, see e.g. [86, 87].

⁹The equivalence is clear by using the relation of partitions with Young diagrams. A partition of $N = l_1 + \dots + l_k$ with $l_i \geq l_{i+1}$ with *at most* k parts can be depicted as a Young diagram with N boxes arranged in k rows of lengths l_i . By conjugating the diagram, i.e. exchanging the roles of rows and columns, it corresponds to a partition of N with arbitrary number of parts, but each part being *at most* k .

the number $p(N, k)$ as the number of partitions of *exactly* k parts (equivalently, partitions with at least one part *exactly equal* to k). One can define a probability measure for the likelihood of a random microstate containing at least one bound state of k D0-branes:

$$f_{N,k} = \frac{p(N, k)}{p(N)}, \quad \sum_{k=1}^N f_{N,k} = 1. \quad (3.9)$$

Defining the quantity

$$X(k) = \frac{k}{\sqrt{N}} - \frac{1}{C} \log n, \quad (3.10)$$

the probability distribution takes the asymptotic form

$$f_{N,k} \sim e^{-\frac{C}{2} e^{-\frac{2}{C} X(k)}} - e^{-\frac{C}{2} e^{-\frac{2}{C} X(k-1)}}. \quad (3.11)$$

The probability was shown in [90] to be maximized at a value

$$k_0 \sim \frac{\sqrt{6}}{\pi} \sqrt{N} L + \frac{6}{\pi^2} (3(L+1)/2 - L^2/4) - \frac{1}{2}, \quad \text{with } L = \log(\sqrt{6N}/\pi). \quad (3.12)$$

This shows that the species scale is exactly given by $N_s = \sqrt{N}$ (and for instance, not smaller), as we had anticipated.

In the previous description, we simply used the microscopic D0-brane combinatorics, and the abstract definition of species scale. In an actual spacetime description of the system, it will be manifest that the D0-brane stack is a small black hole exploring the infinite distance limit of strong coupling, i.e. decompactification to 11d Mtheory. This will allow for the interpretation of the species scales as the 11d Planck scale. We thus turn to the description of the stretched horizon and the species scale from the spacetime perspective.

3.2 Hot D0-branes

In the previous section we have argued that the D0-branes solution develops an species scale-sized stretched horizon. Naively, a direct approach to confirm the existence of a stretched horizon would be to solve the equations of motion including all the higher derivative corrections. However, given the lack of knowledge about the precise form of all such corrections, this is actually not feasible. In this section, we propose an

alternative way to use spacetime equations of motion to explore the finite entropy of the system, in a universal way insensitive to the details of such corrections. We can explore the resolution of the singularity by studying the system at a finite temperature, and determine its properties as a function of N in the limit of vanishing temperature. We will find that the entropy goes as $N^{\frac{1}{2}}$, in perfect agreement with the microstate counting of the previous section. This provides further evidence that the would-be stretched horizon of a stack of D0 branes should have an area that scales as $N^{\frac{1}{2}}$.

We take the near-extremal (finite temperature) solution for D0-branes (see e.g. [52]), with a horizon r_0 . This comes down to introducing factors of

$$f_0 = \left(1 - \frac{r_0^7}{r^7}\right) \quad (3.13)$$

in the metric (3.4), where r_0 is tied to the energy density ϵ of the branes above extremality:

$$r_0^7 \sim \alpha'^4 g_s^2 \epsilon. \quad (3.14)$$

In the limit where $\epsilon \rightarrow 0$, we recover the extremal (zero temperature) stack of D0 branes.

In the spirit of the gauge/gravity correspondence for general D-branes in [52], we will carry out the computation of the entropy in the near-horizon limit

$$r \rightarrow 0 \quad , \quad \alpha' \rightarrow 0 \quad , \quad U = r/\alpha' = \text{fixed} \quad , \quad g_{QM}^2 = (2\pi)^{-2} g_s \alpha'^{-3/2} = \text{fixed}. \quad (3.15)$$

Here g_{QM} is the coupling of the D0-brane worldvolume quantum mechanics. It is useful to define the dimensionless coupling at some energy scale U

$$g_{\text{eff}}^2 \sim g_{QM}^2 N U^{-3} \sim \lambda U^{-3}, \quad (3.16)$$

where $\lambda = g_{QM}^2 N$ is the (dimensionful) 't Hooft coupling.

The solution in the limit becomes

$$\begin{aligned} \frac{ds^2}{\alpha'} &= -\frac{U^{7/2}}{\sqrt{d_0 \lambda}} f_0 dt^2 + \frac{\sqrt{d_0 \lambda}}{U^{7/2}} \left(\frac{dU^2}{f_0} + U^2 d\Omega_8^2 \right), \\ e^\phi &= \frac{(2\pi)^2}{d_0} \frac{1}{N} \left(\frac{\lambda d_0}{U^3} \right)^{7/4} \sim \frac{g_{\text{eff}}^{7/2}}{N}, \end{aligned} \quad (3.17)$$

where

$$d_0 = 240\pi^5, \quad U_0^7 = a_0 g_{QM}^4 \epsilon, \quad a_0 = \frac{4480 \pi^7}{3}. \quad (3.18)$$

The horizon U_0 can be expressed in terms of the temperature as:

$$T^{-1} = \frac{4}{7}\pi\sqrt{\lambda d_0}U_0^{-5/2}. \quad (3.19)$$

We now move on to the Einstein frame to compute the entropy. The Einstein frame spatial metric reads [91]:

$$ds_E^2 = CU^{-7/8} \left(\frac{dU^2}{f_0} + U^2 d\Omega_8^2 \right) \quad \text{with} \quad C = \frac{l_s^2 N^{1/2} d_0^{1/8}}{2\pi\lambda^{3/8}}. \quad (3.20)$$

It is convenient to introduce a new radial variable R , with dimension of length, defined as $U^{9/8} = (R^2/C)$. The metric becomes:

$$ds_E^2 = \frac{256}{81} \frac{dR^2}{1 - \left(\frac{R_H}{R}\right)^{112/9}} + R^2 d\Omega_8^2, \quad (3.21)$$

where R_H has the parametric dependence

$$R_H \sim \left(\frac{T}{\lambda^{1/3}} \right)^{9/40} N^{1/4} l_s. \quad (3.22)$$

So the entropy of the finite temperature black hole in terms of the dimensionless temperature $T/\lambda^{1/3}$ is given by

$$S \sim N^2 \times (T/\lambda^{1/3})^{9/5}. \quad (3.23)$$

Restoring the powers of N hidden in T and λ , we get the scaling behaviour

$$S \sim (U_0^{5/2}/g_{QM}^{5/3})^{9/5} N^{1/2}. \quad (3.24)$$

In the limit $U_0 \rightarrow 0$, one recovers the familiar zero entropy result. However, the point that the above computation shows is that entropy of the system, when cloaked with a fixed horizon size, scales as $N^{1/2}$, in agreement with our discussion in section 3.1.

4 Species Scale Horizon from Emergent Higher Derivative Terms

It is useful to have a more direct intuition, at the level of the geometry, about the appearance of the species scale horizon for D0-branes uncovered in the previous section.

In analogy with the 4d small black holes in section 2.2, we expect this horizon to show up once higher derivative corrections are included in the effective theory, which become relevant precisely at the species scale. Moreover, such new terms are expected to arise from integrating out the towers of light particles associated to the infinite distance limit of the formerly small black hole, in the spirit of Emergence.

In this section we work out in detail this picture for the D0-brane system. We first show that generic 10d R^4 corrections lead to a finite size horizon. We then consider the specific R^4 couplings arising from integrating out a tower of D0-brane states (i.e. 11d KK gravitons), which constitute the SDC tower associated to the infinite distance limit probed by the small black hole solution (decompactification to 11d M-theory). Interestingly we find that these familiar R^4 terms do not suffice to generate the finite size horizon for the system (let alone a species scale one). Turning things around, this suggests that the appearance of the species scale horizon demanded by the swampland considerations requires a crucial role of new physics beyond the usually considered R^4 terms. In fact, we include a specific set of supersymmetric 8-derivative terms involving the RR field strength 2-form, tractable via a lift to 11d Mtheory, and show that they do produce the species scale horizon, thus confirming the swampland expectations.

4.1 Entropy function analysis of general R^4 terms

It is well established that the existence and properties of stretched horizons for 4d small black holes in theories with higher-curvature corrections can be studied using the entropy function formalism [71]. The idea is to use the ansatz for a putative near horizon AdS_2 solution to evaluate an entropy function, and to extremize it with respect to its parameters. In this section we apply a similar logic to the solution of D0-branes in 10d type IIA, and argue that the introduction of higher curvature corrections leads to a stretched horizon for the system.

Before entering the discussion, let us make a general comment. In this section we only include general terms involving only the curvature, but no other fields. This is for tractability, since curvature terms lead to closed expressions which can be analyzed in a model-independent way, and suffice to illustrate the fairly generic appearance of stretched horizons. The discussion of other corrections can be carried out for instance if a specific set of such correction is fixed, as we do in section 4.4 for a set of 8-derivative

corrections including curvature and the RR 2-form field strength.

4.1.1 The D0-brane solution at two-derivative level

We start by recovering the key features of the two-derivative solution of the D0-brane system from the entropy function formalism [71], as warmup for coming sections. As explained above, we introduce an $\text{AdS}_2 \times \mathbf{S}^8$ ansatz:

$$\begin{aligned} ds^2 &= v_1 \left(-r^2 dt^2 + \frac{1}{r^2} dr^2 \right) + v_2 d\Omega_8^2, \\ F_{rt} &= e, \quad \tilde{F}_{\theta\phi_1 \dots \phi_7} = 0, \\ e^\phi &= g_s, \end{aligned} \tag{4.1}$$

where v_1 and v_2 control the AdS and sphere length scales. For the 2-form field strength this introduces electric but not magnetic charge at the horizon. Finally, the last equation sets the value of the string coupling at the horizon.

The entropy function is given (modulo some irrelevant overall factor) by evaluating the action at the near horizon ansatz (4.1) and applying a Legendre transform with respect to e , namely

$$\mathcal{E}(N, g_s, v, \beta, e) = eN - \int_{S^8} d\Omega_8 \sqrt{-g} \mathcal{L}|_h. \tag{4.2}$$

To evaluate this quantity for the ansatz (4.1) we compute

$$\sqrt{-g} = v_1 v_2^4 \sqrt{g_{\Omega_8}} = \frac{v^5}{\beta} \sqrt{g_{\Omega_8}}, \tag{4.3}$$

$$R = -\frac{2}{v_1} + \frac{56}{v_2} = \frac{56 - 2\beta}{v}, \tag{4.4}$$

$$|F_2|^2 = \frac{1}{2!} F_{\mu\nu} F^{\mu\nu} = -\frac{1}{v_1^2} e^2 = -\frac{\beta^2}{v^2} e^2. \tag{4.5}$$

With hindsight, we have introduced new variables $v = v_2$ and $\beta = v_2/v_1$, to describe the overall length scale of $\text{AdS}_2 \times \mathbf{S}^8$ and the relative scale separation. The result is

$$\mathcal{E}(N, g_s, v, \beta, e) = eN - \frac{8\pi^4 v^3 \left(\beta^2 e^2 g_s^{3/2} - 4(\beta - 28)v \right)}{105\beta}. \tag{4.6}$$

The goal is to extremize this function with respect to g_s , v , β and e , expressing the result in terms of the D0-brane charge N . We start with the extremization with

respect to e , which gives

$$\frac{\partial \mathcal{E}}{\partial e} = 0 \quad \rightarrow \quad e = \frac{105N}{16\pi^4 \beta v^3 g_s^{3/2}}. \quad (4.7)$$

Substituting this result, the entropy function reads

$$\mathcal{E}(N, g_s, v, \beta) = \frac{105N^2}{32\pi^4 \beta v^3 g_s^{3/2}} + \frac{32\pi^4 (\beta - 28)v^4}{105\beta}. \quad (4.8)$$

We now extremize with respect to v and β , and get

$$\frac{\partial \mathcal{E}}{\partial v} = \frac{\partial \mathcal{E}}{\partial \beta} = 0 \quad \rightarrow \quad v^7 = \frac{1575N^2}{4096\pi^8 g_s^{3/2}}, \quad \beta = 49. \quad (4.9)$$

Incidentally, the approximately order 1 value of β (taking volume factors into account) is natural from the absence of scale separation in AdS vacua [92]. The resulting entropy function now reads

$$\mathcal{E}(N, g_s) \sim \frac{N^{8/7}}{g_s^{6/7}}. \quad (4.10)$$

We see that there is no well-defined extremum for the parameter g_s , it lies at $g_s \rightarrow \infty$. This simply reflects the fact that the D0-brane solution drives the dilaton to infinity in moduli space at its core, as observed above from the 10d solution. Indeed, we can recover other features of the 10d solution from the entropy function, by studying the scaling of different quantities with g_s . For instance, from (4.9) and (4.4), we have that in the $g_s \rightarrow \infty$ limit

$$v_2 = v \sim \frac{1}{g_s^{3/14}} \rightarrow 0, \quad R \sim g_s^{3/14} \rightarrow \infty. \quad (4.11)$$

Namely, we recover that the horizon volume goes to zero, and the scalar curvature blows up with exactly the same scalings as would be obtained directly from the D0 solution at its core. Note that the g_s we consider in this section is to be compared to the dilaton near the core of the D0 stack in the solution (3.4). Indeed, g_s is the value of the string coupling at the horizon of the would-be black hole, whilst g_s^∞ in (3.4) is the string coupling infinitely far away from the stack of branes.

We therefore recover that the process of extremization of the Wald entropy function of the near-horizon geometry of a black hole electrically charged under the RR 2-form in 10d describes the near-core behaviour of a stack of D0 branes. The fact that the Wald

entropy is only extremized for a value of the string coupling at the core that is blowing up, with a horizon that tends to zero size is exactly the behaviour expected from small black holes, as in Section 2. We conclude from this that the Wald entropy formalism constitutes a reliable description of D0-brane stacks as small black holes. In the next section, we will introduce higher derivative corrections to the 10d supergravity which will affect the Wald entropy in such a way that the same exact minimization procedure will lead to a finite sized black hole instead of a small one.

4.1.2 The D0-brane solution and higher derivative terms

In analogy with the 4d small black hole case, we may expect that the D0-brane solutions can develop a stretched horizon upon the inclusion of suitable higher derivative corrections. In 10d type IIA the first corrections allowed by supersymmetry arise at 8-derivative level. There is a vast literature devoted to the computation of these corrections at tree and one-loop level (see e.g. [76, 93–98]). However, for the present purposes it will be more efficient to assume the presence of general R^4 corrections (the example of supersymmetric R^4 corrections will be discussed explicitly in section 4.3). Also, as explained earlier, we focus on terms involving only curvatures, which allows for a tractable model-independent analysis. The inclusion of terms involving other fields is discussed in section 4.4 for a specific set of supersymmetric 8-derivative terms including the RR 2-form field strength.

We now describe the entropy function computation in the presence of general R^4 corrections. The corrections can be written as a linear combination of all possible contractions between eight inverse metrics and four Riemann tensors with all the indices down. Inspecting the ansatz in (4.1), we see that each inverse metric gives either a $1/v_1$ or a $1/v_2$ contribution, while each Riemann gives either a v_1 or a v_2 . In terms of v and β , this means that any of these contributions will take the form $\beta^a v^{-4}$, with a ranging from -4 to 8 . For instance, the $a = -4$ one corresponds to the case in which all the Riemanns give v_1 and all the metrics give $1/v_2$, while the $a = 8$ one corresponds to all the Riemanns giving v_2 and all the metrics giving $1/v_1$.

Following this reasoning, we can write the R^4 term evaluated at the horizon as

$$\mathcal{L}_{R^4}|_h = -g_s^{1/2} \frac{p_{12}(\beta)}{\beta^4} \frac{1}{v^4}, \quad (4.12)$$

where p_{12} denotes a degree 12 polynomial, and the minus sign is there for future convenience. Note that we already make explicit the dependence on the string coupling, which is fixed to that of the R^4 terms arising at first loop in perturbative Type IIA string theory. Motivated by having $g_s \rightarrow \infty$ at the core of the D0 solution at the two-derivative level, we are neglecting the tree level piece. One could worry that even higher loop contributions would dominate as $g_s \rightarrow \infty$, but they actually vanish. This has been considered previously in the literature from the Type IIA perspective [76, 99, 100] and, as it will become clearer in section 4.4, it is required for a consistent uplift to M-theory in this limit.

Including the extra term in (4.12), the entropy function (4.2) reads

$$\mathcal{E}(N, g_s, v, \beta, e) = eN - \frac{8}{105}\pi^4\beta e^2 g_s^{3/2} v^3 + \frac{16\pi^4 v \left(g_s^{1/2} p_{12}(\beta) + 2(\beta - 28)\beta^4 v^3 \right)}{105\beta^5}. \quad (4.13)$$

The extremization with respect to e is insensitive to the new R^4 term, so the result is again (4.7). Plugging this result, the entropy function reads

$$\mathcal{E}(N, g_s, v, \beta) = \frac{105N^2}{32\pi^4\beta g_s^{3/2} v^3} + \frac{16\pi^4 \left(g_s^{1/2} p_{12}(\beta)v + 2(\beta - 28)\beta^4 v^4 \right)}{105\beta^5}. \quad (4.14)$$

In order to display the effect of the higher derivative term in the dilaton, let us now consider the extremization with respect to g_s . We obtain

$$\frac{\partial \mathcal{E}}{\partial g_s} = 0 \quad \rightarrow \quad g_s = \frac{105}{16\pi^4} \sqrt{\frac{3}{2}} \frac{\beta^2 N}{\sqrt{p_{12}(\beta)} v^2}. \quad (4.15)$$

We see that the new contributions generically stabilize the dilaton. We note that the solution only exist if $p_{12}(\beta_0) > 0$, where β_0 is the yet to be determined value for β extremizing the entropy function. The latter is determined by the extremization with respect to v , namely

$$\frac{\partial \mathcal{E}}{\partial v} = \frac{128\pi^4(\beta - 28)v^3}{105\beta} = 0 \quad \rightarrow \quad \beta = 28. \quad (4.16)$$

Note that we again obtain a value compatible with the absence of scale separation, suggesting that this property is fairly robust against inclusion of higher derivative corrections.

Finally, the extremization with respect to β gives

$$\left. \frac{\partial \mathcal{E}}{\partial \beta} \right|_{\beta=28} = 0 \quad \rightarrow \quad v^4 = \frac{4 p_{12}(28) - 21 p'_{12}(28)}{175616 \cdot 6^{1/4} \pi^2 p_{12}(28)^{1/4}} \sqrt{\frac{5N}{7}}, \quad (4.17)$$

where $p'_{12}(\beta)$ is the derivative of the polynomial with respect to β . Hence, the existence of the solution requires

$$p_{12}(28) > 0 \quad , \quad 4 p_{12}(28) > 21 p'_{12}(28). \quad (4.18)$$

We will see the explicit evaluation of such polynomials for a particular R^4 correction in section 4.3.

In conclusion, under fairly general circumstances, the presence of higher derivative corrections suggests the existence of a stretched horizon for 10d D0-brane solutions. From the above expressions, the scaling of its properties with N is

$$S \sim \sqrt{N} \quad , \quad v^4 \sim \sqrt{N} \quad , \quad \beta = 28 \quad , \quad g_s \sim N^{3/4} \quad , \quad e \sim N^{-1/2}. \quad (4.19)$$

The entropy of the system scales as \sqrt{N} , in agreement with the microstate counting in section 3.1. Notice that this, as well as all the scalings above, rely heavily in the string coupling dependence made explicit in (4.12) and corresponding to the one-loop contribution in perturbative Type IIA string theory. The string coupling at the horizon blows up as $N \rightarrow \infty$, which is consistent with neglecting the tree level contribution in this regime. In the next section we will shed some further light on these scalings, since they lead to the appearance of the species scale.

4.1.3 The Species Scale and Cosmic Censorship

The Species Scale

Here we shortly note that the above horizon scale indeed corresponds to the species scale of the SDC tower associated to the infinite distance limit probed by the solution. This infinite distance limit corresponds to a decompactification to 11d M-theory (consistently with the above mentioned fact that the horizon value of g_s gets strong in the large N regime). The extra dimension is reconstructed by the SDC tower of D0 branes, whose species scale corresponds to the 11d Planck scale. Restoring the 10d Planck scale, we have

$$\Lambda_s \sim M_{11} \sim g_s^{-1/12} M_{10} \sim N^{-1/16} M_{10}, \quad (4.20)$$

where we have used the scaling of g_s with N in equation (4.19). It is indeed easy to show that this coincides with the inverse radius of the stretched horizon. Using the scaling of v with N in (4.19), we have

$$r_h^{-1} \sim v^{-1/2} M_{10} \sim N^{-1/16} M_{10}. \quad (4.21)$$

Let us note that this agreement is easily understood by the fact that the stretched horizon arises from the competition between the classical and the R^4 terms in the effective action. The scale at which these two can compete indeed corresponds to one of the notions of the species scale [47]. The fact that it matches the species scale associated to the tower of D0s can be directly checked in the effective action and, as we explain in section 4.2, can be understood from the emergence of higher derivative terms coming from integrating out the tower of D0s. Here we see that these two notions of species scale also coincide with the one of the smallest BH describable within the EFT, in this case the stretched horizon of the D0 black hole.

The D0-brane bound state size scale

We have emphasized the appearance of a finite size for the system from the space-time viewpoint, via the appearance of a stretched horizon. However, as any quantum system, the D0-brane system has a characteristic size, in particular that associated to the size of the D0-brane bound states. This has been studied in particular in the context of M(atric) theory. In this context, the outcome of [51, 101–103], is an estimate of the size of a bound state of N D0-branes scaling as

$$R \gtrsim N^{1/3} M_{11}^{-1}. \quad (4.22)$$

This grows with N in a way seemingly stronger¹⁰ than the stretched horizon size $r_h \sim M_{11}^{-1}$ discussed above. However, an important observation is that (4.22) is derived in the limit of $g_s \rightarrow 0$ with M_{11} fixed, with g_s the coupling in flat space. Hence, for a proper comparison, in the evaluation of the horizon size $r_h \sim N^{1/16} M_{10}^{-1} \sim g_s^{-1/12} N^{1/16} M_{11}^{-1}$, one should not use the horizon value $g_s \sim N^{3/4}$ in matching M_{10} , but rather consider g_s as a free asymptotic parameter. Thus, in the $g_s \rightarrow 0$ and M_{11} fixed limit, the horizon

¹⁰Note that the stronger growth also holds even if we restrict to the dominant bound state of $k \sim N^{1/2}$ D0-branes, as in section 3.1, which would yield $R \sim N^{1/6}$.

size is parametrically larger than the size of the bound state of gravitons, which are effectively cloaked behind the horizon.

A cosmic censorship interpretation

The above considerations motivate a cosmic censorship interpretation of our result, which provides a different angle to other uses of cosmic censorship in the analysis of swampland constraints (see e.g. [104, 105]). The classical 2-derivative solution displays a singularity. In the full theory, the pathological implications of such singularities must be avoided, either by some desingularization [106], addition of extra degrees of freedom as in orbifolds, or the appearance of a horizon cloaking it. We have shown that in our system the latter occurs via the use of higher derivative interactions, despite the fact that it does not happen in the low-energy two-derivative approximation. Interestingly, this happens in the realm of the effective field theory, on the verge of its validity. This is again tied up to the notion of the species scale as that at which the tower of higher derivative interactions becomes relevant.

4.2 The D0-brane SDC tower and Higher Derivative Emergence

The infinite distance limit in moduli space explored by the core of the solution is the strong coupling limit of type IIA theory, which corresponds to the decompactification limit of 11d M-theory on \mathbf{S}^1 . The tower of states predicted by the swampland distance conjecture corresponds to D0-brane states¹¹, namely the KK momentum states of 11d graviton multiplets. The computation is indeed simpler in this 11d picture, and is a higher-dimensional version of that encountered in section 2.2.3. In fact is a well-established story that this set of states can lead, at the 1-loop level, to the appearance of higher derivative corrections to the 10d effective action, which can be interpreted as tree- and one-loop level (in g_s) terms of type IIA theory [76], as we now discuss.

The terms we are going to compute are the $t_8 t_8 R^4$ terms¹², in principle of the kind

¹¹We expect readers not to confuse the D0-brane states in the SDC tower with the D0-branes sourcing the solution in the previous sections.

¹²As we will explain in section 4.3, the actual 10d type IIA R^4 terms contain a further piece with an $\epsilon_{10}\epsilon_{10}$ structure. This however does not contribute to 4-graviton scattering amplitudes, or conversely is not captured by our 4-graviton 1-loop amplitude (it would require a 5-point computation). This is analogous to the similar statements for the odd-odd structure in the R^4 computation from scattering amplitudes in string theory, see e.g. [107].

included in the analysis in section 4.1.2 (see also section 4.3 for more details on their structure). They are protected by supersymmetry, so only the 1-loop diagram of BPS 11d graviton KK modes can contribute. These can be easily computed from a worldline formalism. Using the general expression (2.24), integrating over continuous momenta, and particularizing for $d = 10$, $n = 4$, we get

$$\mathcal{A}_{10,4} = \frac{\sqrt{\pi}}{2R} \tilde{K} \int_0^\infty \frac{d\tau}{\tau^2} \sum_k e^{-\pi\tau R^{-2}k^2}, \quad (4.23)$$

where the prefactor \tilde{K} contains all the kinematics, and is the linearized approximation to the $t_8 t_8 R^4$ term in the effective action. As in the 4d case, the above expression diverges in two ways: because of the integral and because of the infinite sum. We can perform a Poisson resummation, so that both are nicely combined and we get

$$\frac{1}{\pi^{3/2}} \mathcal{A}_{10,4} = \tilde{K} \int_0^\infty d\hat{\tau} \hat{\tau}^{1/2} \sum_l e^{-\pi\hat{\tau} R^2 l^2} = C \tilde{K} + \frac{\zeta(3)}{\pi R^3} \tilde{K}, \quad (4.24)$$

where $\hat{\tau} = \tau^{-1}$. This Poisson summation trades the KK momentum k for the winding number l of the worldline along the \mathbf{S}^1 . The only divergence is now in the $l = 0$ piece, which has been isolated as the first terms in the second equality; C is an unknown coefficient regularizing the divergence.¹³

As anticipated, the quantum corrections coming from the SDC tower lead to a higher derivative R^4 correction, in principle of the kind invoked to lead to a stretched horizon. There remains to perform a detailed evaluation of these terms and their effect in the entropy function argument, which we postpone to section 4.3.

An important remark is that the R^4 correction obtained in this way is ambiguous. The 4-graviton scattering is an on-shell quantity that is invariant under local field redefinitions $g_{ij} \rightarrow g_{ij} + \delta g_{ij}$, which is usually used to get rid of terms involving Ricci tensors (see e.g. [55]). However, since the $\text{AdS}_2 \times \mathbf{S}^8$ ansatz has a non-vanishing Ricci tensor, doing this is not innocuous for our purposes. We will thus stick to the full $t_8 t_8 R^4$ term, including Ricci terms, as it is the natural kinematic function that appears

¹³In [76], it was fixed by T-duality upon further \mathbf{S}^1 compactification. In our present context, there is no way of fixing it without this kind of extra UV information. It would be interesting if performing the sum over KK modes up to the species scale one could reproduce its precise value, along the lines of [39–43].

in this computation à la Emergence. That this tensor structure gives the full off-shell R^4 correction to the effective action was also suggested in [95]. As evidence for this, it was found that the terms containing no more than one Ricci tensor, which were reliably computed from the vanishing of the worldsheet beta function, precisely match the ones appearing in the $t_8 t_8 R^4$ structure.

Higher Derivative Emergence

When translated to the Type IIA frame, the first and second terms in (4.24) precisely reproduce the $t_8 t_8 R^4$ tensor structure of the 1-loop and tree-level string perturbation theory contributions, respectively. Therefore, this establishes that these can be interpreted as coming from integrating out the full tower of D0-particles, as well as the fast-movers in the massless 10d sector. Let us remark that this result is very much in the spirit of the Emergence Proposal, albeit for higher derivative terms rather than just the kinetic terms.

Note also that the above computation integrates out the full tower of states, i.e. not just up to the species scale as in [39–43]. In fact, interpreting the latter recipe as a regularization procedure to be applied when integrating out infinite number of states, one could hope to recover the right value for C ; we leave this as an interesting open question. On the other hand, notice that integrating the full tower is in the spirit of the recent proposals in [42, 43]. It would be interesting to explore these connections further.

4.3 Supersymmetric 10d R^4 terms are not enough

In this section, we particularize the analysis in section 4.1.2 to the specific R^4 terms appearing in the Type IIA effective action (for a concrete choice fixing all the already mentioned ambiguities concerning these terms). By plugging the $\text{AdS}_2 \times \text{S}^8$ ansatz, we will compute the polynomial appearing in equation (4.12) and check the conditions in equation (4.18).

We will show that, perhaps surprisingly, these supersymmetric R^4 terms do not satisfy the constraints in section 4.1.2 required to generate a finite size horizon. Note that this does not invalidate the analysis of the entropy function, but rather shows that the appearance of the horizon from pure higher curvature terms demands the use of other possibly non-supersymmetric terms. Alternatively, one can maintain the problem

in the realm of supersymmetric corrections, and include higher derivative couplings involving the RR 2-form field strength; these go beyond the analysis in section 4.1.2, and will be discussed in section 4.4.

Hence, we focus on the supersymmetric R^4 corrections in 10d type IIA theory. Following the notation in [98], at the eight derivatives level, the type IIA action gets supplemented by additional terms quartic in the Riemann tensor at both tree level and first loop in the string coupling g_s :

$$S_{\text{IIA}} = S_{\text{IIA}}^{\text{class}} + \frac{1}{3 \cdot 2^{11}} (S^{\text{tree}} + S^{\text{loop}}), \quad (4.25)$$

where S^{tree} and S^{loop} both arise at level α'^3 . In fact, it has been conjectured that these are the only two contributions to this type of term in string perturbation theory [76, 99, 100]. They can be written as follows in the Einstein frame and in 10d Planck units:

$$S^{\text{tree}} = \frac{\zeta(3)}{2} \int d^{10}x \sqrt{-g} e^{-\frac{3}{2}\phi} \left(t_8 t_8 R^4 + \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 \right), \quad (4.26)$$

$$S^{\text{loop}} = \frac{\pi^2}{6} \int d^{10}x \sqrt{-g} e^{\frac{1}{2}\phi} \left(t_8 t_8 R^4 - \frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 \right), \quad (4.27)$$

where the form of the $t_8 t_8 R^4$ and $\epsilon_{10} \epsilon_{10} R^4$ structures can be found in Appendix A. Let us only consider the 1-loop term, which is the leading one as $g_s \rightarrow \infty$.

Evaluating $t_8 t_8 R^4$ and $\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4$ on the $AdS_2 \times S^8$ near-horizon geometry (4.1), one obtains:

$$t_8 t_8 R^4 = \frac{192 (3\beta^4 + 56\beta^2 + 2520)}{v^4}, \quad (4.28)$$

$$\frac{1}{8} \epsilon_{10} \epsilon_{10} R^4 = \frac{161280 (4\beta - 1)}{v^4}. \quad (4.29)$$

Therefore, the 1-loop term can be put in the form (4.12) if

$$p_{12}(\beta) \sim -\beta^4 (3\beta^4 + 56\beta^2 - 3360\beta + 3360), \quad (4.30)$$

where we are ignoring an irrelevant positive-defined prefactor. Now we can check the conditions in (4.18) for the existence of an extremum of the entropy function. The second one is satisfied, while the first one is not. This implies that the familiar R^4 terms used so far do not suffice to generate the stretched horizon for the D0 solution.

This does not invalidate the main claim that the system should develop a species scale horizon. As explained earlier, we expect other higher derivative couplings to contribute non-trivially, and the horizon may very well develop when they are ultimately included. Indeed, we will show in the next section that a simple modification of the above computation promoted to M-theory allows to include couplings involving the RR 2-form F_2 , and lead to a non-trivial species scale stretched horizon for the D0-brane system.

4.4 Including F_2 : Species scale D0-brane horizon from M-theory couplings

In the previous section we showed that 10d supersymmetric R^4 terms do not generate a finite size horizon. Hence, the requirement that there should arise species scale horizon, as demanded by swampland considerations, implies that one should look further. One possibility is to consider other possibly non-supersymmetric pure higher curvature terms. A more attractive avenue is to stay in the realm of supersymmetric terms but include higher derivative couplings involving the RR 2-form field strength F_2 . These go beyond the analysis in section 4.1.2, but, once a specific set of couplings is fixed, are amenable to a direct study again via the entropy function.

In this section we include such terms, completing the analysis at the supersymmetric eight derivatives level of the effective action. The most efficient way of doing this is by encoding these terms in the 11d M-theory effective action, since both the 10d metric and RR 1-form become part of the 11d metric. This is, by encoding the extremal near-horizon ansatz in the 11d metric and plugging it into the M-theory effective action, supplemented by R^4 terms, we can effectively recover the 10d Lagrangian evaluated at the horizon including both R^4 and terms involving F_2 . Let us remark that we will keep the 10d point of view at all times to focus in the presence of a stretched horizon for the D0 solution. This is, here we think about the 11d effective action as just an efficient way of encoding the eight derivatives terms involving F_2 and evaluating them at the extremal near-horizon ansatz.

Our analysis is close to that in [55], with the difference that we emphasize the 10d perspective in the computation. Even though we find analogous results, an important difference in the analysis is that we keep the full $t_8 t_8 R^4$ structure, while only purely Riemannian terms were included in [55]. As already discussed in section 4.2,

the Wald entropy analysis is sensitive to the terms including the Ricci tensor which, via Emergence, are naturally encoded in the $t_8 t_8 R^4$ structure. This is consistent with the results obtained in [95], where the terms in $t_8 t_8 R^4$ including one Ricci tensor were unambiguously shown to appear in the Type IIA effective action.

Following again the notation in [98], and setting the three-form to zero for our purposes, the M-theory effective action can be written at the eight derivatives level as

$$S_M = \frac{1}{2\kappa_{11}^2} \int d^{11}x \sqrt{-\hat{g}} \left(\hat{R} + \frac{(4\pi\kappa_{11}^2)^{2/3}}{(2\pi)^4 \cdot 3^3 \cdot 2^{13}} \left(\hat{t}_8 \hat{t}_8 \hat{R}^4 - \frac{1}{24} \epsilon_{11} \epsilon_{11} \hat{R}^4 \right) \right). \quad (4.31)$$

To simplify the computations, we will take $\kappa_{11} = 1$.¹⁴ The \hat{t}_8 and ϵ_{11} tensors are defined in complete analogy with the Type IIA case. In fact, after writing these terms as contractions of four Riemann tensors, the 11d R^4 term look exactly the same as the 10d one. More details about these structures can be found in Appendix A.

As it is well-known, the circle compactification ansatz from 11 to 10 dimensions is:

$$d\tilde{s}^2 = e^{-\frac{1}{6}\phi} ds^2 + e^{\frac{4}{3}\phi} (dz - C_\mu dx^\mu)^2. \quad (4.32)$$

Plugging this into the effective action recovers the 10d Einstein frame Type IIA effective action up to the eight-derivative level. Instead of doing this and then taking the extremal $AdS_2 \times S^8$ near-horizon ansatz, we perform the computation at the level of M-theory. That is, we first encode the $AdS_2 \times S^8$ ansatz into the 11d metric, and then we plug it into the 11d effective action. This allows us to read off the 10d Lagrangian evaluated at the horizon. For this last step, we need to take into account the relation between the 11d and 10d determinant of the metric appearing in the action. This gives the following relation:

$$\mathcal{L}_{IIA}|_h = g_s^{-1/6} \mathcal{L}_M|_{11d \text{ ansatz}}. \quad (4.33)$$

To encode the extremal $AdS_2 \times S^8$ near-horizon geometry in the 11d metric ansatz, we just plug equation (4.1) into (4.32) to get

$$d\tilde{s}^2 = -g_s^{-1/6} v_1 \left(r^2 dt^2 + \frac{1}{r^2} dr^2 \right) + g_s^{-1/6} v_2 d\Omega_8^2 + g_s^{4/3} (dz - e r dt)^2. \quad (4.34)$$

¹⁴Even though this might seem to imply that the results will be in 11d Planck units, when mapping to the type IIA Einstein frame, the 11d Planck scale gets traded for the 10d one. Thus, the final results are actually in 10d Planck units.

Evaluated on this ansatz, the 11d Ricci scalar reads

$$\hat{R} = g_s^{1/6} \frac{\beta^2 e^2 g_s^{3/2} - 4v(\beta - 28)}{2v^2}, \quad (4.35)$$

where we introduced new variables $v = v_2$ and $\beta = v_2/v_1$ as we did in section 4.1.1. When plugged into (4.33), this recovers the two-derivative piece of the Type IIA Lagrangian evaluated at the horizon (c.f. equation (4.6)).

Similarly, we evaluate the M-theory R^4 terms in this ansatz, getting

$$\begin{aligned} \left(\hat{t}_8 \hat{t}_8 - \frac{1}{24} \epsilon_{11} \epsilon_{11} \right) \hat{R}^4 = \frac{3g_s^{2/3}}{4v^8} & \left(287\beta^8 e^8 g_s^6 - 1392\beta^7 e^6 g_s^{9/2} v \right. \\ & + 32(83\beta^2 + 308) \beta^4 e^4 g_s^3 v^2 \\ & - 768(3\beta^3 + 28\beta - 280) \beta^2 e^2 g_s^{3/2} v^3 \\ & \left. + 256(3\beta^4 + 56\beta^2 - 3360\beta + 3360) v^4 \right). \end{aligned} \quad (4.36)$$

As a check, we can recover the Type IIA R^4 result from the previous section by setting $e = 0$ and plugging (4.36) into (4.33) (c.f. equation (4.28) with the $g_s^{1/2}$ factor appearing in (4.12)).

Putting these two results together in the 11d Lagrangian, using equation (4.33) and plugging the result into the definition of the entropy function in (4.2), we finally obtain:

$$\begin{aligned} \mathcal{E}(N, g_s, v, \beta, e) = eN - \frac{8\pi^4 v^3 \left(\beta^2 e^2 g_s^{3/2} - 4(\beta - 28)v \right)}{105\beta} \\ - \left(\frac{\pi}{2} \right)^{2/3} \frac{g_s^{1/2}}{2580480 v^3 \beta} & \left(287\beta^8 e^8 g_s^6 - 1392\beta^7 e^6 g_s^{9/2} v + 32(83\beta^2 + 308) \beta^4 e^4 g_s^3 v^2 \right. \\ & \left. - 768(3\beta^3 + 28\beta - 280) \beta^2 e^2 g_s^{3/2} v^3 + 256(3\beta^4 + 56\beta^2 - 3360\beta + 3360) v^4 \right). \end{aligned} \quad (4.37)$$

Here we have already performed the integral in equation (4.2), taking into account the determinant of the metric evaluated in the ansatz as given in (4.3).

The entropy function turns out to be rather involved, and we cannot extremize it analytically. Doing so numerically would require fixing N to a set of different values, extremizing the entropy function for each of these and then fitting the dependence on

N of the various quantities (g_s, v, \dots). Instead of doing so, let us perform a change of variables inspired by the scalings found in section 4.1.2. We define

$$g_s = \tilde{g}_s N^{3/4}, \quad v = \tilde{v} N^{1/16}, \quad \beta = \tilde{\beta}, \quad e = \tilde{e} N^{-1/2}. \quad (4.38)$$

Introducing this into the entropy function, we get

$$\begin{aligned} \mathcal{E}(N, \tilde{g}_s, \tilde{v}, \tilde{\beta}, \tilde{e}) = \sqrt{N} & \left[\tilde{e} - \frac{8\pi^4 \tilde{v}^3 \left(\tilde{\beta}^2 \tilde{e}^2 \tilde{g}_s^{3/2} - 4(\tilde{\beta} - 28)\tilde{v} \right)}{105\tilde{\beta}} \right. \\ & - \left(\frac{\pi}{2} \right)^{2/3} \frac{\tilde{g}_s^{1/2}}{2580480 \tilde{v}^3 \tilde{\beta}} \left(287\tilde{\beta}^8 \tilde{e}^8 \tilde{g}_s^6 - 1392\tilde{\beta}^7 \tilde{e}^6 \tilde{g}_s^{9/2} \tilde{v} + 32 \left(83\tilde{\beta}^2 + 308 \right) \tilde{\beta}^4 \tilde{e}^4 \tilde{g}_s^3 \tilde{v}^2 \right. \\ & \left. \left. - 768 \left(3\tilde{\beta}^3 + 28\tilde{\beta} - 280 \right) \tilde{\beta}^2 \tilde{e}^2 \tilde{g}_s^{3/2} \tilde{v}^3 + 256 \left(3\tilde{\beta}^4 + 56\tilde{\beta}^2 - 3360\tilde{\beta} + 3360 \right) \tilde{v}^4 \right) \right]. \end{aligned} \quad (4.39)$$

This change of variables does the important job of factoring out all the N -dependence in the entropy function, which is very non-trivial since all the terms had to conspire to give the same overall \sqrt{N} factor. Thanks to this, we can now focus on the terms in parenthesis and look for its N -independent extrema numerically. If an extremum exist, then we see that the scalings we found in section 4.1.2 are automatically guaranteed. Furthermore, given that the overall factor in front of the entropy function is \sqrt{N} , the scaling of the entropy from the microscopic counting in section 3.1 is also automatically guaranteed.

Finally, we look for extrema of the terms in parenthesis in the previous equation or, equivalently, of $\mathcal{E}(1, \tilde{g}_s, \tilde{v}, \tilde{\beta}, \tilde{e})$. This cannot be done analytically, so we perform a numerical search. The result is the following extremum:

$$\tilde{g}_s \approx 2.90365, \quad \tilde{v} \approx 0.159653, \quad \tilde{\beta} \approx 14.0556, \quad \tilde{e} \approx 0.0479134. \quad (4.40)$$

For these values, one can check that the derivatives of the entropy function are 14 orders of magnitude smaller than the value of function itself. This indicates that, to a very good approximation, this is an extremum of the entropy function. To make this extremum more apparent, we can fix one of the variables to its value at the extremum and display the three-dimensional gradient flow. This is shown in figure 1, where we can see the presence of a non-trivial extremum. As it can be checked also by direct

computation, we also see that the extremum is in fact a saddle point. This fact is not relevant for the Wald entropy formalism, that only requires the presence of an extremum. It would be interesting to understand the thermodynamical stability of this stretched horizon, somewhat along the lines of [55]. However, this analysis presumably requires a systematic inclusion of even higher derivative terms to achieve control of order one factors.




Figure 1. Gradient flows of the entropy function with one of the variables fixed. Lighter/yellow colors denotes bigger values for the gradient, while darker/blue colors denote smaller ones.

In conclusion, we found that the Type IIA action including *all* terms up to the eight derivative level captures the presence of the species scale stretched horizon for the D0 solution. In addition, the entropy associated to the horizon nicely fits the microscopic counting and the near-horizon $\text{AdS}_2 \times \text{S}^8$ is not scale separated. This gives support to the idea that, even though a proper geometric treatment of a stretched horizon requires the inclusion of all the infinite amount of higher derivative operators, the classical level and the (complete) first non-trivial higher derivative level seem to suffice for capturing its presence and its scaling properties with the species scale.

As in the case of 10d R^4 , the new set of higher derivative terms can be understood as arising from integrating out the SDC tower of D0-branes, in the spirit of Emergence. Indeed, since all these couplings are related by 11d Poincaré invariance, the computation of the 4-point amplitude is essentially unchanged. From the 10d perspective, the difference is that instead of computing 4-graviton scattering one should consider processes involving the RR 1-form field. Their worldline vertex operators are directly related [108], hence the structure of the 1-loop integral is essentially unchanged, and the changes only involve the kinematic tensor structure. This computation is directly not available in the literature, but it can be regarded as a 10d decomposition of the computation of the 11d R^4 correction from the 4-point 1-loop superparticle amplitude in [109]. This suffices to show the emergence from the SDC tower of the required higher derivative terms beyond the purely gravitational sector.

5 Conclusions

In this work we have clarified the links among the different avatars of the species scale in the swampland program: (i) as cutoff of the effective theory specially in the presence of the infinite SDC tower of light particles near infinite distance points in moduli space, (ii) the scale at which terms in the infinite series of higher derivative corrections in the gravitational sector start competing with the two-derivative level, (iii) as the size of the smallest black hole which admits a description within the EFT.

Our analysis uses small black holes, whose cores probe infinite distance limits in field space, at which we can characterize the SDC tower of states leading to the species scale in avatar (i). Upon integrating out the tower of states, the theory generates a series

of higher-dimensional operators, naturally leading to avatar (ii) of the species scale. Finally, including these corrections in the black hole solution triggers the appearance of a stretched horizon of a size given by the species scale in avatar (iii).¹⁵

We have provided quantitative examples of this phenomenon, both in a class of 4d small black holes, and in the fascinating case of 10d type IIA D0-branes. The latter case turned out to be highly non-trivial, as the appearance of the species scale horizon required understanding of 10d 8-derivative terms involving not only the curvature but also the RR 2-form field strength. The nice packaging of those terms into an 11d R^4 correction is a tantalizing hint that the species scale can serve as a powerful probe of the UV physics of M-theory.

The 10d D0-brane example displays an interesting feature: the objects making up the small black hole are precisely of the same kind as those in the SDC tower for the corresponding infinite distance limit. This feature arises in the microscopic explanation of the entropy for other black holes in string theory; but the remarkable fact about the D0-brane case is that the higher derivative terms coming from integrating out the tower of D0-branes are crucial to generate a geometric horizon for the D0-brane black hole itself!

This motivates us to entertain the following picture. According to the most radical interpretation of the Emergence Proposal [35–38], not only these higher derivative terms but the very dynamics of gravity would emerge from integrating out the towers of states. In this spirit, the emergence of species scale stretched horizons studied in this work would potentially apply to any black hole horizon. This suggests a possible general picture for the emergence of horizons in quantum gravity: In the UV, the natural quantum system to consider would be that formed by the states in the tower, with no dynamical gravity. When going down to the IR, the species in the tower are integrated out. It is reasonable to expect that this effective description cannot describe the previous physical system in full detail, but it should be able to codify its coarse-grained properties, i.e., its macrostate. How is this made possible? In the IR, gravity emerges and the system would be replaced by a black hole that precisely reproduces

¹⁵It is worth noticing that we consider BPS small black holes, and that the quantum gravity effects we are taking into account come in the form of supersymmetry protected higher derivative corrections. It would be interesting to see how our results might extend to cases with less supersymmetry.

this macrostate. In this sense, the emergence of a horizon would be somehow required for the effective description to be able to keep the macroscopic information about the formerly considered physical system.

It is also tantalizing to speculate that the entropy of the black hole is nothing but a reflection of the entanglement entropy of the species after their degrees of freedom are traced out. This goes in the spirit of ER=EPR [110], the N-portrait picture [111], or of the literature on thermal M(atrix) theory black holes, see e.g. [112–118] (also [119, 120] for more recent developments). On a related note, it would be interesting to understand if and how this system made out of species could radiate and reproduce the evaporation of the black hole, perhaps along the lines of [31].

Back to more earthly matters, our work suggest several interesting questions for further research, for instance:

- We have focused on the interpretations of the species scale in asymptotic regions of moduli space, which are those naturally explored by small black holes. It would be interesting to extend our concept of unification of the three notions of species scale and gain a similar understanding in the interior of moduli space, connecting with the proposal in [30, 47, 48, 121], perhaps using large black hole probes as in [28].

- Even in asymptotic regimes, there is a strong ongoing activity in understanding the precise treatment of the species scale in the process of integrating out the SDC tower of states, with different proposals [39–43]. We hope the appearance of such computations in our setup can serve to clarify the proper physical procedures.

- The non-trivial constraints on higher derivative corrections for the appearance of stretched horizons are reminiscent of similar conditions derived from positivity constraints in the swampland program or in the S-matrix bootstrap, see e.g. [62, 122–124]. It would be interesting to explore this possible connections further.

- The $\text{AdS}_2 \times \mathbf{S}^8$ ansatz in the entropy function analysis suggests, via holography, the possible appearance of IR superconformal behaviour in the worldvolume theory on the D0-brane, once the equivalent to the higher derivative corrections are taken into account. It would be interesting to find further quantitative evidence for this proposal from the field theory side.

- It would be interesting to see whether some of our results extend to lower codimension cases. Higher-dimensional objects that probe infinite distance in field space

and are singular in the EFT (such as the End-of-the-World branes of [20, 24, 26, 125–128]¹⁶) could, in the same way as the small black holes, be shown to develop a species scale sized horizon using a modified version of the entropy function formalism.

- In some instances, higher derivative corrections have been shown to create new singularities on the horizon of large black holes [139]. This is to be put in contrast with our work, where curvature corrections instead smooth out singularities. It would be interesting to better understand the mechanisms that make it such that curvature corrections can seemingly both create and smooth out singularities in the EFT.

- There are interesting recent development regarding the possibility of log corrections to the species scale [41, 48, 85]. In this respect, it is tantalizing that our microscopic considerations in section 3.1 lead to such corrections. It would be interesting to exploit our techniques to shed some light on this question.

We hope to come back to these questions in the near future.

Acknowledgments

We thank Roberta Angius, Alberto Castellano, Lucia Gómez Cordova, Jesús Huertas, Luis Ibáñez, Shota Komatsu, Wolfgang Lerche, Luca Melotti, Fernando Marchesano, Jake McNamara, Miguel Montero, Tomás Ortín and Irene Valenzuela for useful conversations. The work of M.D. and A.U. is supported through the grants CEX2020-001007-S and PID2021-123017NB-I00, funded by MCIN/AEI/10.13039/501100011033 and by ERDF A way of making Europe.

A Tensor Structure of R^4 Corrections

In this appendix we recall the form of the $t_8 t_8 R^4$ and $\epsilon_{10} \epsilon_{10} R^4$ terms arising in 10d type IIA theory (and closely related to those in 11d M-theory). We follow closely the notation in [36].

Let us start with the $t_8 t_8 R^4$ structure, which denotes

$$t_8 t_8 R^4 = t_8^{A_1 \dots A_8} t_8^{B_1 \dots B_8} R_{A_1 A_2 B_1 B_2} \dots R_{A_7 A_8 B_7 B_8}, \quad (\text{A.1})$$

¹⁶For the related topic of solutions in theories with dynamical tadpoles, see [129–132] for early work and [133–138] for related recent developments.

where the t_8 tensor can be written in terms of the metric as¹⁷

$$\begin{aligned}
t_8^{A_1 \dots A_8} = \frac{1}{5} & \left[-2 \left(g^{A_1 A_3} g^{A_2 A_4} g^{A_5 A_7} g^{A_6 A_8} + g^{A_1 A_5} g^{A_2 A_6} g^{A_3 A_7} g^{A_4 A_8} + g^{A_1 A_7} g^{A_2 A_8} g^{A_3 A_5} g^{A_4 A_6} \right) \right. \\
& + 8 \left(g^{A_2 A_3} g^{A_4 A_5} g^{A_6 A_7} g^{A_8 A_1} + g^{A_2 A_5} g^{A_6 A_3} g^{A_4 A_7} g^{A_8 A_1} + g^{A_2 A_5} g^{A_6 A_7} g^{A_8 A_3} g^{A_4 A_1} \right) \\
& \left. - (A_1 \leftrightarrow A_2) - (A_3 \leftrightarrow A_4) - (A_5 \leftrightarrow A_6) - (A_7 \leftrightarrow A_8) \right].
\end{aligned} \tag{A.2}$$

After contracting all the metrics in t_8 with the Riemann tensors in (A.1), the $t_8 t_8 R^4$ structure can be written as:

$$\begin{aligned}
t_8 t_8 R^4 = 12 & \left(R^{A_1 A_2 A_3 A_4} R_{A_1 A_2 A_3 A_4} \right)^2 \\
& + 192 R_{A_1}^{A_5 A_6} R^{A_1 A_2 A_3 A_4} R_{A_2}^{A_7 A_8} R_{A_5 A_7 A_6 A_8} \\
& - 192 R_{A_1 A_2 A_3}^{A_5} R^{A_1 A_2 A_3 A_4} R_{A_4}^{A_6 A_7 A_8} R_{A_5 A_6 A_7 A_8} \\
& + 24 R_{A_1 A_2}^{A_5 A_6} R^{A_1 A_2 A_3 A_4} R_{A_3 A_4}^{A_7 A_8} R_{A_5 A_6 A_7 A_8} \\
& + 384 R_{A_1}^{A_5 A_6} R^{A_1 A_2 A_3 A_4} R_{A_2}^{A_7 A_8} R_{A_4 A_8 A_5 A_7} \\
& - 96 R_{A_1 A_2}^{A_5 A_6} R^{A_1 A_2 A_3 A_4} R_{A_3 A_5}^{A_7 A_8} R_{A_4 A_6 A_7 A_8}.
\end{aligned} \tag{A.3}$$

Similarly, $\epsilon_{10} \epsilon_{10} R^4$ structure denotes

$$\begin{aligned}
\epsilon_{10} \epsilon_{10} R^4 & = \epsilon_{10}^{C_1 C_2 A_1 \dots A_8} \epsilon_{10}^{C_1 C_2 B_1 \dots B_8} R^{B_1 B_2}_{A_1 A_2} \dots R^{B_7 B_8}_{A_7 A_8} \\
& = -2 \cdot 8! \delta^{A_1}_{[B_1} \dots \delta^{A_8}_{B_8]} R^{B_1 B_2}_{A_1 A_2} \dots R^{B_7 B_8}_{A_7 A_8},
\end{aligned} \tag{A.4}$$

where, in the last line, we have used the usual identity for contracting Levi-Civita tensors (c.f. equation (A.1) in [36]). Contracting all the Kronecker delta functions

¹⁷Notice the change of prefactor with respect to equation (B.3) of [36], so that, after expressing this structure as contractions of four Riemann tensors, the result matches that in [95].

with the Riemann tensors in (A.4), the $\epsilon_{10}\epsilon_{10}R^4$ structure can be written as:

$$\begin{aligned}
& \frac{1}{8}\epsilon_{10}\epsilon_{10}R^4 = -12 \left(R^{A_1A_2A_3A_4} R_{A_1A_2A_3A_4} \right)^2 \\
& - 192 R_{A_1}^{A_5} R_{A_3}^{A_6} R^{A_1A_2A_3A_4} R_{A_2}^{A_7} R_{A_4}^{A_8} R_{A_5A_7A_6A_8} \\
& + 192 R_{A_1A_2A_3}^{A_5} R^{A_1A_2A_3A_4} R_{A_4}^{A_6A_7A_8} R_{A_5A_6A_7A_8} \\
& - 24 R_{A_1A_2}^{A_5A_6} R^{A_1A_2A_3A_4} R_{A_3A_4}^{A_7A_8} R_{A_5A_6A_7A_8} \\
& + 384 R_{A_1A_2}^{A_5A_6} R^{A_1A_2A_3A_4} R_{A_3}^{A_7} R_{A_5}^{A_8} R_{A_4A_8A_6A_7} \\
& + 384 R_{A_1}^{A_5} R_{A_3}^{A_6} R^{A_1A_2A_3A_4} R_{A_2}^{A_7} R_{A_5}^{A_8} R_{A_4A_7A_6A_8} - 768 R^{A_1A_2} R_{A_1}^{A_3} R_{A_2}^{A_4} R_{A_3}^{A_5A_6A_7} R_{A_4A_5A_6A_7} \\
& + 384 R^{A_1A_2} R_{A_1}^{A_3A_4A_5} R_{A_2A_3}^{A_6A_7} R_{A_4A_5A_6A_7} + 96 R_{A_1A_2} R^{A_1A_2} R_{A_3A_4A_5A_6} R^{A_3A_4A_5A_6} \\
& - 1536 R^{A_1A_2} R_{A_1}^{A_3A_4A_5} R_{A_2}^{A_6} R_{A_4}^{A_7} R_{A_3A_7A_5A_6} + 768 R^{A_1A_2} R^{A_3A_4} R_{A_1}^{A_5} R_{A_2}^{A_6} R_{A_3A_5A_4A_6} \\
& - 768 R_{A_1}^{A_3} R^{A_1A_2} R_{A_2}^{A_4A_5A_6} R_{A_3A_4A_5A_6} - 32 R R_{A_1A_2}^{A_5A_6} R^{A_1A_2A_3A_4} R_{A_3A_4A_5A_6} \\
& + 128 R R_{A_1}^{A_5} R_{A_3}^{A_6} R^{A_1A_2A_3A_4} R_{A_2A_6A_4A_5} - 768 R^{A_1A_2} R^{A_3A_4} R_{A_1}^{A_5} R_{A_3}^{A_6} R_{A_2A_5A_4A_6} \\
& - 384 R^{A_1A_2} R^{A_3A_4} R_{A_1A_3}^{A_5A_6} R_{A_2A_4A_5A_6} + 1536 R_{A_1}^{A_3} R^{A_1A_2} R^{A_4A_5} R_{A_2A_4A_3A_5} \\
& + 384 R R^{A_1A_2} R_{A_1}^{A_3A_4A_5} R_{A_2A_3A_4A_5} - 24 R^2 R_{A_1A_2A_3A_4} R^{A_1A_2A_3A_4} \\
& - 384 R R^{A_1A_2} R^{A_3A_4} R_{A_1A_3A_2A_4} - 192 \left(R_{A_1A_2} R^{A_1A_2} \right)^2 + 384 R_{A_1}^{A_3} R^{A_1A_2} R_{A_2}^{A_4} R_{A_3A_4} \\
& - 256 R R_{A_1}^{A_3} R^{A_1A_2} R_{A_2A_3} + 96 R^2 R_{A_1A_2} R^{A_1A_2} - 4 R^4.
\end{aligned} \tag{A.5}$$

Evaluating on Mathematica each of the contractions in (A.3) and (A.5) on the $\text{AdS}_2 \times \mathbf{S}^8$ ansatz in (4.1), adding them up, and substituting $v = v_2$ and $\beta = v_2/v_1$, we get to the result reported in equation (4.28).

As we used in section 4.4, both the $\hat{t}_8\hat{t}_8\hat{R}^4$ and the $\frac{1}{24}\epsilon_{11}\epsilon_{11}\hat{R}^4$ structures appearing in the M-theory effective action in equation (4.31) look exactly the same as the terms we just discussed. This is, once written in terms of contractions of 11d Riemman tensors, they precisely reduce to the ones in (A.3) and (A.5). Evaluating these contractions on the 11d metric ansatz in (4.34) with Mathematica, we recover the result in (4.36).

References

- [1] E. Palti, *The Swampland: Introduction and Review*, *Fortsch. Phys.* **67** (2019), no. 6 1900037, [[arXiv:1903.06239](#)].
- [2] M. van Beest, J. Calderón-Infante, D. Mirfendereski, and I. Valenzuela, *Lectures on the Swampland Program in String Compactifications*, [arXiv:2102.01111](#).
- [3] M. Graña and A. Herráez, *The Swampland Conjectures: A Bridge from Quantum Gravity to Particle Physics*, *Universe* **7** (2021), no. 8 273, [[arXiv:2107.00087](#)].
- [4] N. B. Agmon, A. Bedroya, M. J. Kang, and C. Vafa, *Lectures on the string landscape and the Swampland*, [arXiv:2212.06187](#).
- [5] G. Dvali, *Black Holes and Large N Species Solution to the Hierarchy Problem*, *Fortsch. Phys.* **58** (2010) 528–536, [[arXiv:0706.2050](#)].
- [6] G. Dvali and C. Gomez, *Quantum Information and Gravity Cutoff in Theories with Species*, *Phys. Lett. B* **674** (2009) 303–307, [[arXiv:0812.1940](#)].
- [7] G. Dvali and D. Lust, *Evaporation of Microscopic Black Holes in String Theory and the Bound on Species*, *Fortsch. Phys.* **58** (2010) 505–527, [[arXiv:0912.3167](#)].
- [8] G. Dvali and C. Gomez, *Species and Strings*, [arXiv:1004.3744](#).
- [9] G. Dvali, C. Gomez, and D. Lust, *Black Hole Quantum Mechanics in the Presence of Species*, *Fortsch. Phys.* **61** (2013) 768–778, [[arXiv:1206.2365](#)].
- [10] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, *Nucl. Phys.* **B766** (2007) 21–33, [[hep-th/0605264](#)].
- [11] M. Etheredge, B. Heidenreich, S. Kaya, Y. Qiu, and T. Rudelius, *Sharpening the Distance Conjecture in diverse dimensions*, *JHEP* **12** (2022) 114, [[arXiv:2206.04063](#)].
- [12] D. van de Heisteeg, C. Vafa, M. Wiesner, and D. H. Wu, *Bounds on Field Range for Slowly Varying Positive Potentials*, [arXiv:2305.07701](#).
- [13] M. Etheredge, B. Heidenreich, J. McNamara, T. Rudelius, I. Ruiz, and I. Valenzuela, *Running Decompactification, Sliding Towers, and the Distance Conjecture*, [arXiv:2306.16440](#).
- [14] J. Calderón-Infante, A. Castellano, A. Herráez, and L. E. Ibáñez, *Entropy Bounds and the Species Scale Distance Conjecture*, [arXiv:2306.16450](#).

- [15] S. Ferrara, R. Kallosh, and A. Strominger, *$N=2$ extremal black holes*, *Phys. Rev. D* **52** (1995) R5412–R5416, [[hep-th/9508072](#)].
- [16] S. Ferrara and R. Kallosh, *Supersymmetry and attractors*, *Phys. Rev. D* **54** (1996) 1514–1524, [[hep-th/9602136](#)].
- [17] S. Ferrara and R. Kallosh, *Universality of supersymmetric attractors*, *Phys. Rev. D* **54** (1996) 1525–1534, [[hep-th/9603090](#)].
- [18] S. Ferrara, G. W. Gibbons, and R. Kallosh, *Black holes and critical points in moduli space*, *Nucl. Phys. B* **500** (1997) 75–93, [[hep-th/9702103](#)].
- [19] Y. Hamada, M. Montero, C. Vafa, and I. Valenzuela, *Finiteness and the Swampland*, [arXiv:2111.00015](#).
- [20] R. Angius, J. Calderón-Infante, M. Delgado, J. Huertas, and A. M. Uranga, *At the end of the world: Local Dynamical Cobordism*, *JHEP* **06** (2022) 142, [[arXiv:2203.11240](#)].
- [21] R. Angius, J. Huertas, and A. M. Uranga, *Small Black Hole Explosions*, [arXiv:2303.15903](#).
- [22] G. Buratti, J. Calderón, and A. M. Uranga, *Transplanckian axion monodromy!?*, *JHEP* **05** (2019) 176, [[arXiv:1812.05016](#)].
- [23] S. Lanza, F. Marchesano, L. Martucci, and I. Valenzuela, *Swampland Conjectures for Strings and Membranes*, [arXiv:2006.15154](#).
- [24] G. Buratti, M. Delgado, and A. M. Uranga, *Dynamical tadpoles, stringy cobordism, and the SM from spontaneous compactification*, *JHEP* **06** (2021) 170, [[arXiv:2104.02091](#)].
- [25] S. Lanza, F. Marchesano, L. Martucci, and I. Valenzuela, *The EFT stringy viewpoint on large distances*, [arXiv:2104.05726](#).
- [26] G. Buratti, J. Calderón-Infante, M. Delgado, and A. M. Uranga, *Dynamical Cobordism and Swampland Distance Conjectures*, *JHEP* **10** (2021) 037, [[arXiv:2107.09098](#)].
- [27] R. Angius, M. Delgado, and A. M. Uranga, *Dynamical Cobordism and the beginning of time: supercritical strings and tachyon condensation*, *JHEP* **08** (2022) 285, [[arXiv:2207.13108](#)].

- [28] M. Delgado, M. Montero, and C. Vafa, *Black holes as probes of moduli space geometry*, *JHEP* **04** (2023) 045, [[arXiv:2212.08676](#)].
- [29] J. Huertas and A. M. Uranga, *Aspects of dynamical cobordism in AdS/CFT*, *JHEP* **08** (2023) 140, [[arXiv:2306.07335](#)].
- [30] N. Cribiori, D. Lust, and G. Staudt, *Black hole entropy and moduli-dependent species scale*, [arXiv:2212.10286](#).
- [31] N. Cribiori, D. Lust, and C. Montella, *Species Entropy and Thermodynamics*, [arXiv:2305.10489](#).
- [32] G. Dvali and M. Redi, *Black Hole Bound on the Number of Species and Quantum Gravity at LHC*, *Phys. Rev. D* **77** (2008) 045027, [[arXiv:0710.4344](#)].
- [33] P. A. Cano, P. F. Ramírez, and A. Ruipérez, *The small black hole illusion*, *JHEP* **03** (2020) 115, [[arXiv:1808.10449](#)].
- [34] P. A. Cano, A. Murcia, P. F. Ramírez, and A. Ruipérez, *On small black holes, KK monopoles and solitonic 5-branes*, *JHEP* **05** (2021) 272, [[arXiv:2102.04476](#)].
- [35] B. Heidenreich, M. Reece, and T. Rudelius, *The Weak Gravity Conjecture and Emergence from an Ultraviolet Cutoff*, *Eur. Phys. J. C* **78** (2018), no. 4 337, [[arXiv:1712.01868](#)].
- [36] T. W. Grimm, E. Palti, and I. Valenzuela, *Infinite Distances in Field Space and Massless Towers of States*, *JHEP* **08** (2018) 143, [[arXiv:1802.08264](#)].
- [37] B. Heidenreich, M. Reece, and T. Rudelius, *Emergence of Weak Coupling at Large Distance in Quantum Gravity*, *Phys. Rev. Lett.* **121** (2018), no. 5 051601, [[arXiv:1802.08698](#)].
- [38] P. Corvilain, T. W. Grimm, and I. Valenzuela, *The Swampland Distance Conjecture for Kähler moduli*, *JHEP* **08** (2019) 075, [[arXiv:1812.07548](#)].
- [39] F. Marchesano and L. Melotti, *EFT strings and emergence*, *JHEP* **02** (2023) 112, [[arXiv:2211.01409](#)].
- [40] A. Castellano, A. Herráez, and L. E. Ibáñez, *The emergence proposal in quantum gravity and the species scale*, *JHEP* **06** (2023) 047, [[arXiv:2212.03908](#)].
- [41] R. Blumenhagen, A. Gligovic, and A. Paraskevopoulou, *The Emergence Proposal and the Emergent String*, [arXiv:2305.10490](#).

- [42] R. Blumenhagen, N. Cribiori, A. Gligovic, and A. Paraskevopoulou, *Demystifying the Emergence Proposal*, [arXiv:2309.11551](#).
- [43] R. Blumenhagen, N. Cribiori, A. Gligovic, and A. Paraskevopoulou, *The Emergent M-theory Limit*, [arXiv:2309.11554](#).
- [44] S. D. Mathur, *The Fuzzball proposal for black holes: An Elementary review*, *Fortsch. Phys.* **53** (2005) 793–827, [[hep-th/0502050](#)].
- [45] I. Bena and N. P. Warner, *Black holes, black rings and their microstates*, *Lect. Notes Phys.* **755** (2008) 1–92, [[hep-th/0701216](#)].
- [46] B. D. Chowdhury and A. Virmani, *Modave Lectures on Fuzzballs and Emission from the D1-D5 System*, in *5th Modave Summer School in Mathematical Physics*, 1, 2010. [arXiv:1001.1444](#).
- [47] D. van de Heisteeg, C. Vafa, M. Wiesner, and D. H. Wu, *Moduli-dependent Species Scale*, [arXiv:2212.06841](#).
- [48] D. van de Heisteeg, C. Vafa, and M. Wiesner, *Bounds on Species Scale and the Distance Conjecture*, [arXiv:2303.13580](#).
- [49] B. de Wit, J. Hoppe, and H. Nicolai, *On the Quantum Mechanics of Supermembranes*, *Nucl. Phys. B* **305** (1988) 545.
- [50] M. R. Douglas, D. N. Kabat, P. Pouliot, and S. H. Shenker, *D-branes and short distances in string theory*, *Nucl. Phys.* **B485** (1997) 85–127, [[hep-th/9608024](#)].
- [51] T. Banks, W. Fischler, S. H. Shenker, and L. Susskind, *M theory as a matrix model: A Conjecture*, *Phys. Rev. D* **55** (1997) 5112–5128, [[hep-th/9610043](#)].
- [52] N. Izhaki, J. M. Maldacena, J. Sonnenschein, and S. Yankielowicz, *Supergravity and the large N limit of theories with sixteen supercharges*, *Phys. Rev. D* **58** (1998) 046004, [[hep-th/9802042](#)].
- [53] T. Ortiz, H. Samtleben, and D. Tsimpis, *Matrix model holography*, *JHEP* **12** (2014) 096, [[arXiv:1410.0487](#)].
- [54] **Monte Carlo String/M-theory (MCSMC) Collaboration**, S. Pateloudis, G. Bergner, M. Hanada, E. Rinaldi, A. Schäfer, P. Vranas, H. Watanabe, and N. Bodendorfer, *Precision test of gauge/gravity duality in D0-brane matrix model at low temperature*, *JHEP* **03** (2023) 071, [[arXiv:2210.04881](#)].

- [55] A. Sinha and N. V. Suryanarayana, *Extremal single-charge small black holes: Entropy function analysis*, *Class. Quant. Grav.* **23** (2006) 3305–3322, [[hep-th/0601183](#)].
- [56] A. Sen, *Extremal black holes and elementary string states*, *Mod. Phys. Lett. A* **10** (1995) 2081–2094, [[hep-th/9504147](#)].
- [57] A. Dabholkar and S. Nampuri, *Quantum black holes*, *Lect. Notes Phys.* **851** (2012) 165–232, [[arXiv:1208.4814](#)].
- [58] T. Mohaupt, *Black hole entropy, special geometry and strings*, *Fortsch. Phys.* **49** (2001) 3–161, [[hep-th/0007195](#)].
- [59] N. Cribiori, M. Dierigl, A. Gnechchi, D. Lust, and M. Scalisi, *Large and Small Non-extremal Black Holes, Thermodynamic Dualities, and the Swampland*, [arXiv:2202.04657](#).
- [60] N. Cribiori, A. Gnechchi, D. Lüüst, and M. Scalisi, *On the correspondence between black holes, domain walls and fluxes*, *JHEP* **05** (2023) 033, [[arXiv:2302.03054](#)].
- [61] N. Arkani-Hamed, L. Motl, A. Nicolis, and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060, [[hep-th/0601001](#)].
- [62] C. Cheung and G. N. Remmen, *Infrared Consistency and the Weak Gravity Conjecture*, *JHEP* **12** (2014) 087, [[arXiv:1407.7865](#)].
- [63] C. Cheung, J. Liu, and G. N. Remmen, *Proof of the Weak Gravity Conjecture from Black Hole Entropy*, *JHEP* **10** (2018) 004, [[arXiv:1801.08546](#)].
- [64] N. Arkani-Hamed, Y.-t. Huang, J.-Y. Liu, and G. N. Remmen, *Causality, unitarity, and the weak gravity conjecture*, *JHEP* **03** (2022) 083, [[arXiv:2109.13937](#)].
- [65] E. Lauria and A. Van Proeyen, *$\mathcal{N} = 2$ Supergravity in $D = 4, 5, 6$ Dimensions*, vol. 966. 3, 2020.
- [66] J. M. Maldacena, A. Strominger, and E. Witten, *Black hole entropy in M theory*, *JHEP* **12** (1997) 002, [[hep-th/9711053](#)].
- [67] G. Lopes Cardoso, B. de Wit, and T. Mohaupt, *Corrections to macroscopic supersymmetric black hole entropy*, *Phys. Lett. B* **451** (1999) 309–316, [[hep-th/9812082](#)].
- [68] G. Lopes Cardoso, B. de Wit, and T. Mohaupt, *Deviations from the area law for supersymmetric black holes*, *Fortsch. Phys.* **48** (2000) 49–64, [[hep-th/9904005](#)].

- [69] G. Lopes Cardoso, B. de Wit, and T. Mohaupt, *Macroscopic entropy formulae and nonholomorphic corrections for supersymmetric black holes*, *Nucl. Phys. B* **567** (2000) 87–110, [[hep-th/9906094](#)].
- [70] B. Pioline, *Lectures on black holes, topological strings and quantum attractors*, *Class. Quant. Grav.* **23** (2006) S981, [[hep-th/0607227](#)].
- [71] A. Sen, *Black hole entropy function and the attractor mechanism in higher derivative gravity*, *JHEP* **09** (2005) 038, [[hep-th/0506177](#)].
- [72] S.-J. Lee, W. Lerche, and T. Weigand, *Emergent Strings, Duality and Weak Coupling Limits for Two-Form Fields*, [arXiv:1904.06344](#).
- [73] S.-J. Lee, W. Lerche, and T. Weigand, *Emergent Strings from Infinite Distance Limits*, [arXiv:1910.01135](#).
- [74] R. Gopakumar and C. Vafa, *M theory and topological strings. 1.*, [hep-th/9809187](#).
- [75] R. Gopakumar and C. Vafa, *M theory and topological strings. 2.*, [hep-th/9812127](#).
- [76] M. B. Green, M. Gutperle, and P. Vanhove, *One loop in eleven-dimensions*, *Phys. Lett. B* **409** (1997) 177–184, [[hep-th/9706175](#)].
- [77] H. Ooguri and C. Vafa, *Summing up D instantons*, *Phys. Rev. Lett.* **77** (1996) 3296–3298, [[hep-th/9608079](#)].
- [78] M. Gutperle, *A Note on heterotic / type I-prime duality and D0-brane quantum mechanics*, *JHEP* **05** (1999) 007, [[hep-th/9903010](#)].
- [79] A. Collinucci, P. Soler, and A. M. Uranga, *Non-perturbative effects and wall-crossing from topological strings*, *JHEP* **11** (2009) 025, [[arXiv:0904.1133](#)].
- [80] C. Petersson, P. Soler, and A. M. Uranga, *D-instanton and polyinstanton effects from type I' D0-brane loops*, *JHEP* **06** (2010) 089, [[arXiv:1001.3390](#)].
- [81] N. Piazzalunga and A. M. Uranga, *M-theory interpretation of the real topological string*, *JHEP* **08** (2014) 054, [[arXiv:1405.6019](#)].
- [82] E. Gonzalo, L. E. Ibáñez, and A. M. Uranga, *Modular Symmetries and the Swampland Conjectures*, *JHEP* **05** (2019) 105, [[arXiv:1812.06520](#)].
- [83] I. Antoniadis, S. Ferrara, R. Minasian, and K. S. Narain, *R^{*4} couplings in M and type II theories on Calabi-Yau spaces*, *Nucl. Phys. B* **507** (1997) 571–588, [[hep-th/9707013](#)].

- [84] E. Witten, *String theory dynamics in various dimensions*, *Nucl. Phys. B* **443** (1995) 85–126, [[hep-th/9503124](#)].
- [85] N. Cribiori and D. Lüst, *A note on modular invariant species scale and potentials*, [arXiv:2306.08673](#).
- [86] D. Berenstein, *A Toy model for the AdS / CFT correspondence*, *JHEP* **07** (2004) 018, [[hep-th/0403110](#)].
- [87] V. Balasubramanian, J. de Boer, V. Jejjala, and J. Simon, *The Library of Babel: On the origin of gravitational thermodynamics*, *JHEP* **12** (2005) 006, [[hep-th/0508023](#)].
- [88] P. Erdős and J. Lehner, *The distribution of the number of summands in the partitions of a positive integer*, *Duke Math. J.* **8(2)** (1941) 335–345.
- [89] R. E. Crandall, “Probability density for partitions of n with k parts.” <https://www.reed.edu/physics/faculty/wheeler/documents/Quantum%20Mechanics/Miscellaneous%20Essays/Partitions%20&%20Separability/Crandall%20on%20Partitions.pdf>.
- [90] G. Skezeres, *Some asymptotic formulae in the theory of partitions (II)*, *Quart. J. of Math. (Oxford)* **4** (1953) 96–111.
- [91] T. Anous, J. L. Karczmarek, E. Mintun, M. Van Raamsdonk, and B. Way, *Areas and entropies in BFSS/gravity duality*, *SciPost Phys.* **8** (2020), no. 4 057, [[arXiv:1911.11145](#)].
- [92] D. Lüst, E. Palti, and C. Vafa, *AdS and the Swampland*, [arXiv:1906.05225](#).
- [93] M. B. Green and J. H. Schwarz, *Supersymmetrical String Theories*, *Phys. Lett. B* **109** (1982) 444–448.
- [94] D. J. Gross and E. Witten, *Superstring Modifications of Einstein’s Equations*, *Nucl. Phys. B* **277** (1986) 1.
- [95] M. T. Grisaru and D. Zanon, *σ Model Superstring Corrections to the Einstein-hilbert Action*, *Phys. Lett. B* **177** (1986) 347–351.
- [96] M. T. Grisaru, A. E. M. van de Ven, and D. Zanon, *Two-Dimensional Supersymmetric Sigma Models on Ricci Flat Kahler Manifolds Are Not Finite*, *Nucl. Phys. B* **277** (1986) 388–408.
- [97] M. T. Grisaru, A. E. M. van de Ven, and D. Zanon, *Four Loop Divergences for the*

- N=1 Supersymmetric Nonlinear Sigma Model in Two-Dimensions*, *Nucl. Phys. B* **277** (1986) 409–428.
- [98] T. W. Grimm, K. Mayer, and M. Weissenbacher, *Higher derivatives in Type II and M-theory on Calabi-Yau threefolds*, *JHEP* **02** (2018) 127, [[arXiv:1702.08404](#)].
- [99] M. B. Green and P. Vanhove, *D instantons, strings and M theory*, *Phys. Lett. B* **408** (1997) 122–134, [[hep-th/9704145](#)].
- [100] J. G. Russo and A. A. Tseytlin, *One loop four graviton amplitude in eleven-dimensional supergravity*, *Nucl. Phys. B* **508** (1997) 245–259, [[hep-th/9707134](#)].
- [101] J. Polchinski, *M theory and the light cone*, *Prog. Theor. Phys. Suppl.* **134** (1999) 158–170, [[hep-th/9903165](#)].
- [102] L. Susskind, *Holography in the flat space limit*, *AIP Conf. Proc.* **493** (1999), no. 1 98–112, [[hep-th/9901079](#)].
- [103] N. A. Nekrasov, *On the size of a graviton*, in *14th International Workshop on High-Energy Physics and Quantum Field Theory (QFTHEP 99)*, pp. 401–411, 10, 1999. [hep-th/9909213](#).
- [104] T. Crisford, G. T. Horowitz, and J. E. Santos, *Testing the Weak Gravity - Cosmic Censorship Connection*, *Phys. Rev. D* **97** (2018), no. 6 066005, [[arXiv:1709.07880](#)].
- [105] G. T. Horowitz and J. E. Santos, *Further evidence for the weak gravity — cosmic censorship connection*, *JHEP* **06** (2019) 122, [[arXiv:1901.11096](#)].
- [106] I. R. Klebanov and M. J. Strassler, *Supergravity and a confining gauge theory: Duality cascades and chi SB resolution of naked singularities*, *JHEP* **08** (2000) 052, [[hep-th/0007191](#)].
- [107] J. T. Liu and R. Minasian, *Higher-derivative couplings in string theory: dualities and the B-field*, *Nucl. Phys. B* **874** (2013) 413–470, [[arXiv:1304.3137](#)].
- [108] M. B. Green, M. Gutperle, and H. H. Kwon, *Light cone quantum mechanics of the eleven-dimensional superparticle*, *JHEP* **08** (1999) 012, [[hep-th/9907155](#)].
- [109] K. Peeters, J. Plefka, and S. Stern, *Higher-derivative gauge field terms in the M-theory action*, *JHEP* **08** (2005) 095, [[hep-th/0507178](#)].

- [110] J. Maldacena and L. Susskind, *Cool horizons for entangled black holes*, *Fortsch. Phys.* **61** (2013) 781–811, [[arXiv:1306.0533](#)].
- [111] G. Dvali and C. Gomez, *Black Hole’s Quantum N-Portrait*, *Fortsch. Phys.* **61** (2013) 742–767, [[arXiv:1112.3359](#)].
- [112] T. Banks, W. Fischler, I. R. Klebanov, and L. Susskind, *Schwarzschild black holes from matrix theory*, *Phys. Rev. Lett.* **80** (1998) 226–229, [[hep-th/9709091](#)].
- [113] I. R. Klebanov and L. Susskind, *Schwarzschild black holes in various dimensions from matrix theory*, *Phys. Lett. B* **416** (1998) 62–66, [[hep-th/9709108](#)].
- [114] G. T. Horowitz and E. J. Martinec, *Comments on black holes in matrix theory*, *Phys. Rev. D* **57** (1998) 4935–4941, [[hep-th/9710217](#)].
- [115] M. Li, *Matrix Schwarzschild black holes in large N limit*, *JHEP* **01** (1998) 009, [[hep-th/9710226](#)].
- [116] T. Banks, W. Fischler, I. R. Klebanov, and L. Susskind, *Schwarzschild black holes in matrix theory. 2.*, *JHEP* **01** (1998) 008, [[hep-th/9711005](#)].
- [117] L. Susskind, *Matrix theory black holes and the Gross-Witten transition*, [[hep-th/9805115](#)].
- [118] M. Li and E. J. Martinec, *Probing matrix black holes*, [[hep-th/9801070](#)].
- [119] A. V. Smilga, *Comments on thermodynamics of supersymmetric matrix models*, *Nucl. Phys. B* **818** (2009) 101–114, [[arXiv:0812.4753](#)].
- [120] T. Wiseman, *On black hole thermodynamics from super Yang-Mills*, *JHEP* **07** (2013) 101, [[arXiv:1304.3938](#)].
- [121] D. Andriot, *Bumping into the species scale with the scalar potential*, [[arXiv:2305.07480](#)].
- [122] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, and R. Rattazzi, *Causality, analyticity and an IR obstruction to UV completion*, *JHEP* **10** (2006) 014, [[hep-th/0602178](#)].
- [123] B. Bellazzini, C. Cheung, and G. N. Remmen, *Quantum Gravity Constraints from Unitarity and Analyticity*, *Phys. Rev. D* **93** (2016), no. 6 064076, [[arXiv:1509.00851](#)].
- [124] Y. Hamada, T. Noumi, and G. Shiu, *Weak Gravity Conjecture from Unitarity and Causality*, *Phys. Rev. Lett.* **123** (2019), no. 5 051601, [[arXiv:1810.03637](#)].

- [125] R. Blumenhagen, N. Cribiori, C. Kneissl, and A. Makridou, *Dynamical Cobordism of a Domain Wall and its Companion Defect 7-brane*, [arXiv:2205.09782](#).
- [126] R. Blumenhagen, N. Cribiori, C. Kneissl, and A. Makridou, *Dimensional Reduction of Cobordism and K-theory*, *JHEP* **03** (2023) 181, [[arXiv:2208.01656](#)].
- [127] I. Basile, S. Raucci, and S. Thomée, *Revisiting Dudas-Mourad Compactifications*, *Universe* **8** (2022), no. 10 544, [[arXiv:2209.10553](#)].
- [128] R. Blumenhagen, C. Kneissl, and C. Wang, *Dynamical Cobordism Conjecture: Solutions for End-of-the-World Branes*, [arXiv:2303.03423](#).
- [129] E. Dudas and J. Mourad, *Brane solutions in strings with broken supersymmetry and dilaton tadpoles*, *Phys. Lett. B* **486** (2000) 172–178, [[hep-th/0004165](#)].
- [130] R. Blumenhagen and A. Font, *Dilaton tadpoles, warped geometries and large extra dimensions for nonsupersymmetric strings*, *Nucl. Phys. B* **599** (2001) 241–254, [[hep-th/0011269](#)].
- [131] E. Dudas, J. Mourad, and C. Timirgaziu, *Time and space dependent backgrounds from nonsupersymmetric strings*, *Nucl. Phys. B* **660** (2003) 3–24, [[hep-th/0209176](#)].
- [132] E. Dudas, G. Pradisi, M. Nicolosi, and A. Sagnotti, *On tadpoles and vacuum redefinitions in string theory*, *Nucl. Phys. B* **708** (2005) 3–44, [[hep-th/0410101](#)].
- [133] S. Raucci, *On codimension-one vacua and string theory*, *Nucl. Phys. B* **985** (2022) 116002, [[arXiv:2206.06399](#)].
- [134] R. Antonelli and I. Basile, *Brane annihilation in non-supersymmetric strings*, *JHEP* **11** (2019) 021, [[arXiv:1908.04352](#)].
- [135] A. Mininno and A. M. Uranga, *Dynamical Tadpoles and Weak Gravity Constraints*, [arXiv:2011.00051](#).
- [136] I. Basile, *On String Vacua without Supersymmetry: brane dynamics, bubbles and holography*. PhD thesis, Scuola normale superiore di Pisa, Pisa, Scuola Normale Superiore, 2020. [arXiv:2010.00628](#).
- [137] I. Basile, *Supersymmetry breaking, brane dynamics and Swampland conjectures*, *JHEP* **10** (2021) 080, [[arXiv:2106.04574](#)].
- [138] J. Mourad and A. Sagnotti, *A 4D IIB Flux Vacuum and Supersymmetry Breaking. I. Fermionic Spectrum*, [arXiv:2206.03340](#).

- [139] G. T. Horowitz, M. Kolanowski, G. N. Remmen, and J. E. Santos, *Extremal Kerr Black Holes as Amplifiers of New Physics*, *Phys. Rev. Lett.* **131** (2023), no. 9 091402, [[arXiv:2303.07358](https://arxiv.org/abs/2303.07358)].